



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Search for the chiral magnetic effect with isobar collisions
at math

$\sqrt{s_{NN}} = 200$ GeV by the STAR Collaboration at the BNL Relativistic
Heavy Ion Collider

GeV by the STAR Collaboration at the BNL Relativistic
Heavy Ion Collider

M. S. Abdallah et al. (STAR Collaboration)

Phys. Rev. C **105**, 014901 — Published 3 January 2022

DOI: [10.1103/PhysRevC.105.014901](https://doi.org/10.1103/PhysRevC.105.014901)

1 Search for the Chiral Magnetic Effect with Isobar Collisions at $\sqrt{s_{NN}} = 200$ GeV by the
2 STAR Collaboration at RHIC

3 M. S. Abdallah,⁵ B. E. Aboona,⁵⁵ J. Adam,⁶ L. Adamczyk,² J. R. Adams,³⁹ J. K. Adkins,³⁰ G. Agakishiev,²⁸
4 I. Aggarwal,⁴¹ M. M. Aggarwal,⁴¹ Z. Ahammed,⁶⁰ I. Alekseev,^{3,35} D. M. Anderson,⁵⁵ A. Aparin,²⁸
5 E. C. Aschenauer,⁶ M. U. Ashraf,¹¹ F. G. Atetalla,²⁹ A. Attri,⁴¹ G. S. Averichev,²⁸ V. Bairathi,⁵³ W. Baker,¹⁰
6 J. G. Ball Cap,²⁰ K. Barish,¹⁰ A. Behera,⁵² R. Bellwied,²⁰ P. Bhagat,²⁷ A. Bhasin,²⁷ J. Bielcik,¹⁴ J. Bielcikova,³⁸
7 I. G. Bordyuzhin,³ J. D. Brandenburg,⁶ A. V. Brandin,³⁵ I. Bunzarov,²⁸ X. Z. Cai,⁵⁰ H. Caines,⁶³
8 M. Calderón de la Barca Sánchez,⁸ D. Cebra,⁸ I. Chakaberia,^{31,6} P. Chaloupka,¹⁴ B. K. Chan,⁹ F-H. Chang,³⁷
9 Z. Chang,⁶ N. Chankova-Bunzarova,²⁸ A. Chatterjee,¹¹ S. Chattopadhyay,⁶⁰ D. Chen,¹⁰ J. Chen,⁴⁹ J. H. Chen,¹⁸
10 X. Chen,⁴⁸ Z. Chen,⁴⁹ J. Cheng,⁵⁷ M. Chevalier,¹⁰ S. Choudhury,¹⁸ W. Christie,⁶ X. Chu,⁶ H. J. Crawford,⁷
11 M. Csanád,¹⁶ M. Daugherty,¹ T. G. Dedovich,²⁸ I. M. Deppner,¹⁹ A. A. Derevschikov,⁴³ A. Dhamija,⁴¹
12 L. Di Carlo,⁶² L. Didenko,⁶ P. Dixit,²² X. Dong,³¹ J. L. Drachenberg,¹ E. Duckworth,²⁹ J. C. Dunlop,⁶ N. Elsey,⁶²
13 J. Engelage,⁷ G. Eppley,⁴⁵ S. Esumi,⁵⁸ O. Evdokimov,¹² A. Ewigleben,³² O. Eyser,⁶ R. Fatemi,³⁰ F. M. Fawzi,⁵
14 S. Fazio,⁶ P. Federic,³⁸ J. Fedorisin,²⁸ C. J. Feng,³⁷ Y. Feng,⁴⁴ P. Filip,²⁸ E. Finch,⁵¹ Y. Fisyak,⁶ A. Francisco,⁶³
15 C. Fu,¹¹ L. Fulek,² C. A. Gagliardi,⁵⁵ T. Galatyuk,¹⁵ F. Geurts,⁴⁵ N. Ghimire,⁵⁴ A. Gibson,⁵⁹ K. Gopal,²³
16 X. Gou,⁴⁹ D. Grosnick,⁵⁹ A. Gupta,²⁷ W. Guryn,⁶ A. I. Hamad,²⁹ A. Hamed,⁵ Y. Han,⁴⁵ S. Harabasz,¹⁵
17 M. D. Harasty,⁸ J. W. Harris,⁶³ H. Harrison,³⁰ S. He,¹¹ W. He,¹⁸ X. H. He,²⁶ Y. He,⁴⁹ S. Heppelmann,⁸
18 S. Heppelmann,⁴² N. Herrmann,¹⁹ E. Hoffman,²⁰ L. Holub,¹⁴ Y. Hu,¹⁸ H. Huang,³⁷ H. Z. Huang,⁹ S. L. Huang,⁵²
19 T. Huang,³⁷ X. Huang,⁵⁷ Y. Huang,⁵⁷ T. J. Humanic,³⁹ G. Igo,^{9,*} D. Isenhower,¹ W. W. Jacobs,²⁵ C. Jena,²³
20 A. Jentsch,⁶ Y. Ji,³¹ J. Jia,^{6,52} K. Jiang,⁴⁸ X. Ju,⁴⁸ E. G. Judd,⁷ S. Kabana,⁵³ M. L. Kabir,¹⁰ S. Kagamaster,³²
21 D. Kalinkin,^{25,6} K. Kang,⁵⁷ D. Kapukchyan,¹⁰ K. Kauder,⁶ H. W. Ke,⁶ D. Keane,²⁹ A. Kechechyan,²⁸ M. Kelsey,⁶²
22 Y. V. Khyzhniak,³⁵ D. P. Kikola,⁶¹ C. Kim,¹⁰ B. Kimelman,⁸ D. Kincses,¹⁶ I. Kisel,¹⁷ A. Kiselev,⁶
23 A. G. Knospe,³² H. S. Ko,³¹ L. Kochenda,³⁵ L. K. Kosarzewski,¹⁴ L. Kramerik,¹⁴ P. Kravtsov,³⁵ L. Kumar,⁴¹
24 S. Kumar,²⁶ R. Kunnawalkam Elayavalli,⁶³ J. H. Kwasizur,²⁵ R. Lacey,⁵² S. Lan,¹¹ J. M. Landgraf,⁶
25 J. Lauret,⁶ A. Lebedev,⁶ R. Lednicky,^{28,38} J. H. Lee,⁶ Y. H. Leung,³¹ C. Li,⁴⁹ C. Li,⁴⁸ W. Li,⁴⁵ X. Li,⁴⁸
26 Y. Li,⁴⁸ Y. Li,⁵⁷ X. Liang,¹⁰ Y. Liang,²⁹ R. Licenik,³⁸ T. Lin,⁴⁹ Y. Lin,¹¹ M. A. Lisa,³⁹ F. Liu,¹¹ H. Liu,²⁵
27 H. Liu,¹¹ P. Liu,⁵² T. Liu,⁶³ X. Liu,³⁹ Y. Liu,⁵⁵ Z. Liu,⁴⁸ T. Ljubicic,⁶ W. J. Llope,⁶² R. S. Longacre,⁶
28 E. Loyd,¹⁰ N. S. Lukow,⁵⁴ X. F. Luo,¹¹ L. Ma,¹⁸ R. Ma,⁶ Y. G. Ma,¹⁸ N. Magdy,¹² D. Mallick,³⁶ S. Margetis,²⁹
29 C. Markert,⁵⁶ H. S. Matis,³¹ J. A. Mazer,⁴⁶ N. G. Minaev,⁴³ S. Mioduszewski,⁵⁵ B. Mohanty,³⁶ M. M. Mondal,⁵²
30 I. Mooney,⁶² D. A. Morozov,⁴³ A. Mukherjee,¹⁶ M. Nagy,¹⁶ J. D. Nam,⁵⁴ Md. Nasim,²² K. Nayak,¹¹ D. Neff,⁹
31 J. M. Nelson,⁷ D. B. Nemes,⁶³ M. Nie,⁴⁹ G. Nigmatkulov,³⁵ T. Niida,⁵⁸ R. Nishitani,⁵⁸ L. V. Nogach,⁴³
32 T. Nonaka,⁵⁸ A. S. Nunes,⁶ G. Odyniec,³¹ A. Ogawa,⁶ S. Oh,³¹ V. A. Okorokov,³⁵ B. S. Page,⁶ R. Pak,⁶ J. Pan,⁵⁵
33 A. Pandav,³⁶ A. K. Pandey,⁵⁸ Y. Panebratsev,²⁸ P. Parfenov,³⁵ B. Pawlik,⁴⁰ D. Pawlowska,⁶¹ C. Perkins,⁷
34 L. Pinsky,²⁰ R. L. Pintér,¹⁶ J. Pluta,⁶¹ B. R. Pokhrel,⁵⁴ G. Ponimatkin,³⁸ J. Porter,³¹ M. Posik,⁵⁴ V. Prozorova,¹⁴
35 N. K. Pruthi,⁴¹ M. Przybycien,² J. Putschke,⁶² H. Qiu,²⁶ A. Quintero,⁵⁴ C. Racz,¹⁰ S. K. Radhakrishnan,²⁹
36 N. Raha,⁶² R. L. Ray,⁵⁶ R. Reed,³² H. G. Ritter,³¹ M. Robotkova,³⁸ O. V. Rogachevskiy,²⁸ J. L. Romero,⁸
37 D. Roy,⁴⁶ L. Ruan,⁶ J. Rusnak,³⁸ A. K. Sahoo,²² N. R. Sahoo,⁴⁹ H. Sako,⁵⁸ S. Salur,⁴⁶ J. Sandweiss,^{63,*}
38 S. Sato,⁵⁸ W. B. Schmidke,⁶ N. Schmitz,³³ B. R. Schweid,⁵² F. Seck,¹⁵ J. Seger,¹³ M. Sergeeva,⁹ R. Seto,¹⁰
39 P. Seyboth,³³ N. Shah,²⁴ E. Shahaliev,²⁸ P. V. Shanmuganathan,⁶ M. Shao,⁴⁸ T. Shao,¹⁸ A. I. Sheikh,²⁹
40 D. Y. Shen,¹⁸ S. S. Shi,¹¹ Y. Shi,⁴⁹ Q. Y. Shou,¹⁸ E. P. Sichtermann,³¹ R. Sikora,² M. Simko,³⁸ J. Singh,⁴¹
41 S. Singha,²⁶ M. J. Skoby,⁴⁴ N. Smirnov,⁶³ Y. Söhngen,¹⁹ W. Solyst,²⁵ P. Sorensen,⁶ H. M. Spinka,^{4,*}
42 B. Srivastava,⁴⁴ T. D. S. Stanislaus,⁵⁹ M. Stefaniak,⁶¹ D. J. Stewart,⁶³ M. Strikhanov,³⁵ B. Stringfellow,⁴⁴
43 A. A. P. Suaide,⁴⁷ M. Sumner,³⁸ B. Summa,⁴² X. M. Sun,¹¹ X. Sun,¹² Y. Sun,⁴⁸ Y. Sun,²¹ B. Surrow,⁵⁴
44 D. N. Svirida,³ Z. W. Sweger,⁸ P. Szymanski,⁶¹ A. H. Tang,⁶ Z. Tang,⁴⁸ A. Taranenko,³⁵ T. Tarnowsky,³⁴
45 J. H. Thomas,³¹ A. R. Timmins,²⁰ D. Tlusty,¹³ T. Todoroki,⁵⁸ M. Tokarev,²⁸ C. A. Tomkiel,³² S. Trentalange,⁹
46 R. E. Tribble,⁵⁵ P. Tribedy,⁶ S. K. Tripathy,¹⁶ T. Truhlar,¹⁴ B. A. Trzeciak,¹⁴ O. D. Tsai,⁹ Z. Tu,⁶ T. Ullrich,⁶
47 D. G. Underwood,^{4,59} I. Upsal,⁴⁵ G. Van Buren,⁶ J. Vanek,³⁸ A. N. Vasiliev,⁴³ I. Vassiliev,¹⁷ V. Verkest,⁶²
48 F. Videbæk,⁶ S. Vokal,²⁸ S. A. Voloshin,⁶² F. Wang,⁴⁴ G. Wang,⁹ J. S. Wang,²¹ P. Wang,⁴⁸ Y. Wang,¹¹ Y. Wang,⁵⁷
49 Z. Wang,⁴⁹ J. C. Webb,⁶ P. C. Weidenkaff,¹⁹ L. Wen,⁹ G. D. Westfall,³⁴ H. Wieman,³¹ S. W. Wissink,²⁵ J. Wu,¹¹
50 J. Wu,²⁶ Y. Wu,¹⁰ B. Xi,⁵⁰ Z. G. Xiao,⁵⁷ G. Xie,³¹ W. Xie,⁴⁴ H. Xu,²¹ N. Xu,³¹ Q. H. Xu,⁴⁹ Y. Xu,⁴⁹ Z. Xu,⁶
51 Z. Xu,⁹ C. Yang,⁴⁹ Q. Yang,⁴⁹ S. Yang,⁴⁵ Y. Yang,³⁷ Z. Ye,⁴⁵ Z. Ye,¹² L. Yi,⁴⁹ K. Yip,⁶ Y. Yu,⁴⁹ H. Zbroszczyk,⁶¹
52 W. Zha,⁴⁸ C. Zhang,⁵² D. Zhang,¹¹ J. Zhang,⁴⁹ S. Zhang,¹² S. Zhang,¹⁸ X. P. Zhang,⁵⁷ Y. Zhang,²⁶ Y. Zhang,⁴⁸

Y. Zhang,¹¹ Z. J. Zhang,³⁷ Z. Zhang,⁶ Z. Zhang,¹² J. Zhao,⁴⁴ C. Zhou,¹⁸ X. Zhu,⁵⁷ M. Zurek,⁴ and M. Zyzak¹⁷

(STAR Collaboration)

¹Abilene Christian University, Abilene, Texas 79699

²AGH University of Science and Technology, FPACS, Cracow 30-059, Poland

³Alikhanov Institute for Theoretical and Experimental Physics NRC "Kurchatov Institute", Moscow 117218, Russia

⁴Argonne National Laboratory, Argonne, Illinois 60439

⁵American University of Cairo, New Cairo 11835, New Cairo, Egypt

⁶Brookhaven National Laboratory, Upton, New York 11973

⁷University of California, Berkeley, California 94720

⁸University of California, Davis, California 95616

⁹University of California, Los Angeles, California 90095

¹⁰University of California, Riverside, California 92521

¹¹Central China Normal University, Wuhan, Hubei 430079

¹²University of Illinois at Chicago, Chicago, Illinois 60607

¹³Creighton University, Omaha, Nebraska 68178

¹⁴Czech Technical University in Prague, FNSPE, Prague 115 19, Czech Republic

¹⁵Technische Universität Darmstadt, Darmstadt 64289, Germany

¹⁶ELTE Eötvös Loránd University, Budapest, Hungary H-1117

¹⁷Frankfurt Institute for Advanced Studies FIAS, Frankfurt 60438, Germany

¹⁸Fudan University, Shanghai, 200433

¹⁹University of Heidelberg, Heidelberg 69120, Germany

²⁰University of Houston, Houston, Texas 77204

²¹Huzhou University, Huzhou, Zhejiang 313000

²²Indian Institute of Science Education and Research (IISER), Berhampur 760010, India

²³Indian Institute of Science Education and Research (IISER) Tirupati, Tirupati 517507, India

²⁴Indian Institute Technology, Patna, Bihar 801106, India

²⁵Indiana University, Bloomington, Indiana 47408

²⁶Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou, Gansu 730000

²⁷University of Jammu, Jammu 180001, India

²⁸Joint Institute for Nuclear Research, Dubna 141 980, Russia

²⁹Kent State University, Kent, Ohio 44242

³⁰University of Kentucky, Lexington, Kentucky 40506-0055

³¹Lawrence Berkeley National Laboratory, Berkeley, California 94720

³²Lehigh University, Bethlehem, Pennsylvania 18015

³³Max-Planck-Institut für Physik, Munich 80805, Germany

³⁴Michigan State University, East Lansing, Michigan 48824

³⁵National Research Nuclear University MPhI, Moscow 115409, Russia

³⁶National Institute of Science Education and Research, HBNI, Jatni 752050, India

³⁷National Cheng Kung University, Tainan 70101

³⁸Nuclear Physics Institute of the CAS, Rez 250 68, Czech Republic

³⁹Ohio State University, Columbus, Ohio 43210

⁴⁰Institute of Nuclear Physics PAN, Cracow 31-342, Poland

⁴¹Panjab University, Chandigarh 160014, India

⁴²Pennsylvania State University, University Park, Pennsylvania 16802

⁴³NRC "Kurchatov Institute", Institute of High Energy Physics, Protvino 142281, Russia

⁴⁴Purdue University, West Lafayette, Indiana 47907

⁴⁵Rice University, Houston, Texas 77251

⁴⁶Rutgers University, Piscataway, New Jersey 08854

⁴⁷Universidade de São Paulo, São Paulo, Brazil 05314-970

⁴⁸University of Science and Technology of China, Hefei, Anhui 230026

⁴⁹Shandong University, Qingdao, Shandong 266237

⁵⁰Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800

⁵¹Southern Connecticut State University, New Haven, Connecticut 06515

⁵²State University of New York, Stony Brook, New York 11794

⁵³Instituto de Alta Investigación, Universidad de Tarapacá, Arica 1000000, Chile

⁵⁴Temple University, Philadelphia, Pennsylvania 19122

⁵⁵Texas A&M University, College Station, Texas 77843

⁵⁶University of Texas, Austin, Texas 78712

⁵⁷Tsinghua University, Beijing 100084

⁵⁸University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan

⁵⁹Valparaiso University, Valparaiso, Indiana 46383

⁶⁰Variable Energy Cyclotron Centre, Kolkata 700064, India

⁶¹Warsaw University of Technology, Warsaw 00-661, Poland

⁶²Wayne State University, Detroit, Michigan 48201

⁶³Yale University, New Haven, Connecticut 06520

(Dated: November 27, 2021)

The chiral magnetic effect (CME) is predicted to occur as a consequence of a local violation of \mathcal{P} and \mathcal{CP} symmetries of the strong interaction amidst a strong electro-magnetic field generated in relativistic heavy-ion collisions. Experimental manifestation of the CME involves a separation of positively and negatively charged hadrons along the direction of the magnetic field. Previous measurements of the CME-sensitive charge-separation observables remain inconclusive because of large background contributions. In order to better control the influence of signal and backgrounds, the STAR Collaboration performed a blind analysis of a large data sample of approximately 3.8 billion isobar collisions of $^{96}_{44}\text{Ru}+^{96}_{44}\text{Ru}$ and $^{96}_{40}\text{Zr}+^{96}_{40}\text{Zr}$ at $\sqrt{s_{\text{NN}}} = 200$ GeV. Prior to the blind analysis, the CME signatures are predefined as a significant excess of the CME-sensitive observables in Ru+Ru collisions over those in Zr+Zr collisions, owing to a larger magnetic field in the former. A precision down to 0.4% is achieved, as anticipated, in the relative magnitudes of the pertinent observables between the two isobar systems. Observed differences in the multiplicity and flow harmonics at the matching centrality indicate that the magnitude of the CME background is different between the two species. No CME signature that satisfies the predefined criteria has been observed in isobar collisions in this blind analysis.

* Deceased

CONTENTS

135	I. Introduction	5
136	II. Isobar data and blind analysis	6
137	A. Modality of isobar running at RHIC	6
138	B. Detector apparatus and data quality cuts	7
139	C. Blinding of data sets and preparation for analysis	8
140	D. Methods for isobar blind analysis	8
141	E. Quality assurance of the blind data	9
142	F. Methodology of uncertainty estimation	11
143	III. Centrality determination	11
144	IV. Observables for isobar blind analysis	14
145	A. $\Delta\gamma$ and mixed harmonics with second and third order event planes	15
146	B. Relative pseudorapidity dependence of $\Delta\gamma$	17
147	C. Invariant mass dependence of $\Delta\gamma$	17
148	D. $\Delta\gamma$ with spectator and participant planes (approach-I)	18
149	E. $\Delta\gamma$ with participant and spectator planes (approach-II)	19
150	F. R_{Ψ_2} variable	19
151	V. Isobar blind analysis results	20
152	A. $\Delta\gamma$ measurements with TPC event plane (Group-1)	21
153	B. Mixed harmonic measurements (Group-2)	24
154	C. Differential $\Delta\gamma$ measurements in pseudorapidity (Group-2)	28
155	D. Differential $\Delta\gamma$ measurements in invariant mass (Group-3)	29
156	E. CME fraction utilizing spectator and participant planes: approach-I (Group-3)	30
157	F. Ratio of $(\Delta\gamma/v_2)$ between two isobar collisions (Group-4)	33
158	G. CME fraction utilizing spectator and participant planes: approach-II (Group 4)	34
159	H. R_{Ψ_2} measurements (Group-5)	36
160	I. Summary and discussions	37
161	VI. Post blinding	39
162	VII. Conclusion	40
163	VIII. Acknowledgement	40
164	References	40

I. INTRODUCTION

In heavy-ion collisions, an exciting possibility is that regions may be briefly formed in which parity (\mathcal{P}) and charge-parity (\mathcal{CP}) symmetries are locally violated by the strong interaction [1–3]. This would lead to an imbalance between the numbers of right- and left-handed (anti-)quarks. It is demonstrated that if a sufficiently strong (electro-)magnetic field exists in such a region (as it may be in off-center heavy-ion collisions, generated chiefly by the protons in the two nuclei [4–11]) the net effect would be a separation of charges along the direction of the magnetic field [12–14]. This separation of charges is called the Chiral Magnetic Effect (CME). If an observation of the CME could be clearly established in heavy-ion collisions, it would imply the existence of these \mathcal{CP} -violating regions, the restoration of the approximate chiral symmetry in the Quark Gluon Plasma (QGP) medium, and the action of an ultra-strong magnetic field on the collision region (see Refs. [15, 16] for reviews). A precision experimental test of the CME has been an important scientific goal of Brookhaven National Laboratory’s Relativistic Heavy-Ion Collider (RHIC) program over the past decade. CME is also being explored in condensed matter systems [17, 18].

Over the years, extensive efforts have been invested to measure the CME-sensitive charge separation perpendicular to the reaction plane (RP, defined by the collision impact parameter and the beam direction) in heavy-ion collisions [19–29] (also see reviews in Refs. [15, 16, 30–35]). In order to quantify the CME-induced charge transport and other modes of collective motion of the QGP, the azimuthal distribution of final-state particles is often Fourier-decomposed as

$$\frac{dN_\alpha}{d\phi^*} \approx \frac{N_\alpha}{2\pi} [1 + 2v_{1,\alpha} \cos(\phi^*) + 2a_{1,\alpha} \sin(\phi^*) + 2v_{2,\alpha} \cos(2\phi^*) + \dots], \quad (1)$$

where $\phi^* = \phi - \Psi_{\text{RP}}$, with ϕ and Ψ_{RP} being the azimuthal angles of a particle and of the RP, respectively. The subscript α (+ or –) denotes the charge sign of a particle. The coefficients v_1 and v_2 are called “directed flow” and “elliptic flow”, respectively. The v_n are functions of transverse momentum (p_T) and pseudorapidity (η). The coefficient a_1 (with $a_{1,-} = -a_{1,+}$) characterizes the electric charge separation with respect to the RP which is correlated with the direction of magnetic field [5–7]. The most widely used observable in the CME search is the “ γ correlator,” originally proposed in Ref. [36],

$$\gamma_{\alpha\beta} = \langle \cos(\phi_\alpha + \phi_\beta - 2\Psi_{\text{RP}}) \rangle, \quad (2)$$

where ϕ_α and ϕ_β are the azimuthal angles of particles of interest (POIs). Here the averaging $\langle \dots \rangle$ is performed over the pairs of particles and over events. In order to eliminate charge-independent correlation backgrounds mainly from global momentum conservation [37, 38], the difference between the opposite-sign (OS) and same-sign (SS) γ correlators is considered,

$$\Delta\gamma = \gamma_{\text{os}} - \gamma_{\text{ss}}. \quad (3)$$

The $\Delta\gamma$ is sensitive to the preferential emission of positively and negatively charged particles to the opposite sides of the RP. The first measurements of non-zero $\Delta\gamma$ from the STAR (Solenoidal Tracker at RHIC) Collaboration in Au+Au and Cu+Cu collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV are reported in Refs. [19, 20]. In those publications, connections to expectations from CME-driven signals ($\Delta\gamma = 2a_1^2$) and flow-induced background due to resonance decays are identified as possible sources that contribute to $\Delta\gamma$. Subsequent measurements from RHIC [23, 24] and the LHC [21] at different energies have confirmed the observation of non-zero $\Delta\gamma$. Despite the theoretical progress, the quantification of the magnitudes of CME signals in heavy-ion collisions remains a challenge [39–48]. On the other hand, it is understood from phenomenological studies that measurements of $\Delta\gamma$ are dominated by backgrounds that are unrelated to the CME [37, 49–51]. The dominant backgrounds arise from intra-cluster correlations coupled with azimuthal anisotropy [36, 49–55]; namely,

$$\Delta\gamma_{\text{bgd}} = \frac{4N_{2\text{p}}}{N^2} \langle \cos(\phi_\alpha + \phi_\beta - 2\phi_{2\text{p}}) \rangle v_{2,2\text{p}}, \quad (4)$$

where $\phi_{2\text{p}}$ is the azimuthal angle of a correlated 2-particle cluster, $v_{2,2\text{p}}$ is the elliptic flow of such clusters, $N_{2\text{p}}$ is the number of those clusters, and N is the multiplicity of the POIs [34, 36, 49, 55]. An example for this is the correlations among the decay daughters of resonance particles carrying elliptic flow.

Collisions of small systems are often considered to provide a data-driven baseline for a background scenario [25]. In such collisions, the direction of the magnetic field is uncorrelated with azimuthal anisotropies, resulting in nearly vanishing CME-driven signals, while different sources of backgrounds for $\Delta\gamma$ remain [25, 56, 57]. Measurements performed at LHC energies by the CMS Collaboration show similar $\Delta\gamma$ signals for overlapping multiplicities in p +Pb and Pb+Pb collisions [25]. Similar studies are carried out by STAR, with results that show similar (or even larger) values of $\Delta\gamma$ scaled by elliptic anisotropy in p +Au and d +Au collisions as compared to Au+Au collisions [28]. Such

measurements appear to challenge the interpretation of magnetic-field-driven sources of charge separation. However, RP-independent background from three-point correlations $\langle \cos(\phi_1 + \phi_2 - 2\phi_3) \rangle$, where ϕ_1 , ϕ_2 and ϕ_3 are three distinct particles, can be significant in those small-system collisions and peripheral heavy-ion collisions; the same may not be true for more-central collisions [19, 20, 53, 55]. Extrapolation of small-system results as quantitative background baselines for different nucleus-nucleus systems, across the entire range of multiplicity, is not straightforward.

Over the past years, efforts have been dedicated towards developing data-driven methods and observables to isolate possible CME-driven signals from background contributions [22, 26, 27, 58–68], and to applying those methods to existing data. The event-shape engineering (ESE) analyses by the CMS and ALICE Collaborations at the LHC [26, 27] have reported a CME-induced charge separation that is consistent with zero with an upper limit (on the fraction of the $\Delta\gamma$ measurement that is due to CME) of the order of 7% and 26% at 95% confidence level (CL), respectively. Measurements of the pair invariant mass dependence of the $\Delta\gamma$ from STAR [69] have determined an upper limit of 15% at the 95% CL. A recent measurement by the STAR Collaboration using the spectator plane and participant plane analysis [70] has found a signal consistent with zero in peripheral collisions and a hint of finite positive signal in mid-central Au+Au collisions with a 1–3 σ significance. Possible remaining effects from non-flow correlations (two- and multi-particle correlations unrelated to a global symmetry plane) are under investigation [71]. An alternative charge-sensitive variable, $R_{\Psi_2}(\Delta S)$, has been proposed [62, 72, 73] to aid the characterization of CME-driven charge separation. The sensitivity of the $R_{\Psi_2}(\Delta S)$ variable has been studied in different contexts and has also been compared to that for the $\Delta\gamma$ observable [62, 73–81]. In a recent comprehensive investigation of different experimental observables for CME searches, it is found that the $R_{\Psi_2}(\Delta S)$ and $\Delta\gamma$ variables provide similar sensitivities to the CME signal and backgrounds for the two isobars [81].

In order to overcome the large backgrounds, isobar Ruthenium+Ruthenium ($^{96}\text{Ru}+^{96}\text{Ru}$) and Zirconium+Zirconium ($^{96}\text{Zr}+^{96}\text{Zr}$) collisions have been proposed [58]. It is expected that the magnetic field squared would be about 15% larger in Ru+Ru collisions due to its larger atomic number [4, 13], leading to a similar increase in the CME contribution in $\Delta\gamma$, while the same mass number of these two nuclei would lead to similar flow-driven backgrounds. With 1.2 billion minimum-bias (MB) events for each collision system, a 5 σ significance is expected in the CME signal difference between Ru+Ru and Zr+Zr [82]. This expectation is based on the same projection scheme as in Ref. [83], assuming that the CME-related signal fraction is 20% in $\Delta\gamma$.

Although similar, the backgrounds in Ru+Ru and Zr+Zr collisions are not expected to be identical. The difference in the nuclear deformation of the two isobars has been estimated to yield less than 1% difference in $\Delta\gamma$ background in peripheral to mid-central collisions. In more-central collisions from 0–20% centrality, the difference in $\Delta\gamma$ background can be larger than 2% [83, 84]. Further work from sophisticated nuclear structure calculations suggests that the resulting eccentricities (hence the flow-related backgrounds) may differ by 2–3% in mid-central collisions between the two isobars even without deformation [85, 86]. An approximate 4% difference in flow-driven background between these two systems is found in hydrodynamic simulations which include local charge conservation [54]. In order to account for a possible difference in v_2 , one of the variables we will focus on in this paper is the ratio $\Delta\gamma/v_2$, assuming that background proportionality to v_2 is identical between the isobar systems. Note that although v_2 can be precisely measured, the elliptic anisotropy contains non-flow contributions, and the background in $\Delta\gamma$ depends also on other physical processes besides the v_2 (see Eq. (4)). Therefore, it is crucial to minimize background contributions in order to search for the possibly small CME signal. Isobar collisions are considered to be an effective way to achieve that by studying the difference in the CME-sensitive observables between the two isobar systems.

Isobar collisions were acquired at RHIC in 2018 [87]. This paper reports results from a blind analysis [88] performed on the isobar data collected by the STAR Collaboration.

II. ISOBAR DATA AND BLIND ANALYSIS

A. Modality of isobar running at RHIC

The proposal for colliding isobar species is outlined in the 2017-18 RHIC beam use request by the STAR Collaboration [82]. The specific request was for two 3.5-week runs in the year 2018 with collisions of isobar nuclei, $^{96}\text{Ru}+^{96}\text{Ru}$ and $^{96}\text{Zr}+^{96}\text{Zr}$. This proposal is based on the prospect of achieving 5 σ significance in a scenario of a relative difference of the primary CME observable of 2–3% between the two isobar species [82]. It is estimated that with 3.5-week runs it is possible to collect more than 1.2 billion MB events for each species and achieve a statistical precision on the observable difference of about 0.5%. However, a special strategy is needed to minimize the systematic uncertainties. This required a specific plan in synergy with the RHIC Collider Accelerator Department to execute the isobar runs [87].

Studies from previous years using Au+Au and U+U collision data [89] indicate that there are several sources of systematics in the measurements of CME-sensitive observables. Two major sources are: 1) loss of detector acceptance, and 2) variation of luminosity during runs. These effect leads to run-to-run variation of the online trigger efficiency

and charged-particle track reconstruction efficiency in the Time Projection Chamber (TPC) [90]. These two sources can lead to irreducible systematic uncertainties in CME-sensitive observables. In order to keep the systematics due to these two major sources below the aforementioned statistical precision it is necessary to minimize the differences between the run conditions for the two species. Therefore the proposed procedure is to: 1) alternate the isobar species between each store of beam in RHIC, 2) keep long stores with constant beam luminosity, 3) match luminosities between the species, and 4) adjust the luminosity in such a way that the hadronic interaction rate at STAR is close to 10 kHz. With such a strategy, it is estimated that the systematic uncertainties in the ratio of observables could be reduced to about 0.2%. As we discuss later, these conditions were successfully provided by the RHIC facility [87] and this level of precision is indeed achieved in our measurements.

B. Detector apparatus and data quality cuts

STAR was the only operational detector for RHIC running in 2018. The main subsystems used for the analysis of isobar data are the TPC, the Time-of-flight detector (TOF) [91], the Event Plane Detector (EPD) [92], the Zero-Degree Calorimeters (ZDCs) [93], and the Vertex Position Detectors (VPDs) [94].

The TPC is used to detect charged particles within the pseudorapidity range $|\eta| < 1$, with full 2π azimuthal coverage and a transverse momentum lower limit of $p_T > 0.2$ GeV/ c [90]. The TPC is situated inside a magnet which maintained a constant solenoidal field of 0.5 T during the entire isobar runs. The tracking efficiency of the TPC ranges from 85% to 90% as determined using GEANT Monte Carlo (MC) simulations embedded into randomly sampled MB data events [95]. We are able to exploit the advantage of having data sets for two isobars collected under similar run conditions. For example, in the analyses we study the ratios of measurements between the two isobars. We do not apply efficiency corrections because the effects of inefficiency cancel out in these ratios.

For each collision we use the TPC to reconstruct the primary vertex position ($V_{z,\text{TPC}}$) along the beam direction (defined as the z axis) as well as its radial distance from the z axis (V_r). For all analyses, each event is required to have a vertex position within $-35 < V_{z,\text{TPC}} < 25$ cm and $V_r < 2$ cm using a coordinate system with the origin at the TPC center. The asymmetric vertex distribution is a consequence of our online vertex selection. The asymmetry arises due to the timing offset between VPD east and west that measures the z -position of the vertex that we discuss later. This leads to an asymmetric distribution peaked at -5 cm. In order to maximize the statistics we define an asymmetric range of the cut accordingly. To reduce the contamination from secondary charged particles, we require tracks reconstructed in the TPC to have a distance of closest approach (DCA) to the primary vertex of less than 3 cm. We also require each track to have at least 16 ionization points (N_{fits}) in the TPC. To study the effect of track splitting and merging on different v_n coefficients, we carefully study their relative pseudorapidity ($\Delta\eta$) dependence as splitting and merging will result in a peak or a dip, respectively, in this dependence [96–98]. We do not see evidence of track splitting effects; however, we observe a dip at low $\Delta\eta$ due to track merging that is dominant in central events. To minimize track merging, a requirement of $\Delta\eta > 0.05$ is applied. We also do this study for same-sign and opposite-sign pair correlations separately as the possible effects of track merging and splitting are expected to be different between the two cases. During the isobar run in 2018, one of the 24 sectors of the TPC was being used to commission the inner TPC (iTPC) sector and the data from this sector are not used for physics analysis. The loss of tracks due to this sector leads to an identifiable region of depletion in the η - ϕ acceptance map. However, the effect of this acceptance deficit in the final observables is corrected by reconstructing the harmonic flow vectors (Q -vectors) using re-weighting, re-centering, and shifting methods [99]. Such Q -vectors are then used for estimation of different observables and the event plane (EP) in this analysis. It is important to note that this effect is consistently present over the entire period of the run and is common to both the isobar species, and therefore cancels in the ratios of physics observables between Ru+Ru and Zr+Zr.

The MB data sample is collected with a trigger based on information from the VPDs [94]. The VPDs ($4.4 < |\eta| < 4.9$) also provide information on primary collision vertices along the beam direction ($V_{z,\text{VPD}}$). For the selection of good events we require the condition of $|V_{z,\text{TPC}} - V_{z,\text{VPD}}| < 5$ cm (unless otherwise noted). Variations in luminosity are kept to a minimum during the runs, with the dominant part of the MB data set having a variation of luminosity that corresponds to a coincidence of signals from the ZDCs in the range of 9.5–11.5 kHz. The variation of luminosity affects our centrality selection and a correction for this is made. We achieve a trigger efficiency close to 100% for events in which more than 50 tracks are reconstructed per unit pseudorapidity in the TPC (see Sec. III). For events with fewer tracks, the trigger efficiency decreases and a MC Glauber model is used to estimate and correct for such inefficiencies, as discussed in Sec. III.

Our event selection techniques suffer from out-of-time pile-up that requires an offline rejection. About 0.5% of events are identified as pile-up and removed by excluding outliers in the correlation between the number of TPC tracks and the number of those tracks matched with a hit in the TOF detector (the TOF is a fast detector and does not suffer from out-of-time pileup). We also require at least one TPC track matched to the TOF for selecting good

319 events. After all event selection cuts, we analyze approximately 1.9 billion MB events for Ru+Ru and 2.0 billion MB
 320 events for Zr+Zr collisions.

321 Our measurement uses the EPD detector for the first time in collider mode [92]. The EPD is used for measure-
 322 ments of the second- and third-harmonic EPs at forward rapidity. The EPD consists of two segmented scintillator
 323 wheels located at ± 3.75 m from the center of the TPC, along the beam direction, covering an acceptance window
 324 of approximately $2.1 < |\eta| < 5.1$ in pseudorapidity and 2π in azimuth. Each wheel consists of 12 “supersectors” (in
 325 azimuth) that are further divided (radially) into 31 tiles made of plastic scintillator. Each tile is connected to a silicon
 326 photomultiplier via optical fiber. Charged particles emitted in the forward and backward directions produce a signal
 327 distribution with identifiable peaks corresponding to various numbers of minimally ionizing particles in the EPD tiles.
 328 This information in each tile is used to reconstruct the EPs. Further details of the EPD can be found in Ref. [92].

329 The ZDCs and their associated Shower Maximum Detectors (SMDs) are used for determination of the spectator
 330 neutron plane [100, 101]. The ZDCs are Cherenkov-light sampling calorimeters located at forward and backward
 331 angles ($|\eta| > 6.3$) and are each composed of three identical modules. The SMDs are sandwiched between the ZDC
 332 modules and are composed of two planes with scintillator strips aligned with x or y directions perpendicular to the
 333 beam. The SMD information thus can be used to measure the centroid of the hadronic shower produced by the
 334 spectator neutrons in the ZDCs. The x and y positions of the shower centroid ($\langle X, Y \rangle_{\text{ZDCE, W-SMD}}$) calculated on an
 335 event-by-event basis provide spectator-plane reconstruction (see Refs. [102, 103] for details).

336 We do not use the data from the Beam-Beam Counters (BBC) and the Barrel Electromagnetic Calorimeter (BEMC)
 337 in this analysis other than for data quality assurance purposes. The time-dependence of the Q -vectors from the BBCs
 338 are studied to identify bad runs. The number of TPC tracks matched to the BEMC ($N_{\text{trk}}^{\text{BEMC-matched}}$) is also examined
 339 as a function of time to identify outlier runs.


340 C. Blinding of data sets and preparation for analysis

341 The recommendation to perform a blind analysis of the isobar data was initially made by the Nuclear and Particle
 342 Physics Program Advisory Committee at Brookhaven National Laboratory [104]. The procedure to blind the isobar
 343 data is determined and implemented well before the actual data taking. The raw data are made inaccessible to the
 344 analysts to eliminate possible unconscious biases.

345 A total of five institutional groups within the collaboration perform blind analyses of the isobar data. The analysts
 346 from each group focus on a specific analysis method described in Sec. IV. Substantial overlap of some analyses helps to
 347 cross check the results. The details of the blinding procedure and data structure are decided by an Analysis Blinding
 348 Committee (ABC), consisting of STAR members who are not part of the team of analysts. The ABC works in close
 349 collaboration with the data production team to provide the analysts with access only to data in which species-specific
 350 information is disguised or removed, until the final un-blinded analysis step. Before the final step ABC also makes
 351 sure that the information provided to the analysts to perform quality assurance (QA) of the data do not reveal the
 352 species identity.

353 D. Methods for isobar blind analysis

354 The detailed procedure for the blind analysis of isobar data is outlined in Ref. [88] and is strictly followed by the
 355 analysts. As shown in Fig. 1, the blind analysis procedure includes a mock-data challenge to perform a closure test
 356 and three main steps: 1) isobar-mixed analysis, 2) isobar-blind analysis, and 3) isobar-unblind analysis [105].



357 FIG. 1. Flowchart to illustrate the steps of the isobar blind analysis [105]. This is based on the procedure for the isobar blind
 358 analysis outlined in Ref. [88].

359 In the zeroth step preceding the blind analysis, the analysts participated in a mock-data challenge. The purpose
 360 of this step is to familiarize the analysts with the data structures that have been designed for the blind analysis, and

with the techniques to access the data. Feedback is also provided to the ABC to ensure feasibility of the analysis blinding process. Data for Au+Au collisions at $\sqrt{s_{\text{NN}}} = 27$ GeV (collected in 2018 after the isobar run) are used for this step.

The first step of this analysis is referred to as the “isobar-mixed analysis”. In this step the majority of the analysis work is done. Analysts are provided with a data sample where each “run” contains events that are a mixed sample of the two species. The analysis teams then perform QA and a complete analysis of the data. The details of the QA procedure are discussed in the next section. The analysis teams test their analysis code and document their analysis procedures. They are then frozen for the next two steps of the analysis, except for situations as strictly defined at the end of this subsection. An important part of data QA is to reject bad runs and pile-up events. This requires retention of the time ordering of the data. In order to avoid unconscious biases, an automated algorithm for bad run rejection is developed and the corresponding codes are also frozen. The QA algorithm is tested using existing Au+Au and U+U data. In this step the documentation related to the criteria for signatures of the CME in each observable, which we discuss in Sec. IV, is also frozen. From the next steps onwards the analysts can only execute frozen codes. As we discuss later, different groups focus on analysis of specific CME-sensitive observables. In order to check the consistency of the numerical output of the analysis codes from five groups, an exercise is performed in this step. The analysts from different groups are required to estimate a few common observables in the same approach, with exactly the same data, using their own individual codes. The results from different groups are ensured to be numerically identical to each other.

The second step is referred to as the “isobar-blind analysis”. For this the analysts are provided with files, each of which contain data from a single, but blinded, isobar species to perform run-by-run QA. Every file provided to the analyst contains a limited number of events that is determined to be insufficient to allow an identification of the species or the observation of a statistically significant CME signal. A pseudo run-number is used to hide the identity of the species for each file. The mapping between these pseudo run-numbers and the original ones is not revealed to the analysts. The automated algorithms are then used to identify the runs with stable detector performance and to reject bad runs.

The final step is referred to as “isobar-unblind” analysis. In this step, all elements of the data, including species information, are revealed to the analysts and the physics results are produced by the analysts using the previously frozen codes. As mentioned before, analysts from five independent groups participate in the blind analysis. In order to further avoid unconscious biases, analysts from a given group are not allowed to execute their own codes to produce the final results. Instead, a STAR collaborator is identified either from a different blind analysis group or among members not participating in the blind analysis, to run that group’s frozen code. The findings from this step are directly presented in this paper without alteration. A brief discussion of post-blinding analysis results is given near the end of the paper in Sec. VI.

E. Quality assurance of the blind data

Unlike conventional QA, the analysis teams do not have access to the full statistics of the recorded data. In accordance with the blind analysis policy, any form of manual selection or rejection of a part of the data sample is not permitted. This makes the QA of the data analysis challenging. In order to avoid unconscious biases and yet perform an effective clean up of data we develop an automated algorithm with predefined criteria for QA. These algorithms perform three major tasks: 1) identify the regions of the data sample or runs with stable detector performance by studying the time dependence of various quantities, 2) identify regions of the data sample with problematic detector performance or outlier runs, and 3) remove pile-up events.

We study run-by-run variation in the mean value of quantities such as the average multiplicity ($\langle N_{\text{trk}}^{\text{offline}} \rangle$) of tracks from the TPC, basic track level quantities like the distance of closest approach ($\langle \text{DCA} \rangle$), and quantities related to azimuthal acceptance such as mean cosine of the azimuthal angle ($\langle \cos(\phi) \rangle$). The QA procedure is performed over the entire data sample, and separately for the five analysis groups because each group provides a list of such quantities specific to the analysis. For example, Group-3 and Group-4 use the ZDC for EP analysis and therefore need to carefully study the QA variables for quantities related to the ZDC. The analyses of other groups that do not use the ZDC do not need to perform QA related to that detector. Table I lists the common QA variables and criteria, as well as the analysis-specific ones, to reject bad runs.

Data collection for the isobar run took eight weeks and two days. During this time, the acceptance of the detector changed due to the temporary failure of electronics modules or other causes. Thus, periods of stable and uniform operation were identified and each stable period was treated separately for acceptance and track weighting corrections. To identify jumps or boundaries between stable regions we study QA quantities with time or run numbers. We study the first and second order derivatives of quantities with respect to time. The zeros of the first order derivative surrounded by two zeros of the second order derivative defines a run mini-region. From each mini-region we extract

Variable \ Group	Group-1	Group-2	Group-3	Group-4	Group-5
$\langle N_{\text{trk}}^{\text{offline}} \rangle$	■	■	■	■	■
$\langle N_{\text{hits}}^{\text{TOF}} \rangle$	■	□	□	■	□
$\langle N_{\text{trk}}^{\text{TOF-matched}} \rangle$	□	■	■	■	■
$\langle p_T \rangle$	■	□	■	■	□
$\langle \eta \rangle$	■	□	■	■	□
$\langle \text{DCA} \rangle$	■	□	■	■	□
$\langle V_z \rangle$	□	□	■	■	□
$\langle \phi \rangle$	□	□	□	■	□
$\langle N_{\text{fits}} \rangle$	□	□	■	■	□
$\langle Q_{1x} \rangle_{\text{TPC}}$	□	■	■	□	■
$\langle Q_{1y} \rangle_{\text{TPC}}$	□	■	■	□	■
$\langle Q_{2x} \rangle_{\text{TPC}}$	■	■	■	□	■
$\langle Q_{2y} \rangle_{\text{TPC}}$	■	■	■	□	■
$\langle Q_{1x} \rangle_{\text{EPD}}$	□	■	□	□	■
$\langle Q_{1y} \rangle_{\text{EPD}}$	□	■	□	□	■
$\langle Q_{2x} \rangle_{\text{EPD}}$	□	■	□	□	■
$\langle Q_{2y} \rangle_{\text{EPD}}$	□	■	□	□	■
$\langle Q_x \rangle_{\text{BBCE}}$	□	□	■	□	□
$\langle Q_y \rangle_{\text{BBCE}}$	□	□	■	□	□
$\langle Q_x \rangle_{\text{BBCW}}$	□	□	■	□	□
$\langle Q_y \rangle_{\text{BBCW}}$	□	□	■	□	□
$\langle X \rangle_{\text{ZDCE-SMD}}$	□	□	■	□	□
$\langle Y \rangle_{\text{ZDCE-SMD}}$	□	□	■	□	□
$\langle X \rangle_{\text{ZDCW-SMD}}$	□	□	■	□	□
$\langle Y \rangle_{\text{ZDCW-SMD}}$	□	□	■	□	□
$\langle Q_{1x} \rangle_{\text{ZDC}}$	□	□	□	■	□
$\langle Q_{1y} \rangle_{\text{ZDC}}$	□	□	□	■	□
$\langle Q_{2x} \rangle_{\text{ZDC}}$	□	□	□	■	□
$\langle Q_{2y} \rangle_{\text{ZDC}}$	□	□	□	■	□
$\langle N_{\text{trk}}^{\text{BEMC-matched}} \rangle$	□	□	■	□	□

TABLE I. Common and analysis-specific QA variables and criteria used to reject bad runs. ■ Used, □ Unused. See the texts in Sec. II B for the definition of different variables. The Q_{nx} and Q_{ny} refer to the components of the flow Q -vectors that we discuss in the later sections.

416 the local mean and the weighted error. We define regions of stable detector conditions by merging these mini-regions
417 if the mean values of the quantities in adjacent mini-regions are: 1) within five times the weighted error or 2) within
418 one percent of the variation of the local mean. A run is marked as an outlier or bad run in each stable region if the
419 value of the QA quantity is five standard deviations from the local mean. Once the first-round of stable regions are
420 identified and bad runs are removed, the whole process is repeated. Iterations are performed until no additional bad
421 run is identified by the algorithm. The stability of this automated algorithm is tested with existing Au+Au and U+U
422 data sets before the code freeze in step-1 (isobar mixed analysis).

423 In the second step of isobar analysis the blind data set is provided to the analysts that includes all the runs for
424 both species (species identity is blinded) but each run contains only approximately 1% of the entire statistics of that
425 run. Following the methods of the blind analysis, all the files are named by a pseudo-run-number mapped to the
426 original run-number by the production team to ensure the species are blind to the analysts. The analysts prepare the
427 necessary histograms of QA variables with pseudo-run-numbers using the blind data set. A non-analyst then helps to
428 re-map the run-numbers, executes the frozen run-by-run QA algorithm and prepares the final lists of bad runs and
429 stable periods for each group. These numbers are different for different analysis groups because of the difference in
430 the analysis-specific QA variables (see Table I). It is important to note that the QA is performed on the combined
431 data set of two species and not on individual species. By the end of the QA, the automated algorithm identified
432 less than 4% of the data to be discarded from the analysis based on predefined criteria. Since the criteria of pattern
433 recognition to discard the problematic part of the data sample is predefined and frozen prior to the blind analysis,
434 unconscious biases are eliminated.

435 Another automated algorithm is implemented prior to the blind analysis to remove pile-up events. Based on studies
436 of previous data sets it is observed that pile-up events lead to satellites in the correlation between the number of

437 tracks from the TPC ($N_{\text{trk}}^{\text{offline}}$) and the number of TPC tracks matched with TOF ($N_{\text{trk}}^{\text{TOF}}$). For a given window of
 438 $N_{\text{trk}}^{\text{TOF}}$, the distribution of $N_{\text{trk}}^{\text{offline}}$ appears to be described by a double negative binomial distribution with two sets
 439 of widths and means. The wider distribution corresponds to the pile-up events. For each value of $N_{\text{trk}}^{\text{TOF}}$ one can
 440 reduce the pile-up contribution by applying upper and lower cuts of $3(\sigma + \text{Skewness})$ and 4σ respectively, around the
 441 mean value of the narrow distribution. Such a procedure is implemented in the frozen algorithm and used for pile-up
 442 removal in our analysis.

443 In the final step of the analysis when the isobar data are unblinded we check the distributions of energy deposition in
 444 the ZDCs. We find that the Zr+Zr collisions have a significantly larger energy deposition than the Ru+Ru collisions,
 445 consistent with the larger neutron number in the former. We also check the net-charge distributions from the TPC
 446 and find that the Ru+Ru collisions have a larger mean than the Zr+Zr collisions. These checks confirm that the two
 447 species are correctly separated in the unblind sample of the data provided to the analysts.

448 F. Methodology of uncertainty estimation

449 Systematic uncertainties are assessed by varying each of the analysis cuts within a range that is considered as
 450 the reasonable maximum range. This way one estimates the quantity $|\Delta|$ which is the absolute difference between
 451 the magnitudes of an observable with the default cut \mathcal{O}_{def} and with a particular cut variation \mathcal{O}_i . The statistical
 452 fluctuation on this difference is given by $\sigma_{\Delta} = \sqrt{|\sigma_{\text{def}}^2 - \sigma_i^2|}$, where σ_{def} and σ_i are the statistical uncertainties of the
 453 two measurements [106]. If σ_{Δ} is larger than $|\Delta|$, i.e. the change in the result is consistent with statistical fluctuations,
 454 then no systematic uncertainty is considered for this cut variation. Otherwise, the systematic uncertainty is assigned
 455 to be $\sigma_{\text{sys}} = \sqrt{\Delta^2 - \sigma_{\Delta}^2} / \sqrt{12}$ assuming a flat distribution for \mathcal{O}_i . Different sources of systematics are added in
 456 quadrature. For compound observables, such as the $\Delta\gamma/v_2$, systematic uncertainties are assessed as above, treating
 457 the compound observable as a single quantity. This way the (anti-)correlations in the systematic uncertainties in the
 458 component variables are automatically taken into account.

459 All analyses reported in this paper have a common set of cuts and variations for the purpose of systematic uncertainty
 460 determination. As noted above, the events used in all analyses are required to have a primary vertex within $-35 <$
 461 $V_{z,\text{TPC}} < 25$ cm. To estimate the systematic uncertainty due to the acceptance dependence on $V_{z,\text{TPC}}$, results using
 462 only events within $-35 < V_{z,\text{TPC}} < 0$ cm are compared with those from the full $V_{z,\text{TPC}}$ range. A maximum DCA of
 463 3 cm and a minimum N_{fits} of 16 are required for the TPC tracks to be used in the analysis. Systematic uncertainties
 464 are assessed by varying the maximum DCA from 3 cm to 2 cm and the minimum N_{fits} from 16 to 21. In addition to
 465 the common cuts, each analysis has specific cuts described in the corresponding results subsections in Sec. V. At the
 466 end, the systematic uncertainties of all sources are added in quadrature, the value of which is quoted as one standard
 467 deviation.

468 For statistical uncertainty estimations we use the standard error propagation method. We use both analytical
 469 and MC (Bootstrap [107]) approaches to examine the influence of co-variance terms. Such cases may be relevant
 470 for primary CME-sensitive quantities like the ratio of $\Delta\gamma/v_2$. We find that the statistical uncertainties in the ratio
 471 observable ($\Delta\gamma/v_2$) are completely dominated by uncertainties of the numerator (by more than a factor of 50).
 472 Furthermore, the covariance between the numerator and denominator is also negligible, simplifying the statistical
 473 error calculations.

474 III. CENTRALITY DETERMINATION

475 The centrality determination is made at the beginning of the final step of the isobar analysis, using the unblinded
 476 data. This is performed by a team of collaborators who do not take part in the blind analysis of the data, and before
 477 any of the observables are measured.

478 Centrality is defined based on the charged track multiplicity ($N_{\text{trk}}^{\text{offline}}$) from the TPC within the pseudorapidity
 479 acceptance $|\eta| < 0.5$. Each track is required to have a DCA to the primary vertex of less than 3 cm and must be
 480 formed from at least 10 ionization points in the TPC gas volume. The $N_{\text{trk}}^{\text{offline}}$ depends on the tracking efficiency of the
 481 TPC, which in turn depends on the occupancy of the TPC and hence on the collider luminosity, which is monitored
 482 with the ZDC coincidence rate. The $\langle N_{\text{trk}}^{\text{offline}} \rangle$ is found to have a linear dependence on the ZDC coincidence rate.
 483 The parameterization of this dependence is used to correct $N_{\text{trk}}^{\text{offline}}$ for luminosity effect. To this end, $N_{\text{trk}}^{\text{offline}}$ is first
 484 converted to a real number by sampling the range from a half unit below to a half unit above, and the correction is
 485 then applied to the real number. Over the ZDC coincidence rate range of 9.5 kHz to 11.5 kHz, which describes the
 486 dominant part of this data set, the luminosity correction to the multiplicity is less than 0.02% for Ru+Ru collisions
 487 and less than 0.29% for Zr+Zr collisions. This luminosity correction is small owing to the very stable beam conditions
 488 provided by RHIC during the isobar run.

489 The quantity $N_{\text{trk}}^{\text{offline}}$ is further corrected for the acceptance variation as a function of $V_{z,\text{TPC}}$. To obtain the correction
 490 factor, the $N_{\text{trk}}^{\text{offline}}$ distributions, $P(N_{\text{trk}}^{\text{offline}})$, are plotted in 2 cm bins of $V_{z,\text{TPC}}$ in the range $-35 < V_{z,\text{TPC}} < 25$ cm.
 491 These multiplicity distributions in heavy-ion collisions have a characteristic sharp decline at large multiplicity values.
 492 The location of the half-maximum of this decline is measured by fitting this region with an error function. The
 493 correction factor is determined by making the location of the half-maximum point of the given $V_{z,\text{TPC}}$ bin equal to the
 494 one at $-1 < V_{z,\text{TPC}} < 1$ cm (the center of the TPC).




FIG. 2. Distributions of the number of charged particles ($N_{\text{trk}}^{\text{offline}}$) from the TPC in the pseudorapidity acceptance $|\eta| < 0.5$ in Ru+Ru (upper left panel) and Zr+Zr (lower left panel) collisions. The experimental distributions have been corrected for variations in the luminosity and the vertex position $V_{z,\text{TPC}}$, and uncorrected for tracking efficiency. Fits to the experimental distributions (gray circles) are performed by the two-component Glauber model using two sets of Woods-Saxon parameters in Table II (blue crosses for Case-2 and red histograms for Case-3). The Ru+Ru to Zr+Zr ratio of the experimental data, as well as those of the Glauber model fits for Case-2 and Case-3 are shown in the upper right and lower right panels, respectively. The Glauber simulation with the Case-3 nuclear density parameters is used for centrality determination as it provides the best description of the experimental data.

495 Figure 2 shows the luminosity and $V_{z,\text{TPC}}$ corrected distributions $P(N_{\text{trk}}^{\text{offline}})$ in Ru+Ru and Zr+Zr collisions. The
 496 centrality classes in this analysis are defined by fitting the $P(N_{\text{trk}}^{\text{offline}})$ distributions to those obtained from MC Glauber
 497 simulations [108, 109]. In Glauber simulations, the probability of a collision at a given impact parameter (b) and
 498 the corresponding number of participant nucleons (N_{part}) and number of binary nucleon-nucleon collisions (N_{coll})
 499 are obtained by MC sampling. The inputs for this calculation are the nuclear thickness function and the inelastic
 500 nucleon-nucleon cross section ($\sigma_{\text{NN}}^{\text{inel}}$) which is taken to be 42 mb for the current case of $\sqrt{s_{\text{NN}}} = 200$ GeV collisions [110].

501 The nuclear thickness function is the projection of the 3D nuclear density onto the transverse plane (perpendicular
 502 to the z axis). It is obtained by sampling nucleons in the incoming nuclei according to the Woods-Saxon (WS)
 503 distribution defined in the nucleus rest frame with a spherical coordinate system (r is radial position and θ is polar
 504 angle) [111]:

$$\rho(r, \theta) = \frac{\rho_0}{1 + \exp\left[\frac{r - R(1 + \beta_2 Y_2^0(\theta))}{a}\right]}, \quad (5)$$

505 where R is the radius parameter, a is the diffuseness parameter of the nuclear surface, β_2 is the quadrupole deformity
 506 parameter, $Y_2^0(\theta) = \frac{1}{4}\sqrt{\frac{5}{\pi}}(3\cos^2\theta - 1)$, and ρ_0 is the normalization factor. Nuclear density distributions of $^{96}_{44}\text{Ru}$ and

507 $^{96}_{40}\text{Zr}$ are not accurately known [83, 86, 112]. In this work, three sets of WS parameters [83, 113] are investigated. These
 508 sets of parameters are listed in Table II. The first two sets (Case-1 and Case-2) have the same R and a parameters and
 509 different deformations. The parameters are constrained by $e+A$ scattering experiments [114, 115] and calculations
 510 based on a finite-range droplet macroscopic model and the folded-Yukawa single-particle microscopic model [116]. The
 511 charge radius of $^{96}_{44}\text{Ru}$, because of its additional protons, is larger than that of $^{96}_{40}\text{Zr}$. The neutron and proton density
 512 parameters are taken to be the same for both R and a , so Ru is larger than Zr. The third set (Case-3) is from recent
 513 calculations based on energy density functional theory (DFT), assuming the nuclei are spherical [85, 113]. The proton
 514 and neutron distributions are both calculated, and the overall size of Ru is found to be smaller than Zr because of a
 515 significantly thicker neutron skin in the latter. The nucleon distributions are found to be well parameterized by the
 516 halo-type WS distributions (i.e. the neutron a parameter is significantly larger than that for the proton) [113].

TABLE II. The Woods-Saxon parameters used in the Glauber simulations for the centrality determination.

Nucleus	Case-1 [83]			Case-2 [83]			Case-3 [113]		
	R (fm)	a (fm)	β_2	R (fm)	a (fm)	β_2	R (fm)	a (fm)	β_2
$^{96}_{44}\text{Ru}$	5.085	0.46	0.158	5.085	0.46	0.053	5.067	0.500	0
$^{96}_{40}\text{Zr}$	5.02	0.46	0.08	5.02	0.46	0.217	4.965	0.556	0

517 In this analysis we use the simple two-component model for multiparticle production [117]. Several alterna-
 518 tive approaches of multiparticle production have been developed over the years, such as Quark-Glauber [118], IP-
 519 Glasma [119], TRENTO [120] and Shadowed Glauber [121], that improve the two-component model. These approaches
 520 can be investigated in future STAR analyses – for the current work we stick to the two-component nucleon based MC
 521 Glauber model for simplicity. The multiplicity density at a given b , with the corresponding N_{part} and N_{coll} from the
 522 Glauber calculation for each set of the WS parameters, is parameterized by the two-component model [117] as:

$$N_{\text{trk}}^{\text{Glauber}} = n_{pp} [(1-x)N_{\text{part}}/2 + xN_{\text{coll}}], \quad (6)$$

523 where n_{pp} is the average pseudorapidity multiplicity density in zero-bias nucleon-nucleon (NN) collisions, and x is the
 524 relative contribution to multiplicity from hard processes. The multiplicity given by Eq. (6) is the average multiplicity.
 525 Multiplicity fluctuations are taken into account in the following way. $N_{\text{trk}}^{\text{Glauber}}$ is considered to be accumulated by
 526 $(1-x)N_{\text{part}}/2 + xN_{\text{coll}}$ (that is rounded to the closest integer) NN collisions. In each NN collision, the multiplicity n
 527 is obtained by convolution of the negative binomial distribution (NBD)

$$P_{\text{NBD}}(n_{pp}, k; n) = \frac{\Gamma(n+k)}{\Gamma(n+1)\Gamma(k)} \cdot \frac{(n_{pp}/k)^n}{(1+n_{pp}/k)^{n+k}}, \quad (7)$$

528 where Γ is the gamma function and the fluctuation parameter k controls the sharpness of the large multiplicity tail
 529 of the $N_{\text{trk}}^{\text{Glauber}}$ distribution.

530 The Glauber multiplicity distribution obtained in this way is then convolved with a binomial distribution to account
 531 for the tracking inefficiency and acceptance of the TPC. The net effect depends on the TPC hit occupancy and is
 532 modeled as a linear function in the multiplicity [108]. The final $N_{\text{trk}}^{\text{Glauber}}$ distribution is then fitted to the experimental
 533 $N_{\text{trk}}^{\text{offline}}$ distribution, with n_{pp} , k , and x as fit parameters. The fit is performed simultaneously for Ru+Ru and Zr+Zr
 534 datasets with the fit parameters forced to be common for both isobars. Since the peripheral collisions are affected
 535 by trigger inefficiency, the fit range is restricted to $N_{\text{trk}}^{\text{offline}} > 50$. Since the number of simulated events is not equal
 536 to that of data, the Glauber multiplicity distribution is normalized to the data of the corresponding isobar system
 537 within the range from $N_{\text{trk}}^{\text{offline}} = 50$ to 500 before the χ^2 is calculated that is to be minimized in the fit.

538 A simultaneous fit of the $N_{\text{trk}}^{\text{offline}}$ distributions for the two isobars is performed for each set of the WS parameters
 539 for $^{96}_{44}\text{Ru}$ and $^{96}_{40}\text{Zr}$ listed in Table II. The first set of parameters (Case-1) is rejected from further analysis because it
 540 yields the largest χ^2/ndf among the three scenarios. The fit results for Case-2 and Case-3 are shown in Fig. 2 (left
 541 panels), with similar χ^2/ndf values. The $P(N_{\text{trk}}^{\text{offline}})$ distributions shown in Fig. 2 for data are normalized by the
 542 number of events which is approximately 1.9 B for Ru+Ru and 2.0 B for Zr+Zr.

543 In order to further inform the choice of the WS parameters, the ratio of the experimentally measured $N_{\text{trk}}^{\text{offline}}$ distri-
 544 bution for Ru+Ru to the one for Zr+Zr is compared with the same ratio obtained for the MC Glauber calculations.
 545 These ratios are shown in Fig. 2 (right panels) where we have re-binned the histograms to improve the visibility for
 546 data-model comparison. To obtain this figure we take the ratio between the distributions for Ru+Ru and Zr+Zr
 547 shown in Fig.2 (left panels). The same procedure is followed for both data and Glauber. The multiplicity ratio
 548 obtained for Case-3 is in a better agreement with the experimental distribution at $N_{\text{trk}}^{\text{offline}} > 50$, while the ratio for
 549 Case-2 deviates from the experimental ratio, particularly in central collisions. Note that the Case-3 fit ratio does not
 550 fully describe the data on the large multiplicity tail and there is room for future improvement. The larger multiplicity

in central Ru+Ru than in central Zr+Zr collisions is due to the smaller $\sqrt{\langle r^2 \rangle}$, the root-mean-square (RMS) size (and thus a higher energy density) of the $^{96}_{44}\text{Ru}$ nucleus compared to the $^{96}_{40}\text{Zr}$ nucleus, as predicted by DFT [85, 86, 122]. If the radius parameter R is set to be smaller for Ru in the WS density parameterization of Case-2 (and Case-1), then the high multiplicity tails observed in data would also be described [86]. However, it would still fail to describe the subtle shape in the intermediate multiplicity range observed in data [86, 113]. It must be also noted that the non-zero β_2 parameter for $^{96}_{40}\text{Zr}$ as used by Case-2 is not compatible with transition measurements and calculations [123, 124]. Based on the above considerations, the Case-3 WS density parameterization is chosen for our centrality calculations. The fit corresponds to values of MC Glauber parameters $n_{pp} = 2.386$, $k = 3.889$, and $x = 0.123$.

TABLE III. Centrality definition by $N_{\text{trk}}^{\text{offline}}$ ranges (efficiency-uncorrected multiplicity in the TPC within $|\eta| < 0.5$) in Ru+Ru and Zr+Zr collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV. The first column is the centrality range labels we use throughout the paper. The two centrality columns are the actual centrality ranges which are slightly different because of integer edge cuts used for the centrality determination. The mean $\langle N_{\text{trk}}^{\text{offline}} \rangle$ values, the mean number of participants ($\langle N_{\text{part}} \rangle$), and the mean number of binary collisions ($\langle N_{\text{coll}} \rangle$) are also listed. The statistical uncertainties on $\langle N_{\text{trk}}^{\text{offline}} \rangle$ are all significantly smaller than 0.01. The uncertainties on $\langle N_{\text{part}} \rangle$ and $\langle N_{\text{coll}} \rangle$ are systematic.

Centrality label (%)	Ru+Ru					Zr+Zr				
	Centrality(%)	$N_{\text{trk}}^{\text{offline}}$	$\langle N_{\text{trk}}^{\text{offline}} \rangle$	$\langle N_{\text{part}} \rangle$	$\langle N_{\text{coll}} \rangle$	Centrality(%)	$N_{\text{trk}}^{\text{offline}}$	$\langle N_{\text{trk}}^{\text{offline}} \rangle$	$\langle N_{\text{part}} \rangle$	$\langle N_{\text{coll}} \rangle$
0–5	0–5.01	258.–500.	289.32	166.8±0.1	389±10	0–5.00	256.–500.	287.36	165.9±0.1	386±10
5–10	5.01–9.94	216.–258.	236.30	147.5±1.0	323±5	5.00–9.99	213.–256.	233.79	146.5±1.0	317±5
10–20	9.94–19.96	151.–216.	181.76	116.5±0.8	232±3	9.99–20.08	147.–213.	178.19	115.0±0.8	225±3
20–30	19.96–30.08	103.–151.	125.84	83.3±0.5	146±2	20.08–29.95	100.–147.	122.35	81.8±0.4	139±2
30–40	30.08–39.89	69.–103.	85.22	58.8±0.3	89.4±0.9	29.95–40.16	65.–100.	81.62	56.7±0.3	83.3±0.8
40–50	39.89–49.86	44.–69.	55.91	40.0±0.1	53.0±0.5	40.16–50.07	41.–65.	52.41	38.0±0.1	48.0±0.4
50–60	49.86–60.29	26.–44.	34.58	25.8±0.1	29.4±0.2	50.07–59.72	25.–41.	32.66	24.6±0.1	26.9±0.2
60–70	60.29–70.04	15.–26.	20.34	15.83±0.03	15.6±0.1	59.72–70.00	14.–25.	19.34	15.10±0.03	14.3±0.1
70–80	70.04–79.93	8.–15.	11.47	9.34±0.02	8.03±0.04	70.00–80.88	7.–14.	10.48	8.58±0.02	7.12±0.04
20–50	19.96–49.86	44.–151.	89.50	60.9±0.3	96.7±1.0	20.08–50.07	41.–147.	85.68	58.9±0.3	90.3±0.9

The centrality of an event is defined by the percentile of the total cross section. The integer edge cuts are made so that the integrals of the $N_{\text{trk}}^{\text{offline}}$ distributions would be closest to the 5% or 10% mark. For the 0–20% centrality interval the experimental data are used for integration, while the MC Glauber distributions are used for the remaining range. The reason for this choice is because it is certain that the online trigger is fully efficient for collisions more central than 20%.

Table III lists the centrality definition and the corresponding $\langle N_{\text{trk}}^{\text{offline}} \rangle$, $\langle N_{\text{part}} \rangle$ and $\langle N_{\text{coll}} \rangle$ for Ru+Ru and Zr+Zr collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV obtained in this work. Throughout this paper, we label the centralities as in the first column of Table III. Because of the integer edge cuts in the centrality determination, the actual centrality ranges are slightly different, which are also listed in Table III for Ru+Ru and Zr+Zr collisions, respectively. We estimate systematic uncertainties on $\langle N_{\text{part}} \rangle$ and $\langle N_{\text{coll}} \rangle$ by varying the input parameters (R , a) in the MC Glauber simulation and by varying n_{pp} and x in the two-component model. Figure 3 (upper panel) shows the $\langle N_{\text{trk}}^{\text{offline}} \rangle$ as a function of centrality in the two isobar collision systems. The Ru+Ru/Zr+Zr ratio of the mean multiplicities is shown in the lower panel of Fig. 3. The mean multiplicity is larger in Ru+Ru collisions than in Zr+Zr collisions of matching centrality. Note that the shape of this ratio as a function of centrality can be affected by the inexact matching of centralities by integer edge cuts on $N_{\text{trk}}^{\text{offline}}$. The shape may also be influenced by other factors that require further studies.

IV. OBSERVABLES FOR ISOBAR BLIND ANALYSIS

The isobar blind analysis specifically focuses on the following approaches and corresponding observables. The general strategy is to compare results from the two isobar species to search for a statistically significant difference in the observables used. The following subsections describe these approaches and corresponding observables which include: 1) measurements of the second- and higher-order harmonics of the γ correlator, 2) differential measurements of $\Delta\gamma$ (with respect to pseudorapidity gap $\Delta\eta$ and invariant mass m_{inv}) to identify and quantify backgrounds, 3) exploiting the relative charge separation across spectator and participant planes, and 4) the use of the R observable to measure charge separation. The first three approaches are based on the aforementioned three-point correlator and the last employs a different approach. For each observable/approach, we predefine a set of the CME signatures prior to the blind analysis, for which a magnitude of high significance must be observed for an affirmative observation of the CME.




FIG. 3. (Upper) The efficiency-uncorrected mean multiplicity $\langle N_{\text{trk}}^{\text{offline}} \rangle$ from the TPC within $|\eta| < 0.5$ as a function of centrality in Ru+Ru and Zr+Zr collisions. The centrality bins are shifted horizontally for clarity. (Lower) The ratio of the mean multiplicity in Ru+Ru collisions to that in Zr+Zr collisions in matching centrality. The points include statistical uncertainties that are within the marker size.

A. $\Delta\gamma$ and mixed harmonics with second and third order event planes

We rewrite the conventional γ correlator (Eq. (2)) with a more specific notation,

$$\gamma_{112} = \langle \cos(\phi_\alpha + \phi_\beta - 2\Psi_2) \rangle, \quad (8)$$

where ϕ_α and ϕ_β are the azimuthal angles of particles of interest (POIs) and Ψ_2 is the second-order flow plane. Here, the subscripts “1”, “1” and “2” in γ_{112} refer to the harmonics associated with the ϕ_α , ϕ_β and Ψ_2 , respectively. In practice, the flow plane is approximated with the EP (Ψ_{EP}) reconstructed with measured particles, and then the measurement is corrected for the finite EP resolution [125]. The charge-dependent backgrounds in $\Delta\gamma_{112} = \gamma_{112}^{\text{OS}} - \gamma_{112}^{\text{SS}}$ can be broadly understood using the example of resonance decays. If resonances from the event exhibit elliptic flow, their decay daughters could mimic a signal for charge separation across the flow plane with a magnitude proportional to v_2 [36, 49, 51]. Therefore, following Eq. (4), one should study the normalized quantity

$$\frac{\Delta\gamma_{112}}{v_2}, \quad (9)$$

to account for the trivial scaling expected from a purely background scenario. The flowing-resonance picture can be generalized to a larger portion of the event, or even the full event, through the mechanisms of transverse momentum conservation (TMC) [38, 126] and/or local charge conservation (LCC) [51]. In the case of the γ correlator this contribution can be written as

$$\begin{aligned} \gamma_{112} &= \langle \cos(\phi_\alpha + \phi_\beta - 2\Psi_2) \rangle \\ &= \langle \cos(\phi_\alpha - \Psi_2) \cos(\phi_\beta - \Psi_2) \rangle - \langle \sin(\phi_\alpha - \Psi_2) \sin(\phi_\beta - \Psi_2) \rangle \\ &= (\langle v_{1,\alpha} v_{1,\beta} \rangle + B_{\text{IN}}) - (\langle a_{1,\alpha} a_{1,\beta} \rangle + B_{\text{OUT}}). \end{aligned} \quad (10)$$

The CME should dominantly contribute to the $\langle a_{1,\alpha} a_{1,\beta} \rangle$ term. The in-plane $\langle v_{1,\alpha} v_{1,\beta} \rangle$ component represents the charge separation unrelated to the magnetic field direction, and $(B_{\text{IN}} - B_{\text{OUT}})$ denotes the flow-related background.

601 Ideally, the two-particle correlator,

$$\begin{aligned}\delta &= \langle \cos(\phi_\alpha - \phi_\beta) \rangle \\ &= (\langle v_{1,\alpha} v_{1,\beta} \rangle + B_{\text{IN}}) + (\langle a_{1,\alpha} a_{1,\beta} \rangle + B_{\text{OUT}}),\end{aligned}\quad (11)$$

602 should also manifest $\langle a_{1,\alpha} a_{1,\beta} \rangle$, but in reality it could be dominated by short-range two-particle correlation back-
603 grounds (i.e. $B_{\text{IN}} + B_{\text{OUT}}$). Similar to $\Delta\gamma_{112}$, we focus on the difference between the opposite-sign and same-sign δ
604 correlators,

$$\Delta\delta = \delta_{\text{os}} - \delta_{\text{ss}}. \quad (12)$$

605 The background contributions due to the LCC and TMC have a similar characteristic structure that involves the
606 coupling between v_2 and $\Delta\delta$ [37, 38, 51, 126]. This motivates the study of the normalized quantity of $\Delta\gamma$ scaled by
607 v_2 and $\Delta\delta$, defined as:

$$\kappa_{112} \equiv \frac{\Delta\gamma_{112}}{v_2 \Delta\delta}. \quad (13)$$

608 The observation of the CME requires κ_{112} to be larger than $\kappa_{112}^{\text{TMC/LCC}}$. While a reliable estimate of $\kappa_{112}^{\text{TMC/LCC}}$ is
609 still elusive, the comparison of γ_{112} (and κ_{112}) between isobar collisions might give a more definite conclusion on the
610 CME signal.

611 It is intuitive to introduce some variations in the γ correlator to understand the background mechanisms in γ_{112} [26],
612 such as

$$\gamma_{123} = \langle \cos(\phi_\alpha + 2\phi_\beta - 3\Psi_3) \rangle. \quad (14)$$

613 This correlator is expected to be insensitive to the CME, because the correlation is negligible between the magnetic
614 field and the third harmonic plane, Ψ_3 . However, background due to flowing resonances along the Ψ_3 plane can
615 contribute to this observable. In analogy to Eq. (4) one can write:

$$\Delta\gamma_{123,\text{bkgd}} = \frac{4N_{2p}}{N^2} \langle \cos(\phi_\alpha - 2\phi_\beta - 3\phi_{2p}) \rangle v_{3,2p}. \quad (15)$$

616 Therefore, similar to Eq. (9) we also study the scaled quantity

$$\frac{\Delta\gamma_{123}}{v_3}. \quad (16)$$

617 Although the direct comparison of $\Delta\gamma_{112}/v_2$ and $\Delta\gamma_{123}/v_3$ is hard to interpret for a given system [127, 128], it
618 is useful to contrast signal and background scenarios by comparing each quantity between the two isobar systems.
619 When compared between the two isobars, in contrast to $\Delta\gamma_{112}/v_2$ which is driven by differences in both signal and
620 background, $\Delta\gamma_{123}/v_3$ will only be driven by the background difference. Since Ru+Ru has a larger magnetic field
621 than Zr+Zr, the CME expectation for mixed-harmonic measurements would be:

$$\frac{(\Delta\gamma_{112}/v_2)^{\text{Ru+Ru}}}{(\Delta\gamma_{112}/v_2)^{\text{Zr+Zr}}} > 1, \quad (17)$$

$$\frac{(\Delta\gamma_{112}/v_2)^{\text{Ru+Ru}}}{(\Delta\gamma_{112}/v_2)^{\text{Zr+Zr}}} > \frac{(\Delta\gamma_{123}/v_3)^{\text{Ru+Ru}}}{(\Delta\gamma_{123}/v_3)^{\text{Zr+Zr}}}, \quad (18)$$

$$\frac{(\Delta\gamma_{112}/v_2)^{\text{Ru+Ru}}}{(\Delta\gamma_{112}/v_2)^{\text{Zr+Zr}}} > \frac{(\Delta\delta)^{\text{Ru+Ru}}}{(\Delta\delta)^{\text{Zr+Zr}}}. \quad (19)$$

622 The last condition (Eq.19) can be re-written as

$$\frac{\kappa_{112}^{\text{Ru+Ru}}}{\kappa_{112}^{\text{Zr+Zr}}} > 1. \quad (20)$$

623 In general, the algebra relating κ , $\Delta\gamma$, v_2 , and $\Delta\delta$ relies on the symmetry assumption of $\langle \sin(\phi_\alpha - \phi_\beta) \sin n(\phi_\beta - \phi_c) \rangle =$
624 0 , with “ c ” labeling the particle used for EP reconstruction [26] and n representing the harmonic order. One can
625 circumvent this assumption by introducing a slight variant of κ that measures the factorization breaking:

$$k_n = \frac{\Delta \langle \cos(\Delta\phi_{\alpha\beta}) \cos(n\Delta\phi_{\beta c}) \rangle}{v_n^2 \{2\} \Delta\delta_{\alpha\beta}}. \quad (21)$$

626 Here the first “ Δ ” in the numerator denotes the difference between opposite-sign and same-sign measurements of the
 627 quantity inside the average. The quantity $\Delta\phi_{\alpha\beta} = \phi_\alpha - \phi_\beta$ denotes the relative azimuthal angle between charge-
 628 carrying particles, whereas the quantity $\Delta\phi_{\beta c} = \phi_\beta - \phi_c$ is the relative difference between one of the charge-carrying
 629 particles and the particles used for EP reconstruction. The quantity $\Delta\delta_{\alpha\beta}$ in the denominator has the same definition
 630 as Eq. (12). The quantity $v_n\{2\}$ is the n -th order harmonic anisotropy coefficients estimated using two-particle
 631 correlations. The CME is expected to cause an excess charge separation perpendicular to the Ψ_2 plane, whereas the
 632 background-driven charge separations along the Ψ_2 and Ψ_3 planes are proportional to v_2 and v_3 , respectively. Under
 633 these assumptions, one expects the case for the CME to be:

$$\frac{k_2^{\text{Ru+Ru}}}{k_2^{\text{Zr+Zr}}} > \frac{k_3^{\text{Ru+Ru}}}{k_3^{\text{Zr+Zr}}}. \quad (22)$$

634 For simplicity, the notation γ is used in place of γ_{112} in the following subsections (Sec. IV B-E).

635 B. Relative pseudorapidity dependence of $\Delta\gamma$

636 The relative pseudorapidity dependence of azimuthal correlations is widely studied to identify sources of long-range
 637 components that are dominated by early-time dynamics. They are contrasted to late-time correlations that are
 638 restricted by causality to appear as short-range correlations [129]. The same approach can be extended to charge-
 639 dependent correlations which provide the impetus to explore the dependence of $\Delta\gamma$ on the pseudorapidity gap between
 640 the charge-carrying particles $\Delta\eta_{\alpha\beta} = |\eta_\alpha - \eta_\beta|$ in $\langle \cos(\phi_\alpha(\eta_\alpha) + \phi_\beta(\eta_\beta) - 2\Psi_{\text{RP}}) \rangle$. Such measurements have been
 641 performed in STAR with Au+Au and U+U data [89, 130]. The possible sources of short-range correlations due to
 642 photon conversion to e^+e^- , HBT, and Coulomb effects can be identified and described as Gaussian peaks at small
 643 $\Delta\eta_{ab}$, the width and magnitude of which strongly depend on centrality and system size [131]. Going to more peripheral
 644 centrality bins, it becomes harder to identify such components as they overlap with sources of di-jet fragmentation
 645 that dominate both same-sign and opposite-sign correlations. Decomposing different components of $\Delta\gamma$ via study of
 646 $\Delta\eta_{ab}$ -dependence is challenging, although a clear sign of different sources of correlations is visible in the change of
 647 shape of individual same-sign and opposite-sign measurements of the γ correlator [89]. Nevertheless, these differential
 648 measurements of $\Delta\gamma$ in isobar collisions offer the prospects for studying the $\Delta\eta$ dependence of the CME. By comparing
 649 the differential measurements in Ru+Ru and Zr+Zr, it may be possible to extract the $\Delta\eta$ distribution of the CME
 650 signal, thus providing deeper insight into the origin of the phenomenon. The magnetic field driven CME signal is
 651 expected to dominate the long-range component $\Delta\eta_{\alpha\beta} > 1$ of the $\Delta\eta$ dependence like other early stage phenomena
 652 while the background due to resonance decay are expected to be short-range $\Delta\eta_{\alpha\beta} < 1$ [129]. In a CME scenario we
 653 expect the long-range component in Ru+Ru collisions to be larger than in Zr+Zr collisions.

654 C. Invariant mass dependence of $\Delta\gamma$

655 Since resonances present a large background source to the CME, the study of invariant mass (m_{inv}) dependence of
 656 the measured signal is natural and was first introduced in Ref. [63]. If we perform the analysis using pairs of pions,
 657 differential measurement of $\Delta\gamma$ with respect to m_{inv} should show peak-like structures similar to those in the relative
 658 pair multiplicity difference,

$$r = (N_{\text{OS}} - N_{\text{SS}})/N_{\text{OS}}, \quad (23)$$

659 if backgrounds from neutral resonances dominate the measurement. Here N_{OS} and N_{SS} are the numbers of opposite-
 660 sign and same-sign pion pairs, respectively. Indeed, similar peak structures are observed and an analysis utilizing
 661 the m_{inv} dependence and the ESE technique has been performed to extract the possible fraction of the CME signal
 662 in Au+Au collisions [69]. A similar analyses can be applied separately to the individual Ru+Ru and Zr+Zr data to
 663 extract a CME fraction in each system. Such an analysis will be performed in future work.

664 In this analysis we focus on contrasting the two isobar systems. We may gain insight into the mass dependence of
 665 the CME by combining the measurements in Ru+Ru and Zr+Zr collisions. Assuming in this blind analysis that the
 666 physics background is proportional to v_2 only (i.e. everything else is identical between the two isobar systems except
 667 v_2), we have

$$\Delta\gamma^{\text{Ru+Ru}} - a' \Delta\gamma^{\text{Zr+Zr}} = \Delta\gamma_{\text{CME}}^{\text{Ru+Ru}} - a' \Delta\gamma_{\text{CME}}^{\text{Zr+Zr}}, \quad (24)$$

668 where

$$a' = v_2^{\text{Ru+Ru}} / v_2^{\text{Zr+Zr}}. \quad (25)$$

669 The quantity a' can be safely assumed to be independent of m_{inv} , because the two isobar systems are similar. A CME
670 signature would be a positive measurement of the l.h.s. of Eq. (24):

$$\Delta\gamma^{\text{Ru+Ru}} - a'\Delta\gamma^{\text{Zr+Zr}} > 0. \quad (26)$$

671 Because the mass dependence of the CME signal is unlikely to differ between Ru+Ru and Zr+Zr collisions, such
672 a measurement would give unique insight on the mass dependence of the CME. Note Eq. (24) is valid for other
673 independent variables besides m_{inv} , such as the $\Delta\eta$ described in the previous subsection.

674 D. $\Delta\gamma$ with spectator and participant planes (approach-I)

675 This analysis makes use of the fact that the magnetic field driven signal is more correlated to the RP, in contrast to
676 flow-driven backgrounds which are maximal along the participant plane (PP). The idea was first published in Ref. [64]
677 and later discussed in Ref. [65]. It requires measurement of $\Delta\gamma$ with respect to the plane of produced particles, a proxy
678 for the PP, as well as with respect to the plane of spectators, a good proxy for RP. In STAR, the two measurements
679 can be done using Ψ_2 from the TPC and Ψ_1 from the ZDCs, respectively.

680 The approach is based on three main assumptions: 1) the measured $\Delta\gamma$ has contributions from signal and back-
681 ground, which can be decomposed as $\Delta\gamma = \Delta\gamma_{\text{bkg}} + \Delta\gamma_{\text{sig}}$, 2) the background contribution to $\Delta\gamma$ should follow the
682 scaling $\Delta\gamma_{\text{bkg}}\{\text{ZDC}\}/\Delta\gamma_{\text{bkg}}\{\text{TPC}\} = v_2\{\text{ZDC}\}/v_2\{\text{TPC}\}$, and 3) the signal contribution to $\Delta\gamma$ should follow the scaling
683 $\Delta\gamma_{\text{sig}}\{\text{ZDC}\}/\Delta\gamma_{\text{sig}}\{\text{TPC}\} = v_2\{\text{TPC}\}/v_2\{\text{ZDC}\}$. The first one has been known to be a working assumption, widely used
684 for a long time [16, 34]. The second one is borne out by the fact that backgrounds come from particle correlations
685 whose sources are v_2 modulated (see Eq. (4)) [36, 49–51]. The beauty of the method is that, because the TPC and
686 ZDC measurements are performed in identical events, all other factors contributing to $\Delta\gamma$ (such as resonance decay
687 correlations and multiplicity dilution) cancel except v_2 . Nevertheless, non-flow effects could potentially spoil the scal-
688 ing which requires quantitative investigations [71]. The validity of the third assumption is studied and demonstrated
689 in Ref. [64]. The reciprocal stems from fluctuations of RP and PP, whose relative azimuthal angle may be quantified
690 by $a = \langle \cos 2(\Psi_{\text{PP}} - \Psi_{\text{RP}}) \rangle$ [132].

691 Using all three assumptions, one can extract the fraction of a possible CME signal in a fully data-driven way [64],

$$f_{\text{CME}} = \frac{\Delta\gamma_{\text{CME}}\{\text{TPC}\}}{\Delta\gamma\{\text{TPC}\}} = \frac{A/a - 1}{1/a^2 - 1}, \quad (27)$$

692 where

$$A = \Delta\gamma\{\text{ZDC}\}/\Delta\gamma\{\text{TPC}\}, \quad (28)$$

693 and the a parameter can be determined by

$$a = v_2\{\text{ZDC}\}/v_2\{\text{TPC}\}. \quad (29)$$

694 The f_{CME} given by Eq. (27) is the fraction of CME contribution to the $\Delta\gamma\{\text{TPC}\}$ with respect to the TPC EP.

695 Such an analysis has been applied to existing Au+Au data, and a CME signal fraction of the order of 10% has
696 been extracted with a significance of 1–3 σ [70]. We apply the same analysis to the isobar data as part of the blind
697 analysis. The f_{CME} is extracted in each isobar system separately. The case for the CME in this analysis would be

$$f_{\text{CME}}^{\text{Ru+Ru}} > f_{\text{CME}}^{\text{Zr+Zr}} > 0. \quad (30)$$

698 One can get an additional constraint on $f_{\text{CME}}^{\text{Ru+Ru}}$ and $f_{\text{CME}}^{\text{Zr+Zr}}$. Assuming in this blind analysis that the physics
699 background is proportional to v_2 only,

$$(1 - f_{\text{CME}}^{\text{Ru+Ru}})\Delta\gamma^{\text{Ru+Ru}}/v_2^{\text{Ru+Ru}} = (1 - f_{\text{CME}}^{\text{Zr+Zr}})\Delta\gamma^{\text{Zr+Zr}}/v_2^{\text{Zr+Zr}}, \quad (31)$$

700 we obtain

$$f_{\text{CME}}^{\text{Ru+Ru}} = \left(\frac{a'}{A'}\right) f_{\text{CME}}^{\text{Zr+Zr}} + \left(1 - \frac{a'}{A'}\right), \quad (32)$$

701 where

$$A' = \Delta\gamma^{\text{Ru+Ru}}/\Delta\gamma^{\text{Zr+Zr}}, \quad (33)$$

702 and a' is again given by Eq. (25). The quantity a'/A' is the double ratio of

$$a'/A' = (\Delta\gamma^{\text{Zr+Zr}}/v_2^{\text{Zr+Zr}}) / (\Delta\gamma^{\text{Ru+Ru}}/v_2^{\text{Ru+Ru}}). \quad (34)$$

703 The individual measurements of $f_{\text{CME}}^{\text{Ru+Ru}}$ and $f_{\text{CME}}^{\text{Zr+Zr}}$ by Eq. (27) and the constraint on their relationship by Eq. (32)
704 give quantitatively an allowed region of the CME signal fractions.

E. $\Delta\gamma$ with participant and spectator planes (approach-II)

The main objective is to obtain the double ratio $(\Delta\gamma/v_2)_{\text{Ru+Ru}}/(\Delta\gamma/v_2)_{\text{Zr+Zr}}$. As discussed in Ref. [65] an evaluation of the ratio $(\Delta\gamma/v_2)$ does not require knowledge of the reaction plane resolution, which reduces the systematic uncertainty. It also “normalizes” the γ correlator to the elliptic flow value (which is proportional to background) and thus can be used for a direct comparison of the signals in different isobar collisions, even if the values of elliptic flow are slightly different in the two systems. Thus, the double ratio $(\Delta\gamma/v_2)_{\text{Ru+Ru}}/(\Delta\gamma/v_2)_{\text{Zr+Zr}}$, and specifically its deviation from unity, can be directly used for a qualitative detection of the CME signal. To extract the CME signal in this approach the double ratio is fit with the equation:

$$\frac{(\Delta\gamma/v_2)_{\text{Ru+Ru}}}{(\Delta\gamma/v_2)_{\text{Zr+Zr}}} = 1 + f_{\text{CME}}^{\text{Zr+Zr}} [(B_{\text{Ru+Ru}}/B_{\text{Zr+Zr}})^2 - 1], \quad (35)$$

where $f_{\text{CME}}^{\text{Zr+Zr}}$ is the CME fraction in the $\Delta\gamma$ correlator measured in Zr+Zr collisions, and $(B_{\text{Ru+Ru}}/B_{\text{Zr+Zr}})$ is the ratio of the magnetic field strengths in Ru+Ru and Zr+Zr collisions. By default this ratio is taken as the ratio of the nuclear charges, but can be varied to take into account the uncertainties related to the magnetic field determination.

For a non-zero CME signal it is expected that the double ratio $(\Delta\gamma/v_2)_{\text{Ru+Ru}}/(\Delta\gamma/v_2)_{\text{Zr+Zr}}$ would be greater than unity, as the CME signal in Ru+Ru collisions is expected to be about 15% larger than in Zr+Zr collisions [58, 83], and the background difference should be significantly smaller.

For the separate estimates of the CME signal in each of the isobar collisions, the γ correlator and elliptic flow can be also measured using STAR’s two ZDC-SMD Ψ_1 event planes (spectator planes):

$$(\Delta\gamma/v_2)_{\text{ZDC}} = \frac{\Delta\langle\cos(\phi_\alpha + \phi_\beta - \Psi_1^{\text{W}} - \Psi_1^{\text{E}})\rangle}{\langle\cos(2\phi - \Psi_1^{\text{W}} - \Psi_1^{\text{E}})\rangle}, \quad (36)$$

where $\Psi_1^{\text{W(E)}}$ is the event plane determined with ZDC-SMD in the west (east) side of STAR and the west side corresponds to the backward direction. Then this can be used for calculations of the double ratios:

$$\frac{(\Delta\gamma/v_2)_{\text{spectator}}}{(\Delta\gamma/v_2)_{\text{participant}}} = \frac{(\Delta\gamma/v_2)_{\text{ZDC}}}{(\Delta\gamma/v_2)_{\text{TPC}}} = \frac{\Delta\langle\cos(\phi_\alpha + \phi_\beta - \Psi_1^{\text{W}} - \Psi_1^{\text{E}})\rangle/\langle\cos(2\phi - \Psi_1^{\text{W}} - \Psi_1^{\text{E}})\rangle}{\Delta\langle\cos(\phi_\alpha + \phi_\beta - 2\phi_c)\rangle/\langle\cos(2\phi_\alpha - 2\phi_c)\rangle}. \quad (37)$$

To extract the signal, one has to make further assumptions [65]. Following the most plausible scenario of the magnetic field oriented on average perpendicular to the spectator plane, the CME fraction, $f_{\text{CME}}^{\text{TPC}}$, can be extracted via fitting of the results with the equation:

$$\frac{(\Delta\gamma/v_2)_{\text{ZDC}}}{(\Delta\gamma/v_2)_{\text{TPC}}} = 1 + f_{\text{CME}}^{\text{TPC}} \left(\frac{v_2^2\{\text{TPC}\}}{v_2^2\{\text{ZDC}\}} - 1 \right). \quad (38)$$

While the calculation of the double ratio, l.h.s. of Eq. (38), does not require knowledge of the reaction plane resolutions, the quantitative estimate of $f_{\text{CME}}^{\text{TPC}}$ from the double ratio requires v_2 values corrected for the reaction plane resolution. For the correlations relative to the sum of the first harmonic ZDC event planes the corresponding event plane resolution can be extracted directly from the data as $\langle\cos(\Psi_1^{\text{W}} - \Psi_1^{\text{E}})\rangle$.

F. R_{Ψ_2} variable

The $R_{\Psi_2}(\Delta S)$ variable provides an alternate way of measuring charge separation. It is obtained by taking the ratio of two sets of correlation functions [62, 73] defined as:

$$R_{\Psi_2}(\Delta S) = C_{\Psi_2}(\Delta S)/C_{\Psi_2}^\perp(\Delta S). \quad (39)$$

Here the correlation functions $C_{\Psi_2}(\Delta S)$ and $C_{\Psi_2}^\perp(\Delta S)$ quantify charge separation ΔS , perpendicular and parallel (respectively) to the Ψ_2 EP. The suffix “ \perp ” is motivated by the direction of the \vec{B} field. Since the \vec{B} field is nearly perpendicular to the Ψ_2 EP, $C_{\Psi_2}(\Delta S)$ and $C_{\Psi_2}^\perp(\Delta S)$ measure charge separation approximately parallel and perpendicular (respectively) to the \vec{B} field. These correlation functions are further obtained from the ratios of two distributions [62, 73];

$$C_{\Psi_2}(\Delta S) = \frac{N_{\text{real}}(\Delta S)}{N_{\text{shuffled}}(\Delta S)}. \quad (40)$$

Here, $N_{\text{real}}(\Delta S)$ is the distribution of the quantity ΔS that measures the event-by-event average of the charge separation:

$$\Delta S = \frac{\sum_{i=1}^{n^+} w_i^+ \sin(\Delta\phi_i - \Psi_2)}{\sum_{i=1}^{n^+} w_i^+} - \frac{\sum_{i=1}^{n^-} w_i^- \sin(\Delta\phi_i - \Psi_2)}{\sum_{i=1}^{n^-} w_i^-}. \quad (41)$$

Here, n^- and n^+ are the numbers of negatively and positively charged particles in an event, w_i^\pm are weights that account for non-uniformity of the azimuthal acceptance of the TPC and $\phi_i - \Psi_2$. The distribution $N_{\text{shuffled}}(\Delta S)$ is obtained in a way similar to that for $N_{\text{real}}(\Delta S)$ but after random reassignment (shuffling) of the charges of the reconstructed tracks in each event. This randomization makes $N_{\text{shuffled}}(\Delta S)$ insensitive to charge-dependent correlations and ensures identical event property between the numerator and the denominator in Eq. (40). The correlation function $C_{\Psi_2}^\perp(\Delta S)$ is constructed in a way similar to Eq.40 by replacing Ψ_2 with $\Psi_2 + \pi/2$. Both $C_{\Psi_2}(\Delta S)$ and $C_{\Psi_2}^\perp(\Delta S)$ have nearly Gaussian shapes around $\Delta S = 0$.

The final variable $R_{\Psi_2}(\Delta S)$, obtained according to Eq.39 measures the relative charge separation between parallel and perpendicular directions to the \vec{B} field. CME-driven charge separation along the \vec{B} field is expected to lead to concave-shaped $R_{\Psi_2}(\Delta S)$ distributions with stronger CME signals leading to narrower (more concave) distributions [62]. The width of $R_{\Psi_2}(\Delta S)$ reflects the magnitude of charge separation, which is also influenced by particle number fluctuations and resolution of the EP. Both increase the width ($\sigma_{\Delta_{\text{Sh}}}$) of the R-variable. The effect of the particle number fluctuations can be accounted for by scaling ΔS by the width $\sigma_{\Delta_{\text{Sh}}}$ of the N_{shuffled} distribution, i.e., $\Delta S' = \Delta S/\sigma_{\Delta_{\text{Sh}}}$. This re-scaled distribution of $\Delta S'$ can be further corrected for EP resolution. This is done by using a parametrized function $\delta_R = \text{Res} \times e^{(1-\text{Res})^2}$ to correct $\Delta S'$ to $\Delta S'' = \Delta S' \times \delta_R$. Here Res is the EP resolution. This approach of correction has been verified using simulation studies [62] and data-driven tests.

After the analysis code was frozen, the R_{Ψ_3} [62, 74] observable, constructed to be insensitive to CME, was found to have a programming error that failed to convert some integers to floats, along with an issue in azimuthal periodicity (for more details see Ref. [76]). Consequently, R_{Ψ_3} results are not included in this paper.

Assuming Ru+Ru collisions produce a stronger magnetic field than Zr+Zr collisions, a signature for CME-driven charge separation would be indicated by the observation

$$1/\sigma_{R_{\Psi_2}}^{\text{Ru+Ru}} > 1/\sigma_{R_{\Psi_2}}^{\text{Zr+Zr}}, \quad (42)$$

where $\sigma_{R_{\Psi_2}}$ is the Gaussian width of the respective $R(\Delta S'')$ distribution.

V. ISOBAR BLIND ANALYSIS RESULTS

The major background in the primary CME-sensitive observable, $\Delta\gamma_{112}$, is due to elliptic flow, v_2 . Therefore, we first examine the v_2 measurements from the blind analysis. The upper panels of Fig. 4 show a compilation of the v_2 results obtained by different analysis groups using different methods and reported in the following subsections. For clarity, not all results from all groups are shown in Fig. 4. The v_2 values from different methods are expected to be different [99], and we found consistency in results among different groups using the same method. The lower panels of Fig. 4 show the v_2 ratio between Ru+Ru and Zr+Zr collisions. All the ratios, except noticeably the $v_2\{4\}$ and $v_2\{\Psi_{\text{ZDC}}\}$ ratios, fall on a common curve. The ratios are above unity by 2–3% in mid-central collisions and fall off towards peripheral and central collisions, with the exception of the top 5% centrality bin, where the ratios are also above unity by a few percent. The central-collision results are likely due to a larger quadrupole deformity in $^{96}_{44}\text{Ru}$ compared with $^{96}_{40}\text{Zr}$, which needs future investigation. The above-unity ratio in mid-central collisions may originate from the different nuclear structures between the two isobars as predicted by the DFT calculations [85, 86]. These v_2 ratios imply different magnitudes of the CME backgrounds in the two isobar systems, and this effect is taken into account in the $\Delta\gamma_{112}/v_2$ and κ_{112} observables described in Sec. IV. It is often advantageous to study the CME observables with Ψ_{ZDC} , which measures the spectator plane, and is more correlated with the magnetic field than the participant plane. However, the $v_2\{\Psi_{\text{ZDC}}\}$ ratio is significantly larger than unity; this comes primarily from the better alignment of the spectator and participant planes in Ru+Ru than in Zr+Zr collisions, as predicted by the DFT nuclear structure calculations [85]. This would imply that the advantage in using the isobar difference or ratio to search for the CME with Ψ_{ZDC} is limited [85]. The qualitative consistencies between $v_2\{\Psi_{\text{ZDC}}\}$ and $v_2\{4\}$ and between their respective Ru+Ru/Zr+Zr ratios are observed as expected in a Gaussian model description of flow fluctuations [133].

In the following subsections, we discuss the results of the CME-related observables from the five independent analysis groups, each focusing on a few specific observables related to measurements from specific detectors. The




FIG. 4. (Left) Elliptic anisotropy v_2 measurements using different methods in isobar collisions at $\sqrt{s_{NN}} = 200$ GeV as a function of centrality using TPC and EPD detectors. In the upper panels, the solid and open symbols represent measurements for Ru+Ru and Zr+Zr collisions, respectively. The data points are shifted along the x axis for clarity. The lower panels show the v_2 ratios in Ru+Ru over Zr+Zr collisions. The statistical uncertainties are represented by lines and systematic uncertainties by boxes. (Right) The same showing measurements for four particle correlations using TPC and EP determined from ZDC. The data points are shifted horizontally for clarity.

784 detailed implementations differ among the groups with regards to estimation of harmonic flow vectors, re-weighting,
 785 the pseudorapidity gap to reduce non-flow, and correction of non-uniform acceptance. While focusing on various
 786 aspects, four of the five groups have analyzed the $\Delta\gamma/v_2$ observable. Figure 5 compares the $\Delta\gamma/v_2$ measurements
 787 with both the full-event and sub-event methods. The statistical uncertainties are largely correlated among the
 788 groups because the same initial data sample is analyzed; the results are not identical because of the analysis-specific
 789 event selection criteria (see Table.I) and the slightly different methods. Using the Barlow approach [106], we have
 790 verified that the results from different groups are consistent within the statistical fluctuations due to those differences.
 791 Moreover, the final conclusion on the observability of the CME is consistent among all five analysis groups.

792 In addition to the centrality dependence results reported in the following subsections, in order to have the best
 793 statistics, we also quote the final results for the Ru+Ru over Zr+Zr ratio observables for the centrality range of 20–50%.
 794 The choice of this centrality range is determined by two considerations. One is that the mid-central collisions present
 795 the best EP reconstruction resolution as well as the most significant magnetic field strengths (hence the possibly
 796 largest CME signal difference between the isobar species). The other consideration is that the online trigger efficiency
 797 starts to deteriorate from the 50% centrality mark towards more-peripheral collisions (see Sec. III). A compilation of
 798 results from different groups is presented in the summary subsection VI.

799 A. $\Delta\gamma$ measurements with TPC event plane (Group-1)

800 The flow plane for a specific pseudorapidity range is unknown for each event. In practice, we estimate an n^{th} -
 801 harmonic flow plane with the azimuthal angle (Ψ_n) of the flow vector $\vec{Q}_n = (\sum_i^N w_i \cos(n\phi_i), \sum_i^N w_i \sin(n\phi_i))$, where
 802 ϕ_i represents the azimuthal angle of a detected particle, and w_i is a weight (often set to p_T) to optimize the EP
 803 resolution. For example, the v_n measurement with respect to the full TPC EP is denoted by

$$v_n\{\text{TPC EP}\} = \langle \cos(n\phi - n\Psi_n^{\text{TPC}}) \rangle. \quad (43)$$




FIG. 5. Ru+Ru over Zr+Zr ratio of the primary CME-sensitive observable $\Delta\gamma/v_2$ estimated using different methods and by independent analysis groups. The vertical lines represent statistical uncertainties while the rectangular boxes represent systematic uncertainties. The upper panel shows the results using the entire TPC acceptance estimated using event-plane (EP) and three-point correlations (3PC) methods without any η gaps. The lower panel shows the results using a sub-event (SE) method with a gap ($\Delta\eta_{\text{sub}}$) of 0.2. Note the most central data point from Group-4 is for 0–10% centrality. The centrality bins are shifted horizontally for clarity.

804 The corresponding γ_{112} correlator is represented by

$$\gamma_{112}\{\text{TPC EP}\} = \langle \cos(\phi_\alpha + \phi_\beta - 2\Psi_2^{\text{TPC}}) \rangle. \quad (44)$$

805 The two-particle δ correlator is estimated in the same way as defined in Eq. (11). To account for the detector
 806 non-uniformity, both ϕ and Ψ_2^{TPC} have been corrected by the shifting method [99], such that they have uniform
 807 distributions.

808 In this subsection, the POIs (with azimuthal angle represented by ϕ in Eq. (43) or ϕ_α and ϕ_β in Eq. (44)) are taken
 809 from the TPC acceptance of $|\eta| < 1$. By default, the full EP over the same η range is used for the v_2 and $\Delta\gamma_{112}$
 810 measurements, with no η gap between the EP and the POIs or between the two POIs. For each POI or POI pair,
 811 the full EP is re-estimated by excluding the POI or POI pair to remove self-correlation. This approach yields the
 812 smallest statistical uncertainties, with the largest possible number of POIs and the highest possible EP resolution.
 813 The systematic uncertainties due to the lack of an η gap are expected to be canceled to a large extent in the ratio
 814 between the two isobar systems, and this idea has been corroborated by the v_2 ratios in Fig. 4, and will be further
 815 tested in the following discussions of the results using finite η gaps.

816 Figure 6 shows $v_2\{\text{TPC EP}\}$ as a function of centrality for Ru+Ru and Zr+Zr collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV
 817 in the upper panel, and the ratio of Ru+Ru to Zr+Zr in the lower panel. The v_2 ratio averaged over the 20–50%
 818 centrality range is $1.0144 \pm 0.0001(\text{stat.}) \pm 0.0008(\text{syst.})$. Given the statistical and systematic uncertainties, this value is
 819 significantly above unity, and we consider two potential origins: (a) the two nuclei could have different nuclear density
 820 parameters, and (b) non-flow contributions could be different in the two systems. Scenario (b) can be examined using
 821 the measurements with various η gaps: the mean value of the v_2 ratio becomes 1.0146, 1.0149 and 1.0161 for the
 822 two-particle cumulant method ($v_2\{2\}$ defined in Eq. (45)) with no η gap, $\Delta\eta_{\alpha\beta} > 0.05$ and $\Delta\eta_{\alpha\beta} > 0.2$, respectively.
 823 Here $\Delta\eta_{\alpha\beta}$ is the η gap between particles α and β . Since the v_2 ratio is consistently above unity, we exclude the
 824 non-flow explanation. Therefore, the isobar data indicate that the $^{96}_{44}\text{Ru}$ and $^{96}_{40}\text{Zr}$ nuclei have different nuclear density
 825 distributions, yielding a larger eccentricity in Ru+Ru than in Zr+Zr collisions at a given centrality [85]. This results
 826 in the v_2 ratio in the lower panel of Fig. 6 being larger than unity.




FIG. 6. v_2 measured with the full TPC EP for Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV (upper panel) and the ratio of Ru+Ru to Zr+Zr (lower panel). The centrality bins are shifted horizontally for clarity. The border-less horizontal bands denote the statistical uncertainties. The horizontal bands with the dashed border represent the systematic uncertainties.




FIG. 7. $\Delta\delta$ measured for Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV (upper panel) and the ratio of Ru+Ru to Zr+Zr (lower panel). The centrality bins are shifted horizontally for clarity. The border-less horizontal bands denote the statistical uncertainties. The horizontal bands with the dashed border represent the systematic uncertainties.

827 Figure 7 shows $\Delta\delta$ vs centrality for Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV in the upper panel, and the
 828 ratio of Ru+Ru to Zr+Zr in the lower panel. There is no η gap between the two POIs. The $\Delta\delta$ ratio averaged over
 829 the 20–50% centrality range is $0.9851 \pm 0.0003(\text{stat.}) \pm 0.0002(\text{syst.})$, below unity with high measured significance.
 830 The central value of the $\Delta\delta$ ratio changes to 0.9846 and 0.9833 with $\Delta\eta_{\alpha\beta} > 0.05$ and $\Delta\eta_{\alpha\beta} > 0.2$, respectively. Thus
 831 the short-range correlations have a very small impact on the $\Delta\delta$ ratio.

832 Figure 8 shows $\Delta\gamma_{112}$ as a function of centrality measured with the full TPC EP for Ru+Ru and Zr+Zr collisions
 833 at $\sqrt{s_{NN}} = 200$ GeV in the upper panel, and the ratio of Ru+Ru to Zr+Zr in the lower panel. By default, no η gap is
 834 applied between the two POIs or between the EP and the POIs. The $\Delta\gamma_{112}$ ratio averaged over the 20–50% centrality
 835 range is $0.9828 \pm 0.0034(\text{stat.}) \pm 0.0005(\text{syst.})$. When a finite η gap is applied between the two POIs, the central value
 836 of the $\Delta\gamma_{112}$ ratio becomes 0.9822 and 0.9825 with $\Delta\eta_{\alpha\beta} > 0.05$ and $\Delta\eta_{\alpha\beta} > 0.2$, respectively. Therefore, the $\Delta\gamma_{112}$
 837 ratio is insensitive to the short-range correlations.

838 Figure 9 shows κ_{112} vs centrality measured with the full TPC EP for Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$
 839 GeV in the upper panel, and the ratio of Ru+Ru to Zr+Zr in the lower panel. The default κ_{112} ratio averaged
 840 over the 20–50% centrality range is $0.9838 \pm 0.0035(\text{stat.}) \pm 0.0009(\text{syst.})$, which changes to 0.9827 and 0.9831 with
 841 $\Delta\eta_{\alpha\beta} > 0.05$ and $\Delta\eta_{\alpha\beta} > 0.2$, respectively. We conclude that the CME signature predefined in Eq. (20) is not
 842 observed in this blind analysis of the isobar data. It is noteworthy that we have reached a precision better than 0.4%
 843 on these measurements of the ratio between Ru+Ru and Zr+Zr collisions.

844 After unblinding of the isobar species, we observe the multiplicity difference between the two isobar systems at a
 845 given centrality, as shown in Table III. Although the effects of the multiplicity mismatch are largely canceled in the
 846 ratio of $\Delta\gamma_{112}$ over $v_2\Delta\delta$, there could still be residual contributions driving the κ_{112} ratio below unity, which needs
 847 further investigation. Additional discussions on the multiplicity mismatch can be found in Sec. VI.




FIG. 8. $\Delta\gamma_{112}$ measured with the full TPC EP for Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV (upper panel) and the ratio of Ru+Ru to Zr+Zr (lower panel). The centrality bins are shifted horizontally for clarity. The border-less horizontal bands denote the statistical uncertainties. The horizontal bands with the dashed border represent the systematic uncertainties.




FIG. 9. κ_{112} measured with the full TPC EP for Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV (upper panel) and the ratio of Ru+Ru to Zr+Zr (lower panel). The centrality bins are shifted horizontally for clarity. The border-less horizontal bands denote the statistical uncertainties. The horizontal bands with the dashed border represent the systematic uncertainties.

B. Mixed harmonic measurements (Group-2)

While the analysis from the previous group focuses on the EP method, in this subsection: 1) we focus on measurements of harmonic coefficients and charge sensitive correlations using two-point, three-point correlations and the scalar-product method, and 2) we further extend the correlation measurements by requiring one of the particles from the forward EPD.

We measure harmonic flow coefficients $v_n\{2\}$ from the full TPC using two-particle correlations, where

$$v_{n=2,3}^2\{2\}(|\eta| < 1) = \langle \cos(n\phi_1 - n\phi_2) \rangle. \quad (45)$$

In this $v_n^2\{2\}$ measurement from the TPC, we put a cut of $\Delta\eta_{1,2} > 0.05$ to mitigate effects of two track merging and e^+e^- due to photon conversion. For $v_n^2\{2\}$ measurements, we remove the short-range component due to HBT, and Coulomb effects using a double Gaussian fit as described in Ref. [96]. We also estimate harmonic coefficients without such Gaussian subtraction but using a cut of $\Delta\eta_{1,2} > 1$ in Eq. (45). In this paper we denote such measurements as $v_n\{2\}(\Delta\eta > 1)$. In addition we also estimate v_n using sub-event methods $v_n^2\{\text{SP}\} = \langle Q_{n,a} Q_{n,b}^* \rangle$, where the Q -vectors $Q_{n,a}$ and $Q_{n,b}$ are taken from two halves of TPC around $\eta = 0$ separated by a pseudorapidity gap of $\Delta\eta_{\text{sub}} = 0.2$. We denote such measurements as $v_n\{\text{SP}\}(\Delta\eta_{\text{sub}} = 0.2)$.

We present measurements of data from the new EPD detector ($2.1 < |\eta| < 5.1$). We estimate the elliptic and triangular anisotropy of particles at mid-rapidity with respect to the forward participant planes in the EPD by

$$v_{n=2,3}\{\text{SP}\}(\text{TPC-EPD}) \equiv \langle \cos(n\phi - n\Psi_n^{\text{EPD}}) \rangle = \frac{\langle Q_{n,\text{TPC}} Q_{n,\text{EPDE}}^* + Q_{n,\text{TPC}} Q_{n,\text{EPDW}}^* \rangle}{2\sqrt{\langle Q_{n,\text{EPDE}} Q_{n,\text{EPDW}}^* \rangle}}, \quad (46)$$

using the scalar-product (SP) method where Q and Q^* denote the Q -vectors and their complex conjugates [134].




FIG. 10. (Left) Elliptic and triangular anisotropies measured for the two systems using the combination of TPC and EPDs. The boxes represent systematic uncertainties and the lines represent statistical errors. (Right) Compilation of v_3 using different methods and cuts on pseudorapidity separation. Results are shown for individual systems in open symbols for Zr+Zr and solid symbols for Ru+Ru. Results are also shown for the ratio of Ru+Ru to Zr+Zr in open symbols. The centrality bins are shifted horizontally for clarity.

865 The upper and middle panels of Fig. 10 (left) show the centrality dependence of v_2 and v_3 , respectively, with
 866 the two aforementioned approaches. The measurements of these flow harmonics using only TPC and TPC-EPD
 867 are noticeably different, especially in peripheral events for v_2 , and in mid-central and peripheral collisions for v_3 .
 868 A possible explanation for such an observation could be the pseudorapidity dependence of non-flow, de-correlation
 869 and flow fluctuations [133, 135]. Owing to low multiplicity and poor resolution of the third-order EP, EPDs do not
 870 allow for the v_3 measurements beyond 60-70% centrality. A compilation of v_3 results is shown in Fig. 10 (right) to
 871 demonstrate the effect of pseudorapidity separation between POI and EP (or between two POIs).

872 The lower panel (left) presents ratios of the flow harmonics for Ru+Ru over Zr+Zr collisions, with a few interesting
 873 features. First, the v_2 ratio in the most central events (0–5%) is larger than unity with high significance. As mentioned
 874 before, effects due to nuclear deformation can lead to the difference in the shape even in fully-overlapping collisions,
 875 which needs to be confirmed by future studies. Above-unity v_2 ratios are also observed in mid-central collisions.
 876 This is consistent with the expectation of the eccentricity ratios from nuclear density distributions calculated by
 877 DFT [85, 86]. Second, the v_3 ratio is significantly below unity in central events, which is counter intuitive, as v_3 is
 878 supposed to be driven by fluctuations in central collisions. Third, the v_3 ratio significantly deviates from unity in
 879 peripheral events, and this deviation has a dependence on pseudorapidity separation between POI and EP. Thus, we
 880 need a better understanding of the possible differences in the nuclear structure and the deformity of the isobars, when
 881 comparing the two systems at the same centrality. Further exploration along this direction is beyond the scope of
 882 this paper which is primarily focused on the CME blind analysis. These v_n measurements do have implications on
 883 the background contribution to CME that is relevant in the scaled charge separation variables.

884 We perform the measurement of charge separation using the full TPC acceptance ($|\eta| < 1$) in the following way
 885

$$886 \gamma_{112}^{\alpha,\beta}(\eta_\alpha, \eta_\beta)(|\eta| < 1) \equiv \left\langle \cos \left(\phi_\alpha(\eta_\alpha) + \phi_\beta(\eta_\beta) - 2\Psi_2^{|\eta| < 1} \right) \right\rangle = \frac{\langle \cos(\phi_\alpha(\eta_\alpha) + \phi_\beta(\eta_\beta) - 2\phi_c) \rangle}{v_{2,c}\{2\}}. \quad (47)$$

886 The indices “ α, β, c ” denote three distinctly different particles. The subscripts “ α, β ” denote particle pairs with same
 887 (SS) or opposite (OS) sign of electric charges. We use the charge-inclusive reference particle ‘ c ’ as a proxy for the
 888 elliptic flow plane Ψ_2 at midrapidity, and the quantity $v_{2,c}\{2\}$ refers to the two-particle elliptic flow coefficient of the
 889 reference particle ‘ c ’ that we estimate using two-particle correlations as defined in Eq. (45).




FIG. 11. Scaled charge separation across the second and third harmonic EPs obtained using all three particles from the TPC acceptance, divided by the anisotropy coefficient. Results are shown for Ru+Ru and Zr+Zr collisions separately on the upper and middle panels over the centrality range of 0-80%. The centrality bins are shifted horizontally for clarity. The lower panel shows the ratio of various quantities for 20–50% centrality. The border-less horizontal bands over the 20–50% centrality range with different colors represent the statistical uncertainties in the combined centrality for different observables. The horizontal bands with the dashed border represent the systematic uncertainties. The N_{part} scaling is applied in the upper two panels to improve the visibility. The N_{part} scaling is not included in the lower panel for the ratios.




FIG. 12. Scaled charge separation across second and third harmonic EPs scaled by the anisotropy coefficient obtained using all three particles from the TPC acceptance but using a sub-event (SE) from $-1 < \eta < -0.1$ and $0.1 < \eta < 1$. Results are shown for Ru+Ru and Zr+Zr collisions separately in the upper and middle panels over the centrality range of 0-80%. The centrality bins are shifted horizontally for clarity. The lower panel shows the ratio of different quantities for 20–50% centrality. The border-less horizontal bands over 20–50% centrality range with different colors represent the statistical uncertainties on the combined centrality for different observables. The horizontal bands with the dashed border represent the systematic uncertainties. The N_{part} scaling is applied in the upper and middle panels to improve the visibility. The N_{part} scaling is not included in the lower panel for the estimation of ratios.

890 Similarly with respect to the third harmonic plane, we measure

$$891 \quad \gamma_{123}^{\alpha,\beta}(\eta_\alpha, \eta_\beta)(|\eta| < 1) \equiv \left\langle \cos \left(\phi_\alpha(\eta_\alpha) + 2\phi_\beta(\eta_\beta) - 3\Psi_3^{|\eta| < 1} \right) \right\rangle = \frac{\langle \cos(\phi_\alpha(\eta_\alpha) + 2\phi_\beta(\eta_\beta) - 3\phi_c) \rangle}{v_{3,c}\{2\}}. \quad (48)$$

892 Finally we calculate the quantities of interest:

$$893 \quad \Delta\gamma_{1mn} = \gamma_{1mn}^{\text{OS}} - \gamma_{1mn}^{\text{SS}}, \text{ and, } \Delta\gamma_{1mn}/v_n \times N_{\text{part}}. \quad (49)$$

894 The normalization of $\Delta\gamma_{1mn}$ ($m, n = 1, 2$ or $2, 3$) by v_n ($n = 2, 3$) takes into account the flow-driven background due to resonance decays and local charge conservation [36, 51]. The N_{part} scaling compensates for the trivial dilution of correlations expected from superposition of independent sources, and improves the visibility of the data points.

895 The upper and middle panels of Fig. 11 show the CME-sensitive $\Delta\gamma_{112}/v_2$ and the CME-insensitive $\Delta\gamma_{123}/v_3$ (both multiplied by N_{part}), respectively, for individual species. The lower panel presents the ratios of the quantities for the 20–50% centrality bin in Ru+Ru over Zr+Zr collisions. Note that the ratios do not involve N_{part} , whose values are different for the two isobar systems at the same centrality (see Sec. III). The ratio of the quantity $\langle \Delta\gamma_{112}/v_2 \rangle$ is

0.966 \pm 0.005, while the ratio for $\langle\Delta\gamma_{123}/v_3\rangle$ is 0.971 \pm 0.019. The errors quoted here are statistical only. Systematic variation in our measurements for these quantities are not statistically significant, and are estimated to be zero according to the Barlow approach as described in Section-II F. The $\Delta\delta$ ratio in this analysis is 0.9849 \pm 0.0004(stat.) \pm 0.0005(syst.), which is consistent with the value (0.9846 \pm 0.0003 \pm 0.0002) obtained by Group-1 within the statistical and systematic uncertainties. In summary, our observation related to charge separation in the 20–50% centrality is consistent with the following statements:

$$\frac{(\Delta\gamma_{112}/v_2)^{\text{Ru+Ru}}}{(\Delta\gamma_{112}/v_2)^{\text{Zr+Zr}}} < 1, \quad (50)$$

$$\frac{(\Delta\gamma_{112}/v_2)^{\text{Ru+Ru}}}{(\Delta\gamma_{112}/v_2)^{\text{Zr+Zr}}} \approx \frac{(\Delta\gamma_{123}/v_3)^{\text{Ru+Ru}}}{(\Delta\gamma_{123}/v_3)^{\text{Zr+Zr}}}, \quad (51)$$

$$\frac{(\Delta\gamma_{112}/v_2)^{\text{Ru+Ru}}}{(\Delta\gamma_{112}/v_2)^{\text{Zr+Zr}}} < \frac{(\Delta\delta)^{\text{Ru+Ru}}}{(\Delta\delta)^{\text{Zr+Zr}}}. \quad (52)$$

Therefore, our measurements are not consistent with any of the predefined CME signatures as set out in Eqs. (17), (18), and (19).




FIG. 13. Charge separation across the second- and third-order EPs scaled by the anisotropy coefficient obtained using particles from the TPC acceptance and hits from the EPDs. Results are shown for Ru+Ru and Zr+Zr collisions separately on the upper and middle panels over the centrality range of 0–80%. The centrality bins are shifted horizontally for clarity. The lower panel shows the ratio of different quantities for 20–50% centrality. The border-less horizontal bands over 20–50% centrality range with different colors represent the statistical uncertainties on the combined centrality for different observables. The horizontal bands with the dashed border represent the systematic uncertainties. The N_{part} scaling is applied in the upper and middle panels to improve the visibility. The N_{part} scaling is not included in the lower panel for the estimation of ratios.




FIG. 14. Factorization breaking coefficient for the second and third order harmonics measured using particles from the TPC acceptance. Results are shown for Ru+Ru and Zr+Zr collisions separately on the upper and middle panels over the centrality range of 0–80%. The centrality bins are shifted horizontally for clarity. The lower panel shows the ratio of various quantities for 20–50% centrality. The border-less horizontal bands over 20–50% centrality range with different colors represent the statistical uncertainties on the combined centrality for different observables. The horizontal bands with the dashed border represent the systematic uncertainties.

907 We repeat our analysis with sub-events (SE) within the STAR TPC, as described in Eq. (47), and divide the TPC
 908 acceptance into two halves, with $0.1 < \eta < 1$ and $-1 < \eta < -0.1$. When the charge-carrying particles “ α ” and “ β ” are
 909 taken from one half, particle “ c ” is taken from the other. Going from full TPC to sub-event reduces the maximum
 910 value of the relative pseudorapidity $\Delta\eta_{\alpha,\beta}$ from 2 to 0.9. The sub-events are also used for the v_n estimation. In our
 911 blind analysis code we did not include the estimation of $\Delta\delta$ in the acceptance of $0.1 < |\eta| < 1$ which would have been
 912 most appropriate. The $\Delta\delta$ ratio used here are using the full TPC acceptance which is an approximation. The results
 913 from this approach are presented in Fig. 12, and show some noticeable differences from those in Fig. 11, illustrating
 914 the influence of using sub-events. However, the final ratios in the lower panel are consistent with those obtained with
 915 the full TPC, and hence we still do not observe any of the predefined CME signatures (Eqs. (17), (18), and (19)) in
 916 this measurement.

917 We extend our measurement of charge separation with respect to the 2nd harmonic plane from the STAR EPD
 918 using

$$\begin{aligned} \gamma_{112}^{\alpha,\beta}(\eta_\alpha, \eta_\beta)(\text{TPC-EPD}) &\equiv \langle \cos(\phi_\alpha(\eta_\alpha) + \phi_\beta(\eta_\beta) - 2\Psi_2^{\text{EPD}}) \rangle \\ &= \frac{\langle Q_{1,\text{TPC}}^\alpha Q_{1,\text{TPC}}^\beta Q_{2,\text{EPDE}}^* + Q_{1,\text{TPC}}^\alpha Q_{1,\text{TPC}}^\beta Q_{2,\text{EPDW}}^* \rangle}{2\sqrt{\langle Q_{2,\text{EPDE}} Q_{2,\text{EPDW}}^* \rangle}}. \end{aligned} \quad (53)$$

919 Here, we use algebra based on Q -vectors [136], defined as $Q_n = \sum_1^M w_i e^{in\phi} / \sum_1^M w_i$, where M is the number of
 920 particles, and the weight factor w_i accounts for the imperfection in the detector acceptance in bins of η - ϕ , p_T (track-
 921 curvature), $V_{z,\text{TPC}}$, and centrality. After weight correction, the Q -vectors are also “recentered” as $Q - \langle Q \rangle$ for residual
 922 acceptance correction. Similarly we measure charge separation with respect to a 3rd harmonic plane from the EPD
 923 using

$$\begin{aligned} \gamma_{123}^{\alpha,\beta}(\eta_\alpha, \eta_\beta)(\text{TPC-EPD}) &\equiv \langle \cos(\phi_\alpha(\eta_\alpha) + 2\phi_\beta(\eta_\beta) - 3\Psi_3^{\text{EPD}}) \rangle \\ &= \frac{\langle Q_{1,\text{TPC}}^\alpha Q_{2,\text{TPC}}^\beta Q_{3,\text{EPDE}}^* + Q_{1,\text{TPC}}^\alpha Q_{2,\text{TPC}}^\beta Q_{3,\text{EPDW}}^* \rangle}{2\sqrt{\langle Q_{3,\text{EPDE}} Q_{3,\text{EPDW}}^* \rangle}}. \end{aligned} \quad (54)$$

924 Measurements of charge sensitive variables using the combination of the TPC and the EPD are shown in Fig. 13.
 925 Once again these results are consistent with earlier measurements using the full TPC acceptance. Based on the
 926 measurements of ratios of variables in Ru+Ru over Zr+Zr collisions, we do not see any of the predefined CME
 927 signatures (Eqs. (17), (18), and (19)).

928 The quantities k_2 and k_3 that measure factorization breaking defined in Eq. (21) are shown in Fig. 14. Since this
 929 measurement involves several higher-order charge-sensitive correlators, we restrict our measurements to the full TPC
 930 only to achieve the best precision. Averaged over the 20–50% centrality, the k_2 and k_3 ratios are 0.986 ± 0.004 (stat.)
 931 ± 0.002 (syst.) and 0.982 ± 0.019 (stat.) ± 0.006 (syst.), respectively, consistent with each other within one standard
 932 deviation. In other words we find:

$$\frac{k_2^{\text{Ru+Ru}}}{k_2^{\text{Zr+Zr}}} \approx \frac{k_3^{\text{Ru+Ru}}}{k_3^{\text{Zr+Zr}}}. \quad (55)$$

933 The predefined CME signature described in Eq. (22) is not observed.

934 C. Differential $\Delta\gamma$ measurements in pseudorapidity (Group-2)

935 Relative pseudorapidity dependence between the charge-carrying particles ($\Delta\eta_{\alpha,\beta}$) of same-sign and opposite-sign
 936 γ_{112} correlators is shown in Fig. 15 (left) for 20–50% Ru+Ru and Zr+Zr collisions. We show two panels in which
 937 the third particle or the EP is either obtained from TPC or the EPD. The $\Delta\eta_{\alpha,\beta}$ dependence of γ correlator in the
 938 individual isobar species have very similar shapes compared to what is reported in the previous STAR measurement
 939 in Au+Au collisions [130]. Some difference in the shape is observed between measurements using TPC and EPD EPs.
 940 The same is also seen for the $\Delta\eta_{\alpha,\beta}$ dependence of $\Delta\gamma_{112}$ shown in Fig.15 (right). Although interesting dependence is
 941 observed for the individual distributions we do not see any species dependence within the uncertainties of the current
 942 measurements. The expectation for CME was that the long-range part of the $\Delta\eta$ distribution $\Delta\eta_{\alpha,\beta} > 1$ will be
 943 higher for Ru+Ru collisions. No such observation can be made from the results shown in Fig. 15.




FIG. 15. Relative pseudorapidity dependence of the γ (left) and $\Delta\gamma$ (right) correlators shown for two species for same-sign (SS) and opposite-sign (OS) correlations. For clarity, the alternate bins in $\Delta\eta$ are shown. The statistical and systematic errors are shown by vertical lines and square boxes, respectively. Within the uncertainty we do not see any species dependence in these measurements.

D. Differential $\Delta\gamma$ measurements in invariant mass (Group-3)

In order to isolate the resonance background contributions, we report measurements of the $\Delta\gamma$ variable, differential in pair invariant mass m_{inv} . This analysis uses the three-point correlation method to calculate the γ correlators [19, 20],

$$\gamma = \langle \cos(\phi_\alpha + \phi_\beta - 2\phi_c) \rangle / v_{2,c}, \quad (56)$$

where the average $\langle \dots \rangle$ runs over all triplets and over all events. To select good events we require, in addition to those criteria described in Sec. IIB, the VPD primary vertex position to be within $|V_{z,\text{TPC}} - V_{z,\text{VPD}}| < 3$ cm from the one reconstructed by the TPC. The POIs (α and β) are pions within $0.2 < p_T < 1.8$ GeV/ c . They are identified by their specific energy loss in the TPC and their flight time obtained from the TOF detector. The c particles are charged hadrons within $0.2 < p_T < 2.0$ GeV/ c . The POIs and particle c are all taken from $|\eta| < 1$ (self-correlations are avoided) [19, 20]. An η gap of 0.05 is applied between the POIs. No η gap is applied between particle c and either of the POIs. The $v_{2,c}$ of particle c is calculated from two-particle correlations by the $v_2\{2\}$ of Eq. (45) where an η gap of 1.0 is applied between the two particles.

The systematic uncertainties are assessed according to Sec. IIF. In addition, the η gap between the POIs (i.e. between α and β) is varied from 0.05 (default) to 0 (i.e. no gap) and 0.2. The η gap used to determine the $v_{2,c}$ is varied from 1 to 0.5 and 1.4. All systematic uncertainties are added in quadrature.

Figure 16 shows the distributions in the relative pair multiplicity difference of Eq. (23) in Ru+Ru and Zr+Zr collisions in the 20–50% centrality range in the upper panel and their ratio in the lower panel. The ratio has a weak dependence on m_{inv} , with an average value in the 20–50% centrality range of $r^{\text{Ru+Ru}}/r^{\text{Zr+Zr}} = 0.9705 \pm 0.0008$ (stat.). It deviates from unity because the isobar systems do not have the same multiplicity when their centrality defined by cross section percentile is matched (see Sec. III). Note that the r is measured with pion pairs; it does not necessarily equal that of charged hadrons. Also note that the r ratio does not necessarily equal the inverse multiplicity ratio because the difference $N_{\text{OS}} - N_{\text{SS}}$ may not strictly scale with multiplicity.

The upper panel of Fig. 17 shows the $\Delta\gamma$ results in Ru+Ru and Zr+Zr collisions in the 20–50% centrality range as a function of m_{inv} . Resonance peaks are observed in $\Delta\gamma$ corresponding to those in r in Fig. 16. The lower panel shows the $\Delta\gamma$ difference for the isobars after the $\Delta\gamma$ for Zr+Zr is scaled by the v_2 ratio (see Eq. (24)). A constant fit to the




FIG. 16. (a) Distributions in the relative pair multiplicity difference, $r = (N_{OS} - N_{SS})/N_{OS}$, as a function of invariant mass of $\pi^+\pi^-$ pairs in 20–50% Ru+Ru and Zr+Zr collisions and (b) their ratio. Errors shown are statistical. The solid line in the lower panel is a constant fit to the ratio.




FIG. 17. The $\Delta\gamma$ in 20–50% Ru+Ru and Zr+Zr collisions (a) and their difference defined by Eq. (24) (b) as functions of the $\pi^+\pi^-$ invariant mass m_{inv} . The difference in the lower panel would measure the possible CME if the background in $\Delta\gamma$ scales with v_2 only ($a' = v_2^{Ru+Ru}/v_2^{Zr+Zr}$ as defined by Eq. (25)). Error bars are statistical and shaded boxes are systematic uncertainties. The solid line in the lower panel is a constant fit to the data.

968 measured difference in the 20–50% centrality range yields $\Delta\gamma^{Ru+Ru} - a'\Delta\gamma^{Zr+Zr} = (-4 \pm 2 \text{ (stat.)} \pm 6 \text{ (syst.)}) \times 10^{-6}$.
 969 The predefined CME signature of a positive value for this difference (Eq. (26)) is not observed.

970 As described in Sec. IV C, the predefined CME signature described in Eq. (26) explicitly assumes the r ratio to be
 971 unity. Since this assumption is no longer valid for the blind analysis binned in cross-section percentile, as shown in
 972 Fig. 16 lower panel, the relevance of the result in Fig. 17 to the possible CME needs to be reevaluated.

973 E. CME fraction utilizing spectator and participant planes: approach-I (Group-3)

974 The CME signal fraction, f_{CME} , is extracted from two $\Delta\gamma$ measurements in each of the two isobar systems inde-
 975 pendently. One measurement is with respect to the second-order harmonic plane reconstructed from mid-rapidity
 976 particles measured in the TPC, as a proxy for the PP. The other is with respect to the first-order harmonic plane re-
 977 constructed from spectator neutrons measured by the ZDC Shower Maximum Detectors (ZDC-SMDs), as a proxy for
 978 the spectator plane. The details of this spectator-participant plane method to extract f_{CME} is described in Sec. IV D.
 979 To select good events we require, in addition to those criteria described in Sec. II B, the VPD primary vertex position
 980 to be within $|V_{z,TPC} - V_{z,VPD}| < 3$ cm from the one reconstructed by the TPC. In this analysis both the full-event and
 981 sub-event methods are used as in Ref. [70]. The sub-event method is useful to suppress non-flow effects.

982 For the full-event analysis, all three particles are charged hadrons taken from $|\eta| < 1$. The $\Delta\gamma\{\text{TPC}\}$ is calculated
 983 by the three-particle correlation method (Eq. (56)). An η gap of 0.05 is applied between the POIs (α and β); no η gap
 984 is applied between particle c and either of the POIs. The $v_{2,c}$ used in Eq. (56) and the $v_2\{\text{TPC}\}$ needed by Eq. (29)
 985 are equal and are calculated by the two-particle cumulant method of Eq. (45), where no η gap is applied between the
 986 two particles.

For the sub-event analysis, the $\Delta\gamma\{\text{TPC}\}$ and $v_2\{\text{TPC}\}$ are calculated by the EP method (Eqs. (43) and (44)). Each TPC event is divided into two sub-events with $-1 < \eta < -0.05$ and $0.05 < \eta < 1$ (thus an η gap of 0.1 is in between). The POIs are charged particles from one sub-event, and the EP is calculated using charged particles from the other sub-event with $0.2 < p_T < 5$ GeV/c. An η gap of 0.05 is applied between the POIs (α and β). For EP reconstruction, the azimuthal nonuniformity of the efficiency and acceptance of the TPC is corrected by applying a multi-dimensional ϕ -dependent weight. The shifting method is performed to further flatten the EP azimuthal distribution. For $\Delta\gamma\{\text{ZDC}\}$, the POIs are still from one sub-event to keep the same acceptance as that for $\Delta\gamma\{\text{TPC}\}$.

For both the full-event and sub-event methods, the $\Delta\gamma\{\text{ZDC}\}$ and $v_2\{\text{ZDC}\}$ are calculated by the EP method, where the γ correlators are given by

$$\gamma = \langle \cos(\phi_\alpha + \phi_\beta - 2\Psi_{\text{ZDC}}) \rangle, \quad (57)$$

and

$$v_2\{\text{ZDC}\} = \langle \cos 2(\phi - \Psi_{\text{ZDC}}) \rangle. \quad (58)$$

The ZDC EP angle Ψ_{ZDC} is calculated from the Q -vector combined from both ZDC-SMDs. Recentering and shifting are applied to flatten the azimuthal distribution of the reconstructed ZDC EP. The TPC and ZDC EP resolutions are obtained by the conventional method of an iterative procedure [99], using correlations between two η sub-events for the former and between the two ZDC-SMDs for the latter.

The systematic uncertainties are assessed according to Sec. II F. In addition, the η gap between the POIs is varied from 0.05 (default) to 0 (i.e. no gap) and 0.2. For the sub-event method, the η gap between the sub-events is varied from 0.1 (default) to 0.3. All systematic uncertainties are added in quadrature.




FIG. 18. The $\Delta\gamma$ (top row), v_2 (second row), A or a (third row), and A/a (bottom row) as functions of centrality in Ru+Ru (left column) and Zr+Zr (right column) collisions. The $\Delta\gamma$ and v_2 are calculated with respect to the TPC and ZDC harmonic planes. Results are from the full-event method. Error bars are statistical and shaded boxes are systematic uncertainties.

1004

1005

1006

1007

1008

Figure 18 shows the results from the full-event method as functions of centrality in Ru+Ru (left column) and Zr+Zr (right column) collisions. The $\Delta\gamma$ and v_2 with respect to the TPC and ZDC harmonic planes are shown in the first and second rows. The third row shows the ratios A (Eq. (28)) and a (Eq. (29)). The double ratio A/a is shown in the




FIG. 19. The CME signal fraction in the inclusive $\Delta\gamma$ measurement with respect to the TPC EP, f_{CME} , as functions of centrality in Ru+Ru and Zr+Zr collisions at 200 GeV from both the full-event method (a) and the sub-event method (b). Error bars are statistical and shaded boxes are systematic uncertainties.

bottom row. The measured $v_2\{\Psi_{\text{ZDC}}\}$ in central collisions is noticeably large, which is not observed in the results from another group reported in Sec. V F. A difference in the analysis method lies in the way to calculate the ZDC harmonic planes. In this analysis, the first-order harmonic Q -vectors from the two ZDCs are first combined and then the Ψ_{ZDC} is computed. In the analysis in Sec. V F, the correlation is performed with the sum of the two first-order harmonic planes, separately reconstructed in each ZDC. Correspondingly, the EP resolutions are calculated in different ways. The reason for the discrepancy needs further investigation.

We calculate the f_{CME} using Eq. (27). This is the fraction of the CME contribution to the $\Delta\gamma\{\text{TPC}\}$ with respect to the TPC EP. The results from both the full-event and sub-event methods are shown in Fig. 19 as a function of centrality for the two isobar collision systems. The average f_{CME} values in the 20–50% centrality range from the full-event method in Ru+Ru and Zr+Zr collisions are $f_{\text{CME}}^{\text{Ru+Ru}} = 0.29 \pm 0.13$ (stat.) ± 0.01 (syst.) and $f_{\text{CME}}^{\text{Zr+Zr}} = 0.06 \pm 0.08$ (stat.) ± 0.02 (syst.), respectively. The corresponding ratios from the sub-event method are $f_{\text{CME}}^{\text{Ru+Ru}} = 0.12 \pm 0.20$ (stat.) ± 0.00 (syst.) and $f_{\text{CME}}^{\text{Zr+Zr}} = -0.01 \pm 0.12$ (stat.) ± 0.03 (syst.). Systematic variations for $f_{\text{CME}}^{\text{Ru+Ru}}$ are all consistent with statistical fluctuations so a null systematic uncertainty is assigned according to the Barlow prescription [106]. The large statistical uncertainties are dominated by the $\Delta\gamma$ measurements with respect to the ZDCs which have poor EP resolutions.

Figure 20 plots f_{CME} for Ru+Ru collisions on the vertical axis versus f_{CME} for Zr+Zr collisions on the horizontal axis, both at 20–50% centrality, extracted using both the full-event method (left panel) and sub-event method (right panel). An additional constraint is obtained by combining the $\Delta\gamma$ measurements with respect to the TPC EP in both isobar collision systems, as described in Sec. IV D. This is shown in the near-diagonal shaded strip given by Eq. (32) using the measured values for the double ratio (Eq. (34)) of $a'/A' = 1.027 \pm 0.004$ (stat.) ± 0.001 (syst.) and 1.034 ± 0.006 (stat.) ± 0.003 (syst.), for the full-event and sub-event methods, respectively. This would be the correct constraint if the background in $\Delta\gamma$ scales with v_2 only (i.e. the multiplicities are explicitly assumed to be identical between the isobar systems). Since this assumption is no longer valid for the blind analysis as function of the cross-section percentile, the near-diagonal strip does not correctly indicate the allowed CME region. Indeed, as shown in Fig. 20, the present near-diagonal strip does not have overlap with the CME region enclosed by the blue solid lines expected from the magnetic field difference. The relevance of the near-diagonal strip to f_{CME} needs to be




FIG. 20. Constraints on f_{CME} in Ru+Ru vs. f_{CME} in Zr+Zr collisions, both for 20–50% centrality, with 95% confidence level ($\pm 2\sigma$). The left panel shows the full-event results and the right panel shows the sub-event results. The shaded rectangle on the left is the constraint extracted from each individual isobar collision system using the spectator-participant plane method and Eq. (27). The near-diagonal shaded strip would be the constraint extracted from the TPC measurements combining both isobar systems using Eq. (32), provided the background in $\Delta\gamma$ scales with v_2 only ($a' = v_2^{\text{Ru+Ru}}/v_2^{\text{Zr+Zr}}$ as defined by Eq. (25)). The two solid lines indicate the expectation from the magnetic field difference: $f_{\text{CME}}^{\text{Ru+Ru}}/f_{\text{CME}}^{\text{Zr+Zr}} = (1 + \alpha_B)\Delta\gamma^{\text{Zr+Zr}}/\Delta\gamma^{\text{Ru+Ru}} \approx 1 + \alpha_B$, where $\alpha_B = 0.15 \pm 0.05$ is assumed.

revisited in the future by using the properly scaled a' .

F. Ratio of $(\Delta\gamma/v_2)$ between two isobar collisions (Group-4)

One of the main objectives of Group-4 is to obtain the double ratio $(\Delta\gamma/v_2)_{\text{Ru+Ru}}/(\Delta\gamma/v_2)_{\text{Zr+Zr}}$ as a function of centrality. The quantity $(\Delta\gamma/v_2)$ is calculated as

$$(\Delta\gamma/v_2)_{\text{TPC}} = \frac{\Delta\langle\cos(\phi_\alpha + \phi_\beta - 2\phi_c)\rangle}{\langle\cos(2\phi_\alpha - 2\phi_c)\rangle}, \quad (59)$$

where Δ denotes the difference in the γ correlator calculated using opposite-charge and same-charge pairs of particles α and β . The correlator is calculated using the subevents from pseudorapidity windows $0.1 < |\eta| < 1.0$ (default) and $0.2 < |\eta| < 1.0$, with the event plane, or particle “c”, taken from the opposite pseudorapidity window (e.g., when $-0.1 > \eta_{\alpha,\beta} > -1.0$ we take $0.1 < \eta_c < 1.0$ and vice versa) with pseudorapidity gaps between the subevents $\Delta\eta_{\text{sub}} = 0.2$ (for the default case) and $\Delta\eta_{\text{sub}} = 0.3, 0.4$ (for systematic studies). To suppress the non-flow contribution, $\langle\cos(2\phi_\alpha - 2\phi_c)\rangle$ is calculated using the same-charge particles in the default case and using all charged particles when investigating systematic uncertainties. All particles are taken from the transverse momentum region $0.2 < p_T < 2.0$ GeV/c. The results are calculated in 5% centrality bins and then averaged over a wider centrality range using the inverse of squared statistical uncertainty as a weight when needed.

All quantities in this analysis are obtained with the help of the recentered Q vectors and presented as ratios, which greatly reduces the systematic uncertainties. The systematic uncertainty has been estimated from comparison of the results obtained with different η -gaps between the sub-events, using selection criteria on quality of the TPC tracks, and comparing results from events with the event vertex from different sides of the TPC center. In addition, in the estimates of the elliptic flow uncertainties, the results obtained from correlation of unlike-sign charges are also used. All the systematic variations are found to be smaller than the statistical uncertainties.

For a non-zero CME signal the expectation is that the double ratio $(\Delta\gamma/v_2)_{\text{Ru+Ru}}/(\Delta\gamma/v_2)_{\text{Zr+Zr}}$ would be greater than unity, as the CME signal in Ru+Ru collisions is expected to be about 15% larger than in Zr+Zr collisions. The results of our measurements are presented in Fig. 21. The plotted ratio is below unity, which is likely due to a noticeable difference in mean charged multiplicity in collisions of the two isobar species corresponding to the same

1058 centrality. The multiplicity of charged particles in Ru+Ru collisions is observed to be larger than that in Zr+Zr
 1059 collisions as shown in Fig. 3. The drop of the double ratio in most peripheral events is likely due to the sudden change
 1060 in the multiplicity ratio in the corresponding centrality.




FIG. 21. Double ratio $(\Delta\gamma/v_2)_{Ru+Ru}/(\Delta\gamma/v_2)_{Zr+Zr}$ as a function of centrality for isobar collisions, where shaded boxes represent systematic uncertainties.

1061 The quantity $(\Delta\gamma/v_2)$ approximately scales with the inverse of the multiplicity, but no correction for that is
 1062 anticipated in the blind analysis. The fraction of the CME signal contribution to $\Delta\gamma$, if extracted exactly as outlined
 1063 in the blind analysis scheme in the 20 to 50% centrality range, would yield a negative value with an uncertainty of
 1064 about 2% of the $\Delta\gamma$ magnitude.

1065 G. CME fraction utilizing spectator and participant planes: approach-II (Group 4)

1066 For the separate estimates of the CME signal in each of the isobar collisions, the procedure outlined in section IV E,
 1067 Eqs. (36)–(38) was used.

1068 The results obtained in this approach are presented in Fig. 22. We observe that the double ratio, Fig. 22 (left), is very
 1069 close to unity indicating that the signal, is consistent with zero in both isobar collisions. The fraction of the CME signal
 1070 calculated using Eq. (38), is presented in Fig. 22 (right), while elliptic flow calculated relative to the participant (TPC)
 1071 and spectator (ZDC-SMD) planes is presented in Fig. 23. The extracted average CME fraction for 20–50% centrality
 1072 is found to be $f_{CME}^{TPC} = 0.101 \pm 0.123$ (stat.) ± 0.023 (syst.) for Ru+Ru and $f_{CME}^{TPC} = 0.009 \pm 0.088$ (stat.) ± 0.033 (syst.)
 1073 for Zr+Zr. The large statistical uncertainties are dominated by the $\Delta\gamma$ measurements in the ZDCs which have poor
 1074 EP resolutions. The statistical uncertainties on f_{CME} are smaller than those from Group-3 reported in section IV D,
 1075 due to a larger difference in $v_2\{\text{ZDC}\}$ and $v_2\{\text{TPC}\}$ resulting from different approaches of correlating particles at
 1076 midrapidity with signals from the two ZDCs.




FIG. 22. The ratio of $\Delta\gamma/v_2$ (left) and the CME fraction (right) for Ru+Ru and Zr+Zr collisions from spectator/participant plane analysis. Shaded boxes represent systematic uncertainties and hatched areas represent 1σ uncertainties (combined statistical and systematic uncertainties) of the CME fraction for 20–50% centrality, with the mean values indicated by horizontal solid lines. The data of Zr+Zr collisions are shifted horizontally for clarity.




FIG. 23. $v_2\{2\}$, $v_2\{4\}$, and $v_2\{ZDC\}$ for isobar collisions as a function of centrality in the top panel. The data of Zr+Zr collisions are slightly shifted along x-axis for better visibility. Ratios of v_2 between the two systems are plotted in the bottom panel. Open boxes represent systematic uncertainties.

H. R_{Ψ_2} measurements (Group-5)

1077

1078 In this part of the analysis, charged particles with transverse momentum $0.2 < p_T < 2.0$ GeV/ c are used to construct
 1079 Ψ_2 . Each event is subdivided into two sub-events with pseudorapidity $0.1 < \eta < 1.0$ (West) and $-1.0 < \eta < -0.1$
 1080 (East) to obtain Ψ_2^W (West) and Ψ_2^E (East). Afterward, $C_{\Psi_2}(\Delta S)$, $C_{\Psi_2}^\perp(\Delta S)$ and $R_{\Psi_2}(\Delta S)$ are constructed using
 1081 charged particles with $0.35 < p_T < 2.0$ GeV/ c . To avoid potential self-correlations, Ψ_2^E is used for particles within the
 1082 $0.1 < \eta < 1.0$ range and Ψ_2^W for particles within the $-1.0 < \eta < -0.1$ range. Here the ΔS distributions associated
 1083 with the aforementioned quantities are symmetrized around $\Delta S = 0$. The second p_T selection (beginning at 0.35
 1084 GeV/ c) is chosen to minimize the influence of acceptance effects at low p_T while optimizing the statistics.

1085 The sensitivity of the $R_{\Psi_2}(\Delta S'')$ distribution to the potential impact from v_2 -driven background is investigated
 1086 using event-shape selection via fractional cuts on the magnitude of the second harmonic flow Q -vector q_2 relative to
 1087 its maximum value $q_{2,\max}$ at fixed multiplicity [59]. This study is motivated by the fact that v_2 drives background
 1088 sources of CME and the change in q_2 provides a lever-arm to vary v_2 [27, 32]. Therefore, the impact of the v_2 -driven
 1089 charge separation background can be decreased (increased) by choosing events with smaller (larger) q_2 values.




FIG. 24. The q_2 dependence of the $R_{\Psi_2}(\Delta S'')$ distributions for Ru+Ru (a) and Zr+Zr (d) for 20–50% collisions. Panels (b) and (e) show the corresponding q_2 -dependent v_2 values; panels (c) and (f) show the inverse widths ($\sigma_{R_{\Psi_2}}^{-1}$) for distributions in (a) and (d), respectively. The distributions shown in (a) and (d) are symmetrized around $\Delta S'' = 0$. $0.2 < p_T < 2.0$ GeV/ c are used to reconstruct Ψ_2 . A second p_T selection with $0.35 < p_T < 2.0$ GeV/ c are used for particle of interests.

1090 Event-shape selection is performed using three sub-events; $A[\eta < -0.3]$, $B[|\eta| < 0.3]$, and $C[\eta > 0.3]$, following the
 1091 methods described earlier, and with q_2 selections in sub-event B . Figure. 24 shows the q_2 -selected isobar measurements.
 1092 The $R_{\Psi_2}(\Delta S'')$ distributions are given in panels a and d, and the corresponding v_2 values, measured using the two
 1093 sub-event cumulants method [137] and particles with $0.35 < p_T < 2.0$ GeV/ c are shown in panels b and e. The inverse
 1094 widths (panels c and f) are extracted from the distributions shown in (panels a and d). Linear fits to the data in panels
 1095 (b), (c), (e), and (f) indicate that, while v_2 shows a $32.0\% \pm 0.01\%$ increase with q_2 from $q_2=0$ -20% to 60-100%, the
 1096 corresponding inverse width for the $R_{\Psi_2}(\Delta S'')$ distributions show an approximate decrease of $7.0\% \pm 4.0\%$. Further
 1097 studies may be needed to understand the physics behind the observed behavior of the widths of R_{Ψ_2} on q_2 .

1098 The $R_{\Psi_2}(\Delta S'')$ distributions, extracted for several centrality selections in Ru+Ru and Zr+Zr collisions, are shown
 1099 in Fig. 25 (a-d). They indicate centrality-dependent concave-shaped distributions for $R_{\Psi_2}(\Delta S'')$. The corresponding
 1100 inverse widths extracted from these distributions are shown in panel (e). They indicate similar magnitudes for both

1100




FIG. 25. Comparison of the $R_{\Psi_2}(\Delta S'')$ distributions obtained for charged particles in (a) 0-10%, (b) 10-30%, (c) 30-50% and (d) 20-50% collisions in Ru+Ru and Zr+Zr collisions at $\sqrt{s_{NN}} = 200$ GeV. Panel (e) shows the centrality dependence of the inverse widths $\sigma_{R_{\Psi_2}}^{-1}$, extracted from the $R_{\Psi_2}(\Delta S'')$ distributions. Panel (f) shows the ratio of the inverse widths of the two isobars. The distributions shown in (a)-(d) are symmetrized around $\Delta S'' = 0$. $0.2 < p_T < 2.0$ GeV/c are used to reconstruct Ψ_2 . A second p_T selection with $0.35 < p_T < 2.0$ GeV/c are used for particle of interests.

1101 isobars that increase as collisions become more peripheral. The difference between the inverse widths for the two isobars
 1102 is made more transparent in Fig. 25(f), where the ratios $\sigma_{R_{\Psi_2}}^{-1}(\text{Ru+Ru}) / \sigma_{R_{\Psi_2}}^{-1}(\text{Zr+Zr})$ are plotted as a function of
 1103 collision centrality. Note that the systematic uncertainty is negligible compared to the statistical uncertainties for the
 1104 20-50% selection.

1105

I. Summary and discussions

1106 The elliptic flow v_2 coefficients are found to be larger in Ru+Ru than Zr+Zr collisions, by approximately 2% in
 1107 mid-central collisions and by a similar amount in the most central 5% of collisions. The shape and magnitude of the
 1108 $v_2^{\text{Ru+Ru}} / v_2^{\text{Zr+Zr}}$ ratio as a function of centrality are consistent with the corresponding eccentricity ratio predicted by
 1109 DFT calculations [85, 86], which can be parameterized by neutron-halo type WS distributions for the ${}^{96}_{40}\text{Zr}$ nucleus [113].
 1110 Therefore, the current measurements are consistent with the different intrinsic nuclear structures of the two isobars.
 1111 The v_2 difference in central collisions suggests that the ${}^{96}_{44}\text{Ru}$ nucleus is more deformed than the ${}^{96}_{40}\text{Zr}$ nucleus. However,
 1112 the ratio of multiplicity distribution is best described by MC-Glauber simulations without intrinsic shapes for both the
 1113 isobars. Further studies with more sophisticated observables are underway to pin down the nuclear shape difference
 1114 between ${}^{96}_{44}\text{Ru}$ and ${}^{96}_{40}\text{Zr}$. Using the forward detectors EPD and ZDC rather than the TPC to determine the EP leads
 1115 to a noticeable change in the magnitude of v_2 and an even larger change in v_3 . These changes may primarily be
 1116 due to effects of non-flow, longitudinal de-correlation and flow-fluctuations. An interesting observation is that the
 1117 magnitudes of v_3 differ with high significance between the two isobars in both peripheral and central collisions, which
 1118 warrants future investigation.

1119 The primary CME-sensitive observable $\Delta\gamma/v_2$ is analyzed by four independent groups. Prior to the blind analysis,
 1120 the case for observation of a CME signal is predefined to be an excess of $\Delta\gamma/v_2$ in Ru+Ru collisions as compared with
 1121 Zr+Zr collisions. Results from all groups are inconsistent with this expectation, and therefore no conclusive evidence
 1122 of the CME is found in this blind analysis. The analysis from one group uses an alternate CME-sensitive measure,
 1123 namely the R variable. The predefined expectation for the CME for this observable is a larger concavity of the R
 1124 variable in Ru+Ru collisions compared with Zr+Zr collisions. No such observation is found in the data, and therefore




FIG. 26. Compilation of results from the blind analysis. Only results contrasting between the two isobar systems are shown. Results are shown in terms of the ratio of measures in Ru+Ru collisions over Zr+Zr collisions. Solid dark symbols show CME-sensitive measures whereas open light symbols show counterpart measures that are supposed to be insensitive to CME. The vertical lines indicate statistical uncertainties whereas boxes indicate systematic uncertainties. The colors in the background are intended to separate different types of measures. The fact that CME-sensitive observable ratios lie below unity leads to the conclusion that no predefined CME signatures are observed in this blind analysis.

1125 no conclusive evidence of the CME is observed using the R variable in the blind analysis.

1126 Figure 26 presents a compilation of results from the blind analysis for the 20–50% centrality range. In this figure, the
 1127 ratio of the value of each observable in Ru+Ru to its value in Zr+Zr collisions is shown; the statistical and systematic
 1128 uncertainties are shown by lines and boxes, respectively. Included are results for the CME-sensitive observables
 1129 $\Delta\gamma/v_2$, κ , k and $1/\sigma_{R\psi_2}$ using different detector combinations as well as from independent analysis groups. The
 1130 ratio values of $\Delta\gamma/v_2$, κ_{112} , k_2 , and $1/\sigma_{R\psi_2}$ are all less than or consistent with unity, indicating that the predefined
 1131 CME signature is not observed in the isobar blind analysis for any of these observables. This observation is further
 1132 corroborated by the observation that the CME-insensitive quantities $\Delta\gamma_{123}/v_3$ and k_3 have ratios (as shown in the
 1133 figure) consistent with their second-harmonic CME-sensitive counterparts.

1134 In addition to the integrated quantities shown in Fig. 26, we have performed differential measurements of $\Delta\gamma$ with
 1135 $\Delta\eta$ and of $\Delta\gamma$ for pion pairs in invariant mass m_{inv} for both isobar species. No difference in the shape is observed
 1136 between the two species in these differential studies. The mean value of the variable r that measures the relative
 1137 excess of opposite-sign relative to same-sign pion pairs at different values of m_{inv} , is different for the two isobar species,
 1138 being smaller in Ru+Ru collisions; this is qualitatively consistent with the charged hadron multiplicity difference in
 1139 bins of matching centrality between the two isobars.

1140 The comparison of $\Delta\gamma$ measured with respect to the spectator (measured by the ZDC) and participant (measured
 1141 by the TPC) planes is used to extract the CME fraction f_{CME} in each individual species. Two analysis groups used this
 1142 method. Group-3 analyzed both the full-event and sub-event correlations, while Group-4 analyzed only the latter.
 1143 Using the sub-events allows the suppression of non-flow correlations. The sub-event results from the two groups are
 1144 consistent with each other. The statistical uncertainties on f_{CME} from Group-3 are larger than those from Group-4,
 1145 due to a smaller difference in $v_2\{\text{ZDC}\}$ and $v_2\{\text{TPC}\}$ resulting from different approaches of correlating particles at
 1146 midrapidity with signals from two ZDCs (see sections IV D and IV E). All these results give a CME signal fraction
 1147 that is consistent with zero with large statistical uncertainties of approximately 10% (absolute) dominated by the
 1148 ZDC measurements.

1149 The most recent Au+Au results measured by the spectator and participant plane method from STAR indicate a
 1150 possible CME signal fraction of the order of 10% with a significance of 1–3 σ [70]. If the CME signal fraction is also
 1151 10% in isobar collisions, then a 3 σ effect would be expected with the current isobar data sample of approximately 2
 1152 billion MB events each, according to estimations in Ref. [82, 83]. However, it has been pointed out and supported by
 1153 AVFD simulations that the CME signal fraction may be substantially smaller in isobar collisions compared to Au+Au
 1154 collisions [138]. This would imply a substantially smaller significance in this isobar data sample.

VI. POST BLINDING




FIG. 27. Compilation of post-blinding results. This figure is largely the same as Fig. 26 with the following differences: numerical changes in the results from the new run-by-run QA algorithm are treated as an additional systematic uncertainty added in quadrature, and two data points (open markers) have been added on the right to indicate the ratio of inverse multiplicities ($N_{\text{irk}}^{\text{offline}}$) and the ratio of relative pair multiplicity difference (r) as explained in the text.

During the second step of our analysis (the isobar blind analysis) a potential issue was identified related to the predefined criteria of the QA algorithm (as described in Sec. IID). The condition of being within five times the weighted error or one percent of the variation of the local mean, may be too relaxed to identify all the boundaries of stable run periods and outlier runs in some QA variables. When combining the identified run mini-regions, a new algorithm is implemented by 1) removing the “within one percent of the variation of the local mean” condition, and 2) adding a tolerance of “within 2-RMS difference”, which seems to be more effective for some QA variables such as N_{fits} . This new algorithm is again executed in the final step of isobar unblind analysis (Step-3) and all the results using this algorithm are presented in this post-blinding section. No qualitative changes are observed in the final quantities. The numerical changes in the results from this new run-by-run QA algorithm are treated as an additional systematic uncertainty to update Fig.26 and obtain Fig. 27.

Two additional data points are included on Fig. 27 for the following reasons. Most ratio quantities shown in Fig. 26 or Fig.27 have magnitudes that are below unity with high significance, whereas in a purely non-CME scenario with controlled backgrounds, the expectation is that these quantities should be consistent with unity. The reason for these ratios being less than unity is, in part, due to the multiplicity difference in the two isobar systems. As documented in Table III, the multiplicity distributions are different for the two isobar species to the extent that in bins of matching centrality, the mean multiplicity is around 4% lower for mid-central Zr+Zr than for mid-central Ru+Ru collisions. The measured magnitudes of most observables, such as $\Delta\gamma$ and $\Delta\delta$, decrease with increasing multiplicity because of the trivial multiplicity dilution for these per-pair quantities. Therefore, the corresponding ratios of these observables between the two isobar systems will become larger, if taken in bins of matching multiplicity. Under the approximation that background to $\Delta\gamma$ is caused by flowing clusters with the properties of the clusters staying the same and the number of clusters scaling with multiplicity, the value of $\Delta\gamma$ scales with the inverse of multiplicity [20], i.e. $N\Delta\gamma \propto v_2$ with the proportionality presumably equal between the two isobars. Because of this, it may be considered that the proper baseline for the ratio of $\Delta\gamma/v_2$ between the two isobars is the ratio of the inverse multiplicities of the two systems. Analysis with respect to this baseline is not documented in the pre-blinding procedures of this blind analysis, so is not reported as part of the blind analysis. We include this inverse multiplicity ratio as the right-most point in Fig. 27.

It is interesting to note that ordering among the quantities in their magnitudes is observed in Figs. 26 and 27. The $\Delta\gamma/v_2$ ratio has a smaller magnitude than the κ and k ratios. This is consistent with the multiplicity ratio baseline for the former as discussed above, and the fact that the trivial multiplicity dependence cancels in the latter, so its baseline would be unity. On the other hand, the R -variable inverse width $1/\sigma_{R,\psi_2}$ ratio is larger than the $\Delta\gamma/v_2$ ratio. This difference is expected to be driven by: 1) different p_T ranges used for the two quantities, 2) difference in the multiplicity dependence (see, e.g., Ref. [81]), and 3) difference in the non-flow contributions. The scaling relations extracted in Ref. [81] indicate an approximate relation between $1/\sigma_{R,\psi_2}^2$, multiplicity N and $\Delta\gamma$, which would imply

for this analysis $1/\sigma_{R_{\psi_2}}^2 \approx N\Delta\gamma$; an estimate based on the measurements from this analysis indicates this ratio for Ru+Ru over Zr+Zr to be approximately 1.02.

It is not clear that the inverse multiplicity ratio discussed above is the best baseline to use to take into account the multiplicity difference; for example an alternative would be the ratio of excess opposite-sign pairs as quantified by the variable r (see Eq. (4) and Sec. IV C). This r ratio (from Sec. V D for pion pairs) is also shown in Fig. 27. Neither of these baselines would yield the conclusion that a clear CME signal is observed in the analyses presented in Fig. 27.

The baselines for the CME-sensitive observables used in this blind analysis, as discussed above, are only general expectations. The observed multiplicity difference between the isobars requires future CME analyses to better understand the baselines in order to best utilize the precision demonstrated in this analysis.

VII. CONCLUSION

We report an experimental test of the Chiral Magnetic Effect (CME) by a blind analysis of a large statistics data set of isobar $^{96}\text{Ru}+^{96}\text{Ru}$ and $^{96}\text{Zr}+^{96}\text{Zr}$ collisions at nucleon-nucleon center-of-mass energy of 200 GeV, taken in 2018 by the STAR Collaboration at RHIC. The backgrounds are reduced using the difference in observables between the two isobar collision systems. The criteria for a positive CME observation are predefined, prior to the blind analysis, as a significant excess of the CME-sensitive observables in Ru+Ru collisions over those in Zr+Zr collisions. Consistent results are obtained by the five independent groups in this blind analysis. Significant differences in the multiplicity and flow harmonics are observed between the two systems in a given centrality, indicating that the magnitude of the CME background is different between the two species. A precision down to 0.4% is achieved in the relative magnitudes of pertinent observables between the two isobar systems. No CME signature that satisfies the predefined criteria has been observed in isobar collisions in this blind analysis.

VIII. ACKNOWLEDGEMENT

We thank the RHIC Operations Group and RCF at BNL, the NERSC Center at LBNL, and the Open Science Grid consortium for providing resources and support. We are grateful to Oak Ridge National Laboratory for providing the Ru-96, enriched in a special run, and RIKEN, Japan, for providing critical technology for the Zr-96 beam source. This work was supported in part by the Office of Nuclear Physics within the U.S. DOE Office of Science, the U.S. National Science Foundation, the Ministry of Education and Science of the Russian Federation, National Natural Science Foundation of China, Chinese Academy of Science, the Ministry of Science and Technology of China and the Chinese Ministry of Education, the Higher Education Sprout Project by Ministry of Education at NCKU, the National Research Foundation of Korea, Czech Science Foundation and Ministry of Education, Youth and Sports of the Czech Republic, Hungarian National Research, Development and Innovation Office, New National Excellency Programme of the Hungarian Ministry of Human Capacities, Department of Atomic Energy and Department of Science and Technology of the Government of India, the National Science Centre of Poland, the Ministry of Science, Education and Sports of the Republic of Croatia, RosAtom of Russia and German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF), Helmholtz Association, Ministry of Education Culture, Sports, Science, and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS).

-
- [1] D. Kharzeev, R. Pisarski, and M. H. Tytgat, Phys. Rev.Lett. **81**, 512 (1998), arXiv:hep-ph/9804221 [hep-ph].
 - [2] D. Kharzeev and R. D. Pisarski, Phys. Rev. D **61**, 111901 (2000), arXiv:hep-ph/9906401 [hep-ph].
 - [3] P. D. Morley and I. A. Schmidt, Z. Phys. C **26**, 627 (1985).
 - [4] V. Skokov, A. Yu. Illarionov, and V. Toneev, Int. J. Mod. Phys. A **24**, 5925 (2009), arXiv:0907.1396 [nucl-th].
 - [5] A. Bzdak and V. Skokov, Phys. Lett. B **710**, 171 (2012), arXiv:1111.1949 [hep-ph].
 - [6] W.-T. Deng and X.-G. Huang, Phys. Rev. C **85**, 044907 (2012), arXiv:1201.5108 [nucl-th].
 - [7] J. Błoczyński, X.-G. Huang, X. Zhang, and J. Liao, Phys. Lett. B **718**, 1529 (2013), arXiv:1209.6594 [nucl-th].
 - [8] K. Tuchin, Adv. High Energy Phys. **2013**, 490495 (2013), arXiv:1301.0099 [hep-ph].
 - [9] J. Błoczyński, X.-G. Huang, X. Zhang, and J. Liao, Nucl. Phys. A **939**, 85 (2015), arXiv:1311.5451 [nucl-th].
 - [10] L. McLerran and V. Skokov, Nucl. Phys. A **929**, 184 (2014), arXiv:1305.0774 [hep-ph].
 - [11] Y. Sun, Y. Wang, Q. Li, and F. Wang, Phys. Rev. C **99**, 064607 (2019), arXiv:1905.12492 [nucl-th].
 - [12] K. Fukushima, D. E. Kharzeev, and H. J. Warringa, Phys. Rev. D **78**, 074033 (2008), arXiv:0808.3382 [hep-ph].
 - [13] D. E. Kharzeev, L. D. McLerran, and H. J. Warringa, Nucl. Phys. A **803**, 227 (2008), arXiv:0711.0950 [hep-ph].
 - [14] D. Kharzeev, Phys. Lett. B **633**, 260 (2006), arXiv:hep-ph/0406125 [hep-ph].

- 1237 [15] D. E. Kharzeev and J. Liao, *Nature Rev. Phys.* **3**, 55 (2021), arXiv:2102.06623 [hep-ph].
- 1238 [16] D. E. Kharzeev, J. Liao, S. A. Voloshin, and G. Wang, *Prog. Part. Nucl. Phys.* **88**, 1 (2016), arXiv:1511.04050 [hep-ph].
- 1239 [17] Q. Li, D. E. Kharzeev, C. Zhang, Y. Huang, I. Pletikosic, A. V. Fedorov, R. D. Zhong, J. A. Schneeloch, G. D. Gu, and
1240 T. Valla, *Nature Phys.* **12**, 550 (2016), arXiv:1412.6543 [cond-mat.str-el].
- 1241 [18] S. Kaushik, D. E. Kharzeev, and E. J. Philip, *Phys. Rev. B* **99**, 075150 (2019).
- 1242 [19] B. I. Abelev *et al.* (STAR), *Phys. Rev. Lett.* **103**, 251601 (2009), arXiv:0909.1739 [nucl-ex].
- 1243 [20] B. I. Abelev *et al.* (STAR), *Phys. Rev. C* **81**, 054908 (2010), arXiv:0909.1717 [nucl-ex].
- 1244 [21] B. Abelev *et al.* (ALICE), *Phys. Rev. Lett.* **110**, 012301 (2013), arXiv:1207.0900 [nucl-ex].
- 1245 [22] L. Adamczyk *et al.* (STAR), *Phys. Rev. C* **89**, 044908 (2014), arXiv:1303.0901 [nucl-ex].
- 1246 [23] L. Adamczyk *et al.* (STAR), *Phys. Rev. C* **88**, 064911 (2013), arXiv:1302.3802 [nucl-ex].
- 1247 [24] L. Adamczyk *et al.* (STAR), *Phys. Rev. Lett.* **113**, 052302 (2014), arXiv:1404.1433 [nucl-ex].
- 1248 [25] V. Khachatryan *et al.* (CMS), *Phys. Rev. Lett.* **118**, 122301 (2017), arXiv:1610.00263 [nucl-ex].
- 1249 [26] A. M. Sirunyan *et al.* (CMS), *Phys. Rev. C* **97**, 044912 (2018), arXiv:1708.01602 [nucl-ex].
- 1250 [27] S. Acharya *et al.* (ALICE), *Phys. Lett. B* **777**, 151 (2018), arXiv:1709.04723 [nucl-ex].
- 1251 [28] J. Adam *et al.* (STAR), *Phys. Lett. B* **798**, 134975 (2019), arXiv:1906.03373 [nucl-ex].
- 1252 [29] S. Acharya *et al.* (ALICE), *JHEP* **09**, 160 (2020), arXiv:2005.14640 [nucl-ex].
- 1253 [30] D. E. Kharzeev, *Prog. Part. Nucl. Phys.* **75**, 133 (2014), arXiv:1312.3348 [hep-ph].
- 1254 [31] X.-G. Huang, *Rept. Prog. Phys.* **79**, 076302 (2016), arXiv:1509.04073 [nucl-th].
- 1255 [32] J. Zhao, *Int. J. Mod. Phys. A* **33**, 1830010 (2018), arXiv:1805.02814 [nucl-ex].
- 1256 [33] J. Zhao, Z. Tu, and F. Wang, *Nucl. Phys. Rev.* **35**, 225 (2018), arXiv:1807.05083 [nucl-ex].
- 1257 [34] J. Zhao and F. Wang, *Prog. Part. Nucl. Phys.* **107**, 200 (2019), arXiv:1906.11413 [nucl-ex].
- 1258 [35] W. Li and G. Wang, *Ann. Rev. Nucl. Part. Sci.* **70**, 293 (2020), arXiv:2002.10397 [nucl-ex].
- 1259 [36] S. A. Voloshin, *Phys. Rev. C* **70**, 057901 (2004), arXiv:hep-ph/0406311 [hep-ph].
- 1260 [37] A. Bzdak, V. Koch, and J. Liao, *Phys. Rev. C* **83**, 014905 (2011), arXiv:1008.4919 [nucl-th].
- 1261 [38] S. Pratt, S. Schlichting, and S. Gavin, *Phys. Rev. C* **84**, 024909 (2011), arXiv:1011.6053 [nucl-th].
- 1262 [39] D. Kharzeev, A. Krasnitz, and R. Venugopalan, *Phys. Lett. B* **545**, 298 (2002), arXiv:hep-ph/0109253 [hep-ph].
- 1263 [40] B. Muller and A. Schafer, *Phys. Rev. C* **82**, 057902 (2010), arXiv:1009.1053 [hep-ph].
- 1264 [41] K. Liu, *Phys. Rev. C* **85**, 014909 (2012), arXiv:1109.4883 [nucl-th].
- 1265 [42] M. Mace, N. Mueller, S. Schlichting, and S. Sharma, *Phys. Rev. D* **95**, 036023 (2017), arXiv:1612.02477 [hep-lat].
- 1266 [43] M. Mace, S. Schlichting, and R. Venugopalan, *Phys. Rev. D* **93**, 074036 (2016), arXiv:1601.07342 [hep-ph].
- 1267 [44] T. Lappi and S. Schlichting, *Phys. Rev. D* **97**, 034034 (2018), arXiv:1708.08625 [hep-ph].
- 1268 [45] J. Liao, *Pramana* **84**, 901 (2015), arXiv:1401.2500 [hep-ph].
- 1269 [46] Y. Yin and J. Liao, *Phys. Lett. B* **756**, 42 (2016), arXiv:1504.06906 [nucl-th].
- 1270 [47] Y. Jiang, S. Shi, Y. Yin, and J. Liao, *Chin. Phys. C* **42**, 011001 (2018), arXiv:1611.04586 [nucl-th].
- 1271 [48] S. Shi, Y. Jiang, E. Lilleskov, and J. Liao, *Annals Phys.* **394**, 50 (2018), arXiv:1711.02496 [nucl-th].
- 1272 [49] F. Wang, *Phys. Rev. C* **81**, 064902 (2010), arXiv:0911.1482 [nucl-ex].
- 1273 [50] A. Bzdak, V. Koch, and J. Liao, *Phys. Rev. C* **81**, 031901 (2010), arXiv:0912.5050 [nucl-th].
- 1274 [51] S. Schlichting and S. Pratt, *Phys. Rev. C* **83**, 014913 (2011), arXiv:1009.4283 [nucl-th].
- 1275 [52] F. Wang and J. Zhao, *Phys. Rev. C* **95**, 051901 (2017), arXiv:1608.06610 [nucl-th].
- 1276 [53] A. Kovner, M. Lublinsky, and V. Skokov, *Phys. Rev. D* **96**, 096003 (2017), arXiv:1706.02330 [hep-ph].
- 1277 [54] B. Schenke, C. Shen, and P. Tribedy, *Phys. Rev. C* **99**, 044908 (2019), arXiv:1901.04378 [nucl-th].
- 1278 [55] J. Zhao, Y. Feng, H. Li, and F. Wang, *Phys. Rev. C* **101**, 034912 (2020), arXiv:1912.00299 [nucl-th].
- 1279 [56] R. Belmont and J. Nagle, *Phys. Rev. C* **96**, 024901 (2017), arXiv:1610.07964 [nucl-th].
- 1280 [57] D. Kharzeev, Z. Tu, A. Zhang, and W. Li, *Phys. Rev. C* **97**, 024905 (2018), arXiv:1712.02486 [nucl-th].
- 1281 [58] S. A. Voloshin, *Phys. Rev. Lett.* **105**, 172301 (2010), arXiv:1006.1020 [nucl-th].
- 1282 [59] J. Schukraft, A. Timmins, and S. A. Voloshin, *Phys. Lett. B* **719**, 394 (2013), arXiv:1208.4563 [nucl-ex].
- 1283 [60] S. Chatterjee and P. Tribedy, *Phys. Rev. C* **92**, 011902 (2015), arXiv:1412.5103 [nucl-th].
- 1284 [61] V. Koch, S. Schlichting, V. Skokov, P. Sorensen, J. Thomas, S. Voloshin, G. Wang, and H.-U. Yee,
1285 *Chin. Phys. C* **41**, 072001 (2017), arXiv:1608.00982 [nucl-th].
- 1286 [62] N. Magdy, S. Shi, J. Liao, N. Ajitanand, and R. A. Lacey, *Phys. Rev. C* **97**, 061901 (2018),
1287 arXiv:1710.01717 [physics.data-an].
- 1288 [63] J. Zhao, H. Li, and F. Wang, *Eur. Phys. J. C* **79**, 168 (2019), arXiv:1705.05410 [nucl-ex].
- 1289 [64] H.-j. Xu, J. Zhao, X. Wang, H. Li, Z.-W. Lin, C. Shen, and F. Wang, *Chin. Phys. C* **42**, 084103 (2018),
1290 arXiv:1710.07265 [nucl-th].
- 1291 [65] S. A. Voloshin, *Phys. Rev. C* **98**, 054911 (2018), arXiv:1805.05300 [nucl-ex].
- 1292 [66] F. Du, L. E. Finch, and J. Sandweiss, *Phys. Rev. C* **78**, 044908 (2008).
- 1293 [67] L. E. Finch and S. J. Murray, *Phys. Rev. C* **96**, 044911 (2017), arXiv:1801.06476 [hep-ph].
- 1294 [68] A. Tang, *Chin. Phys. C* **44**, 054101 (2020), arXiv:1903.04622 [nucl-ex].
- 1295 [69] J. Adam *et al.* (STAR), (2020), arXiv:2006.05035 [nucl-ex].
- 1296 [70] M. Abdallah *et al.* (STAR), (2021), arXiv:2106.09243 [nucl-ex].
- 1297 [71] Y. Feng, J. Zhao, H. Li, H.-j. Xu, and F. Wang, (2021), arXiv:2106.15595 [nucl-ex].
- 1298 [72] N. N. Ajitanand, R. A. Lacey, A. Taranenko, and J. M. Alexander, *Phys. Rev. C* **83**, 011901 (2011),
1299 arXiv:1009.5624 [nucl-ex].
- 1300 [73] N. Magdy, S. Shi, J. Liao, P. Liu, and R. A. Lacey, *Phys. Rev. C* **98**, 061902 (2018), arXiv:1803.02416 [nucl-ex].

- [74] P. Bozek, Phys. Rev. C **97**, 034907 (2018), arXiv:1711.02563 [nucl-th].
- [75] Y. Sun and C. M. Ko, Phys. Rev. C **98**, 014911 (2018), arXiv:1803.06043 [nucl-th].
- [76] Y. Feng, J. Zhao, and F. Wang, Phys. Rev. C **98**, 034904 (2018), arXiv:1803.02860 [nucl-th].
- [77] L. Huang, M.-W. Nie, and G.-L. Ma, Phys. Rev. C **101**, 024916 (2020), arXiv:1906.11631 [nucl-th].
- [78] S. Shi, H. Zhang, D. Hou, and J. Liao, Phys. Rev. Lett. **125**, 242301 (2020), arXiv:1910.14010 [nucl-th].
- [79] Y. Feng, J. Zhao, H.-j. Xu, and F. Wang, Phys. Rev. C **103**, 034912 (2021), arXiv:2011.01123 [nucl-th].
- [80] N. Magdy, M.-W. Nie, L. Huang, G.-L. Ma, and R. A. Lacey, Phys. Lett. B **811**, 135986 (2020), arXiv:2003.02396 [nucl-ex].
- [81] S. Choudhury et al., (2021), arXiv:2105.06044 [nucl-ex].
- [82] STAR BUR 2018, STAR 2017 BUR https://drupal.star.bnl.gov/STAR/system/files/STAR_BUR_Run1718_v22_0.pdf.
- [83] W.-T. Deng, X.-G. Huang, G.-L. Ma, and G. Wang, Phys. Rev. C **94**, 041901 (2016), arXiv:1607.04697 [nucl-th].
- [84] W.-T. Deng, X.-G. Huang, G.-L. Ma, and G. Wang, Phys. Rev. C **97**, 044901 (2018), arXiv:1802.02292 [nucl-th].
- [85] H.-J. Xu, X. Wang, H. Li, J. Zhao, Z.-W. Lin, C. Shen, and F. Wang, Phys. Rev. Lett. **121**, 022301 (2018), arXiv:1710.03086 [nucl-th].
- [86] H. Li, H.-j. Xu, J. Zhao, Z.-W. Lin, H. Zhang, X. Wang, C. Shen, and F. Wang, Phys. Rev. C **98**, 054907 (2018), arXiv:1808.06711 [nucl-th].
- [87] G. Marr et al., in 10th International Particle Accelerator Conference (2019) pp. 28–32.
- [88] J. Adam et al. (STAR), Nucl. Sci. Tech. **32**, 48 (2021), arXiv:1911.00596 [nucl-ex].
- [89] P. Tribedy (STAR), Nucl. Phys. A **967**, 740 (2017), arXiv:1704.03845 [nucl-ex].
- [90] M. Anderson et al., Nucl. Instrum. Meth. A **499**, 659 (2003), arXiv:nucl-ex/0301015 [nucl-ex].
- [91] W. J. Llope et al., Nucl. Instrum. Meth. A **522**, 252 (2004), arXiv:nucl-ex/0308022 [nucl-ex].
- [92] J. Adams et al., Nucl. Instrum. Meth. A **968**, 163970 (2020), arXiv:1912.05243 [physics.ins-det].
- [93] C. Adler, A. Denisov, E. Garcia, M. J. Murray, H. Strobele, and S. N. White, Nucl. Instrum. Meth. **A470**, 488 (2001), arXiv:nucl-ex/0008005 [nucl-ex].
- [94] W. J. Llope et al., Nucl. Instrum. Meth. A **759**, 23 (2014), arXiv:1403.6855 [physics.ins-det].
- [95] V. Fine and P. Nevski, in 11th International Conference on Computing in High-Energy and Nuclear Physics (2000) pp. 143–146.
- [96] L. Adamczyk et al. (STAR), Phys. Rev. Lett. **116**, 112302 (2016), arXiv:1601.01999 [nucl-ex].
- [97] L. Adamczyk et al. (STAR), Phys. Rev. C **98**, 034918 (2018), arXiv:1701.06496 [nucl-ex].
- [98] L. Adamczyk et al. (STAR), Phys. Lett. B **790**, 81 (2019), arXiv:1701.06497 [nucl-ex].
- [99] A. M. Poskanzer and S. Voloshin, Phys. Rev. C **58**, 1671 (1998), arXiv:nucl-ex/9805001 [nucl-ex].
- [100] C. Adler, H. Strobele, A. Denisov, E. Garcia, M. Murray, and S. White, Nucl. Instrum. Meth. A **461**, 337 (2001).
- [101] SN0448, SN0448: Proposed Addition of a Shower Max Detector to the STAR Zero Degree Calorimeters <https://drupal.star.bnl.gov/STAR/starnotes/public/sn0448>.
- [102] J. Adams et al. (STAR), Phys. Rev. C **73**, 034903 (2006), arXiv:nucl-ex/0510053.
- [103] L. Adamczyk et al. (STAR), Phys. Rev. C **98**, 014915 (2017), arXiv:1712.01332 [nucl-ex].
- [104] NPP PAC report 2017, Recommendations of the Nuclear and Particle Physics Program Advisory Committee, Brookhaven National Laboratory, https://www.bnl.gov/npp/docs/pac0617/PACRecs2017_PAC.pdf.
- [105] P. Tribedy (STAR), J. Phys. Conf. Ser. **1602**, 1 (2020), arXiv:2009.01230 [nucl-ex].
- [106] R. Barlow, in Conference on Advanced Statistical Techniques in Particle Physics (2002) pp. 134–144, arXiv:hep-ex/0207026.
- [107] B. Efron, The Annals of Statistics **7**, 1 (1979).
- [108] B. I. Abelev et al. (STAR), Phys. Rev. C **79**, 034909 (2009), arXiv:0808.2041 [nucl-ex].
- [109] M. L. Miller, K. Reygers, S. J. Sanders, and P. Steinberg, Ann. Rev. Nucl. Part. Sci. **57**, 205 (2007), arXiv:nucl-ex/0701025 [nucl-ex].
- [110] P. A. Zyla et al. (Particle Data Group), PTEP **2020**, 083C01 (2020).
- [111] R. D. Woods and D. S. Saxon, Phys. Rev. **95**, 577 (1954).
- [112] J. Hammelmann, A. Soto-Ontoso, M. Alvioli, H. Elfner, and M. Strikman, Phys. Rev. C **101**, 061901 (2020), arXiv:1908.10231 [nucl-th].
- [113] H.-j. Xu, H. Li, X. Wang, C. Shen, and F. Wang, Phys. Lett. B **819**, 136453 (2021), arXiv:2103.05595 [nucl-th].
- [114] S. Raman, C. W. G. Nestor, Jr, and P. Tikkanen, Atom. Data Nucl. Data Tabl. **78**, 1 (2001).
- [115] B. Pritychenko, M. Birch, B. Singh, and M. Horoi, Atom. Data Nucl. Data Tabl. **107**, 1 (2016), [Erratum: Atom. Data Nucl. Data Tabl. 114, 371–374 (2017)], arXiv:1312.5975 [nucl-th].
- [116] P. Moller, J. R. Nix, W. D. Myers, and W. J. Swiatecki, Atom. Data Nucl. Data Tabl. **59**, 185 (1995), arXiv:nucl-th/9308022.
- [117] D. Kharzeev and M. Nardi, Phys. Lett. B **507**, 121 (2001), arXiv:nucl-th/0012025 [nucl-th].
- [118] S. Eremín and S. Voloshin, Phys. Rev. C **67**, 064905 (2003), arXiv:nucl-th/0302071.
- [119] B. Schenke, P. Tribedy, and R. Venugopalan, Phys. Rev. Lett. **108**, 252301 (2012), arXiv:1202.6646 [nucl-th].
- [120] J. S. Moreland, J. E. Bernhard, and S. A. Bass, Phys. Rev. C **92**, 011901 (2015), arXiv:1412.4708 [nucl-th].
- [121] S. Chatterjee, S. K. Singh, S. Ghosh, M. Hasanujjaman, J. Alam, and S. Sarkar, Phys. Lett. B **758**, 269 (2016), arXiv:1510.01311 [nucl-th].
- [122] H. Li, H.-j. Xu, Y. Zhou, X. Wang, J. Zhao, L.-W. Chen, and F. Wang, Phys. Rev. Lett. **125**, 222301 (2020), arXiv:1910.06170 [nucl-th].
- [123] C. Kremer et al., Phys. Rev. Lett. **117**, 172503 (2016), arXiv:1606.09057 [nucl-ex].

- 1365 [124] T. Togashi, Y. Tsunoda, T. Otsuka, and N. Shimizu, Phys. Rev. Lett. **117**, 172502 (2016), arXiv:1606.09056 [nucl-th].
1366 [125] S. A. Voloshin, A. M. Poskanzer, and R. Snellings, (2008), arXiv:0809.2949 [nucl-ex].
1367 [126] A. Bzdak, V. Koch, and J. Liao, Lect. Notes Phys. **871**, 503 (2013), arXiv:1207.7327 [nucl-th].
1368 [127] S. Choudhury, G. Wang, W. He, Y. Hu, and H. Z. Huang, Eur. Phys. J. C **80**, 383 (2020).
1369 [128] B. Schenke, C. Shen, and P. Tribedy, Phys. Lett. B **803**, 135322 (2020), arXiv:1908.06212 [nucl-th].
1370 [129] A. Dumitru, K. Dusling, F. Gelis, J. Jalilian-Marian, T. Lappi, and R. Venugopalan, Phys. Lett. B **697**, 21 (2011),
1371 arXiv:1009.5295 [hep-ph].
1372 [130] B. I. Abelev *et al.* (STAR), Phys. Rev. C **81**, 054908 (2010), arXiv:0909.1717 [nucl-ex].
1373 [131] G. Agakishiev *et al.* (STAR), Phys. Rev. C **86**, 064902 (2012), arXiv:1109.4380 [nucl-ex].
1374 [132] B. Alver, B. Back, M. Baker, M. Ballintijn, D. Barton, *et al.*, Phys. Rev. C **77**, 014906 (2008), arXiv:0711.3724 [nucl-ex].
1375 [133] S. A. Voloshin, A. M. Poskanzer, A. Tang, and G. Wang, Phys. Lett. B **659**, 537 (2008), arXiv:0708.0800 [nucl-th].
1376 [134] C. Adler *et al.* (STAR), Phys. Rev. C **66**, 034904 (2002), arXiv:nucl-ex/0206001.
1377 [135] P. Bozek, W. Broniowski, and J. Moreira, Phys. Rev. C **83**, 034911 (2011), arXiv:1011.3354 [nucl-th].
1378 [136] A. Bilandzic, R. Snellings, and S. Voloshin, Phys. Rev. C **83**, 044913 (2011), arXiv:1010.0233 [nucl-ex].
1379 [137] J. Jia, M. Zhou, and A. Trzupek, Phys. Rev. C **96**, 034906 (2017), arXiv:1701.03830 [nucl-th].
1380 [138] Y. Feng, Y. Lin, J. Zhao, and F. Wang, Phys. Lett. B **820**, 136549 (2021), arXiv:2103.10378 [nucl-ex].