

This is the accepted manuscript made available via CHORUS. The article has been published as:

First inverse kinematics measurement of resonances in Be^7

(α, n) reaction

relevant to neutrino-driven wind nucleosynthesis using
DRAGON

A. Psaltis, A. A. Chen, R. Longland, D. S. Connolly, C. R. Brune, B. Davids, J. Fallis, R. Giri, U. Greife, D. A. Hutcheon, L. Kroll, A. Lennarz, J. Liang, M. Lovely, M. Luo, C. Marshall, S. N. Paneru, A. Parikh, C. Ruiz, A. C. Shotter, and M. Williams

Phys. Rev. C **106**, 045805 — Published 14 October 2022

DOI: [10.1103/PhysRevC.106.045805](https://doi.org/10.1103/PhysRevC.106.045805)

First inverse kinematics measurement of resonances in ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ relevant to neutrino-driven wind nucleosynthesis using DRAGON

A. Psaltis,^{1,2,*} A.A. Chen,^{1,2} R. Longland,^{3,4} D.S. Connolly,^{5,†} C.R. Brune,⁶ B. Davids,^{5,7} J. Fallis,⁸ R. Giri,⁶ U. Greife,⁹ D.A. Hutcheon,⁵ L. Kroll,^{1,2} A. Lennarz,⁵ J. Liang,^{1,‡} M. Lovely,⁹ M. Luo,¹⁰ C. Marshall,^{3,4,§} S.N. Paneru,^{6,¶} A. Parikh,¹¹ C. Ruiz,^{5,12} A.C. Shotter,¹³ and M. Williams^{5,14}

¹*Department of Physics & Astronomy, McMaster University, Hamilton, Ontario L8S 4M1, Canada*

²*The NuGrid collaboration <https://nugrid.github.io/>*

³*Department of Physics, North Carolina State University, Raleigh, NC, 27695, USA*

⁴*Triangle Universities Nuclear Laboratory, Duke University, Durham, NC, 27710, USA*

⁵*TRIUMF, 4004 Wesbrook Mall, Vancouver, British Columbia V6T 2A3, Canada*

⁶*Department of Physics & Astronomy, Ohio University, Athens, Ohio 45701, USA*

⁷*Department of Physics, Simon Fraser University, Burnaby, British Columbia V5A 1S6, Canada*

⁸*North Island College, 2300 Ryan Rd, Courtenay, BC V9N 8N6, Canada*

⁹*Department of Physics, Colorado School of Mines, Golden, Colorado 80401, USA*

¹⁰*Department of Physics & Astronomy, University of British Columbia, Vancouver, British Columbia V6T 1Z4, Canada*

¹¹*Department de Física, Universitat Politècnica de Catalunya, E-08036 Barcelona, Spain*

¹²*Department of Physics & Astronomy, University of Victoria, Victoria, BC V8W 2Y2, Canada*

¹³*School of Physics, University of Edinburgh EH9 3JZ Edinburgh, United Kingdom*

¹⁴*Department of Physics, University of York, Heslington, York YO10 5DD, United Kingdom*

A possible mechanism to explain the origin of the light p -nuclei in the Galaxy is the nucleosynthesis in the proton-rich neutrino-driven wind ejecta of core-collapse supernovae via the νp -process. However this production scenario is very sensitive to the underlying supernova dynamics and the nuclear physics input. As far as the nuclear uncertainties are concerned, the breakout from the pp -chains via the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction has been identified as an important link which can influence the nuclear flow and therefore the efficiency of the νp -process. However its reaction rate is poorly known over the relevant temperature range, $T = 1.5\text{--}3$ GK. We report on the first direct measurement of two resonances of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction with previously unknown strengths using an intense radioactive ${}^7\text{Be}$ beam from the ISAC facility and the DRAGON recoil separator in inverse kinematics. We have decreased the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction rate uncertainty to $\approx 9.4\text{--}10.7\%$ over the relevant temperature region.

I. INTRODUCTION

The origin of the roughly 35 neutron-deficient stable isotopes with masses $A \geq 74$ — between ${}^{74}\text{Se}$ and ${}^{196}\text{Hg}$ — in the proton-rich side of the valley of stability, known as the “ p -nuclei” is a long-standing puzzle in nuclear astrophysics [1–3]. The p -nuclei were also traditionally referred to as “excluded” nuclei, since they were “shielded” by the s - and the r -process reaction paths [4]. For this reason their observed solar abundances [5], are 1–2 orders of magnitude smaller than their s - and r -process counterparts in the same mass region. It is generally accepted that the p -nuclei in the solar system have been produced by more than one process; however their synthesis mechanism is commonly referred to as “ p -process”.

The photodisintegration of pre-existing neutron-rich seeds, which is one of the most promising nucleosynthesis scenarios of p -nuclei synthesis and is thought to take place in the oxygen–neon layer of core-collapse supernovae (ccSNe), cannot reproduce the solar abundances of the light ${}^{92,94}\text{Mo}$ and ${}^{96,98}\text{Ru}$ isotopes, as well as the rare species ${}^{113}\text{In}$, ${}^{115}\text{Sn}$ and ${}^{138}\text{La}$. Additional astrophysical sites/nucleosynthesis scenarios have been proposed, such as the thermonuclear explosions of Chandrasekhar mass carbon–oxygen white dwarfs (CO WD) [6], which is also supported by Galactic Chemical Evolution (GCE) models and the rp -process in Type I X-ray bursts [7]. It is remarkable that despite the variety of astrophysical models, all these processes can reproduce the solar abundances of most of the p -nuclei to within a factor of 3 [*e.g.* see the sensitivity studies in References 8, 9].

* psaltisa@mcmaster.ca; Present address: Institut für Kernphysik, Technische Universität Darmstadt, 64289 Darmstadt, Germany

† Present address: Los Alamos National Laboratory, Los Alamos, NM 87545, USA

‡ Present address: TRIUMF, 4004 Wesbrook Mall, Vancouver, British Columbia V6T 2A3, Canada

§ Present address: Department of Physics & Astronomy, Ohio University, Athens, Ohio 45701, USA

¶ Facility for Rare Isotope Beams, East Lansing, Michigan 48824, USA

The advancement of multi-dimensional core-collapse supernova simulations with sophisticated neutrino transport methods, see References [10, 11] for the first studies that discussed this and [12, 13] for some more recent results, suggests that the composition of the early innermost ejecta of the neutrino-driven wind that drives the explosion are mostly proton-rich (the electron fraction

Y_e is greater than 0.5)¹ [15, 16], and that gives rise to a new nucleosynthesis scenario, the νp -process [17–19], which can produce the lighter of the p -nuclei.

To summarize the νp -process, the neutrino-driven wind ejects very hot ($T > 10$ GK) and proton-rich material from the proton-neutron star (PNS) (see Figure 1). At these extreme temperatures, the ejecta consist mainly of nucleons from dissociated nuclei. As the wind expands and cools down, nuclear statistical equilibrium (NSE) assembles these nucleons into mainly ^{56}Ni and α particles (which are synthesised via the hot pp -chain sequence [20]) with an excess of free protons. At $T < 3$ –4 GK, ^{56}Ni can rapidly capture free protons. However, the reaction flow cannot move beyond ^{64}Ge , which has a relatively long β^+ half-life of 1.06 min. This issue is resolved by electron antineutrino captures on free protons via the $p(\bar{\nu}_e, e^+)n$ reaction, which produce a small amount of free neutrons, 10^{-11} – 10^{-12} of the total mass. At temperature drops from 3 to 1.5 GK, the much faster (n, p) reaction on ^{56}Ni , followed by a sequence of radiative proton captures, *i.e.* (p, γ) reactions, and further (n, p) reactions bypass ^{64}Ge and similar waiting-points, such as ^{68}Se and ^{72}Kr with half-lives of 35.5 s and 17.1 s, respectively. The reaction flow follows the $Z = N$ line up to the molybdenum region and then moves into more neutron-rich isotopes ($Z < N$) between molybdenum and tin. Finally, as the temperature drops below $T < 1.5$ GK, (p, γ) reactions freeze-out due to the Coulomb barrier, and the produced nuclei decay back to stability, with ^{56}Ni still being the most abundant nucleus in the plasma.




FIG. 1. Simplified schematic of the nucleosynthesis in neutrino-driven wind ejecta. The different stages and outcomes (νp -process and weak r -process) are shown. The figure is adapted from José and Iliadis [21].

¹ A relatively neutron-rich neutrino-driven wind ($0.4 < Y_e < 0.5$), leads to a different nucleosynthesis scenario called the weak r -process, which can produce the lighter heavy elements with $Z = 26$ –47 [14].

The uncertainties of νp -process nucleosynthesis, mainly attributed to the supernova dynamics and the underlying nuclear physics input, have been explored by many groups [14, 22–26] since it was first proposed. The most crucial component for a successful νp -process is the electron fraction Y_e of the ejecta. Recent hydrodynamical studies with proper neutrino transport have shown that Y_e can lie between 0.5 and 0.6 before the onset of νp -processing at $T = 3$ GK. Sensitivity studies have explored a variety of Y_e values, ranging from 0.5 to 0.8, and suggest that a higher Y_e leads to a more efficient νp -process (*i.e.* production of heavier nuclei).

Concerning the nuclear physics input of the νp -process the main uncertainties arise from a handful of reactions, and the nuclear masses along the reaction path. The two most important reactions that dominate the nucleosynthesis in this scenario are the bottleneck $^{56}\text{Ni}(n, p)^{56}\text{Co}$ and triple- α – $^4\text{He}(\alpha\alpha, \gamma)^{12}\text{C}$ – reactions. The former is always the first step of the νp -process and as a result controls the reaction flow, with a smaller rate yielding a more efficient nucleosynthesis, since the free neutrons synthesized from neutrino captures are captured by nuclei with $30 \leq Z \leq 42$, acting as “neutron poisons”.

The triple- α reaction controls the production of α -particles, protons and the ^{56}Ni seed before the onset and during the νp -process. Therefore it controls completely the neutron-to-seed ratio Δ_n , as defined by Pruet *et al.* [19]. Our current knowledge of this reaction, despite its importance, is still limited and bears large experimental uncertainties. The three rates that are most commonly used in nucleosynthesis studies are those from References [27–29]. In addition, Jin *et al.* [30] recently showed that an enhanced triple- α reaction, due to an in-medium width change of the Hoyle state, suppresses the production of p -nuclei in the νp -process.

In the sensitivity study of Wanajo *et al.* [22], some alternative pathways were explored. In particular, the authors found that there are a couple of two-body reaction sequences, namely $^7\text{Be}(\alpha, \gamma)^{11}\text{C}(\alpha, p)^{14}\text{N}$ and $^7\text{Be}(\alpha, p)^{10}\text{B}(\alpha, p)^{13}\text{C}$, which compete with the triple- α reaction, the main link between the pp -chain ($A < 12$) and CNO ($A \geq 20$) region, at the relevant temperature region $T = 1.5$ –3 GK. This competition affects the Δ_n factor and as a result, the reaction flow and the final elemental abundances. The authors studied the sensitivity of the final abundances by multiplying and dividing the $^7\text{Be}(\alpha, \gamma)^{11}\text{C}$ reaction rate by factors of 2 and 10. This rate variation affected the production of light p -nuclei with $90 < A < 110$ up to an order of magnitude. A faster $^7\text{Be}(\alpha, \gamma)^{11}\text{C}$ rate leads to increased production of intermediate-mass nuclei that remove protons from the environment, acting as “proton poisons”. Subsequent studies, such as that by Nishimura *et al.* [26], also acknowledge the importance of the $^7\text{Be}(\alpha, \gamma)^{11}\text{C}$ reaction, but do not provide a quantitative impact in the production of p -nuclei. As we shall discuss in detail in Section II, the $^7\text{Be}(\alpha, \gamma)^{11}\text{C}$ reaction rate is not well known in the relevant temperature range due to unknown res-

onance strengths, and thus an experimental study is required.

In the present work we report on the first inverse kinematics study of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction, using the DRAGON recoil separator and an intense ${}^7\text{Be}$ radioactive ion beam from ISAC. The paper is structured as follows: in Section II we discuss the previous measurements regarding resonances of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction. In Sections III and IV we present the experimental details of the present work along with the analysis procedures, and finally we discuss our results and conclusions in Sections V and VI.

II. PREVIOUS MEASUREMENTS

Our current understanding of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction over the energy region relevant to νp -process nucleosynthesis is based on three experimental studies [31–33].

Figure 2 shows the current level structure of ${}^{11}\text{C}$ along with its mirror ${}^{11}\text{B}$, and in Table I we summarize the resonance parameters for the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction from the $A = 11$ evaluation of Kelley *et al.* [34].

The two lowest-lying energy resonances of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction, which correspond to the $E_x = 8.105$ and 8.420 MeV levels in ${}^{11}\text{C}$, were studied by Hardie *et al.* [31] in forward kinematics at Argonne National Laboratory. The authors used two methods to calculate the resonance strengths: the first was the thick target yield formula (similar to Equation 4), and the second was a complementary relative method which employed the presence of ${}^7\text{Li}$ in the target, and the fact that they were studying the ${}^7\text{Li}(\alpha, \gamma){}^{11}\text{B}$ reaction in the same campaign. More specifically, the relative method provided the resonance strength ratio between the resonances of interest in ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ and the known 660 keV ($E_x = 9.272$ MeV in ${}^{11}\text{B}$) resonance of the ${}^7\text{Li}(\alpha, \gamma){}^{11}\text{B}$ reaction, reported in the same work. The main advantage of this method is that both the ${}^7\text{Li} : {}^7\text{Be}$ ratio in the target and the detector efficiencies are more accurately known than the number of ${}^7\text{Be}$ atoms alone and the absolute efficiencies. Nevertheless, one has to include an extra uncertainty factor from the ${}^7\text{Li}(\alpha, \gamma){}^{11}\text{B}$ resonance. The adopted values for the two resonance strengths in Table I are the weighted averages of the two methods.

Wiescher *et al.* [32] studied the $E_x = 8.654$ and 8.699 MeV levels in ${}^{11}\text{C}$, which correspond to the 1110 and 1155 keV² resonances of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction. They used the ${}^{10}\text{B}(p, \gamma){}^{11}\text{C}$ reaction in forward kinematics employing three different linear accelerators, covering a wide energy range ($E_x = 8 - 10.7$ MeV). In all three experimental setups, several detectors were used, allowing for angular distribution measurements. The authors observed primary γ transitions from the $E_x = 8.654$ and 8.699 MeV

states and calculated the ratio Γ_γ/Γ for them using the cross sections from the γ ray and α -particle channels, $\sigma(p, \gamma)/\sigma(p, \alpha)$.




FIG. 2. Partial level scheme of the mirror nuclei ${}^{11}\text{C}$ and ${}^{11}\text{B}$ adopted from Kelley *et al.* [34], with the addition of the $E_x = 8.900$ MeV state from Yamaguchi *et al.* [33]. The dashed lines indicate isobaric analog states, and next to the ${}^{11}\text{C}$ scheme we present the α separation energy Q_α , the resonances of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction, E_r , in keV and the relevant energy region for νp -process nucleosynthesis.

The most recent study relevant to the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction was performed by Yamaguchi *et al.* [33] using the low-energy radioactive ion beam facility CRIB [35] at CNS in RIKEN, Japan. The ${}^7\text{Be} + \alpha$ resonant scattering and ${}^7\text{Be}(\alpha, p)$ reaction measurements were performed using the thick-target method in inverse kinematics and provided the excitation functions for $E_x = 8.7\text{--}13.0$ MeV. The R -matrix analysis of the data shows two small peaks in the low energy region, between 8.90 and 9.20 MeV. The first one is considered to be the known $5/2^+$ state at 9.20 MeV observed by Wiescher *et al.* [32]. The second one, located at around 8.90 MeV, is regarded by the authors as a new resonance. However, they argue that this spectral feature could also originate from either the

² All resonance energies are expressed in the center of mass system.

TABLE I. Resonance parameters for the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction from Kelley *et al.* [34]. The parameters for the resonance noted with a \diamond are adopted from Yamaguchi *et al.* [33]. Tentative assignments and estimates are presented in parentheses. The resonances noted with a \dagger were studied in the present work. Values noted with a \parallel were adopted from the mirror nucleus ${}^{11}\text{B}$. The proton partial widths Γ_p have been calculated using $C^2S = 1$.

E_x (MeV)	E_r (keV)	J^π	Γ_α	Γ_γ	Γ_p	ℓ	$\omega\gamma$ (eV)
7.4997(15)	-43.9(15)	$3/2^+$	2.2(1.6) eV	1.15 eV $^\parallel$	-	1	...
8.1045(17)	560.5(17)	$3/2^-$	6_{-6}^{+12} eV	0.350(56) eV	-	0	0.331(41)
8.420(2) \dagger	876(2)	$5/2^+$	12.6(38) eV	3.1(13) eV	-	2	3.80(57)
8.654(4) \dagger	1110(4)	$7/2^+$	≤ 5 keV	-	-	3	...
8.699(2) \dagger	1155(2)	$5/2^+$	15(1) keV	1.15(16) eV $^\parallel$	-	1	...
8.900 $^\diamond$	1356	$(9/2^+)$	>8 keV	-	-	(3)	(1.2)
9.645(50)	2101(50)	$(3/2^-)$	210(40) keV	17 eV $^\parallel$	48(9) eV	0	...
9.780(50)	2236(50)	$(5/2^-)$	240(50) keV	1 eV $^\parallel$	520(100) eV	2	...
9.970(50)	2426(50)	$(7/2^-)$	120(20) keV	1 eV $^\parallel$	760(140) eV	2	...
10.083(5)	2539(5)	$9/2^+$	≈ 230 keV	<0.2 eV $^\parallel$	900(180) eV	3	...

$E_x = 8.655$ or the 8.699 MeV states since their the energy uncertainty was quite large in this energy region. Finally, all this information about the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ resonances is summarised in Table I.

The current ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction rate is based on a calculation from NACRE (I and II) [28, 36] and includes contributions only from the 561 and 876 keV resonances, for which experimentally measured strengths exist, and the non-resonant (DC) contribution is adopted with the same parameters as those of the mirror ${}^7\text{Li}(\alpha, \gamma){}^{11}\text{B}$ reaction, for $T < 0.7$ GK. Contributions from the broad resonances at 2101, 2236, 2426 & 2539 keV were also included in NACRE-II and affect the reaction rate for $T > 2$ GK. In NACRE-I the same high energy part of the reaction rate was estimated using Hauser-Feshbach calculations. The rate that was used in the sensitivity study by Wanaajo *et al.* [22] was the one from Angulo *et al.* [28] (NACRE-I), which is uncertain by factors of 1.87–2.54 in the relevant temperature region. The NACRE-II reaction rate is uncertain by factors of 1.76–1.91 for $T = 1.5$ –3 GK. The uncertainties are derived from their Potential Model (PM), which is used to reproduce the experimental astrophysical S -factor data. $S(E) \equiv (E/e^{-2\pi\eta}) \sigma(E)$, where E is the center of mass energy, and η is the Sommerfeld parameter, which is related to the charges and velocities of the interacting particles. More specifically, the uncertainties are calculated by using the maximum and minimum parameters of the PM. It is also worth noting that the sub-threshold resonance at $E_x = 7.50$ MeV ($E_r = -43.9$ keV) has a large contribution at low temperatures, below $T \approx 0.3$ GK and according to Descouvemont [37], it could affect the production of ${}^7\text{Li}$ (fed by the decay of ${}^7\text{Be}$) in classical novae [38].

III. EXPERIMENTAL DETAILS

The measurements of this work were carried out using the DRAGON recoil separator [39] at TRIUMF, Canada's particle accelerator centre in Vancouver, BC.

Intense beams of ${}^7\text{Be}^+$ were produced using the ISOL technique, by bombarding thick ZrC and graphite targets with 55 μA 500 MeV protons from the TRIUMF cyclotron. The $A = 7$ isobars, mainly ${}^7\text{Li}$ and ${}^7\text{Be}$, were extracted from the target through a high-resolution mass separator, and the beryllium ionization was enhanced using the TRIUMF Resonant Ionization Laser Ion Source (TRILIS) [40]. After the ion source, the beam was transported through the ISAC high resolution ($M/\Delta M = 2000$) mass separator and then accelerated through the ISAC-I Radio-Frequency Quadrupole (RFQ) and Drift-Tube Linac (DTL) to energies so that each resonance was centered in the gas target (see Table II for details). The beam energies were chosen in order to cover center-of-mass windows of 1157 ± 24 keV, 1111 ± 13 keV, and 878 ± 17 keV, across the gas target volume. To ensure a pure, contaminant-free radioactive ion beam, an additional carbon stripping foil of 20 $\mu\text{g}/\text{cm}^2$ was placed downstream of the DTL allowing fully stripped ${}^7\text{Be}^{4+}$ to be selected using a bending magnet for transport to DRAGON, thus eliminating the main isobaric contaminant ${}^7\text{Li}$. This technique has also been used in other radioactive beam facilities [41]. Finally, ${}^7\text{Be}^{4+}$ was delivered at the helium-filled windowless gas target of DRAGON at mean intensities of $\approx 1.3 - 5.8 \times 10^8$ pps (see Section IV C for normalisation details of the intensities).

TABLE II. Beam and gas target properties for the two independent measurements of the present study^a.

	E_{beam} (A keV)	E_{lab} (MeV)	I_{beam} $\times 10^8$ (s^{-1})	P_{target} (Torr)
Run 1	464.2(3)	3.249(2)	1.33(7)	7.9(1)
Run 1	442.7(2)	3.099(1)	2.06(8)	5.06(6)
Run 2	441.8(2)	3.093(1)	5.83(2)	4.89(3)
Run 2	351.8(3)	2.463(2)	3.45(12)	5.75(4)

^a The 1110 keV resonance was studied in two independent measurements, due to a low recoil yield in the first measurement.

DRAGON has four main components: (a) the windowless, differentially pumped, recirculating gas target, (b) the γ ray detector array, (c) the electromagnetic mass separator and (d) the recoil detection system, which are shown in the schematic of Figure 3.

The γ ray array consists of 30 BGO scintillator crystals with photo-multiplier tubes (PMTs) covering 89 – 92% of the 4π solid angle [42]. The segmented array allows for the detection of individual prompt γ rays from the radiative capture reactions inside the gas target and the tagging of the associated recoil particles, which provides an additional background reduction in the focal plane detectors.

The DRAGON electromagnetic mass separator consists of two magnetic (M) and two electric dipoles (E) in a MEME configuration. The two-stage separation begins with the first magnetic dipole (MD1), which selects a single charge stage to be transmitted through DRAGON. For our study, we tuned DRAGON to the $q = 2^+$ charge state for all resonances. Recoils that do not have the aforementioned charge are deflected to slits that are located downstream of the MD1. Subsequently, the recoils are led to the first electric dipole (ED1) where they are separated according to mass. ED1 is followed by the second stage of magnetic and electric dipoles (MD2 and ED2), until the beam reaches the focal plane, where the heavy ion detectors are located.




FIG. 3. Schematic representation of the DRAGON recoil separator. The main components are shown.

Close to the focal plane of DRAGON we employed a microchannel plate (MCP) and a double-sided silicon-strip detector (DSSSD). The MCP provided the starting timing signal for a local time-of-flight (TOF) transmission measurement [43]. In the DSSSD, the recoils are stopped, their kinetic energy and position are measured and the stopping timing signal is recorded. In addition, we employed the prompt γ rays from the BGO array as a starting signal for a “separator TOF” measurement for the coincidence analysis (see Section IV B). The data were recorded using a state-of-the-art time-

stamp-based data acquisition system (DAQ) [44].

It is worth mentioning that such a measurement using DRAGON, and any other recoil separator dedicated to resonance strength measurements, is quite challenging, due to geometric considerations. The maximum recoil angle of the reaction at the resonance energies of interest can be calculated using the following equation,

$$\theta_{r,max} = \arctan \left(\frac{E + Q}{\sqrt{2m_1c^2 \left(\frac{m_1+m_2}{m_2} \right) E}} \right) \quad (1)$$

where E is the center of mass energy, Q is the reaction Q value and m_1, m_2 the masses of the projectile and the target nuclei, respectively. This corresponds to a single γ ray emission to the recoil nucleus ground state at 90° in the center-of-mass system.

For the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction maximum recoil angles are $\theta_{r,max} = 42.67$ mrad for the 1155 keV, 43.3 mrad for the 1110 keV and 47.42 mrad for the 876 keV. These numbers far exceed the nominal angular acceptance of DRAGON, ($\theta_{\text{DRAGON}} = \pm 21$ mrad). For this reason, we performed detailed GEANT simulations of DRAGON [45, 46] to extract the transmission of the recoils through the separator ($\eta_{\text{separator}}$), and in addition, the BGO array efficiency (η_{BGO}), which are used for the resonance strength calculations. References [42, 47, 48] provide an in-depth discussion about this approach, and in Section IV F we provide the specifics for the study of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction.

IV. DATA ANALYSIS

We performed yield measurements for three beam energies, corresponding to the 1155, 1110 and 876 keV resonances of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction (see Figure 2). As we have already discussed in Section II, the 876 keV resonance strength has been measured by Hardie *et al.* [31], while the latter two resonances have unknown strengths. Our reasoning to re-measure that resonance is two-fold: on the one hand it is believed to have the greatest impact on the current reaction rate at νp -process energies [36] and on the other hand it will provide one additional demonstration that DRAGON can measure resonance strengths for reactions in which the angular spread of the recoils exceeds its nominal acceptance [47, 48].

A. Thick target yield and resonance strength

The calculation of thermonuclear reaction rates in a laboratory setting requires the determination of the reaction cross section. Instead, what is actually measured in experimental studies is the reaction yield, which can be simply expressed as:

$$Y = \frac{\mathcal{N}_{\mathcal{R}}}{\mathcal{N}_{\mathcal{B}}} \quad (2)$$

where $\mathcal{N}_{\mathcal{R}}$ is the number of reactions that occur and $\mathcal{N}_{\mathcal{B}}$ is the number of incident beam particles. In fact, an experimental setup has a finite detection efficiency, in our case η_{DRAGON} , meaning that it does not detect the total number of reactions, but rather a fraction of it, N_r . According to the analysis mode that we use, singles or coincidences (see Section IV B), η_{DRAGON} can be either $\eta_{DRAGON}^{singles} = \eta_{separator} f_q \eta_{MCP} \eta_{DSSSD} \eta_{live}^{singles}$, or $\eta_{DRAGON}^{coinc} = \eta_{separator} f_q \eta_{BGO} \eta_{MCP} \eta_{DSSSD} \eta_{live}^{coinc}$, respectively. The experimental yield is then given by:

$$Y = \frac{\mathcal{N}_r}{\mathcal{N}_{\mathcal{B}} \eta_{DRAGON}} \quad (3)$$

We can also express the energy-dependent reaction yield as a relation between the cross section $\sigma(E)$ and the target thickness ΔE , or better, the stopping power of the target $\varepsilon(E)$, for beam energy E_{beam} using:

$$Y(E_{beam}) = \int_{E-\Delta E}^E \frac{\sigma(E)}{\varepsilon(E)} dE \quad (4)$$

For narrow resonances with constant stopping power over the resonance width, which can be found in reactions relevant for astrophysics, we can calculate the integral of Equation 4 analytically using a single-level Breit-Wigner (Lorentzian) cross section profile [49]. Specifically, in the case of an infinitely thick target, that is $\Delta E \rightarrow \infty$, or equivalently $\Delta E \gg \Gamma$, we have:

$$Y(E_{beam}) = \frac{\lambda_r^2 \omega \gamma}{2\pi \varepsilon_r} \left[\tan^{-1} \left(\frac{E_{beam} - E_r}{\Gamma/2} \right) + \frac{\pi}{2} \right] \quad (5)$$

where λ_r and ε_r are the de Broglie wavelength and the target stopping power in the center of mass system, E_r is the energy of the resonance, Γ is its width and ΔE is the target thickness. Solving for $E_{beam} = E_r$, we can obtain a simple expression for the reaction yield and the resonance strength $\omega\gamma$:

$$\omega\gamma = \frac{2Y_{\Delta E \rightarrow \infty} \varepsilon}{\lambda_r^2} \frac{m_1}{m_1 + m_2} \quad (6)$$

where the reaction yield $Y_{\Delta E \rightarrow \infty}$ is given by Equation 3 and ε is the target stopping power in the laboratory frame – a discussion on how it is measured in DRAGON experiments can be found in Section IV E.

B. Particle Identification

The first step towards determining the reaction yield and subsequently the strength of a resonance is the identification of the reaction products or recoils. For this,

we employed two distinct methods: a detection of ^{11}C recoils in singles, using the DSSSD and a local TOF (MCP-DSSSD), and in γ -recoil coincidences using the separator TOF (BGO-DSSSD). Figure 4 shows typical particle identification plots for the three resonances of the $^7\text{Be}(\alpha, \gamma)^{11}\text{C}$ reaction both in singles (grey points) and in coincidences (coloured points), with additional software cuts, such as the energy range of the γ rays in the BGO array and the energy deposited in the DSSSD, providing further recoil discrimination. It is evident that for the 876 and 1110 keV resonances the yield is low, but the signal is clear and without any unwanted background, such as unreacted, “leaky” beam. This is consistent with the fact that DRAGON is able to reject unreacted beam particles from (α, γ) reactions very efficiently and has demonstrated a beam suppression of $> 10^{13}$ [50]. In the present experiment, the rejection is higher, due to the use of a fully-stripped beam and the fact that the selected carbon recoils have a very different charge state (4^+ vs 2^+).

In addition to the aforementioned methods, we could also identify the recoils of interest using a timing signal of the 11.8 MHz ISAC-I radio frequency quadrupole (RF) accelerator and the capture of a coincidence γ ray by the BGO array (BGO-RF) [39, 44]. Figure 5 shows the results for each resonance.

C. Beam Normalization

We monitored the ^7Be beam current throughout the experiment using silicon surface barrier detectors (SSB) at well defined laboratory angles of 30° and 57° with respect to the beam axis by detecting the elastically scattered target particles. Due to the low count rate in the SSB detectors, we did not use SSB measurements for short time windows Δt , before and after each yield measurement to calculate the beam normalization \mathcal{R} factor, as is typical in DRAGON experiments (see for example the works in References [39, 51–53]). Instead, we first ensured that the beam current during each run was stable by checking the current on the charge slits after the first magnetic dipole and used the total integrated counts in the SSBs per yield run to calculate the \mathcal{R} factor, which is given by:

$$\mathcal{R} = \frac{I}{|q \cdot e|} \frac{\Delta t}{N_\alpha} \frac{P}{E_b^2} \eta_{target} \quad (7)$$

where I is the average current reading at the upstream Faraday Cup before the gas target, q is the beam charge state (4^+), e is the elementary charge ($e = 1.6 \times 10^{-19}$ C), N_α is the number of scattered α particles detected by the surface barrier detectors during the yield run time Δt , P is the gas target pressure in Torr, E_b is the beam energy in keV/u and η_{target} is the transmission through an empty target. We assume only elastic Rutherford scattering for the target particles and the E_b^2/P factor enters




FIG. 4. Particle identification plot for the ^{11}C recoils for each of the resonances we studied in the present work using the local Time-Of-Flight transmission measurement and the energy deposited in the DSSSD at the focal plane of DRAGON. The coloured and grey points correspond to coincident and singles recoils events, respectively. For the 1110 keV, we show the two independent measurements in separate panels. See the text for details.

Equation 7 to make \mathcal{R} invariant to the chosen beam energy and target pressure [54].

The normalized number of beam particles N_{beam} , is then given by:

$$N_{beam} = \mathcal{R} N_{\alpha} \frac{E_b^2}{P} \quad (8)$$

Table III shows the \mathcal{R} factor results for all the yield measurements of the present work. Note that in our two independent experimental runs we used different SSB gains, threshold settings, and pre-scalers. For this reason, the N_{α} that we use in Equations 7 and 8 to extract N_{beam} are also different.




FIG. 5. Particle identification plot using the BGO versus accelerator RF Time-of-Flight for coincident recoil events. The RF period is $\tau_{RF} = 84.8$ ns.

D. Carbon in helium charge state distribution

DRAGON is tuned to select and transport a single charge state to the final focal plane. For this reason, an accurate knowledge of the recoil charge state distribution (CSD) is necessary to determine the total reaction yield. Since the recoil nucleus, in our case ^{11}C , is unstable, an abundant and stable isotope of the same element is used instead — ^{12}C . The stable ion beam for this measurement was provided from the microwave ion source (MWIS) of the ISAC Off-Line Ion Source (OLIS) [55].

At DRAGON, the charge state distributions can be

TABLE III. Beam normalization results for the yield measurements of the present work^a.

	E_{beam} (A keV)	\mathcal{R} -factor (${}^7\text{Be}/\alpha$)(Torr/keV ²) $\times 10^{13}$ ions	N_{beam}
Run 1	464.2(3)	$1.15(2) \times 10^{11}$	1.07(2)
Run 1	442.6(2)	$1.22(2) \times 10^{11}$	1.76(5)
Run 2	441.8(2)	$1.74(4) \times 10^{10}$	1.53(4)
Run 2	351.8(3)	$2.77(6) \times 10^{10}$	2.12(4)

^a During the two independent experimental runs, we used different SSB gains, threshold settings, and pre-scalers. For this reason the absolute value of the SSB rate is not comparable between these different periods.

determined experimentally by measuring the beam current on Faraday cups before and after the gas target, and comparing it to the current on a Faraday cup downstream from the first magnetic dipole (see Figure 3). We chose to tune DRAGON to the 2⁺ charge state because according to theoretical calculations [56], it is the maximum of the distribution, thus providing the highest recoil yield. Figure 6 shows the results for this charge charge for energies corresponding to the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ resonance strengths.




FIG. 6. Experimentally measured carbon in helium Charge State Distribution. The fit to the experimental data is a Gaussian function, and the band gives the 1 σ confidence interval to the fit.

E. ${}^7\text{Be}$ stopping power in ${}^4\text{He}$

The stopping power ϵ of the gas target is an important component for the calculation of the reaction yield. The advantage of recoil separators, such as DRAGON, is that the stopping power is measured directly and it is not based on semi-empirical estimates that can introduce an additional uncertainty factor in the final result. At DRAGON, the stopping power is measured by varying both the pressure of the gas target and the magnetic field strength needed to center the beam at a momentum dispersed angular focus after the first magnetic dipole. Our

experimental results agree to within 6% with calculations using the SRIM code [57], as shown in Table IV.

TABLE IV. Summary of the ${}^7\text{Be}$ in ${}^4\text{He}$ stopping power measurements. The experimental results are compared with the calculations of SRIM [57]. The units of ϵ are eV/ (10^{15} atoms/cm²).

E_{beam} (A keV)	ϵ_{DRAGON}	ϵ_{SRIM}
464.2(3)	40.7(15)	38.2
442.6(2)	39.7(15)	38.8
441.9(2)	39.5(15)	38.8
351.8(3)	41.5(18)	39.6

F. GEANT simulations of DRAGON

As we have already pointed out in the above, detailed simulations using GEANT are needed to determine the recoil transmission $\eta_{\text{separator}}$ and the efficiency of the BGO array η_{BGO} , which are used to calculate the reaction yield, and subsequently the resonance strength $\omega\gamma$ as part of the recoil detection efficiency of DRAGON, η_{DRAGON} .

The DRAGON GEANT simulation toolkit³ has been extensively used for experimental planning, such as in the study of the ${}^{12}\text{C}(\alpha, \gamma){}^{16}\text{O}$ reaction [58], and its results show agreement with experimental data to within 1–10% [46].

The simulation input file includes all the information GEANT requires to perform the simulation such as the energy, spin, lifetime, and γ branching ratios for each nuclear level and in addition the energy and width of the resonance of interest (see Table V for an overview). For the study of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction, the nuclear information was adopted from the A=11 evaluation of Kelley *et al.* [34]. Specifically for the 1110 keV resonance, since there are no experimentally measured γ branching ratios, we adopted those of the mirror state in ${}^{11}\text{B}$. For the γ ray angular distribution $W(\theta)$, which affects both the transmission of the recoils and the BGO array efficiency [42, 47], we calculated all the possible $W(\theta)$ for each cascade, following the prescription of Rose and Brink [59]. In addition, we changed the gas target pressure in the simulation, in order to obtain the same stopping power as in the experiment (see Section IV E).

We performed simulations for each resonance energy within its uncertainty, ± 2 , ± 4 and ± 2 keV for the 1155, 1110 and 876 keV resonance respectively. The final results used in the data analysis are the averages of these simulations and the systematic uncertainty is attributed mainly to the uncertainty in the γ branching ratios and the range of possible γ angular distributions.

³ The GEANT simulation package of DRAGON can be found at https://github.com/DRAGON-Collaboration/G3_DRAGON.

TABLE V. Settings of the GEANT3 simulation for the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ data analysis. Nuclear properties were adopted from Kelley *et al.* [34]. For the 1110 keV resonance, the branching ratios of the mirror nucleus ${}^{11}\text{B}$ are used.

Quantity	$E_r = 876$ keV			$E_r = 1110$ keV			$E_r = 1155$ keV		
Excited state lifetime	0.030 fs			1.31×10^{-19} s			4.3×10^{-20} s		
Resonance energy (keV)	874-878			1106-1114 keV			1153-1157 keV		
Particle (α) partial width	12.6 eV			5 keV			15 keV		
	E_x^i (MeV)	E_x^f (MeV)	B.R.	E_x^i (MeV)	E_x^f (MeV)	B.R.	E_x^i (MeV)	E_x^f (MeV)	B.R.
γ branching ratios	8.420	0	93	8.654	0	0.9 ± 0.3	8.699	0	42 ± 10
		4.319	7		4.319	86.6 ± 2.3		4.319	42 ± 10
					6.478	12.5 ± 1.1		4.804	2.4 ± 1.5
							6.478	13.6 ± 4.6	

Figure 7 shows the results from a simulation of the $E_r = 1155$ keV resonance. It is evident that transitions with a cascade of multiple γ rays, such as the 8.699 MeV \rightarrow 4.32 MeV provide more favourable conditions for transmission through the separator, since their recoil angular distribution from multiple decay vectors averages out with resulting maximum intensity at lower angles, as Ruiz *et al.* [42] argue.




FIG. 7. Angular distribution for recoils of the $E_r = 1155$ keV resonance that hit the focal plane detector (DSSSD) using GEANT. The contributions from different cascades are shown. The vertical dashed line defines the angular acceptance of DRAGON, $\theta_{\text{DRAGON}} = 21$ mrad.

V. RESULTS

In the following, we shall discuss the results from each resonance studied in the present work and then present the uncertainties and the calculation of the new thermonuclear reaction rate $N_A \langle \sigma v \rangle$.

A. Strength of the 1155 keV resonance

For the highest energy resonance we studied in the present work, 1155 keV, we detected a strong, background-free signal, as is evident in the PID plot of

Figure 4. Clusters of 33 and 49 recoil events in coincidence and singles modes were detected, despite the low recoil transmission through the separator ($\eta_{\text{separator}} = 0.141(28)$, see also Table VIII).

The GEANT simulations we performed for this resonance are in very good agreement with the experimental results, as Figure 8 shows (see also Figure 2 of Ref. [60] for the same resonance). It is worth noting that DRAGON is more sensitive to the detection of recoils that γ decay to the $E_x = 4.319$ MeV state, compared to the ground state, as we discussed in Section IV F.




FIG. 8. Comparison between the experimental data (points) and the GEANT simulation (histograms) for a BGO γ_0 ray energy spectrum from the 1155 keV resonance. The two dominant γ transitions to the $E_x = 4.319$ MeV and the ground state can be clearly seen in the simulation results. We do not depict the weak transition to the $E_x = 4.804$ MeV state. See the text for details.

The final result for the resonance strength is $\omega\gamma_{1155} = 1.73 \pm 0.25(\text{stat.}) \pm 0.40(\text{syst.})$ eV was adopted from the singles analysis and had a smaller uncertainty compared to the coincidence analysis result. However, the two results are in agreement (see also Table VI).

B. Strength of the 1110 keV resonance

The 1110 keV resonance was studied in two independent experimental runs, due to the low recoil yields. We detected $14_{-3.7}^{+4.3}$ and $9_{-2.7}^{+3.8}$ events in singles, with different integrated beam fluxes $1.76(5) \times 10^{13}$ and $1.53(5) \times 10^{13}$ ions, respectively (see also Table III). The asymmetric uncertainty in the amount of detected recoils was calculated according to the prescription of Feldman and Cousins [61] for a poissonian signal with zero background, as we can see from the PID plots of Figure 4.

To account for the asymmetric uncertainties and provide a realistic statistical uncertainty for the number of detected ^{11}C recoils, we proceeded as follows: we first used Fechner’s two-piece normal distribution [62] for the two independent runs, using the Feldman and Cousins [61] prescription for the variances (see Figure 9 - Top). After that, we created a combined probability distribution by calculating averages by randomly sampling from the two individual distributions. The final results for the detected ^{11}C recoils and their respected 1 and 2σ uncertainties are then calculated from the combined distribution. We find $12.1_{-2.5}^{+2.7}$ (1σ) and $_{-4.8}^{+5.3}$ (2σ) events for the 1110 keV resonance in singles, corresponding to $_{-20.7}^{+22.3}\%$ (1σ) and $_{-39.7}^{+43.8}\%$ (2σ) statistical uncertainty, respectively as we show in Figure 9.

The resonance strengths resulting from singles and coincidence analysis are $\omega\gamma_{1110,\text{singles}} = 125_{-25}^{+27}(\text{stat.}) \pm 15(\text{syst.})$ MeV and $\omega\gamma_{1110,\text{coinc}} = 161_{-41}^{+43}(\text{stat.}) \pm 24(\text{syst.})$ MeV, respectively, from which we choose the former as the final result. The large difference compared to the 1155 keV resonance strength – almost an order of magnitude – can be attributed to the difference in the orbital angular momentum, $\ell_\alpha = 1$ for the former state and $\ell_\alpha = 2$ for the latter (see Table I).

C. Strength of the 876 keV resonance

For the lowest energy in this study, the maximum recoil angle is $\theta_{r,\text{max}} = 47$ mrad, which is the largest ever attempted by DRAGON⁴. Nevertheless, one can see a clear signal in the PID plots (see Figures 4 and 5). We detected $13_{-3.7}^{+4.3}$ ^{11}C recoil events corresponding to $_{-28.5}^{+33.1}\%$ 1σ statistical uncertainty, following the Feldman and Cousins [61] prescription for a poissonian signal with zero background.

Our final result for its strength from the singles analysis is $\omega\gamma_{876} = 3.00_{-0.72}^{+0.81}(\text{stat.}) \pm 0.61(\text{syst.})$ eV. We calculated the weighted average of our measurement and the value by Hardie *et al.* [31] to get the adopted resonance strength $\omega\gamma_{876} = 3.61(50)$ eV, which will be used for the calculation of the thermonuclear reaction rate in




FIG. 9. (Top) Individual probability distributions for the detected ^{11}C recoils in singles mode from the two independent measurements of the 1110 keV resonance. (Bottom) Combined probability distribution from the same measurements. The solid line shows the central value, while the dashed and dotted lines show the 1 and 2σ uncertainties, respectively. See the text for details.

Section V E. It is worth noting that for all resonances the results for the strength $\omega\gamma$ agree both in coincidence and singles analysis modes.

Also, from a nuclear structure standpoint, our results for the two previously unknown resonance strengths are in very good agreement with their $^7\text{Li}(\alpha,\gamma)^{11}\text{B}$ analogs [34], namely $E_x = 9.182$ ($7/2^+$) and 9.271 MeV ($5/2^+$), as they are shown in Table VII. The $5/2^-$ state $E_x = 8.921$ in ^{11}B is the only exception, were its resonance strength differs more than two orders of magnitude to its ^{11}C analog.

D. Uncertainties

The uncertainties of the final results of the resonance strengths of this study are of systematic and statistical nature. The former are dominated by the efficiencies of the BGO array (η_{BGO}) and the recoil transmission through the separator ($\eta_{\text{separator}}$), which are inferred from GEANT simulations (Section IV F) [42]. Other sources of systematic uncertainty are the MCP detec-

⁴ The previous largest maximum recoil angle was $\theta_{r,\text{max}} = 33$ mrad in the study of the $^{12}\text{C}(^{16}\text{O},\gamma)^{28}\text{Si}$ reaction [63].

TABLE VI. Resonance strengths $\omega\gamma$ of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction resonances from the literature [31, 34] and the present work (in singles and coincidences modes) that were used for the calculation of the new thermonuclear reaction rate. All results are reported in eV and the statistical and systematic uncertainties are presented separately. The adopted value for the 876 keV resonance strength is the weighted average of our singles measurement and the one from Hardie *et al.* [31]. See the text for details.

E_r (keV)	Literature	Singles	Coincidences	Adopted
561	0.331(41)	0.331(41)
876	3.80(57)	$3.00^{+0.81}_{-0.72}(\text{stat.}) \pm 0.61(\text{syst.})$	$3.91^{+1.29}_{-1.10}(\text{stat.}) \pm 1.18(\text{syst.})$	3.61(50)
1110	...	$0.125^{+0.027}_{-0.025}(\text{stat.}) \pm 0.015(\text{syst.})$	$0.161^{+0.043}_{-0.041}(\text{stat.}) \pm 0.024(\text{syst.})$	0.125(31)
1155	...	$1.73 \pm 0.25(\text{stat.}) \pm 0.40(\text{syst.})$	$1.79 \pm 0.33(\text{stat.}) \pm 0.42(\text{syst.})$	1.73(47)

TABLE VII. Comparison of resonance strengths $\omega\gamma$ for analog states in ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ and ${}^7\text{Li}(\alpha, \gamma){}^{11}\text{B}$ reactions. Literature data were taken from Kelley *et al.* [34].

J^π	Nucleus	E_x (MeV)	$\omega\gamma$ (eV)
$5/2^-$	${}^{11}\text{B}$	8.921(1)	$(8.8 \pm 1.4) \times 10^{-3}$
$5/2^-$	${}^{11}\text{C}$	8.420(2)	3.61(50)
$7/2^+$	${}^{11}\text{B}$	9.182(2)	0.303(26)
$7/2^+$	${}^{11}\text{C}$	8.654(4)	0.125(31)
$5/2^+$	${}^{11}\text{B}$	9.271(2)	1.72(24)
$5/2^+$	${}^{11}\text{C}$	8.699(2)	1.73(47)

tion efficiency, the stopping power measurements and the charge state fractions.

Furthermore, the statistical uncertainties are due to the low recoil detection yield, caused by the very low transmission of the recoils through the separator, but this parameter is well understood and quantified. As we already pointed out, for the 1110 and 876 keV resonances, we used the prescription of Feldman and Cousins [61] for poissonian signals in zero background to extract the statistical uncertainties.

Table VIII shows a detailed breakdown of the uncertainties for each of the three resonances we measured in the present study. Note that for the 1110 keV resonance the uncertainty of the average final result was calculated using the procedure discussed in Section VB.

E. Thermonuclear Reaction Rate

The new ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ thermonuclear reaction rate was calculated using the RatesMC⁵ code [64]. Within the RatesMC framework, each nuclear physics input quantity (*e.g.* resonance energy and resonance strength) has an assigned probability density function (PDF). The code samples these functions randomly many times ($> 10^3$) using a Monte Carlo algorithm and outputs reaction rates and associated rate probability densities. According to the central limit theorem, a random variable that is de-

termined by the product of many factors will be distributed according to a lognormal density function [64–66]. Using a lognormal PDF, the “low”, “recommended”, and “high” Monte Carlo rates are the 16th, 50th (median), and 84th percentile respectively of the cumulative reaction rate distribution. In Table IX we present the adopted thermonuclear reaction rate for ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$.

For our calculation, we used resonance parameters as reported in Kelley *et al.* [34]. More specifically, we included the contribution of the sub-threshold resonance at $E_x = 7.4997$ MeV ($E_r = -43.9$ keV) using a γ partial width from the mirror state in ${}^{11}\text{B}$ ($\Gamma_\gamma = 1.14(4)$ eV), and assumed a reduced α width of 1. According to Descouvemont [37] this resonance can dominate the reaction rate for $T < 0.3$ GK, which can affect the evolution of Population III stars via the hot pp -chains [20] and the production of ${}^7\text{Li}$ in classical novae [38]. In addition to the narrow resonances at 561, 876, 1110 & 1155 keV, we also included contributions from the broad resonances at 2101, 2236, 2426 & 2539 keV (see Table I for details). For the γ partial widths of the latter states, since we used values from the mirror analog ${}^{11}\text{B}$, we assigned them a factor of 2 uncertainty.

The new reaction rate uncertainty has been decreased to $\approx 9.4 - 10.7\%$ over $T = 1.5 - 3$ GK, the relevant temperature window for νp -process nucleosynthesis, compared to factors of 1.76-1.91 of the NACRE-II compilation. While our new rate includes the new measurements of the 1110 and 1155 keV resonance strengths and the updated adopted value for $\omega\gamma_{876}$, it is worth noting that this decrease in the thermonuclear reaction rate is mainly caused from using a Monte Carlo error propagation of the relevant quantities (*e.g.* E_r , $\omega\gamma$ *etc.*) [64], and not by the individual contribution of the previously unmeasured resonance strengths. In Figure 10 we compare the new thermonuclear reaction rate to the NACRE rate [28, 36] and an older reaction rate compilation from Caughlan and Fowler [27] (CF88).

Figure 11 shows the individual resonant contributions to the total ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction rate. For temperatures $T \lesssim 0.2$ GK, the sub-threshold resonance at -43.9 keV dominates the reaction rate, while for $0.2\text{GK} < T < 1.0\text{GK}$ the 561 keV contributes the most, since $\Gamma_\alpha \gg \Gamma_\gamma$ (see Table I) and it is the lowest-lying energy resonance [see the discussion in Ref. 67, Chapter 3]. For the temperatures relevant for the νp -process,

⁵ The RatesMC code to calculate thermonuclear reaction rates can be found at <https://github.com/rlongland/RatesMC>.

TABLE VIII. Values given with uncertainties for the quantities used to calculate the resonance strengths in singles and coincidences modes. The relative errors are quoted in parentheses.

Source	1155 keV	1110 keV (Run 1)	1110 keV (Run 2)	876 keV
Detected recoils, $N_{\text{rec}}^{\text{singles}}$	49(7) (14.3%)	$14^{+4.3}_{-3.7}$ (+31%)	$9^{+3.8}_{-2.7}$ (+42%)	$16^{+4.3}_{-3.8}$ (+27%)
Detected recoils, $N_{\text{rec}}^{\text{coinc}}$	33(6) (18.2%)	$9^{+3.8}_{-2.7}$ (+42%)	$7^{+3.3}_{-2.8}$ (+47%)	$13^{+4.3}_{-3.7}$ (+33%)
Charge state fraction, f_q	0.40(1) (2.5%)	0.41(1) (2.4%)	0.41(1) (2.4%)	0.41(1) (2.4%)
Beam particles, $N_{\text{beam}} \times 10^{13}$	1.07(2) (1.9%)	1.76(5) (2.8%)	1.53(5) (3.2%)	2.12(3) (1.41%)
BGO efficiency, η_{BGO}	0.77(1) (1.3%)	0.81(7) (8.6%)	0.81(7) (8.6%)	0.80(18) (22.5%)
Separator transmission, $\eta_{\text{separator}}$	0.141(28) (19.9%)	0.266(18) (6.8%)	0.266(18) (6.8%)	0.016(3) (18.8%)
MCP efficiency ^a , η_{MCP}	0.545(59) (10.8%)	0.650(61) (9.4%)	0.321(25) (7.8%)	0.351(19) (5.4%)
Live time, $\eta_{\text{live}}^{\text{singles}}$	0.95409(5) (0.005%)	0.95777(5) (0.005%)	0.99099(5) (0.005%)	0.93408(5) (0.005%)
Live time, $\eta_{\text{live}}^{\text{coinc}}$	0.80381(4) (0.005%)	0.80434(4) (0.005%)	0.82156(4) (0.005%)	0.81571 (0.005%)
Stopping power, ϵ (eV/(10^{15} /cm ²))	40.7(15) (3.7%)	39.7(15) (3.8%)	39.5(15) (3.8%)	41.5(18) (4.3%)
Beam energy (A keV)	462.2(3) (0.06%)	442.6(2) (0.05%)	441.8(2) (0.05%)	351.8(3) (0.09%)
Total uncertainty in singles	14.3% (stat.)	$^{+31}_{-26}$ % (stat.)	$^{+42}_{-30}$ % (stat.)	$^{+27}_{-24}$ % (stat.)
(statistical & systematic)	23.2% (syst.)	12.8% (syst.)	11.8% (syst.)	20.2% (syst.)
Total uncertainty in coincidences	18.2% (stat.)	$^{+42}_{-30}$ % (stat.)	$^{+47}_{-40}$ % (stat.)	$^{+33}_{-28}$ % (stat.)
(statistical & systematic)	23.2% (syst.)	15.4% (syst.)	14.6% (syst.)	30.3% (syst.)

^a The MCP efficiency includes both the detection efficiency of the system and also the transmission of the recoils through the thin carbon foil that creates the secondary electrons that the MCP detects.




FIG. 10. The new ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction rate for $T=0.1\text{--}10$ GK compared to the rates by References [27, 28, 36] over the same temperature region.

the 876 keV has a $\approx 60\%$ contribution to the total rate, followed by the 561 keV with $\approx 30\%$. The 1155 keV resonance has a $\approx 10\%$ contribution, while the 1110 keV contributes negligibly to the total reaction rate.




FIG. 11. Resonant contributions to the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ thermonuclear reaction rate. The dotted line at the bottom right corner shows the contribution of the 1110 keV resonance.

The effect of the new reaction rate will be studied extensively in a future publication, taking into account new measurements of the ${}^{10}\text{B}(\alpha, p){}^{13}\text{C}$ reaction [68], which was also included in the sensitivity study of Wanajo *et al.* [22], and the ${}^{59}\text{Cu}(p, \alpha){}^{56}\text{Ni}$ [69] which may be responsible for the Ni–Cu cycle [23].

In addition to the thermonuclear reaction rate, we also calculated the astrophysical S -factor. In Figure 12 we present the astrophysical S -factor for ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ with the individual resonant contributions. Our results agree well with the NACRE-II data, with the exception of the 2101 keV resonance which seems to be mis-placed

TABLE IX. Total thermonuclear reaction rates for ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$. The rate below $T < 0.012$ GK is zero. The rates are expressed in units of $\text{cm}^3 \text{mol}^{-1} \text{s}^{-1}$. Columns 2,3 and 4 list the 16th, 50th and 86th percentiles of the total rate probability density (PDF) at given temperatures. “*f.u.*” is the factor uncertainty, and is obtained from the 16th and 84th percentiles.

T (GK)	Low	Median	High	f.u.
0.012	3.463×10^{-34}	8.835×10^{-34}	2.348×10^{-33}	2.590
0.013	4.535×10^{-33}	1.153×10^{-32}	3.066×10^{-32}	2.588
0.014	4.594×10^{-32}	1.168×10^{-31}	3.097×10^{-31}	2.586
0.015	3.758×10^{-31}	9.552×10^{-31}	2.528×10^{-30}	2.584
0.016	2.563×10^{-30}	6.511×10^{-30}	1.725×10^{-29}	2.583
0.018	7.618×10^{-29}	1.935×10^{-28}	5.129×10^{-28}	2.579
0.020	1.407×10^{-27}	3.572×10^{-27}	9.463×10^{-27}	2.576
0.025	4.793×10^{-25}	1.208×10^{-24}	3.195×10^{-24}	2.568
0.030	4.012×10^{-23}	1.008×10^{-22}	2.665×10^{-22}	2.560
0.040	2.466×10^{-20}	6.147×10^{-20}	1.616×10^{-19}	2.544
0.050	2.309×10^{-18}	5.693×10^{-18}	1.495×10^{-17}	2.528
0.060	7.225×10^{-17}	1.766×10^{-16}	4.628×10^{-16}	2.513
0.070	1.117×10^{-15}	2.705×10^{-15}	7.081×10^{-15}	2.498
0.080	1.057×10^{-14}	2.548×10^{-14}	6.647×10^{-14}	2.483
0.090	6.993×10^{-14}	1.681×10^{-13}	4.372×10^{-13}	2.468
0.100	3.570×10^{-13}	8.501×10^{-13}	2.201×10^{-12}	2.453
0.110	1.472×10^{-12}	3.490×10^{-12}	8.984×10^{-12}	2.439
0.120	5.172×10^{-12}	1.216×10^{-11}	3.106×10^{-11}	2.424
0.130	1.580×10^{-11}	3.679×10^{-11}	9.412×10^{-11}	2.410
0.140	4.322×10^{-11}	9.999×10^{-11}	2.548×10^{-10}	2.396
0.150	1.078×10^{-10}	2.473×10^{-10}	6.275×10^{-10}	2.382
0.160	2.482×10^{-10}	5.656×10^{-10}	1.429×10^{-9}	2.367
0.180	1.116×10^{-9}	2.459×10^{-9}	6.115×10^{-9}	2.310
0.200	4.918×10^{-9}	9.604×10^{-9}	2.208×10^{-8}	2.096
0.250	5.348×10^{-7}	6.305×10^{-7}	7.894×10^{-7}	1.251
0.300	2.666×10^{-5}	3.051×10^{-5}	3.506×10^{-5}	1.147
0.350	4.586×10^{-4}	5.215×10^{-4}	5.955×10^{-4}	1.142
0.400	3.813×10^{-3}	4.334×10^{-3}	4.923×10^{-3}	1.139
0.450	1.950×10^{-2}	2.209×10^{-2}	2.507×10^{-2}	1.137
0.500	7.106×10^{-2}	8.022×10^{-2}	9.077×10^{-2}	1.135
0.600	4.816×10^{-1}	5.413×10^{-1}	6.105×10^{-1}	1.130
0.700	1.860×10^0	2.083×10^0	2.340×10^0	1.125
0.800	5.136×10^0	5.726×10^0	6.390×10^0	1.118
0.900	1.139×10^1	1.265×10^1	1.401×10^1	1.111
1.000	2.178×10^1	2.407×10^1	2.651×10^1	1.105
1.250	7.217×10^1	7.907×10^1	8.638×10^1	1.095
1.500	1.633×10^2	1.792×10^2	1.953×10^2	1.094
1.750	2.946×10^2	3.236×10^2	3.538×10^2	1.095
2.000	4.578×10^2	5.041×10^2	5.522×10^2	1.098
2.500	8.379×10^2	9.262×10^2	1.021×10^3	1.103
3.000	1.235×10^3	1.370×10^3	1.516×10^3	1.107
3.500	1.612×10^3	1.801×10^3	2.003×10^3	1.116
4.000	1.955×10^3	2.195×10^3	2.468×10^3	1.129
5.000	2.515×10^3	2.876×10^3	3.332×10^3	1.167
6.000	2.924×10^3	3.398×10^3	4.121×10^3	1.210
7.000	3.202×10^3	3.812×10^3	4.777×10^3	1.250
8.000	3.386×10^3	4.112×10^3	5.325×10^3	1.284
9.000	3.503×10^3	4.324×10^3	5.765×10^3	1.312
10.000	3.558×10^3	4.460×10^3	6.079×10^3	1.335




FIG. 12. The astrophysical S -factor for the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction based on our RatesMC calculations. Contributions from different resonances are shown. The narrow resonances at 561, 876 & 1110 keV are shown with the vertical dotted lines. The astrophysical S -factor from NACRE-II [36] is also shown.

to lower energies [see Figure 42 in Ref. 36].

VI. DISCUSSION AND CONCLUSIONS

In the present work we performed the first inverse kinematics study of the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction to measure unknown resonance strengths at energies relevant to νp -process nucleosynthesis. We report the first measurement of the 1155 & 1110 keV resonances with strengths of $1.73 \pm 0.25(\text{stat.}) \pm 0.40(\text{syst.})$ eV and $125_{-25}^{+27}(\text{stat.}) \pm 15(\text{syst.})$ MeV, respectively. We also re-measured the 876 keV resonance strength ($\omega\gamma_{876} = 3.00_{-0.72}^{+0.81}(\text{stat.}) \pm 0.61(\text{syst.})$ eV) and our result agrees with the measurement of Hardie *et al.* [31] ($\omega\gamma = 3.80(57)$ eV).

As we have also demonstrated in Psaltis *et al.* [47], the present work shows that DRAGON is capable of handling measurements in which the maximum recoil cone angle exceeds its acceptance, after a systematic study of the BGO array efficiency and its transmission using extensive GEANT simulations. That opens a new avenue for future experiments using DRAGON, that were previously thought to be inaccessible due to large maximum recoil angles (*e.g.* ${}^{18}\text{O}(\alpha, \gamma){}^{22}\text{Ne}$, ${}^{20}\text{Ne}(\alpha, \gamma){}^{24}\text{Mg}$ and others).

The new ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction rate is constrained to 9.4-10.7 % for $T = 1.5$ -3 GK which is sufficient for nucleosynthesis calculations. The effect of the rate, along with other measured reactions relevant to nucleosynthesis in neutrino-driven winds will be explored in a subsequent study. According to the work of Wanajo *et al.* [22], the ${}^7\text{Be}(\alpha, \gamma){}^{11}\text{C}$ reaction rate in the relevant energies can affect the number of the neutron-to-seed ratio Δ_n , regulating the νp -process efficiency in synthesizing neutron-deficient isotopes. This is a particularly interesting result, since most recent self-consistent 3D core-collapse

supernova simulations favour proton-rich conditions [70]. In addition, a rigorous study of the astrophysical conditions of the proton-rich neutrino driven ejecta, and how they produce different nucleosynthesis outputs, using all the up-to-date nuclear physics input is desired.

The intense ${}^7\text{Be}$ RIBs produced with carbide targets can be utilized for more demanding experiments, such as ${}^7\text{Be}(p, \gamma)$ and ${}^7\text{Be}$ α -scattering. Pure graphite targets bombarded by protons at $100 \mu\text{A}$ (or a UC_x target at $40 \mu\text{A}$) could produce as much as 10^{10} s^{-1} of ${}^7\text{Be}$.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the beam delivery and ISAC operations groups at TRIUMF. In particular, we thank F. Ames, T. Angus, A. Gottberg, S. Kiy, J. Lassen and O. Shelbaya for all their help during the experiment. The authors thank the anonymous referee for useful comments that improved the manuscript. The core operations of TRIUMF are supported via a contribution from the federal government through the National Research Council of Canada, and the Govern-

ment of British Columbia provides building capital funds. Authors from McMaster University are funded by the National Sciences and Engineering Research Council of Canada (NSERC). DRAGON is funded through NSERC grant SAPPJ-2019-00039. AP also acknowledges support from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)-Project No. 279384907-SFB 1245, and the State of Hesse within the Research Cluster ELEMENTS (Project ID 500/10.006). RL and CM acknowledge support from the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under Grant Nos. DE-SC0017799 and DE-FG02-97ER41042. CRB, RG and SP acknowledge support from the U.S. Department of Energy, under grants number DE-FG02-88ER40387 and DE-NA0003883. Authors from Colorado School of Mines acknowledge support from U.S. Department of Energy Office of Science DE-FG02-93ER40789 grant. Authors from the UK are supported by the Science and Technology Facilities Council (STFC). This work benefited from discussions at the “Nuclear Astrophysics at Rings and Recoil Separators” Workshop supported by the National Science Foundation under Grant No. PHY-1430152 (JINA Center for the Evolution of the Elements).

-
- [1] M. Arnould and S. Goriely, *Phys. Rep.* **384**, 1 (2003).
 - [2] T. Rauscher *et al.*, *Rep. Prog. Phys.* **76**, 066201 (2013).
 - [3] M. Pignatari *et al.*, *Int. J. Mod. Phys. E* **25**, 1630003 (2016).
 - [4] A. G. W. Cameron, *Publ. Astron. Soc. Pac* **69**, 201 (1957).
 - [5] K. Lodders *et al.*, in *Solar system* (Springer, 2009) pp. 712–770.
 - [6] C. Travaglio *et al.*, *Astrophys. J* **795**, 141 (2014).
 - [7] H. Schatz *et al.*, *Phys. Rep.* **294**, 167 (1998).
 - [8] W. Rapp *et al.*, *Astrophys. J* **653**, 474 (2006).
 - [9] T. Rauscher *et al.*, *Mon. Not. R. Astron. Soc.* **463**, 4153 (2016).
 - [10] L. Hüdepohl *et al.*, *Phys. Rev. Lett.* **104**, 251101 (2010).
 - [11] T. Fischer *et al.*, *Astron. Astrophys.* **517**, A80 (2010).
 - [12] O. Just *et al.*, *Mon. Not. R. Astron. Soc.* **453**, 3386 (2015).
 - [13] E. O’Connor, *Astrophys. J Supplement Series* **219**, 24 (2015).
 - [14] A. Arcones and F. Montes, *Astrophys. J* **731**, 5 (2011).
 - [15] S. Wanajo *et al.*, *Astrophys. J* **852**, 40 (2018).
 - [16] D. Vartanyan *et al.*, *Mon. Not. R. Astron. Soc.* **482**, 351 (2019).
 - [17] C. Fröhlich *et al.*, *Phys. Rev. Lett* **96**, 142502 (2006).
 - [18] S. Wanajo, *Astrophys. J* **647**, 1323 (2006).
 - [19] J. Pruet *et al.*, *Astrophys. J* **644**, 1028 (2006).
 - [20] M. Wiescher *et al.*, *Astrophys. J* **343**, 352 (1989).
 - [21] J. José and C. Iliadis, *Rep. Prog. Phys.* **74**, 096901 (2011).
 - [22] S. Wanajo *et al.*, *Astrophys. J* **729**, 46 (2011).
 - [23] A. Arcones, C. Fröhlich, and G. Martínez-Pinedo, *Astrophys. J* **750**, 18 (2012).
 - [24] C. Fröhlich and T. Rauscher, in *AIP Conference Proceedings*, Vol. 1484 (American Institute of Physics, 2012) pp. 232–239.
 - [25] M. Jacobi, *Influence of astrophysical and nuclear physics uncertainties on the nucleosynthesis in proton-rich neutrino-driven winds*, Master’s thesis, Technische Universität Darmstadt (2018).
 - [26] N. Nishimura *et al.*, *Mon. Not. R. Astron. Soc.* **489**, 1379 (2019).
 - [27] G. R. Caughlan and W. A. Fowler, *At. Data Nucl. Data Tables* **40**, 283 (1988).
 - [28] C. Angulo *et al.*, *Nucl. Phys. A* **656**, 3 (1999).
 - [29] H. O. Fynbo *et al.*, *Nature* **433**, 136 (2005).
 - [30] S. Jin *et al.*, *Nature* **588**, 57 (2020).
 - [31] G. Hardie *et al.*, *Phys. Rev. C* **29**, 1199 (1984).
 - [32] M. Wiescher *et al.*, *Phys. Rev. C* **28**, 1431 (1983).
 - [33] H. Yamaguchi *et al.*, *Phys. Rev. C* **87**, 034303 (2013).
 - [34] J. Kelley *et al.*, *Nucl. Phys. A* **880**, 88 (2012).
 - [35] Y. Yanagisawa *et al.*, *Nucl. Instrum. Methods Phys. Res A* **539**, 74 (2005).
 - [36] Y. Xu *et al.*, *Nucl. Phys. A* **918**, 61 (2013).
 - [37] P. Descouvemont, *Nucl. Phys. A* **584**, 532 (1995).
 - [38] M. Hernanz, J. José, A. Coc, and J. Isern, *Astrophys. J Lett.* **465**, L27 (1996).
 - [39] D. Hutcheon *et al.*, *Nucl. Instrum. Methods Phys. Res A* **498**, 190 (2003).
 - [40] J. Lassen *et al.*, *Hyperfine Interact.* **162**, 69 (2005).
 - [41] M. Gaelens *et al.*, *Nucl. Instrum. Methods Phys. Res B* **204**, 48 (2003).
 - [42] C. Ruiz *et al.*, *Eur. Phys. J. A* **50**, 99 (2014).
 - [43] C. Vockenhuber *et al.*, *Nucl. Instrum. Methods Phys. Res A* **603**, 372 (2009).
 - [44] G. Christian *et al.*, *Eur. Phys. J. A* **50**, 75 (2014).

- [45] D. G. Gigliotti *et al.*, *Nucl. Instrum. Methods Phys. Res B* **204**, 671 (2003).
- [46] D. G. Gigliotti, *Efficiency calibration measurement and GEANT simulation of the DRAGON BGO gamma ray array at TRIUMF*, *Master's thesis*, University of Northern British Columbia, Prince George, Canada (2004).
- [47] A. Psaltis *et al.*, *Nucl. Instrum. Methods Phys. Res A* **987**, 164828 (2021).
- [48] A. Psaltis, *Radiative alpha capture on ${}^7\text{Be}$ with DRAGON at vp -process nucleosynthesis energies*, *Ph.D. thesis*, McMaster University (2020).
- [49] W. A. Fowler *et al.*, *Rev. Mod. Phys.* **20**, 236 (1948).
- [50] S. Sjøe *et al.*, *Nucl. Instrum. Methods Phys. Res A* **700**, 179 (2013).
- [51] U. Hager *et al.*, *Phys. Rev. C* **85**, 035803 (2012).
- [52] D. Connolly *et al.*, *Phys. Rev. C* **97**, 035801 (2018).
- [53] M. Williams *et al.*, *Phys. Rev. C* **102**, 035801 (2020).
- [54] J. M. D'Auria *et al.*, *Phys. Rev. C* **69**, 065803 (2004).
- [55] K. Jayamanna, *Hyperfine Interact.* **225**, 51 (2014).
- [56] W. Liu *et al.*, *Nucl. Instrum. Methods Phys. Res A* **496**, 198 (2003).
- [57] J. F. Ziegler *et al.*, *Nucl. Instrum. Methods Phys. Res B* **268**, 1818 (2010).
- [58] C. Matei *et al.*, *Phys. Rev. Lett* **97**, 242503 (2006).
- [59] H. Rose and D. Brink, *Rev. Mod. Phys.* **39**, 306 (1967).
- [60] A. Psaltis *et al.* (2022), (Accepted).
- [61] G. J. Feldman and R. D. Cousins, *Phys. Rev. D* **57**, 3873 (1998).
- [62] K. F. Wallis, *Statistical Science* **29**, 106 (2014).
- [63] D. Lehbertz *et al.*, *Phys. Rev. C* **85**, 034333 (2012).
- [64] R. Longland *et al.*, *Nucl. Phys. A* **841**, 1 (2010).
- [65] J. H. Gaddum, *Nature* **156**, 463 (1945).
- [66] C. Iliadis *et al.*, *J Phys G Nucl Part Phys* **42**, 034007 (2015).
- [67] C. Iliadis, *Nuclear physics of stars* (John Wiley & Sons, 2015).
- [68] Q. Liu *et al.*, *Phys. Rev. C* **101**, 025808 (2020).
- [69] J. S. Randhawa *et al.*, *Phys. Rev. C* **104**, L042801 (2021).
- [70] R. Bollig *et al.*, *Astrophys. J* **915**, 28 (2021).