



This is the accepted manuscript made available via CHORUS. The article has been published as:

Statistical analysis for the neutrinoless double- β -decay matrix element of ${}^{48}\text{Ca}$

M. Horoi, A. Neacsu, and S. Stoica

Phys. Rev. C **106**, 054302 — Published 2 November 2022

DOI: [10.1103/PhysRevC.106.054302](https://doi.org/10.1103/PhysRevC.106.054302)

A Statistical Analysis for the Neutrinoless Double-Beta Decay Matrix element of ^{48}Ca

M. Horoi,¹ A. Neacsu,^{2,3} and S. Stoica²

¹*Department of Physics, Central Michigan University, Mount Pleasant, MI 48859, USA*

²*International Center for Advanced Training and Research in Physics (CIFRA), Magurele, Romania*

³*Horia Hulubei National Institute for Physics and Nuclear Engineering (IFIN-HH), Magurele, Romania*

(Dated: October 18, 2022)

Neutrinoless double beta decay ($0\nu\beta\beta$) nuclear matrix elements (NME) are the object of many theoretical calculation methods, and are very important for analysis and guidance of a large number of experimental efforts. However, there are large discrepancies between the NME values provided by different methods. In this paper we propose a statistical analysis of the ^{48}Ca $0\nu\beta\beta$ NME using the interacting shell model, emphasizing the range of the NME probable values and its correlations with observables that can be obtained from the existing nuclear data. Based on this statistical analysis with three independent effective Hamiltonians we propose a common probability distribution function for the $0\nu\beta\beta$ NME, which has a range of (0.45 - 0.95) at 90% confidence level of, and a mean value of 0.68.

I. INTRODUCTION

The study of the double-beta decay (DBD) is currently a hot research topic since it is viewed as one of the most promising approaches to clarify important, as yet unknown, properties of neutrinos and to explore physics beyond the Standard Model (SM) [1, 2]. Two scenarios are possible for this process to occur: i) two-neutrino double-beta ($2\nu\beta\beta$) transitions (with emission of two electrons/positrons and two anti-neutrinos/neutrinos), which conserve the lepton number and are allowed by the SM and ii) double-beta decay transitions without emission of neutrinos ($0\nu\beta\beta$), which violate the lepton number conservation and are only allowed by theories beyond SM (BSM).

Neutrinoless DBD was not experimentally detected so far, but its measurement would provide us with important information about lepton number violating (LNV) processes; neutrino properties (neutrino absolute mass scale and mass hierarchy, neutrino nature as Dirac or Majorana fermion, number of neutrino flavours); CP and Lorentz symmetries violation; constraining of different BSM mechanisms that may contribute to this decay mode, etc. The most common mechanism investigated is the light left-handed (LH) Majorana neutrinos exchange between two nucleons, but once a LNV operator is introduced in the Lagrangian, several other mechanisms are also allowed, such as: the exchange of light and heavy neutrinos in left-right symmetric models, the exchange of supersymmetric particles, DBD with the emission of Majorons, etc.

The DBD half-life equations can be expressed, in a good approximation, as a product of some factors. The $2\nu\beta\beta$ half-life is a product of a phase space factor (PSF), which depends on the atomic charge and energy released in the decay, and a nuclear matrix element (NME) related to the nuclear structure of the parent and daughter nuclei. The $0\nu\beta\beta$ half-life, besides the PSF and NME factors, also contains a LNV factor, related to the particular BSM mechanism that may contribute to the decay. If several mechanisms are considered, the inverse half-life

can be written as a sum of all the individual contributions and their interference terms [2–8]. Using the experimental limits of the $0\nu\beta\beta$ decay half-lives and the theoretical values of PSF and NME, one can constrain the LNV parameters and the associated BSM scenarios, usually under the assumption that only one mechanism contributes at one time.

There is currently significant progress in the DBD experiments (in terms of the amount of source material, decreasing background and improvement in the detection techniques), leading to the expectation that the next generation of experiments will be able to cover the entire region of the neutrino inverted mass hierarchy [9]. Concurrently, the progress of the theoretical methods now provides us with accurate PSF values for all the double-beta decay modes and transitions. [10–12]. Thus, at present, the uncertainty in the DBD calculations remains mostly the NME evaluation.

There are several nuclear structure methods for the NME calculation, the most used being: Shell model methods [13–23], pnQRPA methods [24–30], IBA methods [31, 32], Energy Density Functional method [33], PHFB [34], Coupled-Cluster method (CC) [35], in-medium generator coordinate method (IM-GCM) [36], and valence-space in-medium similarity renormalization group method (VS-IMSRG) [37]. Each of these methods have their strengths and weakness, largely discussed over time in the literature, and the current situation is that there are still significant differences between NME values calculated with different methods, and sometimes, even between NME values calculated with the same methods (see for example the review [9]). For the $2\nu\beta\beta$ decay the NME are products of two Gamow-Teller (GT) transition amplitudes, and most of the nuclear methods overestimate them, in comparison with experiment. This drawback is often treated by introducing a quenching factor that multiplies the GT operator and reduces its strength. This is equivalent to using a quenched axial vector constant, instead of its bare value $g_A = 1.27$.

For the $0\nu\beta\beta$ decay the NME calculation is more complicated, since besides the GT transitions, other tran-

sitions may contribute as well. Also, the NME values calculated by different methods may differ by factors of 3-4 for most relevant isotopes, and up to 7-8 in the case of ^{48}Ca (see e.g. Fig. 5 of Ref. [9], and Refs. [35, 36]). Uncertainties in the NME values are further amplified when predicting half-lives, since they enter at the power of two in the lifetime formulas. In addition, there is no measured lifetime for this decay mode to compare with, and these uncertainties in the NME computation reflect in the interpretation of the DBD data and planning the performances of the DBD experiments.

The shell model-based methods have some advantages such as the inclusion of all correlations between nucleons around the Fermi surface, preserving all symmetries of the nuclear many-body problem, and the use of widely tested nucleon-nucleon (NN) interactions. For different mass regions of nuclei, one uses several different effective NN effective Hamiltonians that are appropriate for the corresponding model spaces. These effective Hamiltonians are usually obtained by starting with a theoretical Bruekner G-Matrix Hamiltonian that is further fine-tuned to describe the experimental energy levels for a large number of nuclei that can be investigated in the corresponding model spaces. These effective Hamiltonians are described by a small number of single particle energies and a finite number of two-body matrix elements. As a by-product, the wave functions produced by these Hamiltonians can be used to describe and predict observables, such as the electromagnetic transition probabilities, Gamow-Teller transitions probabilities, nucleon occupation probabilities, spectroscopic factors, etc, using relative simple changes of the transition operators in terms of effective charges and quenching factors. These effective charges and quenching factors are calibrated to the existing data. For $0\nu\beta\beta$ NME such calibrations are not yet possible due to the lack of data. However, different existing effective Hamiltonians for nuclei involved in a given $0\nu\beta\beta$ decay produce smaller ranges of the NME. In addition, some recent ab-initio methods, such as IM-SRG [36, 37], build on the modern advances in the shell model by providing ab-initio derived effective Hamiltonians and effective transition operators, and they can provide some guidance for calibrating the shell model $0\nu\beta\beta$ NME.

It would be thus interesting to study the robustness of the $0\nu\beta\beta$ NME to small changes of the parameters of different effective shell model Hamiltonians and to examine how the NME changes are correlated with other observables. In this work, we propose a statistical analysis of $0\nu\beta\beta$ NME of ^{48}Ca calculated with the interacting shell-model using three independent effective Hamiltonians (FPD6, GXPF1A, KB3G), emphasizing the range of the NME probable values and their correlations with several observables that can be compared to existing nuclear data. Based on this statistical analysis we propose a common probability distribution function for the $0\nu\beta\beta$ NME. We apply our analysis to ^{48}Ca , which is the lightest DBD isotope and thus more accessible to ab-initio

calculations. We only consider in this work the standard light LH neutrino exchange mass mechanism, which is most likely to contribute to the $0\nu\beta\beta$ decay process.

The paper is organized as follows. In section II the calculation methods of the observables and the statistical model are presented. Then, in section III we present the results and discussions on their relevance, and in section IV we end with conclusions and outlook. Finally we included an Appendix with a short presentation of the Gram-Charlier A series that we used in our statistical model.

II. THE STATISTICAL MODEL

	Exp.	Error	FPD6	GXPF1A	KB3G
$0\nu\beta\beta$ NME	N/A	N/A	0.79	0.559	0.693
$2\nu\beta\beta$ NME	0.035 [38]	0.003	0.062	0.050	0.045
^{48}Ca B(E2) \uparrow	0.008 [39]	0.001	0.007	0.006	0.05
^{48}Ca 2+	3.832 [40]	0.15	3.658	3.735	4.238
^{48}Ca 4+	4.503 [40]	0.15	4.134	4.264	4.231
^{48}Ca 6+	7.953 [40]	0.15	7.396	7.705	7.831
^{48}Ca Occ(Nf5)	0.032*	0.395#	0.117	0.032	0.112
^{48}Ca Occ(Nf7)	7.892*	0.395#	7.693	7.892	7.795
^{48}Ca Occ(Np1)	0.009*	0.395#	0.029	0.009	0.024
^{48}Ca Occ(Np3)	0.067*	0.395#	0.161	0.067	0.070
$^{48}\text{Ca} \rightarrow ^{48}\text{Sc}$ GT	1.09 [41]	0.28	1.01	1.226	0.510
^{48}Ti B(E2) \uparrow	0.063 [39]	0.003	0.064	0.052	0.052
^{48}Ti 2+	0.984 [40]	0.150	1.118	1.010	0.985
^{48}Ti 4+	2.296 [40]	0.150	2.492	2.168	2.214
^{48}Ti 6+	3.333 [40]	0.150	3.425	2.922	3.046
^{48}Ti Occ(Nf5)	0.168*	0.277#	0.310	0.168	0.263
^{48}Ti Occ(Nf7)	5.535*	0.277#	5.253	5.535	5.416
^{48}Ti Occ(Np1)	0.048*	0.277#	0.068	0.048	0.061
^{48}Ti Occ(Np3)	0.248*	0.277#	0.369	0.248	0.260
^{48}Ti Occ(Pf5)	0.032*	0.092#	0.101	0.032	0.097
^{48}Ti Occ(Pf7)	1.839*	0.092#	1.672	1.839	1.763
^{48}Ti Occ(Pp1)	0.010*	0.092#	0.031	0.010	0.021
^{48}Ti Occ(Pp3)	0.119*	0.092#	0.196	0.119	0.120
$^{48}\text{Ti} \rightarrow ^{48}\text{Sc}$ GT	0.014 [41]	0.005	0.050	0.032	0.056

Table I. Experimental data, experimental errors, and the calculated values using 3 effective Hamiltonians for the observables analyzed. The data for occupation probabilities (*) is not available and it was replaced with the GXPF1A results and errors (#) of 5% of the highest nucleon species occupation.

We plan to investigate the effect of small, random variation of the shell model effective Hamiltonian on the neutrinoless double beta decay NME of ^{48}Ca , and the NME correlations with other calculated observables, such as 2^+ energies, B(E2) \uparrow values, $2\nu\beta\beta$ matrix elements, Gamow-Teller transition probabilities, neutron and proton occupation probabilities, etc.

To achieve that goal we selected a number of often used effective Hamiltonians describing nuclei around

^{48}Ca in the fp -shell ($0f_{7/2}$, $0f_{5/2}$, $1p_{3/2}$, and $1p_{1/2}$ orbitals for both protons and neutrons), and added small random contributions to their two-body matrix elements (TBME). For this project we only considered the FPD6 Hamiltonian [42], the KB3G Hamiltonian [15], and GXPF1A Hamiltonian [43, 44] as starting effective Hamiltonians. In order to maintain the magicity of ^{48}Ca we decided to keep the single particle (s.p.) energies in the perturbed effective Hamiltonians the same as in the starting Hamiltonians. Certainly, this choice is limiting our analysis to shell model techniques, the fp model space, and only three effective Hamiltonians that although widely used have their limitations. One could, in principle use a larger model space, such as sd - pf , or consider ab-initio derived effective Hamiltonians, such as those proposed in [36, 37, 45]. However, this would be beyond the limited purpose of this paper.

One important decision to be made about the random contributions to the starting Hamiltonians is the choice of their maximum amplitude (range). In this work we were guided by the analysis of the USDA/USDB effective Hamiltonians [46] where one starts with an underlying G-matrix and modifies linear combinations of two-body matrix elements in a fine-tuning procedure [46] until the root mean square (RMS) deviation of the calculated energies vs the experimental ones shows some signs of convergence. In this fine-tuning process one would not want to change the TBME too much from the original G-matrix values, because the over-fitted TBME could result in unitary changes of the s.p. wave functions that may produce slightly better energies, but incorrect observables. For USDA, for example, the RMS deviation of the TBME was about 300 keV, while a small improvement in the overall energies given by USDB resulted in an additional change of 100 keV if the RMS deviation of the TBME. This analysis suggests that an additional RMS of about 100 keV would not dramatically change the quality of the TBME in the sd -shell and we extended this choice to the fp -shell. An analysis of the TBME for all three starting Hamiltonians listed above indicates that a $\pm 10\%$ range for the random contributions would suffice. Another interesting study would be to consider selectively modifying only specific parts of the effective Hamiltonian that would correspond to certain properties, such as pairing, deformation, or the ones that affect the rotational properties of the daughter nucleus. This would imply a different analysis that could relate changes in the values of the observables to changes of the matrix elements of the effective Hamiltonian with specific spin and isospin, and we plan to do such investigations in a future paper.

In the analysis we included as observables the $0\nu\beta\beta$ NME, the $2\nu\beta\beta$ NME, the Gamow-Teller probability to reach the first 1^+ state in ^{48}Sc from the ground states (g.s.) of the parent, ^{48}Ca , and of the daughter, ^{48}Ti , the energies of the 2^+ , 4^+ and 6^+ states of the parent and daughter, the $B(E2)\uparrow$ transition probabilities to the first 2^+ state of parent and daughter, the neutron occupa-

tion probabilities of the pf states of the parent, and neutron and proton occupation probabilities of the pf states of the daughter nucleus. The experimental occupation probabilities for the nuclei relevant for the ^{48}Ca $0\nu\beta\beta$ decay are not available, but we include synthetic (calculated) values in the analysis because the corresponding occupation probabilities are available for other nuclei of interest for $0\nu\beta\beta$ decay [47–49], and it might be interesting to see if they have any correlations with the $0\nu\beta\beta$ NME. All in all, there are 24 observables included in our statistical analysis. Some studies also look at other observables that could be of relevance to the double-beta decay analysis, such as the related one muon capture (OMC) rates [50]. Recent references for the OMC data are [51] and [52]. However, one muon capture rate calculations are very complex, depending on interfering contributions to the decay amplitude (see e.g. Ref. [50]), and are very sensitive to the effective Hamiltonian used (see e.g. Ref. [53]). Given this complexity we decided to postpone the inclusion of the OMC rates in our observables list.

The main goals are: (i) for each starting effective Hamiltonian find correlations between $0\nu\beta\beta$ NME and the other observables that are accessible experimentally; (ii) find theoretical ranges for each observables; (iii) establish the shape of different distributions for each observables and starting Hamiltonians; (iv) use this information to find weights of contributions from different starting Hamiltonians to the "optimal" distribution of the $0\nu\beta\beta$ NME; (v) find an "optimal" value of the $0\nu\beta\beta$ NME and its predicted probable range (theoretical error). One should mention that similar studies for other observables were recently proposed [54].

The $0\nu\beta\beta$ NME is related to the half-life of the respective process [17] by

$$\left(T_{1/2}^{0\nu}\right)^{-1} = G_{0\nu}(E_0, Z)g_A^4 |M_{0\nu}|^2 |\langle \eta_l \rangle|^2, \quad (1)$$

where $G_{0\nu}$ and $M_{0\nu}$ are the PSF and nuclear matrix elements for the 0ν decay, g_A is the axial vector coupling constant, and $\langle \eta_l \rangle \equiv \langle m_{\beta\beta} \rangle / m_e c^2$ is a BSM parameter associated with the light neutrino exchange mechanisms. Here we only consider the contribution from the light LH neutrino exchange mechanism, which is likely to contribute to the $0\nu\beta\beta$ decay. The methodology of calculating the $0\nu\beta\beta$ NME, $M_{0\nu}$, within the shell model was extensively described elsewhere [17, 18, 23] and it will not be repeated here. Suffices to say that it includes a short range correlation function that can be viewed as an effective modification of the bare operator (see below). We should mention that we do not include the recently proposed contact term contribution to the $0\nu\beta\beta$ operator [55–57], which we believe that's more appropriate for a chiral EFT approach to the short-range correlations in nuclei. In addition, its associated weak coupling strength is somewhat uncertain and needs to be obtained consistently from the data (see e.g. [45]). However, we "cali-

brate” our short-range correlator to the ab-initio results [37, 45].

The $2\nu\beta\beta$ NME is related to the half-life of the respective process [16] by

$$\left(T_{1/2}^{2\nu}\right)^{-1} = G_{2\nu}(E_0, Z)g_A^4 |m_e c^2 M_{2\nu}|^2. \quad (2)$$

Here, $G_{2\nu}$ is the appropriate PSF, and $M_{2\nu}$ can be calculated with

$$M_{2\nu} = \sum_k \frac{q^2 \langle 0_f^+ | \sigma\tau^- | 1_k^+ \rangle \langle 1_k^+ | \sigma\tau^- | 0_i^+ \rangle}{E_k - E_0}, \quad (3)$$

where the summation is on the 1_k^+ states in ^{48}Sc and $E_0 = Q_{\beta\beta}/2 + \Delta M(^{48}\text{Sc} - ^{48}\text{Ca})$

Often, the shell model calculations of the $0\nu\beta\beta$ NME are described as using the “bare” transition operator. This characterization is unfortunate, while the transition operator (see e.g. Eq. (7-12) of Ref. [17]) contains the bare operator from the underlying theory of $0\nu\beta\beta$ decay, modified by a phenomenological effective short-range correlation function, $1 + f(r)$, which is quenching the $0\nu\beta\beta$ NME. Therefore, the short-range modification of the bare operator acts as an effective operator. In practice the parameters of an effective operator need to be calibrated to the data. Given that the short-range correlator has

radial dependence, its calibration has been only done relative to some ab-initio results. The standard Miller-Spencer short-range correlator [17, 58] produces the highest quenching of the NME, while the CD-Bonn parameterization of the short-range correlator [59] produces little to no quenching. A direct renormalization of the $0\nu\beta\beta$ NME by a similarity renormalization group (SRG) evolution of the NME of the bare operator from 200 to 10 major harmonic oscillator shells using CDBonn two body wave functions, indicates that using a phenomenological CDBonn parametrization of the short-range correlator is a reasonable approach [60]. More recent ab-initio calculations of the $0\nu\beta\beta$ NME using the $N^3\text{LO}$ Hamiltonian provides more quenched values, more consistent with the shell model results base on Miller-Spencer parametrization of the short-range correlator. In an effort to calibrate the effective operator used in shell model calculations to the latest ab-initio results we used the Miller-Spencer correlator in this study.

The other observables used in this study including the excited state energies, the GT strengths to the first 1^+ state ^{48}Sc , the $\text{B}(\text{E}2)\uparrow$ to the first 2^+ state in the parent and daughter, as well as the s.p. occupation probabilities, are calculated in the standard way. Here we use in all cases the same effective charges ($e_p = 1.5$ and $e_n = 0.5$) for the $\text{B}(\text{E}2)\uparrow$, and the same quenching factor ($q = 0.74$) for the the GT strengths and $M_{2\nu}$.

III. RESULTS

The experimental data used in this study listed in Table I and in the legends of the rightmost column in Figures 1/2 - 5/6 are taken from Ref. [40] (excitation energies of the 2^+ , 4^+ and 6^+ states of ^{48}Ca and ^{48}Ti in MeV), Ref. [38] ($2\nu\beta\beta$ NME in MeV^{-1}), Ref. [39] ($\text{B}(\text{E}2)\uparrow$ in $e^2 b^2$), and Ref. [41] for the GT transition probabilities to the first excited 1^+ state in ^{48}Sc . The experimental errors for the excitation energies are very small, and for the calculation the χ^2 we use the typical theoretical RMS value of 150 keV [43]. The experimental occupation probabilities are not available, and we took as reference the GXPF1A results assuming a uniform error that we choose to be 5% of the highest occupation for a each nucleon species in the fp -shell. In the Figures Occ (Nf7) designates the neutron occupation probability of the $f_{7/2}$ s.p. orbital, Occ (Pf3) designates the proton occupation probability of the $p_{3/2}$ s.p. orbital, etc.

Figures 1/2 - 5/6 show the main results of this study. The leftmost columns indicates the 24 observables discussed in section II, including the $0\nu\beta\beta$ NME. The middle column shows the scatter plots of the correlation of each variable with the $0\nu\beta\beta$ NME, and the last column

shows the distribution of each observable when the random term is added to the respective effective Hamiltonian. The legends in column with correlations show the standard Pearson correlator R , and in the last columns the legends include the mean, standard deviations, and the skewness (normalized 3rd moment) of the distributions, as well as the result for the starting interactions (FPD6, GXPF1A, and KB3G) and the experimental values when available. Figures 1/2 present the results for the FPD6 effective Hamiltonian, Figures 3/4 show the results for the GXPF1A effective Hamiltonian, and Figures 5/6 present the results for the KB3G effective Hamiltonian. For each starting effective Hamiltonian we use 20,000 random Hamiltonians produced by the procedure described in section II.

The results in Figures 1/2 - 5/6 indicate strong correlations between the $0\nu\beta\beta$ NME and the $2\nu\beta\beta$ NME. This behavior is expected due to the GT transition operator that dominates the results for both NME. Alternative approaches of obtaining these NME, e.g. QRPA calculations, are calibrating parts of their nuclear Hamiltonian, such as the isoscalar particle-particle interaction g_{pp} , to describe the experimental value of the $2\nu\beta\beta$ NME and to approximately restore the isospin symmetry, thus induc-

	$0\nu\beta\beta$ NME	$2\nu\beta\beta$ NME	^{48}Ca B(E2) \uparrow	^{48}Ca 2+	^{48}Ca 4+	^{48}Ca 6+	^{48}Ca Occ (Nf5)	^{48}Ca Occ (Nf7)	^{48}Ca Occ (Np1)	^{48}Ca Occ (Np3)	$^{48}\text{Ca} \rightarrow$ ^{48}Sc GT	^{48}Ti B(E2) \uparrow
$0\nu\beta\beta$ NME	1.00	0.90	0.43	0.22	0.24	0.18	0.62	-0.58	0.30	0.41	0.12	-0.06
$2\nu\beta\beta$ NME	0.90	1.00	0.38	0.28	0.30	0.17	0.44	-0.47	0.25	0.37	0.10	-0.18
^{48}Ca B(E2) \uparrow	0.43	0.38	1.00	-0.35	-0.27	-0.31	0.55	-0.91	0.53	0.92	-0.17	0.32
^{48}Ca 2+	0.23	0.28	-0.35	1.00	0.96	0.49	-0.06	0.32	-0.23	-0.42	0.51	-0.38
^{48}Ca 4+	0.24	0.30	-0.27	0.96	1.00	0.44	-0.01	0.26	-0.17	-0.36	0.54	-0.33
^{48}Ca 6+	0.18	0.17	-0.31	0.49	0.44	1.00	-0.22	0.25	-0.19	-0.20	0.26	-0.22
^{48}Ca Occ(Nf5)	0.62	0.44	0.55	-0.06	-0.01	-0.22	1.00	-0.77	0.38	0.42	-0.04	0.12
^{48}Ca Occ(Nf7)	-0.58	-0.47	-0.91	0.32	0.26	0.25	-0.77	1.00	-0.58	-0.90	0.17	-0.30
^{48}Ca Occ(Np1)	0.30	0.25	0.53	-0.23	-0.17	-0.19	0.38	-0.58	1.00	0.48	-0.11	0.19
^{48}Ca Occ(Np3)	0.41	0.37	0.92	-0.42	-0.36	-0.20	0.42	-0.90	0.48	1.00	-0.22	0.34
$^{48}\text{Ca} \rightarrow$ ^{48}Sc GT	0.12	0.10	-0.17	0.51	0.54	0.26	-0.04	0.17	-0.11	-0.22	1.00	-0.27
^{48}Ti B(E2) \uparrow	-0.06	-0.18	0.32	-0.38	-0.34	-0.22	0.12	-0.30	0.19	0.34	-0.27	1.00
^{48}Ti 2+	0.80	0.89	0.37	0.32	0.32	0.20	0.47	-0.47	0.24	0.35	0.23	-0.24
^{48}Ti 4+	0.78	0.85	0.35	0.35	0.35	0.25	0.45	-0.45	0.23	0.34	0.22	-0.19
^{48}Ti 6+	0.75	0.80	0.42	0.28	0.28	0.24	0.48	-0.52	0.26	0.42	0.03	0.13
^{48}Ti Occ(Nf5)	0.14	-0.03	0.39	-0.16	-0.11	-0.36	0.75	-0.53	0.25	0.24	-0.00	0.27
^{48}Ti Occ(Nf7)	0.12	0.32	-0.51	0.47	0.42	0.37	-0.37	0.53	-0.33	-0.49	0.20	-0.82
^{48}Ti Occ(Np1)	-0.13	-0.29	0.27	-0.34	-0.29	-0.24	0.10	-0.26	0.56	0.25	-0.02	0.65
^{48}Ti Occ(Np3)	-0.25	-0.38	0.43	-0.51	-0.47	-0.26	0.03	-0.37	0.21	0.50	-0.28	0.88
^{48}Ti Occ(Pf5)	0.22	0.01	0.20	0.00	0.03	-0.08	0.56	-0.35	0.15	0.12	0.21	0.27
^{48}Ti Occ(Pf7)	0.21	0.41	-0.16	0.24	0.20	0.16	-0.10	0.16	-0.11	-0.16	-0.01	-0.85
^{48}Ti Occ(Pp1)	-0.11	-0.29	0.13	-0.19	-0.16	-0.14	0.06	-0.13	0.30	0.12	0.16	0.60
^{48}Ti Occ(Pp3)	-0.33	-0.49	0.11	-0.28	-0.25	-0.16	-0.09	-0.06	0.05	0.14	-0.09	0.89
$^{48}\text{Ti} \rightarrow$ ^{48}Sc GT	0.15	0.40	0.23	-0.03	-0.02	-0.24	0.15	-0.20	0.13	0.19	-0.55	0.03

Table II. Correlation matrix for the 24 observables described in the text using the FPD6 starting Hamiltonian (continues in Table III).

	^{48}Ti 2+	^{48}Ti 4+	^{48}Ti 6+	^{48}Ti Occ (Nf5)	^{48}Ti Occ (Nf7)	^{48}Ti Occ (Np1)	^{48}Ti Occ (Np3)	^{48}Ti Occ (Pf5)	^{48}Ti Occ (Pf7)	^{48}Ti Occ (Pp1)	^{48}Ti Occ (Pp3)	$^{48}\text{Ti} \rightarrow$ ^{48}Sc GT
$0\nu\beta\beta$ NME	0.80	0.78	0.75	0.14	0.12	-0.13	-0.25	0.22	0.21	-0.11	-0.33	0.15
$2\nu\beta\beta$ NME	0.89	0.85	0.80	-0.03	0.32	-0.29	-0.38	0.01	0.41	-0.28	-0.48	0.39
^{48}Ca B(E2) \uparrow	0.37	0.35	0.42	0.39	-0.51	0.27	0.43	0.19	-0.16	0.13	0.11	0.23
^{48}Ca 2+	0.32	0.35	0.27	-0.16	0.47	-0.34	-0.51	0.00	0.24	-0.19	-0.28	-0.03
^{48}Ca 4+	0.32	0.35	0.28	-0.11	0.42	-0.29	-0.47	0.03	0.20	-0.16	-0.25	-0.02
^{48}Ca 6+	0.20	0.25	0.24	-0.36	0.37	-0.24	-0.26	-0.07	0.16	-0.14	-0.16	-0.24
^{48}Ca Occ(Nf5)	0.47	0.45	0.48	0.75	-0.37	0.10	0.03	0.56	-0.10	0.06	-0.09	0.15
^{48}Ca Occ(Nf7)	-0.47	-0.45	-0.52	-0.53	0.53	-0.26	-0.37	-0.35	0.16	-0.13	-0.06	-0.20
^{48}Ca Occ(Np1)	0.24	0.23	0.26	0.25	-0.33	0.56	0.21	0.15	-0.11	0.30	0.05	0.13
^{48}Ca Occ(Np3)	0.35	0.34	0.42	0.24	-0.49	0.25	0.50	0.12	-0.16	0.12	0.14	0.19
$^{48}\text{Ca} \rightarrow$ ^{48}Sc GT	0.23	0.22	0.03	0.00	0.20	-0.02	-0.28	0.21	-0.01	0.16	-0.09	-0.55
^{48}Ti B(E2) \uparrow	-0.24	-0.19	0.13	0.27	-0.82	0.65	0.88	0.27	-0.85	0.60	0.89	0.03
^{48}Ti 2+	1.00	0.96	0.85	0.16	0.25	-0.24	-0.42	0.20	0.36	-0.19	-0.51	0.28
^{48}Ti 4+	0.96	1.00	0.89	0.16	0.21	-0.21	-0.36	0.22	0.29	-0.15	-0.44	0.24
^{48}Ti 6+	0.85	0.89	1.00	0.20	-0.01	-0.09	-0.09	0.22	0.08	-0.07	-0.18	0.35
^{48}Ti Occ(Nf5)	0.16	0.16	0.20	1.00	-0.64	0.34	0.22	0.81	-0.42	0.35	0.18	-0.05
^{48}Ti Occ(Nf7)	0.25	0.21	-0.01	-0.64	1.00	-0.74	-0.88	-0.55	0.86	-0.66	-0.80	0.12
^{48}Ti Occ(Np1)	-0.24	-0.21	-0.09	0.34	-0.74	1.00	0.66	0.41	-0.75	0.88	0.67	-0.22
^{48}Ti Occ(Np3)	-0.42	-0.36	-0.09	0.22	-0.88	0.66	1.00	0.19	-0.82	0.56	0.90	-0.10
^{48}Ti Occ(Pf5)	0.20	0.22	0.22	0.81	-0.55	0.41	0.19	1.00	-0.57	0.56	0.26	-0.30
^{48}Ti Occ(Pf7)	0.36	0.29	0.08	-0.42	0.86	-0.75	-0.82	-0.57	1.00	-0.81	-0.94	0.31
^{48}Ti Occ(Pp1)	-0.19	-0.15	-0.07	0.35	-0.66	0.88	0.56	0.56	-0.81	1.00	0.67	-0.35
^{48}Ti Occ(Pp3)	-0.51	-0.44	-0.18	0.18	-0.80	0.67	0.90	0.26	-0.94	0.67	1.00	-0.23
$^{48}\text{Ti} \rightarrow$ ^{48}Sc GT	0.28	0.24	0.35	-0.05	0.12	-0.22	-0.10	-0.30	0.31	-0.35	-0.23	1.00

Table III. Correlation matrix: continuation of Table II.

ing correlation with the $2\nu\beta\beta$ NME. In the shell model approach, the Hamiltonian remains unchanged, and all

symmetries are enforced. Therefore, we conclude that the strong correlations between the $0\nu\beta\beta$ NME and the $2\nu\beta\beta$ NME are genuine.

Interestingly, the correlations between the $0\nu\beta\beta$ NME and the Gamow-Teller (GT) transitions probabilities to the first 1^+ state in ^{48}Sc are much reduced. One explanation of this phenomena is based on the fact the distributions of the GT strength from the parent and daughter (see last column in Figures 1/2 - 5/6) are asymmetric in opposite direction, thus diminishing the correlation effects. A quick look to the full correlation matrix in Table II shows that the GT strengths to the first 1^+ state in ^{48}Sc from ^{48}Ca and ^{48}Ti are anti-correlated with a correlation coefficient of about -0.5.

Other observables that have relatively high (anti)correlations with the $0\nu\beta\beta$ NME are the en-

ergies of the 2^+ , 4^+ and 6^+ states in ^{48}Ti , and the neutron occupation probabilities in ^{48}Ca . Overall, the correlators R with the $2\nu\beta\beta$ NME are around 0.9, the ones with the energies of the 2^+ , 4^+ and 6^+ states in ^{48}Ti are about 0.77, and correlators with the $0f_{5/2}$ occupation probability is about 0.6, while the occupation probability of the $0f_{7/2}$ is anti-correlated with the $0\nu\beta\beta$ NME, $R \approx -0.6$.

The correlations between $0\nu\beta\beta$ NME and the energies of the $2+$, $4+$ and $6+$ states in the daughter could be related to the mismatch of the pairing effects between the parent and daughter, given that the correlations with the B(E2) values are weak. This conclusion could be strengthened when one studies the effects of selectively changing the pairing two body matrix elements of the effective Hamiltonian.

Additional interesting information can be extracted from the full correlation matrix for all 24 observables. Tables II - III show the full correlation matrix evaluated for the FPD6 starting Hamiltonian. It is interesting to analyze which other observables are correlated with those that are directly correlated with the $0\nu\beta\beta$ NME. We already discussed the correlations between the GT strengths and the $2\nu\beta\beta$ NME. In addition, one can observe that the 2^+ , 4^+ and 6^+ in ^{48}Ti are correlated with the neutron f and p states occupancies in ^{48}Ca , which in their turn are correlated with some of the neutron states occupancies in ^{48}Ti . Also, some of the neutron occupation probabilities in ^{48}Ti are correlated to the B(E2) \uparrow values. It is also observed that three of the neutron occupancies in ^{48}Ca correlate with the $0\nu\beta\beta$ and $2\nu\beta\beta$ NME, while the fourth anti-correlates. One could understand the (anti)correlation occupancies of the four orbitals because three of them constrain the 4th as the number of particles is fixed. These observations highlight the importance of a reliable experimental investigation of the occupation probabilities for these nuclei. In some cases, most noticeably the B(E2) \uparrow PDF plots shown in Figures 1 - 6, there appears to be some significant fluctuations in the density and height of the bars. This is simply a plotting artifact due to rounding errors when displaying a very narrow range of small values that are mapped on a small number of bins.

It would be interesting to extract some information about possible range and mean value of the $0\nu\beta\beta$ NME based on this statistical analysis. First, it is clear that the value of all observables are quite stable to reasonably small changes of the effective Hamiltonian. No hints of any wild departure from the main values that would indicate some phase transitions are found. This seems to be a consequence of the preservation of nuclear many-body symmetries in the shell model. One can further try using

the distributions of all available effective Hamiltonians to draw conclusions on some optimal values for the $0\nu\beta\beta$ NME and its range (error). One direct approach would be to superpose the distributions of the NME produced in Figures 1/2 - 5/6 with some weighting factors W_H ,

$$P(x) = W_{FPD6}P_{FPD6}(x) + W_{GXPF1A}P_{GXPF1A}(x) + W_{KB3G}P_{KB3G}(x), \quad (4)$$

where x is the value of the $0\nu\beta\beta$ NME. The normalized weights W_H can be inferred using, for example, the likelihood probability $\propto \exp(-\chi^2/2)$ for independent observables, and the Bayesian Model Averaging for the correlated observables that dominate the contributions to the weights (see e.g. section 4 of [61]). Based on the data we show in Table II, we get the following χ^2 for each starting effective Hamiltonian: 7.9 for FPD6, 4.8 for GXPF1A, and 7.3 for KB3G. Unfortunately, there is no experimental data for the occupation probabilities that seem to correlate directly and indirectly with the $0\nu\beta\beta$ NME. Therefore, here we present the results of a "democratic" approach in which all W_H are 0.33. Fig. 7 shows the probability distribution functions (PDF) for the three starting effective Hamiltonians and their weighted sum. To calculate each PDF we use the Gram-Charlier A series expansion [62] (see Appendix for detail), based on the first four normalized moments of the distributions presented in Figures 1/2 - 5/6. Based on the results of our statistical analysis summarized in Fig. 7 (see "weighted sum" curve) one can infer that with 90% confidence the $0\nu\beta\beta$ NME lies in the range between 0.45 and 0.95, with a mean value of about 0.68.




Figure 1. (Color online) Correlations scattered plots and PDFs for the FPD6 starting Hamiltonian.




Figure 2. (Color online) Continuation of Figure 1.




Figure 3. (Color online) Correlations scattered plots and PDFs for the GXPF1A starting Hamiltonian.




Figure 4. (Color online) Continuation of Figure 3.




Figure 5. (Color online) Correlations scattered plots and PDFs for the KB3G starting Hamiltonian.




Figure 6. (Color online) Continuation of Figure 5.




Figure 7. PDF of the $0\nu\beta\beta$ NME distributions for the FPD6, GXPF1A and KB3G Hamiltonians and their weighted sum (see text for details).

IV. CONCLUSION AND OUTLOOK

In conclusion, we developed a statistical model for analyzing the distribution of the $0\nu\beta\beta$ NME of ^{48}Ca using the interactive shell model in the fp -shell model space. In the analysis we started from three widely used effective Hamiltonians for the low part of the fp -shell, FPD6, GXPF1A and KB3G, to which we added a random contributions to the TBME of $\pm 10\%$. Using sample sizes of 20,000 points we analyzed for each starting effective Hamiltonian: (i) the correlations between $0\nu\beta\beta$ NME and the other observables that are accessible experimentally; (ii) the theoretical ranges for each observables; (iii) the shape of different distributions for each observables and starting Hamiltonians; (iv) the weighted contributions from different starting Hamiltonians to the "optimal" distribution of the $0\nu\beta\beta$ NME; (v) an "optimal" value of the $0\nu\beta\beta$ NME and its predicted probable range (theoretical error).

We found that the $0\nu\beta\beta$ NME correlates strongly with the $2\nu\beta\beta$ NME, but much less with the Gamow-Teller strengths to the first 1^+ state in ^{48}Sc . We also found that the $0\nu\beta\beta$ NME exhibits reasonably strong correlations with the energies of the 2^+ , 4^+ and 6^+ states in ^{48}Ti , and with the neutron occupation probabilities in ^{48}Ca . We also found that there are additional correlations between observables, such as the energies of the 2^+ , 4^+ and 6^+ states in ^{48}Ti and the neutron occupation probabilities, as well as between $B(E2)\uparrow$ values in ^{48}Ti and proton and neutron occupation probabilities, which can indirectly influence the $0\nu\beta\beta$ NME. Therefore, we conclude that reliable experimental values of the occupa-

tion probabilities in ^{48}Ti and ^{48}Ca would be useful for this analysis, potentially helpful to reduce the uncertainties of the $0\nu\beta\beta$ NME.

Based on this statistical analysis with three independent effective Hamiltonians we propose a common probability distribution function for the $0\nu\beta\beta$ NME, which has a range (theoretical error) of (0.45 - 0.95) at 90% confidence level, and a mean value of 0.68. We also hope that the present analysis will help ab-initio studies, such as [35–37] to better identify correlations and further reduce the uncertainties of the $0\nu\beta\beta$ NME

Acknowledgements. MH acknowledges support from the US Department of Energy grant DE-SC0022538 "Nuclear Astrophysics and Fundamental Symmetries". SS and AN acknowledge support by grants of Romanian Ministry of Research, Innovation and Digitalization through the projects CNCS – UEFISCDI number 99/2021 within PN-III-P4-ID-PCE-2020-2374 and CNCS – UEFISCDI number TE 12/2022 within PN-III-P1-1.1-TE-2021-0343.

Appendix: Gram-Charlier A series

In order to have a good representation for the PDF of the $0\nu\beta\beta$ NME, we consider small deviation from normal distribution via the Gram-Charlier A series [62]. This is given by:

$$P(x) \approx \frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\frac{(x-\mu)^2}{2\sigma^2}\right] \cdot \left[1 + \frac{\mu_3}{3!} He_3((x-\mu)/\sigma) + \frac{\mu_4-3}{4!} He_4((x-\mu)/\sigma)\right] \quad (\text{A.1})$$

where $He_k(y)$ are the Chebyshev-Hermite polynomials, $He_3(y) = y^3 - 3y$ and $He_4(y) = y^4 - 6y^2 + 3$, μ and σ are the mean and variance of a probability density function (PDF), and μ_k , with $k = 3, 4$ are normalized moments of the same PDF, $P(x)$:

$$\mu_k = \int \left(\frac{x-\mu}{\sigma}\right)^k P(x) dx, \quad (\text{A.2})$$

In practice, we use the sample moments, μ_3 (skewness) and $\mu_4 - 3$ (kurtosis), which in the limit of very large sample sizes become very close to the underlying moments.

-
- [1] F. T. Avignone, III, S. R. Elliott, and J. Engel, *Rev. Mod. Phys.* **80**, 481 (2008).
 [2] J. D. Vergados, H. Ejiri, and F. Simkovic, *Rep. Prog.*

- Phys.* **75**, 106301 (2012).
 [3] M. Doi, T. Kotani, and E. Takasugi, *Prog. Theor. Phys. Suppl.* **83**, 1 (1985).

- [4] W. Rodejohann, *J. Phys. G* **39**, 124008 (2012).
- [5] F. F. Deppisch, M. Hirsch, and H. Pas, *J. Phys. G* **39**, 124007 (2012).
- [6] M. Horoi and A. Neacsu, *Phys. Rev. D* **93**, 113014 (2016), [arXiv:arXiv:1511.00670 \[hep-ph\]](#) [hep-ph].
- [7] A. Neacsu and M. Horoi, *Advances in High Energy Physics* **2016** (2016).
- [8] F. Ahmed, A. Neacsu, and M. Horoi, *Physics Letters B* **769**, 299 (2017).
- [9] J. Engel and J. Menéndez, *Reports on Progress in Physics* **80**, 046301 (2017).
- [10] J. Kotila and F. Iachello, *Phys. Rev. C* **85**, 034316 (2012).
- [11] S. Stoica and M. Mirea, *Phys. Rev. C* **88**, 037303 (2013).
- [12] M. Mirea, T. Pahomi, and S. Stoica, *Rom. Rep. Phys.* **67**, 872 (2015).
- [13] E. Caurier, A. Poves, and A. P. Zuker, *Phys. Lett. B* **252**, 13 (1990).
- [14] E. Caurier, F. Nowacki, A. Poves, and J. Retamosa, *Phys. Rev. Lett.* **77**, 1954 (1996).
- [15] E. Caurier, G. Martínez-Pinedo, F. Nowacki, A. Poves, and A. P. Zuker, *Rev. Mod. Phys.* **77**, 427 (2005).
- [16] M. Horoi, S. Stoica, and B. A. Brown, *Phys. Rev. C* **75**, 034303 (2007).
- [17] M. Horoi and S. Stoica, *Phys. Rev. C* **81**, 024321 (2010).
- [18] M. Horoi, *Phys. Rev. C* **87**, 014320 (2013).
- [19] M. Horoi and B. A. Brown, *Phys. Rev. Lett.* **110**, 222502 (2013).
- [20] R. A. Sen'kov and M. Horoi, *Phys. Rev. C* **90**, 051301(R) (2014).
- [21] A. Neacsu and M. Horoi, *Phys. Rev. C* **91**, 024309 (2015).
- [22] M. Horoi and A. Neacsu, *Phys. Rev. C* **93**, 024308 (2016).
- [23] M. Horoi and A. Neacsu, *Phys. Rev. C* **98**, 035502 (2018).
- [24] J. Suhonen and O. Civitarese, *Phys. Rep.* **300**, 123 (1998).
- [25] F. Simkovic, G. Pantis, J. D. Vergados, and A. Faessler, *Phys. Rev. C* **60**, 055502 (1999).
- [26] S. Stoica and H. Klapdor-Kleingrothaus, *Nucl. Phys. A* **694**, 269 (2001).
- [27] V. Rodin, A. Faessler, F. Simkovic, and P. Vogel, *Nucl. Phys. A* **766**, 107 (2006).
- [28] M. Kortelainen and J. Suhonen, *Phys. Rev. C* **75**, 051303(R) (2007).
- [29] A. Faessler, V. Rodin, and F. Simkovic, *J. Phys. G* **39**, 124006 (2012).
- [30] F. Simkovic, V. Rodin, A. Faessler, and P. Vogel, *Phys. Rev. C* **87**, 045501 (2013).
- [31] J. Barea and F. Iachello, *Phys. Rev. C* **79**, 044301 (2009).
- [32] J. Barea, J. Kotila, and F. Iachello, *Phys. Rev. C* **87**, 014315 (2013).
- [33] T. R. Rodriguez and G. Martinez-Pinedo, *Phys. Rev. Lett.* **105**, 252503 (2010).
- [34] P. K. Rath, R. Chandra, K. Chaturvedi, P. Lohani, P. K. Raina, and J. G. Hirsch, *Phys. Rev. C* **88**, 064322 (2013).
- [35] S. Novario, P. Gysbers, J. Engel, G. Hagen, G. R. Jansen, T. D. Morris, P. Navrátil, T. Papenbrock, and S. Quaglioni, *Phys. Rev. Lett.* **126**, 182502 (2021).
- [36] J. M. Yao, B. Bally, J. Engel, R. Wirth, T. R. Rodríguez, and H. Hergert, *Phys. Rev. Lett.* **124**, 232501 (2020).
- [37] A. Belley, C. G. Payne, S. R. Stroberg, T. Miyagi, and J. D. Holt, *Phys. Rev. Lett.* **126**, 042502 (2021).
- [38] A. Barabash, *UNIVERSE* **6**, 159 (2020).
- [39] B. Pritychenko, M. Birch, B. Singh, and M. Horoi, *ATOMIC DATA AND NUCLEAR DATA TABLES* **107**, 1 (2016).
- [40] J. Chen, *Nuclear Data Sheets* **179**, 1 (2022).
- [41] E.-W. Grewe, D. Frekers, S. Rakers, T. Adachi, C. Bäumer, N. T. Botha, H. Dohmann, H. Fujita, Y. Fujita, K. Hatanaka, K. Nakanishi, A. Negret, R. Neveling, L. Popescu, Y. Sakemi, Y. Shimbara, Y. Shimizu, F. D. Smit, Y. Tameshige, A. Tamii, J. Thies, P. vonBrentano, M. Yosoi, and R. G. T. Zegers, *Phys. Rev. C* **76**, 054307 (2007).
- [42] A. Richter, M. G. van der Merwe, R. E. Julies, and B. A. Brown, *Nucl. Phys. A* **523**, 325 (1991).
- [43] M. Honma, T. Otsuka, B. A. Brown, and T. Mizusaki, *Phys. Rev. C* **69**, 034335 (2004).
- [44] M. Honma, T. Otsuka, B. A. Brown, and T. Mizusaki, *Eur. Phys. J. A* **25 Suppl.** **1**, 499 (2005).
- [45] R. Wirth, J. M. Yao, and H. Hergert, *Phys. Rev. Lett.* **127**, 242502 (2021).
- [46] B. A. Brown and W. A. Richter, *Phys. Rev. C* **74**, 034315 (2006).
- [47] J. P. Schiffer, S. J. Freeman, J. A. Clark, C. Deibel, C. R. Fitzpatrick, S. Gros, A. Heinz, D. Hirata, C. L. Jiang, B. P. Kay, A. Parikh, P. D. Parker, K. E. Rehm, A. C. C. Villari, V. Werner, and C. Wrede, *Phys. Rev. Lett.* **100**, 112501 (2008).
- [48] B. P. Kay, J. P. Schiffer, S. J. Freeman, T. Adachi, J. A. Clark, C. M. Deibel, H. Fujita, Y. Fujita, P. Grabmayr, K. Hatanaka, D. Ishikawa, H. Matsubara, Y. Meada, H. Okamura, K. E. Rehm, Y. Sakemi, Y. Shimizu, H. Shimoda, K. Suda, Y. Tameshige, A. Tamii, and C. Wrede, *Phys. Rev. C* **79**, 021301(R) (2009).
- [49] B. P. Kay, T. Bloxham, S. A. McAllister, J. A. Clark, C. M. Deibel, S. J. Freedman, S. J. Freeman, K. Han, A. M. Howard, A. J. Mitchell, P. D. Parker, J. P. Schiffer, D. K. Sharp, and J. S. Thomas, *Phys. Rev. C* **87**, 011302(R) (2013).
- [50] M. Kortelainen and J. Suhonen, *Journal of Physics G: Nuclear and Particle Physics* **30**, 2003 (2004).
- [51] D. Zinatulina, V. Brudanin, V. Egorov, C. Petitjean, M. Shirchenko, J. Suhonen, and I. Yutlandov, *Phys. Rev. C* **99**, 024327 (2019).
- [52] I. H. Hashim and H. Ejiri, *Frontiers in Astronomy and Space Sciences* **8** (2021), 10.3389/fspas.2021.666383.
- [53] T. Siiskonen, J. Suhonen, V. Kuz'min, and T. Teterova, *Nuclear Physics A* **635**, 446 (1998).
- [54] J. M. R. Fox, C. W. Johnson, and R. N. Perez, *Phys. Rev. C* **101**, 054308 (2020).
- [55] V. Cirigliano, W. Dekens, J. de Vries, M. L. Graesser, E. Mereghetti, S. Pastore, and U. van Kolck, *Phys. Rev. Lett.* **120**, 202001 (2018).
- [56] V. Cirigliano, W. Dekens, J. de Vries, M. Hoferichter, and E. Mereghetti, *Phys. Rev. Lett.* **126**, 172002 (2021).
- [57] V. Cirigliano, W. Dekens, J. de Vries, M. L. Graesser, E. Mereghetti, S. Pastore, M. Piarulli, U. van Kolck, and R. B. Wiringa, *Phys. Rev. C* **100**, 055504 (2019).
- [58] G. Miller and J. E. Spencer, *Ann. Phys. (NY)* **100**, 562 (1976).
- [59] F. Šimkovic, A. Faessler, H. Mütter, V. Rodin, and M. Stauf, *Phys. Rev. C* **79**, 055501 (2009).
- [60] M. Horoi, *Bull. Amer. Phys. Soc.* **61**, 123 (2016).
- [61] V. Kejzlar, L. Neufcourt, W. Nazarewicz, and P.-G. Reinhard, *Journal of Physics G: Nuclear and Particle Physics* **47**, 094001 (2020).
- [62] H. Cramer, *Mathematical Methods of Statistics* (Princeton University Press, Princeton, 1957).