

This is the accepted manuscript made available via CHORUS. The article has been published as:

J/ψ production and elliptic flow in relativistic heavy-ion collisions

Taesoo Song, Che Ming Ko, Su Houng Lee, and Jun Xu

Phys. Rev. C **83**, 014914 — Published 31 January 2011

DOI: [10.1103/PhysRevC.83.014914](https://doi.org/10.1103/PhysRevC.83.014914)

J/ψ production and elliptic flow in relativistic heavy-ion collisions

Taesoo Song,^{1,*} Che Ming Ko,^{2,†} Su Houn Lee,^{3,‡} and Jun Xu^{1,§}

¹*Cyclotron Institute, Texas A&M University, College Station, TX 77843-3366, USA*

²*Cyclotron Institute and Department of Physics and Astronomy,*

Texas A&M University, College Station, TX 77843-3366, USA

³*Institute of Physics and Applied Physics, Yonsei University, Seoul 120-749, Korea*

Using the two-component model for charmonium production, which includes contributions from both initial hard nucleon-nucleon scattering and from regeneration in the quark-gluon plasma, we study the nuclear modification factor R_{AA} and elliptic flow v_2 of J/ψ in relativistic heavy-ion collisions. For the expansion dynamics of produced hot dense matter, we introduce a schematic fireball model with its transverse acceleration determined from the pressure gradient inside the fireball and azimuthally anisotropic expansion parameterized to reproduce measured v_2 of light hadrons. We assume that light hadrons freeze out at the temperature of 120 MeV while charmonia at 160 MeV, similar to the kinetic and chemical freeze-out temperatures in the statistical model, respectively. For the properties of charmonia in the quark-gluon plasma, we use the screening mass between their charm and anticharm quarks and their dissociation cross sections given by the perturbative QCD (pQCD) in the leading order and up to the next-to-leading order, respectively. For the relaxation time of charm and anticharm quarks in the quark-gluon plasma, we also use the one calculated in the leading order pQCD. Modeling the effect of higher-order corrections in pQCD by introducing multiplicative factors to the dissociation cross sections of charmonia and the elastic scattering cross sections of charm and anticharm quarks, we find that this effect is small for the R_{AA} of J/ψ as they suppress the number of initially produced J/ψ but enhance the number of regenerated ones. The higher-order corrections increase, however, the v_2 of J/ψ . Our results suggest that the v_2 of J/ψ can play an important role in discriminating between J/ψ production from initial hard collisions and from regeneration in the quark-gluon plasma.

PACS numbers:

I. INTRODUCTION

One of the signatures suggested for the formation of the quark-gluon plasma (QGP) in relativistic heavy-ion collisions is the suppressed production of J/ψ [1]. This was based on the idea that once the QGP is formed in heavy-ion collisions, the J/ψ produced from initial hard scattering would be dissociated in the QGP because of the color screening between its charm and anticharm quark, reducing thus the number of produced J/ψ . However, recent results from Lattice Quantum Chromodynamics (LQCD) calculations have suggested that J/ψ can exist above the critical temperature for the phase transition between the QGP and the hadron gas (HG) [2, 3]. This thus not only reduces the dissociation of J/ψ but also can lead to the production of J/ψ from the recombination of charm and anticharm quarks in QGP [4, 5]. The latter effect becomes more important when the collision energy is high as more charm and anticharm quarks are produced from initial hard scattering.

Experimentally, suppressed production of J/ψ in heavy-ion collisions compared with that is expected from initial hard scattering has been observed at both Super

Proton Synchrotron (SPS) and Relativistic Heavy Ion Collider (RHIC) [6–9]. Many theoretical explanations have been proposed to understand these experimental observations [10–13]. Among them are the two-component model [11] and the statistical model [13]. In the two-component model, some of initially produced J/ψ were found to survive during the collision if the J/ψ dissociation temperature is sufficiently high [11]. In the statistical model, it is assumed, however, that all initially produced J/ψ are dissociated in QGP and the detected J/ψ are entirely due to regeneration from charm and anticharm quarks in the QGP [13]. Both models have been able to successfully reproduce the measured nuclear modification factor R_{AA} of J/ψ as a function of the number of participants [11, 13, 14] as well as of the transverse momentum of J/ψ [15, 16]. In the two-component model, the number of regenerated J/ψ is smaller than that in the statistical model, because not all charm quarks produced in a collision are thermalized in QGP, the finally produced J/ψ include, however, those produced from initial hard scattering and having survived thermal decay in the produced hot dense matter. Although the transverse momentum of thermally produced J/ψ in the statistical model is smaller than that of J/ψ produced from initial hard scattering, it is enhanced by the transverse flow. Therefore, it is difficult to discriminate the two models based on the number of finally produced J/ψ and their transverse momentum distribution.

Previously, two of us have studied in the two-component model the R_{AA} of J/ψ as a function of the

*Electronic address: songtsoo@yonsei.ac.kr

†Electronic address: ko@comp.tamu.edu

‡Electronic address: suhoun@yonsei.ac.kr

§Electronic address: xujun@comp.tamu.edu

number of participants in heavy-ion collisions [14]. In that study, the initial local temperature of produced QGP was determined from the local entropy density, which was taken to be proportional to the linear combination of the number of participants and the number of binary collisions in that region. The initial fireball thus separates into two regions with temperatures higher and lower than the dissociation temperature of J/ψ . While the survival rate of initially produced J/ψ vanishes in the high temperature region, its value in the low temperature region is finite and depends on the temperature. The different values of the J/ψ survival rate in the two regions were found to give rise to a kink in the dependence of the R_{AA} of J/ψ on the number of participants as seen in the experimental data. In the above mentioned work, both the screening mass between charm and anticharm quark and the relaxation factor of charm quarks in QGP were calculated in the leading order in pQCD, while the dissociation cross sections of charmonia in QGP were calculated up to the next-to-leading order in pQCD. Since the strong coupling constant is found to be large up to around $2T_c$ in LQCD, the QGP remains nonperturbative below this temperature and higher-order corrections in pQCD may not be negligible. The higher-order corrections are expected to enhance in QGP both the thermal decay rate of charmonia, which would decrease the number of initially produced J/ψ , and the relaxation of charm quarks, which would increase the number of regenerated J/ψ . The cancelation of these two effects is probably one of the reasons that both the two-component model and the statistical model can reproduce the measured R_{AA} of J/ψ as a function of the number of participants in heavy-ion collisions.

In the present work, we improve the two-component model in two ways. Firstly, instead of assuming that the transverse acceleration of produced fireball is constant and vanishes at a certain time as in Ref. [14], the acceleration in the present work is obtained from the pressure inside the fireball and continues to be present until the fireball freezes out. Secondly, we take into account the azimuthally anisotropic expansion in noncentral collisions by using a simple parametrization with a transverse-momentum-dependent flow velocity to reproduce the measured v_2 of light hadrons. This schematic fireball expansion model is then applied to study the R_{AA} of J/ψ as a function of both the number of participants and the transverse momentum of J/ψ as well as to study the v_2 of J/ψ . For the chemical and thermal freeze-out temperatures, we follow those used in the statistical model. In this model, the chemical freeze-out occurs at temperature around 160 MeV, after which chemical equilibrium is reached and no additional effective chemical reactions are present and the kinetic freeze-out takes place later when the temperature drops to 120 MeV. Since the kinetic freeze-out depends on the elastic scattering cross sections of hadrons and, unlike light hadrons, the elastic scattering cross sections of J/ψ in hadronic matter are small, we assume that the kinetic freeze-out temper-

ature for J/ψ is the same as the chemical freeze-out temperature. Finally, we take into account the higher-order corrections by multiplying the pQCD cross sections with arbitrary constants to study their effects on the R_{AA} and v_2 of J/ψ . We find that although the fraction of J/ψ produced from regeneration increases with increasing larger higher-order corrections, the resulting R_{AA} of J/ψ remains essentially unchanged. On the other hand, the v_2 of J/ψ at low transverse momentum becomes more negative with larger higher-order corrections. Therefore, studying the v_2 of J/ψ can help discriminate between the two J/ψ production mechanisms of from initial hard scattering and from regeneration in the QGP.

This paper is organized as follows. In Sec. II, we briefly describe the two-component model used in Ref. [14]. We then introduce in Sec. III the schematic model for the expanding fireball produced in heavy-ion collision to obtain the time evolution of the thermal properties as well as the spatial and kinematic information of the fireball for both central and non-central collisions. In Sec. IV, the R_{AA} of J/ψ as a function of the number of participants and the transverse momentum of J/ψ are calculated with the schematic model. Both the Cronin effect for the initial transverse momentum broadening and the leakage effect on initially produced J/ψ are considered. Results on the elliptic flow of J/ψ are given in Sec. V. In Sec. VI, higher-order corrections are introduced by multiplying the pQCD results with constant factors, and their effects on the R_{AA} and v_2 of J/ψ are investigated. Finally, the summary and conclusions are given in Sec. VII. Also, details on the azimuthal angle dependence of the flow velocity are given in the Appendix.

II. THE TWO-COMPONENT MODEL FOR J/ψ PRODUCTION

In the two-component model for J/ψ production in relativistic heavy-ion collisions, some of initially produced J/ψ from hard scattering can survive through the collisions and contribute to the final observed J/ψ in experiments, besides those regenerated from the charm and anticharm quarks in the produced QGP. In this Section, we briefly describe the method used in Ref [14] to evaluate these two contributions.

A. directly produced J/ψ

The number of initially produced J/ψ in a heavy-ion collision at impact parameter b can be determined by using the Glauber model, that is

$$N_{J/\psi}^{AA} = \sigma_{J/\psi}^{NN} A^2 T_{AA}(\vec{b}), \quad (1)$$

where $\sigma_{J/\psi}^{NN}$ is the cross section for J/ψ production in a $p + p$ collision, which has a value of $0.774 \mu\text{b}$ per unit rapidity near midrapidity at collision energy $\sqrt{s} = 200$

GeV [17], and A is the mass number of colliding nuclei. The overlap function T_{AA} in the above equation is defined by

$$T_{AA} = \int d^2s T_A(\vec{s}) T_A(\vec{b} - \vec{s}) \quad (2)$$

in terms of the thickness function

$$T_A(\vec{s}) = \int_{-\infty}^{\infty} dz \rho_A(\vec{s}, z), \quad (3)$$

where $\rho_A(\vec{s}, z)$ is the nucleon distribution in the colliding nucleus.

The survival probability of an initially produced J/ψ in relativistic heavy-ion collisions is given by

$$S_{\text{th}}(\vec{b}, \vec{s}) = \exp \left\{ - \int_{\tau_0}^{\tau_{\text{cf}}} \Gamma(\tau') d\tau' \right\}, \quad (4)$$

where τ_0 is the time the J/ψ is produced from an initial hard scattering and τ_{cf} is the time the J/ψ freezes out from the hot dense matter. The J/ψ survival probability depends on its thermal decay width $\Gamma(T)$ in the produced hot dense matter,

$$\Gamma(T) = \sum_i \int \frac{d^3k}{(2\pi)^3} v_{\text{rel}}(k) n_i(k, T) \sigma_i^{\text{diss}}(k, T). \quad (5)$$

In the above, the summation is over the particle species i , which includes the quarks and gluons in QGP as well as the baryons and mesons in HG, with its density denoted by $n_i(k, T)$; v_{rel} is the relative velocity between the J/ψ and the particle; and σ_i^{diss} is the dissociation cross section of the J/ψ by the particle.

Since some directly produced J/ψ are from the decay of directly produced excited charmonium states χ_c and ψ' , the survival probability of an initially produced J/ψ from thermal decay is the linear combination of the survival rates for J/ψ , χ_c , and ψ' , that is

$$S_{\text{th}}(\vec{b}, \vec{s}) = 0.67 S_{\text{th}}^{J/\psi}(\vec{b}, \vec{s}) + 0.25 S_{\text{th}}^{\chi_c}(\vec{b}, \vec{s}) + 0.08 S_{\text{th}}^{\psi'}(\vec{b}, \vec{s}), \quad (6)$$

where we have assumed that 25% and 8% of J/ψ are, respectively, from the feed-down of χ_c and ψ' as usually adopted in the literature [18]. We note that in Ref [14] the dissociate cross sections of charmonia in QGP were calculated in pQCD up to the next-to-leading order and those in HG were calculated via the factorization formula with the Bethe-Salpeter amplitude for the bound charmonium states [19, 20].

B. regenerated J/ψ

For regenerated J/ψ , their number is similarly determined as in the statistical model, that is

$$N_{\text{reg } J/\psi}^{AA} = \gamma^2 \left\{ n_{J/\psi} S_{\text{th}-H}^{J/\psi} + Br(\chi_c \rightarrow J/\psi) n_{\chi_c} S_{\text{th}-H}^{\chi_c} + Br(\psi' \rightarrow J/\psi) n_{\psi'} S_{\text{th}-H}^{\psi'} \right\} V R. \quad (7)$$

In the above, $n_{J/\psi}$, n_{χ_c} , and $n_{\psi'}$ are, respectively, the equilibrium densities of J/ψ , χ_c , and ψ' at chemical freeze-out when the system has volume V . The charm fugacity parameter γ takes into account the fact that the total charm quark number is not in chemical equilibrium. It is determined by

$$N_{c\bar{c}}^{AA} = \left\{ \frac{1}{2} \gamma n_0 \frac{I_1(\gamma n_0 V)}{I_0(\gamma n_0 V)} + \gamma^2 n_h \right\} V, \quad (8)$$

where I_0 and I_1 are modified Bessel functions with n_0 and n_h being, respectively, the densities of open and hidden charmed hadrons, and $N_{c\bar{c}}^{AA}$ is the total charm quark pairs produced from initial hard scattering given by

$$N_{c\bar{c}}^{AA} = \sigma_{c\bar{c}}^{NN} A^2 T_{AA}(\vec{b}), \quad (9)$$

where $\sigma_{c\bar{c}}^{NN}$ is the cross section for $c\bar{c}$ pair production in a $p + p$ collision, which is 63.7 μb per unit rapidity near midrapidity at $\sqrt{s} = 200$ GeV in perturbative QCD [21]. The factor R in Eq. (7) takes into account the nonequilibrium effect of charm quarks in QGP and is evaluated according to

$$R = 1 - \exp \left\{ - \int_{\tau_0}^{\tau_{\text{QGP}}} d\tau \Gamma_c(T(\tau)) \right\}, \quad (10)$$

where τ_0 and τ_{QGP} are, respectively, the time charm quark pairs are produced from initial hard scattering and the time QGP phase ends, and $\Gamma_c(T)$ is the thermal scattering width of charm quarks in QGP and is calculated in Ref [14] using the leading-order pQCD.

III. A SCHEMATIC MODEL FOR FIREBALL EXPANSION

In Ref. [14], it was assumed for simplicity that the fireball formed in heavy-ion collisions expands transversely with constant acceleration of 0.1 c^2/fm until certain time. Besides, the expansion is assumed to be azimuthally isotropic even in noncentral collisions, neglecting thus the initial elliptic shape of the collision geometry. Since the transverse acceleration of the fireball is expected to be a smooth and continuous function of time and also to be azimuthally anisotropic in non-central collisions, we improve the treatment of the acceleration of the fireball by using the thermodynamical consideration and also introduce a simple parametrization with a transverse-momentum-dependent flow velocity for the anisotropic expansion.

A. Central collisions

To treat dynamically the transverse acceleration of the fireball formed in relativistic heavy-ion collisions, we consider the pressure inside the fireball. In terms of the energy density e , entropy density s , and temperature T of

the fireball as well as the chemical potential μ_i and number density n_i of particle species i in the fireball at a certain time, the pressure p is given by

$$p = sT - e + \sum_i \mu_i n_i. \quad (11)$$

The outward force due to the pressure is then

$$F = pA, \quad (12)$$

where A is the area of the cylindrical side of the fireball. Letting M be the inertia mass against the transverse expansion of the fireball, the transverse acceleration is then

$$a_T = \frac{(p - p_f)A}{M}, \quad (13)$$

where p_f is the pressure of the fireball at freeze-out, if we take into consideration of the fact that the acceleration should vanish at freeze-out, as particles undergo only free streaming afterwards. It is assumed in the above that the fireball is globally in thermal equilibrium. We have thus ignored the fact that in more realistic hydrodynamic approach only local thermal equilibrium is achieved and each fluid element of the fireball moves collectively. Assuming the transverse acceleration a_T acts in the local frame, the transverse velocity of the boundary of the fireball in the laboratory frame at " $i+1$ "th time step can then be obtained from its value at " i "th time step by adding the effect of acceleration relativistically, i.e.,

$$v^{i+1} = \frac{v^i + \Delta v}{1 + v^i \Delta v}, \quad (14)$$

where $\Delta v = a_T^i \Delta t$.

For the thermal quantities in Eq. (11), we use the quasiparticle picture for the QGP, which assumes that strongly interacting massless partons (quarks and gluons) can be substituted by noninteracting massive partons to reproduce the energy density and pressure extracted from LQCD. In this approach, the pressure, energy density, and entropy density are given, respectively, by [22]

$$\begin{aligned} p(T) &= \sum_i \frac{g_i}{6\pi^2} \int_0^\infty dk f_i(T) \frac{k^4}{E_i} - B(T) \\ &\equiv p_0(T) - B(T) \\ e(T) &= \sum_i \frac{g_i}{2\pi^2} \int_0^\infty dk k^2 f_i(T) E_i + B(T) \\ s(T) &= \sum_i \frac{g_i}{2\pi^2 T} \int_0^\infty dk f_i(T) \frac{\frac{4}{3}k^2 + m_i^2(T)}{E_i}. \end{aligned} \quad (15)$$

In the above, $m_i(T)$, g_i , and μ_i are, respectively, the thermal mass, degeneracy factor and chemical potential of parton species i with μ_i obtained from the flavor conservation [23]. The parton distribution function is denoted by

$$f_i(T) = \frac{1}{\exp[(E_i - \mu_i)/T] \pm 1} \quad (16)$$

with the plus and minus signs in the denominator for quarks and gluons, respectively, and $E_i = (m_i^2 + k^2)^{1/2}$. For the bag pressure $B(T)$, it is obtained from the relation $s = \partial p / \partial T$ such that

$$B(T) = B_0 + \sum_i \int_{T_c}^T dT \frac{\partial p_0}{\partial m_i(T)} \frac{\partial m_i(T)}{\partial T}, \quad (17)$$

where $T_c = 170$ MeV is the critical temperature for the QGP to HG phase transition and B_0 is the bag pressure at T_c with its value taken to be 0.1 times the energy density of QGP at T_c to keep the pressure continuous at T_c . From the above thermal quantities, the parton thermal masses can be determined by fitting the QGP equation of state from LQCD as given in details in Ref. [14]. For thermal quantities in the HG phase, they are calculated in the resonance gas model by including all mesons of masses less than 1.5 GeV and all baryons of masses less than 2.0 GeV, and for simplicity they are assumed to be noninteracting. Figure 1 shows the resulting energy density and pressure, divided by temperature to the fourth power, as functions of temperature. We note that the pressure at the thermal freeze-out temperature 120 MeV is $7.75 \times 10^{-5} \text{ GeV}^4$.




FIG. 1: (color online) Energy density (solid line) and pressure (dashed line) divided by temperature to the fourth power in the quasiparticle model for QGP and the resonance gas model for HG.

The inertia mass M in Eq. (13) is a free parameter in our model and is determined by fitting the average transverse momenta of light hadrons at thermal freeze out to the measured data. Assuming particles freeze out at certain proper time, their number is then

$$N = \int d\sigma^\mu J_\mu = \int d\sigma^\mu u_\mu n$$

$$\begin{aligned}
&= \int \frac{d^3p}{(2\pi)^3} \int d\sigma \cdot u e^{-p \cdot u/T} \\
&= \int \frac{d^3p}{(2\pi)^3 E} \int d\sigma \cdot p e^{-p \cdot u/T}, \quad (18)
\end{aligned}$$

where σ^μ and J_μ are the hypersurface vector of the fireball and the current density vector of particles, respectively; u_μ and n are, respectively, the 4-velocity of the fluid element and the number density of particles. In obtaining the last step of the above equation, we have used the Boltzmann distribution for evaluating the number density and the fact that the 4-velocity of a fluid element is equal to the average velocity of particles in the fluid element. Ignoring the transverse boost and keeping only the longitudinal boost, the hypersurface vector becomes

$$\begin{aligned}
d\sigma^\mu &= (\partial\tau/\partial t, \vec{0}, \partial\tau/\partial z) \tau d\eta r dr d\phi \\
&= (\cosh \eta, \vec{0}, \sinh \eta) \tau d\eta r dr d\phi, \quad (19)
\end{aligned}$$

where

$$\eta = \frac{1}{2} \ln \frac{t+z}{t-z} \approx \frac{1}{2} \ln \frac{1+v_z}{1-v_z}.$$

For the 4-velocity u_μ of the fluid element, both boosts are included, with the transverse boost followed by the longitudinal boost, leading to

$$u_\mu = (\cosh \eta \cosh \rho, \cos \phi \sinh \rho, \sin \phi \sinh \rho, \sinh \eta \cosh \rho),$$

where ρ is the transverse rapidity defined as $\rho = \tanh(v_T)$ with v_T being the transverse velocity. The transverse rapidity of the fluid element is assumed to be proportional to the radial distance from the center of the fireball, based on the result from hydrodynamics [24]. For the 4-momentum p_μ of a particle, it is

$$p_\mu = (m_T \cosh y, p_T \cos \varphi, p_T \sin \varphi, m_T \sinh y),$$

where y and $m_T = \sqrt{m^2 + p_T^2}$ are, respectively, the rapidity and the transverse energy. In general, three different azimuthal angles are required: one for the hypersurface vector σ_μ ; one for the 4-velocity u_μ of the fluid element; and the other for the momentum vector p_μ . In central collisions, the azimuthal angle for σ_μ is, however, the same as that for u_μ . This is also the case in non-central collisions if only the radial flow is considered and this is assumed in present study.

Substituting Eqs. (19) into Eq. (18) leads to [25]

$$\begin{aligned}
\frac{dN}{dy dp_T^2} &= \frac{1}{2(2\pi)^3} \int_0^{2\pi} d\varphi \int d\sigma \cdot p e^{-p \cdot u/T} \\
&= \frac{\tau m_T}{2(2\pi)^3} \int dA_T \int_0^{2\pi} d\varphi e^{p_T \sinh \rho \cos(\phi - \varphi)/T} \\
&\quad \times \int_{-\infty}^{\infty} d\eta \cosh(\eta - y) e^{-m_T \cosh \rho \cosh(\eta - y)/T} \\
&= \frac{\tau m_T}{(2\pi)^2} \int dA_T I_0 \left[\frac{p_T \sinh \rho}{T} \right] K_1 \left[\frac{m_T \cosh \rho}{T} \right], \quad (20)
\end{aligned}$$

where A_T is the transverse area, and I_n and K_n are modified Bessel functions. From Eq. (18), the averaged transverse momentum of particles is given by

$$\langle p_T \rangle = \frac{\int dp_T^2 p_T (dN/dy dp_T^2)}{\int dp_T^2 (dN/dy dp_T^2)}. \quad (21)$$

We find that with the inertia mass $M = 570$ GeV, the transverse velocity of fireball at freeze-out is 0.71, and the average transverse momenta of pions, kaons and protons are 462, 668 and 953 MeV/c, respectively, which are comparable to measured values of 451 ± 33 , 670 ± 78 and 949 ± 85 MeV/c for π^+ , K^+ and proton, respectively, in Au+Au collisions as $\sqrt{s} = 200$ GeV and of 0-5% centrality [26]. We note that the measured average transverse momenta of π^- , K^- and antiproton have similar values as those for their antiparticles.




FIG. 2: (color online) Transverse velocity (a) and temperature (b) of fireball in central Au+Au collisions at $\sqrt{s} = 200$ GeV using transverse acceleration from the thermodynamical approach (solid line) or constant acceleration (dotted line) as functions of time.

In Fig. 2(a), we show the transverse velocity of the fireball obtained with the acceleration using the above thermodynamical approach as a function of time and compare it with that using the constant acceleration as in Ref. [14]. The transverse velocity is seen to increase rapidly at first and then slowly in the thermodynamical approach, whereas it increases linearly with time and stays constant after a certain time in the constant acceleration approach. For the temperature of the fireball, shown in Fig. 2(b), it evolves in time similarly in the two cases in spite of their different evolutions in the transverse velocity. In the thermodynamic approach, the QGP phase terminates at $t = 5.4$ fm/c, the mixed phase at $t = 6.9$ fm/c, and the HG phase reaches the freeze-out temperature at $t = 18.7$ fm/c. In the case of constant acceleration, the corresponding times are 5.6 fm/c,

7.0 fm/c, and 18.3 fm/c, respectively, and are thus similar. Therefore, the modified acceleration introduced in the present study is not expected to change the results in Ref. [14] on the participant number dependence of the R_{AA} of J/ψ that was obtained with the constant acceleration.

B. Non-central collisions

In noncentral collisions, the length of the overlapping region of two colliding nuclei is shorter in the direction of impact parameter than in its perpendicular direction on the transverse plane, which are designated, respectively, as the x -direction and the y -direction. The difference in the overlapping lengths along these two directions leads to difference in the pressure gradient in these two directions, and as a result fluid velocities along these two directions are different as well, with the fluid velocity in the x -direction higher than the one in the y -direction. To include such an anisotropic expansion of the fireball, we add a term proportional to the eccentricity of the fireball to the acceleration [Eq. (13)] as follows

$$\begin{aligned} a_x &= a_T(1 + z\epsilon) \\ a_y &= a_T(1 - z\epsilon), \end{aligned} \quad (22)$$

where the eccentricity ϵ is defined as

$$\epsilon = \frac{R_y - R_x}{R_y + R_x},$$

and z is a fitting parameter to take into account the different accelerations in and perpendicular to the reaction plane in non-central collisions. In the above, R_x and R_y are, respectively, the half length of the fireball in the x -direction and in the y -direction on the transverse plane. If the shape of the overlapping region is a perfect ellipse, they then correspond to the semiminor and semimajor of the ellipse, respectively. For simplicity, we transform the initial shape of the fireball into a ellipse by keeping the same transverse area and the same R_x to R_y ratio. The velocities in the x -direction and in the y -direction are then, respectively,

$$\begin{aligned} v_x^{i+1} &= \frac{v_x^i + \Delta v_x}{1 + v_x^i \Delta v_x} \\ v_y^{i+1} &= \frac{v_y^i + \Delta v_y}{1 + v_y^i \Delta v_y}, \end{aligned} \quad (23)$$

where $\Delta v_x = a_x^i \Delta t$ and $\Delta v_y = a_y^i \Delta t$. Assuming that the fireball keeps the elliptic shape on transverse plane during expansion, the fluid velocity, which is assumed to be along the radial direction, as a function of the azimuthal angle ϕ is then

$$v(\phi) = R^3(\phi) \left[\frac{v_x \cos^2 \phi}{R_x^3} + \frac{v_y \sin^2 \phi}{R_y^3} \right], \quad (24)$$

where

$$R(\phi) = \left[\frac{\cos^2 \phi}{R_x^2} + \frac{\sin^2 \phi}{R_y^2} \right]^{-1/2} \quad (25)$$

is the radial distance from the center of the fireball to its surface along the azimuthal angle ϕ . Details in the derivation of the above result are given in the Appendix. For the transverse acceleration a_T in Eq. (22), it is obtained in the same way as in central collisions. The constant z in Eq. (22) is determined by requiring that the final v_2 of light hadrons, given by the expectation value of $\cos(2\phi)$ as

$$\begin{aligned} v_2(p_T) &= \frac{\int d\phi \cos(2\phi) (dN/dy d^2 p_T)}{\int d\phi (dN/dy d^2 p_T)} \\ &= \frac{\int dA_T \cos(2\phi) I_2(p_T \sinh \rho/T) K_1(m_T \cosh \rho/T)}{\int dA_T I_0(p_T \sinh \rho/T) K_1(m_T \cosh \rho/T)}, \end{aligned} \quad (26)$$

reproduces the experimental data.




FIG. 3: (color online) Time evolution of temperature (a), v_x and v_y (b), R_x and R_y (c) for Au+Au collisions at $\sqrt{s} = 200$ GeV and impact parameter $b = 9$ fm.

In Fig. 3, we show the time evolution of the temperature of the fireball and of v_x , v_y , R_x and R_y for Au+Au collisions at $\sqrt{s} = 200$ GeV and impact parameter $b = 9$ fm. These results are obtained with the anisotropic constant $z = 2.2$ in Eq. (22) and the inertia mass

$M = 108 \text{ GeV}/c^2$ in Eq. (13). It is seen that the QGP phase ends at $t = 2.8 \text{ fm}/c$, the mixed phase lasts until $t = 3.8 \text{ fm}/c$, and the freeze-out temperature is reached at $t = 11.1 \text{ fm}/c$. The flow velocities v_x and v_y are $0.51 c$ and $0.27 c$ at the phase transition to HG and reach $0.76 c$ and $0.57 c$ at freeze-out, respectively. We note that the length of R_x becomes slightly longer than that of R_y at freeze-out.




FIG. 4: (color online) Elliptic flow v_2 of pions (solid line), kaons (dashed line), and protons (dotted line) for Au+Au collisions at $\sqrt{s} = 200 \text{ GeV}$ and impact parameter $b = 9 \text{ fm}$ before the high p_T correction (a) and after the correction (b). Experimental data from collisions at 20-60% centrality [27] are shown by rectangles, circles, and triangles for pions, kaons, and protons, respectively.

The elliptic flow v_2 of pions (solid line), kaons (dashed line) and protons (dotted line) at $b = 9 \text{ fm}$, obtained from Eq. (26) and using the results of Fig. 3, are shown in Fig. 4(a). Except in the low p_T region, they are seen to deviate from the experimental data from collisions at 20-60% centrality [27], shown by rectangles, circles, and triangles for pions, kaons, and protons, respectively. The reason for this deviation is because high- p_T particles are not likely to be completely thermalized as assumed in our fireball model. To take into the non-equilibrium effect, we multiply $\exp[-C(p_T/n)]$ to $\Delta v = (v_x - v_y)/2$,

where $C = 1.14 \text{ GeV}^{-1}$ is a fitting parameter and n is the number of constituent quarks in a hadron, as described in the Appendix. Figure 4(b) shows the resulting v_2 of light hadrons after this high p_T correction, and they are seen to reproduce very well the experimental data at all p_T . Since the high p_T correction factor modifies the Boltzmann distribution in Eq. (18), it can be considered as an effective way to include the viscous effect on the fireball expansion. In the quark-counting picture for hadron scattering, the ratio of the baryon to the meson scattering cross sections is $3/2$. This then leads to a smaller viscous effect on baryons than mesons, which is consistent with the quark number dependence in the above high p_T correction factor.




FIG. 5: (color online) Participant number dependence of average transverse momenta of π^+ (solid line), K^+ (dashed line), and proton (dotted line) in Au+Au collisions at $\sqrt{s} = 200 \text{ GeV}$ from the fireball model with corresponding experimental data shown by rectangles, circles and triangles.

The average transverse momenta of pions, kaons, and protons in the case of $b = 9 \text{ fm}$ are 433 , 619 and $874 \text{ MeV}/c$, respectively, which are comparable to measured values of 436 ± 35 , 655 ± 54 , and $901 \pm 88 \text{ MeV}/c$ in collisions at 30-40% centrality as well as 426 ± 35 , 636 ± 54 and $868 \pm 88 \text{ MeV}/c$ in collisions at 40-50% centrality for π^+ , K^+ and proton, respectively [26]. For other values of the impact parameter, it is found that the inertia mass parameterized as $M = 0.36 N_{\text{part}}^{1.24}$ with N_{part} being the number of participants in a collision, works well as shown in Fig. 5, where the participant number dependence of the average transverse momenta of π^+ , K^+ and protons obtained with the parameterized inertia mass are compared with the experimental data.

IV. NUCLEAR MODIFICATION FACTOR OF J/ψ

The nuclear modification factor R_{AA} of J/ψ is a measure of its production in heavy-ion collisions relative to that in $p+p$ collisions multiplied by the number of binary collisions. Its value is less than one if there is a suppression and greater than one if there is an enhancement. Besides effects due to the produced hot dense matter, the R_{AA} of J/ψ is also affected by two other important effects, namely the Cronin effect and the leakage effect, which have been neglected in Ref. [14].

A. the Cronin effect

Charmonium is produced mainly through the gluon-gluon fusion in high-energy nuclear collisions. If a gluon in one nucleus is scattered by the nucleon in the other colliding nucleus before it produces the charmonium, the gluon obtains an additional transverse momentum, and the charmonium produced from the gluons then also acquires this additional transverse momentum. As a result, the distribution of the transverse momentum of J/ψ becomes different from that in $p+p$ collisions, and this is called the Cronin effect. Taking into account of this effect, the transverse momentum distribution of J/ψ in an A+A collision at impact parameter \vec{b} is related to the distribution $f_{p+p}(\vec{p}_T)$ in $p+p$ collisions by

$$f_{A+A}(\vec{p}_T, \vec{b}) = \frac{A^2}{\pi} \int d\vec{s} \frac{T(\vec{s})T_A(\vec{b}-\vec{s})}{\delta p_T^2(\vec{b}, \vec{s})} \times \int d^2\vec{p}_T' \exp \left[-\frac{p_T'^2}{\delta p_T^2(\vec{b}, \vec{s})} \right] f_{p+p}(|\vec{p}_T - \vec{p}_T'|), \quad (27)$$

with

$$\delta p_T^2(\vec{b}, \vec{s}) = \frac{\langle \delta p_T^2 \rangle}{\lambda_{gN}} L_{gN}(\vec{b}, \vec{s})$$

where $\langle \delta p_T^2 \rangle$ is the average of added squared transverse momentum by one gluon-nucleon collision, λ_{gN} is the mean free path of a gluon in uniform nuclear matter, $\langle L_{gN} \rangle$ is the average path length of the gluon in uniform nuclear matter before it produces a charmonium. The prefactor $\langle \delta p_T^2 \rangle / \lambda_{gN}$ is fitted to the experimental data of $p+A$ or $d+A$ collisions and $\langle L_{gN} \rangle$ is calculated with the Glauber model according to

$$\langle L_{gN} \rangle = \frac{\int d^2b d^2s dz_A dz_B L(\vec{b}, \vec{s}, z_A, z_B) K(\vec{b}, \vec{s}, z_A, z_B)}{\int d^2b d^2s dz_A dz_B K(\vec{b}, \vec{s}, z_A, z_B)}, \quad (28)$$

where

$$L(\vec{b}, \vec{s}, z_A, z_B) = \int_{-\infty}^{z_A} dz \frac{\rho_A(\vec{s}, z)}{\rho_o} + \int_{z_B}^{\infty} dz \frac{\rho_B(\vec{b}-\vec{s}, z)}{\rho_o},$$

$$K(\vec{b}, \vec{s}, z_A, z_B) = \rho_A(\vec{s}, z_A) \rho_B(\vec{b}-\vec{s}, z_B) \times \exp \left(-\sigma_{\text{abs}} \left[\int_{z_A}^{\infty} dz \rho_A(\vec{s}, z) + \int_{-\infty}^{z_B} dz \rho_B(\vec{b}-\vec{s}, z) \right] \right).$$

In the above, z_A and z_B are longitudinal positions where J/ψ is produced from the center of nucleus A and from the center of nucleus B, respectively; ρ_A and ρ_B are the distribution functions of nucleons in nucleus A and in nucleus B, respectively, and they are normalized to their mass numbers; ρ_o is the saturation density of nuclear matter. The function $L(\vec{b}, \vec{s}, z_A, z_B)$ is the sum of the path length in nucleus A of a gluon in nucleus B and the path length in nucleus B of a gluon in nucleus A, which correspond to the first term and the second term on the right hand side of $L(\vec{b}, \vec{s}, z_A, z_B)$, respectively. The probability that a charmonium is produced at the position z_A from the center of nucleus A and z_B from the center of nucleus B is given by $K(\vec{b}, \vec{s}, z_A, z_B)$. The exponential factor in $K(\vec{b}, \vec{s}, z_A, z_B)$ is due to the effect of nuclear absorption or nuclear destruction as produced J/ψ can be absorbed or destroyed by nucleons with the absorption cross section σ_{abs} before it gets out of the colliding nuclei. We note that the integration range for the Cronin effect is from $-\infty(\infty)$ to $z_A(z_B)$ and for nuclear absorption from $z_A(z_B)$ to $\infty(-\infty)$ in nucleus A(B), as the Cronin effect is present before the production of the primordial charmonium and the nuclear absorption happens after the creation of the J/ψ . In the present work, L_{gN} is calculated at each point on the transverse plane as follows:

$$L_{gN}(\vec{b}, \vec{s}) = \frac{\int dz_A dz_B L(\vec{b}, \vec{s}, z_A, z_B) K(\vec{b}, \vec{s}, z_A, z_B)}{\int dz_A dz_B K(\vec{b}, \vec{s}, z_A, z_B)}. \quad (29)$$

The Cronin effect thus depends on the transverse position at which the J/ψ is produced.

The average squared transverse momentum $\langle p_T^2 \rangle_{AA}$ of J/ψ in A+A collisions is then modified by the Cronin effect as

$$\langle p_T^2 \rangle_{AA} = \langle p_T^2 \rangle_{pp} + \frac{\langle \delta p_T^2 \rangle}{\lambda_{gN}} \langle L_{gN} \rangle, \quad (30)$$

where $\langle p_T^2 \rangle_{pp}$ is the average squared transverse momentum of charmonium in $p+p$ collisions. For center of mass energy at $\sqrt{s_{NN}} = 200$ GeV, it is known that $f_{p+p}(\vec{p}_T) = \alpha(1 + p_T^2/\beta)^{-6}$ with $\beta = 4.1$ GeV² [17], and we use $\langle \delta p_T^2 \rangle / \lambda_{gN} = 0.1$ GeV²/fm and $\sigma_{\text{abs}} = 1.5$ mb for Au+Au collisions [15].

B. the leakage effect

In Ref. [14], it was assumed that charmonia produced inside the initial fireball of quark-gluon plasma could not get out and charmonia produced in the hadronic corona

outside the quark-gluon plasma could not get into the fireball. In realistic situation, it is possible for charmonia of high p_T to escape from the fireball during its time evolution, leading to the so-called leakage effect. To take into account this effect, we consider the thermal decay width of a charmonium inside the fireball, i.e., its transverse position (x and y) satisfies $[x/R_x(t)]^2 + [y/R_y(t)]^2 < 1$,

$$\Gamma(x, y, t) = \sum_i g_i \int \frac{d^3 k}{(2\pi)^3} v_{\text{rel}}(k) n_i(k, T) \sigma_i^{\text{diss}}(k, T), \quad (31)$$

where i denotes the particle species that dissociates the charmonium and g_i and n_i are its degeneracy and density, respectively; v_{rel} is the relative velocity between the charmonium and the particle; σ_i^{diss} is the dissociation cross section of charmonium by the particle. The charmonium thermal decay width vanishes once it moves out of the fireball when its transverse position satisfies $[x/R_x(t)]^2 + [y/R_y(t)]^2 > 1$. Furthermore, we set the charmonium decay width to infinite if it is produced in a region with temperatures higher than its dissociation temperature.

C. The R_{AA} of J/ψ




FIG. 6: (color online) R_{AA} of J/ψ as a function of the number of participants in Au+Au collisions at $\sqrt{s} = 200$ GeV.

With the two-component model described in Sec. II and the fireball acceleration determined thermodynamically, results for the R_{AA} of J/ψ as a function of the number of participants including both the Cronin and the leakage effect are shown in Fig. 6 by the solid line. Since the Cronin effect only changes the transverse momentum




FIG. 7: (color online) Nuclear modification factor R_{AA} of J/ψ as a function of transverse momentum in collisions of 0-20% (a), 20-40% (b), 40-60% (c) and 60-92% (d) centralities in Au+Au collisions at $\sqrt{s} = 200$ GeV. Dashed and dotted lines are, respectively, the R_{AA} from initially produced J/ψ and regenerated J/ψ , and the solid line is the sum of the two.

of J/ψ , not the number of J/ψ , it does not affect the R_{AA} of J/ψ . The leakage effect, on the other hand, may increase R_{AA} or decrease R_{AA} , depending on whether J/ψ initially produced in the corona can move into the fireball and on whether J/ψ initially produced inside the fireball can escape to the outside. The present results are found to be hardly distinguishable from those obtained in Ref. [14] based on the constant acceleration of the fireball and without including the Cronin and leakage effects as the lifetime of QGP differs very little in the two cases and the leakage effect also turns out to be small.

Figure 7 shows the R_{AA} of J/ψ as a function of transverse momentum obtained with the fireball acceleration calculated thermodynamically. It is seen that the R_{AA} of initially produced J/ψ , given by the dashed line, increases with transverse momentum because of the Cronin and leakage effects. On the contrary, the R_{AA} of regenerated J/ψ decreases with transverse momentum at high p_T as shown by the dotted line.

In Figure 8, we show the participant number dependence of the average squared transverse momentum of J/ψ in Au+Au collisions from the two-component model. Our results reproduce the experimental data for peripheral and mid-central collisions. For central collisions, the two-component model overestimates the measured average squared transverse momentum of J/ψ . One possible reason for this is that we may have overestimated the diffusion of charm quarks in the fireball. The number of charm quarks produced in heavy-ion collision is proportional to the number of binary collisions of nucleons,




FIG. 8: (color online) Average squared transverse momentum of J/ψ in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV from the two-component model (solid line) and experimental data (filled squares).

which is more concentrated in the central region of the fireball. The number of light quarks is, on the other hand, proportional to the number of participants. Since the relative concentration of the number of binary collisions to the number of participants is larger in central collisions than in semi-central or peripheral collisions, charm quarks are initially more concentrated in the central region than light quarks. If the diffusion of charm quarks is slow, they would remain near the central region even at hadronization and their transverse velocity would then be smaller than those of light quarks.

V. ELLIPTIC FLOW OF J/ψ

The elliptic flow v_2 of J/ψ in the two-component model can be different from that of light hadrons as a result of following two effects. Firstly, J/ψ 's are produced from two different mechanisms: the initial hard scattering and the regeneration, while light hadrons are produced only through regeneration. Secondly, the freeze-out temperature for J/ψ could be different from that of light hadrons. Since initially produced J/ψ 's have transverse momenta that are azimuthally isotropic, they are expected to have vanishing v_2 . However, the anisotropic expansion of the fireball could give rise to nonzero v_2 for J/ψ . In non-central collisions, the size of the fireball is shorter in the x -direction than in the y -direction. As a result, the survival rate of J/ψ moving in the x -direction will be higher than that moving in the y -direction. In this case, the v_2 of J/ψ is positive, because the number of J/ψ mov-

ing in the x -direction is more than that moving in the y -direction. On the contrary, the v_2 of regenerated J/ψ is negative at low p_T for the following reason. Assuming particles are locally thermalized, they then have a thermal distribution in the local frame. Since the thermal distribution is peaked at $p^2 = 2T(T + \sqrt{T^2 + m^2})$, where T and m are the temperature and mass of the particle, respectively, more massive particles thus peak at higher momenta. If the thermalized matter is boosted with a certain velocity, the peak is shifted to higher momentum in the laboratory frame. In noncentral collisions, the difference in the length of the overlapping region in the x -direction and the y -direction leads to a difference in the pressure gradient and as a result a difference in fluid velocity along these two directions. Because the length of the overlapping region is shorter in the x -direction, the fluid velocity is higher and the peak of the momentum distribution is more shifted in this direction. As the peak shifts to a higher momentum, the number density at momenta lower than the peak momentum becomes smaller than the number density in the unshifted or less shifted momentum distribution, while the number density at momenta higher than the peak becomes larger. As a result, the number density of J/ψ moving in the x -direction is smaller than that of J/ψ moving in the y -direction at small p_T , and this becomes opposite at high p_T , leading thus to a negative v_2 .




FIG. 9: (color online) Elliptic flow v_2 of J/ψ for Au+Au collisions at $\sqrt{s} = 200$ GeV and $b = 9$ fm in the two-component model. The dashed, dotted and solid lines are, respectively, the v_2 of initially produced, regenerated and total J/ψ .

Figure 9 shows the results from the two-component model for the v_2 of initially produced, regenerated and total J/ψ in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and $b = 9$ fm. These results are obtained with the high- p_T correction to the flow velocity as in the case for light

hadrons. It is seen that the v_2 of initially produced J/ψ is essentially zero, whereas the v_2 of regenerated J/ψ has a minimum value of -0.028 at $p_T = 1.1$ GeV. The v_2 of total J/ψ is slightly negative, because only 12% of J/ψ are from regeneration for collisions at $b = 9$ fm in the two-component model. We note that the v_2 of J/ψ has been recently measured by the PHENIX Collaboration for Au+Au collisions at 20-60% centralities [28, 29] and it has a value of -0.15 ± 0.12 between $p_T = 1$ and 2 GeV, which is much much more negative than our value.

VI. EFFECT OF HIGHER-ORDER PQCD CORRECTIONS

In our two-component model, the screening mass between charm quarks and the thermal decay widths of charmonia as well as the relaxation factor of charm quarks in QGP are calculated in pQCD. Specifically, the dissociation cross section of charmonium used for determining its thermal decay width is calculated up to the next-to-leading order, and the elastic cross section of charm quarks for calculating the relaxation factor, and the screening mass are calculated in the leading order. Since it is known that the pQCD is not reliable up to around $2T_c$, corrections due to higher orders are expected to be important for the temperature regime considered in the present study. The higher-order correction for the dissociation cross sections of charmonia is expected to suppress the survival probability of charmonia in hot dense matter while the correction for the relaxation factor of charm quarks would enhance the regeneration of charmonia. Therefore, there could be a balance between the dissociation and the regeneration of charmonia in calculations including higher-order corrections. To include higher-order corrections, we simply introduce in the present study some multiplication factors to the dissociation cross sections of charmonia and to the elastic cross section of charm quarks as following:

$$\begin{aligned} \sigma'_{J/\psi+q(g) \rightarrow c+\bar{c}+X} &= A\sigma_{J/\psi+q(g) \rightarrow c+\bar{c}+X} \\ \sigma'_{c+q(g) \rightarrow c+q(g)} &= B\sigma_{c+q(g) \rightarrow c+q(g)}, \end{aligned} \quad (32)$$

where A and B are arbitrary constants.

In Fig. 10, the R_{AA} of J/ψ is shown as a function of the number of participants. Solid, dashed, dotted, and dash-dotted lines correspond, respectively, to results obtained in the two-component model with $(A,B)=(1,1)$, $(2,2)$, $(5,4)$, and $(10,6.5)$ for the multiplicative factors used to include high-order effects in pQCD. These results show that higher-order corrections do not change much the value of the R_{AA} of J/ψ even with a large correction factor of ten. This is due to the balance between the dissociation and the regeneration of J/ψ as mentioned previously, with the suppression of J/ψ caused by enhanced dissociation compensated by the enhancement of regenerated J/ψ . However, as the correction factor becomes large, the kink in the R_{AA} curve around $N_{\text{part}} = 170$ becomes less noticeable. To understand this, we note that




FIG. 10: (color online) Nuclear modification factor R_{AA} of J/ψ as a function of the number of participants in Au+Au collisions at $\sqrt{s} = 200$ GeV with different values of the multiplicative parameters $(A,B)=(1,1)$, $(2,2)$, $(5,4)$ and $(10,6.5)$ for including the effect of higher-order corrections in pQCD. Corresponding results are shown by solid, dashed, dotted, and dot-dashed lines, respectively.

the kink in the curve is caused by the discontinuity in the survival probability of J/ψ in the region of the fireball where the temperature is higher than the dissociation temperature and in the region where the temperature is lower than the dissociation temperature. As higher-order corrections become large, the thermal decay width of J/ψ increases and the survival probability of J/ψ in the low temperature region becomes comparable to that in the high temperature region. As a result, the kink disappears when higher-order corrections are large.

Figure 11 shows the fraction of regenerated J/ψ among total J/ψ in Au+Au collisions at $\sqrt{s} = 200$ GeV in the two-component model. Without higher-order corrections (solid line), the regenerated J/ψ is about 31% of total J/ψ produced in central collisions. Including higher-order corrections, the percentage increases to 52, 81 and 96%, respectively, for multiplicative factors $(A,B)=(2,2)$, $(5,4)$ and $(10,6.5)$, shown by dashed, dotted, and dot-dashed lines, respectively. We note that the fraction of regenerated J/ψ is smaller in semi-central and peripheral collisions than in central collisions.

Figure 12 shows the R_{AA} of J/ψ as a function of the transverse momentum in the same collisions at 0-20% (a), 20-40% (b), 40-60% (c) and 60-92% (d) centralities. The lines in each figure correspond, respectively, to multiplicative factors $(A,B)=(1,1)$, $(2,2)$, $(5,4)$ and $(10,6.5)$ for the higher-order corrections. It is seen that the R_{AA} does not change much even with large higher-order corrections as a result of the large transverse flow of regener-




FIG. 11: (color online) Fraction of regenerated J/ψ among total J/ψ in Au+Au collisions at $\sqrt{s} = 200$ GeV. Solid, dashed, dotted, and dot-dashed lines correspond, respectively, to multiplicative factors $(A,B)=(1,1)$, $(2,2)$, $(5,4)$ and $(10,6.5)$ for higher-order corrections.

ated J/ψ . As shown in Fig. 7, the p_T dependence of the R_{AA} of initially produced J/ψ is similar to that of regenerated J/ψ except at high p_T where the R_{AA} of initially produced J/ψ increases while that of regenerated J/ψ decreases. As a result, the R_{AA} at high p_T decreases when higher-order corrections are included, and this seems to contradict with the p_T dependence of measured R_{AA} , which is an increasing function at high p_T . Although higher-order corrections are small at high p_T , they are relatively large at low p_T , and this seems to reproduce the observed decreasing R_{AA} with increasing p_T , when p_T is low, in peripheral collisions as shown in Fig. 12(d). This is mainly due to the fact that in peripheral collisions, the time between the thermalization of fireball and the freeze-out of J/ψ is short, so the transverse flow is not sufficiently developed to give a large transverse flow velocity to the regenerated J/ψ .

In Fig. 13, we show the average squared transverse momentum of J/ψ before and after including higher-order corrections. It is seen that higher-order corrections have very little effect on the results.

Figure 14 shows the v_2 of J/ψ in the same collisions at $b = 9$ fm without and with higher-order corrections. Different from the R_{AA} of J/ψ , the change in v_2 due to higher-order corrections is significant. For increasing large correction factors listed in the above, the fraction of regenerated J/ψ among total J/ψ is 12%, 23%, 42% and 63%, respectively. This can be easily understood from the fact that the v_2 of initially produced J/ψ is very small. In terms of the v_2 of initially produced J/ψ and that of regenerated J/ψ , the v_2 of total J/ψ is given




FIG. 12: (color online) R_{AA} of J/ψ as a function of transverse momentum in Au+Au collisions at $\sqrt{s} = 200$ GeV and centralities of 0-20% (a), 20-40% (b), 40-60% (c) and 60-92% (d). Solid, dashed, dotted, and dot-dashed lines correspond, respectively, to multiplicative factors $(A,B)=(1,1)$, $(2,2)$, $(5,4)$ and $(10,6.5)$ for the higher-order corrections.

by

$$v_2 = \frac{\int \cos(2\varphi) dN}{\int dN} = \frac{\int \cos(2\varphi) dN_R + \int \cos(2\varphi) dN_I}{\int dN_R + \int dN_I} \approx \frac{\int \cos(2\varphi) dN_R}{\int dN_R + \int dN_I} = \frac{\int \cos(2\varphi) dN_R}{\int dN_R} \frac{N_R}{N}, \quad (33)$$

where N , N_I and N_R are, respectively, the total number of J/ψ , the number of initially produced J/ψ and that of regenerated J/ψ . Equation (33) shows that the v_2 of total J/ψ in heavy-ion collisions can be approximated by the v_2 of regenerated J/ψ multiplied by its fraction among total J/ψ . The double-dot-dashed line in Fig. 14 is the v_2 of J/ψ if all J/ψ are produced from regeneration. It is worthwhile to point out that even in this extreme case, the v_2 of J/ψ still only reaches the upper error bar of experimental data, which has a value of -0.03 between $p_T = 1$ and 2 GeV [28, 29]. If the experimental data on v_2 of J/ψ is correct, it then means that the fraction of regenerated J/ψ is not small even in semi-central collisions.

VII. SUMMARY AND CONCLUSIONS

We have investigated in the two-component model the R_{AA} of J/ψ as a function of transverse momentum and the number of participants as well as the v_2 of J/ψ in relativistic heavy-ion collisions. The study was based on a schematic model for the expansion of the fireball formed




FIG. 13: (color online) Average squared transverse momentum of J/ψ in Au+Au collisions at $\sqrt{s} = 200$ GeV. Solid, dashed, dotted, and dot-dashed lines correspond, respectively, to multiplicative factors $(A,B)=(1,1)$, $(2,2)$, $(5,4)$ and $(10,6.5)$ for the higher-order corrections.

in these collisions. In this model, the transverse acceleration of the fireball was assumed to be proportional to the outgoing force given by the product of the pressure of the hot dense matter inside the fireball and the surface area of the fireball, and inversely proportional to an inertia mass, which was taken to be a free parameter to fit the experimental data on the transverse momentum spectra of light hadrons. The anisotropic expansion in non-central collisions was modeled by simply adding or subtracting an additional term proportional to the eccentricity of the fireball to the acceleration in x -direction or y -direction. The coefficient of the additional term was determined such that the v_2 of light hadrons was reproduced after including the high- p_T correction to the flow velocity due to the non-equilibrium effect. Both the Cronin effect due to the initial transverse momentum broadening of gluons in nuclei and the leakage effect due to the escape of initially produced J/ψ from produced hot dense matter on the R_{AA} and v_2 of J/ψ were included. For the kinetic freeze-out temperature of J/ψ , it was taken to be the same as the chemical freeze-out temperature, because of the much smaller elastic cross section of J/ψ than that of light hadrons, which normally has a lower kinetic freeze-out temperature than the chemical freeze-out temperature. For Au+Au collisions at $\sqrt{s} = 200$ GeV, the model was shown to give a good description of participant number dependence of the R_{AA} of J/ψ as in Ref. [14]. It could also describe the increase of J/ψ R_{AA} with increasing transverse momenta as a result of the inclusion of the Cronin and leakage effects in our study.




FIG. 14: (color online) Elliptic flow v_2 of J/ψ in Au+Au collisions at $\sqrt{s} = 200$ GeV and $b = 9$ fm with higher-order corrections. Solid, dashed, dotted, and dot-dashed lines correspond, respectively, to multiplicative factors $(A,B)=(1,1)$, $(2,2)$, $(5,4)$ and $(10,6.5)$ for the higher-order corrections. The double-dot-dashed line is the v_2 of J/ψ if they are totally from regeneration.

For the J/ψ elliptic flow, its value was found to be small as J/ψ was mostly produced from initial hard scattering in the two-component model.

We have also studied the higher-order corrections in pQCD by simply multiplying separate constant factors to the dissociation cross sections of charmonia and to the elastic cross section of charm quarks. The former is related to the thermal decay widths of charmonia and the latter to the relaxation factor of charm quarks. Higher-order corrections were, however, not applied to the screening mass. We found that higher-order corrections enhanced both the dissociation of charmonia and the relaxation of charm quarks. The former decreases the number of initially produced J/ψ while the latter increases the number of regenerated J/ψ . As a result, the total number of J/ψ was found to change very little after including higher-order corrections. However, the fraction of initially produced J/ψ and that of regenerated ones were affected significantly by higher-order corrections with larger higher-order corrections leading to a larger fraction of regenerated J/ψ among total J/ψ . This is the reason why the R_{AA} of J/ψ as a function of the number of participants was insensitive to the higher-order corrections, and both the two-component model and the statistical model can describe successfully the dependence of R_{AA} on the number of participants. Applying same higher-order corrections to the study of the transverse momentum dependence of the R_{AA} of J/ψ , we found that it was also insensitive to the fraction of

regenerated J/ψ among the total J/ψ except at high p_T , as the small transverse momentum of regenerated J/ψ is compensated by the large transverse flow velocity. On the other hand, the v_2 of J/ψ is sensitive to the fraction of regenerated J/ψ because the v_2 of initially produced J/ψ is almost zero. If the elastic cross section of J/ψ in QGP is not small, the initially produced J/ψ can then acquire nonnegligible v_2 through scattering with thermal partons. However, initially produced J/ψ is expected to have a smaller binding energy and larger radius because of the high temperature. The loosely bound J/ψ is thus more likely to be dissociated rather than scattered elastically when colliding with thermal partons. For regenerated J/ψ , they were assumed to be produced from thermalized charm quarks in QGP. It is not clear yet how close charm quarks are from thermal equilibrium in heavy-ion collisions. Recent experimental data on non-photonic single electrons, which are produced through the decay of heavy mesons, suggest that charm quarks are thermalized significantly [30]. If charm quarks are thermalized, resulting thermally produced J/ψ would share their thermal properties and thus acquire appreciable v_2 . Also, if there is an abrupt decrease in the mass of charmonia near T_c as suggested in Ref. [32], the yield of regenerated J/ψ would be enhanced and so is its contribution to v_2 .

We have used in the present study an equation of state with a first-order phase transition between the strongly coupled quark-gluon plasma and the hadronic resonance gas. Recent lattice QCD calculations imply, however, that the QCD phase transition is a crossover in the high temperature and low net baryon density matter produced in heavy-ion collisions at RHIC. Although a crossover transition would increase the duration of the transition between the quark-gluon plasma and the hadronic matter, it is not expected to affect much our results as the quark-gluon plasma has a smaller average volume during

the crossover phase transition than during the first-order phase transition, resulting thus in a small difference in the four volume.

Presently, the v_2 of J/ψ has large error bars in the experimental data. If refined and precise data on the v_2 of J/ψ become available, they are expected to play an important role in discriminating between the two production mechanisms for J/ψ , i.e., initial production and regeneration.

Acknowledgements

This work was supported in part by the U.S. National Science Foundation under Grant No. PHY-0758115, the Welch Foundation under Grant No. A-1358, the Korean Ministry of Education through the BK21 Program, and the Korea Research Foundation under Grant No. KRF-2006-C00011.

Appendix

Let the semimajor and the semiminor of an elliptic fireball produced in heavy-ion collisions be R_y and R_x , respectively, and the fluid velocities along these two axes be v_y and v_x , respectively, at a certain time during the evolution of the fireball. A short time Δt later, the semimajor becomes $R'_y = R_y + v_y \Delta t$ and the semiminor $R'_x = R_x + v_x \Delta t$. Suppose that a certain point on the ellipse, $(x/R_x)^2 + (y/R_y)^2 = 1$, moves to a slightly expanded ellipse, $(x/R'_x)^2 + (y/R'_y)^2 = 1$, in radial direction keeping the same polar angle ϕ . Then the velocity of the point $v(\phi)$ satisfies

$$\frac{\left[\{R(\phi) + v(\phi)\Delta t\} \cos \phi \right]^2}{\left[R_x + v_x \Delta t \right]^2} + \frac{\left[\{R(\phi) + v(\phi)\Delta t\} \sin \phi \right]^2}{\left[R_y + v_y \Delta t \right]^2} = 1, \quad (34)$$

where

$$R(\phi) = \left[\frac{\cos^2 \phi}{R_x^2} + \frac{\sin^2 \phi}{R_y^2} \right]^{-1/2}$$

is the distance from the center of the ellipse to the point on the ellipse along the polar angle ϕ . Expanding Eq. (34) with respect to the infinitesimal time interval Δt gives

$$\frac{\{R(\phi) \cos \phi\}^2}{R_x^2} + \frac{\{R(\phi) \sin \phi\}^2}{R_y^2} + \Delta t \left[v(\phi) \left(\frac{2R(\phi) \cos^2 \phi}{R_x^2} + \frac{2R(\phi) \sin^2 \phi}{R_y^2} \right) - \frac{2R(\phi)^2 v_x \cos^2 \phi}{R_x^3} - \frac{2R(\phi)^2 v_y \sin^2 \phi}{R_y^3} \right] = 1,$$

leading thus to the following expanding velocity of the

surface of the fireball

$$v(\phi) = R^3(\phi) \left[\frac{v_x \cos^2 \phi}{R_x^3} + \frac{v_y \sin^2 \phi}{R_y^3} \right]. \quad (35)$$

Defining $v = (v_x + v_y)/2$ and $\Delta v = (v_x - v_y)/2$, Eq. (35) can be rewritten as

$$v(\phi) = R^3(\phi) \left[\frac{(v + \Delta v) \cos^2 \phi}{R_x^3} + \frac{(v - \Delta v) \sin^2 \phi}{R_y^3} \right]. \quad (36)$$

Because high- p_T particles are unlikely to follow the col-

lective motion, both v and Δv can be functions of p_T . For simplicity, we modify only Δv by multiplying it with the factor $\exp[-C(p_T/n)]$, where C is a fitting parameter [31].

-
- [1] T. Matsui and H. Satz, Phys. Lett. **B178**, 416 (1986).
 - [2] M. Asakawa and T. Hatsuda, Phys. Rev. Lett. **92**, 012001 (2004).
 - [3] S. Datta, F. Karsch, P. Petreczky and I. Wetzorke, Phys. Rev. D **69**, 094507 (2004).
 - [4] P. Braun-Munzinger and J. Stachel, Phys. Lett. B **490**, 196 (2000).
 - [5] R. L. Thews, M. Schroedter and J. Rafelski, Phys. Rev. C **63**, 054905 (2001).
 - [6] M. C. Abreu *et al.* [NA50 Collaboration], Phys. Lett. **B410**, 337 (1997).
 - [7] R. Arnaldi *et al.* [NA60 Collaboration], Nucl. Phys. A **783**, 261 (2007).
 - [8] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **98**, 232301 (2007).
 - [9] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **96**, 012304 (2006).
 - [10] B. Zhang, C. M. Ko, B. A. Li, Z. w. Lin and B. H. Sa, Phys. Rev. C **62**, 054905 (2000).
 - [11] L. Grandchamp and R. Rapp, Nucl. Phys. **A709**, 415 (2002).
 - [12] L. Yan, P. Zhuang and N. Xu, Phys. Rev. Lett. **97**, 232301 (2006).
 - [13] A. Andronic, P. Braun-Munzinger, K. Redlich and J. Stachel, Nucl. Phys. **A789**, 334 (2007).
 - [14] T. Song, W. Park and S. H. Lee, Phys. Rev. C **81**, 034914 (2010).
 - [15] X. Zhao and R. Rapp, Phys. Lett. **B664**, 253 (2008).
 - [16] S. V. Akkelin, P. Braun-Munzinger and Yu. M. Sinyukov, Phys. Rev. C **81**, 034912 (2010).
 - [17] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **98**, 232002 (2007).
 - [18] L. A. Linden Levy [PHENIX Collaboration], J. Phys. G **36**, 064013 (2009).
 - [19] Y. Park, K. I. Kim, T. Song, S. H. Lee and C. Y. Wong, Phys. Rev. C **76**, 044907 (2007).
 - [20] T. Song, Y. Park, S. H. Lee and C. Y. Wong, Phys. Lett. **B659**, 621 (2008).
 - [21] M. Cacciari, P. Nason and R. Vogt, Phys. Rev. Lett. **95**, 122001 (2005).
 - [22] P. Levai and U. W. Heinz, Phys. Rev. C **57**, 1879 (1998).
 - [23] P. Braun-Munzinger, I. Heppe and J. Stachel, Phys. Lett. **B465**, 15 (1999).
 - [24] D. Teaney, J. Lauret and E. V. Shuryak, arXiv:nucl-th/0110037.
 - [25] R. J. Fries, B. Muller, C. Nonaka and S. A. Bass, Phys. Rev. C **68**, 044902 (2003).
 - [26] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. C **69**, 034909 (2004).
 - [27] S. Afanasiev *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **99**, 052301 (2007).
 - [28] C. Silvestre [PHENIX Collaboration], J. Phys. G **35**, 104136 (2008).
 - [29] D. Krieg and M. Bleicher, Eur. Phys. J. A **39**, 1 (2009).
 - [30] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **98**, 172301 (2007).
 - [31] Y. Oh, Z. W. Lin and C. M. Ko, Phys. Rev. C **80**, 064902 (2009).
 - [32] K. Morita and S. H. Lee, Phys. Rev. Lett. **100**, 022301 (2008).