

This is the accepted manuscript made available via CHORUS. The article has been published as:

Exclusive neutral pion electroproduction in the deeply virtual regime

E. Fuchey *et al.* (Jefferson Lab Hall A Collaboration)

Phys. Rev. C **83**, 025201 — Published 2 February 2011

DOI: [10.1103/PhysRevC.83.025201](https://doi.org/10.1103/PhysRevC.83.025201)

Exclusive Neutral Pion Electroproduction in the Deeply Virtual Regime

E. Fuchey,¹ A. Camsonne,^{1,2} C. Muñoz Camacho,^{3,1} M. Mazouz,^{4,5} G. Gavalian,⁶ E. Kuchina,⁷ M. Amarian,⁶ K. A. Aniol,⁸ M. Beaumel,³ H. Benaoum,⁹ P. Bertin,^{1,2} M. Brossard,¹ M. Canan,⁶ J.-P. Chen,² E. Chudakov,² B. Craver,¹⁰ F. Cusanno,¹¹ C.W. de Jager,² A. Deur,² C. Ferdi,¹ R. Feuerbach,² J.-M. Fieschi,¹ S. Frullani,¹¹ M. Garçon,³ F. Garibaldi,¹¹ O. Gayou,¹² R. Gilman,⁷ J. Gomez,² P. Gueye,¹³ P.A.M. Guichon,³ B. Guillon,⁴ O. Hansen,² D. Hayes,⁶ D.W. Higinbotham,² T. Holmstrom,¹⁴ C.E. Hyde,^{6,1} H. Ibrahim,⁶ R. Igarashi,¹⁵ F. Itard,¹ X. Jiang,⁷ H.S. Jo,¹⁶ L.J. Kaufman,¹⁷ A. Kelleher,¹⁴ A. Kolarkar,¹⁸ G. Kumbartzki,⁷ G. Laveissiere,¹ J.J. LeRose,² R. Lindgren,¹⁰ N. Liyanage,¹⁰ H.-J. Lu,¹⁹ D.J. Margaziotis,⁸ Z.-E. Meziani,²⁰ K. McCormick,⁷ R. Michaels,² B. Michel,¹ B. Moffit,¹⁴ P. Monaghan,¹² S. Nanda,² V. Nelyubin,¹⁰ M. Potokar,²¹ Y. Qiang,¹² R.D. Ransome,⁷ J.-S. Réal,⁴ B. Reitz,² Y. Roblin,² J. Roche,² F. Sabatié,³ A. Saha,² S. Sirca,²¹ K. Slifer,¹⁰ P. Solvignon,²⁰ R. Subedi,²² V. Sulkosky,¹⁴ P.E. Ulmer,⁶ E. Voutier,⁴ K. Wang,¹⁰ L.B. Weinstein,⁶ B. Wojtsekhowski,² X. Zheng,²³ and L. Zhu²⁴

(The Jefferson Lab Hall A Collaboration)

¹Clermont Université, Université Blaise Pascal, CNRS/IN2P3,

Laboratoire de Physique Corpusculaire, 63000 Clermont-Ferrand, France

²Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

³CEA Saclay, IRFU/SPhN, Gif-sur-Yvette, 91191, France

⁴LPSC, Université Joseph Fourier, CNRS/IN2P3, INPG, 38026 Grenoble, France

⁵Faculté des sciences de Monastir, 5000 Tunisia

⁶Old Dominion University, Norfolk, Virginia 23508, USA

⁷Rutgers, The State University of New Jersey, Piscataway, New Jersey 08854, USA

⁸California State University, Los Angeles, Los Angeles, California 90032, USA

⁹Syracuse University, Syracuse, New York 13244, USA

¹⁰University of Virginia, Charlottesville, Virginia 22904, USA

¹¹INFN/Sezione Sanità, 00161 Roma, Italy

¹²Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

¹³Hampton University, Hampton, Virginia 23668, USA

¹⁴College of William and Mary, Williamsburg, Virginia 23187, USA

¹⁵University of Saskatchewan, Saskatchewan, SK, Canada, S7N 5C6

¹⁶Institut de Physique Nucléaire CNRS-IN2P3, Orsay, France

¹⁷University of Massachusetts Amherst, Amherst, Massachusetts 01003, USA

¹⁸University of Kentucky, Lexington, Kentucky 40506, USA

¹⁹Department of Modern Physics, University of Science and Technology of China, Hefei 230026, China

²⁰Temple University, Philadelphia, Pennsylvania 19122, USA

²¹Institut Jozef Stefan, University of Ljubljana, Ljubljana, Slovenia

²²Kent State University, Kent, Ohio 44242, USA

²³Argonne National Laboratory, Argonne, Illinois 60439, USA

²⁴University of Illinois, Urbana, Illinois 61801, USA

We present measurements of the $ep \rightarrow ep\pi^0$ cross section extracted at two values of four-momentum transfer $Q^2 = 1.9 \text{ GeV}^2$ and $Q^2 = 2.3 \text{ GeV}^2$ at Jefferson Lab Hall A. The kinematic range allows to study the evolution of the extracted cross section as a function of Q^2 and W . Results are confronted with Regge inspired calculations and GPD predictions. An interpretation of our data within the framework of semi-inclusive deep inelastic scattering is also discussed.

PACS numbers: 13.60.Hb, 13.60.Le, 13.87.Fh, 14.20.Dh, 25.30.Rw

I. INTRODUCTION

The past decade has shown a strong evolution of the study of hadron structure through exclusive processes, allowing access to the three-dimensional structure of hadrons. Exclusive processes include Deeply Virtual Compton Scattering (DVCS) and Deeply Virtual Meson Production (DVMP). This document focuses on the latter, and more precisely on neutral pion production. We present measurements of the differential cross section for the forward exclusive electroproduction reaction

$ep \rightarrow ep\pi^0$, through virtual photo-absorption. A diagram of this process, including definitions of the kinematic variables, is presented in Fig. 1. Results will be presented for four kinematics. Two of them are defined by the same value of $x_{Bj} = 0.36$ and are called Kin2 (at $Q^2 = 1.9 \text{ GeV}^2$) and Kin3 (at $Q^2 = 2.3 \text{ GeV}^2$). The two remaining ones are defined by the same value of $Q^2 = 2.1 \text{ GeV}^2$ and are called KinX2 (at $x_{Bj} = 0.40$) and KinX3 (at $x_{Bj} = 0.33$).

The behavior of the cross section will be compared to different models that are available to describe π^0 electro-




FIG. 1: Diagram of the forward π^0 electroproduction reaction (top), and of the dominant π^0 decay mode (bottom). The kinematic invariants of this reaction are defined as: $Q^2 = -(k - k')^2$, $x_{Bj} = Q^2/(2p \cdot q)$, $t = (q - q')^2$, $W^2 = s = M_p + Q^2(1/x_{Bj} - 1)$, and $t_{min} = \frac{(Q^2 - m_\pi^2)^2}{4s} - (|q^{CM}| - |q'^{CM}|)^2$, with $|q^{CM}|$ and $|q'^{CM}|$ the norms of \vec{q} , \vec{q}' in the $p\pi^0$ final state Center-of-Mass Frame.

production, including the Regge Model and the Generalized Parton Distribution (GPD) framework.

Forward photo-production at asymptotically high energies can be described by Regge theory, which exploits the analytic properties of the scattering amplitude in the limit $t/s \rightarrow 0$ [1]. Previous analyses have applied Regge phenomenology to exclusive photo- and electroproduction in the kinematic range presented here [2, 3]. Recent computations with Regge-inspired models exist for our kinematics. These models include ρ , ω and b meson exchange as well as π^\pm rescattering. Among these, there is the t -channel meson exchange (TME) model by J.M. Laget *et al.*. A brief description of this model has been given in [4], and it is described extensively in [5, 6]. Another Regge-inspired computation by S. Ahmad, G. Goldstein and S. Liuti [7] is available for our kinematics. Recent JLab Hall C experiments studying the Q^2 -dependence of charged pion electroproduction, with a longitudinal-transverse separation were analyzed using the TME formalism [8].

In the Bjorken limit $Q^2 \rightarrow \infty$, and $t/Q^2 \ll 1$ at fixed x_{Bj} , the scattering amplitude is dominated by the leading order (or leading twist) amplitude of GPDs and the pion Distribution Amplitude (DA) [9–11]. The GPDs are light-cone matrix elements of non-local bilinear quark and gluon operators [12–14], unifying the elastic electro-weak form factors with the forward parton distributions of deep inelastic lepton scattering. Cross sections predictions within the GPDs framework exist for the longitudinal cross section σ_L [10, 11]. With the definitions of [9–11], the cross sections are predicted to scale as $\sigma_L \sim Q^{-6}$ and $\sigma_T \sim Q^{-8}$. Thus at sufficiently high Q^2 , σ_L will dominate over σ_T . Beam spin asymmetries for forward exclusive π^0 electroproduction have been measured for $Q^2 > 1 \text{ GeV}^2$ [4]. We performed measurements at two Q^2 values at fixed x_B in order to test these predictions of Q^2 -dependence.

An interpretation of exclusive data with semi-inclusive mechanisms also exists to explain transverse cross sec-




FIG. 2: (Color online) Distribution of $H(e, e'\pi^0)X$ events in the $[x_{Bj}, Q^2]$ plane, for Kin2 ($x_{Bj} = 0.36$, $Q^2 = 1.9 \text{ GeV}^2$) and Kin3 ($x_{Bj} = 0.36$, $Q^2 = 2.3 \text{ GeV}^2$). Events for KinX2 ($x_{Bj} = 0.40$, $Q^2 = 2.1 \text{ GeV}^2$) and KinX3 ($x_{Bj} = 0.33$, $Q^2 = 2.1 \text{ GeV}^2$) are bounded by the two horizontal lines.

tions of hard exclusive charged pion electroproduction [15].

In the second section details of the experiment are presented, while the third section is devoted to the calibration of the calorimeter. The formalism of π^0 electroproduction by Drechsel and Tiator [16] is presented in the fourth section with a special emphasis on the expressions for the hadronic tensors. The fifth section is devoted to the extraction of the cross sections and the sixth and seventh sections to the radiative corrections and the evaluation of the systematic errors. Finally, our results are presented in section eight, with a discussion and conclusions in sections nine and ten.

II. EXPERIMENT

The present data were acquired as part of Jefferson Lab Hall A experiment E00-110 [17]. Additional details about the experimental configuration, calibrations, and analysis can be found in [18, 19]. This paper reports on the analysis of the triple coincidence $H(e, e'\gamma\gamma)X$ events. A 5.75 GeV electron beam was incident on a 15 cm liquid hydrogen target, for a typical luminosity of $10^{37} \text{ cm}^{-2} \text{ s}^{-1}$. Electrons were detected in a High Resolution Spectrometer (HRS), photons in a 132 element PbF_2 calorimeter, each measuring $3 \times 3 \text{ cm}^2 \times 20X_0$. The high resolution allows to accurately define:

- the virtual photon, having the kinematics centered at a fixed $x_{Bj} = 0.36$ and two values of $Q^2 = 1.9$ and 2.3 GeV^2 , as shown in Fig. 2.
- the real photon momentum unit vector, thanks to the vertex resolution of the HRS, and the position resolution of the electromagnetic calorimeter.

The validation threshold for the data acquisition trigger was set to about 1 GeV for each photon cluster. For

the exclusive $\pi^0 \rightarrow \gamma\gamma$ events, the minimum distance between the centroids of the two clusters that guarantees separation is about 10 cm. This is achieved by the minimal opening angle $\approx 2m_\pi/E_\pi$ and the distance from the center of the target to the calorimeter front face $L = 110$ cm. The achieved coincidence resolving time between the scattered electron and either photon cluster is 0.6 ns, rms. Fig. 2 shows the distribution of $H(e, e'\pi^0)X$ events in the $[x_{Bj}, Q^2]$ plane, for missing mass squared $M_X^2 = (q + p - q')^2 \leq 1.15 \text{ GeV}^2$.

The analysis relies only on two specific qualities of the experiment:

- Thanks to the resolution of the spectrometer and the calorimeter, one can use the missing-mass squared to ensure exclusivity. The exclusive sample is selected by putting a cut on the missing-mass squared at the proton plus the pion mass squared.
- For exclusive events, the reconstruction of the invariant momentum transfer t and t_{min} relies on the positions of the reconstructed photons, of which the resolution is better than that of the energy. From this, a resolution in t better than that in the energy is obtained. All data are presented as a function of $t_{min} - t$, which is directly linked to the angle of the pion production relative to the virtual photon direction in the center of mass θ_π^{CM} : $t_{min} - t = 2q^{CM}q'^{CM}(1 - \cos\theta_\pi^{CM})$.

In the $ep \rightarrow e'\gamma_1\gamma_2 X$ reaction, there are six four-vectors, equivalent to 24 independent kinematic variables. The measured four-vectors k , p , and k' , and four-momentum conservation, reduce the number of independent variables to eight. The measurement of the two directional vectors $\hat{k}(\gamma_1) = \vec{q}_1/q_1$ and $\hat{k}(\gamma_2) = \vec{q}_2/q_2$ from the target vertex (reconstructed by the HRS) to the two cluster positions in the calorimeter provides four more kinematic constraints. Finally, the hypothesis that the observed calorimeter showers are due to photons ($m_{q_1} = m_{q_2} = 0$) provides two more kinematic constraints. The remaining two unknowns, which we express as $m_{\gamma\gamma}^2 = (q_1 + q_2)^2$ and M_X^2 , are determined by the previous constraints plus the energy of the two photons. Fig. 3 displays the distribution of the $H(e, e'\gamma\gamma)X$ events in the $[M_X^2, m_{\gamma\gamma}]$ plane, for Kin3. The upper left panel of this figure shows a clear correlation between the two variables in the exclusive region ($M_X^2 \simeq M_p^2$). This is a consequence of resolution fluctuations in the energies E_1 and E_2 of the two photons issued from a π^0 , which correlate fluctuations in M_X^2 and $m_{\gamma\gamma}$. The missing mass in the right-hand panels is obtained by an empirical adjustment:

$$M_X^2|_{Corr} = M_X^2|_{Raw} + C \times (m_{\gamma\gamma} - m_\pi), \quad (1)$$

with $C = 13 \text{ GeV}$.

This transformation produces a noticeable improvement in the M_X^2 distribution (lower right panel of Fig. 3).




FIG. 3: (a), (b): Distributions of $H(e, e'\gamma\gamma)X$ events within cuts in $[M_X^2, m_{\gamma\gamma}]$ plane for Kin3. (a): Raw distribution showing a clear correlation between these two variables. (b): The same distribution after a rotation around (M_p^2, m_{π^0}) to improve the M_X^2 resolution. (c), (d): Projections on the M_X^2 axis of the $[M_X^2, m_{\gamma\gamma}]$ distributions shown respectively in (a), (b). The lower right panel shows that the resolution is indeed improved by the rotation.

III. CALIBRATION

We performed elastic $H(e, e'_{Calo} p_{HRS})$ calibrations at the beginning, middle, and end of the experiment [20]. The calorimeter was retracted to a position at 5.5 m from the target, in order to optimize the electron coverage in the calorimeter with the proton acceptance of the HRS. These data were used for the block calibration. After calibration the calorimeter energy resolution was observed to be 2.4 % at 4.2 GeV with a position resolution of 2 mm at 110 cm from the target. The elastic data also provided a consistency check on the efficiency of the detectors and all associated electronics from the observation that the elastic cross section agreed with the Kelly form-factor parametrization [21] at the 1.1 % level. During the experiment, the light output from the PbF_2 blocks decreased by up to 20 %, strongly correlated with the distance of the blocks from the beam line. We attribute this to radiation damage of the blocks. In addition, seven blocks, at random positions, showed much higher radiation damage. One explanation could be a poorer crystal quality of those crystals. We adjusted the calibration of each block, assuming an independent linear dose *versus* attenuation curve. In addition to radiation damage, each crystal received a pile-up of low-energy photons in random coincidence, resulting in a degradation of the energy resolution, and in a shift in the calibration as a function of its distance to the beam line. This effect was taken




FIG. 4: Projection on the calorimeter of the virtual photons γ^* within cuts for Kin3. Also shown is the block relabeling used for the calorimeter calibration described in the text. The calorimeter is viewed from the rear, with the downstream beam passing to the right.

into account through successive steps:

- For each block the position of the reconstructed missing-mass squared peak was centered at M_p^2 through an energy calibration of the experimental data;
- A GEANT simulation generated a sharper resolution in missing mass than the experimental data for each calorimeter block. For each block the energy of the simulation was calibrated together with a simultaneous energy smearing, in order to center the reconstructed missing-mass peak position at M_p^2 , and to equate the resolution of the simulation to that of the experimental data.

These calibrations are explained in the following paragraphs.

We consider only the 90 blocks of the inner calorimeter (see Fig. 4 for the labeling), indexed by μ . We will assume that the energy of the photon is driven by the block where the shower makes the largest energy deposit. The 90 distributions of missing-mass squared $(M_X^2)_\mu^i = (k + P - k' - q_\mu - q_\nu)_i^2 = (E_X^2)_i - (P_X^2)_i$ are built with all events i involving block μ . Note that for each event i , the reconstructed missing-mass squared appears in two distributions. To compare these distributions, two estimators are constructed: the mean $\langle M_X^2 \rangle_\mu$ and the sigma σ_μ of a Gaussian fitted to these distributions, over a limited range ($0.62 \text{ GeV}^2 < (M_X^2)_\mu < 1.09 \text{ GeV}^2$). The calorimeter is calibrated using

$$\Delta M_X^2 = -2\Delta q_\mu \left(E_X - \frac{\vec{P}_X \cdot \vec{q}_\mu}{|q_\mu|} \right), \quad (2)$$

with $\Delta M_X^2 = \langle (M_X^2)_\mu \rangle - M_p^2$. Neglecting the P_X term compared to E_X between the parentheses, we obtain an energy correction:

$$q_\mu^i \rightarrow q_\mu^i + \Delta q_\mu^i = q_\mu^i + \frac{\Delta M_X^2}{2(E_X)_i}. \quad (3)$$

We recall here that each event involves two blocks. The reconstructed missing mass of one block is then influenced by contributions from all other blocks. Because of this, several iterations are necessary.

Then, the missing-mass distribution of each block for simulated events is adjusted to get the same missing-mass position and resolution as the experimental missing-mass distribution. The missing-mass cut applied to ensure exclusivity is the same for simulation and data, and if the resolution is better for simulation, applying such a cut will remove more experimental events than simulation particularly near the beam where the noise degrades the experimental resolution. This gives a spurious contribution to the $\cos \phi_\pi$ term which has to be removed by smearing the simulation resolution. To this purpose, the momentum of each event i at the n^{th} iteration contributing to the M_X^2 distribution of the block μ is changed from $(q_\mu^i)_{n-1}$ to $(q_\mu^i)_n$ with a sampling from a gaussian distribution:

$$(\vec{q}_\mu)_n^i = \frac{(\vec{q}_\mu)_{n-1}^i}{|q_\mu|_{n-1}^i} \text{Gauss} \left((q_\mu)_n^i, \frac{\Delta \sigma_\mu}{\sqrt{2}} \right) \quad (4)$$

where

$$\Delta \sigma_\mu = \sqrt{(\sigma_\mu)_{\text{data}}^2 - (\sigma_\mu)_{\text{simu}}^2}, \quad (\sigma_\mu)_{\text{data}} > (\sigma_\mu)_{\text{simu}} \quad (5)$$

$$\Delta \sigma_\mu = 0, \quad (\sigma_\mu)_{\text{data}} < (\sigma_\mu)_{\text{simu}} \quad (6)$$

and $(q_\mu)_n^i$ is given by equation (3), except we put $\Delta M_X^2 = (\langle (M_X^2)_\mu \rangle_{\text{simu}} - \langle (M_X^2)_\mu \rangle_{\text{data}})/2$ in this case. The factor 2 in the denominator of ΔM_X^2 is used to ensure a smooth convergence.

	$\langle m - m_{\pi^0} \rangle \text{ (GeV)}$	$\sqrt{\langle (m - m_{\pi^0})^2 \rangle} \text{ (GeV)}$
KIN 3		
data	-0.00081	0.0088
simulation	+0.00072	0.0089
KIN 2		
data	-0.00017	0.0079
simulation	+0.00191	0.0085

TABLE I: Mean deviation and resolution width of the $\pi^0 \rightarrow \gamma\gamma$ reconstruction of the data and simulation. Events are selected by $M_X^2 < 1.15 \text{ GeV}^2$ and calorimeter threshold $E_{Thr} = 1.0 \text{ GeV}$.

The results of these iterations are shown in Fig. 5. Fig. 6 and Table I illustrate the quality of the final calibration adjustments. The calibration of the missing-mass squared was cross-checked by comparing the invariant-mass distribution of both photons in each event. Table I lists the mean values of these distributions with respect to the pion mass, and their resolution. The agreement of the calibration with the data is at the 1.9 MeV level, while the widths of these distributions agree to better than 1 MeV.




FIG. 5: (Color online) Different iterations of the calibration, for Kin2. The differences, between simulation and data, of the missing-mass peak position (a) and resolution (b) are shown before calibration (crosses), after calibration (open circles), and at a random iteration during calibration (asterisks).




FIG. 6: (Color online) (a): Raw $H(e, e'\pi^0)X$ missing-mass distribution for Kin3 (solid histogram), compared to the simulation (dashed histogram), and the difference between the two (dotted histogram). (b): $H(e, e'\pi^0)X$ missing-mass distribution at different values for the calorimeter threshold, corrected with a factor $1/(1 - 2(E_{\text{thr}}/|p_{\pi}|))$. This correction adds to the distribution all π^0 events missed because of the threshold value.

IV. CROSS-SECTION ANALYSIS

In order to extract the differential cross section, it is advantageous to incorporate all model-independent kinematic dependences of the differential cross section into the experimental simulation. To this end, we express the differential cross section in terms of structure functions as described in the paper of Drechsel and Tiator [16] directly related to bilinear combinations of the CGLN helicity amplitudes [22]. We define the differential phase-space elements $d^3\Phi_e = dQ^2 dx_{Bj} d\phi_e$ and $d^5\Phi = d^3\Phi_e d[t_{\min} - t] d\phi_\pi$ and the equivalent real photon

energy in the CM frame $k_\gamma^{CM} = (W^2 - M_p^2)/2W$. Here $t_{\min} = \frac{(Q^2 - m_\pi^2)^2}{4s} - (|q^{CM}| - |q'^{CM}|)^2$ with $|q^{CM}|$ and $|q'^{CM}|$ the norms of \vec{q} , \vec{q}' in the Center-of-Mass Frame.

All these quantities are defined using the convention of Drechsel and Tiator [16]:

- \hat{z} axis along the virtual photon
- $\hat{y} = (\hat{k}_i \wedge \hat{k}_f) / \sin \theta_e$ orthogonal to the leptonic plane
- $\hat{x} = \hat{y} \wedge \hat{z}$

To lowest order in the fine structure constant α , the differential cross section for an electron of helicity h is:

$$\frac{d^5\sigma(h)}{d^5\Phi} = \Gamma \frac{d^2\sigma_v(h)}{dtd\phi_\pi} \quad (7)$$

$$\Gamma = \frac{\alpha}{2\pi^2} \frac{k'}{k} \frac{k_\gamma}{Q^2} \frac{1}{1 - \epsilon} \quad (8)$$

with $k_\gamma = (W^2 - M_p^2)/2M_p$ and k and k' the energies of the incident and scattered electron, respectively. The virtual photo-absorption cross section is expanded as

$$\begin{aligned} \frac{d^2\sigma_v(h)}{dtd\phi_\pi} = & \frac{1}{2q^{CM}k_\gamma^{CM}} \{ R_T + \epsilon_L R_L + \epsilon R_{TT} \cos 2\phi_\pi \\ & + \sqrt{2\epsilon_L(1 + \epsilon)} R_{TL} \cos \phi_\pi \\ & + h \sqrt{2\epsilon_L(1 - \epsilon)} R_{TL'} \sin \phi_\pi \} \end{aligned} \quad (9)$$

where $q^{CM} = |\vec{q}| \times M_p / W$ is the CM virtual photon three-momentum, $\epsilon = 1/[1 + 2(q^2/Q^2) \tan^2 \theta_e/2]$ is the degree of linear polarization of the virtual photons, and $\epsilon_L/\epsilon = 4M_p^2 x_{Bj}^2 / Q^2$. The response functions are defined as functions of the usual hadronic tensor $W^{\mu\nu}$:

$$R_T = \frac{W_{xx} + W_{yy}}{2} \quad (10)$$

$$R_L = W_{zz} \quad (11)$$

$$\cos \phi_\pi R_{TL} = -\Re W_{xz} \quad (12)$$

$$\sin \phi_\pi R_{TL'} = -\Im W_{yz} \quad (13)$$

$$\cos 2\phi_\pi R_{TT} = \frac{W_{xx} - W_{yy}}{2} \quad (14)$$

The interference terms R_{TL} and $R_{TL'}$ have a leading $\sin \theta_\pi^{CM}$ dependence, and the linear polarization interference term R_{TT} has a leading $\sin^2 \theta_\pi^{CM}$ dependence. For this reason, we define reduced structure functions r_Λ , which remove this phase-space dependence, which are directly related to bilinear combinations of the CGLN helicity amplitudes F_i [22]:

$$\begin{pmatrix} r_{TL} \\ r_{TL'} \end{pmatrix} = \frac{1}{\sin \theta_\pi^{CM}} \begin{pmatrix} R_{TL} \\ R_{TL'} \end{pmatrix} \quad (15)$$

$$r_{TT} = \frac{R_{TT}}{\sin^2 \theta_\pi^{CM}} \quad (16)$$

$$r_L = R_L \quad (17)$$

$$r_T = R_T \quad (18)$$

Since our kinematics cover a wide range in x_{Bj} as well as in Q^2 , we also have to include the Q^2 and the W dependence of the hadronic tensor $(W_{xx} + W_{yy})/2 + \epsilon_L W_{zz} = r_T + \epsilon_L r_L$. We perform a preliminary extraction of the cross section on the kinematic points Kin2 and Kin3 (respectively KinX2 and KinX3) to get an estimate of the Q^2 (respectively W) dependence of the hadronic tensor. The extracted Q^2 and W dependences are then introduced explicitly in the formalism to perform a second “definitive” extraction. The dependence is modeled in the form $(Q^2)^n$ and W^δ . With the first iteration, the cross sections changed by 3%, but with a second iteration the cross sections changed only by 0.3%.

The results will be presented as four separated cross sections following the usual decomposition found in the literature:

$$\begin{aligned} \frac{d^2\sigma_v}{dt d\phi_\pi} = & \frac{1}{2\pi} \left\{ \frac{d\sigma_T}{dt} + \epsilon_L \frac{d\sigma_L}{dt} \right. \\ & + \sqrt{2\epsilon_L(1+\epsilon)} \frac{d\sigma_{TL}}{dt} \cos\phi_\pi \\ & + \epsilon \frac{d\sigma_{TT}}{dt} \cos 2\phi_\pi \\ & \left. + h\sqrt{2\epsilon_L(1-\epsilon)} \frac{d\sigma_{TL'}}{dt} \sin\phi_\pi \right\}. \end{aligned} \quad (19)$$

V. EXTRACTION

We define a compact notation that summarizes (9) in the form:

$$\frac{d^5\sigma}{d^5\Phi} = \sum_{\Lambda} \frac{d^3\Gamma_{\Lambda}}{d^3\Phi_e} r_{\Lambda} = \sum_{\Lambda} \mathcal{F}_{\Lambda}(x_v) r_{\Lambda} \quad (20)$$

with $\mathcal{F}_{\Lambda}(x_v)$ containing all the kinematic dependence, $\Lambda \in \{T+\epsilon_L L, TL, TT, TL'\}$ and x_v summarizing all variables k, Q^2, x_{Bj}, W, t , considered at the vertex. $T + \epsilon_L L$ reflects the fact that we used only one incident energy and consequently, we were not able to disentangle $d\sigma_T$ and $d\sigma_L$. This notation will be convenient to use for the presentation of the extraction process.

The experimental data used for the analysis have the kinematical coverage shown in Fig. 2. The analysis includes a complete simulation of the resolution and acceptance of the HRS, the external and internal radiative effects on the incident and scattered electron, and a GEANT based simulation of the acceptance and response of the PbF₂ array. Simulation events are generated uniformly in the target vertex v along the beam line, and uniformly in a phase space $\Delta^5\Phi$. This results in well defined values of θ_π^{CM} in each bin. The Δt bins are the same in the generation and experimental phase spaces, but resolution and radiative effects can cause migration




FIG. 7: (Color online) Raw $H(e, e'\gamma\gamma)X$ distribution in the $[t_{min} - t, Q^2]$ plane with cuts for Kin3. The vertical lines delimit the bins we chose in $t_{min} - t$ for our analysis. Superimposed is the $(t_{min} - t)$ resolution for each alternate bin, showing that each bin is larger than the resolution.

of events from one bin to one of its neighbors (Fig. 7). Rather than extracting average cross sections in the experimental bins, we use the simulation and the theoretical form of eq. (20) to directly extract differential cross sections from the experimental yields.

We divide the acceptance into 24 equal bins in $\phi_\pi \in [0, 2\pi]$ and 8 bins in $t_{min} - t \in [0, 0.3]$ GeV² for both the helicity dependent and independent parts of the cross section. A bin j_d in the kinematic variables reconstructed by the detector is defined by the limits $\phi_\pi \in [\phi(j_d), \phi(j_d) + \Delta\phi(j_d)]$, $(t_{min} - t) \in [(t_{min} - t)(j_d), (t_{min} - t)(j_d) + \Delta(t_{min} - t)(j_d)]$, etc. The statistics $\Delta N(j_d)$ in a bin j_d are determined by the physical cross section at the vertex convoluted with the detector response:

$$\begin{aligned} \Delta N(j_d) = & \mathcal{L}u \times \int_{\Delta x_d} dx_d \int_{\Delta x_v} dx_v \mathcal{R}(x_d, x_v) \sum_{\Lambda} \mathcal{F}_{\Lambda}(x_v) r_{\Lambda} \end{aligned} \quad (21)$$

where:

- x_v summarizes the reaction vertex variables,
- x_d summarizes the reaction vertex variables as reconstructed in the detector,
- Δx_d summarizes the range of integration for bin j_d ,
- Δx_v summarizes the range of integration for all bins j_v ,
- $\mathcal{L}u$ is the integrated luminosity,
- $\mathcal{R}(x_d, x_v)$ is the probability distribution for an event originating at the vertex with kinematics x_v to be reconstructed by the detector with vertex kinematics x_d . This expresses the effects of detector resolution, internal and external radiation, detector efficiency, and anything else that could migrate

events from vertex kinematics x_v to the detector kinematics x_d .

For the analysis and simulation, the integral is split into a sum over the bins Δx_v in the kinematic variables at the reaction vertex:

$$\Delta N(j_d) = \mathcal{L}u \int_{\Delta x_d} dx_d \sum_{j_v} \int_{\Delta x_v \in \text{bin } j_v} dx_v \mathcal{R}(x_d, x_v) \sum_{\Lambda} \mathcal{F}_{\Lambda}(x_v) r_{\Lambda} \quad (22)$$

Because the functions $\mathcal{F}_{\Lambda}(x_v)$ contain the main part of the dependence on the variables at the vertex, the quantity r_{Λ} in a bin Δx_v will be assimilated to its average $\langle r_{\Lambda} \rangle_{x_v} \equiv r_{v,\Lambda}$ in this bin. Then, the last equation can be summarized in a vector notation:

$$\Delta N(j_d) = \sum_{j_v} K_{j_d, j_v}^{\Lambda} r_{j_v, \Lambda} \quad (23)$$

with:

$$K_{j_d, j_v}^{\Lambda} = \mathcal{L}u \int_{\Delta x_d} \int_{\Delta x_v \in \text{bin } j_v} \mathcal{R}(x_d, x_v) \mathcal{F}_{\Lambda}(x_v) dx_d dx_v. \quad (24)$$

We then replace the integration by a summation over the simulated events i :

$$K_{j_d, j_v}^{\Lambda} = \mathcal{L}u \sum_{i \in \{j_v, j_d\}} \frac{\mathcal{F}_{\Lambda}(x_v)}{N_{\text{gen}}} \Delta^5 \Phi. \quad (25)$$

where the sum is over events originating in vertex bin j_v and reconstructed in bin j_d . N_{gen} is the number of events generated in the simulation, $\Delta^5 \Phi$ is the total phase space factor. The matrices K_{j_d, j_v}^{Λ} are constructed from simulation events, summed over all events within cuts. We define $N_d = N^+ + N^-$ with N^+ (N^-) the number of counts within cuts with positive (negative) electron helicity. The cuts are the same for simulation and data (Table II). The cuts and the corrections are summarized in Tables II and III, respectively.

A χ^2 is built, assuming that the statistical error on the simulation is much smaller than the statistical error of the data:

$$\chi^2 = \sum_{j_d} \frac{\left(N_d - \sum_{j_v} K_{j_d, j_v}^{\Lambda} r_{j_v, \Lambda} \right)^2}{N_d} \quad (26)$$

The minimization of χ^2 with respect to the unknown quantities $r_{j_v, \Lambda}$ results in a linear system from which the $r_{j_v, \Lambda}$ are extracted. To be fully consistent, one of the two quantities in the numerator has to be corrected for some instrumental systematic effects (Table III). Note that all vertex bins populate experimental bins, but the detector bin at the largest experimental bin in $(t_{\min} - t)$ can receive contributions from larger values of $(t_{\min} - t)$, not generated in the simulation. Hence, although we extract an $r_{j_v, \Lambda}$ value for the last bin, we do not include it in our

Spectrometer Cuts	
$-6.0 \text{ cm} < v < +7.5 \text{ cm}$	
$ x_{\text{HRS plane}} < 3.5 \text{ cm}$ (Horizontal collimator)	
$ y_{\text{HRS plane}} < 7.0 \text{ cm}$ (Vertical collimator)	
$ k' - p_{\text{HRS}} /p_{\text{HRS}} < 4.5\%$	
$r > +0.005 \text{ m}$	
Calorimeter Cuts	
$-15.0 \text{ cm} < x_{\text{calo}} < +12.0 \text{ cm}$	
$ y_{\text{calo}} < 15.0 \text{ cm}$	
Physics Cuts	
$105 \text{ MeV} < m_{\gamma\gamma} < 165 \text{ MeV}$	

TABLE II: Cuts applied in the primary extraction. r is the value of the so-called r -function. The r -function defines the distance of the particle from the acceptance bound, and is positive (negative) if the particle is in (out of) the acceptance [23]. The M_X^2 and E_{Thr} optimizations are presented in Table IV.

Correction	Kin3	Kin2
Multi-tracks in HRS	1.079	1.099
Triple Cluster in Calo	1.035	1.020
Radiative correction	0.91 ± 0.02	0.91 ± 0.02

TABLE III: Correction factors applied in the data analysis. The radiative correction factor is the combination of the virtual radiative correction factors (vertex renormalization and vacuum polarization) and the cut-off independent real radiation effects (Section VI).

results, its role is only to populate the lower $(t_{\min} - t)$ bins.

The average values of the kinematic variables Q^2 , ϵ , x_{Bj} , W , t , t_{\min} , *etc.* in a bin at the vertex are:

$$\bar{x}_{j_v} = \frac{\sum_{i \in \Delta x_v} x_v K_{j_d, j_v}^{\Lambda} r_{j_v, \Lambda}}{\sum_{i \in \Delta x_v} K_{j_d, j_v}^{\Lambda} r_{j_v, \Lambda}}. \quad (27)$$

Because the $r_{j_v, \Lambda}$ are by construction constant over the bin Δx_v and the integrals of \mathcal{F}_{TL} , \mathcal{F}_{TT} , and $\mathcal{F}_{TL'}$ cancel when integrating over ϕ_{π} , we can write:

$$\bar{x}_{j_v} = \frac{\sum_{i \in \Delta x_v} x_v K_{j_d, j_v}^{T+\epsilon_L L}}{\sum_{i \in \Delta x_v} K_{j_d, j_v}^{T+\epsilon_L L}}. \quad (28)$$

These values are summarized in Table VII for quantities independent of the $(t_{\min} - t)$ bin and in Table VIII for quantities depending of the $(t_{\min} - t)$ bin.

Finally, the cross sections at the point \bar{x}_{j_v} in a bin j_v are obtained by:

$$\frac{d\sigma_{\Lambda}}{dt} = \mathcal{F}_{\Lambda}(\bar{x}_{j_v}) r_{j_v, \Lambda} \quad (29)$$

The results are displayed in Tables IX and X. The first table shows the results for the two kinematics Kin2 and Kin3, that cover the full kinematic range of the experiment, resulting in two domains of different Q^2 , at constant x_{Bj} . The second table shows the results for the two kinematics KinX2 and KinX3, that only cover the domain between the two horizontal lines on Fig. 2, in order to have two domains of different x_{Bj} at constant Q^2 .

VI. RADIATIVE CORRECTIONS

The external radiative effects on the incident electron, and internal real radiative effects at the vertex are treated in the equivalent radiator approximation [24, 25]. Pre-radiation is modeled by generating an event-by-event energy loss ΔE_{in} of the incident electron (E_0) following a distribution ($b \simeq 4/3$):

$$I_{in}(E_0, \Delta E_{in}, t_{in}) = \frac{bt_{in} + \delta_S/2}{\Delta E_{in}} \left[\frac{\Delta E_{in}}{E_0} \right]^{bt_{in} + \delta_S/2} \quad (30)$$

with

$$\delta_S = \frac{2\alpha}{\pi} \left[\ln \frac{Q^2}{m_e^2} - 1 \right] \quad (31)$$

where t_{in} is the event-by-event target thickness (in radiation lengths) traversed by the electron before the scattering vertex. The Schwinger term δ_S models the internal pre-radiation. The scattered energy at the vertex is $E'_v = E_0 - \Delta E_{in} - Q^2/(2M_p x_{Bj})$. Internal post-radiation is modeled by a similar distribution in the post-radiated energy ΔE_{out} :

$$I_{out} = \frac{\delta_S/2}{\Delta E_{out}} \left[\frac{\Delta E_{out}}{E'_v} \right]^{\delta_S/2} \quad (32)$$

These radiative effects are treated within the peaking approximation. External post-radiation by the scattering electron is modeled with the GEANT3 simulation. Kinematic shifts (*e.g* in either the norm and direction of \vec{q}) from external and internal radiations are fully included in the simulation and thereby unfolded from the extracted cross sections.

In addition to these radiative effects incorporated into our Monte Carlo, we correct the data for internal virtual radiation (vacuum polarization and vertex renormalization effects) as well as the cut-off independent effect of un-resolvable soft real radiation. These contributions are

calculated by the following terms, respectively [26]:

$$\begin{aligned} \delta_{\text{Vacuum}} &= \frac{2\alpha}{3\pi} \left[\ln \left(\frac{Q^2}{m_e^2} \right) - \frac{5}{3} \right] \\ \delta_{\text{Vertex}} &= \frac{\alpha}{\pi} \left[\frac{3}{2} \ln \left(\frac{Q^2}{m_e^2} \right) - 2 - \frac{1}{2} \ln^2 \left(\frac{Q^2}{m_e^2} \right) + \frac{\pi^2}{6} \right] \\ \delta_{\text{Real},0} &= \frac{\alpha}{\pi} \left[-\frac{1}{2} \ln^2 \left(\frac{E}{E'} \right) + \frac{1}{2} \ln^2 \left(\frac{Q^2}{m_e^2} \right) - \frac{\pi^2}{3} \right. \\ &\quad \left. + Sp \left(\cos^2 \frac{\theta_e}{2} \right) \right], \end{aligned} \quad (33)$$

where $Sp(\cos^2 \theta_e/2)$ is the Spence function. After an approximate resummation, the correction we apply to the raw counts (to obtain the equivalent Born approximation cross section) is

$$\text{RadCorr} = e^{-\delta_{\text{Vertex}} - \delta_{\text{Real},0}} (1 - \delta_{\text{Vacuum}}/2)^2 \quad (34)$$

The numerical values for our kinematics are tabulated in Table III.

VII. SYSTEMATIC ERRORS

Two classes of inclusive hadronic electro-production channels compete with the exclusive $H(e, e'\pi^0)p$ reaction: the $H(e, e'\pi^0)N\pi, N\pi\pi, \dots$ channels, with a threshold at $M_X^2 = (M_p + m_\pi)^2 = 1.15 \text{ GeV}^2$ and the $H(e, e'\pi^0)\gamma p$ channel. The first class includes N^* and non-resonant $N\pi$ production in the final state, and diffractive $\rho^+ \rightarrow \pi^+\pi^0$ production via the $ep \rightarrow e\rho^+n$ reaction. All these channels can be observed in a missing-mass squared distribution (Fig. 6). The $H(e, e'\pi^0)\gamma p$ channel originates from the diffractive $ep \rightarrow ep\omega$ reaction, with a 8.5 % branching-ratio decay channel [27]. In our acceptance, the $(e, e'\pi^0)$ missing-mass squared threshold for exclusive ω electroproduction is 1.0 GeV^2 , thus slightly lower than the $N\pi$ threshold of 1.15 GeV^2 . However, based on $ep \rightarrow ep\omega$ measurements performed by [28], the expected background of $\omega\pi^0\gamma$ events for $M_X^2 < 1.15 \text{ GeV}^2$ is less than 1% of the exclusive $H(e, e'\pi^0)p$ yield in all $t_{min} - t$ bins.

The systematic errors in the extraction method are due to the cut on the missing-mass squared M_X^2 and on the calorimeter threshold E_{Thr} . The stability of the results is checked by varying each cut in turn. The variation in the estimator

$$R = \sum_{\text{Bin}=0}^6 (r_T + \epsilon_L r_L) \quad (35)$$

is used to quantify the systematic errors.

- For the exclusivity (M_X^2) cut, we consider the stability interval from 0.9 to 1.10 GeV^2 in the M_X^2 cut. At the high end we expect the cross section to have contributions from inelastic final states (Fig.




FIG. 8: (Color online) Total cross section integrated over $t_{min} - t$ and ϕ_π , for Kin3, as a function of the M_X^2 cut. The vertical lines indicate, from left to right, the minimal, optimal and maximal M_X^2 cut values of the stability domain.

6). At the low end, we are removing roughly half of the statistics, and we become progressively more sensitive to the experimental lineshape. The stability of the exclusivity cut (*e.g.* for Kin 3) is plotted in Figs. 8, respectively. The cuts and variation are listed in Tables IV and V. In each case, this study is performed with E_{Thr} fixed at 1.0 GeV.

- For the calorimeter threshold E_{Thr} , the stability of R is expected when the software threshold is fixed above the hardware threshold. Above the hardware threshold, the cut is directly correlated with the $\pi^0 \rightarrow \gamma\gamma$ decay phase space, and the number of events decreases linearly with E_{Thr} . This comes from the isotropic decay of the pion, leading to a flat energy distribution of each decay photon. Fig. 9 shows for Kin2, the quantity R along with the raw number of counts. The stability is indeed no longer observed when the statistics are not linear with the threshold, meaning the hardware threshold competes with the analysis threshold. The same behavior is shown for Kin3. For both kinematics, the systematic error coming from the calorimeter threshold is evaluated as $\pm 1\%$.

The optimal cut is set in the middle of the stability interval (see Figs. 8 and 9). The stability interval bounds and the optimal values for the M_X^2 cut and E_{Thr} are listed in Table IV for both kinematics.

The reduced structure functions r_Λ are extracted at the optimal value of the cuts. For the structure functions implied in ϕ_π dependences, systematic errors are taken as the RMS difference between the r_Λ computed at the optimum cuts and the r_Λ computed at each of the four extremities of the stability domain.

All instrumental sources of systematic errors are shown along with the analysis systematic errors in Table V. Since all sources of systematic errors are independent, we added them quadratically. This total systematic error is included in Tables IX and X.




FIG. 9: (Color online) (a): Total cross section integrated over $t_{min} - t$ and ϕ_π , for Kin2, as a function of E_{Thr} . The vertical lines indicate, from left to right, the minimal, optimal and maximal E_{Thr} values of the stability domain (see Table IV for Kin3 values). (b): The number of events as a function of E_{Thr} . The stability domain for E_{Thr} shows the statistics linearly decreasing with E_{Thr} .

Variable	Minimum	Optimum	Maximum
Kin3 / KinX3			
M_X^2 cut (GeV ²)	0.90	1.00	1.10
E_{Thr} (GeV)	1.20	1.275	1.35
Kin2 / KinX2			
M_X^2 cut (GeV ²)	0.90	1.00	1.10
E_{Thr} (GeV)	1.00	1.075	1.15

TABLE IV: Values of the M_X^2 cut and E_{Thr} defining the global cross-section stability domain. Minimum and Maximum are the bounds of this domain, and Optimum is the cut value set in the middle of the stability interval.

VIII. RESULTS

The exclusive π^0 electroproduction cross section, and in particular the ϕ_π dependences of its separated components were extracted for Kin2, Kin3, KinX2 and KinX3. Our statistics allowed us to achieve, for the ϕ_π -independent cross section, a statistical precision of 3% for Kin2 and Kin3, and of 5% for KinX2 and KinX3. This difference is due to the fact that we could use the full statistics for Kin2 and Kin3, whereas less than half of statistics were available for KinX2 and KinX3.

Fig. 10 shows $\sigma_T + \epsilon_L \sigma_L$ and Fig. 11 σ_{TL} , σ_{TT} , and $\sigma_{TL'}$ plotted as a function of $t_{min} - t$, both for Kin2 and Kin3. Fig. 12 shows $\sigma_T + \epsilon_L \sigma_L$ and Fig. 13 σ_{TL} , σ_{TT} and $\sigma_{TL'}$, both for KinX2 and KinX3.

We performed fits proportional to $\sin \theta_\pi^{CM}$ for σ_{TL} and $\sigma_{TL'}$, and proportional to $\sin^2 \theta_\pi^{CM}$ for σ_{TT} . These fits, including statistical and systematic errors, are shown as bands in Figs. 11 and 13, and in Tables XI and XII. Their reduced χ^2 are below 1.05 for the Q^2 -dependent data, and below 0.75 for the x_{Bj} -dependent data. This

	Kin3	Kin2
	KinX3	KinX2
Exclusivity cut	1.5 %	3.0 %
Calorimeter threshold	1.0 %	
HRS acceptance	2.2 %	
Radiative corrections	1.5 %	
Target length	0.5 %	
Hadronic tensor integration	0.3 %	
Multi tracks corrections	0.1 %	
3 clusters corrections	0.1 %	
Luminosity	0.1 %	
Dead time	0.1 %	
Particle identification	0.1 %	
Total Quadratic	3.3 %	4.2 %
Beam polarization	2.0 %	
Total Quadratic	3.9 %	4.6 %

TABLE V: Experimental systematic errors. The first “Total Quadratic” row shows the quadratic sum of all experimental helicity-independent systematic errors. The second “Total Quadratic” row shows the quadratic sum of all experimental systematic errors including helicity-dependent effects.




FIG. 10: (a) separated $H(e, e' \pi^0)p$ cross section $\sigma_T + \epsilon_L \sigma_L$ as a function of $t_{\min} - t$ for $x_{Bj} = 0.36$. Error bars represent statistical errors only. (b): ratio of $\sigma_T + \epsilon_L \sigma_L$ for the two kinematics as a function of $t_{\min} - t$. The fit of this ratio (dashed line) indicates the Q^2 -dependence of the cross section.

confirms that the main t -dependence of $\sigma_{TL, TL'}$, and σ_{TT} is given by $\sin \theta_{\pi}^{CM}$ and $\sin^2 \theta_{\pi}^{CM}$ respectively.

The lower panel of Fig. 10 (respectively Fig. 12) also shows the Q^2 -dependence (respectively x_{Bj} -dependence) for the total cross section $\sigma_T + \epsilon_L \sigma_L$. To investigate a Q^2 - or a x_{Bj} -dependence, the ratio of $\sigma_T + \epsilon_L \sigma_L$ for the two kinematics is plotted as a function of $t_{\min} - t$. This ratio is found to be independent of t , thus the value of this ratio is fitted by a constant at the x_{Bj} - and Q^2 - values for the two kinematics.




FIG. 11: σ_{TL} (a), σ_{TT} (b) and $\sigma_{TL'}$ (c) $H(e, e' \pi^0)p$ cross-section components as a function of $t_{\min} - t$ for the two Q^2 -values. Kin2 is represented by the open circles, and Kin3 by the solid circles. Error bars represent statistical errors only. The bands (light for Kin2 and dark for Kin3) show fits proportional to $\sin \theta_{\pi}^{CM}$, $\sin^2 \theta_{\pi}^{CM}$, and $\sin \theta_{\pi}^{CM}$, respectively. Refer to Table IX for more detailed cross-section values, statistical and systematic errors.




FIG. 12: (a): Separated $H(e, e' \pi^0)p$ cross section $\sigma_T + \epsilon_L \sigma_L$ as a function of $t_{\min} - t$ for $Q^2 = 2.1 \text{ GeV}^2$. Error bars represent statistical errors only. (b) Ratio of $\sigma_T + \epsilon_L \sigma_L$ for the two kinematics as a function of $t_{\min} - t$. The fit of this ratio (dashed line) indicates the W -dependence of the cross section.

The dependence of $\sigma_T + \epsilon_L \sigma_L$ in Figs. 10 and 12 yields the following conclusions:

- The ratio $[\sigma_T + \epsilon_L \sigma_L]_{\text{Kin3}} / [\sigma_T + \epsilon_L \sigma_L]_{\text{Kin2}}$ is flat in $t_{\min} - t$ with a reduced χ^2 of 0.33. The ratio is found to be 0.633 ± 0.009 , indicating a Q^2 -dependence of the total cross section of about $1/Q^{4.5}$.
- The ratio $[\sigma_T + \epsilon_L \sigma_L]_{\text{KinX3}} / [\sigma_T + \epsilon_L \sigma_L]_{\text{KinX2}}$ is also flat in $t_{\min} - t$ with a reduced χ^2 of 0.56. This ratio is found to be 0.660 ± 0.015 , indicating




FIG. 13: σ_{TL} (a), σ_{TT} (b) and $\sigma_{TL'}$ (c) $H(e, e'\pi^0)p$ cross-section components as a function of $t_{min} - t$ for the two x_{Bj} -values. KinX2 is represented by the open circles, and KinX3 by the solid circles. Error bars represent statistical errors only. The bands (light for KinX2 and dark for KinX3) show fits proportional to $\sin^2 \theta_{\pi}^{CM}$, $\sin^2 \theta_{\pi}^{CM}$, and $\sin^2 \theta_{\pi}^{CM}$, respectively. Refer to Table X for more detailed cross-section values, statistical and systematic errors.

a W -dependence of the total cross section of about $1/W^{3.5}$.

The Q^2 - and W -dependences of the relevant quantities ($\sigma_T + \epsilon_L \sigma_L$, σ_T , and σ_L , with our conventions (*i.e.* Drechsel-Tiator) and VGG conventions) have been summarized in Table VI.

Quantity	Q^2 -dependence	W -dependence
$\sigma_T + \epsilon_L \sigma_L$	$(Q^2)^{-2.39 \pm 0.08}$	$(W)^{-3.48 \pm 0.11}$
σ_L (Drechsel-Tiator)	$(Q^2)^{-0.50 \pm 0.13}$	$(W)^{-0.46 \pm 0.57}$
σ_L (VGG)	$(Q^2)^{-1.50 \pm 0.08}$	$(W)^{1.28 \pm 2.52}$

TABLE VI: Q^2 - and W -dependences for the total cross section and the longitudinal cross section with Drechsel-Tiator conventions and with VGG conventions. For σ_L , the dependences have been evaluated neglecting σ_T . The Q^2 - and W -dependences of σ_T alone (*i.e.* assuming $\sigma_L = 0$) are the same as the Q^2 - and W -dependences of $\sigma_T + \epsilon_L \sigma_L$.

IX. DISCUSSION

In the domain in $t_{min} - t$ where we extracted cross sections, the r_A values from Equations (15) and (16) are constant within statistics, as evidenced by the fits in Figs. 11 and 13.

The data we extracted (see previous section) yield two conclusions with regard to the available models:

- The t -channel meson exchange model of J.M. Laget (Fig. 14), is able to describe $\sigma_T + \epsilon_L \sigma_L$ and $\sigma_{TL'}$, but neither σ_{TL} nor σ_{TT} [6];

- the Q^2 -dependence of the cross section (Fig. 10 and Table VI) demonstrates that we are far from the QCD leading twist prediction of $d\sigma_L/dt$, which behaves as $1/Q^6$. On the other hand, it is similar to the Q^2 -dependence of the transverse cross section for charged pion electroproduction published by Hall C [8].




FIG. 14: (color online) New calculations, at Kin2 (left panels) and Kin3 (right panels) of the t -channel meson exchange model, including charge pion rescattering with πN and $\pi \Delta$ intermediate states [6]. Dashed lines: pole contributions and Pomeron cut alone. Dash-dotted lines: without $\rho \Delta$ cuts. Full lines: $\rho \Delta$ cuts included.

Moreover, the π^0 has no charge, no spin, so a direct coupling with a virtual photon is suppressed, which removes the pion-pole contribution to the longitudinal cross section. This suggests that the transverse $ep \rightarrow ep\pi^0$ cross section is likely to dominate, and transverse $ep \rightarrow en\pi^+$ cross sections have already been described by quark fragmentation mechanisms usually used to describe semi inclusive processes.

T. Horn *et al.* measured the exclusive π^+ electroproduction cross section at $Q^2 = 1.60$ and 2.45 GeV^2 , with σ_T and σ_L separation [8]. The t -channel meson exchange model by J.M. Laget reproduces the σ_L component. However, the σ_T component does not follow the TME model prediction. M. Kaskulov *et al.* performed PYTHIA-JETSET calculations using the Lund model applied to π^+ transverse cross sections at Hall C kinematics [15]. In this model, the virtual photon strikes a quark, with a probability given by the structure functions. Due to this, the hadronic system fragments into two jets. The jet engendered by the single quark gives a pion, and the one engendered by the remainder of the nu-

cleon gives the final neutron. These calculations applied to Hall C π^+ transverse cross sections are in excellent agreement with the data. This gives evidence that the π^+ transverse cross sections at $Q^2 > 1 \text{ GeV}^2$ above the resonance region is described by a partonic process. This suggests that the present π^0 data could similarly be described by incoherent scattering on the partonic structure of the nucleon target.

For these reasons, we consider our data within the context of Semi-Inclusive Deep Inelastic Scattering (SIDIS). We can try to fit our data with a SIDIS formalism written by Anselmino *et al.* [29]. The equation (38) of [29] gives the cross section for semi-inclusive production of a pion (valid for any hadron):

$$\begin{aligned} \frac{d^5 \sigma^{\ell p \rightarrow \ell \pi X}}{dx_{Bj} dQ^2 dz_\pi d^2 p_{\pi\perp}} &\simeq \sum_q \frac{2\pi\alpha^2 e_q^2}{Q^4} f_q(x_{Bj}) D_q^h(z_\pi) \\ &\times \left[1 + (1-y)^2 - 4 \frac{(2-y)\sqrt{1-y}\langle k_\perp^2 \rangle z_\pi p_{\pi\perp}}{\langle p_{\pi\perp}^2 \rangle \sqrt{Q^2}} \cos \phi_\pi \right] \\ &\times \frac{1}{\pi \langle p_{\pi\perp}^2 \rangle} e^{-p_{\pi\perp}^2 / \langle p_{\pi\perp}^2 \rangle} \end{aligned} \quad (36)$$

where $y = p \cdot q / p \cdot k$, and $z_\pi = p \cdot p_\pi / p \cdot q$ is the fraction of the reaction energy carried by the measured hadron, and the quantities between angle brackets are the standard deviations of transverse momentum distributions, which are approximated as gaussian. $\langle k_\perp^2 \rangle$ stands for the parton transverse momentum in the proton, and $\langle p_{\pi\perp}^2 \rangle = \langle p_\perp^2 \rangle + z_\pi^2 \langle k_\perp^2 \rangle$ for the measured transverse momentum of the observed hadron, where $\langle p_\perp^2 \rangle$ stands for the transverse momentum of the hadron with respect to the direction of the struck quark. The idea is to adjust the ratio of the $\cos \phi_\pi$ over constant term in brackets of equation (36) by adjusting only the parameter $\langle p_\perp^2 \rangle / \langle k_\perp^2 \rangle$:

$$\frac{\sqrt{2\epsilon_L(1+\epsilon)}\sigma_{TL}}{\sigma_T + \epsilon_L\sigma_L} = \frac{4(2-y)\sqrt{1-y}z_\pi p_{\pi\perp}}{\left(\frac{\langle p_\perp^2 \rangle}{\langle k_\perp^2 \rangle} + z_\pi^2\right)\sqrt{Q^2}(1+(1-y)^2)} \quad (37)$$

Two conclusions arise from the fits shown in Fig. 15:

- the minus sign affecting the $\cos \phi_\pi$ term in the SIDIS model is in agreement with the σ_{TL} ;
- $\langle p_\perp^2 \rangle$ must be equal to $\sim 5.0 \times \langle k_\perp^2 \rangle$ to reproduce the data.

The authors of [29] adjusted their model to semi-inclusive data. They give $\langle k_\perp^2 \rangle = 0.25 \text{ GeV}^2$ and $\langle p_\perp^2 \rangle = 0.20 \text{ GeV}^2$, giving a ratio $\langle p_\perp^2 \rangle / \langle k_\perp^2 \rangle \sim 0.8$. However, they extracted these values in the inclusive region, implying a high multiplicity of particles, whereas in our data, the multiplicity of particles is unity. Typically, Anselmino *et al.* fit their model with data covering the range $0.1 < z_h < 1.0$, with most of the statistics within $z_h < 0.4$, whereas our data are within $z_h > 0.9$. Furthermore, Kaskulov *et al.* [15] used a value of 1.4 GeV^2 for the RMS transverse momentum of partons in their




FIG. 15: Ratio $\frac{\sqrt{2\epsilon_L(1+\epsilon)}\sigma_{TL}}{\sigma_T + \epsilon_L\sigma_L}$ for Kin2 (open circles) and Kin3 (solid circles) plotted as a function of $p_{\pi\perp}$. Error bars represent statistical errors only. We fitted to each kinematics a model by Anselmino *et al.* in [29] using $\langle p_\perp^2 \rangle / \langle k_\perp^2 \rangle$ as a free parameter, where $\langle k_\perp^2 \rangle$ is the intrinsic transverse momentum of quarks, and $\langle p_\perp^2 \rangle$ is the transverse momentum transferred during the hadronization process. The reduced χ^2 of the fits are 2.12 for Kin3 and 2.65 for Kin2.

fit of the Hall C π^+ data. The exclusive limit of SIDIS could be defined by a SIDIS-inspired model applicable to data at $z_\pi \rightarrow 1.0$ or, more practically, when the measured hadron carries such a large fraction z_π of the total energy of the reaction that it does not allow the production of another particle.

The HERMES and COMPASS collaborations have published $\langle \cos 2\phi_\pi \rangle$ moments of π^+ and π^- SIDIS, including z_h up to 0.7 [30]. However, it is not possible to make a direct comparison to our σ_{TT} π^0 data as the π^+ and π^- moments on the proton have different sign and magnitude for Boer-Mulders effect. On the other hand, the higher twist Cahn effect, which also contributes to σ_{TT} , does not give by itself a satisfying description of σ_{TT} .

X. CONCLUSIONS

We extracted the separated differential π^0 cross section at Jefferson Lab, Hall A, at four kinematic settings: Kin2 and Kin3 with a 3 % statistical precision, and KinX2 and KinX3 with a 5 % statistical precision. We studied the Q^2 -dependence of the hadronic tensor with the two first settings, and the x_{Bj} -dependence with the latter two.

The shape and order of magnitude of $\sigma_T + \epsilon_L\sigma_L$ indicate that the t -channel meson-exchange model is not able to reproduce the total π^0 cross section.

Table VI summarizes the contradiction between our data and the leading twist QCD prediction for high Q^2 . Instead of a $\sim Q^{-6}$ dependence we find, under the assumption that σ_T is negligible (which is very unlikely), a Q^{-3} dependence for σ_L .

On the other hand, the cross section extracted may

show an analogy with the formalism of SIDIS at the exclusive limit. Our $ep \rightarrow ep\pi^0$ data, and Hall C $ep \rightarrow ep\pi^+$ data are important bases for studying the applicability of the SIDIS concepts to exclusive data.

To improve the understanding of our data, we plan to run another π^0 experiment in 2010, at two beam energies, allowing us to disentangle $\epsilon_L \frac{d\sigma_L}{dt}$ from $\frac{d\sigma_T}{dt}$.

We acknowledge the essential work of the JLab accelerator division and the Hall A technical staff. This work was supported by DOE contract DOE-AC05-06OR23177 under which the Jefferson Science Associates, LLC, operates the Thomas Jefferson National Accelerator Facility. We acknowledge additional grants from DOE, NSF and the French CNRS, ANR, and Commissariat à l'Énergie Atomique.

-
- [1] G. R. Goldstein and J. F. Owens, Phys. Rev. **D7**, 865 (1973).
 - [2] M. Guidal, J. M. Laget, and M. Vanderhaeghen, Nucl. Phys. **A627**, 645 (1997).
 - [3] M. Guidal, J. M. Laget, and M. Vanderhaeghen, Phys. Lett. **B400**, 6 (1997).
 - [4] R. De Masi et al. (CLAS), Phys. Rev. **C77**, 042201 (2008).
 - [5] J. M. Laget, Phys. Rev. **C76**, 052201 (2007).
 - [6] J. M. Laget (2010), 1004.1949.
 - [7] S. Ahmad, G. R. Goldstein, and S. Liuti, Phys. Rev. **D79**, 054014 (2009).
 - [8] T. Horn et al., Phys. Rev. **C78**, 058201 (2008).
 - [9] J. C. Collins, L. Frankfurt, and M. Strikman, Phys. Rev. **D56**, 2982 (1997).
 - [10] K. Goeke, M. V. Polyakov, and M. Vanderhaeghen, Prog. Part. Nucl. Phys. **47**, 401 (2001).
 - [11] M. Vanderhaeghen, P. A. M. Guichon, and M. Guidal, Phys. Rev. **D60**, 094017 (1999).
 - [12] D. Mueller, D. Robaschik, B. Geyer, F. M. Dittes, and J. Horejsi, Fortschr. Phys. **42**, 101 (1994).
 - [13] X.-D. Ji, Phys. Rev. Lett. **78**, 610 (1997).
 - [14] A. V. Radyushkin, Phys. Rev. **D56**, 5524 (1997).
 - [15] M. M. Kaskulov, K. Gallmeister, and U. Mosel, Phys. Rev. **D78**, 114022 (2008).
 - [16] D. Drechsel and L. Tiator, J. Phys. **G18**, 449 (1992).
 - [17] P. Y. Bertin, C. E. Hyde, R. Ransome, and F. Sabatié (2000), jefferson Lab Proposal, E00-110.
 - [18] C. Muñoz Camacho, Ph.D. thesis, Université Paris VI, France (2005).
 - [19] C. Munoz Camacho et al. (Jefferson Lab Hall A), Phys. Rev. Lett. **97**, 262002 (2006).
 - [20] M. Mazouz and V. E., Hall A Technical Note Jlab-TN-05-033, Jefferson Lab (2005), URL <http://hallaweb.jlab.org/publications/Technotes/technote.html>.
 - [21] J. J. Kelly, Phys. Rev. **C70**, 068202 (2004).
 - [22] G. F. Chew, M. L. Goldberger, F. E. Low, and Y. Nambu, Phys. Rev. **106**, 1345 (1957).
 - [23] M. Rvachev, Hall A Technical Note Jlab-TN-01-055, Jefferson Lab (2001), URL <http://hallaweb.jlab.org/publications/Technotes/technote.html>.
 - [24] L. W. Mo and Y.-S. Tsai, Rev. Mod. Phys. **41**, 205 (1969).
 - [25] Y.-S. Tsai, Rev. Mod. Phys. **46**, 815 (1974).
 - [26] M. Vanderhaeghen et al., Phys. Rev. **C62**, 025501 (2000).
 - [27] W. M. Yao et al. (Particle Data Group), J. Phys. **G33**, 1 (2006).
 - [28] L. Morand et al. (CLAS), Eur. Phys. J. **A24**, 445 (2005).
 - [29] M. Anselmino et al., Phys. Rev. **D71**, 074006 (2005).
 - [30] V. Barone, S. Melis, and A. Prokudin (2009), 0912.5194.

	Q^2 dependence		x_{Bj} dependence	
	Kin3	Kin2	KinX3	KinX2
N_{π^0}	15516	23429	5952	9860
N_{Gen}	2.14×10^9	2.14×10^9		
$\int \mathcal{L} dt$	$5.10 \times 10^9 \text{ nb}^{-1}$	$2.99 \times 10^9 \text{ nb}^{-1}$		
$Q^2 \text{ (GeV}^2\text{)}$	2.350 ± 0.002	1.941 ± 0.010	2.155 ± 0.268	2.073 ± 0.001
x_{Bj}	0.368 ± 0.001	0.368 ± 0.005	0.335 ± 0.045	0.394 ± 0.003
$W \text{ (GeV)}$	2.217 ± 0.004	2.055 ± 0.012	2.272 ± 0.072	2.016 ± 0.008
$t_{min} \text{ (GeV}^2\text{)}$	-0.173 ± 0.001	-0.170 ± 0.005	-0.137 ± 0.048	-0.199 ± 0.003
ϵ	0.649 ± 0.002	0.769 ± 0.003	0.648 ± 0.001	0.768 ± 0.003
$E_0 \text{ (GeV)}$	5.752 ± 0.001	5.753 ± 0.001	5.752 ± 0.001	5.753 ± 0.001
$E' \text{ (GeV)}$	2.348 ± 0.007	2.937 ± 0.020	2.321 ± 0.029	2.951 ± 0.016
$q^{lab} \text{ (GeV)}$	3.734 ± 0.007	3.143 ± 0.017	3.732 ± 0.009	3.151 ± 0.014
$p_{\pi}^{CM} \text{ (GeV)}$	0.904 ± 0.002	0.806 ± 0.007	0.937 ± 0.043	0.783 ± 0.005
$k_{\gamma}^{CM} \text{ (GeV)}$	0.910 ± 0.002	0.813 ± 0.007	0.942 ± 0.042	0.790 ± 0.005

TABLE VII: Average quantities weighted with the cross section for the four kinematics of the experiment. Errors are the maximal deviation of the values in the seven $t_{min} - t$ bins, compared to the averages listed.

Q^2 dependence			x_{Bj} dependence		
$t_{min} - t \text{ (GeV}^2\text{)}$	$\sin \theta_{\pi}^{CM}$	$\sin^2 \theta_{\pi}^{CM}$	$t_{min} - t \text{ (GeV}^2\text{)}$	$\sin \theta_{\pi}^{CM}$	$\sin^2 \theta_{\pi}^{CM}$
Kin3			KinX3		
0.0095	0.077	0.007	0.0095	0.076	0.007
0.0298	0.144	0.021	0.0297	0.143	0.020
0.0546	0.194	0.038	0.0545	0.193	0.037
0.0844	0.241	0.058	0.0843	0.240	0.058
0.1188	0.285	0.081	0.1188	0.284	0.081
0.1583	0.328	0.108	0.1579	0.326	0.106
0.2063	0.372	0.139	0.2057	0.370	0.137
Kin2			KinX2		
0.0094	0.085	0.008	0.0094	0.085	0.008
0.0296	0.159	0.026	0.0296	0.160	0.026
0.0541	0.215	0.046	0.0542	0.216	0.047
0.0839	0.267	0.071	0.0840	0.268	0.072
0.1179	0.315	0.099	0.1181	0.316	0.100
0.1576	0.362	0.131	0.1579	0.364	0.133
0.2050	0.410	0.168	0.2051	0.412	0.170

TABLE VIII: Values for $t_{min} - t$, $\sin \theta_{\pi}^{CM}$ and $\sin^2 \theta_{\pi}^{CM}$, weighted by the cross section.

Q^2 dependence		
	Kin3 $x_{Bj} = 0.369, Q^2 = 2.350\text{GeV}^2$	Kin2 $x_{Bj} = 0.368, Q^2 = 1.941\text{GeV}^2$
$t_{min} - t$ GeV^2	$d\sigma_T/dt + \epsilon_L d\sigma_L/dt$ nb/GeV^2	
0.010	$377 \pm 10 \pm 12$	$571 \pm 10 \pm 24$
0.030	$381 \pm 12 \pm 12$	$600 \pm 12 \pm 25$
0.054	$403 \pm 10 \pm 13$	$641 \pm 12 \pm 27$
0.084	$425 \pm 11 \pm 14$	$673 \pm 15 \pm 28$
0.118	$418 \pm 11 \pm 14$	$645 \pm 16 \pm 27$
0.158	$395 \pm 13 \pm 13$	$636 \pm 25 \pm 27$
0.206	$384 \pm 13 \pm 13$	$628 \pm 36 \pm 26$
	$d\sigma_{TL}/dt$	
0.010	$-13 \pm 23 \pm 10$	$17 \pm 19 \pm 13$
0.030	$38 \pm 26 \pm 24$	$-43 \pm 22 \pm 12$
0.054	$-25 \pm 22 \pm 11$	$-23 \pm 21 \pm 12$
0.084	$-26 \pm 25 \pm 13$	$-19 \pm 27 \pm 14$
0.118	$-75 \pm 24 \pm 9$	$-103 \pm 30 \pm 21$
0.158	$-91 \pm 30 \pm 8$	$-185 \pm 52 \pm 43$
0.206	$-123 \pm 31 \pm 10$	$-189 \pm 74 \pm 34$
	$d\sigma_{TT}/dt$	
0.010	$-12 \pm 23 \pm 14$	$-39 \pm 19 \pm 7$
0.030	$-25 \pm 27 \pm 15$	$-110 \pm 24 \pm 13$
0.054	$-74 \pm 22 \pm 4$	$-141 \pm 22 \pm 17$
0.084	$-64 \pm 25 \pm 14$	$-174 \pm 28 \pm 17$
0.118	$-124 \pm 24 \pm 16$	$-319 \pm 29 \pm 23$
0.158	$-137 \pm 29 \pm 15$	$-352 \pm 45 \pm 53$
0.206	$-134 \pm 30 \pm 15$	$-343 \pm 57 \pm 68$
	$d\sigma_{TL'}/dt$	
0.010	$9 \pm 49 \pm 20$	$31 \pm 51 \pm 15$
0.030	$119 \pm 55 \pm 21$	$136 \pm 61 \pm 24$
0.054	$129 \pm 46 \pm 12$	$61 \pm 56 \pm 41$
0.084	$151 \pm 51 \pm 30$	$123 \pm 68 \pm 20$
0.118	$153 \pm 47 \pm 17$	$120 \pm 69 \pm 24$
0.158	$87 \pm 54 \pm 23$	$142 \pm 91 \pm 36$
0.206	$127 \pm 51 \pm 15$	$76 \pm 99 \pm 80$

TABLE IX: Separated cross section values from Equation (19) (first quoted value) with statistic errors (second quoted value) and systematic errors (third quoted value) for each of the seven considered bins.

x_{Bj} dependence		
	KinX3 $x_{Bj} = 0.335, Q^2 = 2.155\text{GeV}^2$	KinX2 $x_{Bj} = 0.394, Q^2 = 2.073\text{GeV}^2$
$t_{min} - t$ GeV^2	$d\sigma_T/dt + \epsilon_L d\sigma_L/dt$ nb/GeV^2	
0.010	$439 \pm 19 \pm 14$	$635 \pm 17 \pm 26$
0.030	$437 \pm 22 \pm 14$	$703 \pm 21 \pm 29$
0.054	$457 \pm 18 \pm 15$	$683 \pm 19 \pm 28$
0.084	$442 \pm 21 \pm 14$	$688 \pm 23 \pm 29$
0.118	$466 \pm 22 \pm 15$	$682 \pm 23 \pm 28$
0.158	$407 \pm 29 \pm 13$	$662 \pm 34 \pm 28$
0.205	$406 \pm 34 \pm 13$	$591 \pm 44 \pm 25$
	$d\sigma_{TL}/dt$	
0.010	$20 \pm 46 \pm 38$	$-26 \pm 30 \pm 22$
0.030	$2 \pm 50 \pm 17$	$-100 \pm 37 \pm 61$
0.054	$-28 \pm 43 \pm 15$	$-88 \pm 32 \pm 54$
0.084	$-37 \pm 50 \pm 19$	$-68 \pm 38 \pm 487$
0.118	$-74 \pm 55 \pm 27$	$-170 \pm 40 \pm 562$
0.158	$-188 \pm 80 \pm 27$	$-155 \pm 63 \pm 657$
0.205	$-174 \pm 90 \pm 32$	$-228 \pm 82 \pm 738$
	$d\sigma_{TT}/dt$	
0.010	$-16 \pm 44 \pm 16$	$-63 \pm 33 \pm 18$
0.030	$-44 \pm 50 \pm 32$	$-83 \pm 41 \pm 22$
0.054	$-63 \pm 42 \pm 15$	$-153 \pm 36 \pm 24$
0.084	$-114 \pm 47 \pm 8$	$-186 \pm 43 \pm 78$
0.118	$-156 \pm 50 \pm 18$	$-327 \pm 44 \pm 109$
0.158	$-244 \pm 66 \pm 35$	$-247 \pm 65 \pm 141$
0.205	$-124 \pm 69 \pm 42$	$-444 \pm 82 \pm 183$
	$d\sigma_{TL'}/dt$	
0.010	$68 \pm 97 \pm 35$	$-23 \pm 84 \pm 138$
0.030	$12 \pm 109 \pm 39$	$112 \pm 100 \pm 104$
0.054	$236 \pm 88 \pm 19$	$50 \pm 90 \pm 63$
0.084	$126 \pm 99 \pm 26$	$211 \pm 104 \pm 95$
0.118	$119 \pm 93 \pm 22$	$3 \pm 106 \pm 111$
0.158	$246 \pm 106 \pm 89$	$78 \pm 136 \pm 126$
0.205	$177 \pm 104 \pm 30$	$62 \pm 146 \pm 146$

TABLE X: Separated cross section values from Equation (19) (first quoted value) with statistic errors (second quoted value) and systematic errors (third quoted value) for each of the first seven bins in $t_{min} - t$ for $1.95 \text{ GeV}^2 < Q^2 < 2.25 \text{ GeV}^2$.

Q^2 dependence
Kin3 $x_{Bj} = 0.368, Q^2 = 2.350$ (GeV ²)
$\frac{W_{xx}-W_{yy}}{2} = [-562 \pm 62 \pm 32] \times \sin^2 \theta_h^{CM} \cos 2\Phi_h$ nb
$\Re(W_{xz}) = [97 \pm 18 \pm 8] \times \sin \theta_h^{CM} \cos \Phi_h$ nb
$\Im(W_{xz}) = [-206 \pm 35 \pm 13] \times \sin \theta_h^{CM} \sin \Phi_h$ nb
Kin2 $x_{Bj} = 0.368, Q^2 = 1.941$ (GeV ²)
$\frac{W_{xx}-W_{yy}}{2} = [-1024 \pm 58 \pm 51] \times \sin^2 \theta_h^{CM} \cos 2\Phi_h$ nb
$\Re(W_{xz}) = [82 \pm 17 \pm 11] \times \sin \theta_h^{CM} \cos \Phi_h$ nb
$\Im(W_{xz}) = [-142 \pm 38 \pm 19] \times \sin \theta_h^{CM} \sin \Phi_h$ nb

TABLE XI: Φ_h -dependent hadronic tensor parameterization for constant x_{Bj} . The first error is the statistical error, the second is the systematic error.

x_{Bj} dependence
KinX3 $x_{Bj} = 0.335, Q^2 = 2.155$ (GeV ²)
$\frac{W_{xx}-W_{yy}}{2} = [-770 \pm 135 \pm 63] \times \sin^2 \theta_h^{CM} \cos 2\Phi_h$ nb
$\Re(W_{xz}) = [121 \pm 43 \pm 17] \times \sin \theta_h^{CM} \cos \Phi_h$ nb
$\Im(W_{xz}) = [-278 \pm 69 \pm 28] \times \sin \theta_h^{CM} \sin \Phi_h$ nb
KinX2 $x_{Bj} = 0.394, Q^2 = 2.073$ (GeV ²)
$\frac{W_{xx}-W_{yy}}{2} = [-1003 \pm 86 \pm 153] \times \sin^2 \theta_h^{CM} \cos 2\Phi_h$ nb
$\Re(W_{xz}) = [163 \pm 24 \pm 72] \times \sin \theta_h^{CM} \cos \Phi_h$ nb
$\Im(W_{xz}) = [-101 \pm 58 \pm 55] \times \sin \theta_h^{CM} \sin \Phi_h$ nb

TABLE XII: Φ_h -dependent hadronic tensor parameterization for constant Q^2 . The first error is the statistical error, the second is the systematic error.