



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Near-yrast, medium-spin structure of ^{143}Xe

T. Rząca-Urban, W. Urban, J. A. Pinston, A. G. Smith, and I. Ahmad

Phys. Rev. C **83**, 067301 — Published 17 June 2011

DOI: [10.1103/PhysRevC.83.067301](https://doi.org/10.1103/PhysRevC.83.067301)

The near-yrast, medium-spin structure of ^{143}Xe

T. Rzaca-Urban,¹ W. Urban,^{1,2} J. A. Pinston,³ A. G. Smith,⁴ and I. Ahmad⁵

¹*Faculty of Physics, University of Warsaw, ul. Hoża 69, PL-00-681 Warsaw, Poland*

²*Institut Laue-Langevin, 6 rue J. Horowitz, F-38042 Grenoble Cedex 9, France*

³*LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3,*

Institut National Polytechnique de Grenoble, F-38026 Grenoble Cedex, France

⁴*School of Physics and Astronomy, The University of Manchester, M13 9PL Manchester, UK*

⁵*Argonne National Laboratory, Argonne, IL 60439, USA*

(Dated: May 23, 2011)

Excited states in ^{143}Xe , populated in spontaneous fission of ^{248}Cm have been studied by means of γ spectroscopy using the Eurogam2 Ge array. We have identified three rotational bands in ^{143}Xe , a decoupled band originating from the $i_{13/2}$ neutron excitation, a strongly- coupled band based on the $5/2^-$ ground state and a decoupled band based on the 322.9-keV level with spin 9/2. The new excitation scheme of ^{143}Xe is compared to Quasiparticle Rotor Model calculations, performed with a reflection symmetric potential.

PACS numbers: 21.10.Tg, 23.20.Lv, 25.85.Ca, 27.70.+q

The $5/2^-$ spin and parity of the ground state in ^{143}Xe is well established [1] and together with the $3/2^+$ ground state in ^{143}Cs it can explain the 1^- and 4^- spin and parity of the ground state and the 92.2-keV isomer in ^{144}Cs , respectively [2]. However, the neutron configuration corresponding to this state is a puzzle. One would expect for the ground state of ^{143}Xe spin and parity of $1/2^+$ or $3/2^-$, corresponding to the $1/2[660]$ or $3/2[521]$ orbitals, respectively (see Fig.[2] in Ref.[3]). The fact that it is $5/2^-$ suggests that the Nilsson scheme may not be valid in ^{143}Xe because of too low a deformation (we note that the ground state of ^{141}Xe also has spin and parity $5/2^-$). Alternatively, it is possible that the deformation of ^{143}Xe is not too low but that octupole correlations present in this region [4–7] change the order of the Nilsson orbitals, as proposed by Leander *et al* [3].

In ^{143}Xe one might expect a deformed band based on the ground state, by analogy with the g.s. band observed in the ^{145}Ba isotope [5]. Observation of such a band in ^{143}Xe should help to identify the ground state configuration. By analogy with ^{145}Ba one may also expect a low-lying $13/2^+$ level (in ^{143}Xe) originating from the $\nu i_{13/2}$ orbital, with a rotational band on top of it. The properties of rotational bands in ^{143}Xe should allow the deformation to be determined and the validity of the Nilsson scheme for this nucleus to be verified. The study of the evolution of the deformation along the N=89 isotonic chain should also help predict the properties of excited states in N=89 isotones with $Z < 54$, ^{141}Te and ^{139}Sn .

Excited levels in ^{143}Xe were identified and studied by Bentaleb *et al* [8] in a measurement of prompt- γ rays following spontaneous fission of ^{248}Cm using the Eurogam1 detector array. In Ref. [8] a band was found on top of the 741.0-keV level, which resembles the $13/2^+$ band in ^{145}Ba but has a different decay pattern. The limited information reported in [8] is not sufficient to provide a reliable interpretation of ^{143}Xe . Therefore, we reinvestigated this nucleus using data from a measurement of the γ rays following spontaneous fission of ^{248}Cm , performed




FIG. 1: γ spectra double gated on lines in ^{143}Xe .

with the Eurogam2 detector array [9]. Our measurement provided an order of magnitude more data than the run of Ref.[8]. More details about our experiment and data analysis can be found in our earlier paper [10].

We gated on γ lines of ^{143}Xe reported in Ref.[8] and found new lines, which has enabled a rearrangement of the decay scheme proposed in [8]. In Fig.1 (a) we show a spectrum doubly gated on the 78.9-keV ground-state transition and the 244.0-keV line, which is a self-gating doublet [8]. In the spectrum one observes other lines reported in [8], several lines from the complementary Mo isotopes [11, 12] and a new line at 380.1-keV. Figure 1(b) shows a spectrum doubly gated on the 78.9-keV and the 380.1-keV lines in which the known 244.0-keV line and new lines at 528.4- and 600.5-keV are seen. Lines from the complementary ^{102}Mo and ^{103}Mo isotopes are also presented but no other known lines from ^{143}Xe . This indicates that the 380.1-keV line belongs to a new cascade feeding the 323-keV level in ^{143}Xe reported in Ref.[8].

In Fig.2(a) we show a spectrum doubly gated on the




FIG. 4: Angular correlations for $\gamma - \gamma$ cascades in ^{143}Xe , as measured in the present work.

TABLE I: Properties of γ transitions and excited levels in ^{143}Xe , as measured in this work.

E_γ (keV)	I_γ (rel.)	Multipolarity	E_{lev}^{ini} (keV)	I_{lev}^{ini}
78.9 (1)	100 (15)	M1+E2, $\Delta I=1$	78.9	$7/2^-$
174.1 (2)	14 (2)		497.1	$11/2^-$
193.8 (1)	59 (5)		497.1	$11/2^-$
206.0 (5)			703.0	$13/2$
224.4 (1)	73 (5)		303.3	$9/2^-$
244.0 (2)	57 (7)	$\Delta I=1$	322.9	$9/2$
244.2 (2)	83 (6)	$\Delta I=1$	741.3	$13/2$
249.1 (3)	12 (4)		1006.2	$(15/2^-)$
259.9 (2)	25 (5)		757.1	$(13/2^-)$
267.2 (1)	66 (5)	E2, $\Delta I=2$	1008.5	$17/2$
279.0 (5)			1284.7	$(17/2^-)$
303.2 (2)	8 (3)		303.3	$9/2^-$
303.4 (3)	20 (5)		1006.2	$(15/2^-)$
380.1 (1)	41 (6)	E2, $\Delta I=2$	703.0	$13/2$
418.2 (1)	80 (5)	E2, $\Delta I=2$	497.1	$11/2^-$
420.7 (1)	50 (4)	E2, $\Delta I=2$	1429.2	$21/2$
454.0 (2)	15 (5)		757.1	$(13/2^-)$
509.0 (3)	14 (4)		1006.2	$(15/2^-)$
527.6 (3)	10 (5)		1284.6	$(17/2^-)$
528.4 (1)	23 (4)	E2, $\Delta I=2$	1231.4	$17/2$
534.0 (2)	33 (4)	E2, $\Delta I=2$	1963.2	$(25/2)$
600.5 (3)	6 (2)		1831.9	
600.7 (4)	17 (6)		2563.9	

been deduced, where possible, from the conversion coefficient estimates, from angular correlations and from the observed intensity branching ratios. When assigning spins to excited levels, we assumed that spins are growing with excitation energy, which is commonly observed for excited states populated in spontaneous fission [15].

The new data indicate that the 741.3-keV level has spin $13/2$. Its parity could not be deduced directly but the systematics of the $13/2^+$ relative to the $5/2^-$ excitation energy in the $N=89$ isotones, shown in Fig.5, strongly suggests that the parity of the 741.3-keV level is positive. At neutron number $N = 89$ the Fermi level approaches the $1/2^+[660]$ Nilsson orbital originating from the $i_{13/2}$




FIG. 5: Positions of band heads in $N=89$ isotones. The data are taken from Ref.[21].




FIG. 6: Total aligned angular momentum in the band on top of the of 741.3-keV level in ^{143}Xe , calculated assuming $K=1/2$ for this band.

intruder shell. In Fig. 6 we show the total aligned angular momentum for the band on top of the 741.3-keV level in ^{143}Xe , relative to the ground-state band in the ^{142}Xe core. The aligned momentum in this band of $i = 6.5\hbar$ indicates uniquely the $i_{13/2}$ origin of this band and is consistent with the population of the $1/2^+[660]$ orbital.

In Fig.5 we also show positions of the $7/2_1^-$ and $9/2_1^-$ levels in the $N = 89$ isotones. The 78.8-keV level in ^{143}Xe fits well the trend for the $7/2_1^-$ levels, supporting its spin and parity assignment. In heavier $N = 89$ isotones this level is the first excited level in the band based on the $5/2^- [523]$ level and we propose the same interpretation for the 78.8 keV $7/2^-$ in ^{143}Xe . The $9/2^-$ level at 303.3 keV in ^{143}Xe follows the trend for the $9/2^-$ members of the ground-state configuration in $N = 89$ isotones and it is the second excited level in the $5/2^- [523]$ band.

The structure of the $N = 89$ isotones clearly differs from the structure of the $N = 85$ and $N = 87$ isotones. In $^{139}\text{Xe}_{85}$ [16] and $^{141}\text{Xe}_{87}$ [7] one observes decoupled bands originating from the $\nu f_{7/2}$ level and at $N = 87$ there are also decoupled bands originating from the $\nu h_{9/2}$ level. In contrast, at $N = 89$ strongly coupled bands are present. Their presence indicates an increase of nuclear deformation between $N = 87$ and $N = 89$.

The low position of the $5/2^- [523]$ configuration at $N = 89$ is unexpected, as mentioned before. This intriguing observation has also been discussed in Ref.[18], where the authors concluded that octupole correlations are responsible for lowering the $5/2^- [523]$ configuration. Such




FIG. 7: Comparison of the experimental and calculated energies of excited states in ^{143}Xe , as obtained in the present work. Calculations are normalized to experiment at the $5/2^-$ level.

an effect has been predicted [3] and observed in ^{145}Ba [5], the $N = 89$ isotone of ^{143}Xe . Ref.[18] suggested that the reflection-asymmetric orbitals persist at $N = 89$ down to the proton number $Z = 54$. Interestingly, we have found [17] that $B(E1)/B(E2)$ branching ratios in ^{142}Xe increase by an order of magnitude, compared to lighter Xe isotopes. It is then likely that octupole correlations may play a role in ^{143}Xe .

To test the proposed configurations in ^{143}Xe we performed Quasiparticle Rotor Model (QPRM) calculations with a reflection-symmetric potential, using the codes GAMPN, ASYRMO and PROBI [19]. We used a deformation of $\epsilon_2 = 0.15$ for both the positive-parity levels and the negative-parity levels, an inertia parameter $a = 23.3$ keV and a Coriolis attenuation parameter $\xi = 0.70$. Standard values for the κ and μ parameters of the ls and l^2 terms were used [20]. To calculate the γ -decay pattern we took a collective g factor for the core of $g_R = Z/A$ and an effective value of the free neutron g factor of $g_s^{eff} = g_s^{free}$.

The experimental and calculated excitation energies in

^{143}Xe are compared in Fig. 7. The energy of the $13/2^+$ level is reproduced well with the reflection-symmetric potential. Therefore, the mechanism proposed in Ref.[3] may not be needed to explain the relative positions of the $5/2^-$ and $13/2^+$ levels in ^{143}Xe . In our calculations a complex mixing of four orbitals, two with $K=1/2$ ($1/2[541]$ and $1/2[530]$) and two with $K=3/2$ ($3/2[532]$ and $3/2[521]$) produces the $5/2^-$ low-energy solution.

Finally, we comment on the 322.9-keV level in ^{143}Xe , with spin (9/2) and unknown parity. Two low-lying $9/2^-$ levels, originating from the $\nu f_{7/2}$ and $\nu h_{9/2}$ orbitals are expected in $N = 89$ isotones. Strong mixing between these orbitals repels solutions with the same spin. In Fig. 7 the $9/2_1^-$ and $9/2_2^-$ levels are calculated more than 400 keV apart. This suggests that the 322.9-keV level in ^{143}Xe should not have negative parity. Positive parity for the 322.9-keV level would indicate the presence of octupole correlations in ^{143}Xe , because it is not likely that a $9/2^+$ at 322.9 keV would correspond to the $9/2^+$ member of the $i_{13/2}$ decoupled band, expected about 100 keV below the $13/2^+$, as observed in heavier $N = 89$ isotones (see Fig.5). The $9/2^+$ member of the $i_{13/2}$ band in ^{143}Xe is calculated at 662 keV, fitting well the trend (open circle in Fig. 7).

The present data do not provide a clear answer about the role of octupole correlations in ^{143}Xe . On one hand QPRM calculations with a reflection-symmetric potential can reproduce the $5/2^-$ spin and parity for the ground state. On the other hand, a rather poor reproduction of other negative-parity levels suggests that this may be an accidental result and that the mechanism proposed in Ref.[3] is valid. Finding of the parity of the $9/2_2$ level might resolve this problem. A systematic study of the properties of analogous $9/2_2$ levels in the $N=89$ isotones should help achieving this goal.

This work was partly supported by the U.S. Department of Energy, Office of Nuclear Physics, under contract No. DE-AC02-06CH11357. The authors are grateful to the Office of Basic Energy Sciences, US Department of Energy, through the transplutonium element production facilities at the Oak Ridge National Laboratory, for the use of the ^{248}Cm .

-
- [1] J. K. Tuli, Nucl. Data Sheets **94**, 605 (2001).
[2] T. Rzaca-Urban *et al.*, Phys. Rev. C **80**, 064317 (2009).
[3] G. A. Leander *et al.*, Phys. Lett. **B152**, 284 (1985).
[4] W. R. Phillips *et al.*, Phys. Rev. Lett **57**, 3257 (1986).
[5] M. A. Jones *et al.*, Nucl. Phys. **A605**, 133 (1996).
[6] W. Urban *et al.*, Nucl. Phys. **A613**, 107 (1997).
[7] W. Urban *et al.*, Eur. Phys. J. A **8**, 5 (2000).
[8] M. Bentaleb *et al.*, Z. Phys. A **354**, 143 (1996).
[9] P. J. Nolan *et al.*, Ann.Rev.Nuc.Part.Sci.**44**, 561 (1994).
[10] W. Urban *et al.*, Z. Phys. A **358**, 145 (1997).
[11] D. DE Frenne, Nucl. Data Sheets **110**, 1745 (2009).
[12] D. DE Frenne, Nucl. Data Sheets **110**, 2081 (2009).
[13] M. A. Jones *et al.*, Rev. Sci. Instrum. **69**, 4120 (1998).
[14] T. Kibédi *et al.*, Nucl. Instr. Meth. **A 589**, 202 (2008).
[15] I.Ahmad, W.R.Phillips, Rep.Prog.Phys. **58**, 1415 (1995).
[16] W. Urban *et al.*, Phys. Rev. C **66**, 044302 (2002).
[17] W. Urban *et al.*, Eur. Phys. J A **16**, 303 (2003).
[18] W. Borchers *et al.*, Phys. Lett. **B216**, 7 (1989).
[19] P. Semmes and I. Ragnarsson, 1991 (unpublished).
[20] T.Bengtsson, I.Ragnarsson, Nucl. Phys.**A436**, 14 (1985).
[21] ENSDF data base (at www.nndc.bnl.gov).