

This is the accepted manuscript made available via CHORUS. The article has been published as:

# Multiple $\beta^{-}$ decaying states in $^{194}\text{Re}$ : Shape evolution in neutron-rich osmium isotopes

N. Al-Dahan *et al.*

Phys. Rev. C **85**, 034301 — Published 5 March 2012

DOI: [10.1103/PhysRevC.85.034301](https://doi.org/10.1103/PhysRevC.85.034301)

# Multiple $\beta^-$ Decaying states in $^{194}\text{Re}$ : Shape Evolution in Neutron-Rich Osmium Isotopes

N. Al-Dahan,<sup>1,2</sup> P.H. Regan,<sup>2,\*</sup> Zs. Podolyák,<sup>2</sup> P.M. Walker,<sup>2</sup> N.Alkhomashi,<sup>3</sup> G.D. Dracoulis,<sup>4</sup> G. Farrelly,<sup>2</sup> J. Benlliure,<sup>5</sup> S.B. Pietri,<sup>2,6</sup> R.F. Casten,<sup>7</sup> P.D. Stevenson,<sup>2</sup> W. Gelletly,<sup>2</sup> S.J. Steer,<sup>2</sup> A.B. Garnsworthy,<sup>2,8</sup> E. Casarejos,<sup>5,9</sup> J. Gerl,<sup>6</sup> H.J. Wollersheim,<sup>6</sup> J. Grebosz,<sup>10</sup> M. Górska,<sup>6</sup> I. Kojouharov,<sup>6</sup> H. Schaffner,<sup>6</sup> A. Algara,<sup>11,12</sup> G. Benzoni,<sup>13</sup> A. Blazhev,<sup>14</sup> P. Boutachkov,<sup>6</sup> A.M. Bruce,<sup>15</sup> I.J. Cullen,<sup>2</sup> A.M. Denis Bacelar,<sup>15</sup> A.Y. Deo,<sup>2</sup> M.E. Estevez,<sup>5</sup> Y. Fujita,<sup>16</sup> R. Hoischen,<sup>17</sup> R. Kumar,<sup>18</sup> S. Lalkovski,<sup>15</sup> Z. Liu,<sup>19</sup> P.J. Mason,<sup>2</sup> C. Mihai,<sup>20</sup> F. Molina,<sup>10</sup> D. Mücher,<sup>14</sup> B. Rubio,<sup>11</sup> A. Tamii,<sup>21</sup> S. Tashenov,<sup>6</sup> J.J. Valiente-Dobón,<sup>22</sup> and P.J. Woods<sup>19</sup>

<sup>1</sup>Department of Physics, College of Science, University of Kerbala, Kerbala, Iraq

<sup>2</sup>Department of Physics, University of Surrey, Guildford, GU2 7XH, UK

<sup>3</sup>King Abdulaziz City for Science and Technology (KACST), P.O.Box6086, Riyadh11442, Saudi Arabia

<sup>4</sup>Department of Nuclear Physics, Australian National University, Canberra, ACT 0200, Australia

<sup>5</sup>Universidad de Santiago de Compostela, Santiago de Compostela, Spain

<sup>6</sup>GSI, Planckstrasse 1, D-64291 Darmstadt, Germany

<sup>7</sup>Wright Nuclear Structure Laboratory, Yale University, New Haven, Connecticut 06520-8120, USA

<sup>8</sup>TRIUMF, 4004 Wesbrook Mall, Vancouver, BC, V6T 2A3, Canada

<sup>9</sup>University of Vigo, E-36310, Spain

<sup>10</sup>The Henryk Niewodniczański Institute of Nuclear Physics, PL-31-342, Kraków, Poland

<sup>11</sup>IFIC, CSIC-Universidad de Valencia, 46071 Valencia, Spain

<sup>12</sup>Institute of Nuclear Research of the Hungarian Academy of Sciences, Debrecen, H-4001, Hungary

<sup>13</sup>INFN sezione di Milano, I-20133 Milano, Italy

<sup>14</sup>Institut für Kernphysik, Universität zu Köln, 50937 Köln, Germany

<sup>15</sup>School of Environment and Technology, University of Brighton, Brighton, BN2 4GJ, UK

<sup>16</sup>Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

<sup>17</sup>Department of Physics, Lund University, S-22100 Lund, Sweden

<sup>18</sup>Inter-University Accelerator Centre, New Delhi 110067, India

<sup>19</sup>School of Physics and Astronomy, University of Edinburgh, Edinburgh, EH9 3JZ, UK

<sup>20</sup>Horia Hulubei National Institute of Physics and Nuclear Engineering, R-76900 Bucharest, Romania

<sup>21</sup>Research Centre for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047, Japan

<sup>22</sup>INFN-Laboratori Nazionali di Legnaro, 35020 Legnaro, Italy

$\beta$  decays from heavy, neutron-rich nuclei with  $A \sim 190$  have been investigated following their production via the relativistic projectile fragmentation of an  $E/A = 1$  GeV  $^{208}\text{Pb}$  primary beam on a  $\sim 2.5$  g/cm<sup>2</sup>  $^9\text{Be}$  target. The reaction products were separated and identified using the GSI FRagment Separator (FRS) and stopped in the RISING active stopper. Gamma decays were observed and correlated with these secondary ions on an event-by-event basis such that  $\gamma$  ray transitions following from both internal (isomeric) and  $\beta$  decays were recorded. A number of discrete,  $\beta$ -delayed  $\gamma$ -ray transitions associated with  $\beta$  decays from  $^{194}\text{Re}$  to excited states in  $^{194}\text{Os}$  have been observed, including previously reported decays from the yrast  $I^\pi = (6^+)$  state. Three previously unreported  $\gamma$ -ray transitions with energies 194, 349 and 554 keV are also identified; these transitions are associated with decays from higher spin states in  $^{194}\text{Os}$ . The results of these investigations are compared with theoretical predictions from Nilsson Multi Quasi Particle (MQP) calculations. Based on lifetime measurements and the observed feeding pattern to states in  $^{194}\text{Os}$ , it is concluded that there are three  $\beta^-$ -decaying states in  $^{194}\text{Re}$ .

PACS numbers: 25.70.Mn, 21.60.-n and 29.30.Kv

## I. INTRODUCTION

Neutron-rich nuclei with  $A \sim 190$  show a wide variation of nuclear structural properties and are characterised by the presence of different ground-state shapes, for example prolate, oblate, triaxial /  $\gamma$ -soft and spherical [1–3]. The lighter isotopes of elements in this region are prolate deformed in their ground states and with the ad-

dition of more and more neutrons the shape becomes oblate [4]. At the  $N = 126$  closed shell, the nuclei become spherical [5, 6].

The Os-Pt region is known to be structurally very complex [7–9] with evidence of oblate,  $\gamma$ -soft and triaxial deformations. As such, it offers a sensitive testing ground for nuclear models [10, 11]. The Os isotopes exhibit a transition between deformed and spherical nuclei with increasing neutron number [3, 12]. These characteristics suggest that the study of the neutron-rich Os isotopes may help us to understand the interplay between complex nuclear excitation modes [12]. The nucleus  $^{194}\text{Os}$

---

\*Corresponding author: p.regan@surrey.ac.uk

has two neutrons more than the heaviest stable Osmium isotope,  $^{192}\text{Os}$ . The study of the low-lying energy spectrum of  $^{194}\text{Os}$  helps to illuminate how nuclear shapes are changing approaching the phase / shape transition region [3, 12] for Os isotopes.

The projectile fragmentation technique provides opportunities to observe nuclei far from stability [13]. Fragmentation has proved to be an efficient tool for producing exotic nuclear species and when combined with high sensitivity  $\gamma$ -ray detection arrays, structural information can be gained for otherwise inaccessible nuclei [14–17]. The highest sensitivity is achieved with both isomeric and  $\beta$ -delayed  $\gamma$ -ray spectroscopy techniques. Information on the excited states in nuclei populated in this way can be obtained when only a few hundred nuclei of interest are produced [18, 19].

The research presented in this paper involves the production and structural investigation of the most neutron-rich isotopes of the elements Rhenium (Re,  $Z = 75$ ) and Osmium (Os,  $Z = 76$ ) studied to date and the correlation of the decay of  $^{194}\text{Re}$  nuclei with  $\gamma$  rays from transitions in the daughter  $^{194}\text{Os}$  nuclei. These nuclei were produced in projectile fragmentation reactions and identified using the FRS [20] at the GSI laboratory in Darmstadt, Germany. The decays of these nuclei have been studied following both beta and isomer-delayed  $\gamma$ -ray spectroscopy using the RISING [20, 21]  $\gamma$ -ray spectrometer. The  $\beta$ -delayed studies utilised the RISING active stopper which enabled positional and temporal correlations to be determined between the implantation of a specific exotic isotope and its subsequent  $\beta$  decay. Some results from this same experiment on the decay of tantalum isotopes have been reported in reference [22].

## II. EXPERIMENTAL DETAILS

The nuclei of interest were produced following the interaction of a  $^{208}\text{Pb}$  primary beam with an energy of 1 GeV per nucleon from the SIS-18 synchrotron at GSI, with a  $^9\text{Be}$  target of thickness  $2.54\text{ g/cm}^2$  located at the entrance of the FRS. The typical length of the primary beam spill was about 1 second.

To maximise the number of fully stripped ( $q = Z$ ) nuclei passing through the FRS, niobium foils of  $223\text{ mg/cm}^2$  and  $108\text{ mg/cm}^2$  thicknesses were placed after the target and the degrader at the intermediate focal plane (S2). Three FRS settings are discussed in the current work, namely those centred on the transmission of fully stripped  $^{190}\text{Ta}$ ,  $^{192}\text{Ta}$  and  $^{194}\text{Re}$  ions. Note, that unless stated otherwise, the results shown in this work are from the summed data from the three settings outlined in Table 1.

In this experiment the active stopper consisted of three  $5\text{ cm} \times 5\text{ cm}$  Double-sided Silicon Strip Detectors (DSSSDs; see Refs. [23, 24]), each with 16 individual strips on the front and back faces. Figure 1 shows a schematic of the detector configuration at the final focal




FIG. 1: Schematic of the detector configuration at the final focal plane of the FRS used in the present work, for the March 2007 experiment. Three of the possible six positions (black boxes) in the active stopper were occupied by DSSSDs in this particular experiment. MW= MultiWire detectors; Sci= Scintillator detectors; MUSIC= Multi Sampling Ionising Chamber detectors. The secondary ions were transmitted from left to right in this figure. The label “4” indicates that the detectors are placed at the final focal plane of the FRS.

plane of the FRS at GSI which was used in this work.

The active stopper was viewed by the Stopped RISING  $\gamma$ -ray spectrometer, which consists of fifteen, seven element germanium cluster detectors with a measured full-energy peak efficiency of  $\sim 15\%$  at 661 keV [25]. In the experiment described in this work, the active stopper [23] was used for the first time in conjunction with the RISING  $\gamma$ -ray array. The ions of interest were implanted in the active stopper detector set-up. This highly-pixelated Silicon detector stack allows for correlation in time and space of the signal from the implanted ion and subsequent signals produced by  $\beta$  decays.

The technique of correlating the implanted ions with their subsequent  $\beta$  decay is based on the measurement of two main parameters: (i) the identification of the implantation position in the DSSSD detectors and (ii) the correlation time between the implanted ions and subsequent  $\beta$  particle detected in the same or any of 8 neighbouring pixels. In the current experiment, the FRS was operated in a monochromatic mode using an aluminium wedge-shaped degrader [20, 26] at the intermediate focal plane (S2), which had the effect of distributing the implanted ions across a relatively wide area on the DSSSDs. This approach has two advantages; firstly it minimises the probability of having multiple implantations in the same pixel during a typical correlation time; and second it minimises the distribution range of the ions of a given species within the active stopper. Thus selected isotopes were stopped in a single layer in the active stopper silicon detectors. The ions were implanted in the active stopper

TABLE I: Experimental parameter details for the FRS settings, centred on the transmissions of  $^{190}\text{Ta}$ ,  $^{192}\text{Ta}$  and  $^{194}\text{Re}$  respectively.

Setting	$B\rho_1$ (Tm)	$B\rho_2$ (Tm)	S2 degrader Thickness (mg/cm <sup>2</sup> )	S4 degrader Thickness (mg/cm <sup>2</sup> )	Beam Intensity (ion/spill)	Spill Repetition (s)	Total Collection Time (hours)
$^{190}\text{Ta}$	13.08	9.59	5050	3320	$10^8$	20	62
$^{192}\text{Ta}$	13.23	9.75	5050	3450	$10^9$	15	66
$^{194}\text{Re}$	13.05	9.45	5050	3040	$4 \times 10^8$	7	6

and their positions determined by measuring the implantation strip positions as  $(x, y)$  coordinates. The absolute measurement of the implantation time was made with a digital, (absolute) time stamp. Valid implanted events were identified by the production of a high-energy signal in the active stopper detector ( $> 10$  MeV) measured using the logarithmic part of the pre-amplifier response. For the isomer measurements, the implant- $\gamma$ -ray correlation time for the  $\gamma$ -ray electronic Digital Gamma Finder (DGF) modules was fixed to be between  $0 \rightarrow 400$   $\mu\text{s}$ . After this time the  $\gamma$ -ray DGF electronics were closed for that event. If a subsequent decay signal was detected in the DSSSD, the decay trigger logic gave a signal for the  $\gamma$ -ray electronics to be opened again. For such events, the DGF electronics were opened for a further 400  $\mu\text{s}$  period following the  $\beta$  trigger. To measure the  $\beta$ -decay signal, the linear region of the DSSSD pre-amplifiers was used. For each pixel where an implant occurred and for a specific period of time after the implantation, the energy deposited above the threshold (150 keV [23]) was considered to arise from the emission of a  $\beta$  particle from the radioactive ion implanted in the same pixel. Due to the finite  $\beta$ -range in the DSSSDs and to increase the efficiency of measuring ion-correlated  $\beta$ -particle decay events in the off-line software analysis, a matching program was developed to record the decays in the pixels directly neighbouring that of the original ion implant. Therefore, the correlation algorithm considered implantations and decays in the same pixel and decays in the 8 directly neighbouring pixels (i.e. the  $\beta$  particles were not always measured in the same pixel as the one where their radioactive mother ion was implanted). Each subsequent  $\beta^-$  particle in the same pixel could also be correlated in the time regime, until the implantation of a new ion. The average ion-implantation rate across the full DSSSD implantation array (i.e., across all pixels) was between 0.2 and 0.3 ions per second.

In this work, the  $\beta$ -delayed  $\gamma$ -ray spectroscopy of the  $^{194}\text{Re} \rightarrow ^{194}\text{Os}$  decay is discussed. The difference between applying the ion- $\beta$  correlation only between (i) the implantations and decays in the same pixel and (ii) decays in the same pixel plus the 8 neighbouring pixels for the  $\beta$ -delayed  $\gamma$ -ray spectrum for the  $^{194}\text{Re}$  decay is shown in figure 2. Panel (b) of figure 2 has poorer statistics but a better signal-to-noise ratio than the spectrum in panel (a). For this reason, the corre-




FIG. 2: (a)  $\beta$ -delayed  $\gamma$ -ray spectrum of  $^{194}\text{Os}$  using the implantations and decays in the same pixel plus decays in any of the 8 neighbouring pixels; (b)  $\beta$ -delayed  $\gamma$ -ray spectrum of  $^{194}\text{Os}$  using the implantations and decays in the same pixel and (c) shows the  $\beta$ -delayed  $\gamma$ -ray spectrum for excited states of  $^{190}\text{W}$  to confirm the approach of using implantations and decays in the same pixel. This decay has been reported separately in Ref. [22].

lation algorithm applied subsequently to these data was only between the implantations and decays in the same pixel in the DSSSDs. This approach was applied to all  $\beta$ -delayed  $\gamma$ -ray spectra for the nuclei in the particle identification plot. To confirm this approach, good agreement was obtained from the comparison between the  $\beta$ -delayed  $\gamma$ -ray spectrum for excited states in  $^{190}\text{W}$  from the current work with the previously reported results for this nucleus from isomer spectroscopy [27, 28] as shown in figure 2 panel (c). This result was published separately in reference [22].

### A. Particle Identification

The particle identifications in the current work were made on an event-by-event basis. There are two stages for the particle identification of the heavy ions in this




FIG. 3: (Color online) Charge state selection plot of nuclei centred on the transmission of fully stripped ions from summed data, using monochromatic mode.




FIG. 4: (Color online) Particle identification plot for the summed data from the  $^{190}\text{Ta}$ ,  $^{192}\text{Ta}$  and  $^{194}\text{Re}$  centred settings for fully stripped ions ( $\Delta q = 0$ ).

type of study. The first step is a selection on in-flight changes in charge state. The FRS selection is dependent on the charge-changing properties of each ion as it passes through the detector and degrader materials of the FRS. “Fully stripped” ions correspond to no charge state change between the first and second sections of the FRS ( $\Delta q = 0$ ), while “hydrogen-like” ions correspond to the pick-up of one electron at the intermediate focal plane ( $\Delta q = -1$ ), and “helium-like” ions correspond to the pick-up of two electrons ( $\Delta q = -2$ ). Charge state selection was achieved by using the position of the ion in plastic scintillators at the intermediate and final focal plane of the FRS to define the velocities and magnetic

rigidities of the secondary ions. The difference in magnetic rigidity between the first and second stage of the fragment separator can be used to estimate the energy loss of the ion through the degrader. This information together with the energy loss of the ions as measured at the final focal plane using the MUSIC detectors allows a degree of charge state discrimination. This technique is of particular use in the case of heavy neutron-rich nuclei (see Ref. [29, 30] for more details). For the separation of different charge states, it is necessary to derive how  $A/q$  and  $B\rho$  depend on each other.

According to the charge state calculations using the GLOBAL code [31], 95.4% of the  $^{194}\text{Re}$  ions were fully stripped exiting the target and 83.6% after the intermediate focal plane. For the FRS settings centred on the transmission of  $^{190}\text{Ta}$ ,  $^{192}\text{Ta}$  and  $^{194}\text{Re}$ , the identification of the three charge state groups is shown in figure 3

The mass-to-charge ratio of the ions, ( $A/q$ ), was determined from their Time-Of-Flight (TOF) and magnetic rigidity measurements in the second part of the FRS. The TOF in the second stage of the FRS was determined by measuring the time difference of the ions as they passed between two plastic scintillators which were placed (i) before the S2 degrader and (ii) at the final focal plane of the FRS. The change in the magnetic rigidity of ions was measured before and after they passed through the degrader at S2 which was used to obtain information on any change in charge state. The energy deposited by the identified fragments, which gives information on the atomic number ( $Z$ ), was measured as they passed through two Multi Sampling Ionization Chambers (MUSIC). By determining  $A/q$ , the charge state change, the position at the final focal plane and  $Z$ , an unambiguous event-by-event identification can be obtained. Further details of the particle identification technique are given in Refs. [6, 32–34]. The transmitted ions were slowed down in a variable thickness aluminium degrader at the final focal plane and finally implanted into the active stopper. Figure 4 presents the particle identification plot for the summed data assuming fully stripped ions ( $\Delta q = 0$ ). In this figure, the atomic number,  $Z$ , which is calculated from the energy loss of the ions in the MUSIC detector, is plotted versus the TOF in the second half of the FRS, which is related via the magnetic rigidity to the  $\frac{A}{q}$  of the transmitted ions [30].

## B. Confirming the Ion Identification

The secondary ions were separated, selected and identified primarily using their  $A/q$  values. The ions were implanted in the three DSSSDs. The previously reported isomeric decays in  $^{188}\text{Ta}$  [16],  $^{190}\text{W}$  [27, 28],  $^{192}\text{Re}$  [16, 35] and  $^{193}\text{Re}$  [16, 35] were all clearly identified as well as evidence for isomeric decays in  $^{187}\text{Hf}$ ,  $^{189,190}\text{Ta}$  and  $^{191}\text{W}$ , reported previously in Ref. [22]. The particle identification procedure provided an independent validation for the  $\gamma$ -ray energy and timing setups. Figure 5 shows the




FIG. 5:  $\gamma$ -ray spectra from isomeric decays in (a)  $^{187}\text{Hf}$  for a time range  $\Delta t = (0.03 - 12.68)\mu\text{s}$ , (b)  $^{188}\text{Ta}$  for a time range  $\Delta t = (0.05 - 25.63)\mu\text{s}$ , (c)  $^{189}\text{Ta}$  for a time range  $\Delta t = (0.1 - 10.6)\mu\text{s}$ , (d)  $^{190}\text{Ta}$  for a time range  $\Delta t = (0.03 - 3.33)\mu\text{s}$ , (e)  $^{190}\text{W}$  for a time range  $\Delta t = (0.05 - 90.28)\mu\text{s}$ , (f)  $^{191}\text{W}$  for a time range  $\Delta t = (0.03 - 3.33)\mu\text{s}$ , (g)  $^{192}\text{Re}$  for a time range  $\Delta t = (0.05 - 90.28)\mu\text{s}$  and (h)  $^{192}\text{Re}$  for a time range  $\Delta t = (0.15 - 90.35)\mu\text{s}$ , from the present work [16, 22].

final  $\gamma$ -ray energy spectra corresponding to decays from isomeric states which are identified in the summed data from the  $^{190}\text{Ta}$ ,  $^{192}\text{Ta}$  and  $^{194}\text{Re}$  centred settings, for fully stripped ions.

### III. RESULTS

Gamma rays emitted following the  $\beta^-$ -decay of  $^{194}\text{Re}$  ( $Z = 75$ ) to excited states in the daughter nucleus  $^{194}\text{Os}$  ( $Z = 76$ ) were measured. The  $\beta$ -delayed  $\gamma$ -ray spectrum of  $^{194}\text{Os}$  for ion- $\beta$  correlation times (in the same pixel) of  $\Delta t (\text{implant} - \beta) = 0 - 50$  s is shown in figure 2(b). The previously reported decays from the yrast  $2^+$ ,  $4^+$  and  $6^+$  states in  $^{194}\text{Os}$  with energies 218 [12, 36, 37], 383 [12, 37] and 530 [37] keV together with the characteristic Osmium  $K_{\alpha 1}$ ,  $K_{\alpha 2}$  and  $K_{\beta 1}$  X rays with energies 63, 61 and 71 keV respectively are clearly identified in this spectrum. A discrete  $\gamma$ -ray transition with energy 478 keV is also observed, consistent with the previously reported decay [12] from the  $I^\pi = 0_2^+$  state in  $^{194}\text{Os}$ . Three previously unreported transitions in  $^{194}\text{Os}$  with energies 194, 349 and 554 keV are also observed. To verify that these three transitions are associated with decays in the  $^{194}\text{Os}$  daughter nucleus,  $\beta$ -delayed  $\gamma$ -ray spectra for all the nuclei in the particle identification plot of fully stripped ions from the summed data were examined for short ( $\Delta t(\text{implant} - \beta) = 0 \rightarrow 30$  s) and long ( $\Delta t(\text{implant} - \beta) = 0 \rightarrow 120$  s) correlation times. The  $\gamma$ -ray transitions with energies 194, 349 and 554 keV are

only associated with the decay of  $^{194}\text{Re}$  into  $^{194}\text{Os}$ .

The decay half-lives of the mother nuclei were deduced from the time correlation between the implantation time of the identified fragments in the DSSSD detector and the subsequent  $\beta$ -decay in the same pixel. The time correlation was determined using the same DGF time stamping system used in the isomer decay analysis, providing a resolution of 25 ns [23]. The time differences were histogrammed and used to generate a  $\beta$  decay curve for the identified fragments. These data could then also be further gated by the condition that specific discrete  $\gamma$  rays associated with decay in the daughter nuclei were also observed. The decay curves were fitted to single-component exponential decays using a least square fitting minimisation method and assuming a constant background level. Using the  $\beta^-$ -correlation only approach, the half-life measurement for  $^{192}\text{Re}$  of 16(2)s is consistent with the literature value for this decay of 16(1) s [38, 39]. The decay time curves for both  $^{192}\text{Re}$  and  $^{194}\text{Re}$  from the current work for ion- $\beta$ -correlations are shown in figure 6 panels (a) and (b) respectively for the summed data from the  $^{190}\text{Ta}$ ,  $^{192}\text{Ta}$  and  $^{194}\text{Re}$  centred settings.

$^{194}\text{Re}$  is the heaviest isotope of this element studied to date. A single component exponential fit to the decay function for times between ion implantation and subsequent  $\beta^-$  decay in the same pixel for  $^{194}\text{Re}$  ions is shown in figure 6. This simple fitting procedure yields a fitted value for the apparent decay life for  $^{194}\text{Re}$  of 6(1) s. However, it is important to note that this represents only pure ion- $\beta$  correlation times, with no selection (via additional gamma-ray gating) of which states were populated in the  $^{194}\text{Os}$  daughter nucleus following the decay of the  $^{194}\text{Re}$  mother.

The  $\gamma$ -ray gated  $\beta$ -decay time spectra associated with all the identified transitions in the  $\beta$ -delayed  $\gamma$ -ray spectrum of  $^{194}\text{Os}$  were measured for different correlation times between  $\Delta t (\text{implant} - \beta) = 0 \rightarrow 1000$  s, see figure 7. These data are summarised in Table II and suggest that there are (at least) *three* separate  $\beta^-$  decaying states from  $^{194}\text{Re}$  into  $^{194}\text{Os}$ . The experimental arguments for this interpretation are outlined as follows. For the 530 keV transition (see figure 7(c)), there is intensity associated with the decay transition after 160 s while in the case of the 478 keV transition (see figure 7 (d)), no such intensity is apparent above the constant, random background level after 40 s. The  $^{194}\text{Re}$  decay half-life associated with feeding to the 478 keV transition was measured to be 5(1) s, consistent within the experimental uncertainties of the half-life value from the decay time for the pure ion- $\beta$  correlation shown in figure 6 (b). There is evidence for a decay with  $T_{1/2}=5(1)$  s (see figures 6(b) and 7 (d)) which feeds the  $0_2^+$  states in  $^{194}\text{Os}$ . Note that by the usual  $\beta$  decay selection rules, this also implies a significant degree of feeding in the  $\beta$  decay of  $^{194}\text{Re}$  *direct to the*  $I^\pi=0^+$  *ground state* of  $^{194}\text{Os}$ , which is not associated with any co-incident,  $\beta$ -delayed  $\gamma$ -ray transition. This is consistent with the observation of a half-life of 6(1) s obtained in figure 6 (b) for the correlations between ions






FIG. 6: (Color online) Decay time curve for  $\beta$ -decay of (a)  $^{192}\text{Re}$  and (b)  $^{194}\text{Re}$  from ion- $\beta$  time correlations for the summed data from the  $^{190}\text{Ta}$ ,  $^{192}\text{Ta}$  and  $^{194}\text{Re}$  centred settings. By using a least squared fit the data were fitted to a single exponential decay function plus a constant background level.

and beta-particles only. A second decay with a half life of approximately 25(8) s was measured from the  $\beta$ -delayed  $\gamma$ -ray projection gated by the 349 keV transition, see figure 7 (f). There is also evidence for a third decaying state in  $^{194}\text{Re}$  with a longer half life of approximately 100 s which feeds the  $6_1^+$  state in  $^{194}\text{Os}$ . The long half-life explains the “non-observation” of the 530 keV transition for short correlation times in the  $\beta$ -delayed  $\gamma$ -ray spectrum of  $^{194}\text{Os}$ , see figure 8. By investigating the difference in ion- $\beta$  decay profiles gated on specific  $\gamma$ -ray transitions in the  $^{194}\text{Os}$  daughter, information could be inferred which leads to the conclusion that there are at least two and probably three individual states in  $^{194}\text{Re}$  which  $\beta^-$  decay, each with a different decay half-life<sup>1</sup>. The three previously unreported  $\gamma$ -ray transitions with energies 194, 349 and 554 keV have been independently related to transitions in associated with decays from high-spin states in  $^{194}\text{Os}$  identified in studies of deep inelastic reactions by Dracoulis *et al.*, [41]. Dracoulis *et al.*, placed these transitions in a level scheme of  $^{194}\text{Os}$  as decays from states with spins / parities  $I^\pi \sim 10^- - 11^-$  [41]. Figure 9 shows the section of the partial experimental level scheme for  $^{194}\text{Os}$  observed by Dracoulis *et al.*, [41] of relevance to the present work.

The  $\beta^-$  decay half-life for  $^{192}\text{Re}$  extracted from the ion-




FIG. 7: (Color online)  $\beta$ -delayed  $\gamma$ -ray gated time spectra for the transitions 218, 383, 530, 478, 194, 349 and 554 keV associated with decay in  $^{194}\text{Os}$  for different correlation times  $\Delta t(\text{implant} - \beta) = 0 \rightarrow 1000$  s from the current work. By using a least squared fit the data were fitted to a single exponential decay function plus a constant background level.

$\beta$  correlations from the present work is 16(2) s. By contrast with the  $^{194}\text{Re}$  decay case, the gamma-gated spectrum associated with this decay shows no clear evidence of feeding in the current work to previously reported, low-lying states in  $^{192}\text{Os}$  [42]. Figure 2a does show a small cluster of counts just below the expected energy of the yrast  $2^+$  state in  $^{192}\text{Os}$  (206 keV), which might represent weak evidence for some (possibly indirect) feeding, the statistics are not sufficient in the current work to demonstrate a direct branch. The finite statistics obtained in the current work favours an interpretation of a single decaying state, probably of low-spin ( $0^+$  or  $0^-$ ) which decays predominantly direct to the ground state of  $^{192}\text{Os}$ .

## IV. DISCUSSION

### A. Multi Quasi Particle (MQP) calculations for $^{194}\text{Re}$

Multi Quasi Particle (MQP) calculations have been performed to predict the spins and parities of the low-lying states in the  $^{194}\text{Re}$  mother nucleus. These have been performed for both the prolate and oblate deformations in this nucleus assuming axial symmetry, see Table II. The MQP calculations predict that in the case of the  $^{194}\text{Re}$  nucleus there is a low-lying prolate configuration with  $I^\pi = 11^-$ , arising from the coupling of the neutron orbital  $13/2^+[606]\uparrow$  to the proton orbital  $9/2^-[514]\uparrow$ . For oblate deformation, the lowest predicted energy states from these calculations have  $I^\pi = 0^+, 1^-$  and  $1^+$  arising

<sup>1</sup> We are aware of recent results of mass measurements from the GSI Storage Ring that are consistent with  $^{194}\text{Re}$  having two long-lived metastable states at excitation energies less than 1 MeV with lifetimes in the seconds range, in addition to the  $\beta^-$  decaying ground state [40].





TABLE III: Results of the Multi Quasi Particle (MQP) calculations for the  $^{194}\text{Re}$  mother nucleus of the spin and the parity of low-lying states in this nucleus for axially symmetric prolate and oblate deformations. The quadrupole ( $\varepsilon_2$ ) and the hexadecapole ( $\varepsilon_4$ ) deformation parameters are taken from Ref. [43]<sup>a</sup>.

Prolate						Oblate					
$I^\pi$	Neutron orbital ( $\nu$ )	Proton orbital ( $\pi$ )	Energy <sup>b</sup> (keV)	$V_R^c$ (keV)	Net <sup>d</sup> (keV)	$I^\pi$	Neutron orbital ( $\nu$ )	Proton orbital ( $\pi$ )	Energy <sup>b</sup> (keV)	$V_R^c$ (keV)	Net <sup>d</sup> (keV)
6 <sup>+</sup> 3 <sup>+</sup>	3/2 <sup>-</sup> [501]↑	9/2 <sup>-</sup> [514]↑	22		(22)	1 <sup>+</sup> 0 <sup>+</sup>	1/2 <sup>+</sup> [660]↑	1/2 <sup>+</sup> [411]↓	64 64	+70 -70	134 -6
7 <sup>+</sup> 6 <sup>+</sup>	13/2 <sup>+</sup> [606]↑	1/2 <sup>+</sup> [411]↓	91 91	+70 -70	161 21	2 <sup>+</sup> 1 <sup>+</sup>	1/2 <sup>+</sup> [660]↑	3/2 <sup>+</sup> [402]↓	2 2	+70 -70	72 -68
9 <sup>+</sup> 4 <sup>+</sup>	13/2 <sup>+</sup> [606]↑	5/2 <sup>+</sup> [402]↑	24 24	-84 +84	-60 108	3 <sup>+</sup> 0 <sup>+</sup>	3/2 <sup>+</sup> [651]↑	3/2 <sup>+</sup> [402]↓	174 174	+70 -70	244 104
10 <sup>+</sup> 3 <sup>+</sup>	13/2 <sup>+</sup> [606]↑	7/2 <sup>+</sup> [404]↓	337 337	+63 -63	400 274	1 <sup>-</sup> 0 <sup>-</sup>	1/2 <sup>+</sup> [660]↑	1/2 <sup>-</sup> [550]↑	0 0	-70 +70	-70 +70
2 <sup>-</sup> 1 <sup>-</sup>	3/2 <sup>-</sup> [501]↑	1/2 <sup>+</sup> [411]↓	67		(67)	2 <sup>-</sup> 1 <sup>-</sup>	3/2 <sup>+</sup> [651]↑	1/2 <sup>-</sup> [550]↑	173 173	-71 +71	102 244
4 <sup>-</sup> 1 <sup>-</sup>	3/2 <sup>-</sup> [501]↑	5/2 <sup>+</sup> [402]↑	0		(0)						
5 <sup>-</sup> 0 <sup>-</sup>	5/2 <sup>-</sup> [503]↓	5/2 <sup>+</sup> [402]↑	264		(264)						
11 <sup>-</sup> 2 <sup>-</sup>	13/2 <sup>+</sup> [606]↑	9/2 <sup>-</sup> [514]↑	46 46	-71 +71	-25 117						
12 <sup>-</sup> 1 <sup>-</sup>	13/2 <sup>+</sup> [606]↑	11/2 <sup>-</sup> [505]↑	304 304	-71 +71	233 375						

<sup>a</sup>  $\varepsilon_2 = 0.125$ ,  $\varepsilon_4 = 0.067$  for the prolate deformation; <sup>a</sup>  $\varepsilon_2 = -0.158$ ,  $\varepsilon_4 = 0.033$  for the oblate deformation.

<sup>b</sup> Energy of the state

<sup>c</sup> Empirical Residual interactions.

<sup>d</sup> Sum of Energy and  $V_R$  (keV)

### C. Systematics of $^{194}\text{Os}$ collective states

The energies of the lowest excited states can be used to infer information about the quadrupole character of the nuclei [48]. Systematics related to the yrast 2<sup>+</sup>, 4<sup>+</sup>, 6<sup>+</sup>, the second 2<sup>+</sup> and 0<sup>+</sup> states (2<sub>2</sub><sup>+</sup> and 0<sub>2</sub><sup>+</sup>) as well as the calculated 6<sup>+</sup> states using the Anharmonic Vibrator Model (AVM), (6<sub>Th</sub><sup>+</sup>) through the osmium isotopic chain are presented in figure 10. The energy of 6<sub>Th</sub><sup>+</sup> state was calculated by using the following relation

$$6_{Th}^+ = (3 \times E(2_1^+)) + (3 \times \varepsilon) \quad (1)$$

where  $\varepsilon$  is calculated assuming the AVM using the following expression [49].

$$\varepsilon = E(4_1^+) - (2 \times E(2_1^+)) \quad (2)$$

where  $E(2_1^+)$  and  $E(4_1^+)$  are the excitation energies of the first 2<sup>+</sup> and 4<sup>+</sup> excited states respectively. It is convenient to look at  $\varepsilon/E(2_1^+)$ . Clearly, for a harmonic vibrator,  $\varepsilon/E(2_1^+) = 0$  while for a pure rigid axial rotor it is 4/3.

By using the  $E(2_1^+)$  data, information about the collective character of the nuclei can be inferred. For the

Osmium isotopes, the yrast 2<sup>+</sup>, 4<sup>+</sup>, 6<sup>+</sup> and 6<sub>Th</sub><sup>+</sup> energies increase with increasing neutron number,  $N$ , while the excitation energy of the  $I^\pi = 2_2^+$  decreases up to  $N = 116$  and increases for heavier  $N$  values, as shown in figure 10. The excitation energy of the 0<sub>2</sub><sup>+</sup> states decreases from an energy of approximately 1 MeV in the N=110 to 116 isotones ( $^{184-192}\text{Os}$ ) and drops dramatically to approximately 700 keV in  $^{194}\text{Os}$ , close in energy to the 2<sub>2</sub><sup>+</sup> and yrast 4<sup>+</sup> states. This gives rise to a near-energy triplet of 0<sub>2</sub><sup>+</sup>, 2<sub>2</sub><sup>+</sup> and yrast 4<sup>+</sup> states in  $^{194}\text{Os}$  which is consistent with a change in shape from the deformed gamma-soft rotor observed for lighter Os isotopes, to a more quadrupole vibrational behaviour at N=118. The shape evolution in this region of the nuclear chart has been studied by Nomura et al., [4]. In particular, IBM calculations have been used to predict the energy sequence and transition rates in the osmium isotopes with increasing neutron number approaching N=120 with parameters based on a mapping of predicted ground state deformations calculated using a Gogny energy density functional. These calculations predict a more gamma-soft rotor behaviour for  $^{194}\text{Os}$  than is observed experimentally, with a higher-lying excited 0<sub>2</sub><sup>+</sup> state which is not reproduced by the experimental data. The ratio of  $\varepsilon/E(2_1^+)$  decreases with  $N$  for both Tungsten and Osmium isotopes, see figure 11. As seen in the figure,  $\varepsilon/E(2_1^+)$  has near rotor values for the light




FIG. 8: (Color online)  $\beta$ -delayed  $\gamma$ -ray spectra showing transitions in (a)  $^{192}\text{Os}$  for correlation time  $\Delta t(\text{implant} - \beta) = 0 \rightarrow 100$  s from the decay of  $^{192}\text{Re}$ , (b)  $^{194}\text{Os}$  for correlation time  $\Delta t(\text{implant} - \beta) = 0 \rightarrow 10$  s, (c)  $^{194}\text{Os}$  for correlation time  $\Delta t(\text{implant} - \beta) = 0 \rightarrow 50$  s and (d)  $^{194}\text{Os}$  for correlation time  $\Delta t(\text{implant} - \beta) = 40 \rightarrow 400$  s, from the decay of  $^{194}\text{Re}$  of fully stripped ions from the combination of  $^{190}\text{Ta}$ ,  $^{192}\text{Ta}$  and  $^{194}\text{Re}$  settings. Previously unreported transitions are labelled in red (color online).




FIG. 9: Partial energy level scheme for  $^{194}\text{Os}$  (from [44]) showing decay gamma rays observed in the present work following the  $\beta^-$  decay of  $^{194}\text{Re}$ .

Os and W isotopes and decreases with increasing neutron number. For the heaviest isotopes known to date, it is passing through a transitional region, apparently en route towards vibrator values.




FIG. 10: (Color online) Systematics of the  $E(2_1^+)$ ,  $E(0_2^+)$ ,  $E(2_2^+)$ ,  $E(4_1^+)$ ,  $E(6_1^+)\text{Exp}$  and  $E(6_1^+)\text{Th}$ . energies for the neutron number  $N=110-122$  for Osmium isotopes.




FIG. 11: (Color online) The ratio of the calculated anharmonicity  $\epsilon$  values normalised to the excitation energy of the yrast state  $2^+$ ,  $(\epsilon/E(2_1^+))$  for Osmium isotopes with  $N=110-122$ . [38].

## V. CONCLUSIONS

A number of discrete,  $\beta$ -delayed  $\gamma$ -ray transitions associated with  $\beta$  decays from  $^{194}\text{Re}$  to excited states in  $^{194}\text{Os}$  have been observed, including previously reported decays from the yrast  $I^\pi = (6^+)$  state in this nucleus. Three previously unreported  $\gamma$ -ray transitions with energies 194, 349 and 554 keV are also identified. The results suggest that there are three separate  $\beta$ -decaying states in the  $^{194}\text{Re}$  which feed the :

- (1)  $0_2^+$  state and the  $0^+$  ground state of  $^{194}\text{Os}$  with  $T_{1/2}=5(1)$  s.

- (2)  $(11^-)$  high spin state in  $^{194}\text{Os}$  ( $T_{1/2}=25(8)$  s).  
 (3) and possibly to the  $(6^+)$  yrast state in  $^{194}\text{Os}$  ( $T_{1/2}=100(10)$  s).

Nilsson Multi Quasi Particle calculations for prolate and oblate (axially symmetric) deformations predict a variety of possible low-lying candidate states in  $^{194}\text{Re}$  which would be expected to  $\beta$  decay to excited states in  $^{194}\text{Os}$ . Specifically, both prolate (high-spin) and oblate (low-spin) configurations are predicted for  $^{194}\text{Re}$  with spin/parities consistent with the expected decays which are observed to populate discrete levels in the  $^{194}\text{Os}$  daughter nucleus. The excitation energy levels for states in  $^{194}\text{Os}$  when compared with systematics for the Osmium isotopic chain are consistent with a  $\gamma$ -soft nucleus,

as predicted by contemporary TRS and HF mean-field calculations for this nucleus.

## VI. ACKNOWLEDGMENTS

The excellent work of the GSI accelerator staff is acknowledged. This work is supported by the EPSRC(UK), STFC(UK), AWE plc, the Iraqi Government by Ministry of Higher Education and Scientific Research, University of Kerbala, College of Science, Department of Physics, The German BMBF under Grant 06KY205I and the US DOE under grant No. DE-FG02-91ER-40609.

- 
- [1] P. D. Stevenson *et al.*, Phys. Rev. C **72**, 047303 (2005).  
 [2] P. Sarriguren, R. Rodríguez-Guzmán, and L. M. Robledo, Phys. Rev. C **77**, 064322 (2008).  
 [3] L. M. Robledo, R. Rodríguez-Guzmán, and P. Sarriguren, J. Phys. G: Nucl. and Part. Phys. **36**, 115104 (2009).  
 [4] K. Nomura, T. Otsuka, R. Rodríguez-Guzmán, L.M. Robledo, P. Sarriguren, P.H. Regan, P.D. Stevenson and Z. Podolyák, Phys. Rev. C **83**, 054303 (2011).  
 [5] K. H. Maier *et al.*, Phys. Rev. C **30**, 1702 (1984).  
 [6] S. J. Steer *et al.*, Phys. Rev. C **78**, 061302(R) (2008).  
 [7] S. W. Yates *et al.*, Nucl. Phys. **A222**, 301 (1974).  
 [8] R. D. Bagnel *et al.*, Phys. Lett. **66B**, 129 (1977).  
 [9] F. T. Baker *et al.*, Nucl. Phys. **A258**, 43 (1976).  
 [10] C. Y. Wu *et al.*, Nucl. Phys. **A607**, 178 (1996).  
 [11] C. Wheldon *et al.*, Phys. Rev. C **59**, 2334(R) (1999).  
 [12] E. R. Flynn and D. G. Burke, Phys. Rev. C **17**, 501 (1978).  
 [13] H. Geissel, G. Münzenberg, and K. Riisager, Annu. Rev. Nucl. Part. Sci. **45**, 163 (1995).  
 [14] J. Benlliure *et al.*, Nucl. Phys. A **87**, 660 (1999).  
 [15] K. Krumbholz *et al.*, Z. Phys. A **11**, 350 (1995).  
 [16] M. Caamaño *et al.*, Eur. Phys. J. A **23**, 201 (2005).  
 [17] Zs. Podolyák *et al.*, Phys. Rev. C **79**, 031305(R) (2009).  
 [18] M. Pfützner *et al.*, Phys. Lett. B **444**, 32 (1998).  
 [19] Zs. Podolyák *et al.*, Euro. Phys. J. Special Topics **150**, 165 (2007).  
 [20] H. Geissel *et al.*, Nucl. Instr. Meth. B **70**, 286 (1992).  
 [21] M. Górska *et al.*, Acta Phys. Pol. B **38**, 1219 (2007).  
 [22] N. Alkhomashi *et al.*, Phys. Rev. C **80**, 064308 (2009).  
 [23] R. Kumar *et al.*, Nucl. Instr. Meth. A **598**, 754 (2009).  
 [24] P. H. Regan *et al.*, Int. J. of Mod. Phys. E **17**, 8 (2008).  
 [25] S. Pietri *et al.*, Nucl. Instr. Meth. B **261**, 1079 (2007).  
 [26] G. Münzenberg, Nucl. Instr. Meth. **B70**, 265 (1992).  
 [27] Zs. Podolyák *et al.*, Phys. Lett. B **491**, 225 (2000).  
 [28] G. F. Farrelly *et al.*, Acta Phys. Polo. B **40**, 885 (2009).  
 [29] S. J. Steer *et al.*, Acta Phys. Pol. B **38**, 1283 (2007).  
 [30] J. Benlliure *et al.*, Nuc. Phys. A **674**, 578 (1999).  
 [31] C. Scheidenberger *et al.*, Nucl. Instr. Meth. B **142**, 441 (1998).  
 [32] P. H. Regan *et al.*, Nucl. Phys. A **787**, 491 (2007).  
 [33] Zs. Podolyák *et al.*, Phys. Lett. B **672**, 116 (2009).  
 [34] S. Pietri *et al.*, Nucl. Instr. Meth. B **261**, 1079 (2007).  
 [35] S. J. Steer *et al.*, Int. J. of Mod. Phys. E **18**, 1002 (2009).  
 [36] R. F. Casten *et al.*, Phys. Lett. B **76**, 280 (1978).  
 [37] C. Wheldon *et al.*, Phys. Rev. C **63**, 011304(R) (2000).  
 [38] <http://www.nndc.bnl.gov/chart/>  
 Online database, National Nuclear Data Center,  
 (Site Visited: March, 2010).  
 [39] C. M. Baglin, Nucl. Data Sheets **84**, 717 (1998).  
 [40] M. Reed, P.M. Walker *et al.*, to be published.  
 [41] G. D. Dracoulis *et al.*, to be published (2011).  
 [42] C. Baglin *et al.*, Nucl. Data Sheet **84**, 717 (1998).  
 [43] P. Möller *et al.*, Atomic data and Nucl. Data Tables **59**, 185 (1995).  
 [44] G. D. Dracoulis, to be published (2011).  
 [45] W. Nazarewicz, M. A. Riley, and J. D. Garrett, Nucl. Phys. A **512**, 61 (1990).  
 [46] E. B. Suckling and P. D. Stevenson, Private Communication (2009).  
 [47] S. Raman, C. W. Nestor, JR., and P. Tikkanen, Atom. Data and Nucl. Data Tables **78**, 1 (2001).  
 [48] R. F. Casten, *Nuclear Structure from a Simple Perspective* (Oxford Science Publications, Oxford New York, 2000).  
 [49] N. V. Zamfir and R. F. Casten, Phys. Rev. Lett. **75**, 1280 (1995).