

This is the accepted manuscript made available via CHORUS. The article has been published as:

Questioning tests of hadron production models based on minimum-bias distributions on transverse energy

Thomas A. Trainor

Phys. Rev. C **91**, 044905 — Published 14 April 2015

DOI: [10.1103/PhysRevC.91.044905](https://doi.org/10.1103/PhysRevC.91.044905)

Questioning tests of hadron production models based on minimum-bias distributions on transverse energy

Thomas A. Trainor

CENPA 354290, University of Washington, Seattle, WA 98195

(Dated: August 22, 2014)

A recent study reports measurements of transverse-energy E_t distributions at mid-rapidity for several high-energy nuclear collision systems. The E_t data are analyzed in the context of constituent-quark (CQ) participants estimated with a Glauber-model simulation. The study concludes that systematic variations of hadron and E_t yields previously interpreted in terms of a two-component soft+hard model (TCM) of hadron production including a dijet (hard) contribution are actually the result of CQ participant trends with only soft production. It is claimed that deviations from linear scaling with the number of nucleon participants of hadron yields vs A - A centrality do not actually arise from dijet production as previously assumed. In the present study I examine the new E_t data in the context of the TCM and compare those results with previous differential spectrum and minimum-bias correlation analysis. I present substantial evidence supporting a significant dijet contribution to all high-energy nuclear collisions consistent with the TCM and conclude that the E_t data, given their systematic uncertainties, fail to support claimed CQ model interpretations.

PACS numbers: 12.38.Qk, 13.87.Fh, 25.75.Ag, 25.75.Bh, 25.75.Ld, 25.75.Nq

I. INTRODUCTION

Some features of the hadronic final state of high-energy nuclear collisions may carry significant information about collision dynamics and production mechanisms. In order of successive integrations (and therefore reduced information) are multiparticle correlations, single-particle spectra, minimum-bias (MB) distributions of event number on integrated particle-number or momentum/energy yields within acceptance windows and centrality distributions of such quantities. This study considers MB distributions within a larger context provided by differential measures. I consider what information can be extracted from MB distributions on transverse energy E_t and multiplicity n_{ch} within some angular acceptance.

Reference [1] studies mid-rapidity MB distributions on E_t in the context of soft hadron production and a constituent-quark (CQ) model (CQM). The study concludes that a two-component model (TCM) including soft (projectile dissociation) and hard (jet-related) hadron production conventionally used to describe MB trends on E_t and n_{ch} serves as a proxy for the real mechanism dominated by soft hadron production from QCD color strings connecting participant CQs, with no significant dijet contribution. The arguments supporting that interpretation are based in part on the historical development of hadronic physics since the 1960s and in part on apparent consistency between MB distributions on n_{ch} and E_t and Monte Carlo simulation of MB distributions on the number of conjectured CQ participants N_{qp} [2].

The CQ narrative deviates strongly from the body of data represented by the TCM as it has developed during the past thirteen years of Relativistic Heavy Ion Collider (RHIC) operation. Some studies of spectra and correlations from $\sqrt{s_{NN}} = 200$ GeV p - p collisions [3–5] reveal a strong dijet contribution quantitatively consistent with perturbative QCD (pQCD) predictions that com-

bine measured jet spectra describing Sp̄pS calorimeter data [6] with measured fragmentation functions (FFs) from the Large Electron-Positron (LEP) collider [7, 8]. The p - p dijet systematics can be extrapolated to A - A centrality dependence via a Glauber A - A geometry model to form a TCM reference. Deviations from the p - p extrapolation reference may reveal new physics. In particular, features of the MB distribution on n_{ch} are described in detail by combining the p - p TCM with modification of FFs according to a simple pQCD prescription [9].

In this study I review claims that a soft mechanism following CQ scaling describes all mid-rapidity hadron and E_t production. I summarize experimental evidence from p - p and Au-Au data that supports a soft+hard two-component model. I consider the structure of E_t data from Ref. [1] and find that the reported MB distribution on E_t for Au-Au collisions is actually consistent with the full TCM, including measured strong centrality variation of a jet-related hard component. I find no necessity to invoke CQ scaling. Certain aspects of p - p and A - A data appear to falsify that conjecture.

The parameter x as used in this study has two conventional meanings. In the context of QCD hadron structure x is a parton momentum fraction. In the context of the TCM for nuclear collisions $x \approx n_h/n_{pp}$ is the fraction of p - p (N - N) multiplicity included in the hard component.

This article is arranged as follows: Sec. II introduces two-component and CQ models for E_t and hadron production. Sec. III reviews relevant analysis methods for high-energy nuclear collisions. Sec. IV introduces PHENIX E_t data and CQ interpretations. Sec. V summarizes the phenomenology of 200 GeV p - p collisions. Sec. VI summarizes the phenomenology of 62.4 and 200 GeV Au-Au collisions. Sec. VII presents a set of challenges for the CQ model based on material in the previous two sections. Secs. VIII and IX present Discussion and Summary. Three appendices review TCM descrip-

tions of MB distributions, TCM energy dependence and the algebraic structure of joint MB distributions.

II. E_t AND HADRON PRODUCTION MODELS

This study compares two-component (significant dijet component) and CQ (no significant dijet component) models for transverse-energy and hadron production near mid-rapidity. Reference [1] favors a CQM in which hadron production is dominated by soft processes (conjectured fragmentation of color strings connecting CQs) with systematics determined by CQ participant number N_{qp} . The role of dijets is assumed to be negligible. Supporting arguments for the CQM are derived from centrality dependence of E_t and n_{ch} production and from the structure of MB distributions on those quantities. The TCM alternative includes dijet production from collisions of small- x gluons as a significant component. Advocates of the CQM assert that the TCM serves only as a proxy for the CQM in describing E_t and hadron production. In this section I introduce the two models.

A. The CQM and soft hadron production

Mid-rapidity measurements of $dn_{ch}/d\eta$ and $dE_t/d\eta$ are reported to give “excellent characterization of the nuclear geometry...and are sensitive to the underlying reaction dynamics...” [1]. For example, $dn_{ch}/d\eta$ is found to deviate from strict proportionality to the number of participant nucleons N_{part} (linearity) expected for soft hadron production. That deviation has been characterized conventionally by the two-component (soft+hard) model, with dijet production as the hard component.

However, for the CQM it is assumed that mid-rapidity hadron and E_t production are dominated by low- p_t soft hadrons and should therefore be insensitive to hard processes. An alternative model based on the number of CQ participants N_{qp} includes only soft-hadron production (no dijets), with produced n_{ch} and E_t proportional to the number of color strings that connect constituent quarks. The relation $dn_{ch}/d\eta \propto N_{qp}$ is assumed [2] based on data from Ref. [10] that cover only the most-central 40% of the Au-Au total cross section.

Reference [1] claims to establish the same proportionality for $dE_t/d\eta$, but the analysis and its interpretation are constrained by critical assumptions: “...possible models motivated by the fact that half the momentum of a nucleon is carried by [small- x] gluons when probed at high Q^2 in hard-scattering are not considered...we limit our comparison to the nucleon and [large- x] constituent-quark participant models...widely used since the 1970’s....” The study concludes that “...the success of the two component model [of hadron and E_t production] is not because there are some contributions proportional to N_{part} and some proportional to $N_{coll} [\equiv N_{bin}]$, but rather because a particular linear com-

bination of N_{part} and N_{coll} turns out to be an empirical proxy for the nuclear geometry of the number of constituent quark participants, N_{qp} in A+A collisions.” Thus, the TCM “...does not represent a hard-scattering component in E_t distributions.” Reference [1] rejects any statistically-significant role for dijets in HE nuclear collisions.

B. The TCM and importance of small- x gluons

A key assumption in the argument of Ref. [1] is exclusion of the role of small- x partons (mainly gluons) in hadron and E_t production at mid-rapidity. However, that assumption contradicts descriptions of hadron small- x structure derived from deep-inelastic scattering (e.g. parton distribution functions or PDFs) that have been greatly refined over the past two decades. It is likely that small- x gluons dominate hadron production in high-energy nuclear collisions, both as sources of “soft” hadron production and, via large-angle parton-parton scattering, as sources of “hard” dijet production. The latter can be predicted quantitatively via pQCD calculations given the flux of small- x gluons [11].

The TCM has had two principal manifestations at RHIC: (a) phenomenological description of A-A production with TCM parameter x fitted to more-central A-A data [12] and (b) extrapolation (based on the Glauber model of A-A collisions) of a detailed TCM for p - p collisions including direct dijet manifestations in spectra and correlations compatible with pQCD calculations. The extrapolation from p - p provides a *Glauber linear superposition* (GLS) reference for the TCM in A-A collisions [3–5]. Reference (b) serves as a null hypothesis: A-A centrality trends inconsistent with the GLS reference (e.g., requiring a variable x) may indicate novel physics [11, 13].

The number of constituent-quark participants N_{qp} is determined by a Glauber-model simulation of A-A collisions assuming three CQs per nucleon, with cross sections adjusted to be self-consistent. By hypothesis CQs reside near $x = 1/3$ (dressed quarks or valons [14]) whereas according to modern QCD the partons most responsible for mid-rapidity hadron production at RHIC energies should reside near $x = 0.01$. Within the same QCD context the small- x structure of hadrons should be universal, not influenced by the configuration of valence quarks.

In this study I present several examples from p - p collision systematics that exclude the CQ hypothesis. I summarize detailed relations among yields, spectra and correlations that are describe accurately by a TCM reference derived from p - p data quantitatively related to pQCD calculations. The p - p reference is then modified in a simple manner, again consistent with pQCD, to describe a broad array of data from more-central Au-Au collisions. I relate the TCM to model and data MB distributions and demonstrate accurate correspondence.

III. ANALYSIS METHODS

I briefly summarize the kinematic variables, spaces and methods required to describe high-energy nuclear collisions, especially related to A - A centrality measurement and yield, spectrum and correlation measurements. Additional method descriptions can be found in Refs. [3–5, 13, 15–19].

A. Kinematic variables and spaces

High-energy nuclear collisions are described efficiently within a cylindrical coordinate system (p_t, η, ϕ) , where p_t is the transverse momentum, ϕ is the azimuth angle from a reference direction and pseudorapidity $\eta = -\ln[\tan(\theta/2)] \approx \sin(\pi/2 - \theta)$ is a measure of the polar angle, the approximation being valid near $\eta = 0$. A bounded detector angular acceptance is denoted by intervals $(\Delta\eta, \Delta\phi)$ in the primary single-particle space (η, ϕ) .

Although scalar momentum p_t is directly measured by particle detectors in this study I prefer to use an alternative measure. To provide better visual access to low-momentum structure and to simplify the description of jet-related spectrum hard components (defined below) I present single-particle (SP) spectra in terms of transverse rapidity $y_t = \ln[(m_t + p_t)/m_h]$ with transverse mass $m_t = \sqrt{p_t^2 + m_h^2}$ and rest mass $m_h = m_\pi$ assumed for unidentified hadrons. The statistical measure $\langle p_t \rangle$ (both event-wise and ensemble means) is retained for analysis of spectra [3, 13], fluctuations [20] and correlations [21, 22].

The main kinematic quantity for this study is transverse energy E_t as measured by an electromagnetic calorimeter (EMCal) and integrated within some angular acceptance. I adopt the PHENIX notation convention $dE_t/d\eta \rightarrow \langle E_t \rangle$. I also consider E_t per hadron $\langle e_t \rangle = E_t / n_{ch}$ within some acceptance.

B. A - A centrality measures

The A - A centrality evolution of n_{ch} , E_t and other quantities is a major issue for this study. A - A centrality is conventionally measured at RHIC by Glauber-simulated participant-nucleon number N_{part} . However, alternative measures offer better access to the more-peripheral centrality variation required to test relevant hypotheses.

Minimum-bias distributions of A - A cross section σ on participant number N_{part} and N - N binary collision number N_{bin} are accurately described by power-law trends, leading to simple parametrizations in terms of the fractional cross section σ/σ_0 [23]

$$(N_{part}/2)^{1/4} = 0.5^{1/4} \frac{\sigma}{\sigma_0} + (N_{part,max}/2)^{1/4} \left(1 - \frac{\sigma}{\sigma_0}\right)$$

$$N_{bin}^{1/6} = 0.5^{1/6} \frac{\sigma}{\sigma_0} + N_{bin,max}^{1/6} \left(1 - \frac{\sigma}{\sigma_0}\right), \quad (1)$$

with $N_{part,max} = 382$ and $N_{bin,max} = 1136$ for 200 GeV Au-Au collisions. Those parametrizations describe Glauber simulations (with $\sigma_{NN} = 42$ mb) at the percent level (e.g., within 1% of the 200 GeV N_{part} values in Table V of Ref. [1]). I use the same values for all energies above $\sqrt{s_{NN}} \approx 30$ GeV as purely geometric centrality measures. The preferred centrality measure for data plots is mean participant path length $\nu = 2N_{bin}/N_{part}$ which provides good visual access to the more-peripheral data required to test the N - N linear-superposition hypothesis.

In Ref. [1] the number of conjectured participant quarks N_{qp} is also determined with a Glauber Monte Carlo simulation. The CQ cross section is defined so as to sum to the N - N cross section at each collision energy, and the number of participant CQs is then determined vs A - A centrality. Simulation results are shown in Fig. 1 (left panel) as the ratio N_{qp}/N_{part} with open points (N_{part} from Ref. [1] Table V) and solid points [N_{part} from Eq. (1)]. The solid curve is a simple parametrization used below for illustration. The open and solid triangles represent estimated ratios for 200 GeV p - p collisions. The solid point is obtained by extrapolating the solid curve. The open point is discussed in Secs. IV C and VII B.




FIG. 1: (a) Ratio N_{qp}/N_{part} derived from a Glauber Monte Carlo simulation of 200 GeV Au-Au collisions from Ref. [1], with N_{part} values from that reference (open circles) and from Eq. (1) (solid points) with a parametrization used in the present study (curve). The two values for 200 GeV p - p collisions (triangles) are discussed in Secs. IV C and VII B. (b) The ratio parametrization from the left panel vs participant pathlength ν (solid curve). Also plotted are the TCM for 200 GeV Au-Au with fixed $x = 0.1$ (upper dashed curve) and $(2/N_{part})dn_{ch}/d\eta$ trends with variable x for three energies (lower dashed curve and dash-dotted and dotted curves). Those curves are scaled down relative to data by factor 1.35.

In Sec. V of Ref. [1] the N_{qp} trend (solid curves) is said to be similar to that for the charged-particle yield n_{ch} per participant nucleon, possibly explaining the measured n_{ch} data trend without recourse to a TCM dijet contribution. In Fig. 1 (right panel) $(2/N_{part})dn_{ch}/d\eta$ distributions for three energies are scaled down by common factor 1.35. The upper dashed line that does approximate the N_{qp} trend is the conventional TCM for 200 GeV assuming fixed parameter value $x \approx 0.1$ for all centralities. The lower dashed curve is the TCM with parameter x varying from 0.015 (GLS extrapolation from p - p col-

lisions) to 0.095 (derived empirically from more-central Au-Au collisions) as required by the measured spectrum and angular-correlation data described below. The lowest two curves show the same variable- x TCM model derived from 130 and 62.4 GeV data. Deviations from the p - p GLS reference value $x \approx 0.015$ have been related quantitatively to modification of parton fragmentation to jets in more-central 200 GeV Au-Au collisions [11].

Figure 1 (right panel) suggests that detailed comparisons between measured hadron-production trends and the N_{qp} trend for more-peripheral A - A collisions may already falsify the CQM, as discussed in Sec. VII C. But that centrality interval is typically de-emphasized at RHIC. More-differential methods may provide a definitive picture, as discussed in Secs. V and VI.

C. The two-component model

The TCM of hadron production in high-energy nuclear collisions includes (near mid-rapidity) a soft component (SC) attributed to projectile-nucleon fragmentation (dissociation) with A - A centrality dependence proportional to N_{part} and a hard component (HC) attributed to large-angle-scattered parton fragmentation to dijets proportional to N_{bin} . The TCM has been applied successfully to single-particle spectra and their integrated yields [3, 13] and to two-particle correlations [4, 5, 15, 16] from p - p and A - A collision systems at several energies.

The soft component of spectra appears to be universal across a broad range of collision systems and energies, reflecting local charge conservation (unlike-sign pair correlations dominate) and an m_t spectrum described by a Lévy distribution with fixed parameters [3, 13]. The hard component of p - p spectra is quantitatively consistent with predictions [11] derived from measured jet (parton) energy spectra [6] and fragmentation functions (FFs) [24]. The hard component of p - p angular correlations on angle differences is consistent with theoretical expectations for dijets: (a) a same-side (SS) 2D peak at the origin corresponds to intrajet correlations and (b) an away-side (AS) 1D peak on azimuth corresponds to interjet (back-to-back jet pairs) correlations.

A TCM reference (null hypothesis) for A - A collisions can be defined by extrapolation from p - p collision data assuming linear superposition of N - N collisions according to the Glauber model, the reference then described appropriately as Glauber linear superposition [15]. Testing the GLS reference in A - A collisions requires accurate measurements especially for more-peripheral collisions. Deviations from the GLS reference in more-central A - A collisions may then signal new physics. Monte Carlos such as HIJING (with no jet quenching [25]) based on models of p - p collisions (PYTHIA [26], HERWIG [27]) should follow the GLS model derived from p - p data for all Au-Au centralities, but they do not [15]. The peripheral region is typically de-emphasized in RHIC analysis, for instance by plotting data vs N_{part} where the more-

peripheral 50% of the total cross section is confined to $N_{part} < 50$ or 13% of the total axis interval. Rather than extrapolate upward from p - p collisions to establish a GLS reference the conventional approach at RHIC has been to derive a fixed value of TCM model parameter x from more-central Au-Au data and then compare only the more-central region with A - A models [10, 12].

The hard components of spectra and angular correlations from more-peripheral Au-Au collisions are indeed consistent with extrapolation of measured p - p structure but undergo substantial changes with increasing Au-Au centrality above a *sharp transition* (ST) on centrality near $\nu \approx 3$ or $\sigma/\sigma_0 \approx 0.5$ [11, 13, 15]. Accurate spectrum and MB measurements reveal that the effective TCM x from 200 GeV Au-Au increases with centrality from a smaller p - p value in more-peripheral collisions to six times that value in more-central collisions, the change corresponding mainly to evolution of parton FFs [11]. The strong variation of x has implications for the structure of MB event-frequency or cross-section distributions on n_{ch} and E_t , as discussed in App. A 1.

D. Single-particle spectra in p-p and A-A collisions

Single-particle p_t or y_t spectra from high-energy p - p collisions can be grouped in multiplicity classes based on the total charge multiplicity within some angular acceptance $\Delta\eta$. Variation of the spectrum shape with multiplicity can be employed to decompose spectra into two components with fixed shapes independent of multiplicity: “soft” (scaling approximately as n_{ch}) and “hard” (scaling approximately as n_{ch}^2) [3]. The soft and hard spectrum components integrated within some angular acceptance yield soft n_s and hard n_h multiplicity components, with $n_{ch} = n_s + n_h$. The quantitative correspondence of the p - p spectrum hard component to a pQCD jet description was established in Ref. [11].

The p - p spectrum TCM serves in turn as the baseline or GLS reference for an A - A spectrum TCM in which some model parameters extrapolated from p - p phenomenology are permitted to vary to accommodate the A - A spectrum data. Whereas n_{ch} is the control or “centrality” parameter for p - p collisions participant path length ν is the relevant centrality parameter for A - A collisions.

E. Correlations in p-p and A-A collisions

Angular correlations on (η, ϕ) and correlations on transverse-rapidity space $y_t \times y_t$ have been studied in detail for p - p and A - A collisions [4, 5, 15, 16, 28, 29]. A two-component model for correlations in p - p collisions provides the basis for a mathematical model of angular correlations [4, 5] which is then applied to A - A collisions with minor modifications [15, 16]. The SC produces correlations only on η , only for unlike-sign (US) charge pairs,

and only below 0.5 GeV/c [4, 5]. It falls to zero amplitude by mid-centrality in 200 GeV Au-Au collisions.

The hard component of angular correlations is fully consistent with expectations for dijet correlations below the sharp transition on centrality. Above the ST the same-side 2D peak (intrajet correlations) becomes elongated on η and narrows significantly on ϕ [15, 16]. The volume of the SS 2D peak is consistent with the SP spectrum hard component in terms of number of integrated fragment pairs, corresponding quantitatively to the pQCD-predicted number of dijets and their fragments [30]. p_t angular correlations analogous to particle-number angular correlations are described in Sec. VID.

Particle-number correlations on transverse-rapidity space $y_t \times y_t$ from p - p collisions also exhibit distinct soft and hard components [4, 5]. The 2D soft component lies almost completely below $p_t = 0.5$ GeV/c. The 2D hard component lies almost completely above that point and, projected onto 1D y_t , is consistent with the SP y_t spectrum hard component. The phenomenology of y_t correlations has been explored in detail [4, 5, 28]. Above the ST on Au-Au centrality the hard component splits, with one part (mainly pions) moving to smaller y_t and the other part (mainly protons) moving to larger y_t , again in agreement with measured SP spectrum trends [13].

IV. PHENIX MINIMUM-BIAS E_t DATA

I first consider the analysis and alternative interpretation of MB distributions on E_t presented in Ref. [1]. The basic data consist of MB distributions integrated within some fiducial angular acceptance and corrected to a reference acceptance as shown in Fig. 3 and corrected E_t production vs centrality as in Table V of that paper. For the present study I emphasize the 200 GeV p - p and Au-Au data and do not consider d-Au data or Additive Quark Model scaling. As noted, to simplify the notation I adopt the PHENIX convention $dE_t/d\eta \leftrightarrow \langle E_t \rangle$.

A. PHENIX EMCal angular acceptance

The absolute size of the detector angular acceptance has important consequences for MB distributions. In Ref. [1] a distinction is established between the fiducial (physical) detector angular acceptance and the defined reference acceptance ($\Delta\eta = 1, \Delta\phi = 2\pi$). The PHENIX EMCal has a substantially smaller angular acceptance than some comparable detectors. For instance, the effective acceptance of the PHENIX EMCal is $1/(8.4 \times 1.2) \approx 1/10$ the STAR TPC or barrel EMCal acceptance. Certain aspects of data obtained within the fiducial acceptance such as mean values can be corrected to a reference acceptance. But the effects of acceptance on other aspects such as statistical fluctuations may persist as biases in the corrected data. Two issues are relevant for

this study: (a) void probabilities and (b) structure of the *terminus* or central- A - A end (tail) of MB distributions.

In the average non-single-diffractive (NSD) p - p collision approximately 10 particles (charged + neutral) may fall within the STAR fiducial EMCal/TPC acceptance (with $\Delta\eta \approx 2$), but only 1 particle would fall within the PHENIX fiducial acceptance. The Poisson void probability for STAR is $\exp(-10) \approx 10^{-5}$, but for PHENIX the void probability is $p_0 \approx 0.35$ (as in Table X of Ref. [1]). Reference [1] notes that “The importance of taking account of p_0 ...cannot be overemphasized.” For the PHENIX acceptance that statement is well justified.

MB data from two detectors corrected to a common reference acceptance may be quite different in the limit of central collisions where the tail of the distribution reflects fluctuations in central collisions determined by the actual or fiducial acceptance. The tail width relative to the terminus mean $\sigma_n/\bar{n}_0 \approx 1/\sqrt{\bar{n}_0}$ (assuming Poisson statistics) for STAR relative to PHENIX should be $1/\sqrt{10} \approx 1/3$. Rescaling the MB distribution to the reference acceptance will not change that ratio. Thus, the terminus widths of MB distributions from PHENIX should be about three times larger than the equivalent from STAR. That issue is discussed further in App. A 2.

B. $\langle E_t \rangle$ centrality trends

Figure 2 (left panel) shows ratio $(2/N_{part})dE_t/d\eta$ from Au-Au collisions for three energies vs centrality measured by mean participant pathlength ν . The $dE_t/d\eta$ data and uncertainties are from Tables V-VII of Ref. [1] while the $N_{part}/2$ and ν values are from Eq. (1). The uncertainties include total point-to-point plus common offset systematic uncertainties. The $\langle E_t \rangle$ centrality trends are described as varying “nonlinearly” with N_{part} , interpreted to imply inconsistency with the participant scaling expected from soft production and therefore apparently requiring hard scattering and the TCM.

Figure 2 (right panel) shows ratio $(2/N_{qp})dE_t/d\eta$ vs centrality obtained by dividing the data in the left panel by ratio N_{qp}/N_{part} from Fig. 1. The ratio data above $\nu = 2$ appear to be constant within systematic uncertainties, suggesting that mid-rapidity hadron production in Au-Au collisions actually occurs by a soft process scaling with the number of CQ participants, not with nucleon participants. Values for 200 GeV p - p collisions derived from data in Table X of Ref. [1] are plotted at $\nu = 1.25$ (\approx NSD N - N collisions). The upper p - p datum is said to confirm the self-consistency of the CQ analysis. The two p - p results are related to p - p N_{qp} values in Fig. 1 and are discussed in Secs. IV C and VII B.

Figure 3 (left panel) shows the TCM for $dn_{ch}/d\eta$ data from three collision energies similarly scaled by N_{qp} . The dotted curve represents the conventional TCM with fixed $x \approx 0.1$, while the other curves represent variable- x models that describe spectrum yields accurately. The dotted curve appears to be constant within typical sys-




FIG. 2: (a) Per-participant transverse-energy production measured by $(2/N_{part})dE_t/d\eta$ vs centrality for 200, 130 and 62.4 GeV Au-Au collisions. The data and uncertainties (stat + sys) are from Tables V-VII of Ref. [1]. The N_{part} and ν values are from Eq. (1). The solid triangle is the corrected p - p value from Table X of Ref. [1]. (b) Data from the left panel divided by ratio N_{qp}/N_{part} from Fig. 1. The p - p points are obtained from the point in the left panel with N_{qp}/N_{part} ratio values 1.95 (solid point, extrapolation of the Fig. 1 solid curve) and 1.4 (open point, separate p - p Glauber constituent-quark simulation reported in Ref. [1]).

tematic uncertainties, seeming to confirm a claim by Ref. [10] that the increase in $(2/N_{part})dn_{ch}/d\eta$ described by the fixed- x TCM arises not from increasing contributions by hard processes but rather from increasing CQ number (and associated soft hadrons) relative to participant nucleons. However the lowest three curves, representing measured production trends, consistent below the ST with the GLS reference extrapolated from p - p measurements, deviate substantially from the CQM for more-peripheral collisions and may falsify that model (see Sec. VII C). The two points are p - p values derived from the two p - p values of N_{qp} from Fig. 1. The preferred value $N_{qp} = 2.8$ from a p - p Glauber simulation leads to the open point which is far above the trend for Au-Au collisions.

As a logical next step I bypass the conjectured geometry model parameters by calculating $\langle e_t \rangle = (dE_t/d\eta)/(dn_{tot}/d\eta)$, where n_{tot} represents all hadrons contributing to $\langle E_t \rangle$, approximated in this case by $n_{tot} \approx 1.5n_{ch}$ to account for π^0 s. Figure 3 (right panel) shows $\langle e_t \rangle$ vs Au-Au centrality (points with uncertainties) for three collision energies. The n_{tot} values are derived from the full TCM with variable x that describes accurately the measured charged-particle spectrum integrals [13]. Also shown is the $\langle e_t \rangle$ value for 200 GeV p - p collisions (solid triangle) derived from the PHENIX corrected $\langle E_t \rangle$ value for p - p collisions from Table X ($2.6 \times 0.65 = 1.7$ GeV). That value is consistent with NSD p - p charge-particle $\langle p_t \rangle \approx 0.4$ GeV/c given $dn_{ch}/d\eta \approx 2.5$ with additional factor 1.5 to approximate total hadrons.

If free-streaming hadrons emerge from a thermalized medium, $\langle e_t \rangle$ or $\langle p_t \rangle$ is expected to relate to a temperature T that may vary with A - A centrality. However, the concept of an equilibrated temperature for Au-Au collisions below the sharp transition that appear to exhibit




FIG. 3: (a) The relation of measured charged-hadron production to Glauber-simulated number of constituent quarks in Au-Au collisions for three energies derived from the curves in Fig. 1. The points are for 200 GeV p - p collisions. (b) Per-hadron mean E_t defined as $\langle e_t \rangle = dE_t/d\eta / dn_{tot}/d\eta$, with n_{tot} (all hadrons contributing to E_t) approximated by $1.5n_{ch}$ assuming that pions dominate (approximately 85% of all hadrons). The $dE_t/d\eta$ data and uncertainties are taken from Tables V-VII of Ref. [1]. The $dn_{tot}/d\eta$ trends are derived from STAR data represented by TCM parametrizations with varying x in Fig. 1 (right panel). The solid curve is a prediction from the TCM for 200 GeV Au-Au collisions [11]. Other curves are described in the text.

negligible particle rescattering is questionable.

The $\langle e_t \rangle$ centrality trend can be predicted from the TCM based on measured spectrum properties [13] as

$$\langle e_t \rangle(\nu) \approx [1 - x(\nu)]\langle e_t \rangle_{soft} + x(\nu)\nu \langle e_t \rangle_{hard}(\nu), \quad (2)$$

where $x(\nu)$ is the same centrality-dependent parameter that describes integrated yields $n_{ch}(\nu)$, $\langle e_t \rangle_{soft}$ is the fixed mean value for soft hadron production and $\langle e_t \rangle_{hard}(\nu)$ is the mean value for the hard component that varies strongly with centrality due to modification of parton fragmentation. Below the Au-Au ST x and $\langle e_t \rangle_{hard}$ have fixed values derived from p - p collision data.

To facilitate comparisons I approximate all hadrons as pions. The effect of the pion mass is small. $\langle e_t \rangle$ is only a few percent larger than $\langle p_t \rangle$ for $\langle e_t \rangle \approx 0.5$ GeV so I increase measured $\langle p_t \rangle$ values by 0.05 GeV/c accordingly. The $\langle p_t \rangle_{soft}$ value for NSD p - p collisions (see Fig. 4 – right panel) is 0.385 ± 0.02 GeV/c, and the $\langle p_t \rangle_{hard}$ value is 1.2 ± 0.1 GeV/c. In Fig. 3 the horizontal hatched band estimates $\langle e_t \rangle_{soft}$ and the dash-dotted curve represents the GLS reference $\langle e_t \rangle$ prediction. I observe that the hadron hard-component multiplicity increases much faster than GLS above the sharp transition (vertical band ST). If $\langle e_t \rangle_{hard}$ remained fixed while $x(\nu)$ increased according to $dn_{ch}/d\eta$ measurements the predicted $\langle e_t \rangle$ trend would be the dashed curve. However, spectrum analysis reveals that $\langle p_t \rangle_{hard}$ falls from 1.2 GeV/c below the ST to 0.6 GeV/c for more-central Au-Au collisions. If that variation is also incorporated the TCM prediction for $\langle e_t \rangle$ becomes the solid curve.

The PHENIX 200 and 130 GeV Au-Au data trends are consistently 35% above the solid curve, whereas the $\langle e_t \rangle$ value for p - p collisions (solid triangle) is consistent with

STAR $\langle p_t \rangle$ data (hatched band). It is notable that the point-to-point variation of PHENIX $\langle e_t \rangle$ data follows the TCM prediction closely. The dotted curve is the TCM solid curve multiplied by factor 1.35. The TCM was not otherwise adjusted to accommodate those data.

C. Minimum-bias distributions on E_t

Given those results for the Au-Au centrality dependence of E_t mean values, Ref. [1] extends the CQ treatment to full MB distributions on E_t . Several hypotheses are considered in which E_t production scales with some element number N_X , where X can represent N - N binary collisions, participant (wounded) nucleons, constituent quarks or participant-quark-related strings. A model MB distribution on an observable such as E_t is generated by folding a MB distribution on N_x with model conditional MB distributions on E_t given N_x elements. The result is described as the Extreme Independent Model.

The A - A MB distribution on N_X is determined by Glauber Monte Carlo simulations. The measured MB distribution on E_t for p - p collisions is used to infer an equivalent for element type X denoted $f_{1X}(E_t)$. Conditional distribution $P_{N_X}(E_t)$ for N_X elements is obtained by convoluting $f_{1X}(E_t)$ multiple times and combining with calculated weights. Finally, the A - A MB distribution on E_t is obtained by convoluting the MB distribution on N_X with the $P_{N_X}(E_t)$ according to Eq. (7) of Ref. [1].

The model chosen for $f_{1X}(E_t)$ is the gamma distribution, for which a simple parameter change conveniently describes the results of multiple convolutions. Aside from the algebraic convenience the gamma distribution on E_t does not describe the p - p MB distribution well; the tail is poorly represented. The importance of accounting for void events (N - N events with no E_t in the fiducial acceptance, described by probability p_0) is emphasized.

The Au-Au Glauber MB distributions on N_{part} and N_{bin} are accurately represented by Eq. (1). The MB distribution on N_{qp} is shown in Fig. 9 of Ref. [1]. Figure 10 (b) of Ref. [1] demonstrates an unfolding of the p - p MB distribution to obtain $f_{1qp}(E_t)$. The decomposition is based on w_n numbers in Table XIII equivalent to $N_{qp} = 2.8$ for p - p collisions. That value can be contrasted with $N_{qp} = 4$ obtained by extrapolating the N_{qp}/N_{part} trend in Fig. 1 to $\nu = 1.25$ (N - N collisions). The difference is discussed in Sec. VII B. The MB endpoint on N_{part} is 382, whereas that on N_{qp} is approximately $382 \times 2.8 = 1070$. The CQM strongly disagrees with the tail of the p - p data distribution as noted. Those results are combined per Eq. (7) or (12) to obtain the model MB distribution compared with Au-Au data in Fig. 11. The model is said to describe the MB data well. The “excellent agreement” is interpreted to support the CQM.

D. The TCM as expressed within a CQM context

In the CQM context a significant role for dijet production in the MR is considered unlikely based on historical experience: “This [hard scattering in the TCM] seems to contradict the extensive measurements of N_{ch} and E_t distributions in p + p collisions described in Sec. II [of Ref. [1]] which show that these distributions represent measurements of the soft-multiparticle physics that dominates the p + p inelastic cross section” [1]. Reference [1] proposes relations between the TCM and (a) production centrality trends and (b) MB distributions on E_t based on its Eqs. (6,21) interpreted to demonstrate the unsuitability of the TCM for description of data.

The conjectured TCM for E_t production includes a single-component factor $\langle E_t \rangle^{pp}$ to represent p - p collisions

$$\langle E_t \rangle^{AA} = \langle E_t \rangle^{pp} [(1-x)N_{part}/2 + xN_{bin}], \quad (3)$$

implying that E_t production for p - p soft and hard components is the same (a common factor). If TCM parameter x in Eq. (3) also describes hadron yields then the equation reduces to $\langle e_t \rangle^{AA} = \langle e_t \rangle^{pp}$ and E_t production per hadron is assumed to be independent of A - A centrality. But those assumptions are contrary to the TCM with variable x and contradicted by differential spectrum and correlation data. Equation (3) can be contrasted with Eq. (2) (and see App. A) where soft and hard $\langle e_t \rangle$ components in p - p collisions are quite different, and the $\langle e_t \rangle$ hard component is allowed to vary with Au-Au centrality consistent with measured spectrum HC evolution [13].

The relation of the TCM to MB distributions is considered to be especially problematic. Analogies are drawn from conjectured TCM descriptions of production centrality trends. It is first assumed that the data MB distribution is the weighted sum of the limiting cases (MB on N_{part} and N_{bin}) by analogy with the TCM for integrated yields [e.g. Eq. (3) above]. One example in Figs. 17 and 18 of Rev. [1] is represented by the first line of

$$\begin{aligned} \frac{d\sigma}{dE_t} &= (1-x) \left(\frac{dE_t}{dN_{part}} \right)^{-1} \frac{d\sigma}{dN_{part}} \\ &+ x \left(\frac{dE_t}{dN_{bin}} \right)^{-1} \frac{d\sigma}{dN_{bin}}, \\ \text{or} &= \left[\frac{(1-x)dE_t}{dN_{part}} \right]^{-1} \frac{d\sigma}{dN_{part}} \\ &+ \left(\frac{x dE_t}{dN_{bin}} \right)^{-1} \frac{d\sigma}{dN_{bin}}, \end{aligned} \quad (4)$$

with the same value $dE_t/dN_x \rightarrow \langle E_t \rangle^{pp} \approx 1.8$ GeV for N_{part} and N_{bin} . Figure 19 of Ref. [1] shows an alternative hypothesis, scaling the limiting cases horizontally on $\langle E_t \rangle$ by fractions x and $1-x$ as in the second line. Since neither expression corresponds to the MB data the TCM is rejected. But the apparent disagreements between CQM conjectured TCM implementations and MB data come

not from inherent failings of the TCM but from incorrect implementations, as discussed further in Sec. VIII C.

The TCM does appear to describe production vs centrality trends accurately, as demonstrated in Sec. VI E. But that model is said to be “...just a proxy for the correct description of the underlying physics, because $dE_T^{AA}/d\eta$ is strictly proportional to N_{qp} ...” The TCM is described as an “empirical proxy” referring to the conventional RHIC TCM implementation with fixed $x \approx 0.1$ fitted to more-central Au-Au data [12]. But the TCM for production centrality dependence is not inferred empirically from a relation like Eq. (6) of Ref. [1]. It is predicted from measured p - p TCM trends and the Glauber model that serves as a null hypothesis (GLS) for more-central A - A collisions. The proper TCM for MB distributions describes MB data accurately as demonstrated in App. A, whereas the CQM cannot produce a GLS equivalent.

I now examine comprehensive yield, spectrum and correlation systematics from RHIC p - p and Au-Au collisions in the context of the TCM. I demonstrate that the TCM provides a remarkably accurate description of an extensive data phenomenology. Those data include dijet production trends that are quantitatively consistent with pQCD theory and thus inconsistent with the CQM.

V. p - p COLLISIONS AND MINIJETS

The TCM derived from p - p spectrum data and extrapolated to A - A centrality provides a reference for A - A collisions representing Glauber linear superposition. The hard-component contribution to the p - p TCM is attributed to MB dijets. Because of the steepness of the QCD parton spectrum most dijets appear near the most-probable jet energy 3 GeV (defining minijets) [6]. Issues relating to p - p collision centrality, hadron and dijet production and the TCM are discussed in [31].

A. Soft and hard events and yield n_{ch} components

p - p collisions can be classified as soft or hard event types. Soft and hard events are distinguished from soft and hard components of ensemble-averaged yields, spectra and correlations. Soft events include no jet structure within the acceptance and therefore only a soft component. Hard events include at least one minimum-bias dijet within the angular acceptance and therefore both soft and hard spectrum and correlation components.

I observe that dijet production in p - p collisions scales approximately as n_{ch}^2 [3] and is most directly related to the multiplicity soft component n_s . For a given n_s the dijet number within some η acceptance $\Delta\eta$ is $n_j(n_s) = \Delta\eta f(n_s)$, with dijet frequency $f(n_s)$ per unit η scaled from non-single-diffractive (NSD) p - p collisions. The relation between soft and hard components n_s and n_h and dijet rate $n_j(n_s)$ is established in Sec. V C. The

Poisson probabilities for soft and hard events are then respectively $P_s(n_s) = \exp(-n_j)$ and $P_h(n_s) = 1 - P_s(n_s)$. For small n_j $P_h \approx n_j$. For the PHENIX acceptance the probability of a dijet in NSD p - p collisions is $P_h \approx 0.004$.

The yields n_x defined here correspond to spectrum integrals within some angular acceptance 2π and $\Delta\eta$. For each multiplicity class defined in terms of an n_{ch} interval I have $n_s + n_h = n_{ch}$ averaged over all events. For soft events $n_s'' = n_{ch}$ and for hard events $n_s' + n_h' = n_{ch}$. I then obtain the following relations:

$$\begin{aligned} n_{ch} &= n_s + n_h = P_s n_{ch} + P_h (n_s' + n_h') \\ n_s &= P_s n_{ch} + P_h n_s' \quad \text{and} \quad n_h = P_h n_h' \end{aligned} \quad (5)$$

B. p - p single-particle spectra

Single-particle spectra from 200 GeV p - p collisions plotted on y_t for several n_{ch} classes reveal a composite spectrum structure represented by two fixed functional forms [unit-integral soft and hard components $\hat{S}_0(y_t)$ and $\hat{H}_0(y_t)$] with amplitudes scaling approximately as n_{ch} and n_{ch}^2 [3]. The combination defines the TCM for y_t spectra from p - p collisions conditional on measured n_{ch} integrated within some η acceptance $\Delta\eta$ described by

$$\begin{aligned} \rho(y_t, n_{ch}) &\equiv \frac{dn_{ch}}{y_t dy_t \Delta\eta} = S(y_t, n_{ch}) + H(y_t, n_{ch}) \\ &= \rho_s(n_{ch}) \hat{S}_0(y_t) + \rho_h(n_{ch}) \hat{H}_0(y_t), \end{aligned} \quad (6)$$

where $\rho_s = n_s/\Delta\eta$ and $\rho_h = n_h/\Delta\eta$ are soft and hard angular densities, and $\rho_0(n_{ch}) = n_{ch}/\Delta\eta$ is the corresponding total charge density. Soft component $\hat{S}_0(y_t)$ is defined as the limiting form as $\rho_s \rightarrow 0$ of spectra normalized as ρ/ρ_s . Hard component $\hat{H}_0(y_t)$ models data hard components $H(y_t, n_{ch})/\rho_s$ obtained by subtracting soft-component model $\hat{S}_0(y_t)$ from those normalized spectra.

The hard component of 1D SP spectra, interpreted as a manifestation of MB dijet structure, is quantitatively consistent with jet-related two-particle correlations and pQCD predictions [3, 11, 13, 30]. The fixed hard-component spectrum shape $\hat{H}_0(y_t)$ is predicted by measured e^+e^- fragmentation functions convoluted with a minimum-bias pQCD parton (dijet) spectrum with lower bound near 3 GeV and spectrum integral $\sigma_{dijet} = 4 \pm 1$ mb [11].

C. p - p minimum-bias dijet production

Equation (6) integrated over some angular acceptance $\Delta\eta$ becomes

$$\frac{dn_{ch}}{y_t dy_t} = n_s \hat{S}_0(y_t) + n_h \hat{H}_0(y_t). \quad (7)$$

The multiplicity trend for the extracted hard components reported in Ref. [3] implies that $n_h/n_s = \alpha n_s$ with $\alpha \approx$

0.005 within $\Delta\eta = 1$ or

$$\begin{aligned}\rho_h(n_s) &= 0.005\rho_s^2 \\ &\equiv f(n_s)\epsilon(\Delta\eta)2\bar{n}_{ch,j},\end{aligned}\quad (8)$$

where the second line represents the jet hypothesis from Ref. [3] and defines dijet frequency $f = dn_j/d\eta$ (dijet number per p - p event per unit η) with mean dijet fragment multiplicity $2\bar{n}_{ch,j}$ into 4π acceptance. Factor $\epsilon(\Delta\eta) \in [0.5, 1]$ represents the average fraction of a dijet that appears in hard events within acceptance $\Delta\eta$ [11].

Scaling from the trend in Eq. (8) I can estimate $f(n_s)$ for NSD p - p collisions

$$f_{NSD} = \frac{0.005}{\epsilon(\Delta\eta)2\bar{n}_{ch,j}}\rho_{s,NSD}^2 \approx 0.02 \pm 0.005 \quad (9)$$

assuming $\rho_{s,NSD} \approx 2.5$ for 200 GeV NSD p - p collisions and $\epsilon(\Delta\eta)2\bar{n}_{ch,j} \approx 1.5$ from p - \bar{p} measurements [3, 24]. That result can be compared with a pQCD prediction for p - p NSD collisions $f_{NSD} = \sigma_{dijet}/\sigma_{NSD} \times \Delta\eta_{4\pi} \approx 4/36.5 \times 5 = 0.022$ [11]. Thus, from a TCM analysis of the n_{ch} dependence of p - p y_t spectra I obtain the model functions $\hat{S}_0(y_t)$ and $\hat{H}_0(y_t)$, the soft and hard hadron densities ρ_s and $\rho_h \approx (0.006 \pm 0.001)\rho_s^2$ and the dijet η density per p - p event $f(n_s) = n_j(n_s)/\Delta\eta \approx 0.0035\rho_s^2$.

D. p - p mid-rapidity production trends

Figure 4 (left panel) shows the data trend that lead to Eq. (8). The points are determined by running integrals of spectrum data to obtain integrated multiplicities n_s and n_h with minimal model bias. The ratios do not depend on a spectrum model except for the soft-component Lévy distribution used to extrapolate the y_t spectra to zero momentum. The fixed soft model is common to all multiplicity classes. The solid line is $n_h/n_s = 0.005n_s/\Delta\eta$ for acceptance $\Delta\eta = 1$. The coefficient changes to 0.006 for the STAR TPC acceptance with $\Delta\eta = 2$. Given an accurate TCM reference for p - p SP spectra I can extract an analytic $\langle p_t \rangle$ trend that may provide a baseline for $\langle e_t \rangle$ data from Au-Au collisions.

Figure 4 (right panel) shows $\langle p_t \rangle$ values also extracted as limits of running integrals of SP spectra. The solid line is the TCM weighted-mean expression [see Eq. (2)] $\langle p_t \rangle = [(1-x) \times 0.38 + x \times 1.2]$ GeV/c with $x = n_h/n_s \approx 0.005n_s/\Delta\eta$, where the weights are approximately n_s/n_{ch} and n_h/n_{ch} , 0.385 GeV/c is an estimate of soft component $\langle p_t \rangle_{soft}$ and 1.2 GeV/c estimates hard component $\langle p_t \rangle_{hard}$, slightly above the hard-component peak mode at 1 GeV/c due to its skewness (QCD power-law tail). The uncertainty in the soft-component $\langle p_t \rangle$ (hatched band) is dominated by uncertainty in the spectrum extrapolation to zero momentum, since 30% of the spectrum integral lies below 0.2 GeV/c.

In conventional descriptions of scattered-parton fragmentation to collimated jets and projectile-hadron fragmentation (dissociation) to hadrons along the collision




FIG. 4: (a) Hard/soft multiplicity ratio n_h/n_s (points) vs soft component n_s consistent with a linear trend (line). Those data are the basis for Eq. (8). Assuming that n_s represents the density of small- x participant partons (gluons) and n_h represents dijet production by parton scattering, an eikonal model of p - p collision geometry (analogous to the Glauber model of A - A collisions) would predict the $n_s^{1/3}$ trend (dashed curve). (b) The $\langle p_t \rangle$ trend predicted by the p - p spectrum TCM (line) and as determined by direct spectrum integration (points). The hatched band represents the uncertainty in extrapolating spectra to zero momentum.

axis there is a direct correspondence between partons at the bottom of a fragmentation cascade and the final-state hadron distribution denoted by Local Parton-Hadron Duality (LPHD) [32]. In the LPHD description the small- x partons described by a parton distribution function or PDF should correspond quantitatively to soft hadron production at mid-rapidity, and dijet production at mid-rapidity should correspond to counterpropagating fluxes of small- x partons within the colliding hadron projectiles.

The detailed correspondence between soft and hard spectrum components represented by Fig. 4 suggests that (a) small- x gluons represented by n_s provide the underlying degree of freedom in high-energy p - p collisions and (b) the dijet production trend represented by $n_h \propto n_s^2$ falsifies the eikonal approximation as applied to p - p collisions. According to the Glauber model applied to A - A collisions and based on the eikonal approximation the expectation for p - p collisions should be $n_h/n_s \propto n_s^{1/3}$ (dashed curve in Fig. 4 – left panel). The data trend implies that each participant parton in one projectile can interact with any participant in the other projectile [31].

E. p - p two-particle correlations

Two-particle correlations for 200 GeV p - p collisions are fully consistent with the SP y_t spectrum results described above and follow the same TCM quantitatively. The correlation measure $\Delta\rho/\sqrt{\rho_{ref}}$ is a 2D density proportional to the number of correlated pairs per final-state hadron [15] and is analogous to ratio n_h/n_s assuming $n_h \rightarrow$ correlated-pair number.

Figure 5 (left panel) shows $y_t \times y_t$ correlations for 200 GeV NSD p - p collisions. The logarithmic interval

$y_t \in [1, 4.5]$ corresponds to $p_t \in [0.15, 6]$ GeV/c. The two peak features correspond to TCM soft and hard components. The 2D hard component with mode near $y_t = 2.7$ (1 GeV/c) corresponds quantitatively to the 1D SP spectrum hard component modeled by $\hat{H}_0(y_t)$. The soft component is consistent with longitudinal fragmentation (dissociation) of projectile nucleons.




FIG. 5: (Color online) (a) Minimum-bias correlated-pair density on 2D transverse-rapidity space $y_t \times y_t$ from 200 GeV p - p collisions showing soft (smaller y_t) and hard (larger y_t) components as peak structures. (b) Correlated-pair density on 2D angular difference space $(\eta_\Delta, \phi_\Delta)$. Hadrons are selected with $p_t \approx 0.6$ GeV/c ($y_t \approx 2$). Nevertheless, features expected from dijets are observed: (a) same-side 2D peak representing intrajet correlations and (b) away-side 1D peak on azimuth representing interjet (back-to-back jet) correlations.

Figure 5 (right panel) shows 2D angular correlations on difference variables (e.g. $\eta_\Delta = \eta_1 - \eta_2$). The hadron p_t values for that plot are constrained to lie near 0.6 GeV/c (just above $y_t = 2$), corresponding to the saddle between soft and hard components in the left panel. Although the hadron p_t is very low the structures expected for jet angular correlations are still clearly evident: a SS 2D peak at the origin representing intrajet correlations and a 1D peak on azimuth corresponding to interjet correlations between jets from parton collision partners. The soft component, a narrow 1D Gaussian on η_Δ including only unlike-sign charge pairs, is excluded by the $p_t > 0.5$ GeV/c cut. There are no “long-range” correlations on η corresponding to the soft component.

VI. A-A HARD-COMPONENT SYSTEMATICS

Given the detailed reference TCM from p - p collisions I now consider TCM evolution with A - A centrality and some implications for hadron and E_t production. The inferred A - A spectrum soft component is consistent with that derived from p - p spectra and appears to represent a universal transverse property of fragmentation.

A. A-A integrated yields

The TCM for p - p collisions is controlled by the SC multiplicity n_s , conjectured to represent the number of participant small- x gluons. As noted, HC production

proportional to number of participant-parton binary collisions scales as n_s^2 . The ratio of binary collisions to participants is then proportional to the number of participants, $n_h/n_s \propto n_s$ [31].

By analogy the TCM for A - A collisions is controlled by the SC scaling with participant multiplicity N_{part} . The A - A hard component HC should scale with the number of N - N binary collisions N_{bin} estimated in the Glauber model (based on the eikonal approximation) to vary as $N_{bin} \propto N_{part}^{4/3}$. For yields and spectra scaled by number of participant pairs $N_{part}/2$ the HC should vary proportional to $\nu \equiv 2N_{bin}/N_{part} \approx N_{part}^{1/3}$, consistent with the eikonal approximation (whereas p - p collisions do not follow the eikonal approximation). The TCM for A - A yields within some acceptance $\Delta\eta$ is

$$n_{ch}(\nu) = n_s \frac{N_{part}}{2} + n_h(\nu) N_{bin} \quad (10)$$

with the limiting case of Eq. (5). For the GLS reference n_h retains the fixed value from NSD p - p collisions. I now focus on HCs for A - A spectra and correlations.

B. Au-Au single-particle spectra

Figure 6 (left panel) shows Au-Au spectrum HC data for identified pions from 0-12% central 200 GeV Au-Au collisions (points). Hadron spectrum densities have the form $\rho_{0h} = d^2 n_h / 2\pi y_t dy_t d\eta$. Also shown are the p - p TCM HC model $H_{pp}(y_t)$ derived from p - p collisions (dashed curve) and the GLS prediction for central Au-Au (dash-dotted curve, 5 times dashed curve). The centrality-dependent data HC (solid dots) is obtained by

$$\nu H_{AA}(y_t, \nu) = (2/N_{part}) \rho_{0\pi}(y_t, \nu) - S_{NN}(y_t), \quad (11)$$

where $S_{NN}(y_t)$ is by definition the limiting case of $(2/N_{part}) \rho_{0h}(y_t, \nu)$ as $\nu \rightarrow 0$ (no dijet production) [13]. As noted, $S_{NN}(y_t)$ is consistent with the shape of the SC model $\hat{S}_0(y_t)$ inferred from p - p collisions. For the GLS reference H_{AA} retains the fixed form $H_{NN} \approx H_{pp}$ independent of A - A centrality and the HC scales with ν .

Above $y_t = 4$ ($p_t \approx 4$ GeV/c) the central-Au-Au pion spectrum is suppressed by factor 5, consistent with conventional R_{AA} measurements, and thus happens to coincide with the p - p hard component. At lower momenta (below $y_t = 3.3$ or $p_t = 2$ GeV/c) the hard-component data rise far above the GLS reference, which fact is obscured by R_{AA} due to severe bias by inclusion of the spectrum SC in that measure. The solid curve is a pQCD calculation based on a measured MB parton spectrum folded with parametrized FFs [11]. The same calculation for p - p collisions (dashed curve) describes the p - p hard-component data (open circles) quantitatively at the 10% level. For more-central Au-Au collisions the FF parametrization is modified (one parameter is changed by 10%). The parameter change is equivalent to changing a gluon splitting function in the DGLAP equations [9].

The theory description (solid curve) is then accurate over the entire y_t acceptance. On that basis there can be little doubt that the TCM pion spectrum HC is jet related even in central Au-Au collisions.




FIG. 6: (Color online) (a) Hard components from 0-12% central 200 GeV Au-Au collisions for identified pions (solid points [13]) and from 200 GeV p - p collisions (open points [3]). The dashed and dash-dotted curves show the GLS references for p - p and Au-Au collisions. The solid curve is a pQCD description of Au-Au HC data assuming a slight modification of e^+e^- fragmentation functions [11]. (b) Spectra for identified protons (green curves) from several centralities of Au-Au collisions. The red dotted curve is the TCM soft component. The dashed curve is the TCM proton hard component inferred from Au-Au spectrum systematics. The dash-dotted curves show GLS references for peripheral and central Au-Au collisions. The data hard components for more-central collisions show an increase relative to GLS above the HC peak mode at $y_t = 2.7$ (1 GeV/c) in contrast to the identified-pion result.

Figure 6 (right panel) shows full-spectrum data for identified protons. The proton spectra (thin solid curves) are compared with inferred spectrum SC $S_{NN}(y_t)$ (dotted curve) common to all centralities. The dashed curve is the p - p -equivalent proton HC model H_{NN} derived as the limiting case of the Au-Au HC for $\nu \rightarrow 1$ (isolated N - N collisions). It is remarkable that the proton HC amplitude near $y_t = 2.7$ ($p_t \approx 1$ GeV/c) is comparable to the pion HC amplitude at the same point, although the spectrum SCs are very different [13].

The proton HC evolution with Au-Au centrality is formally similar to that for pions in the sense that strong reduction at larger y_t is compensated by enhancement at some smaller y_t , but for protons the enhancement occurs only above 1 GeV/c (approximately the proton mass). The dash-dotted curves marked GLS are defined as $S_{NN} + \nu H_{NN}$ with $\nu = 1.25, 5.8$. The spectrum data for protons in central collisions cross the GLS curve at $y_t \approx 4$ (4 GeV/c), whereas the crossing for pions is at $y_t \approx 3$ (1.3 GeV/c). And below 1 GeV/c the proton HC is consistent with *no FF modification*, follows the GLS.

The difference in evolution between proton and pion HCs fully accounts for the so-called baryon-meson (B/M) puzzle conventionally attributed to CQ coalescence. These results suggest that the B/M puzzle is actually an aspect of parton fragmentation possibly related to hadron fragment mass [13]. The centrality evolution of proton

spectra also strongly supports interpretation in terms of minimum-bias jets.

C. Au-Au number angular correlations

Figure 7 shows 2D number angular correlations from 200 GeV Au-Au collisions for all $p_t > 0.15$ GeV/c and for peripheral (left panel, 85-95%) and central (right panel, 0-5%) collisions. In both cases a SS 2D peak and AS 1D peak on azimuth are evident as for p - p collisions. Absent the restrictive p_t cuts invoked for Fig. 5 (right panel) a narrow 2D peak attributed to conversion-electron pairs and Bose-Einstein correlations (at the origin) and a 1D peak on η_Δ (soft component, left panel only) appear. In the right panel the SS 2D jet peak in central collisions is strongly elongated on η_Δ . A nonjet quadrupole component proportional to $\cos(2\phi_\Delta)$ is not evident for these centralities but does appear for mid-central data. The z -axis zeros are defined by 2D-model fits to data in which it is assumed that the AS 1D peak is positive definite and the nonjet quadrupole has zero mean [15].




FIG. 7: (Color online) (a) Angular correlations from 85-95% central 200 GeV Au-Au collisions equivalent to NSD N - N collisions. The soft component is a 1D peak on η_Δ . The hard components are a SS 2D peak and AS 1D peak on azimuth. Conversion electrons and Bose-Einstein correlations contribute a narrow exponential peak at the origin confined to $p_t < 0.5$ GeV/c. (b) The same for 0-5% central Au-Au collisions. The SS 2D peak is strongly elongated on η_Δ . The soft component falls to zero by mid-centrality. The nonjet quadrupole (v_2) amplitude is negligible [18, 19].

Figure 8 shows the centrality dependence of the SS 2D peak (left panel) and AS 1D peak (right panel) amplitudes for 200 GeV (solid points) and 62.4 GeV (open points). The GLS reference for those per-hadron fit-model parameters is the dashed curves defined by $A_{X,pp} \nu / [1 + 0.02(\nu - 1)]$, where the numerator is consistent with the HC from p - p correlations and the denominator is the GLS hadron-yield centrality trend. The data follow the GLS trend up to the ST at $\nu \approx 3$ ($\sigma/\sigma_0 \approx 0.5$). Within a single centrality bin the slopes increase dramatically, and the amplitudes continue to increase rapidly until $\nu \approx 5$ beyond which the data appear to fall off.

In the left panel the 62.4 GeV A_{2D} data multiplied by factor 1.6 coincide with the 200 GeV data. In Ref. [22] it was observed that the energy dependence of p_t cor-




FIG. 8: (a) The SS 2D peak amplitude vs centrality measured by pathlength ν for two collision energies [15]. The 62.4 GeV data coincide with 200 GeV data when multiplied by factor 1.6, as discussed in the text. The data follow the GLS reference (dashed curve) up to $\nu = 3$ (sharp transition or ST) after which the slope changes by factor 3 or more. (b) Equivalent results for the AS 1D peak amplitude. The 62 and 200 GeV data correspond for more-peripheral collisions without additional factor. That relation is expected for dijets.

relation amplitudes attributed to MB jets appeared to vary as $\log(\sqrt{s_{NN}})$ and to extrapolate to zero amplitude near $\sqrt{s_{NN}} = 10$ GeV suggesting an energy dependence $\approx \log(\sqrt{s_{NN}}/Q_0)$ with energy scale $Q_0 \approx 10$ GeV. The jet-related amplitudes in the left panel are consistent with that conjecture since $\log(200/9)/\log(62.4/9) = 1.6$. The energy trend suggests that dijet production drops to zero near 10 GeV due to kinematic constraints on parton (mainly gluon) fragmentation to charged hadrons [11].

The AS peak amplitudes for two energies in the right panel coincide over most centralities with no scale factor. That is expected if the energy dependence is due to the increased kinematic range of colliding small- x partons with increasing collision energy. On 2D dijet rapidity space (y_{z1}, y_{z2}) a dijet is represented by a single point. I speculate that while the kinematic boundaries of that space may expand with increasing collision energy the dijet density remains approximately the same, and the AS 1D peak represents the unchanging 2D dijet density. The SS 2D peak amplitude represents a 1D projection of that space onto its difference axis and therefore does scale with $\log(\sqrt{s})$ changes in the kinematic boundary on y_z .

D. Au-Au p_t angular correlations

The plots in Figs. 5 (right) and 7 represent hadron number angular correlations. Angular correlations of hadron p_t can be obtained either by inversion of the scale dependence of $\langle p_t \rangle$ fluctuations [17, 21] or by direct pair counting. The same jet-related correlation structures are observed, with minor quantitative differences in the SS 2D peak structure. Such results indicate that $\langle p_t \rangle$ fluctuations, once expected to reveal critical fluctuations of temperature near a QCD phase boundary, are actually

dominated by a MB jet (minijet) contribution.




FIG. 9: (Color online) Upper: p_t angular correlations for (a) 85-95% and (b) 10-20% central 200 GeV Au-Au collisions inferred by inverting $\langle p_t \rangle$ fluctuation scale dependence [21]. AS dipole and nonjet quadrupole components of 2D model fits to the data have been subtracted. Lower: Results for the same collision systems but the p_t correlations are obtained by direct pair counting rather than fluctuation inversion. Improved angular resolution is evident as well as unfiltered statistical fluctuations.

Figure 9 (upper panels) shows p_t angular correlations for (a) 85-95% and (b) 10-20% central 200 GeV Au-Au collisions obtained by scale inversion of $\langle p_t \rangle$ fluctuations [17]. Such 2D angular correlations are described by a standard fit model including several elements [15, 21]. In this case fitted AS dipole and nonjet quadrupole components have been subtracted to isolate the SS 2D peak structure. A similar analysis of HIJING Monte Carlo data confirms a jet interpretation for that structure [33].

Figure 9 (lower panels) shows p_t angular correlations for the same collision systems obtained by direct pair counting, confirming the results in the upper panels obtained by fluctuation scale inversion. The inversion process smooths the data (regularization): statistical fluctuations are reduced but angular resolution is also reduced. The SS 2D peak for p_t correlations is narrower than that observed for number correlations [compare panel (c) with Fig. 5 – right]. That difference is expected for jet correlations, since fewer fragments with larger momenta are found closer to the jet thrust axis and more fragments with smaller momenta appear at larger angles.

Two features of the SS 2D peak in p_t correlations are especially notable. The negative-going region on either side of the peak near the origin suggests the possibility of a recoil component. These are covariance densities, and negative covariance corresponds to anticorrelation of momenta. The other feature is the “ridge” especially obvious in panel (d). The strong p_t correlations at

larger η_Δ argue against η elongation of the SS 2D peak in more-central Au-Au collisions residing within a soft background. Related studies of number angular correlations with p_t cuts indicate that the SS peak η elongation persists up to 4 GeV/c hadron momentum [19], a trend not likely to arise from a soft process.

Thus, in several ways the hard components of 2D number and p_t angular correlations from Au-Au collisions follow trends expected for dijets. Dijet production closely follows the predicted GLS trend over half the Au-Au total cross section then undergoes a substantial change within a small centrality interval, described as the sharp transition. Above the ST the correlation structure continues to follow expectations for parton fragmentation to jets, but with fragmentation substantially modified.

E. Correspondence of spectra and correlations

Further support for a jet interpretation of the TCM HC is provided by quantitative correspondence between spectrum and correlation data based on a dijet hypothesis. In Fig. 10 (left panel) the solid curve describes the number of dijets (at least one jet) appearing within angular acceptance $\Delta\eta = 2$ (the STAR TPC acceptance) as a function of mean participant-nucleon pathlength ν . The p - p value $n_j(NSD) \approx 0.04$ is obtained from Eq. (9). The solid curve is the Au-Au GLS reference $n_j(\nu) = n_j(NSD)N_{bin}$. The hatched band labeled ST indicates the position of the 200 GeV Au-Au sharp transition, which happens to correspond to dijet η density $f \approx 1$. The dashed curve and left-hand hatched band are discussed in App. B3.

Figure 10 (right panel) shows 200 GeV HC yields predicted from jet-related angular correlations summed with a measured soft-component contribution S_{NN} (solid curve [30]) compared to yields obtained directly from spectra (points [13]). The volume of the SS 2D peak in angular correlations is processed using the dijet number per event in the left panel to derive the number of correlated pairs per dijet. The number of correlated pairs is converted to the number of fragments per dijet and multiplied by n_j to obtain the predicted fragment number per event νH_{AA} . When that HC is added to SC S_{NN} I obtain the TCM prediction for charged-hadron yields. The agreement with integrated yields from hadron spectra (points) is within systematic uncertainties (upper hatched band). The dash-dotted line shows the conventional 200 GeV Au-Au TCM with fixed $x = 0.095$ [12]. The lower hatched band is the GLS reference TCM with $x = 0.015$ derived from p - p spectrum data. The dashed curve shows a prediction based on CGC theory [12, 34]. This comparison provides additional strong support for a dijet interpretation of the HC of spectra and correlations for all Au-Au centralities.




FIG. 10: (a) Dijet number within angular acceptance $\Delta\eta = 2$ at mid-rapidity for 200 GeV Au-Au collisions scaling with N_{bin} from Eq. (1) (GLS reference, solid curve). The horizontal hatched band indicates the measured dijet rate for NSD p - p collisions in the same acceptance. The upper-right hatched band indicates the sharp transition (ST) near $\nu = 3$ for jet correlations in Au-Au collisions. The dashed curve and left-most hatched band are discussed in App. B3. (b) Per-participant hadron production measured by $(2/N_{part})dn_{ch}/d\eta$ vs ν for 200 GeV Au-Au collisions (solid points) inferred from analysis of identified-hadron spectra [13]. The dash-dotted line is the conventional TCM with fixed $x = 0.095$ [12]. The solid curve is a prediction obtained by analysis of jet-related angular correlations based on the n_j solid curve in the left panel [30]. The dashed curve is a prediction derived from color-glass condensate theory [12, 34].

VII. CHALLENGING THE CQM

I now confront the conjectured CQM with experimental evidence summarized in the previous two sections and arguments based on the historical development of QCD and logical structure.

A. Arguments from historical concepts

Some supporting arguments for the CQM appear to be based on historical concepts of hadron production in HE nuclear collisions. It is notable that in Ref. [1] 31 references appear after 1990 whereas 48 appear before that date [note the relation to LEP commissioned in 1989 and the Hadron-Electron Ring Accelerator (HERA) commissioned in 1992]. The HERA e - p collider experiments provided a large volume of accurate data on hadron (proton) internal structure, including parton distribution functions (PDFs). The LEP e^+e^- collider experiments provided a large volume of accurate data on parton FFs over a broad range of dijet energies and parton types. The two data volumes, combined with pQCD theory, accurately and comprehensively describe hadron production in HE nuclear collisions, including dijets [11, 24].

Historical issues can be considered in a context where the full 4π solid angle is divided into a limiting-fragmentation (LF) region near the beam rapidity and a mid-rapidity (MR) region near the collision center of momentum. With the discovery of partons (mainly va-

lence quarks) in lower-energy e - p collisions at SLAC and the initial development of QCD the search commenced for hard parton scattering as a transport process from LF to MR in HE collisions. Hard parton scattering to jets should occur rarely, and the evidence should appear as transport from LF to MR. Jet-related hadron production in the MR should be sparse and highly structured.

As recounted in Ref. [1], contrary to expectations copious E_t and hadron production was actually observed in the MR. However, the production appeared to be weakly structured on angle and approximately thermal in terms of the fixed relation of E_t to hadron multiplicity. That unexpected result lead to the conclusion that production in the MR must be dominated by soft processes such as QCD string fragmentation [35], with negligible jet contributions. The strings were thought to connect colored valence quarks/diquarks or dressed constituent quarks (valons [14]). That historical context apparently prompted Ref. [1] to the conclusion that inference of copious (semi)hard scattering to the MR (minijets) “seems to contradict the extensive measurements...[of E_t and n_{ch}]...which show that these distributions represent measurements of the soft-multiparticle physics that dominates...” p - p collisions. As a consequence, the study in Ref. [1] is restricted to “...the nucleon and constituent-quark participant models of soft-multiparticle production widely used since the 1970’s...”

B. Arguments from p - p phenomenology

The importance of projectile small- x partons (mainly gluons) to hadron production in the MR has already been mentioned in Sec. IIB. Here I review some detailed aspects of p - p data phenomenology pertaining to that issue. The CQM must contend with the accurately-measured n_{ch} dependence of p - p collisions and copious dijet production for larger event multiplicities. Aside from substantial MB dijet production at mid-rapidity another major issue is the dominant role of fluctuations.

The integrated MR charge $\rightarrow n_{ch}$ in 200 GeV p - p collisions fluctuates event-wise over a large interval (at least a factor ten), which is problematic for a CQM with 2-3 participant quarks N_{qp} per projectile and hadrons dominated by string fragments proportional to that number. The resolved-dijet production scales approximately as n_{ch}^2 , leading to a factor 100 variation in dijet number. For 200 GeV p - p collisions with $dn_{ch}/d\eta \approx 25$ (10 \times NSD) the mean number of dijets within $\Delta\eta = 2$ is approximately two *per event*, and fragments from resolved jets comprise 10% of all hadrons. A string/CQ model cannot generate dijets at that level (or at all) and cannot describe the large range of multiplicity fluctuations, whereas measured projectile small- x structure and corresponding dijet production based on perturbative QCD can do so given event-wise fluctuations in the parton splitting cascade within a proton [31].

Figure 10 of Ref. [1] provides another indicator.

The gamma-distribution model function representing the CQM (solid curve) deviates strongly from the tail of the p - p data distribution that represents large fluctuations in E_t production. When applied in Eq. (7) to construct the CQ MB model function as a convolution the result actually fails to represent the MB data. Close examination of Fig. 11 at the terminus reveals that the CQM (red curve) has a much larger slope (smaller fluctuation width) than the data (blue points). The apparent agreement of the MB distributions arises directly from the relation shown in Fig. 1 (right panel) of the present study. The substantial deviations for more-peripheral collisions that may falsify the CQM are not visible in the conventional (and insensitive) semilog plotting format.

Use of a Glauber Monte Carlo to estimate the p - p N_{qp} as in Fig. 1 seems problematic. In Sec. VD I note that the trend $N_{bin} \propto N_{part}^2$ (with $N_{part} \sim n_s$) that describes dijet production in p - p collisions in terms of small- x parton participants is inconsistent with the eikonal approximation which corresponds to $N_{bin} \propto N_{part}^{4/3}$. But the Glauber model is based on the eikonal approximation and cannot therefore be used to model p - p collisions in terms of conjectured CQ participants. The value $N_{qp} = 2.8$ inferred from a p - p Glauber simulation is inconsistent with the value $N_{qp} \approx 4$ obtained in Fig. 1 by extrapolating the Au-Au trend (where the eikonal approximation and Glauber model are valid) to p - p collisions for $\nu \approx 1$. That is a significant issue because the value $N_{qp} = 2.8$ is supposed to indicate self-consistency of the CQM as in Sec. VIII-B of Ref. [1].

C. Arguments from A - A phenomenology

The CQM appears to be relevant because the simulated N_{qp} centrality trend approximates the A - A centrality trends for n_{ch} and E_t , at least for more-central collisions. However, major problems arise for N_{qp} from more-differential analysis. Below the sharp transition near $\nu = 3$ Au-Au spectra and correlations follow a TCM reference (GLS) quantitatively equivalent to that describing p - p collisions, with n_s replaced by N_{part} as the SC control parameter and n_h/n_s replaced by ν as a measure of N - N binary collisions [15]. The number of resolved dijets n_j having correlation structure consistent with in-vacuum pQCD jets increases from 0.04 per event to about 2 per event within the STAR TPC acceptance, scaling with the number of N - N binary collisions as expected for dijets. The same jet correlation structure appears in both number and p_t angular correlations. Substantial deviations from the N_{qp} trend for more-peripheral collisions are already evident in Fig. 1 (right panel). If the CQM is excluded by the dijet presence in p - p collisions it is excluded in more-peripheral Au-Au: the spectrum HC remains quantitatively related to pQCD dijets.

In more-central collisions above the sharp transition the dijet contribution is altered quantitatively, but the spectrum HC is still described by pQCD modulo a simple

modification of FFs that conserves the parton energy and remains consistent with the general form of the DGLAP equations. The same basic correlation structure persists even in central collisions and still scales with N_{bin} , with *no actual reduction from the expected dijet number*. In central 200 GeV Au-Au collisions I find that 1/3 of the final state is included within resolved dijets [15, 30], dramatically inconsistent with the CQM

Reference [1] states that the form of $(2/N_{part})dn_{ch}/d\eta$ vs N_{part} is statistically equivalent at 200 GeV and 2.76 TeV (ALICE data) “...although the jet cross section increases by a very large factor.” The conclusion is drawn that hadron production must then be dominated by soft processes with no significant dijet contribution. But careful examination of the comparison reveals several issues, as established in App. B2. The shape of the centrality trend for PHENIX data is substantially distorted by fluctuations for more-central collisions due to the small angular acceptance. The sharp transition at 200 GeV near $\nu = 3$ appears to be significantly lower at 2.76 GeV. When the comparison is made by scaling the ALICE data down by the same factor 2.1 the more-peripheral 2.76 TeV data fall well below the measured 200 GeV NSD p - p numbers. Thus, a combination of several mitigating factors may conspire in the original comparison to give a false impression. I find that a TCM scaled up by factors 1.8 and 1.8^2 derived from RHIC energy systematics below 200 GeV predicts the ALICE data trend within rather small data uncertainties. The TCM is actually strongly supported by that comparison.

Reference [1] argues that dijets are strongly suppressed in more-central Au-Au collisions (jet quenching). So a TCM increase in hadron production due to dijets would be inconsistent with jet suppression inferred from R_{AA} data. However, spectrum data presented in Sec. VIB reveal that the actual jet-related hadron production integrated over the full p_t acceptance *increases* with centrality by up to a factor six. The noted suppression at higher p_t is compensated by a much larger enhancement at lower p_t (so as to conserve the parton energy), still within the collimated jet structure [11, 13]. That conclusion is supported by jet-related angular correlation trends [15].

D. Arguments from p-A phenomenology

Several lessons from p-A collision data that appear to support the TCM are drawn in Ref. [1]. A projectile nucleon (participant) interacting with a target nucleus may be excited (wounded) only once and thereafter does not change its internal state in subsequent N - N collisions. Because of time dilation the excited nucleon cannot fragment within the target nucleus. The projectile (p) hemisphere is independent of A , and the target (A) hemisphere depends on ν , the mean path length of the projectile in the target. The formation-time argument appears to eliminate the possibility for rescattering of secondary partons or hadrons: “This feature [formation

time] immediately eliminates the possibility of a cascade in the nucleus from rescattering of the secondary products” [1].

Certain implications follow for A - A collisions. Absence of rescattering would explain why 3 GeV minijets survive intact without significant loss in central Au-Au collisions [15]. It would also explain the absence of collective (radial and elliptic) flows indicated by recent observations [13, 18, 19]. Soft hadron production by linearly-independent projectile-nucleon dissociation would then scale as N_{part} with properties independent of A - A centrality. Dijet production would scale as the number of N - N binary collisions N_{bin} . And the number of dijets per participant nucleon would scale with ν adopted from the p- A phenomenology. But that is exactly the A - A TCM.

E. Arguments from logical structure

Reference [1] seeks to verify what may be called hypothesis A: Almost all hadron and transverse momentum/energy production in the MR results from soft hadron production following N_{qp} scaling. The paper establishes that A implies certain data trends B: (i) $(2/N_{qp})dE_t/d\eta$ should be approximately constant with Au-Au centrality and (ii) the MB distribution on N_{qp} may be used to generate a model distribution on E_t that should describe MB data. Evidence apparently supporting B is provided and it is concluded that hypothesis A is valid. I note that (i) and (ii) are mathematically equivalent, and results B rely on an approximate relation between N_{qp} and $dn_{ch}/d\eta$ that may be accidental.

The logical structure of such an argument is known to be questionable.¹ Other hypotheses A' may also imply B, and other results B' may falsify A. Observation of B would indeed imply \bar{A} , falsification of the hypothesis. But observation of B does not imply or require A, nor does it falsify alternatives A' (e.g., the TCM). Examples of phenomena B' that falsify A include the n_{ch} systematics of dijet production in p - p collisions, agreement with the GLS trend below the ST in Au-Au collisions, strong jet-related correlations for all collision systems, and the central role of N_{bin} HC scaling in all collision systems.

In contrast, the TCM is implied by modern QCD theory combined with the measured properties of hadrons (PDFs from HERA) and dijets (FFs from LEP). The TCM is required by the phenomenology of p - p and A - A collisions, including differential spectrum and correlation measurements. The TCM is not an empirical description of data (as conventionally represented). It is a quantitative prediction based on QCD theory and data from elementary collisions in the form of the GLS reference. In contrast, the CQM makes no such quantitative predic-

¹ The structure is an example of “undistributed middle” fallacy.

tions and fails to describe data trends for more-peripheral A - A collisions. The CQM assumes no hard scattering, so copious evidence for hard scattering from spectra and correlations appears to falsify the CQM.

F. Summary

With the premise that hadron production in the MR must be controlled by large- x degrees of freedom in the LF represented by conjectured constituent quarks, that hadron production from jets must be sharply reduced with increasing A - A centrality as implied by R_{AA} data, and that “...possible models motivated by the fact that half of the momentum of a nucleon is carried by gluons when probed at high Q^2 in hard scattering are not considered...” [1], the CQM seems to be the only alternative. But the description of high-energy nuclear collisions during the past twenty years has become increasingly reliant on just the small- x partons excluded from the CQM. And the improvement in analysis methods applicable to dijet structure since first RHIC operation is substantial.

I do observe that a soft component compatible with projectile nucleon dissociation plays a major role in hadron production complementary to dijet production, but the SC is easily described by a few universal parameters, with form independent of A - A centrality. Without a dijet contribution the CQM must rely on string fragmentation as the sole transport mechanism from LF to MR. The CQ concept and string fragmentation are non-perturbative, and the CQM can thus make no connection to pQCD theory. Direct evidence from several differential methods for copious dijet production in the MR, described quantitatively by pQCD for all A - A centralities, then appears to exclude the CQM. In contrast, the TCM describes a large variety of data at the few-percent level over a broad range of collision energies from SPS to LHC.

VIII. DISCUSSION

In this section I consider further aspects of hadron production mechanisms and methods intended to reveal them, including measure sensitivity, model energy dependence, the utility of MB distribution modeling and the role of fluctuations in MB distributions and production centrality trends.

A. Sensitivity to hadron and E_t production

The main question addressed in this study relates to the sensitivity of certain aspects of data to collision mechanisms. What mechanisms (or models thereof) can be tested by what aspects of the data? The ability to test models relies on the amount of information in the data, which in turn depends strongly on how the data are presented (statistical analysis and plotting format).

The greatest information is carried by multiparticle correlations within narrow multiplicity or centrality bins. Successive integrations (and therefore information loss) lead to two-particle correlations (a limiting case), fluctuations (integrals of correlations), spectra, full MB distributions and yields within centrality bins. The information carried by MB distributions is small compared to that in differential 2D correlations and 1D spectra.

Aside from the inherent information content of the data the choice of plotting format can determine the fraction of such information that is visually accessible. The peripheral centrality region is conventionally disregarded, by the choice of centrality interval for actual measurements (the top 40, 65 or 80% are common choices) and plotting format (N_{part} de-emphasizes the peripheral region). In contrast, a per-participant measure plotted vs ν provides an ideal format for testing the TCM hypothesis.

Use of p_t instead of y_t de-emphasizes the low- p_t region where major jet-related variations with A - A centrality occur. Spectrum ratio R_{AA} nominally measuring jet structure is insensitive to jets below 4 GeV/c because of the spectrum SC which dominates the ratio below that point, whereas TCM analysis of spectra reveals large HC contributions extending down to 0.5 GeV/c or lower. MB distributions plotted in a semilog format on E_t , N_{part} or n_{ch} convey *no significant information* visually. The same data plotted in a power-law format over the full centrality range convey all available information, including variation of TCM parameter x with centrality and fluctuations for central ($b = 0$) A - A collisions and possibly p - p collisions.

In Ref. [1] the key data plots are Figs. 3, 5, 6, 7 and 16, with 3 and 16 being semilog plots vs E_t . Such data are said to “provide excellent characterization of the nuclear geometry...and are sensitive to the underlying reaction dynamics....” But that claim is not born out by the E_t data. Figs. 5 and 6 on N_{part} are replotted on ν as Figs. 2 and 3 of the present paper. (Fig. 7 is effectively an integral of Fig. 6 and as such carries less information.) Whereas the ST is a prominent feature of the centrality dependence of correlations, spectra and integrated hadron yields from Au-Au collisions consistent with the TCM and QCD, there is no evidence for the ST in the E_t data as plotted, no sensitivity to that important new phenomenon. Part of the problem is the large systematic uncertainties, especially for more-peripheral collisions, presumably arising from uncertainty in collision centrality and therefore correct values for N_{part} or N_{qp} . Thus, although the TCM and CQM strongly disagree in more-peripheral collisions the data presented in Ref. [1] cannot test the model differences.

The composite nature of E_t presents another issue. The E_t integrated within some acceptance can be factored into the integrated hadron multiplicity n_{ch} and mean E_t per hadron denoted by $\langle e_t \rangle$. A similar argument pertains to $\langle p_t \rangle$ fluctuations [20]. Given the information extracted from n_{ch} distributions I seek unique aspects of E_t production beyond hadron multiplicities. Such in-

formation may be carried in per-hadron mean values as shown in Fig. 3. Unfortunately, the large systematic uncertainties preclude effective model tests, especially regarding detailed structure predicted by the TCM and accessed by differential spectrum analysis. One may conclude that in the CQM context there is no new information from E_t measurements beyond hadron multiplicities.

B. Utility of MB distribution modeling

Reference [1] models the MB distribution on E_t based on convolution of a model for the E_t distribution from p - p collisions and a MB distribution on N_{qp} derived from a Glauber simulation. The procedure is described as the Extreme Independent Model. The model is compared to data in Fig. 11 and “excellent agreement” is reported. However, one can question the utility of such a procedure.

Equation (7) of Ref. [1] represents a convolution integral that gives a MB distribution on E_t given conditional distributions on E_t for a given number of N_x collisions and a MB distribution on N_x , with the form

$$\frac{d\sigma}{dE_t} = \sum_{N_x} \frac{d\sigma}{dN_x} P(E_t|N_x) \quad (12)$$

where the $P(E_t|N_x)$ are derived by convoluting gamma distributions fitted to p - p data. In the CQM the implicit assumption is invoked that hadron and E_t production remain the same for all A - A centralities. Only N_{qp} varies.

There is an exact equivalence between a MB frequency distribution on quantity X integrated within some angular acceptance and the centrality variation of the produced X . A running integral of the MB distribution gives $\sigma(X)$ vs X . Combined with $N_{part}/2$ and ν vs σ from a Glauber MC as in Eq. (1) the running integral gives $(2/N_{part})X$ vs ν . That procedure is used in App. A 2 to determine the distortion effect of fluctuations at the terminus of the MB distribution on the per-participant production centrality trend, as in Fig. 15 (right panel).

N_{qp} is observed to have approximately the same trend as n_{ch} assuming fixed TCM parameter $x \approx 0.1$ as in Fig. 1 (right panel). Thus, the MB distribution on N_{qp} is equivalent to that on n_{ch} within a rescaling of the x axis. In App. C I obtain a correspondence between MB distributions $d\sigma/dE_t \approx (dE_t/dN_{qp})^{-1} d\sigma/dN_{qp}$, where Jacobian $dE_t/dN_{qp} \approx 2/3$ GeV is approximately independent of N_{qp} as indicated in Fig. 2 (right panel). The same argument holds for the relation between n_{ch} and N_{qp} , where the Jacobian is $dN_{qp}/dn_{ch} \approx 1.5$. Finally, I have $dE_t/dn_{ch} \approx 1$ GeV to close the circle.

Thus, the form of the CQM MB distribution on E_t is determined to good approximation by that on n_{ch} and the equivalent on N_{qp} . Any MB distribution within that “family” can be generated from the TCM by the method described in App. A 1. The only difference is the Jacobian factor. The terminus half-maximum values are then related as follows (referring to figures in Ref. [1]):

$N_{part}/2 = 191$ [Eq. (1)], $N_{qp} = 2 \times 2.8 \times 191 = 1070$ (Fig. 9) $E_t = 0.66 \times 1070 = 710$ (Fig. 11 – upper panel). The comparison between CQM distribution and data in Fig. 11 of Ref. [1] is said to indicate “excellent agreement,” including the matched terminus positions. But the general MB model shape *must* describe the data as implied by Fig. 1 (right panel), and the endpoints are determined by Jacobians already established elsewhere.

The only unique information in a MB distribution within this family of kinematic quantities, beyond their production trends on centrality, is the slope at the terminus representing fluctuations in central collisions. Close examination of the terminus shapes in Fig. 11 reveals substantial disagreement. The model slope ($\propto 1/\sigma_{E_t}$) is much larger than the data, indicating that fluctuations are underestimated. But the CQM assumes the same production mechanisms for p - p collisions and central Au-Au collisions, whereas the correlation structure in the two cases is very different [15], and fluctuations reflect (are integrals of) that correlation structure [17].

There are substantial differences between the CQM and the TCM (and the data it describes accurately), but those differences (reflected for instance by significant variation of the Jacobians noted above) are concealed by the insensitive semilog format invoked for conventional MB distribution plots. Thus, I can conclude that the Extreme Independent Model of Ref. [1] does not reveal additional information beyond what is accessible from production centrality trends, visually suppresses significant model differences within the semilog plotting format and cannot describe fluctuations in central Au-Au collisions. Its utility is therefore questionable.

C. The TCM MB distribution in a CQM context

In Sec. IV D an argument is presented that the TCM fails to model MB data distributions and should therefore be rejected. However, the TCM implementation is incorrect as demonstrated in this subsection. The general problem relates to the 3D space (X, N_x, N_y) where X is some produced quantity such as E_t and N_x , N_y are parameters modeling A - A geometry. The MB density on (X, N_x, N_y) may be normalized to the total cross section σ_0 or to unity. For this discussion I choose the latter and discuss probabilities in terms of $P = \sigma/\sigma_0$.

The 3D space can be projected to three 1D marginal spaces. One marginal space dP/dX is directly measurable and the other two are approximated by Glauber simulations. The running integrals in those spaces provide parametric relations, e.g. $X(P)$, $N_x(P)$ and $N_y(P)$. Mean values of the three quantities are parametrically related through common parameter P to define a curve or locus of means in the 3D space. Those relations provide the basis for inferred production centrality trends such as shown in Fig. 10 (right panel). I choose one model parameter $N_x \rightarrow N_{part}$ as the basic degree of freedom for the TCM. I empirically observe simple power-law trends

such as $N_{part}^{1/4}(P) \propto P$ as expressed in Eqs. (1). Given that framework I consider details of the TCM as manifested in different contexts.

The MB distribution on X can be related to the MB distribution on one of the geometry parameters through a Jacobian derived from the parametric relations, as in Eq. (C3). That representation is advantageous because of the simple power-law structures on N_{part} and N_{bin} and the TCM formulation of the parametric relations as in Eq. (11). The result is the correct form of the TCM for the MB distribution on X as described in App. A, with direct comparisons to data. I find that the TCM accurately describes all the experimentally accessible features of MB data distributions to their uncertainty limits.

The conjectured relations in Eq. (4) represent an alternative implementation of the TCM for MB distributions. Dividing both sides by σ_0 I obtain probability densities with some interesting features. First, the density in one marginal space is related to a sum of densities in two other marginal spaces. In terms of probabilities that implies a dichotomy: In each event either participants *or* binary collisions are related to E_t . Second, the Jacobians relating E_t to N_{part} and N_{bin} are both assumed to be equal to fixed value $\langle E_t \rangle^{pp}$. Neither aspect seems plausible. I conclude that Eq. (4) is an improper implementation of the TCM for MB distributions, and TCM comparisons with MB data as in Ref. [1] are misleading.

D. CQM, TCM and mean- p_t trends at the LHC

The CQM assumes that charge number n_{ch} , p_t and E_t production near midrapidity are proportional to the number of constituent-quark participants N_{qp} representing string fragmentation as the underlying mechanism [1]. The trend of N_{qp} with A - A centrality revealed by a Glauber MC simulation then produces the characteristic centrality trends in those quantities previously attributed to the TCM (with fixed x parameter) including dijet production by hard scattering. N_{qp} scaling then implies that ratios $\langle p_t \rangle$ and $\langle E_t \rangle$ should be nearly constant with A - A centrality, and also implies that $\langle p_t \rangle$ should be nearly constant with varying n_{ch} in p - p collisions. Recent LHC data [36] contradict the CQM expectation but are accurately described by the TCM [37] as summarized below.

Ensemble $\langle p_t \rangle$ is defined by $\langle p_t \rangle = P_t/n_{ch}$ where total p_t denoted by P_t is integrated within some angular acceptance and n_{ch} is the total charge integrated within the same acceptance. Just as $n_{ch} = n_s + n_h$ in the TCM we assume $P_t = P_{t,s} + P_{t,h}$. We then have $P_t = n_s \langle p_t \rangle_s + n_h \langle p_t \rangle_h$ and the TCM for p - p $\langle p_t \rangle$ is

$$\begin{aligned} \langle p_t \rangle_{pp}(n_s) &= \frac{n_s \langle p_t \rangle_s + n_h \langle p_t \rangle_h}{n_s + n_h} \\ &= \frac{\langle p_t \rangle_s + x(n_s) \langle p_t \rangle_h(n_s, \sqrt{s})}{1 + x(n_s)}, \end{aligned} \quad (13)$$

where $x(n_s) = n_h/n_s$ is defined in Sec. V D. The two $\langle p_t \rangle$

components can be inferred from spectrum model functions $S_0(y_t)$ and $H_0(y_t)$ or from $\langle p_t \rangle$ vs n_{ch} data. The p - p $\langle p_t \rangle$ data show rapid increase with n_{ch} , inconsistent with the CQM but quantitatively consistent with the TCM.




FIG. 11: (a) $\langle p_t \rangle$ vs n_{ch} data for several energies from the SPS, RHIC and LHC [3, 6, 36]. The curves through data are defined by Eq. (13). (b) $\langle p_t \rangle$ hard components inferred from the data in panel (a).

Figure 11 (left panel) shows LHC p - p $\langle p_t \rangle$ data (top three data sets [36]) and those from UA1 [6] and STAR [3] (lower three data sets). The curves through data are defined by Eq. (13). The soft component $\langle p_t \rangle_s$ has fixed value 0.385 GeV/c in the full p_t acceptance but shifts upward with partial p_t acceptance as for the LHC data (to ≈ 0.5 GeV/c). The shift in $\langle p_t \rangle_s$ can be calculated given the *effective* p_t acceptance. The comparison between Eq. (13) and data then determines hard component $\langle p_t \rangle_h(n_s, \sqrt{s})$. Figure 11 (right panel) shows $\langle p_t \rangle_h(n_s, \sqrt{s})$ values inferred for several collision energies.




FIG. 12: (a) n_{ch} -averaged hard-component $\langle p_t \rangle_h(\sqrt{s})$ values vs jet spectrum width Δy_{max} demonstrating the linear correlation that confirms a dijet origin for the p - p spectrum hard component, as established in Ref. [11]. (b) p -Pb and Pb-Pb $\langle p_t \rangle$ data from the LHC [36] (points) compared to the $\langle p_t \rangle$ TCM (curves). The p -Pb data for larger n_{ch} are well described by a Glauber linear superposition (GLS) form of the TCM consistent with transparent p -Pb collisions (no rescattering). For smaller n_{ch} the p -Pb data correspond quantitatively to p - p collisions at the same energy.

Figure 12 (left panel) shows the average $\langle p_t \rangle_h(\sqrt{s})$ values inferred from the data in Fig. 11 (right panel) (dashed lines) plotted vs the quantity $\Delta y_{max}(\sqrt{s})$ measuring the

widths of MB jet spectra vs p - p collision energy [38]. The linear relation demonstrates that there is an exact correspondence between p - p p_t spectrum hard components and jet spectrum widths as expected if spectrum hard components are convolutions of jet fragmentation functions and MB jet spectra, as demonstrated in Ref. [11].

Figure 12 (right panel) shows an extension of the p - p $\langle p_t \rangle$ TCM of Eq. (13) to A - A collisions (solid curve) compared with corresponding LHC data. Again, the TCM provides an excellent data description and furnishes new insights into collision mechanisms [37]. Conventional Monte Carlo models cannot describe the LHC data [36]. One can compare the right panel of Fig. 12 with Fig. 3 (right panel) to see that the LHC Pb-Pb data are consistent with the Au-Au downturn to the TCM p - p value (solid triangle) but inconsistent with the CQM.

IX. SUMMARY

Based on comparisons of Monte Carlo simulations for the number of constituent-quark participants N_{qp} with the measured centrality trends of integrated multiplicity n_{ch} and transverse energy E_t it is argued that the two-component (soft+hard) model (TCM) of mid-rapidity production, including a substantial dijet contribution, is actually a proxy for a constituent-quark (CQ) model (CQM) in which dijets play no significant role. Further support for the CQM is derived from comparison of an Extreme Independent Model of minimum-bias (MB) distributions based on N_{qp} with MB data and comparisons of RHIC and LHC production trends that seem very similar despite the large energy difference. Hadron and E_t production near mid-rapidity is seen as arising from fragmentation of QCD color strings joining CQs residing at large momentum fraction x within projectile nucleons.

In the present study I confront arguments supporting the CQM with contrasting differential evidence from yields, spectra and correlations that provide strong support for the TCM and the major role played by dijets in p - p and A - A collisions near mid-rapidity. The TCM relates directly to pQCD predictions and provides detailed quantitative descriptions of a broad range of phenomena in correlations, spectra, yields and MB distributions.

Differential spectrum and correlation measurements from 200 GeV p - p collisions reveal a dijet contribution quantitatively consistent with pQCD predictions based on measured jet spectra and fragmentation functions, the production depending on the number of small- x partons (mainly gluons) represented by a soft multiplicity component n_s . The dijet rate (hard component) is proportional to n_s^2 representing parton-parton binary collisions and indicating that the eikonal approximation is not valid for p - p collisions. Fluctuations in the soft multiplicity over a ten-fold range correspond to variations in the dijet rate over a hundred-fold range, including multiple dijets per collision at the upper end. The p - p phenomenology provides the reference for a TCM in A - A collisions repre-

sented Glauber linear superposition of N - N collisions (GLS).

In A - A collisions the role of soft multiplicity n_s is assumed by participant-nucleon number N_{part} , and N - N binary collisions are represented by N_{bin} with ratio n_h/n_s replaced by $\nu = 2N_{bin}/N_{part}$. The A - A TCM based on those parameters is observed to follow the GLS reference over half the Au-Au total cross section for 62.4 and 200 GeV collisions, with dijet structure as in p - p collisions. In more-central Au-Au collisions, above a sharp transition in the hard-component centrality trend, the dijet structure is observed to change quantitatively but remains consistent with a slightly-modified pQCD description. Although some dijet properties change the number of dijets remains proportional to N_{bin} . No dijets are lost.

The measured energy dependence of the TCM for Au-Au collisions below 200 GeV can be used to predict data trends at LHC energies. The hadron production vs centrality for Pb-Pb at 2.76 TeV is described within small data uncertainties by the extrapolated TCM.

In contrast the CQM is excluded by data in several ways. The p - p multiplicity dependence of spectrum and correlation structure, which clearly requires a dijet contribution, cannot be described by the soft-only CQM. Careful comparison of N_{qp} with integrated n_{ch} Au-Au centrality trends for more-peripheral collisions reveals substantial discrepancies (factor six difference in slopes below the sharp transition). The detailed evolution of Au-Au spectrum structure and angular correlations cannot be explained by a homogeneous model of soft production. The exact correspondence of jet-related correlated pair number and spectrum integrals based on a pQCD dijet frequency is also incompatible with the CQM.

A conjectured algebraic relation between the TCM and MB distributions invoked to reject that model is found to be invalid. The correct algebraic relation between the TCM for production centrality trends and MB distributions is established in the present study. The TCM description of MB data is accurate and provides new insights into detailed structure, including consequences of production fluctuations in central A - A collisions. In contrast, the Extreme Independent Model relating the CQM to MB distributions is shown to be misleading. The apparent agreement with data results from the insensitivity of the conventional semilog plotting format. The only real model test is the role of fluctuations in central collisions (shape of MB tail structure), which the CQM fails.

Can MB distributions on E_t and n_{ch} test hadron production models? The answer is yes provided several conditions are met: (i) The MB plotting format must make all information visually accessible, (ii) distortions from production fluctuations in more-central collisions must be understood, (iii) systematic uncertainties relating to centrality determination in more-peripheral collisions must be controlled, (iv) a proper linear-superposition reference extrapolated from p - p collisions must be established and (v) a mathematically correct TCM model of MB distributions must be defined to extract quantitative data.

That being said, analysis of MB distributions however accurate must compete with more-differential methods (spectra and correlations) that provide much more information about collision dynamics. Collision models have already been rigorously tested by a combination of highly-differential spectrum and correlation analysis. The TCM prevails as a general framework within which model details can be evaluated, and dijet production is an essential feature of any model of nuclear collisions.

I conclude that the TCM is not a misleading proxy for a more-legitimate soft model based on constituent quarks. Instead, the CQM is a proxy based on a single accidental relation and is falsified by a complex of differential analysis results described accurately by a TCM consistent with the pQCD description of dijets.

This material is based upon work supported by the U.S. Department of Energy Office of Science, Office of Nuclear Physics under Award Number DE-FG02-97ER41020.

Appendix A: TCM and MB distributions

Figures 17-19 of Ref. [1] show conjectures about a relation between the TCM and MB data distributions. The figures show comparisons between MB data distributions on E_t and the extreme cases of pure participant or N - N binary-collision scaling (Figs. 17 and 18) as discussed in Sec. IV D. Since both hypotheses fail to describe the data the TCM is rejected. But the descriptions fail not from problems with the TCM but from misapplication of the TCM to MB distributions as discussed in Sec. VIII C. In this Appendix I derived the correct relation between the TCM and MB distributions on E_t and n_{ch} .

1. The TCM and MB distribution structure

Since $\langle e_t \rangle$ in Fig. 3 (right panel) is nearly constant with centrality the power-law treatment of MB distributions on n_{ch} in Ref. [23] is a good approximation to the present case on E_t . I assume some acceptance $\Delta\eta$ with integrated total charge n_{ch} . Reference [23] notes that the MB distribution on N_{part} is almost exactly a power law $\propto N_{part}^{-3/4}$ and is therefore approximately constant on $N_{part}^{1/4}$ within some bounded interval (rectangular). The structure of the measured MB distribution on $n_{ch}^{1/4}$ (deviations from the rectangular participant-scaling distribution) may directly reveal three aspects of hadron production: (a) fluctuations in p - p collisions, (b) fluctuations in central A - A collisions and (c) the parameter x in the TCM, measured respectively by the slopes on the left and right ends and in the central part of the MB distribution.

The data MB distribution can be predicted quantitatively from the TCM in the following way. I assume the usual TCM expression for the charge yield $(2/N_{part})n_{ch} = n_{pp}[1 + x(\nu - 1)]$, where n_{pp} is the charge

yield for p - p (N - N) collisions. Then invoking Eq. (C3) with power-law variables I obtain

$$\frac{d\sigma}{dn_{ch}^{1/4}} = \left[\frac{dn_{ch}^{1/4}}{d(N_{part}/2)^{1/4}} \right]^{-1} \frac{d\sigma}{d(N_{part}/2)^{1/4}}, \quad (\text{A1})$$

where the first factor is a Jacobian that can be derived from the basic TCM yield expression and the second factor is the rectangular MB distribution representing participant scaling. The MB distribution on $n_{ch}^{1/4}$ is then

$$\frac{d\sigma}{dn_{ch}^{1/4}} = \frac{[1 + x(\nu - 1)]^{3/4}}{1 + x(\nu - 1) + x\nu/3} \times \frac{d\sigma}{n_{pp}^{1/4} d(N_{part}/2)^{1/4}}, \quad (\text{A2})$$

where I have invoked the approximation $\nu \approx (N_{part}/2)^{1/3}$ [23]. If x is constant for all centralities the first factor (inverse Jacobian) is approximately a straight line on ν with negative slope ($\approx 1 - 7x\nu/12$) reflecting some fractional contribution to $n_{ch}(N_{part})$ from N - N binary-collision scaling. The x inferred from differential spectrum data increases by a factor six from peripheral ($x \approx 0.015$) to central ($x \approx 0.1$) Au-Au collisions [3, 13, 15, 30]. The slope of the first factor then varies accordingly. Note that in Eq. (A5) of Ref. [23] the quantity $\nu^{5/4}/3$ should be $\nu/3$ as in Eq. (A2).




FIG. 13: (a) MB distributions on $n_{ch}^{1/4}$ derived from Glauber Monte Carlo simulations [23] assuming participant scaling (dotted lines) and the conventional TCM with fixed $x = 0.1$ (dashed lines) according to Eq. (A2). (b) The same trends plotted in the conventional MB semilog format on n_{ch} .

Figure 13 (left panel) shows MB distributions on $n_{ch}^{1/4}$ for (a) participant scaling (dotted lines, $x = 0$) and (b) the TCM describing more-central 200 GeV Au-Au collisions (dashed lines, fixed $x = 0.1$ for all centralities). Both distributions integrate to the Au-Au total cross section 7.2 barns. The sloping dashed line represents Eq. (A2). In that plotting format the relation of distribution (b) to collision geometry and production mechanisms is easy to demonstrate (as discussed below). Figure 13 (right panel) shows the same distributions in the conventional semilog plotting format, with Jacobian factor $dn^{1/4}/dn = 1/4n^{3/4}$. Connections to the underlying

particle production mechanisms and collision geometry are not visually accessible. The straightforward TCM implementation in Eq. (A2) and Fig. 13 is dramatically different from the conjectures in Figs. 17-19 of Ref. [1].

Figure 14 (left panel) shows a comparison between STAR 130 GeV h^- data (solid curve [39]) and the corresponding TCM of Eq. (A2) (dashed curve) with constant $x = 0.08$ and $n_{pp} = 2.25/2$ (approximate h^- η density near $\eta = 0$ for 130 GeV). The dotted curve represents participant scaling. A similar comparison appears in Fig. 3 of Ref. [23]. This semilog plotting format obscures essential data features relating to the TCM.




FIG. 14: (a) MB distribution of negative hadrons h^- from 130 GeV Au-Au collisions (solid curve [39]) compared to power-law participant scaling (dotted curve) and the TCM with fixed $x = 0.08$ (dashed curve). The dash-dotted curve is discussed in the text. (b) The same distributions plotted in power-law format. Information in the data is only the more-central mean slope (reflecting the TCM x value for more-central collisions) and the slope at the terminus reflecting hadron production fluctuations in Au-Au collisions with $N_{part,max} = 382$. The step in the dashed curve at mid-centrality is a schematic model of the ST (see the text).

Figure 14 (right panel) shows the same distributions in the power-law format. The dotted lines represent participant scaling and the dashed lines (for more-central collisions) represent Eq. (A2) with $x = 0.08$. Each distribution integrates to $\sigma_0 = 7.2$ barns. The model curves extend over the full fractional centrality interval $\sigma/\sigma_0 \in [0, 1]$. The MB data and TCM agree to a few percent, well within systematic uncertainties. The TCM depends only on parameters $n_{pp} \equiv 2.25/2$ and $x = 0.08$ and otherwise was not fitted to the MB h^- data. The right-hand dash-dotted line in the right panel is a good approximation to the tail structure in the left panel when transformed to that semilog format (dash-dotted curve).

The power-law format provides a precise estimate of the end-point multiplicity as the half-max point on the terminus (steeply falling part) of the MB distribution at the right end. That quantity is not accessible with the conventional semilog plotting format. For these data the endpoint lies at $n_{h-} = 4.15^4 = 300$. From the corresponding Au-Au participant endpoint $N_{part}/2 = 191$ I obtain $(2/N_{part})n_{h-} = 1.57$. Given $n_{h-,pp} = 2.25/2 = 1.12$ for 130 GeV NSD p - p collisions I obtain $1+x(\nu-1) = 1.4$ at $\nu \approx 6$, implying $x \approx 0.08$. Thus, the terminus end-

point position accurately corresponds to TCM parameter x and the slope of the dashed curve described by Eq. (A2), at least for more-central collisions.

Also included in the right panel is a schematic model of the ST observed at 62.4 and 200 GeV. The dashed curve for more-peripheral collisions follows the GLS trend described by Eq. (A2) with $x = 0.02$. Extrapolated to central collisions the endpoint in that case would be $[1.1 \times (2.25/2) \times 191]^{1/4} = 3.92$ or $n_{h-} = 236$. Near the ST at mid-centrality parameter x increases from 0.02 to 0.08 (short vertical line), with endpoint $n_{h-} = 300$ as noted.

From these comparisons I find that one learns little from MB distributions in the conventional semilog plotting format but a substantial amount from data in the power-law format. Relative to the power-law Glauber model one can estimate p - p fluctuations, central A - A fluctuations and centrality evolution of TCM parameter x . The consequence of dijet production in A - A collisions (TCM hard component) is redistribution of the A - A cross section to larger integrated yields according to Eq. (A2).

2. Production fluctuations in central collisions

Fluctuations in particle and E_t/p_t production produce a distortion of MB distributions. Differences in detector angular acceptances can lead to corresponding differences in MB distributions. The terminus slope in the power-law format (e.g. Fig. 14, right-hand dash-dotted line) represents production fluctuations in central collisions ($b \approx 0$). The terminus slope m is proportional to $1/\sigma_n$ measuring charge multiplicity fluctuations (the variance depends on the angular acceptance). The mean slope of the right-hand terminus is $|m| \approx 10$, and the Gaussian r.m.s. on $n_{h-}^{1/4}$ is $\sigma_n = 2/\sqrt{2\pi}|m| \approx 0.08$. The corresponding fluctuation manifestation on n_{h-} is estimated by the dash-dotted curve in the left panel, with relative r.m.s. on n_{h-} $4 \times \sigma_n/300^{1/4} \approx 0.08$ or 8% for charge-number correlations within acceptance $\Delta\eta = 1$. Production fluctuations in p - p collisions from the terminus sketched by the left dash-dotted line aren't accessible because these MB data do not extend low enough on $n_{h-}^{1/4}$.

The terminus fluctuation structure in the MB distribution has major consequences for plots of production vs centrality. I can determine fluctuation effects in the latter using the exact correspondence between a power-law MB distribution and the corresponding plot of production on centrality.

Figure 15 (left panel) shows the TCM model for 200 GeV Au-Au collisions (dash-dotted lines) with $x = 0.095$. The dotted curve represents that curve folded with a Gaussian (approximated by an error function) having a fluctuation width corresponding to the STAR TPC acceptance (see Fig. 14 – right panel). For the PHENIX EMCal acceptance the corresponding width is three times larger (solid curve). I transform MB fluctuation models to production centrality trends as follows. The running integral of a curve in the left panel yields $\sigma(n_{ch}^{1/4})$ vs

$n_{ch}^{1/4}(\sigma)$. I divide $n_{ch}(\sigma)$ by $N_{part}(\sigma)/2$ and convert σ/σ_0 to ν using Eqs. (1) to obtain the required production centrality dependence $(2/N_{part})dn_{ch}/d\eta$ vs ν .




FIG. 15: (a) Models of MB frequency distributions on n_{ch} in the power-law format including no fluctuations (dash-dotted curve), fluctuations for the STAR TPC acceptance (dotted curve) and for the fiducial PHENIX EMCAL acceptance (solid curve), both corrected to the reference acceptance. (b) Measured production centrality trends for STAR (solid dots) [13] and PHENIX (open squares) and “RHIC average” (open triangles) [40] compared to integrated model MB distributions from the left panel (same line styles). The large effect of angular acceptance on fluctuation distortions for more-central collisions is demonstrated.

Figure 15 (right panel) shows the integration results as the solid and dotted curves plotted above $\nu = 4.5$. The solid curve below $\nu = 4.5$ represents the ST trend as derived from jet-related angular correlations and shown in Fig. 10 (right panel). Corresponding PHENIX and “RHIC average” production data are plotted as open squares and open triangles respectively [40] compared to STAR spectrum integrals (solid dots) [13]. This exercise demonstrates that the ten-times smaller PHENIX EMCAL angular acceptance results in a substantial fluctuation distortion or bias for more-central collisions, equivalent to about 18% relative fluctuations compared to 6% for STAR within $\Delta\eta = 2$. The consequence for STAR data is a single most-central point falling slightly above the fixed- x TCM dash-dotted line.

Comparing the right panels of Figs. 14 and 15 I observe that a MB distribution on produced quantity X is more differential than the X production centrality trend, but also includes substantial statistical noise. The running integral leading to the production centrality trend does filter the noise to better reveal the information. But the same noise reduction may be accomplished by rebinning the MB distribution (uniform bins in the power-law format). Both formats convey the same information.

Appendix B: TCM vs Collision energy

Reference [1] presents arguments that the TCM, including its hard component representing dijet production, is actually a proxy for soft production scaling with the number of constituent-quark participants. LHC

hadron production data [36] are invoked to argue that the centrality trend for hadron production from 200 GeV Au-Au is essentially the same as that from 2.76 TeV Pb-Pb collisions, although “the jet cross section increases by a very large factor.” Thus, hadron production at both energies must be dominated by soft processes with negligible dijet contribution. However, the detailed structure of the data trends and the predicted energy evolution of the TCM actually support the TCM interpretation.

1. Predicting the energy evolution of the TCM

Two features of 2D angular correlations at RHIC have been observed to scale with collision energy as $\log(\sqrt{s_{NN}}/Q_0)$. For the nonjet quadrupole $Q_0 \approx 13.5$ GeV [18]. For dijet production *per final-state hadron* in p_t correlations $Q_0 \approx 10$ GeV [22]. A jet-production lower bound near $Q_0 \approx 10$ GeV is consistent with the observed jet-energy lower bound 2 - 3 GeV for parton fragmentation to charged hadrons [11]. For jet-related number correlations I observe that $\log(\sqrt{s_{NN}}/9 \text{ GeV})$ predicts factor 1.6 in Fig. 8 (left panel) relating dijet production per hadron in Au-Au collisions at 200 vs 62.4 GeV [15] and predicts factor 2.2 relating equivalent dijet structure in p - p collisions at 200 GeV vs 7 TeV [41]. The corresponding factor relating 200 GeV to 2.76 TeV is 1.8.

Hadron production in p - p collisions inferred from y_t spectrum n_{ch} dependence is described in Sec. VD: (a) small- x gluons estimated by soft yield n_s provide the common underlying degree of freedom in high-energy N - N collisions, and (b) the dijet production trend is indicated by hard yield $n_h \propto n_s^2$. If the relative density of small- x gluons produced at a given N - N collision energy scales proportional to $\log(\sqrt{s}/Q_0)$ and $n_h/n_s \propto n_s$ then observed dijet energy trends are explained quantitatively.

The TCM centrality trend for 200 GeV Au-Au collisions is defined by $2.5[1 + x(\nu - 1)]$ with GLS $x \approx 0.02$ below the ST and modified-FF $x \approx 0.095$ above the ST. According to the above argument the TCM for 2.76 TeV is predicted by scaling up the soft component (increased small- x gluons) by factor 1.8 and the hard component *relative to the soft component* by another factor 1.8. The absolute increase in dijet production should then be factor $1.8^2 = 3.24$ (increased collisions of small- x gluons).

2. Charged-hadron production at two energies

Figure 16 (left panel) shows per-participant-pair charge production data from 2.76 TeV Pb-Pb collisions (inverted solid triangles) and p - p collisions (upright solid triangle) [36]. To accommodate the 6% larger atomic number of lead the path length is scaled up by 2% since $\nu \propto N_{part}^{1/3}$ is a good approximation. The TCM limiting cases are defined as described above by $1.8 \times 2.5[1 + 1.8 \times x(\nu - 1)]$ with $x = 0.02$ (dashed line) and $x = 0.095$ (dash-dotted line). The solid curve is the 200 GeV centrality

trend inferred from jet-related correlations (reflecting the ST) scaled up in the same way. Given the relation to data the 2.76 GeV hadron production data are *predicted* by a TCM based on the energy dependence of soft and hard production below 200 GeV. The TCM was not adjusted to accommodate the 2.76 TeV data.




FIG. 16: (a) Hadron production vs centrality for 2.76 TeV Pb-Pb collisions (inverted solid triangles [36]) compared to TCM trends (dash-dotted and dashed lines) extrapolated from the 200 GeV TCM based on measured RHIC energy trends as described in the text. The upright solid triangle denotes p - p data. (b) Hadron production vs centrality for 200 GeV Au-Au collisions (open triangles [40]) compared to a TCM prediction from analysis of jet-related angular correlations (dotted curve) [30]. The dash-dotted line indicates the conventional 200 GeV Au-Au TCM with fixed $x = 0.095$ [12]. The dashed line indicates a GLS extrapolation from p - p collisions with $x = 0.015$. The solid triangles and solid line are the solid triangles and dash-dotted line in the left panel scaled down by factor 2.1. The two line slopes differ by factor 1.55.

Figure 16 (right panel) shows the “RHIC average” hadron production data appearing in Fig. 15 (open triangles, reported in Ref. [40]) compared to the 200 GeV TCM. The inverted solid triangles are the 2.76 TeV data in the left panel scaled down by factor 2.1. The apparent similarity has been invoked to support the conclusion that particle production follows the same general trend at two very different energies and therefore must be due exclusively to a soft process, inconsistent with the TCM [1]. In that comparison the 200 GeV data were actually scaled up by factor 2.1 to match the 2.76 TeV data, but this reversed-scaling comparison is equivalent.

The claimed similarity of data trends at two different energies is interpreted to imply a common soft hadron-production mechanism with no significant dijets from hard processes, but the comparison is misleading. The scaled 2.76 TeV data actually strongly disagree with the 200 GeV production data from Ref. [13] (dotted curve) and the 200 GeV TCM in several ways: (i) The average slope of the scaled higher-energy data (solid line) is $1.8^2/2.1 = 1.55$ times larger than the STAR 200 GeV data trend and TCM (dash-dotted line). The difference greatly exceeds what is allowed by various systematic uncertainties. (ii) The apparent agreement for more-central data results from the substantial fluctuation bias in the PHENIX data, whereas that bias is small for the ALICE

acceptance. The unbiased limiting case is the TCM dash-dotted line. (iii) The downscaled data for peripheral 2.76 TeV Pb-Pb and p - p collisions fall a factor $1.8/2.1 = 0.86$ below the 200 GeV NSD value 2.5 and inferred S_{NN} , again well beyond data uncertainties. In contrast, the 200 GeV TCM scaled appropriately for small- x parton participants and parton-parton binary collisions *predicts* a TCM in the left panel that describes the 2.76 TeV data within their uncertainties. Thus, the recent LHC data strongly favor a TCM based on small- x parton energy dependence extrapolated from RHIC data trends.

This energy comparison has interesting implications. The hadron production data at 2.76 TeV are described accurately by a TCM derived from 200 GeV data with two adjustments: The soft component is multiplied by factor 1.8, the hard component by factor $1.8^2 = 3.24$. The factor 1.8 is inferred from a $\log(\sqrt{s}/Q_0)$ trend (with $Q_0 \approx 9$ GeV) inferred from RHIC data below 200 GeV and interpreted to represent production of participant small- x gluons (soft component) in p - p or N - N collisions which may then collide as binary pairs to produce dijets (hard component). The result in Fig. 16 (left panel) suggests that modifications to jet formation in Pb-Pb collisions at 2.76 TeV are *essentially identical* to those in Au-Au collisions at 62.4 and 200 GeV. The only difference may be a shift of the ST to more-peripheral collisions. Substantial FF modifications above the ST observed at RHIC energies apparently remain unchanged.

3. Possible energy evolution of the sharp transition

The sharp transition first observed for minimum-bias jet-related angular correlations in Au-Au collisions and indicated by the hatched box ST in Fig. 16 (right panel) [15] is a remarkable new feature of hadron production systematics. Below the ST hadron production (spectra and correlations) follows the GLS reference extrapolated from p - p collisions. Above the ST hadron production within the hard component increases dramatically, the increase attributed mainly to modified parton fragmentation [11]. While fragment yields are suppressed at larger p_t as indicated by ratio R_{AA} they are much more enhanced at lower p_t (the increase concealed by the properties of R_{AA}) but remain within the correlated jet structure and conserve the parton energy [13, 30]. The result is an increase in hadron fragments *within intact jets* of up to a factor 6 and no reduction in dijet number [11].

It could be argued that the most significant information in Fig. 16 (left panel) is the possible migration of the ST downward in centrality with increasing collision energy. In Fig. 10 (left panel) the ST at 200 GeV (right hatched box) corresponds to a dijet η density per Au-Au collision (solid curve) of approximately unity. The dashed curve is the solid curve scaled up by factor $1.8^2 \approx 3.25$ corresponding to the predicted absolute increase in dijet yield from 200 GeV to 2.76 TeV. The left hatched box indicates the centrality for the same dijet

density per Au-Au collision given the increased dijet rate per N - N collision. Comparing the two panels of Fig. 16 I find that the 2.76 TeV data are consistent with a shift in the ST downward from $\nu \approx 3$ to $\nu \approx 2$ consistent with the exercise in Fig. 10. That correspondence suggests that the ST and modified parton fragmentation may be determined by the dijet density within A - A collisions.

Appendix C: Joint MB distributions

MB distributions such as Fig. 2 of Ref. [1] are related to a system of joint, conditional and marginal distributions of cross-section σ or event-number density on some measured kinematic quantity X (e.g. n_{ch} or E_t) and one or more possible A - A model parameters N_x derived from the 3D space (X, N_x, N_y) by projection. The 3D density distribution is typically tightly correlated (localized) as expressed by a locus of mean values $(\bar{X}, \bar{N}_x, \bar{N}_y)$ realized for instance by Eq. (10) for N_{part} , N_{bin} and $X = n_{ch}$. Projections onto subspaces and running integrals within subspaces are used to determine the locus of mean values and detailed features of projected MB distributions.

It is useful to establish an algebraic context for such projections, for example based on the integrated E_t within some angular acceptance vs a model parameter N_x such as $N_{part}/2$ or N_{qp} . The 2D joint MB distribution $F(E_t, N_x)$ may be normalized to unity (joint probability distribution) or to A - A total cross section σ_0 . Applying the chain rule for joint probabilities I obtain

$$\begin{aligned} F(E_t, N_x) &= \hat{G}(E_t|N_x)F(N_x) \\ &= \hat{H}(N_x|E_t)F(E_t), \end{aligned} \quad (C1)$$

where for instance $\hat{G}(E_t|N_x)$ is the unit-normal conditional probability density distribution on E_t given a specific N_x condition, and $F(E_t)$ is the marginal projection of 2D $F(E_t, N_x)$ onto E_t with

$$F(E_t) = \int dN_x \hat{G}(E_t|N_x)F(N_x). \quad (C2)$$

Equation (C2) describes Fig. 2 of Ref. [1] where $\hat{G}(E_t|N_x)$ represents the individual lower curves and $F(E_t)$ is the upper solid curve. In Eq. (7) of Ref. [1] $\sigma_{BA} w_n \leftrightarrow F(N_x)$ and $P_n(E_t) \leftrightarrow \hat{G}(E_t|N_x)$. For generality I treat $F(N_x)$ as a density on continuous N_x which could be a sum of delta functions at integer values.

Each conditional distribution $\hat{G}(E_t|N_x)$ is a density on E_t peaked near some conditional mean value $\bar{E}_t(N_x)$. On the other hand, for a given E_t the ensemble of $\hat{G}(E_t|N_x)$ can be seen as a distribution on N_x peaked near some mean value $\bar{N}_x(E_t)$. The conditional distribution in the argument of the integral in Eq. (C2) can be converted to a density on N_x with a suitable Jacobian $J(E_t, N_x)$. Treating $J(E_t, N_x)\hat{G}(E_t|N_x)$ as a weight function on N_x the integral in Eq. (C2) can be interpreted as a weighted average of $F(N_x)/J(E_t, N_x)$ at given E_t approximated

by that ratio evaluated at $\bar{N}_x(E_t)$. With $J(E_t, N_x) = dE_t/dN_x$ I then obtain

$$F(E_t) = [dE_t/dN_x]_{\bar{N}_x}^{-1} F[\bar{N}_x(E_t)], \quad (C3)$$

where the Jacobian dE_t/dN_x is a variable quantity derived from the locus (curve) of mean values $\bar{E}_t(N_x)$. Analysis of MB distributions indicates that in cases relevant to A - A collision geometry the details of conditional distributions $\hat{G}(E_t|N_x)$, other than the conditional mean values $\bar{E}_t(N_x)$, are not relevant except at the ends (termini) of the $F(N_x)$ distribution [23].

-
- [1] S. S. Adler *et al.* (PHENIX Collaboration), Phys. Rev. C **89**, 044905 (2014).
 - [2] S. Eremín and S. Voloshin, Phys. Rev. C **67**, 064905 (2003).
 - [3] J. Adams *et al.* (STAR Collaboration), Phys. Rev. D **74**, 032006 (2006).
 - [4] R. J. Porter and T. A. Trainor (STAR Collaboration), J. Phys. Conf. Ser. **27**, 98 (2005).
 - [5] R. J. Porter and T. A. Trainor (STAR Collaboration), PoS **CFRNC2006**, 004 (2006).
 - [6] C. Albajar *et al.* (UA1 Collaboration), Nucl. Phys. B **309**, 405 (1988).
 - [7] D. Buskulic *et al.* (ALEPH Collaboration), Z. Phys. C **55**, 209 (1992).
 - [8] M. Z. Akrawy *et al.* (OPAL Collaboration) Phys. Lett. B, **247**, 617 (1990).
 - [9] N. Borghini and U. A. Wiedemann, hep-ph/0506218.
 - [10] B. B. Back *et al.* (PHOBOS Collaboration), Phys. Rev. C **65**, 061901 (2002).
 - [11] T. A. Trainor, Phys. Rev. C **80**, 044901 (2009).
 - [12] D. Khazeev and M. Nardi, Phys. Lett. B **507**, 121 (2001).
 - [13] T. A. Trainor, Int. J. Mod. Phys. E **17**, 1499 (2008).
 - [14] R. C. Hwa and M. S. Zahir, Phys. Rev. D **23**, 2539 (1981).
 - [15] G. Agakishiev, *et al.* (STAR Collaboration), Phys. Rev. C **86**, 064902 (2012).
 - [16] J. Adams *et al.* (STAR Collaboration), Phys. Rev. C **73**, 064907 (2006).
 - [17] T. A. Trainor, R. J. Porter and D. J. Prindle, J. Phys. G **31**, 809 (2005).
 - [18] D. T. Kettler (STAR collaboration), Eur. Phys. J. C **62**, 175 (2009).
 - [19] D. Kettler (STAR Collaboration), J. Phys. Conf. Ser. **270**, 012058 (2011).
 - [20] J. Adams *et al.* (STAR Collaboration), Phys. Rev. C **71**, 064906 (2005).
 - [21] J. Adams *et al.* (STAR Collaboration), J. Phys. G **32**, L37 (2006).
 - [22] J. Adams *et al.* (STAR Collaboration), J. Phys. G **33**, 451 (2007).
 - [23] T. A. Trainor and D. J. Prindle, hep-ph/0411217.
 - [24] T. A. Trainor and D. T. Kettler, Phys. Rev. D **74**, 034012 (2006).
 - [25] X.-N. Wang, Phys. Rev. D **46**, R1900 (1992); X.-N. Wang and M. Gyulassy, Phys. Rev. D **44**, 3501 (1991).
 - [26] T. Sjöstrand and M. van Zijl, Phys. Rev. D **36**, 2019 (1987); T. Sjöstrand, Comput. Phys. Commun. **82**, 74 (1994); T. Sjöstrand, L. Lönnblad, S. Mrenna and P. Skands, hep-ph/0308153.
 - [27] M. Bahr, S. Gieseke, M. A. Gigg, D. Grellscheid, K. Hamilton, O. Latunde-Dada, S. Platzer and P. Richardson *et al.*, Eur. Phys. J. C **58**, 639 (2008).
 - [28] E. W. Oldag (STAR Collaboration), J. Phys. Conf. Ser. **446**, 012023 (2013).
 - [29] J. Adams *et al.* (STAR Collaboration), Phys. Lett. B **634**, 347 (2006).
 - [30] T. A. Trainor and D. T. Kettler, Phys. Rev. C **83**, 034903 (2011).
 - [31] T. A. Trainor, Phys. Rev. D **87**, 054005 (2013).
 - [32] Ya. I. Azimov, Yu. L. Dokshitzer, V. A. Khoze, S. I. Troyan, Z. Phys. C **27**, 65 (1985), Z. Phys. C **31**, 213 (1986).
 - [33] Q. J. Liu, D. J. Prindle and T. A. Trainor, Phys. Lett. B **632**, 197 (2006).
 - [34] T. A. Trainor, J. Phys. G **39**, 095102 (2012).
 - [35] B. Andersson, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **7**, 1 (1997).
 - [36] K. Aamodt *et al.* (ALICE Collaboration), Phys. Rev. Lett. **106**, 032301 (2011).
 - [37] T. A. Trainor, arXiv:1403.6494.
 - [38] T. A. Trainor, arXiv:1403.3685.
 - [39] C. Adler *et al.* (STAR Collaboration), Phys. Rev. Lett. **87**, 112303 (2001).
 - [40] S. S. Adler *et al.* (PHENIX Collaboration), Phys. Rev. C **71**, 034908 (2005), [Erratum-ibid. C **71**, 049901 (2005)].
 - [41] T. A. Trainor and D. T. Kettler, Phys. Rev. C **84**, 024910 (2011).