



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

ϕ meson production in the forward/backward rapidity region in Cu + Au collisions at $\sqrt{s_{NN}}=200$ GeV

A. Adare *et al.* (PHENIX Collaboration)

Phys. Rev. C **93**, 024904 — Published 4 February 2016

DOI: [10.1103/PhysRevC.93.024904](https://doi.org/10.1103/PhysRevC.93.024904)

1 ϕ meson production in the forward/backward rapidity region in Cu+Au collisions at
2 $\sqrt{s_{NN}} = 200$ GeV

3 A. Adare,¹³ C. Aidala,^{39,44} N.N. Ajitanand,⁶⁴ Y. Akiba,^{58,59} R. Akimoto,¹² J. Alexander,⁶⁴ M. Alfred,²³
4 H. Al-Ta'ani,⁵² K.R. Andrews,¹ A. Angerami,¹⁴ K. Aoki,^{32,58} N. Apadula,^{28,65} E. Appelt,⁶⁹ Y. Aramaki,^{12,58}
5 R. Armendariz,⁸ H. Asano,^{35,58} E.C. Aschenauer,⁷ E.T. Atomssa,⁶⁵ T.C. Awes,⁵⁴ B. Azmoun,⁷ V. Babintsev,²⁴
6 M. Bai,⁶ X. Bai,¹¹ N.S. Bandara,⁴³ B. Bannier,⁶⁵ K.N. Barish,⁸ B. Bassalleck,⁵¹ A.T. Basye,¹ S. Bathe,^{5,59}
7 V. Baublis,⁵⁷ C. Baumann,^{7,45} S. Baumgart,⁵⁸ A. Bazilevsky,⁷ M. Beaumier,⁸ S. Beckman,¹³ R. Belmont,^{13,44,69}
8 J. Ben-Benjamin,⁴⁶ R. Bennett,⁶⁵ A. Berdnikov,⁶¹ Y. Berdnikov,⁶¹ D. Black,⁸ D.S. Blau,³⁴ J.S. Bok,^{52,72}
9 K. Boyle,⁵⁹ M.L. Brooks,³⁹ D. Broxmeyer,⁴⁶ J. Bryslawskij,⁵ H. Buesching,⁷ V. Bumazhnov,²⁴ G. Bunce,^{7,59}
10 S. Butsyk,^{39,51} S. Campbell,^{14,28,65} P. Castera,⁶⁵ C.-H. Chen,^{59,65} C.Y. Chi,¹⁴ M. Chiu,⁷ I.J. Choi,^{25,72}
11 J.B. Choi,¹⁰ S. Choi,⁶³ R.K. Choudhury,⁴ P. Christiansen,⁴¹ T. Chujo,⁶⁸ O. Chvala,⁸ V. Cianciolo,⁵⁴ Z. Citron,^{65,70}
12 B.A. Cole,¹⁴ Z. Conesa del Valle,³⁶ M. Connors,⁶⁵ N. Cronin,^{46,65} N. Crossette,⁴⁶ M. Csanád,¹⁷ T. Csörgő,⁷¹
13 S. Dairaku,^{35,58} T.W. Danley,⁵³ A. Datta,^{43,51} M.S. Daugherty,¹ G. David,⁷ M.K. Dayananda,²⁰ K. DeBlasio,⁵¹
14 K. Dehmelt,⁶⁵ A. Denisov,²⁴ A. Deshpande,^{59,65} E.J. Desmond,⁷ K.V. Dharmawardane,⁵² O. Dietzsch,⁶² L. Ding,²⁸
15 A. Dion,^{28,65} P.B. Diss,⁴² J.H. Do,⁷² M. Donadelli,⁶² L. D'Orazio,⁴² O. Drapier,³⁶ A. Drees,⁶⁵ K.A. Drees,⁶
16 J.M. Durham,^{39,65} A. Durum,²⁴ Y.V. Efremenko,⁵⁴ T. Engelmores,¹⁴ A. Enokizono,^{54,58,60} H. En'yo,^{58,59}
17 S. Esumi,⁶⁸ K.O. Eyser,⁷ B. Fadem,⁴⁶ N. Feege,⁶⁵ D.E. Fields,⁵¹ M. Finger,⁹ M. Finger, Jr.,⁹ F. Fleuret,³⁶
18 S.L. Fokin,³⁴ J.E. Frantz,⁵³ A. Franz,⁷ A.D. Frawley,¹⁹ Y. Fukao,^{32,58} T. Fusayasu,⁴⁸ K. Gainey,¹ C. Gal,⁶⁵
19 P. Gallus,¹⁵ P. Garg,³ A. Garishvili,⁶⁶ I. Garishvili,^{38,66} H. Ge,⁶⁵ F. Giordano,²⁵ A. Glenn,³⁸ X. Gong,⁶⁴
20 M. Gonin,³⁶ Y. Goto,^{58,59} R. Granier de Cassagnac,³⁶ N. Grau,^{2,14} S.V. Greene,⁶⁹ M. Grosse Perdekamp,²⁵
21 Y. Gu,⁶⁴ T. Gunji,¹² L. Guo,³⁹ H. Guragain,²⁰ H.-Å. Gustafsson,^{41,*} T. Hachiya,⁵⁸ J.S. Haggerty,⁷ K.I. Hahn,¹⁸
22 H. Hamagaki,¹² J. Hamblen,⁶⁶ H.F. Hamilton,¹ R. Han,⁵⁶ S.Y. Han,¹⁸ J. Hanks,^{14,65} C. Harper,⁴⁶ S. Hasegawa,²⁹
23 T.O.S. Haseler,²⁰ K. Hashimoto,^{58,60} E. Haslum,⁴¹ R. Hayano,¹² X. He,²⁰ T.K. Hemmick,⁶⁵ T. Hester,⁸ J.C. Hill,²⁸
24 R.S. Hollis,⁸ W. Holzmann,¹⁴ K. Homma,²² B. Hong,³³ T. Horaguchi,⁶⁸ Y. Hori,¹² D. Hornback,⁵⁴ T. Hoshino,²²
25 N. Hotvedt,²⁸ J. Huang,^{7,39} S. Huang,⁶⁹ T. Ichihara,^{58,59} R. Ichimiya,⁵⁸ H. Iinuma,³² Y. Ikeda,^{58,68} K. Imai,^{29,35,58}
26 Y. Imazu,⁵⁸ M. Inaba,⁶⁸ A. Iordanova,⁸ D. Isenhower,¹ M. Ishihara,⁵⁸ A. Isinhue,⁴⁶ M. Issah,⁶⁹ D. Ivanishchev,⁵⁷
27 Y. Iwanaga,²² B.V. Jacak,⁶⁵ S.J. Jeon,⁴⁷ M. Jezghani,²⁰ J. Jia,^{7,64} X. Jiang,³⁹ D. John,⁶⁶ B.M. Johnson,⁷ T. Jones,¹
28 K.S. Joo,⁴⁷ D. Jouan,⁵⁵ D.S. Jumper,²⁵ J. Kamin,⁶⁵ S. Kanda,^{12,32} S. Kaneti,⁶⁵ B.H. Kang,²¹ J.H. Kang,⁷²
29 J.S. Kang,²¹ J. Kapustinsky,³⁹ K. Karatsu,^{35,58} M. Kasai,^{58,60} D. Kaway,^{43,59} A.V. Kazantsev,³⁴ T. Kempel,²⁸
30 J.A. Key,⁵¹ V. Khachatryan,⁶⁵ P.K. Khandai,³ A. Khanzadeev,⁵⁷ K.M. Kijima,²² B.I. Kim,³³ C. Kim,³³ D.J. Kim,³⁰
31 E.-J. Kim,¹⁰ G.W. Kim,¹⁸ M. Kim,⁶³ Y.-J. Kim,²⁵ Y.K. Kim,²¹ B. Kimelman,⁴⁶ E. Kinney,¹³ Á. Kiss,¹⁷
32 E. Kistenev,⁷ R. Kitamura,¹² J. Klatsky,¹⁹ D. Kleinjan,⁸ P. Kline,⁶⁵ T. Koblesky,¹³ L. Kochenda,⁵⁷ M. Kofarago,¹⁷
33 B. Komkov,⁵⁷ M. Konno,⁶⁸ J. Koster,^{25,59} D. Kotchetkov,⁵³ D. Kotov,^{57,61} A. Král,¹⁵ F. Krizek,³⁰ G.J. Kunde,³⁹
34 K. Kurita,^{58,60} M. Kurosawa,^{58,59} Y. Kwon,⁷² G.S. Kyle,⁵² R. Lacey,⁶⁴ Y.S. Lai,¹⁴ J.G. Lajoie,²⁸ A. Lebedev,²⁸
35 D.M. Lee,³⁹ G.H. Lee,¹⁰ J. Lee,¹⁸ K.B. Lee,^{33,39} K.S. Lee,³³ S. Lee,⁷² S.H. Lee,⁶⁵ S.R. Lee,¹⁰ M.J. Leitch,³⁹
36 M.A.L. Leite,⁶² M. Leitgab,²⁵ B. Lewis,⁶⁵ X. Li,¹¹ S.H. Lim,⁷² L.A. Linden Levy,¹³ H. Liu,³⁹ M.X. Liu,³⁹ B. Love,⁶⁹
37 D. Lynch,⁷ C.F. Maguire,⁶⁹ Y.I. Makdisi,⁶ M. Makek,^{70,73} A. Manion,⁶⁵ V.I. Manko,³⁴ E. Mannel,^{7,14} Y. Mao,^{56,58}
38 T. Maruyama,²⁹ H. Masui,⁶⁸ M. McCumber,^{13,39,65} P.L. McGaughey,³⁹ D. McGlinchey,^{13,19} C. McKinney,²⁵
39 N. Means,⁶⁵ A. Meles,⁵² M. Mendoza,⁸ B. Meredith,²⁵ Y. Miake,⁶⁸ T. Mibe,³² A.C. Mignerey,⁴² K. Miki,^{58,68}
40 A. Milov,⁷⁰ D.K. Mishra,⁴ J.T. Mitchell,⁷ Y. Miyachi,^{58,67} S. Miyasaka,^{58,67} S. Mizuno,^{58,68} A.K. Mohanty,⁴
41 S. Mohapatra,⁶⁴ P. Montuenga,²⁵ H.J. Moon,⁴⁷ T. Moon,⁷² Y. Morino,¹² A. Morreale,⁸ D.P. Morrison,^{7,†}
42 M. Moskowitz,⁴⁶ S. Motschwiller,⁴⁶ T.V. Moukhanova,³⁴ T. Murakami,^{35,58} J. Murata,^{58,60} A. Mwai,⁶⁴ T. Nagae,³⁵
43 S. Nagamiya,^{32,58} K. Nagashima,²² J.L. Nagle,^{13,‡} M. Naglis,⁷⁰ M.I. Nagy,^{17,71} I. Nakagawa,^{58,59} H. Nakagomi,^{58,68}
44 Y. Nakamiya,²² K.R. Nakamura,^{35,58} T. Nakamura,⁵⁸ K. Nakano,^{58,67} C. Nattrass,⁶⁶ P.K. Netrakanti,⁴ J. Newby,³⁸
45 M. Nguyen,⁶⁵ M. Nihashi,^{22,58} T. Niida,⁶⁸ S. Nishimura,¹² R. Nouicer,^{7,59} T. Novák,^{31,71} N. Novitzky,^{30,65}
46 A.S. Nyanin,³⁴ C. Oakley,²⁰ E. O'Brien,⁷ C.A. Ogilvie,²⁸ H. Oide,¹² M. Oka,⁶⁸ K. Okada,⁵⁹ J.D. Orjuela Koop,¹³
47 J.D. Osborn,⁴⁴ A. Oskarsson,⁴¹ M. Ouchida,^{22,58} K. Ozawa,^{12,32} R. Pak,⁷ V. Pantuev,^{26,65} V. Papavassiliou,⁵²
48 B.H. Park,²¹ I.H. Park,¹⁸ J.S. Park,⁶³ S. Park,⁶³ S.K. Park,³³ S.F. Pate,⁵² L. Patel,²⁰ M. Patel,²⁸ H. Pei,²⁸
49 J.-C. Peng,²⁵ H. Pereira,¹⁶ D.V. Perepelitsa,^{7,14} G.D.N. Perera,⁵² D.Yu. Peressounko,³⁴ J. Perry,²⁸ R. Petti,^{7,65}
50 C. Pinkenburg,⁷ R. Pinson,¹ R.P. Pisani,⁷ M. Proissl,⁶⁵ M.L. Purschke,⁷ H. Qu,^{1,20} J. Rak,³⁰ B.J. Ramson,⁴⁴
51 I. Ravinovich,⁷⁰ K.F. Read,^{54,66} K. Reygers,⁴⁵ D. Reynolds,⁶⁴ V. Riabov,^{50,57} Y. Riabov,^{57,61} E. Richardson,⁴²
52 T. Rinn,²⁸ N. Riveli,⁵³ D. Roach,⁶⁹ G. Roche,^{40,*} S.D. Rolnick,⁸ M. Rosati,²⁸ S.S.E. Rosendahl,⁴¹ Z. Rowan,⁵

53 J.G. Rubin,⁴⁴ M.S. Ryu,²¹ B. Sahlmueller,^{45,65} N. Saito,³² T. Sakaguchi,⁷ H. Sako,²⁹ V. Samsonov,^{50,57}
 54 S. Sano,¹² M. Sarsour,²⁰ S. Sato,²⁹ T. Sato,⁶⁸ M. Savastio,⁶⁵ S. Sawada,³² B. Schaefer,⁶⁹ B.K. Schmoll,⁶⁶
 55 K. Sedgwick,⁸ J. Seele,⁵⁹ R. Seidl,^{58,59} Y. Sekiguchi,¹² A. Sen,^{20,66} R. Seto,⁸ P. Sett,⁴ A. Sexton,⁴² D. Sharma,^{65,70}
 56 A. Shaver,²⁸ I. Shein,²⁴ T.-A. Shibata,^{58,67} K. Shigaki,²² H.H. Shim,³³ M. Shimomura,^{28,49,68} K. Shoji,^{35,58}
 57 P. Shukla,⁴ A. Sickles,^{7,25} C.L. Silva,^{28,39} D. Silvermyr,^{41,54} C. Silvestre,¹⁶ K.S. Sim,³³ B.K. Singh,³ C.P. Singh,³
 58 V. Singh,³ M. Skolnik,⁴⁶ M. Slunečka,⁹ M. Snowball,³⁹ T. Sodre,⁴⁶ S. Solano,⁴⁶ R.A. Soltz,³⁸ W.E. Sondheim,³⁹
 59 S.P. Sorensen,⁶⁶ I.V. Sourikova,⁷ P.W. Stankus,⁵⁴ P. Steinberg,⁷ E. Stenlund,⁴¹ M. Stepanov,^{43,52,*} A. Ster,⁷¹
 60 S.P. Stoll,⁷ M.R. Stone,¹³ T. Sugitate,²² A. Sukhanov,⁷ T. Sumita,⁵⁸ J. Sun,⁶⁵ J. Sziklai,⁷¹ E.M. Takagui,⁶²
 61 A. Takahara,¹² A. Taketani,^{58,59} R. Tanabe,⁶⁸ Y. Tanaka,⁴⁸ S. Taneja,⁶⁵ K. Tanida,^{35,58,59,63} M.J. Tannenbaum,⁷
 62 S. Tarafdar,^{3,70} A. Taranenko,^{50,64} E. Tennant,⁵² H. Themann,⁶⁵ D. Thomas,¹ R. Tieulent,²⁰ A. Timilsina,²⁸
 63 T. Todoroki,^{58,68} M. Togawa,⁵⁹ L. Tomášek,²⁷ M. Tomášek,^{15,27} H. Torii,^{12,22} C.L. Towell,¹ R. Towell,¹
 64 R.S. Towell,¹ I. Tserruya,⁷⁰ Y. Tsuchimoto,²² K. Utsunomiya,¹² C. Vale,⁷ H.W. van Hecke,³⁹ M. Vargyas,¹⁷
 65 E. Vazquez-Zambrano,¹⁴ A. Veicht,¹⁴ J. Velkovska,⁶⁹ R. Vértesi,⁷¹ M. Virius,¹⁵ A. Vossen,²⁵ V. Vrba,^{15,27}
 66 E. Vznuzdaev,⁵⁷ X.R. Wang,^{52,59} D. Watanabe,²² K. Watanabe,^{58,60,68} Y. Watanabe,^{58,59} Y.S. Watanabe,^{12,32}
 67 F. Wei,^{28,52} R. Wei,⁶⁴ J. Wessels,⁴⁵ S. Whitaker,²⁸ A.S. White,⁴⁴ S.N. White,⁷ D. Winter,¹⁴ S. Wolin,²⁵
 68 C.L. Woody,⁷ R.M. Wright,¹ M. Wysocki,^{13,54} B. Xia,⁵³ L. Xue,²⁰ S. Yalcin,⁶⁵ Y.L. Yamaguchi,^{12,58,65}
 69 R. Yang,²⁵ A. Yanovich,²⁴ J. Ying,²⁰ S. Yokkaichi,^{58,59} J.H. Yoo,³³ J.S. Yoo,¹⁸ I. Yoon,⁶³ Z. You,^{39,56}
 70 G.R. Young,⁵⁴ I. Younus,^{37,51} H. Yu,⁵⁶ I.E. Yushmanov,³⁴ W.A. Zajc,¹⁴ A. Zelenski,⁶ S. Zhou,¹¹ and L. Zou⁸

(PHENIX Collaboration)

¹Abilene Christian University, Abilene, Texas 79699, USA

²Department of Physics, Augustana University, Sioux Falls, South Dakota 57197, USA

³Department of Physics, Banaras Hindu University, Varanasi 221005, India

⁴Bhabha Atomic Research Centre, Bombay 400 085, India

⁵Baruch College, City University of New York, New York, New York, 10010 USA

⁶Collider-Accelerator Department, Brookhaven National Laboratory, Upton, New York 11973-5000, USA

⁷Physics Department, Brookhaven National Laboratory, Upton, New York 11973-5000, USA

⁸University of California-Riverside, Riverside, California 92521, USA

⁹Charles University, Ovocný trh 5, Praha 1, 116 36, Prague, Czech Republic

¹⁰Chonbuk National University, Jeonju, 561-756, Korea

¹¹Science and Technology on Nuclear Data Laboratory, China Institute of Atomic Energy, Beijing 102413, P. R. China

¹²Center for Nuclear Study, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan

¹³University of Colorado, Boulder, Colorado 80309, USA

¹⁴Columbia University, New York, New York 10027 and Nevis Laboratories, Irvington, New York 10533, USA

¹⁵Czech Technical University, Žitkova 4, 166 36 Prague 6, Czech Republic

¹⁶Dapnia, CEA Saclay, F-91191, Gif-sur-Yvette, France

¹⁷ELTE, Eötvös Loránd University, H-1117 Budapest, Pázmány P. s. 1/A, Hungary

¹⁸Ewha Womans University, Seoul 120-750, Korea

¹⁹Florida State University, Tallahassee, Florida 32306, USA

²⁰Georgia State University, Atlanta, Georgia 30303, USA

²¹Hanyang University, Seoul 133-792, Korea

²²Hiroshima University, Kagamiyama, Higashi-Hiroshima 739-8526, Japan

²³Department of Physics and Astronomy, Howard University, Washington, DC 20059, USA

²⁴IHEP Protvino, State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, 142281, Russia

²⁵University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA

²⁶Institute for Nuclear Research of the Russian Academy of Sciences, prospekt 60-letiya Oktyabrya 7a, Moscow 117312, Russia

²⁷Institute of Physics, Academy of Sciences of the Czech Republic, Na Slovance 2, 182 21 Prague 8, Czech Republic

²⁸Iowa State University, Ames, Iowa 50011, USA

²⁹Advanced Science Research Center, Japan Atomic Energy Agency, 2-4

Shirakata Shirane, Tokai-mura, Naka-gun, Ibaraki-ken 319-1195, Japan

³⁰Helsinki Institute of Physics and University of Jyväskylä, P.O.Box 35, FI-40014 Jyväskylä, Finland

³¹Károly Róberts University College, H-3200 Gyöngös, Mátraiút 36, Hungary

³²KEK, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan

³³Korea University, Seoul, 136-701, Korea

³⁴National Research Center "Kurchatov Institute", Moscow, 123098 Russia

³⁵Kyoto University, Kyoto 606-8502, Japan

³⁶Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS-IN2P3, Route de Saclay, F-91128, Palaiseau, France

³⁷Physics Department, Lahore University of Management Sciences, Lahore 54792, Pakistan

³⁸Lawrence Livermore National Laboratory, Livermore, California 94550, USA

³⁹Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

⁴⁰LPC, Université Blaise Pascal, CNRS-IN2P3, Clermont-Fd, 63177 Aubiere Cedex, France

- 113 ⁴¹Department of Physics, Lund University, Box 118, SE-221 00 Lund, Sweden
 114 ⁴²University of Maryland, College Park, Maryland 20742, USA
 115 ⁴³Department of Physics, University of Massachusetts, Amherst, Massachusetts 01003-9337, USA
 116 ⁴⁴Department of Physics, University of Michigan, Ann Arbor, Michigan 48109-1040, USA
 117 ⁴⁵Institut für Kernphysik, University of Muenster, D-48149 Muenster, Germany
 118 ⁴⁶Muhlenberg College, Allentown, Pennsylvania 18104-5586, USA
 119 ⁴⁷Myongji University, Yongin, Kyonggido 449-728, Korea
 120 ⁴⁸Nagasaki Institute of Applied Science, Nagasaki-shi, Nagasaki 851-0193, Japan
 121 ⁴⁹Nara Women's University, Kita-uoya Nishi-machi Nara 630-8506, Japan
 122 ⁵⁰National Research Nuclear University, MEPhI, Moscow Engineering Physics Institute, Moscow, 115409, Russia
 123 ⁵¹University of New Mexico, Albuquerque, New Mexico 87131, USA
 124 ⁵²New Mexico State University, Las Cruces, New Mexico 88003, USA
 125 ⁵³Department of Physics and Astronomy, Ohio University, Athens, Ohio 45701, USA
 126 ⁵⁴Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA
 127 ⁵⁵IPN-Orsay, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, BP1, F-91406, Orsay, France
 128 ⁵⁶Peking University, Beijing 100871, P. R. China
 129 ⁵⁷PNPI, Petersburg Nuclear Physics Institute, Gatchina, Leningrad region, 188300, Russia
 130 ⁵⁸RIKEN Nishina Center for Accelerator-Based Science, Wako, Saitama 351-0198, Japan
 131 ⁵⁹RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, New York 11973-5000, USA
 132 ⁶⁰Physics Department, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima, Tokyo 171-8501, Japan
 133 ⁶¹Saint Petersburg State Polytechnic University, St. Petersburg, 195251 Russia
 134 ⁶²Universidade de São Paulo, Instituto de Física, Caixa Postal 66318, São Paulo CEP05315-970, Brazil
 135 ⁶³Department of Physics and Astronomy, Seoul National University, Seoul 151-742, Korea
 136 ⁶⁴Chemistry Department, Stony Brook University, SUNY, Stony Brook, New York 11794-3400, USA
 137 ⁶⁵Department of Physics and Astronomy, Stony Brook University, SUNY, Stony Brook, New York 11794-3800, USA
 138 ⁶⁶University of Tennessee, Knoxville, Tennessee 37996, USA
 139 ⁶⁷Department of Physics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan
 140 ⁶⁸Center for Integrated Research in Fundamental Science and Engineering, University of Tsukuba, Tsukuba, Ibaraki 305, Japan
 141 ⁶⁹Vanderbilt University, Nashville, Tennessee 37235, USA
 142 ⁷⁰Weizmann Institute, Rehovot 76100, Israel
 143 ⁷¹Institute for Particle and Nuclear Physics, Wigner Research Centre for Physics, Hungarian
 144 Academy of Sciences (Wigner RCP, RMKI) H-1525 Budapest 114, POBox 49, Budapest, Hungary
 145 ⁷²Yonsei University, IPAP, Seoul 120-749, Korea
 146 ⁷³University of Zagreb, Faculty of Science, Department of Physics, Bijenička 32, HR-10002 Zagreb, Croatia
 147 (Dated: December 14, 2015)

148 The PHENIX experiment at the Relativistic Heavy Ion Collider has measured ϕ meson production
 149 and its nuclear modification in asymmetric Cu+Au heavy-ion collisions at $\sqrt{s_{NN}} = 200$ GeV at both
 150 forward Cu-going direction ($1.2 < y < 2.2$) and backward Au-going direction ($-2.2 < y < -1.2$),
 151 rapidities. The measurements are performed via the dimuon decay channel and reported as a
 152 function of the number of participating nucleons, rapidity, and transverse momentum. In the most
 153 central events, 0%–20% centrality, the ϕ meson yield integrated over $1 < p_T < 5$ GeV/ c prefers a
 154 smaller value, which means a larger nuclear modification, in the Cu-going direction compared to
 155 the Au-going direction. Additionally, the nuclear-modification factor in Cu+Au collisions averaged
 156 over all centrality is measured to be similar to the previous PHENIX result in d +Au collisions for
 157 these rapidities.

158 PACS numbers: 25.75.Dw

* Deceased

† PHENIX Co-Spokesperson: morrison@bnl.gov

‡ PHENIX Co-Spokesperson: jamie.nagle@colorado.edu

I. INTRODUCTION

159

160 The Relativistic Heavy Ion Collider (RHIC) accelerator and its four experiments have previously provided extensive
 161 experimental evidence to confirm the formation of a deconfined state of nuclear matter, referred to as the quark-gluon
 162 plasma (QGP), in the initial stages of high-energy heavy-ion collisions [1–4]. Currently, a major objective in the field
 163 of high-energy nuclear physics is to characterize the properties of the QGP in a quantitative way. The ϕ meson is
 164 a useful probe for studying the QGP properties, because it is sensitive to several aspects of the collision, including
 165 modifications of strangeness production in bulk matter [5–7]. Due to its small inelastic cross section for interaction
 166 with nonstrange hadrons [6, 8], the ϕ meson is less affected by late hadronic rescattering and may reflect the initial
 167 evolution of the system. Being composed of a nearly pure strange anti-strange ($s\bar{s}$) state, the ϕ meson puts additional
 168 constraints on models of quark recombination in the QGP.

169 The study of the QGP typically involves comparisons of different observables measured in nucleus-nucleus (A+B)
 170 collisions and in proton-proton ($p+p$) collisions at the same center-of-mass energy. Modifications in the A+B collisions
 171 with respect to $p+p$ collisions could be interpreted as being due to the hot nuclear matter (HNM) – possibly QGP –
 172 being produced. However, nuclear modifications could be present in the initial state of the collisions even if no QGP is
 173 produced. These effects, typically referred to as cold nuclear matter (CNM), may include the modification of parton
 174 distribution functions (PDFs) in a nucleus [9], initial-state energy loss [10], and the Cronin effect, which is often
 175 attributed to multiple scattering of the incoming parton inside the target nucleus [11, 12]. CNM effects can be probed
 176 with d +Au collisions. PHENIX has previously measured ϕ meson production in d +Au collisions at forward, mid- and
 177 backward rapidities [13]. Suppression was observed in the forward (d -going) direction, where small- x partons from
 178 the Au nucleus are probed, and an enhancement was seen in the backward (Au-going) direction. Similar behavior
 179 was previously observed for inclusive charged hadrons and open heavy flavor in d +Au collisions [14, 15], potentially
 180 indicating similar particle production and modification mechanisms.

181 The rapidity dependence y of particle production in asymmetric collisions with a smaller-A projectile and a large-A
 182 target, provides a way to investigate both hot and cold nuclear-matter effects. Previous J/ψ meson data in Cu+Au
 183 collisions [16] showed that the ratio of forward ($1.2 < y < 2.2$, or Cu-going) to backward ($-2.2 < y < -1.2$, or
 184 Au-going) J/ψ modification was comparable in both sign and magnitude to that expected from CNM effects. The ϕ
 185 meson is composed of lighter closed flavor ($s\bar{s}$) and its production from 1.0 GeV/ c to 5.0 GeV/ c involves a mix of soft
 186 and hard processes and would provide a link between heavy flavor and lighter mesons. Comparison of the ϕ meson
 187 production in Cu+Au and d +Au systems and to J/ψ production in Cu+Au collisions may shed light on the mixture
 188 of HNM and CNM effects on ϕ -meson production.

189 The production of ϕ mesons has already been measured at PHENIX in $p+p$, d +Au, Cu+Cu, and Au+Au at
 190 midrapidity [17–19] and in $p+p$ and d +Au at forward and backward rapidities [13, 20] over a wide range in p_T .
 191 Previous measurements from Au+Au and Cu+Cu collisions [18] in a similar momentum range were found to be
 192 consistent with HNM effects and exhibited large flow anisotropies. The STAR Collaboration has also previously
 193 measured ϕ meson production at midrapidity in Cu+Cu and Au+Au collisions [21, 22]. ϕ meson production has also
 194 been measured by the ALICE Collaboration at large rapidity in $p+p$ and p +Pb collisions [23] and at midrapidity in
 195 Pb+Pb collisions [24].

196 In this paper, the production of ϕ mesons is determined at forward and backward rapidities via dimuons recon-
 197 structed in the PHENIX muon spectrometers in Cu+Au collisions at $\sqrt{s_{NN}} = 200$ GeV recorded in 2012. The particle
 198 multiplicity at these rapidities in heavy-ion collisions results in large combinatorial backgrounds and produces a
 199 challenging environment for ϕ meson measurements. Previous measurements were thus limited to smaller collision
 200 species. A procedure for removing the background is detailed and a measurement of the ϕ meson nuclear modification
 201 factor R_{CuAu} in Cu+Au collisions at forward and backward rapidities is presented versus y , p_T , and the number of
 202 participating nucleons.

203

II. EXPERIMENTAL SETUP

204 The PHENIX detector is described in detail in [25], and a schematic of the 2012 setup is shown in Fig. 1. This
 205 analysis uses the dimuon decay channel of the ϕ meson. The detectors relevant for this measurement are forward and
 206 backward muon spectrometers [26], the two beam-beam counters (BBCs) [27], the silicon vertex tracker (VTX) [28],
 207 and the forward silicon vertex detector (FVTX) [29].

208 This study used minimum bias (MB) events triggered by the BBCs. The BBCs comprise two arrays of 64 Čerenkov
 209 counters covering the pseudorapidity range $3.1 < |\eta| < 3.9$. The MB trigger required two or more counters firing on
 210 each side and a z -vertex selection around the nominal center of the detector acceptance [16]. The MB trigger fired on
 211 $93 \pm 3\%$ of the 5.2 ± 0.2 b total inelastic Cu+Au cross section. In this case, the z -vertex was measured by the BBCs
 212 with a resolution of $\sigma_z \approx 0.5$ – 2.0 cm, depending on the event multiplicity.




FIG. 1. (color online) The 2012 setup of the PHENIX detector.

213 The collision point is determined in x , y and z by the two vertex detectors, VTX and FVTX, with a resolution of
 214 better than 100 microns. The VTX and FVTX detectors were installed in 2011 and 2012 to provide precise particle
 215 vertexing and tracking in the central and forward/backward rapidities. Covering approximately the same rapidity
 216 range as the existing muon spectrometers, the FVTX is composed of two endcaps, each with four stations that are
 217 perpendicular to the beamline and composed of silicon mini-strip sensors that have a 75 micron pitch in the radial
 218 direction and lengths in the ϕ direction varying from 3.4 mm to 11.5 mm. The VTX, which surrounds the collision
 219 region at PHENIX, comprises four layers of silicon sensors. The inner two layers and outer two layers are composed
 220 of 30 pixel ladders and 44 stripixel ladders, respectively.

221 The muon system is separated into the north and south muon arms. Each arm comprises four subcomponents: an
 222 absorber material, a magnet, a muon tracker (MuTr), and a muon identifier (MuID). Initially, the absorbers were
 223 composed of 19 cm copper and 60 cm iron, but an additional 36.2 cm of stainless steel was added in 2010 to help
 224 decrease the hadronic background. Following the absorber in each muon arm is the MuTr, which comprises three
 225 sets of cathode strip chambers in a radial magnetic field with an integrated bending power of 0.8 T·m. The final
 226 component is the MuID, which comprises five alternating steel absorbers and Iarocci tubes to further reduce the
 227 number of punch-through hadrons that can be mistakenly identified as muons. The backplates of the magnets provide
 228 the first absorber layer for the muon identifier systems. The backplate of the south muon magnet is 10 cm shorter
 229 than the backplate of the north muon magnet, resulting in less total absorber material in the south arm than the
 230 north arm, and thus a slightly different momentum acceptance. The muon spectrometers cover the pseudorapidity
 231 range $1.2 < |\eta| < 2.2$ over the full azimuth. Muon candidates are identified by reconstructed tracks in the MuTr
 232 matched to MuID tracks, where at least one of the tracks from a pair of muon candidates in the same event penetrates
 233 through to the last MuID plane. The minimum momentum needed for a muon to reach the last MuID plane is ~ 3
 234 GeV/ c .

235 III. DATA ANALYSIS

236 A. Dataset and quality cuts

237 In this analysis, ϕ meson candidates are selected from two reconstructed muons in the RHIC Cu+Au dataset from
 238 2012. The ϕ meson invariant yields are then measured and used to calculate the nuclear modification factor R_{CuAu} ,
 239 which is compared to results from other systems. For this analysis, 4.73 billion ($\mathcal{L} = 0.97 \text{ nb}^{-1}$) sampled MB events
 240 were used within ± 10 cm z -vertex and 0%–93% centrality. The total inelastic cross section for Cu+Au collisions at
 241 200 GeV was estimated by a Glauber simulation to be 5.2 ± 0.2 b.

242 A set of quality assurance cuts is applied to the data to select good muon candidates and improve the signal-to-
 243 background ratio. These cuts are summarized in Table I. The collision z -vertex is required to be within ± 10 cm of the
 244 center of the interaction region along the beam direction, as measured with the BBCs. The MuTr tracks are required
 245 to match the MuID tracks at the first MuID layer in both position and angle. In addition, only dimuon candidates in

TABLE I. Quality cuts for ϕ meson signal extraction in Cu+Au collisions.

Variable	Au-going	Cu-going	Meaning
$ z_{\text{vtx}} (\text{cm})$	< 10	< 10	Collision vertex along the beam direction as measured by the BBCs
$pDG0$ (GeV/c·cm)	< 90	< 50	Track momentum times the spatial difference between the MuTr track and MuID track at the first MuID layer
$pDDG0$ (GeV/c·radian)	< 30	< 45	Track momentum times the slope difference between the MuTr track and MuID track at the first MuID layer
Track χ^2	< 5	< 10	χ^2/NDF of the μ track
Lastgap	one track ≥ 2 other track ≥ 4	one track ≥ 2 other track ≥ 4	Last MuID plane that the μ track penetrated
nidhits	$>(2 \times \text{lastgap} - 1)$	$>(2 \times \text{lastgap} - 1)$	Number of hits in the MuID, out of the maximum 10
ntrhits	> 11	> 10	Number of hits in the MuTr, out of the maximum 16
χ^2_{vtx}	< 4	< 7	χ^2/NDF of the dimuon track with the vertex
Dimuon p_T (GeV/c)	$1 - 5$	$1 - 5$	Transverse momentum of the dimuon pair
$ p_z (\text{GeV}/c)$	> 2.4	> 2.5	Momentum of the μ along the beam axis

246 which at least one track penetrated to the final MuID layer are selected. Furthermore, the track is required to have
247 greater than a minimum number of possible hits in the MuTr and MuID, and a maximum allowed χ^2 is applied to
248 both the track and vertex determination. There is a minimum allowed single muon momentum along the beam axis,
249 p_z , which is reconstructed and energy-loss corrected at the collision vertex. Finally, this analysis is restricted to the
250 dimuon p_T range of $1 - 5$ GeV/c. This limitation is due to the large backgrounds and small acceptance at low p_T and
251 small statistics at high p_T , preventing signal extraction of the ϕ meson. The events are sorted into centrality classes
252 using the combined charge from both BBCs [16]. The number of binary collisions N_{coll} and number of participating
253 nucleons N_{part} are extracted from a Glauber simulation [16].

254 B. Background subtraction

255 The PHENIX muon spectrometers have a small acceptance for ϕ mesons. Going from the most peripheral centrality
256 bin, 40%–93%, to the most central bin, 0%–20%, the signal-to-background ratio decreases from 0.28 to 0.067 in the
257 Cu-going direction ($1.2 < y < 2.2$) and from 0.37 to 0.090 in the Au-going direction ($-2.2 < y < -1.2$). Due to the
258 very low signal-to-background ratio, particularly in the most central events, the background subtraction is of crucial
259 importance. Accordingly, several different background subtraction methods were explored and compared.

260 The invariant mass distribution is formed by combining muon candidate tracks of opposite charge. This unlike-sign
261 invariant mass spectrum contains the ϕ , ρ and ω mesons as well as both uncorrelated and correlated backgrounds.
262 The uncorrelated backgrounds come from random combinatorial associations of muon candidates, while the correlated
263 backgrounds arise from open charm decay (e.g., $D\bar{D}$ where both decay semileptonically to muons), open beauty decay,
264 η meson and ω meson Dalitz decays and the Drell-Yan process. These correlated backgrounds are described in Sec.
265 III C. The uncorrelated combinatorial background is accounted for via two methods: (1) like-sign dimuons and (2)
266 event mixing.

267 First, the uncorrelated combinatorial background is estimated through the like-sign background subtraction tech-
268 nique, which is generally associated with the assumption that the like-sign dimuon pairs come purely from combina-
269 torial processes without any correlation between muons. It follows that the like-sign distribution can be subtracted
270 from the unlike-sign distribution according to the relationship described in Eq. 1

$$N_{+-} = FG_{+-} - FG_{\pm\pm}, \quad (1)$$

where N_{+-} is the uncorrelated background subtracted signal and FG_{+-} and $FG_{\pm\pm}$ are the unlike-sign and like-sign dimuon pairs, respectively, corresponding to pairs formed within the same event. The like-sign distribution $FG_{\pm\pm}$ is normalized to a quantity that is more precise and not sensitive to differences in the detector acceptance between like-sign and unlike-sign pairs. This background normalization is described in Eq. 2 [30]

$$FG_{\pm\pm} = (FG_{++} + FG_{--}) \frac{2\sqrt{\int FG_{++} dm \int FG_{--} dm}}{\int (FG_{++} + FG_{--}) dm}, \quad (2)$$

where m is the dimuon invariant mass, and the integration is carried out in the range $0.2 < m < 5.0 \text{ GeV}/c^2$.

In parallel to the like-sign technique, the uncorrelated background is also estimated through the event mixing technique. In the standard event mixing method, muons from different events are randomly associated to produce a background distribution of uncorrelated dimuon pairs. Events were mixed with partners from within the same 2%-centrality and 1-cm z -vertex bins in order to minimize the systematic uncertainties. The mixed-event background distributions (BG) were generated with about 8 times higher statistics than the actual background and then normalized to match the same-event foreground (FG). The normalization factor also accounts for slightly different multiplicities from mixing of slightly different events. Although a mass-dependent technique was developed for this analysis, a standard event mixing technique is described in advance. In previous PHENIX analyses, the normalization factor α was calculated as described in Eq. 3

$$\alpha = \sqrt{\frac{\int FG_{++} dm \int FG_{--} dm}{\int BG_{++} dm \int BG_{--} dm}}, \quad (3)$$

where FG_{++} and FG_{--} are the like-sign pairs from the same event and BG_{++} and BG_{--} are the like-sign pairs from mixed events.




FIG. 2. (color online) The event mixing normalization factor α versus mass. This factor shows a dependence on mass, particularly in the low mass region. The error function, which was used to fit α , can also be seen in the plot along with the fit parameters and goodness of fit.

After subtracting and fitting the resonances as well as the remaining correlated background, the yields from mixed-event background subtraction are consistent with the yields from the like-sign technique within statistical uncertainties. The event mixing technique is used in this analysis due to the statistical limitations of the like-sign technique. The differences between the like-sign and event mixing techniques are used to determine one component of the systematic uncertainty on the yield, as described later in Sec. III F.

In this method, each term in the square root of Eq. 3 was integrated over all mass, introducing a mass-independent normalization factor [16, 31]. Dimuons from same events are less likely to be reconstructed in close proximity to each other than those in mixed events, resulting in a larger relative number of mixed-event dimuons at low mass, where the opening angle is small, than at higher mass. Therefore, the normalization factor, which is simply a ratio of the

296 like-sign same-event dimuons to like-sign mixed-event dimuons, drops at lower masses. Because this normalization
 297 factor depends on mass, particularly in the ϕ meson region, it became necessary to introduce a mass-dependent
 298 normalization, as described in Eq. 4, rather than the more commonly used mass-integrated normalization from Eq. 3.

$$\alpha(m) = \sqrt{\frac{FG_{++}(m)FG_{--}(m)}{BG_{++}(m)BG_{--}(m)}} \quad (4)$$

302 This mass-dependent normalization factor is then fit as a function of mass, and the fit function – rather than
 303 the integrated normalization factor – is multiplied to the unlike-sign mixed-event background to get the normalized
 304 background spectrum $BG_{+-}^{\text{normalized}}$,

$$BG_{+-}^{\text{normalized}}(m) = \alpha(m) \times BG_{+-}(m). \quad (5)$$

305 Several fitting functions were tested, including a polynomial and an error function. The error function, which is
 306 used in the final analysis, is described in Eq. 6, where $g(m)$ is the error function and p_0 , p_1 and p_2 are free parameters
 307 of the fit. A plot of the normalization factor as a function of mass fit with an error function is shown in Fig. 2.

$$g(m) = p_0 \times \text{Erf}\left(\frac{m - p_1}{p_2}\right) \quad (6)$$

308 The application of event mixing to describe and subtract backgrounds in the ϕ meson mass region is shown in Fig. 3,
 309 where the open squares represent the mixed-event background and the closed circles are the unlike-sign spectrum.
 310 Before background subtraction, the $\rho + \omega$, ϕ and J/ψ peaks are clearly seen.




FIG. 3. (color online) The unlike-sign spectra and combinatorial background described with event mixing for $1.2 < y < 2.2$ (Cu-going direction) and $-2.2 < y < -1.2$ (Au-going direction). The $\rho + \omega$, ϕ and J/ψ peaks are clearly visible before background subtraction. The mass bin width is 71 MeV as marked on the vertical axis.

308 C. Signal extraction and correlated background

309 After the mixed-event background subtraction, there is still some correlated background remaining. In previous
 310 PHENIX analyses, it was shown that heavy flavor (charm and beauty) contributions were negligible in the ϕ meson
 311 mass region for $p+p$ and $d+Au$ collisions at 200 GeV [13, 20]. Simulation studies showed that η meson Dalitz decays
 312 are one possible contributor to the correlated background. The correlated background is well described by the function
 313 in Eq. 7

$$f(m) = \exp(a \cdot m) + b + c \cdot m, \quad (7)$$

314 where a , b and c are free parameters of the fit $f(m)$. Accordingly, the correlated background in real data are also fit
 315 with the function described in Eq. 7, as shown in Fig. 4, where the mass distribution after mixed-event background
 316 subtraction is shown. Several other fit functions and fit ranges were tested and used to estimate a systematic
 317 uncertainty.

318 The ϕ and ω meson signals are each described by a Gaussian and the signal from the ρ meson by a Breit-Wigner
 319 distribution, as shown in Fig. 4, along with the correlated background description. The ϕ meson mass resolution is
 320 ~ 90 MeV/ c^2 . The PHENIX muon arms are not able to resolve the ρ and ω peaks separately, so a combined fit is
 321 made. All fit parameters are constrained but allowed to vary, except the ratio of the yield of ρ mesons to that of
 322 $\rho + \omega$, which is set as a constant based on the expected ratio between their cross sections and branching ratios. The
 323 data are binned as a function of p_T , y and centrality over the range $1 < p_T < 5$ GeV/ c , $1.2 < |y| < 2.2$, and 0%–93%
 324 centrality.




FIG. 4. (color online) The dimuon mass spectra for $1.2 < y < 2.2$ (Cu-going direction) and $-2.2 < y < -1.2$ (Au-going direction) after subtracting mixed events and fitting the ϕ and $\rho + \omega$ peaks and the remaining correlated background. The mass bin width is 71 MeV as marked on the vertical axis.

D. Detector acceptance and reconstruction efficiency

325
 326 The product of detector acceptance and reconstruction efficiency, $A\varepsilon_{\text{rec}}$, of dimuon decays of ϕ mesons is determined
 327 by the full event reconstruction of the ϕ meson signal obtained from PYTHIA 6.42 [32], run through a full GEANT3 [33]
 328 simulation of the 2012 PHENIX detector setup, and embedded in the MB real-data background. The embedded
 329 simulated events are then reconstructed in the same manner as data with the same cuts applied as in the real data
 330 analysis. The background subtraction and signal extraction are also handled in the exact same manner as in real
 331 data. The $A\varepsilon_{\text{rec}}$ is then calculated as the number of reconstructed ϕ meson candidates divided by the number of ϕ
 332 mesons generated in PYTHIA, both within an appropriate kinematic bin. As previously mentioned, the south arm has
 333 a smaller amount of absorber material, causing a larger acceptance in the south arm (Au-going direction) than in the
 334 north arm (Cu-going direction). In addition, the $A\varepsilon_{\text{rec}}$ has a centrality and p_T dependence. Specifically, for the lower
 335 p_T bin (1-2.5 GeV/ c), $A\varepsilon_{\text{rec}} = 1.21 \times 10^{-3}$ in the Cu-going direction and 1.86×10^{-3} in the Au-going direction, while
 336 for the higher p_T bin (2.5-5 GeV/ c), $A\varepsilon_{\text{rec}} = 1.69 \times 10^{-2}$ in the Cu-going direction and 1.81×10^{-2} in the Au-going
 337 direction. The centrality dependence is not as strong, with the values going from $A\varepsilon_{\text{rec}} = 2.23 \times 10^{-3}$ in the Cu-going
 338 direction and 2.37×10^{-3} in the Au-going direction at 0%–20% centrality to $A\varepsilon_{\text{rec}} = 2.41 \times 10^{-3}$ in the Cu-going
 339 direction and 3.83×10^{-3} in the Au-going direction at 40%–93% centrality.

TABLE II. Systematic uncertainties included in the invariant yield calculations.

Type	Origin	Value
A	Signal extraction	2–31%
B	MuID efficiency	2%
B	MuTr efficiency	2%
B	$A\varepsilon_{\text{rec}}$	13%
B	ϕ candidate selection	3%
B	Like-sign background subtraction	5%
C	MB trigger	3%

E. Invariant yields and nuclear modification factors

The invariant yield is calculated according to the relation:

$$BR \frac{d^2N}{dydp_T} = \frac{1}{\Delta y \Delta p_T} \frac{N}{A\varepsilon_{\text{rec}} N_{\text{evt}}}, \quad (8)$$

where BR is the branching ratio to dimuons ($BR(\phi \rightarrow \mu^+ \mu^-) = (2.89 \pm 0.19) \times 10^{-4}$ [34]), N_{evt} is the number of sampled MB events within the relevant centrality selection ($N_{\text{evt}} = 4.73 \times 10^9$ for the 0%–93% selection), N is the number of observed ϕ mesons, and Δy and Δp_T are the bin widths in y and p_T , respectively. To evaluate the nuclear matter effects on ϕ meson production in Cu+Au collisions, the ϕ meson yields in Cu+Au collisions are compared to those measured in $p+p$ collisions at the same energy after scaling by the number of nucleon-nucleon collisions in the Cu+Au system, N_{coll} . This ratio is called the nuclear modification factor R_{CuAu} , and is defined as:

$$R_{\text{CuAu}} = \frac{\frac{d^2 N_{\text{CuAu}}}{dydp_T}}{N_{\text{coll}} \times \frac{d^2 N_{pp}}{dydp_T}}. \quad (9)$$

The $p+p$ reference data used in the R_{CuAu} are from Ref. [20]. Because the rapidity and p_T binning in the Cu+Au analysis differs from that in the $p+p$ analysis, the $p+p$ invariant yields were re-measured using the same binning as the Cu+Au yields and in a manner similar to Ref. [20]. The sampled luminosity of the $p+p$ data used in this analysis corresponds to $\mathcal{L} = 14.1 \text{ pb}^{-1}$ [20].

F. Systematic uncertainties

The systematic uncertainties associated with this measurement are categorized as Type-A, Type-B or Type-C. Type-A refers to point-to-point uncorrelated uncertainties that allow the data points to move independently with respect to one another. They are added in quadrature with the statistical uncertainties and represented on the plots as an error bar. Type-B uncertainties are correlated point-to-point, which means the points move coherently. All sources of Type-B uncertainty are added in quadrature and displayed as boxes around the data points. Finally, Type-C refers to the global uncertainties which allow the data points to move together by an identical multiplicative factor. The Type-C uncertainties are given in the legends of the plots.

Several systematic uncertainties are evaluated for this analysis. For the signal extraction uncertainty, different fits and parameters are tested for the background normalization factor, the correlated background, the $\rho + \omega$ signal, and the ϕ meson signal. This is done separately for each kinematic bin, and a 2-31% systematic uncertainty is assigned, with the largest uncertainty on yields extracted from the most central events. This is because the high multiplicity in central collisions results in large combinatorial backgrounds and a very small signal-to-background ratio. It is important to note here that the signal extraction uncertainty was primarily dominated by the fluctuations in the correlated background. The $p+p$ reference uncertainty comes from the uncertainty on the ϕ yields in the $p+p$ reference [20]. There is a 4% systematic uncertainty from the MuID efficiency and a 2% uncertainty from the

TABLE III. Systematic uncertainties included in the nuclear modification factor calculations.

Type	Origin	Value
A	Signal extraction	2–31%
A	$p+p$ reference (integrated centrality only)	5–13%
B	MuID efficiency	4%
B	MuTr efficiency	2%
B	$A\varepsilon_{\text{rec}}$	13%
B	ϕ candidate selection	3%
B	Like-sign background subtraction	5%
B	N_{coll} (centrality bins only)	5–10%
C	MB trigger	10%
C	N_{coll} (integrated centrality only)	5%
C	$p+p$ reference (centrality bins only)	11%

TABLE IV. Invariant yield as a function of centrality for $1 < p_T < 5$ GeV/ c and $1.2 < |y| < 2.2$. The first value represents the statistical and Type-A systematic uncertainties, while the second is the systematic uncertainty of Type-B. An additional $\pm 3\%$ Type-C global systematic uncertainty also applies to the yields. The last column summarizes the forward/backward ratio shown in Fig. 11. The forward/backward ratio has no Type-C systematic uncertainty.

Centrality Bin	$\langle N_{\text{part}} \rangle$	$BR \frac{dN}{dy}$ (Cu-going)	$BR \frac{dN}{dy}$ (Au-going)	forward/backward ratio
0%–20%	154.8 ± 4.1	$(7.3 \pm 7.5 \pm 1.1) \times 10^{-5}$	$(3.4 \pm 1.0 \pm 0.5) \times 10^{-4}$	$0.2^{+0.3}_{-0.2} \pm < 0.1$
20%–40%	80.4 ± 3.3	$(1.2 \pm 0.3 \pm 0.2) \times 10^{-4}$	$(1.2 \pm 0.3 \pm 0.2) \times 10^{-4}$	$1.0^{+0.4}_{-0.3} \pm 0.1$
40%–93%	19.5 ± 0.5	$(1.5 \pm 0.6 \pm 0.2) \times 10^{-5}$	$(2.7 \pm 0.7 \pm 0.4) \times 10^{-5}$	$0.6^{+0.4}_{-0.3} \pm 0.1$

368 MuTr efficiency in $p+p$ collisions [20]. In Cu+Au collisions, the MuTr efficiency uncertainty remains the same, while
369 the MuID efficiency uncertainty drops down to 2% [16]. For the $A\varepsilon_{\text{rec}}$ uncertainty, the p_T and y distributions in
370 PYTHIA are changed to match the slope of the distributions in real data, and allowed to vary over the range of the
371 error bars in data, yielding a 13% systematic uncertainty. Real data and simulation inconsistencies in each of the
372 muon identification cuts listed in Table I are also evaluated. They can affect the yields by 3%, which is assigned
373 as a systematic uncertainty on the ϕ meson candidate selection. The like-sign background subtraction uncertainty
374 of 5% comes from differences in the yields when using the like-sign method or the event mixing method. The N_{coll}
375 uncertainty of 5–10% arises from the fact that N_{coll} carries a statistical uncertainty itself. Finally, the MB trigger
376 efficiency uncertainty was 10% in the $p+p$ reference [20] and 3% in Cu+Au collisions [16]. All of these systematic
377 uncertainties are tabulated in Tables II and III.

378 IV. RESULTS

379 The invariant yields for $1 < p_T < 5$ GeV/ c ϕ mesons are calculated as a function of centrality, y and p_T as described
380 in Eq. 8. The results are summarized in Tables IV – VI. Similarly, the nuclear modification factors are formed from

TABLE V. Invariant yield as a function of p_T for 0%–93% centrality and $1.2 < |y| < 2.2$. The first error represents the statistical and Type-A systematic uncertainties, while the second is the systematic uncertainty of Type-B. An additional $\pm 5.8\%$ Type-C global systematic uncertainty also applies.

p_T^{min} (GeV/ c)	p_T^{max} (GeV/ c)	$BR \frac{d^2N}{dy dp_T}$ (Cu-going) (GeV/ c) $^{-1}$	$BR \frac{d^2N}{dy dp_T}$ (Au-going) (GeV/ c) $^{-1}$
1.0	2.5	$(2.7 \pm 0.8 \pm 0.4) \times 10^{-5}$	$(5.5 \pm 0.9 \pm 0.8) \times 10^{-5}$
2.5	5.0	$(1.8 \pm 1.0 \pm 0.3) \times 10^{-7}$	$(4.1 \pm 1.0 \pm 0.6) \times 10^{-7}$

TABLE VI. Invariant yield as a function of rapidity for 0%–93% centrality and $1 < p_T < 5$ GeV/ c . The first error represents the statistical and Type-A systematic uncertainties, while the second is the systematic uncertainty of Type-B. An additional $\pm 5.8\%$ Type-C global systematic uncertainty also applies.

$ y ^{\min}$	$ y ^{\max}$	$BR \frac{dN}{dy}$ (Cu-going)	$BR \frac{dN}{dy}$ (Au-going)
1.8	2.2	$(6.4 \pm 3.1 \pm 0.9) \times 10^{-5}$	$(1.1 \pm 0.2 \pm 0.2) \times 10^{-4}$
1.2	1.8	$(5.3 \pm 2.3 \pm 0.8) \times 10^{-5}$	$(1.1 \pm 0.3 \pm 0.2) \times 10^{-4}$

TABLE VII. Nuclear modification factors as a function of centrality for $1 < p_T < 5$ GeV/ c and $1.2 < |y| < 2.2$. The first error represents the statistical and Type-A systematic uncertainties, while the second is the systematic uncertainty of Type-B. An additional $\pm 15\%$ Type-C global systematic uncertainty also applies.

Centrality Bin	$\langle N_{\text{coll}} \rangle$	R_{CuAu} (Cu-going)	R_{CuAu} (Au-going)
0%–20%	313.8 ± 28.4	$0.4 \pm 0.4 \pm 0.1$	$1.7 \pm 0.5 \pm 0.3$
20%–40%	129.3 ± 12.4	$1.4 \pm 0.4 \pm 0.3$	$1.4 \pm 0.3 \pm 0.3$
40%–93%	21.6 ± 1.0	$1.1 \pm 0.5 \pm 0.2$	$1.9 \pm 0.5 \pm 0.3$

381 the invariant yields using Eq. 9 and tabulated in Tables VII – IX.

382 Fig. 5 shows the invariant yield as a function of the number of participating nucleons N_{part} . In Fig. 6, the
 383 dependence of the invariant yield on transverse momentum p_T is shown. The invariant yield as a function of rapidity
 384 is plotted in Fig. 7. More ϕ mesons are produced in the Au-going direction ($-2.2 < y < -1.2$) than in the Cu-going
 385 direction ($1.2 < y < 2.2$). This may be explained by the larger multiplicity in the Au-going direction coupled with a
 386 mixture of both HNM and CNM effects.

387 Although the invariant yields are interesting on their own, the nuclear modification factor is studied in order to
 388 evaluate the effects of hot and cold nuclear matter on ϕ meson production in Cu+Au collisions at $\sqrt{s_{NN}} = 200$ GeV.

389 The nuclear modification factor as a function of N_{part} is shown in Fig. 8. There is a dependence of R_{CuAu} on
 390 both centrality and rapidity. In the Au-going direction, the R_{CuAu} is greater than unity for all centralities. The
 391 rapidity dependence is similar to the trend observed by PHENIX for $\phi \rightarrow \mu^+ \mu^-$ in d +Au collisions [13] as well as
 392 measurements made by the ALICE Collaboration at large rapidity in p +Pb collisions at 5.02 TeV at the Large Hadron
 393 Collider [23], where an enhancement was observed in the Pb-going direction while the p -going direction was either
 394 suppressed or consistent with unity depending on the p_T range.

395 To further understand the relative roles of different nuclear matter effects in this collision system, the transverse
 396 momentum dependence of the nuclear modification factor is shown in Fig. 9. The data points are placed at the mean
 397 p_T of the bin. Here, the nuclear modification is calculated over integrated centrality, but it should be noted that
 398 the data are dominated by central collisions. There is an enhancement at low p_T in the Au-going direction. In the
 399 Cu-going direction, R_{CuAu} is consistent with unity. The enhancement in the Au-going direction is similar in scale to
 400 that observed in the Au-going direction in d +Au collisions [13], indicating similar nuclear modification between the
 401 two collision systems.

402 Fig. 10 shows the nuclear modification factor R_{CuAu} as a function of y for two rapidity regions, $1.2 < |y| < 1.8$ and
 403 $1.8 < |y| < 2.2$. The data points are placed at the mean y of the bin. As in Fig. 9, the nuclear modification factor
 404 is inclusive of centrality. The rapidity-dependence of R_{CuAu} is similar to the trend observed in previous ϕ meson
 405 measurements in $p(d)$ +Au collisions. In particular, ϕ meson production is enhanced in the Au-going direction. None
 406 of the Cu-going points show significant suppression given the statistical uncertainties. For comparison, the PHENIX
 407 J/ψ meson results in the same Cu+Au dataset from Ref. [16] are also shown in Fig. 10. While the closed charm shows
 408 suppression at both forward and backward rapidity for $1.2 < |y| < 2.2$, the closed strangeness is enhanced at backward

TABLE VIII. Nuclear modification factors as a function of p_T for 0%–93% centrality and $1.2 < |y| < 2.2$. The first error represents the statistical and Type-A systematic uncertainties, while the second is the systematic uncertainty of Type-B. An additional $\pm 11\%$ Type-C global systematic uncertainty also applies.

p_T^{\min} (GeV/ c)	p_T^{\max} (GeV/ c)	R_{CuAu} (Cu-going)	R_{CuAu} (Au-going)
1.0	2.5	$1.1 \pm 0.4 \pm 0.2$	$2.3 \pm 0.4 \pm 0.4$
2.5	5.0	$0.6 \pm 0.4 \pm 0.1$	$1.4 \pm 0.3 \pm 0.2$

TABLE IX. Nuclear modification factors as a function of rapidity for 0%–93% centrality and $1 < p_T < 5$ GeV/ c . The first error represents the statistical and Type-A systematic uncertainties, while the second is the systematic uncertainty of Type-B. An additional $\pm 11\%$ Type-C global systematic uncertainty also applies.

$ y ^{\min}$	$ y ^{\max}$	R_{CuAu} (Cu-going)	R_{CuAu} (Au-going)
1.8	2.2	$1.2 \pm 0.6 \pm 0.2$	$2.1 \pm 0.5 \pm 0.3$
1.2	1.8	$0.7 \pm 0.3 \pm 0.1$	$1.4 \pm 0.4 \pm 0.2$




FIG. 5. (color online) Invariant yield as a function of the number of participating nucleons for $1.2 < |y| < 2.2$ and $1 < p_T < 5$ GeV/ c . The centrality bins are 0%–20%, 20%–40% and 40%–93%, and the data points are placed at the mean N_{part} calculated from a Glauber simulation. The data points for the Cu-going direction, $1.2 < y < 2.2$, are shifted along the x-axis to $N_{\text{part}} + 3$ to make the points visible, while the Au-going direction, $-2.2 < y < -1.2$, remains unshifted. The values are shown in Table IV.

409 rapidity. In Cu+Au collisions, the J/ψ meson yield is strongly suppressed in the Au-going direction compared to the
 410 ϕ meson yield at the same rapidity. This is similar to the differences previously observed between J/ψ and ϕ meson
 411 nuclear modification in d +Au collisions [13]. These differences could be attributed to a larger J/ψ break up cross
 412 section, effects in the higher-energy-density backward-rapidity region, or changes between soft and hard production
 413 mechanisms between the two mesons.

414 The forward and backward differences can be quantified by the ratio of the yield values for the forward rapidity
 415 (Cu-going direction) to the backward rapidity (Au-going direction). Fig. 11 shows the forward/backward ratio as a
 416 function of participating nucleons for $1.2 < |y| < 2.2$ and $1 < p_T < 5$ GeV/ c . The Type-C and Type-B systematic
 417 uncertainties, except for the $A\varepsilon_{\text{rec}}$ uncertainty, cancel when taking this ratio. The remaining systematic uncertainties
 418 are the Type-A signal extraction uncertainty and the Type-B $A\varepsilon_{\text{rec}}$ uncertainty. The difference in suppression between
 419 the forward and backward rapidity is more noticeable in the most central collisions, 0%–20%. In this centrality bin,
 420 the probability of observing the forward/backward ratio greater than or equal to unity was found to be p -value=1.2%,
 421 corresponding to a statistical significance of 2.3σ . The particle multiplicity for central collisions should be about 20%
 422 higher in the Au-going direction than in the Cu-going direction [35], however, the much smaller ratio observed may
 423 indicate that increased recombination effects or additional thermal strangeness production may also occur at higher
 424 energy density. In central collisions, the forward/backward ratio in ϕ production (~ 0.2) is smaller than that in J/ψ
 425 production (~ 0.8) in Cu+Au collisions [16].




FIG. 6. (color online) Invariant yield as a function of transverse momentum for $1.2 < |y| < 2.2$ and 0%–93% centrality. The p_T bins are $1 < p_T \leq 2.5$ and $2.5 < p_T < 5$ GeV/c, and the data points are placed at the mean p_T of the bin. The Cu-going direction corresponds to the forward rapidity, $1.2 < y < 2.2$, while the Au-going direction corresponds to the backward rapidity, $-2.2 < y < -1.2$. The values are shown in Table V.




FIG. 7. (color online) Invariant yield as a function of rapidity for $1 < p_T < 5$ GeV/c and 0%–93% centrality. The rapidity bins are $1.2 < |y| < 1.8$ and $1.8 < |y| < 2.2$ and the data points are placed at the mean y of the bin. The Cu-going direction covers the region $1.2 < y < 2.2$, while the Au-going direction covers the region $-2.2 < y < -1.2$. The values are shown in Table VI.

V. SUMMARY

426

427 In summary, ϕ meson production and its nuclear modification have been measured in Cu+Au collisions at $\sqrt{s_{NN}} =$
 428 200 GeV for $1.2 < |y| < 2.2$ and $1.0 < p_T < 5.0$ GeV/c via the dimuon decay channel. This first measurement of ϕ
 429 meson production and its nuclear modification in a heavy-ion system at forward/backward rapidity at RHIC extends
 430 measurements of ϕ from smaller systems, $p+p$ and $d+Au$, in the forward and backward rapidity. The invariant yields
 431 and nuclear modification factors have been presented here as a function of N_{part} , p_T and rapidity.




FIG. 8. (color online) The nuclear modification factor R_{CuAu} as a function of the number of participating nucleons for $1.2 < |y| < 2.2$ and $1 < p_T < 5$ GeV/c. The centrality bins are 0%–20%, 20%–40% and 40%–93%, and the data points are placed at the mean N_{part} calculated from a Glauber simulation. The data points for the Cu-going direction, $1.2 < y < 2.2$, are shifted along the x-axis to $N_{\text{part}}+3$ to make the points visible, while the data points for the Au-going direction, $-2.2 < y < -1.2$, remain unshifted. The values are shown in Table VII.




FIG. 9. (color online) The nuclear modification factor R_{CuAu} as a function of transverse momentum for $1.2 < |y| < 2.2$ and 0%–93% centrality. The p_T bins are $1 < p_T \leq 2.5$ and $2.5 < p_T < 5$ GeV/c, and the data points are placed at the mean p_T of the bin. The Cu-going direction corresponds to the forward rapidity, $1.2 < y < 2.2$, while the Au-going direction corresponds to the backward rapidity, $-2.2 < y < -1.2$. The values are shown in Table VIII.

432 The ϕ meson yields in Cu+Au collisions are found to be generally smaller in the Cu-going direction than in
 433 the Au-going direction. This is most pronounced in the most central events, 0%–20%, and at low momentum,
 434 1.0–2.5 GeV/c. In central collisions (0%–20%), the forward/backward ratio is below unity at a confidence level
 435 of 99%. It has been shown that these results follow a trend similar to what was seen previously at PHENIX in




FIG. 10. (color online) The nuclear modification factor R_{CuAu} as a function of rapidity for $1 < p_T < 5$ GeV/c and 0%–93% centrality. The rapidity bins are $1.2 < |y| < 1.8$ and $1.8 < |y| < 2.2$ and the data points are placed at the mean y of the bin. The values are shown in Table IX. Also included are previous PHENIX results for ϕ mesons in d +Au collisions [13] represented by open circles and J/ψ mesons in Cu+Au collisions [16] represented by open triangles. Positive rapidity, $1.2 < y < 2.2$, corresponds to the Cu-going and d-going directions, while negative rapidity, $-2.2 < y < -1.2$, is the Au-going direction.




FIG. 11. (color online) The forward/backward ratio as a function of the number of participating nucleons for $1 < p_T < 5$ GeV/c and $1.2 < |y| < 2.2$. The values are shown in Table IV. The Cu-going direction covers positive rapidity, $1.2 < y < 2.2$, while the Au-going direction covers negative rapidity, $-2.2 < y < -1.2$.

436 d +Au at the same rapidity and energy [13] as well as the ALICE measurement in p +Pb collisions at larger rapidity
 437 ($-4.46 < y < -2.96$ and $2.03 < y < 3.53$) and higher energy ($\sqrt{s_{NN}} = 5.02$ TeV) [23]. While this agreement could
 438 imply a role for CNM effects on ϕ production in Cu+Au collisions, the production of ϕ in heavy-ion collisions for these
 439 kinematics is expected to have substantial contributions from HNM effects as which were demonstrated to dominate

440 previous measurements at midrapidity for both Cu+Cu and Au+Au collisions [18]. A competition between CNM
 441 and HNM production mechanisms appears relevant for ϕ production at forward rapidity for heavy-ion collisions and
 442 a comprehensive description is needed from soft and hard physics models. Although the ϕ meson is sensitive to both
 443 CNM and HNM effects, this study was statistically limited, a factor that also affects the precise determination of the
 444 systematic uncertainties. A high statistics measurement and theory calculations are both needed in order to make
 445 conclusions about the various physics processes that might be at play here, including modifications of strangeness
 446 production in bulk matter and quark recombination.

447 ACKNOWLEDGMENTS

448 We thank the staff of the Collider-Accelerator and Physics Departments at Brookhaven National Laboratory and
 449 the staff of the other PHENIX participating institutions for their vital contributions. We acknowledge support from
 450 two Offices within the Office of Science of the Department of Energy: The Office of Nuclear Physics and the Office
 451 of Science Graduate Student Research award program, which is administered by the Oak Ridge Institute for Science
 452 and Education, and support from the National Science Foundation, Abilene Christian University Research Council,
 453 Research Foundation of SUNY, and Dean of the College of Arts and Sciences, Vanderbilt University (U.S.A), Ministry
 454 of Education, Culture, Sports, Science, and Technology and the Japan Society for the Promotion of Science (Japan),
 455 Conselho Nacional de Desenvolvimento Científico e Tecnológico and Fundação de Amparo à Pesquisa do Estado de
 456 São Paulo (Brazil), Natural Science Foundation of China (P. R. China), Croatian Science Foundation and Ministry of
 457 Science, Education, and Sports (Croatia), Ministry of Education, Youth and Sports (Czech Republic), Centre National
 458 de la Recherche Scientifique, Commissariat à l'Énergie Atomique, and Institut National de Physique Nucléaire et de
 459 Physique des Particules (France), Bundesministerium für Bildung und Forschung, Deutscher Akademischer Austausch
 460 Dienst, and Alexander von Humboldt Stiftung (Germany), National Science Fund, OTKA, Károly Róbert Univer-
 461 sity College, and the Ch. Simonyi Fund (Hungary), Department of Atomic Energy and Department of Science and
 462 Technology (India), Israel Science Foundation (Israel), Basic Science Research Program through NRF of the Ministry
 463 of Education (Korea), Physics Department, Lahore University of Management Sciences (Pakistan), Ministry of Ed-
 464 ucation and Science, Russian Academy of Sciences, Federal Agency of Atomic Energy (Russia), VR and Wallenberg
 465 Foundation (Sweden), the U.S. Civilian Research and Development Foundation for the Independent States of the
 466 Former Soviet Union, the Hungarian American Enterprise Scholarship Fund, and the US-Israel Binational Science
 467 Foundation.

-
- 468 [1] I. Arsene *et al.* (BRAHMS Collaboration), “Quark gluon plasma and color glass condensate at RHIC? The Perspective
 469 from the BRAHMS experiment,” Nucl. Phys. A **757**, 1 (2005).
 470 [2] B. B. Back *et al.*, “The PHOBOS perspective on discoveries at RHIC,” Nucl. Phys. A **757**, 28 (2005).
 471 [3] J. Adams *et al.* (STAR Collaboration), “Experimental and theoretical challenges in the search for the quark gluon plasma:
 472 The STAR Collaboration’s critical assessment of the evidence from RHIC collisions,” Nucl. Phys. A **757**, 102 (2005).
 473 [4] K. Adcox *et al.* (PHENIX Collaboration), “Formation of dense partonic matter in relativistic nucleus-nucleus collisions at
 474 RHIC: Experimental evaluation by the PHENIX Collaboration,” Nucl. Phys. A **757**, 184 (2005).
 475 [5] P. Koch, B. Muller, and J. Rafelski, “Strangeness in Relativistic Heavy Ion Collisions,” Phys. Rept. **142**, 167 (1986).
 476 [6] A. Shor, “ ϕ meson production as a probe of the quark gluon plasma,” Phys. Rev. Lett. **54**, 1122 (1985).
 477 [7] A. Andronic, “An overview of the experimental study of quark-gluon matter in high-energy nucleus-nucleus collisions,”
 478 *Proceedings, 26th International Symposium on Lepton Photon Interactions at High Energy (LP13)*, Int. J. Mod. Phys.
 479 **A29**, 1430047 (2014).
 480 [8] C. M. Ko and D. Seibert, “What can we learn from a second ϕ meson peak in ultrarelativistic nuclear collisions?” Phys.
 481 Rev. C **49**, 2198 (1994).
 482 [9] J. J. Heckman, J. Seo, and C. Vafa, “Phase Structure of a Brane/Anti-Brane System at Large N ,” J. High Energy Phys.
 483 **07 (2007)** 073.
 484 [10] I. Vitev, “Initial state parton broadening and energy loss probed in d +Au at RHIC,” Phys. Lett. B **562**, 36 (2003).
 485 [11] J. W. Cronin, Henry J. Frisch, M. J. Shochet, J. P. Boymond, R. Mermod, P. A. Piroue, and R. L. Sumner, “Production
 486 of Hadrons with Large Transverse Momentum at 200 GeV, 300-GeV, and 400-GeV,” *High energy physics. Proceedings,
 487 17th International Conference, ICHEP 1974, London, England, July 01-July 10, 1974*, Phys. Rev. D **11**, 3105 (1975).
 488 [12] A. Accardi and M. Gyulassy, “Cronin effect versus geometrical shadowing in $d + Au$ collisions at RHIC,” Phys. Lett. B
 489 **586**, 244 (2004).
 490 [13] A. Adare *et al.* (PHENIX Collaboration), “ ϕ meson production in $d + Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV,”
 491 arXiv:1506.08181.

- 492 [14] S. S. Adler *et al.* (PHENIX Collaboration), “Nuclear Modification Factors for Hadrons at Forward and Backward Rapidities
493 in Deuteron-Gold Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. Lett.* **94**, 082302 (2005).
- 494 [15] A. Adare *et al.* (PHENIX Collaboration), “System-size dependence of open-heavy-flavor production in nucleus-nucleus
495 collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **90**, 034903 (2014).
- 496 [16] A. Adare *et al.* (PHENIX Collaboration), “Nuclear matter effects on J/ψ production in asymmetric Cu + Au collisions
497 at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **90**, 064908 (2014).
- 498 [17] S. S. Adler *et al.* (PHENIX Collaboration), “Production of ϕ mesons at midrapidity in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions
499 at RHIC,” *Phys. Rev. C* **72**, 014903 (2005).
- 500 [18] A. Adare *et al.* (PHENIX Collaboration), “Nuclear modification factors of ϕ mesons in d +Au, Cu+Cu and Au+Au
501 collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **83**, 024909 (2011).
- 502 [19] A. Adare *et al.* (PHENIX Collaboration), “Measurement of neutral mesons in p + p collisions at $\sqrt{s} = 200$ GeV and scaling
503 properties of hadron production,” *Phys. Rev. D* **83**, 052004 (2011).
- 504 [20] A. Adare *et al.* (PHENIX Collaboration), “Low-mass vector-meson production at forward rapidity in p + p collisions at
505 $\sqrt{s} = 200$ GeV,” *Phys. Rev. D* **90**, 052002 (2014).
- 506 [21] B. I. Abelev *et al.* (STAR Collaboration), “Partonic flow and ϕ -meson production in Au+Au collisions at $\sqrt{s_{NN}} = 200$,”
507 *Phys. Rev. Lett.* **99**, 112301 (2007).
- 508 [22] B. I. Abelev *et al.* (STAR Collaboration), “Energy and system size dependence of ϕ meson production in Cu+Cu and
509 Au+Au collisions,” *Phys. Lett. B* **673**, 183 (2009).
- 510 [23] J. Adam *et al.* (ALICE Collaboration), “ ϕ -meson production at forward rapidity in p -Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV
511 and in pp collisions at $\sqrt{s} = 2.76$ TeV,” arXiv:1506.09206.
- 512 [24] B. B. Abelev *et al.* (ALICE Collaboration), “ $K^*(892)^0$ and $\phi(1020)$ production in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,”
513 *Phys. Rev. C* **91**, 024609 (2015).
- 514 [25] K. Adcox *et al.* (PHENIX Collaboration), “PHENIX detector overview,” *Nucl. Instrum. Methods Phys. Res., Sec. A* **499**,
515 469 (2003).
- 516 [26] H. Akikawa *et al.* (PHENIX Collaboration), “PHENIX muon arms,” *Nucl. Instrum. Methods Phys. Res., Sec. A* **499**, 537
517 (2003).
- 518 [27] M. Allen *et al.* (PHENIX Collaboration), “PHENIX inner detectors,” *Nucl. Instrum. Methods Phys. Res., Sec. A* **499**, 549
519 (2003).
- 520 [28] A. Taketani *et al.* (PHENIX Collaboration), “Silicon vertex tracker for RHIC PHENIX experiment,” *Technology and
521 instrumentation in particle physics. Proceedings, 1st International Conference, TIPP09, Tsukuba, Japan, March 12-17,
522 2009*, *Nucl. Instrum. Methods Phys. Res., Sec. A* **623**, 374 (2010).
- 523 [29] C. Aidala *et al.*, “The PHENIX Forward Silicon Vertex Detector,” *Nucl. Instrum. Methods Phys. Res., Sec. A* **755**, 44
524 (2014).
- 525 [30] C. Baglin *et al.* (NA38 Collaboration), “The Production of J/ψ in 200-GeV/nucleon Oxygen Uranium Interactions,” *Phys.
526 Lett. B* **220**, 471 (1989).
- 527 [31] A. Adare *et al.* (PHENIX Collaboration), “ J/ψ suppression at forward rapidity in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,”
528 *Phys. Rev. C* **84**, 054912 (2011).
- 529 [32] T. Sjostrand, P. Eden, C. Friberg, L. Lonnblad, G. Miu, S. Mrenna, and E. Norrbin, “High-energy physics event generation
530 with PYTHIA 6.1,” *Comput. Phys. Commun.* **135**, 238 (2001).
- 531 [33] R. Brun, F. Carminati, and S. Giani, “GEANT Detector Description and Simulation Tool,” CERN-W5013 (1994).
- 532 [34] J. Beringer *et al.* (Particle Data Group Collaboration), “Rev. of Particle Phys. (RPP),” *Phys. Rev. D* **86**, 010001 (2012).
- 533 [35] L.-W. Chen and C. M. Ko, “Anisotropic flow in Cu+Au collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys. Rev. C* **73**, 014906 (2006).