



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Spectroscopy of the neutron-rich hypernucleus $_{\Lambda}^{7}\text{He}$ from electron scattering

T. Gogami *et al.* (HKS (JLab E05-115) Collaboration)

Phys. Rev. C **94**, 021302 — Published 12 August 2016

DOI: [10.1103/PhysRevC.94.021302](https://doi.org/10.1103/PhysRevC.94.021302)

Spectroscopy of the neutron-rich hypernucleus, ${}^7_{\Lambda}\text{He}$ by electron beam

T. Gogami,^{1,*} C. Chen,² D. Kawama,¹ P. Achenbach,³ A. Ahmidouch,⁴ I. Albayrak,² D. Androic,⁵ A. Asaturyan,⁶ R. Asaturyan,^{6,†} O. Ates,² P. Baturin,⁷ R. Badui,⁷ W. Boeglin,⁷ J. Bono,⁷ E. Brash,⁸ P. Carter,⁸ A. Chiba,¹ E. Christy,² S. Danagoulian,⁴ R. De Leo,⁹ D. Doi,¹ M. Elaasar,¹⁰ R. Ent,¹¹ Y. Fujii,¹ M. Fujita,¹ M. Furic,⁵ M. Gabrielyan,⁷ L. Gan,¹² F. Garibaldi,¹³ D. Gaskell,¹¹ A. Gasparian,⁴ Y. Han,² O. Hashimoto,^{1,†} T. Horn,¹¹ B. Hu,¹⁴ Ed.V. Hungerford,¹⁵ M. Jones,¹¹ H. Kanda,¹ M. Kaneta,¹ S. Kato,¹⁶ M. Kawai,¹ H. Khanal,⁷ M. Kohl,² A. Liyanage,² W. Luo,¹⁴ K. Maeda,¹ A. Margaryan,⁶ P. Markowitz,⁷ T. Maruta,¹ A. Matsumura,¹ V. Maxwell,⁷ A. Mkrtchyan,⁶ H. Mkrtchyan,⁶ S. Nagao,¹ S.N. Nakamura,¹ A. Narayan,¹⁷ C. Neville,⁷ G. Niculescu,¹⁸ M.I. Niculescu,¹⁸ A. Nunez,⁷ Nuruzzaman,¹⁷ Y. Okayasu,¹ T. Petkovic,⁵ J. Pochodzalla,³ X. Qiu,¹⁴ J. Reinhold,⁷ V.M. Rodriguez,¹⁹ C. Samanta,²⁰ B. Sawatzky,¹¹ T. Seva,⁵ A. Shichijo,¹ V. Tadevosyan,⁶ L. Tang,^{2,11} N. Taniya,¹ K. Tsukada,¹ M. Veilleux,⁸ W. Vulcan,¹¹ F.R. Wesselmann,²¹ S.A. Wood,¹¹ T. Yamamoto,¹ L. Ya,² Z. Ye,² K. Yokota,¹ L. Yuan,² S. Zhamkochyan,⁶ and L. Zhu²

(HKS (JLab E05-115) Collaboration)

¹Graduate School of Science, Tohoku University, Sendai, Miyagi 980-8578, Japan

²Department of Physics, Hampton University, Hampton, VA 23668, USA

³Institute for Nuclear Physics, Johannes Gutenberg-University, D-55099 Mainz, Germany

⁴Department of Physics, North Carolina A&T State University, Greensboro, NC 27411, USA

⁵Department of Physics & Department of Applied Physics, University of Zagreb, HR-10000 Zagreb, Croatia

⁶A.I.Alikhanyan National Science Laboratory, Yerevan 0036, Armenia

⁷Department of Physics, Florida International University, Miami, FL 27411, USA

⁸Department of Physics, Computer Science & Engineering, Christopher Newport University, Newport News, VA, USA 23606

⁹Istituto Nazionale di Fisica Nucleare, Sezione di Bari and University of Bari, I-70126 Bari, Italy

¹⁰Department of Physics, Southern University at New Orleans, New Orleans, LA 70126, USA

¹¹Thomas Jefferson National Accelerator Facility (JLab), Newport News, VA 23606, USA

¹²Department of Physics, University of North Carolina at Wilmington, Wilmington, NC 28403, USA

¹³INFN, Sezione Sanità and Istituto Superiore di Sanità, 00161 Rome, Italy

¹⁴Nuclear Physics Institute, Lanzhou University, Gansu 730000, China

¹⁵Department of Physics, University of Houston, Houston, Texas 77204, USA

¹⁶Department of Physics, Yamagata University, Yamagata, 990-8560, Japan

¹⁷Mississippi State University, Mississippi State, Mississippi 39762, USA

¹⁸Department of Physics and Astronomy, James Madison University, Harrisonburg, VA 22807, USA

¹⁹Escuela de Ciencias y Tecnología, Universidad Metropolitana, San Juan, 00928, Puerto Rico

²⁰Department of Physics & Astronomy, Virginia Military Institute, Lexington, Virginia 24450, USA

²¹Department of Physics, Xavier University of Louisiana, New Orleans, LA 70125, USA

The missing mass spectroscopy of the ${}^7_{\Lambda}\text{He}$ hypernucleus was performed, using the ${}^7\text{Li}(e, e'K^+){}^7_{\Lambda}\text{He}$ reaction at JLab Hall-C. The Λ binding energy of the ground state ($1/2^+$) was determined with a smaller error than that of the previous measurement, being $B_{\Lambda} = 5.55 \pm 0.10^{\text{stat.}} \pm 0.11^{\text{sys.}}$ MeV. The experiment also provided new insight into charge symmetry breaking in p -shell hypernuclear systems. Finally, a peak at $B_{\Lambda} = 3.65 \pm 0.20^{\text{stat.}} \pm 0.11^{\text{sys.}}$ MeV was observed and assigned as a mixture of $3/2^+$ and $5/2^+$ states, confirming the “glue-like” behavior of Λ , which makes an unstable state in ${}^6\text{He}$ stable against neutron emission.

Nuclear physicists explore the low energy behavior of the strongly interacting many-body systems, extracting an effective potential which can be used for nuclear structure and interaction calculations. Effective potential techniques can also be applied to hypernuclear systems, as the lifetime of a hyperon in a nucleus is much greater than the relaxation time associated with strong interactions. On the other hand, the two-body potentials for the hyperon-nucleon interaction, YN , are not determined as well as those for NN , due to the experimental difficulties

of producing and detecting hyperons in free scattering experiments. However, embedding a hyperon within the nuclear medium (hypernucleus) does allow extraction of effective potentials from detailed measurements of hypernuclear energy levels and transitions.

Although many species of Λ hypernuclei with masses $A \leq 209$ have been observed [1, 2], more systematic and precise data are still needed for further insight into the ΛN interaction. Nowadays, experimental studies of Λ hypernuclei use: 1) hadron beams at J-PARC [3, 4], 2) heavy ion beams at GSI [5–7], 3) heavy ion colliders at RHIC [8] and LHC [9], and 4) electron beams at MAMI [10, 11] and JLab [12–21]. The different production mechanisms are complementary and allow to use their specific sensitivities to excite particular structures

* Current address: Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047, Japan

† Deceased

which highlight nuclear features of interest.

One such feature of interest is charge symmetry breaking (CSB) in Λ hypernuclei. The difference in ground-state binding energies in the $A = 3$ non-strange nuclei (${}^3\text{He}$ and ${}^3\text{H}$) is 0.7638 ± 0.0003 MeV [22]. There remains a binding-energy difference of 0.081 MeV after accounting for the 0.683 MeV Coulomb correction [23]. In s -shell hypernuclei, a large CSB, $\Delta B_\Lambda({}^4_\Lambda\text{He} - {}^4_\Lambda\text{H}; 0^+) = B_\Lambda({}^4_\Lambda\text{He}; 0^+) - B_\Lambda({}^4_\Lambda\text{H}; 0^+) = (2.39 \pm 0.03) - (2.04 \pm 0.04) = +0.35 \pm 0.06$ MeV, is found by comparing the ground state binding energies between ${}^4_\Lambda\text{H}$ and ${}^4_\Lambda\text{He}$ [24]. Though the Coulomb effect of the core nuclei, ${}^3\text{He}$ and ${}^3\text{H}$, is already subtracted in the $\Delta B_\Lambda({}^4_\Lambda\text{He} - {}^4_\Lambda\text{H}; 0^+)$ calculation, the binding energy difference of the above hypernuclear isospin doublet is due, in part, to differences in the Coulomb energy caused by contraction of the nucleus as a result of the additional Λ binding. The Coulomb-energy correction was predicted as $\Delta B_C = 0.02\text{--}0.08$ MeV [25–27], and thus $\Delta B_\Lambda({}^4_\Lambda\text{He} - {}^4_\Lambda\text{H}; 0^+) + \Delta B_C \simeq 0.4$ MeV is attributed to the ΛN CSB for the 0^+ state in the $A = 4$ iso-doublet hypernuclear system. This difference in the binding energy is approximately five times larger than for $A = 3$ non-strange nuclei. Recent gamma-ray measurement indicates that little binding energy difference exists between the (1^+) excited states [4] though it was believed that the 1^+ excited states had as large CSB as the ground states [28–30]. These residual differences are difficult to explain by Coulomb energy alone. A detailed discussion of hypernuclear CSB [31–33], in addition to other topics of interest, has been recently published [34].

CSB in p -shell hypernuclear systems is predicted to be smaller than in s -shell systems [31]. Hence, differences in Λ binding energies between p -shell mirror hypernuclei are predicted to be less than a few 100 keV [31]. Previous experiments at JLab Hall-C measured Λ binding energies of ${}^7_\Lambda\text{He}$ [18], ${}^9_\Lambda\text{Li}$ [35], ${}^{10}_\Lambda\text{Be}$ [21], ${}^{12}_\Lambda\text{B}$ [12, 13, 19], ${}^{28}_\Lambda\text{Al}$ [16], and ${}^{52}_\Lambda\text{V}$ [35] via the $(e, e'K^+)$ reaction. The present paper reports a new result for Λ binding energy of ${}^7_\Lambda\text{He}$ with an improved systematic error, and is compared to its isotopic mirror hypernuclei. In addition, due to improved statistics, the experiment extracted the first observation of a peak corresponding to the excited states ($3/2^+$, $5/2^+$) of ${}^7_\Lambda\text{He}$.

CSB in hypernuclear p -shell systems can be studied by comparing the Λ binding energies for $A = 7$, isotriplet ($T = 1$) Λ hypernuclei, which are the simplest p -shell hypernuclear systems, ${}^7_\Lambda\text{He}$ ($\alpha + n + n + \Lambda$), ${}^7_\Lambda\text{Li}^*$ ($\alpha + p + n + \Lambda$) and ${}^7_\Lambda\text{Be}$ ($\alpha + p + p + \Lambda$). The isospin of the ground state of ${}^7_\Lambda\text{Li}$ is $T = 0$. Thus, an excited state of ${}^7_\Lambda\text{Li}$ with $T = 1$ should be compared with the isotriplet partners. The ground state binding energies of ${}^7_\Lambda\text{Li}$ ($T = 0$) and ${}^7_\Lambda\text{Be}$ were measured to be 5.58 ± 0.03 MeV and 5.16 ± 0.08 MeV, respectively, by the emulsion experiments [36]. The binding energy of ${}^7_\Lambda\text{Li}^*$ ($T = 1$) is obtained as 5.26 ± 0.03 MeV by using information of the energy spacing of $E_x({}^7_\Lambda\text{Li}^*; T = 1, 1/2^+) = 3.88$ MeV measured by the γ -ray spectroscopy [37] and the excitation

energy of $E_x({}^6\text{Li}^*; T = 1) = 3.56$ MeV [38]. The ground-state ($1/2^+$) Λ binding energy of ${}^7_\Lambda\text{He}$ using the $(e, e'K^+)$ reaction at JLab Hall-C (JLab E01-011), was successfully determined to be $B_\Lambda = 5.68 \pm 0.03^{\text{stat.}} \pm 0.25^{\text{sys.}}$ MeV [18]. As a result, the measured energies of $A = 7$, $T = 1$ hypernuclei differ from a cluster model prediction which used a phenomenological ΛN CSB potential which was constructed to reproduce the energies of ${}^4_\Lambda\text{He}$ and ${}^4_\Lambda\text{H}$ [39]. The error on the Λ binding energy of ${}^7_\Lambda\text{He}$ was larger than for other Λ hypernuclei, and was dominated by systematic contributions. Therefore the present experiment (JLab E05-115) focused on the determination of the Λ binding energy of ${}^7_\Lambda\text{He}$ with particular emphasis in reducing the systematic error.

The core nucleus, ${}^6\text{He}$ ($\alpha + n + n$) in ${}^7_\Lambda\text{He}$ is known as a typical neutron-halo nucleus. The first-excited state energy of ${}^6\text{He}$ (2^+) was measured to be 0.824 MeV above the $\alpha + n + n$ breakup threshold, having a decay width of $\Gamma = 0.113$ MeV [38]. The corresponding states of ${}^7_\Lambda\text{He}$ ($3/2^+$, $5/2^+$), in which a Λ resides in the s -orbit, are predicted to be stable against neutron-emission breakup [39–41] due to the attractive ΛN interaction. In addition, the existence of isomeric states in ${}^7_\Lambda\text{He}$ [42–44] was speculated from widely scattered binding energy obtained by the emulsion experiment although it had not been confirmed yet experimentally. The production cross section for a sum of these states ($3/2^+$, $5/2^+$) with the (γ, K^+) reaction at small K^+ scattering angle was predicted to be $\approx 60\%$ of that for the ground state ($1/2^+$) [40]. Although a small structure was observed in the spectrum which might correspond to the $3/2^+$ and $5/2^+$ states, a lack of statistics prevented confirmation of the observation of these states in the previous measurement [18]. The present experiment acquired five times higher statistics and can now confirm observation of these states and thus the “glue-like” behavior of the Λ . This paper reports the observation of the ground state ($1/2^+$), and for the first time, the observation of the $3/2^+$ and $5/2^+$ states.

The $(e, e'K^+)$ reaction was used for Λ hypernuclear production. Electroproduction is related to photoproduction through a virtual photon produced in the (e, e') reaction [45–47]. In the geometry for JLab E05-115, the virtual photon can be treated as almost real since the square of the four momentum transfer, $Q^2 (= -q^2 > 0)$ is quite small ($Q^2 \simeq 0.01$ [GeV/c] 2). The experimental kinematics can be found in [19]. We used a continuous wave electron beam with an energy of $E_e = 2.344$ GeV, provided by the CEBAF (Continuous Electron Beam Accelerator Facility) at JLab. The electron beam was transported to the experimental target which was installed at the entrance of a charge separation dipole magnet (splitter magnet; SPL). The K^+ s ($p_{K^+}^{\text{center}} = 1.200$ GeV/c) and scattered electrons ($p_{e'}^{\text{center}} = 0.844$ GeV/c) were bent in opposite directions by SPL, and were analyzed with the High Resolution Kaon Spectrometer (HKS) [48, 49] and High Resolution Electron Spectrometer (HES), respectively. Details for the experimental setup are described in [19, 21, 48]. One important feature of the present ex-

periment is the excellent resolution of $\Delta p/p \simeq 2 \times 10^{-4}$ (FWHM) for both K^+ and e' at approximately 1 GeV/c, due to the optics of the SPL+HKS+HES system. Thus, an energy resolution of about 0.5 MeV (FWHM) was obtained for hypernuclear spectroscopy [19, 21].

The positions and angles of the K^+ s and scattered electrons at reference planes in the magnetic spectrometers were measured by particle detectors. This information was converted to momentum vectors at the target position with backward transfer matrices (BTM) representing the optical systems for the SPL+HES and SPL+HKS, respectively, in order to reconstruct the missing mass (M_{HYP}). Once the missing mass was obtained, the Λ binding energy (B_Λ) was calculated as $B_\Lambda(^A_\Lambda Z) = M(^{A-1}Z) + M_\Lambda - M_{\text{HYP}}(^A_\Lambda Z)$, where Z denotes the proton number, and $M(^{A-1}Z)$ and M_Λ are the masses of a core nucleus and a Λ .

The energy scale calibration was performed by optimizing the backward transfer matrices (BTM) of the magnetic spectrometer systems [19]. The BTM optimization is also correlated with energy resolution in the resulting hypernuclear spectra. For the BTM optimization, events of Λ and Σ^0 production from the 0.45 g/cm² polyethylene target were used along with events from the production of the $^{12}_\Lambda\text{B}$ ground state from a 0.0875 g/cm² natural carbon target. Systematic errors, which originated from the BTM optimization process, needed to be estimated carefully since the BTM optimization mainly determines the accuracy of the binding energy (B_Λ) and excitation energy (E_Λ) of a Λ hypernucleus. In order to estimate the achievable energy accuracy, a fully-modeled Monte Carlo simulation was performed. As a result, it was found that B_Λ and E_Λ could be determined with accuracies of < 0.09 MeV and < 0.05 MeV, respectively, by this optimization method. Another major contribution to the uncertainty on the binding energy is due to energy loss corrections for the particles in the target. This contribution was also estimated by the Monte Carlo simulation taking into account the target thickness uncertainty. Finally, the total systematic errors for B_Λ and E_Λ were estimated as 0.11 MeV and 0.05 MeV, respectively [19, 21].

An enriched ^7Li target (purity of 99%) with a thickness of 0.208 g/cm² was used for the $^7_\Lambda\text{He}$ production. The nominal beam current for the production run of $^7_\Lambda\text{He}$ was 35 μA , and the total incident charge on the ^7Li target was 4.839 C ($\simeq 3 \times 10^{19}$ electrons). Figure 1 shows the obtained binding energy ($-B_\Lambda$) spectrum with an ordinate axis of $(\frac{d\sigma}{d\Omega_K})$ as defined in [21]. For the binding energy calculation, the nuclear masses of $M(^7\text{Li}) = 5605.54$ MeV and $M(^6\text{He}) = 6533.83$ MeV [50] were used. Events from quasi-free Λ production were distributed in the region of $-B_\Lambda > 0$. The distribution of the accidental $e'K^+$ -coincidence events in the spectrum was obtained by the mixed events analysis [18]. This method provides an accidental coincidence spectrum with high statistics thus reducing the statistical uncertainty caused by background subtraction.

Figure 2 shows the spectrum of the Λ binding en-




FIG. 1. The binding energy ($-B_\Lambda$) spectrum of $^7_\Lambda\text{He}$ with an ordinate axis of $(\frac{d\sigma}{d\Omega_K})$ defined in [21].




FIG. 2. The spectrum of the binding energy ($-B_\Lambda$) and the excitation energy ($E_\Lambda \equiv -(B_\Lambda - B_\Lambda(\#1))$) for the $^7\text{Li}(e, e'K^+)^7_\Lambda\text{He}$ reaction with an ordinate axis of $(\frac{d\sigma}{d\Omega_K})$ after the accidental $e'K^+$ -coincidence distribution was subtracted. The curve is a fit with two Voigt functions.

ergy ($-B_\Lambda$) and the excitation energy ($E_\Lambda \equiv -(B_\Lambda - B_\Lambda(\#1))$) of $^7_\Lambda\text{He}$ after the accidental $e'K^+$ -coincidence distribution was subtracted. In order to find peak candidates, a peak search by tests of statistical significance defined as $S/\sqrt{S+N}$ was applied. The statistical significance was calculated for each bin of the histogram, and the tests for robustness used several settings of bin size to find peak candidates, taking into account the energy resolution. As a result, two peak candidates were found with peak separation of $\geq 3\sigma$ as labeled by #1 and #2 in Fig. 2. The statistical significance for peak #1 is

7.5σ in a range of -7.0 to -4.0 MeV, which is larger than that of the previous measurement (5.5σ) [18]. The two peak candidates were fitted by Voigt functions (convolution of Gaussian and Lorentzian functions) to obtain B_Λ (E_Λ) and the differential cross section for each peak. The fitting results are given in Table I. The en-

TABLE I. Fitting results for the Λ binding energy, excitation energy (E_Λ), and $\left(\frac{d\sigma}{d\Omega_K}\right)$ defined in [21] for ${}^7\text{Li}(e, e'K^+){}_\Lambda^7\text{He}$. Errors are statistical and systematic.

Peak	Number of events	B_Λ [MeV] (E_Λ [MeV])	$\left(\frac{d\sigma}{d\Omega_K}\right)$ [nb/sr]
#1	413 ± 38	$5.55 \pm 0.10 \pm 0.11$ (0.0)	$10.7 \pm 1.0 \pm 1.8$
#2	239 ± 22	$3.65 \pm 0.20 \pm 0.11$ ($1.90 \pm 0.22 \pm 0.05$)	$6.2 \pm 0.6 \pm 1.1$

ergy resolution was obtained to be 1.3 MeV (FWHM) which is consistent with the estimation by the Monte Carlo simulation, although our previously published result for ${}_\Lambda^12\text{B}$ [19] was better (FWHM $\simeq 0.54$ MeV). In the Monte Carlo simulation, it was found that our BTMs have a momentum dependence on the z -displacement (beam direction) from the interaction point. This dependence significantly contributes to the energy resolution, adding a kinematical contribution due to the large recoil of the light hypernuclear system. The length in z -direction of the ${}^7\text{Li}$ target (4.0 mm) was longer than that of the natural ${}^{12}\text{C}$ target (0.5 mm) used for a measurement of ${}^{12}\text{C}(e, e'K^+){}_\Lambda^12\text{B}$ [19]. Thus, the peak width for ${}_\Lambda^7\text{He}$ increased with respect to ${}_\Lambda^12\text{B}$ result as the simulation indicated.

Peak #1 is considered as the ground state of ${}_\Lambda^7\text{He}$ (${}^6\text{He}[J_C; E_x] \otimes j^\Lambda = [0^+; \text{g.s.}] \otimes s_{1/2}^\Lambda = 1/2^+$). The Λ binding energy of the $1/2^+$ state was obtained to be $5.55 \pm 0.10^{\text{stat.}} \pm 0.11^{\text{sys.}}$ MeV which is consistent with the previous result ($5.68 \pm 0.03^{\text{stat.}} \pm 0.25^{\text{sys.}}$ MeV) [18] but with improved uncertainty. For the previous results, the statistical error is smaller since the energy resolution for ${}_\Lambda^7\text{He}$ spectrum is better while the systematic error dominates. In the present result, on the other hand, statistical and systematic errors are balanced, reducing the total uncertainty by optimizing the target thickness and the energy-calibration method.

Figure 3 shows the measured Λ binding energies of $A = 7$, $T = 1$ hypernuclei with statistical error bars, as compared to a theoretical prediction by a four-body cluster model [39]. Colored boxes on the results of ${}_\Lambda^7\text{He}$ indicate systematic errors on B_Λ . In the cluster model prediction [39], a phenomenological even-state CSB potential was introduced to reproduce the binding energies of ${}_\Lambda^4\text{He}$ and ${}_\Lambda^4\text{H}$. This was applied to the $A = 7$, $T = 1$ hypernuclear system. Binding energy predictions without and with the phenomenological CSB potential are shown by solid and dashed lines, respectively, in Fig. 3. The




FIG. 3. Measured Λ binding energies of ${}_\Lambda^7\text{He}$ (present result and [18]), ${}_\Lambda^7\text{Li}^*$ [36, 37], and ${}_\Lambda^7\text{Be}$ [36] for the $1/2^+$ state with statistical error bars. Colored boxes on the experimental results of ${}_\Lambda^7\text{He}$ indicate systematic errors on B_Λ . Solid and dashed lines represent theoretical calculations without and with a phenomenological even-state ΛN CSB potential, which reproduces Λ binding energies of ${}_\Lambda^4\text{He}$ and ${}_\Lambda^4\text{H}$, by a four-body cluster model [39].

present result seems to favor the energy prediction without the phenomenological CSB potential. This is also the case for the other experimental data in the $A = 7$, $T = 1$ system. This comparison suggests that a phenomenological CSB potential needs further consideration. It is possible to introduce a strong odd-state CSB potential in addition to one for the even-state in order to make the experiment and theoretical prediction more consistent [39, 51], although the validity of a strong odd-state interaction can be questioned [52]. It was suggested that the CSB interaction needs inclusion of explicit ΛN - ΣN coupling [31]. It seems clear that further systematic studies, with more precise data particularly for the p -shell hypernuclei are needed.

Peak #2 was obtained at $B_\Lambda(\#2) = 3.65 \pm 0.20^{\text{stat.}} \pm 0.11^{\text{sys.}}$ MeV with a differential cross section of $6.2 \pm 0.6^{\text{stat.}} \pm 1.1^{\text{sys.}}$ nb/sr. Figure 4 shows a B_Λ comparison between the obtained results and theoretical predictions [40, 41] with energy levels of the core nucleus, ${}^6\text{He}$ [38]. Energy levels of the first excited doublet ($3/2^+$, $5/2^+$) are predicted to be approximately 1.7 MeV above the ground state ($1/2^+$) [40, 41]. Moreover, a ratio of the differential cross section of a sum of $3/2^+$ and $5/2^+$ states to that of the ground state ($1/2^+$) is predicted to be approximately 0.6. The value of E_Λ and the ratio of $\left(\frac{d\sigma}{d\Omega_K}\right)$ for peak #1 to peak #2 are $E_\Lambda = 1.90 \pm 0.20^{\text{stat.}} \pm 0.11^{\text{sys.}}$ MeV and $\simeq 0.58$, respectively. The results are consistent with the above theoretical predictions, and thus, peak #2 is interpreted as ${}^6\text{He}[J_C; E_x] \otimes j^\Lambda = [2^+; 1.8 \text{ MeV}] \otimes s_{1/2}^\Lambda = (3/2^+, 5/2^+)$.




FIG. 4. Obtained energy levels of ${}^7_{\Lambda}\text{He}$ with theoretical predictions [40, 41]. Reported energy levels for ${}^6\text{He}$ is also shown [38].

The $3/2^+$ and $5/2^+$ states of ${}^7_{\Lambda}\text{He}$ are more than 2.3 MeV below the ${}^5_{\Lambda}\text{He}+n$ breakup energy [36, 38], which is the lowest neutron emission breakup, as shown in Fig. 4. On the other hand, the 2^+ state of ${}^6\text{He}$, which corresponds to the $3/2^+$ and $5/2^+$ states of ${}^7_{\Lambda}\text{He}$, was reported to be 0.824 MeV above the $\alpha+n+n$ energy [38] ($E = 1.797 \pm 0.025$ MeV) meaning that this state is not stable against neutron emission. Therefore, the present result of the peak for the $3/2^+$ and $5/2^+$ states in ${}^7_{\Lambda}\text{He}$ confirms the Λ “glue-like” role, making an unbound nucleus bound.

The successful observation of the first excited doublet in ${}^7_{\Lambda}\text{He}$ opens a door to study unstable states of light nuclei. For instance, energy-level predictions of the second 2^+ state (2_2^+) in a neutron-halo nucleus, ${}^6\text{He}$ are largely different depending on models as shown in [53]. Recently the excitation energy of 2_2^+ state of ${}^6\text{He}$ was measured to be $E_x = 2.6 \pm 0.3$ MeV with a width of $\Gamma = 1.6 \pm 0.4$ MeV by the two-neutron transfer reaction, $p({}^8\text{He}, t){}^6\text{He}$ [53]. This measurement would exclude several theoretical models. However, the 2_2^+ energy was derived from a spectral decomposition by fitting to a spectrum in which a few of major states are overlapping because of their large decay widths ($\Gamma =$ a few MeV).

On the other hand, the $3/2_2^+$ and $5/2_2^+$ states in ${}^7_{\Lambda}\text{He}$, which correspond to the 2_2^+ state in ${}^6\text{He}$, are predicted to be much narrower [41] due to the additional binding of the Λ . Therefore, a measurement of the $3/2_2^+$ and $5/2_2^+$ states in ${}^7_{\Lambda}\text{He}$ combined with a realistic cluster calculation may provide better understanding of the 2_2^+ state in ${}^6\text{He}$. The differential cross section of the sum of the $3/2_2^+$ and $5/2_2^+$ states in ${}^7_{\Lambda}\text{He}$ is predicted to be approximately

16% of that for the ground state [41]. Consequently, the observation of the $3/2_2^+$ and $5/2_2^+$ states in ${}^7_{\Lambda}\text{He}$ is promising for future spectroscopy at JLab using the $(e, e'K^+)$ reaction.

In summary, the spectroscopy of Λ hypernuclei was performed with a new magnetic spectrometer system, SPL+HKS+HES at JLab Hall-C via the $(e, e'K^+)$ reaction. A spectroscopic measurement of a neutron rich hypernucleus, ${}^7_{\Lambda}\text{He}$, was performed with an enriched ${}^7\text{Li}$ target, and the hypernuclear structure was successfully observed with an energy resolution of 1.3 MeV FWHM.

The ground state energy ($1/2^+$) of ${}^7_{\Lambda}\text{He}$ was determined to be $B_{\Lambda} = 5.55 \pm 0.10^{\text{stat.}} \pm 0.11^{\text{sys.}}$ MeV which was consistent with the result of the previous measurement and improved the total error. The Λ binding energy provides insight into charge symmetry breaking (CSB) effects of the ΛN interaction by comparison with the bindings of isotopic mirror hypernuclei in the $A = 7$, $T = 1$ system (${}^7_{\Lambda}\text{Li}^*$, ${}^7_{\Lambda}\text{Be}$). Further systematic investigation with better precision, particularly for p -shell hypernuclei, are necessary in order to deepen our understanding of ΛN CSB. The $(e, e'K^+)$ reaction at JLab provides a unique method to measure the absolute Λ binding energies of p -shell hypernuclei or heavier with less than a few 100 keV accuracy.

The first excited doublet ($3/2^+$, $5/2^+$) in ${}^7_{\Lambda}\text{He}$, which correspond to the 2^+ state in ${}^6\text{He}$, was successfully observed for the first time. A peak for a sum of the $3/2^+$ and $5/2^+$ was determined to be $B_{\Lambda} = 3.65 \pm 0.20^{\text{stat.}} \pm 0.11^{\text{sys.}}$ MeV with the differential cross section of $\left(\frac{d\sigma}{d\Omega_K}\right) = 6.2 \pm 0.6^{\text{stat.}} \pm 1.1^{\text{sys.}}$ nb/sr. The peak for the $3/2^+$ and $5/2^+$ states was found to be approximately 2.3 MeV below the lowest neutron emission energy. The result shows that the 2^+ state in ${}^6\text{He}$, which is an unstable state for the $\alpha+n+n$ breakup, becomes stable against neutron-emission breakup once a Λ is bound in the nucleus, owing to the additional binding of the Λ .

We thank the JLab staffs of the physics, accelerator, and engineering divisions for support of the experiment. Also, we thank E. Hiyama, M. Isaka, D.J. Millener, and T. Motoba for valuable exchanges related to their theoretical works. The program at JLab Hall-C is supported by JSPS KAKENHI Grant No. 12002001, No. 15684005, No. 16GS0201, and No. 24-4123 (Grant-in-Aid for JSPS fellow), JSPS Core-to-Core Program No. 21002, and JSPS Strategic Young Researcher Overseas Visits Program for Accelerating Brain Circulation No. R2201. This work is supported by U.S. Department of Energy contracts No. DE-AC05-84ER40150, No. DE-AC05-06OR23177, No. DE-FG02-99ER41065, No. DE-FG02-97ER41047, No. DE-AC02-06CH11357, No. DE-FG02-00ER41110, and No. DE-AC02-98CH10886, and US-NSF contracts No. 013815 and No. 0758095.

-
- [1] O. Hashimoto and H. Tamura, *Prog. Part. Nucl. Phys.* **57**, 564653 (2006).
- [2] A. Feliciello and T. Nagae, *Rep. Prog. Phys.* **78**, 096301 (2015).
- [3] *List of proposed experiments*, http://j-parc.jp/researcher/Hadron/en/Experiments_e.html
- [4] T.O. Yamamoto *et al.* (J-PARC E13 Collaboration), *Phys. Rev. Lett.* **115**, 222501 (2015).
- [5] T.R. Saito *et al.*, *Nucl. Phys. A* **881**, 218–227 (2012).
- [6] C. Rappold *et al.* (HypHI Collaboration), *Phys. Rev. C* **88**, 041001(R) (2013).
- [7] C. Rappold, *et al.*, *Phys. Lett. B* **747**, 129–134 (2015).
- [8] Y. Zhu (for the STAR Collaboration), *Nucl. Phys. A* **904–905**, 551c–554c (2013).
- [9] ALICE Collaboration (J. Adam *et al.*), *Phys. Lett. B* **754**, 360 (2016).
- [10] A. Esser, S. Nagao, F. Schulz, P. Achenbach *et al.* (A1 Collaboration), *Phys. Rev. Lett.* **114**, 232501 (2015).
- [11] F. Schulz, P. Achenbach *et al.* (A1 Collaboration), *Nucl. Phys. A* in Press, doi:10.1016/j.nuclphysa.2016.03.015 (2016).
- [12] T. Miyoshi *et al.* (HNSS Collaboration), *Phys. Rev. Lett.* **90**, 23 (2003).
- [13] L. Yuan *et al.* (HNSS Collaboration), *Phys. Rev. C* **73**, 044607 (2006).
- [14] M. Iodice *et al.* (Jefferson Lab Hall A Collaboration), *Phys. Rev. Lett.* **99**, 052501 (2007).
- [15] F. Cusanno *et al.* (Jefferson Lab Hall A Collaboration), *Phys. Rev. Lett.* **103**, 202501 (2009).
- [16] O. Hashimoto *et al.*, *Nucl. Phys. A* **835**, 121–128 (2010).
- [17] F. Cusanno *et al.*, *Nucl. Phys. A* **835**, 129–135 (2010).
- [18] S.N. Nakamura, A. Matsumura, Y. Okayasu, T. Seva, V.M. Rodriguez, P. Baturin *et al.* (HKS (JLab E01-011) Collaboration), *Phys. Rev. Lett.* **110**, 012502 (2013).
- [19] L. Tang, C. Chen, T. Gogami, D. Kawama, Y. Han *et al.* (HKS (JLab E05-115 and E01-011) Collaborations), *Phys. Rev. C* **90**, 034320 (2014).
- [20] G.M. Urciuoli *et al.* (Jefferson Lab Hall A Collaboration), *Phys. Rev. C* **91**, 034308 (2015).
- [21] T. Gogami, C. Chen, D. Kawama *et al.* (HKS (JLab E05-115) Collaboration), *Phys. Rev. C* **93**, 034314 (2016).
- [22] J.H.E. Mattauch, W. Thiele and A.H. Wapstra, *Nucl. Phys.* **67**, 1 (1965).
- [23] R.A. Brandenburg, S.A. Coon and P.U. Sauer, *Nucl. Phys. A* **294**, 3, 305 (1978).
- [24] D.H. Davis, *Nucl. Phys. A* **754**, 3c–13c (2005).
- [25] R.H. Dalitz and F. Von Hippel, *Phys. Lett.* **10**, 1 (1964).
- [26] J.L. Friar and B.F. Gibson, *Phys. Rev. C* **18**, 908 (1978).
- [27] A.R. Bodmer and Q.N. Usmani *Phys. Rev. C* **4**, 31 (1985).
- [28] M. Bedjidian *et al.*, *Phys. Lett. B* **62**, 467 (1976).
- [29] M. Bedjidian *et al.*, *Phys. Lett. B* **83**, 252 (1979).
- [30] A. Kawachi, Ph.D. thesis, University of Tokyo, Tokyo, Japan, 1997.
- [31] A. Gal, *Phys. Lett. B* **744**, 352–357 (2015).
- [32] D. Gazda and A. Gal, *Phys. Rev. Lett.* **116**, 122501 (2016).
- [33] D. Gazda and A. Gal, *Nucl. Phys. A* in Press, doi:10.1016/j.nuclphysa.2016.05.015 (2016).
- [34] A. Gal, E.V. Hungerford, and D.J. Millener, *arXiv* 1605.00557, to be published *Rev. Mod. Phys.* (2016).
- [35] T. Gogami, Ph.D. thesis, Tohoku University, Sendai, Japan, 2014.
- [36] M. Jurič *et al.*, *Nucl. Phys. B* **52**, 1 (1973).
- [37] H. Tamura *et al.*, *Phys. Rev. Lett.* **84**, 5963 (2000).
- [38] D.R. Tilley, C.M. Cheves, J.L. Godwin, G.M. Hale, H.M. Hofmann, J.H. Kelley, C.G. Sheu, H.R. Weller, *Nucl. Phys. A* **708**, 3 (2002).
- [39] E. Hiyama, Y. Yamamoto, T. Motoba, M. Kamimura, *Phys. Rev. C* **80**, 054321 (2009).
- [40] O. Richter, M. Sotona and J. Žofka, *Phys. Rev. C* **43**, 6 (1991).
- [41] E. Hiyama, M. Isaka, M. Kamimura, T. Myo, T. Motoba, *Phys. Rev. C* **91**, 054316 (2015).
- [42] J. Pniewski and M. Danysz, *Phys. Lett.* **1**, 142 (1962).
- [43] R.H. Dalitz and A. Gal, *Nucl. Phys. B* **1**, 1 (1967).
- [44] J. Pniewski and Z. Szymański, D.H. Davis, and J. Sacton, *Nucl. Phys. B* **2**, 317 (1967).
- [45] M. Sotona and S. Frullani, *Prog. Theor. Phys. Suppl.* **177**, 151 (1994).
- [46] Ed V. Hungerford, *Prog. Theor. Phys. Suppl.* **117**, 135 (1994).
- [47] G. Xu and Ed V. Hungerford, *Nucl. Instrum. Methods Phys. Res. A* **501**, 602–614 (2003).
- [48] T. Gogami *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **729**, 816–824 (2013).
- [49] Y. Fujii *et al.*, *Nucl. Instrum. Methods Phys. Res. A* **795**, 351–363 (2015).
- [50] G. Audi, A.H. Wapstra and C. Thibault, *Nucl. Phys. A* **729**, 337–676 (2003).
- [51] E. Hiyama and Y. Yamamoto, *Prog. Theor. Phys.* **128**, 1, 105 (2012).
- [52] E. Hiyama, *Nucl. Phys. A* **914**, 130 (2013).
- [53] X. Mougeot *et al.*, *Phys. Lett. B* **718**, 441 (2012).