



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Three-body model for the two-neutron emission of ^{16}Be

A. E. Lovell, F. M. Nunes, and I. J. Thompson

Phys. Rev. C **95**, 034605 — Published 10 March 2017

DOI: [10.1103/PhysRevC.95.034605](https://doi.org/10.1103/PhysRevC.95.034605)

Three-body model for the two-neutron emission of ^{16}Be

A. E. Lovell,^{1,2} F. M. Nunes,^{1,2} and I. J. Thompson³

¹*National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, MI 48824, USA*

²*Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA*

³*Lawrence Livermore National Laboratory, L-414, Livermore, CA 94551, USA*

(Dated: February 17, 2017)

Background: While diproton emission was first theorized in 1960 and first measured in 2002, it was first observed only in 2012. The measurement of ^{14}Be in coincidence with two neutrons suggests that ^{16}Be does decay through the simultaneous emission of two strongly correlated neutrons.

Purpose: In this work, we construct a full three-body model of ^{16}Be (as $^{14}\text{Be} + n + n$) in order to investigate its configuration in the continuum and in particular, the structure of its ground state.

Method: In order to describe the three-body system, effective n - ^{14}Be potentials were constructed, constrained by the experimental information on ^{15}Be . The hyperspherical R-matrix method was used to solve the three-body scattering problem, and the resonance energy of ^{16}Be was extracted from a phase-shift analysis.

Results: In order to reproduce the experimental resonance energy of ^{16}Be within this three-body model, a three-body interaction was needed. For extracting the width of the ground state of ^{16}Be , we use the full width at half maximum of the derivative of the three-body eigenphase shifts and the width of the three-body elastic scattering cross section.

Conclusions: Our results confirm a dineutron structure for ^{16}Be , dependent on the internal structure of the subsystem ^{15}Be .

PACS numbers: 24.10.Eq, 24.30.Gd

Keywords: dineutron, two-neutron decay, hyperspherical harmonics, R-matrix, rare isotopes

I. INTRODUCTION

Exotic nuclei are found across the nuclear chart. Proton and neutron halos are found near the proton and neutron dripline, respectively, not only in the lightest mass nuclei but also possibly in nuclei as heavy as neon [1]. Two-nucleon halo systems can be Borromean, where, if we think of these nuclei in terms of a core plus two neutrons or protons, the three-body system is bound but each of the two-body subsystems is unbound [2] (Ch. 9). Unsurprisingly, beyond the dripline, novel structures can give rise to exotic decay paths.

Two-proton decay was first theorized in 1960 [3]. When two nucleons decay from a nucleus A , there are three possible mechanisms. First, A can decay through the simultaneous emission of the two valence nucleons, in a true three-body decay. If there is a state in the $A-1$ nucleus below the ground state of the parent nucleus, the two nucleons are likely to be emitted sequentially from A , stepping through the intermediate $A-1$ state. However, if the ground state in the $A-1$ nucleus is energetically inaccessible to the emission of one nucleon and there is correlation between the two nucleons before the decay, dinucleon emission is the likely alternative.

Because of the Coulomb interaction, the diproton phenomena is extremely hard to observe; the two protons are repelled from one another as soon as they exit the nucleus, making it difficult to reconstruct the diproton angular correlations that were present in the parent nucleus. Nevertheless, it has been observed in many nuclei. The dineutron decay, on the other hand, poses its own challenges. The neutron dripline is harder to reach than the proton dripline, and the statistics for neutron-rich nuclear decays beyond the neutron dripline, involving two-

neutron coincidences, are very low. In both, dineutron and diproton decay, the differentiation between a correlated decay and an uncorrelated three-body decay is made based on model considerations and therefore is not free from ambiguity.

Two-proton emission from the ground state of a nucleus was experimentally observed for the first time in ^{45}Fe [4, 5], over forty years after the initial prediction. Since then, many examples of two-proton emission have been seen from ground states [6–8], as well as from excited states [9]. Because the relevant degrees of freedom are those related to the emission of the two protons from the parent nucleus, three-body models have been used to theoretically describe these decays. Different structural configurations of the parent nucleus give rise to different values for its width and half-life, as well as different ways of sharing the energy between the three particles. It is only through the comparison of model calculations to the data that insights into the nature of the decay can be obtained [10, 11].

In comparison to the large number of two-proton emitters that have been studied experimentally and theoretically, two-neutron emitters have not been as well investigated. In one of the first theoretical studies of two-neutron emission, Grigorenko [12] discussed the existence of one-, two-, and four-neutron emitters, as well as comparisons of their widths in a three-body framework. Recently, a few cases of two-neutron emission have been observed [13–15]. The first of these was observed in a 2012 experiment at the National Superconducting Cyclotron Laboratory [13] through the decay of ^{16}Be to ^{14}Be plus two neutrons. As the ground state energy of ^{16}Be was found to be 1.35 MeV (with a width of 0.8 MeV) and a lower limit of 1.54 MeV had previously been placed on

the ground state of ^{15}Be [16], ^{16}Be is an ideal candidate for simultaneous two-neutron emission. Depending on the width of the ground state of ^{15}Be , sequential neutron decay from ^{16}Be to ^{14}Be could be energetically inaccessible. A later experiment [17] determined that the lowest state in ^{15}Be is an $l = 2$ state at 1.8 MeV with a width of 575 ± 200 keV.

Although comparisons of the ^{16}Be data in [13] to dineutron, sequential, and three-body decay models showed the data best matched the dineutron emission, there was some controversy over this finding [18, 19]. Extreme models were used to show the difference between dineutron emission and a three-body decay. The dineutron was modeled as a cluster and the decay as a two-body $^{16}\text{Be} \rightarrow ^{14}\text{Be} + 2n$, in an s-wave relative motion. The three-body breakup corresponded to phase space only. A more realistic, full three-body model ($^{14}\text{Be} + n + n$) is necessary to help clarify the mode of decay of this exotic nucleus. Several three-body models have been successfully used to describe the continuum states of ^{26}O [20, 21] but no application to ^{16}Be is thus far available. This is the goal of the present study.

This paper is organized into the following sections. In Section II, we introduce the three-body hyperspherical R-matrix theory used in this work. In Section III, details about the two- and three-body potentials are presented, as well as a convergence study of our calculations. Our results, assuming either a $1d_{5/2}$ or a $2s_{1/2}$ ground state for ^{15}Be , are discussed in Section IV, and in Section V, we discuss the consequences of these models. Finally, we conclude in Section VI.

II. THEORETICAL FRAMEWORK

In this work, the ^{16}Be system is assumed to take the form of *core* + n + n and therefore should satisfy the three-body Schrödinger equation:

$$(T_r + T_s + V_{cn_1} + V_{cn_2} + V_{nn} + V_{3b})\Psi = E_{3B}\Psi, \quad (1)$$

where \vec{r} and \vec{s} are the standard Jacobi coordinates, as shown in Figure 1, where r is the distance between two of the bodies, and s is the distance between the third body and the center of mass of the first two. V_{cn_i} and V_{nn} are the pairwise interactions. Typically, when the degrees of freedom in the core are frozen, the final three-body system becomes under-bound. Traditionally, three-body interactions are then introduced to take into account the additional binding needed to reproduce the experimental ground state. This is the role of V_{3b} in Eqn. (1).

Eqn. (1) is a 6-dimensional equation, where the coordinates \vec{r} and \vec{s} do not separate due to the fact that the pairwise interactions depend on both. The hyperspherical harmonic method makes a particular choice of coordinates and basis functions such that this three-body Schrödinger equation becomes a set of 1-dimensional coupled hyper-radial equations. This is briefly described here.




FIG. 1: Three Jacobi coordinate systems, a) Jacobi X system, b) Jacobi Y system, and c) Jacobi T system. Because the two neutrons are identical, the X and Y coordinate systems are identical.

A. Hyperspherical harmonic method

For a three-body system, there are three sets of Jacobi coordinates that can be defined, Figure 1. We will use i to denote one of the three Jacobi systems, X, Y, or T. We assume the T coordinate system for convenience ($i = 3$, which we omit through the rest of this work). Now, \vec{x} and \vec{y} are the scaled Jacobi coordinates [2] (Ch. 9), defined by

$$\vec{x} = \frac{\vec{r}}{\sqrt{2}} \quad (2)$$

and

$$\vec{y} = \sqrt{\frac{2A_3}{A_3 + 2}} \vec{s}, \quad (3)$$

where A_3 is the mass number of the core. From here, we can define the hyperspherical coordinates

$$\rho^2 = x^2 + y^2, \quad (4)$$

and

$$\tan\theta = \frac{x}{y}. \quad (5)$$

Note that ρ is invariant among the three Jacobi coordinate systems, but θ depends on i . Using these coordinates, the kinetic energy operator can be written as:

$$T = -\frac{\hbar^2}{2m} \left[\frac{1}{\rho^5} \frac{\partial}{\partial \rho} \left(\rho^5 \frac{\partial}{\partial \rho} \right) + \frac{1}{\rho^2 \sin^2 2\theta} \frac{\partial}{\partial \theta} \left(\sin^2 2\theta \frac{\partial}{\partial \theta} \right) - \frac{L_x^2}{\rho^2 \sin^2 \theta} - \frac{L_y^2}{\rho^2 \cos^2 \theta} \right], \quad (6)$$

where m is the unit mass, here $m = 938.0$ MeV/ c^2 .

We perform the standard partial wave decomposition of the wavefunction,

$$\Psi^{JM} = \sum_{l_x l_y l_s j I} \psi_{l_x l_y}^{l_s j I J}(x, y) \left\{ ([Y_{l_x} \otimes Y_{l_y}]_l \otimes [X_{\sigma_1} \otimes X_{\sigma_2}]_s)_j \otimes \phi_I \right\}_{JM}, \quad (7)$$

where l is the total orbital angular momentum, l_x is the relative orbital angular momentum in the $2n$ system, l_y is the relative orbital angular momentum in the $core + (2n)$ system, I is the spin of the core, S is the total spin of the two neutrons, and j is the total angular momentum of the two neutrons relative to the core. Next we expand the part dependent on (x, y) in hyperspherical functions $\varphi_K^{l_x l_y}(\theta)$,

$$\psi_{l_x l_y}^{lSjIJ}(x, y) = \rho^{-5/2} \sum_{K=0}^{K_{max}} \chi_{K l_x l_y}^{lSjIJ}(\rho) \varphi_K^{l_x l_y}(\theta) \quad (8)$$

where $\varphi_K^{l_x l_y}(\theta)$ is set to an eigenfunction of the angular operator in Eqn. (6) with eigenvalue $K(K+4)$. Its explicit form is:

$$\varphi_K^{l_x l_y}(\theta) = N_K^{l_x l_y} (\sin\theta)^{l_x} (\cos\theta)^{l_y} P_n^{l_x+1/2, l_y+1/2}(\cos 2\theta), \quad (9)$$

where $P_n^{l_x+1/2, l_y+1/2}(\cos 2\theta)$ are the Jacobi Polynomials, and $N_K^{l_x l_y}$ is a normalization factor resulting from the condition:

$$\int_0^{\pi/2} \varphi_K^{l_x l_y}(\theta) \varphi_{K'}^{l_x l_y}(\theta) \sin^2 \theta \cos^2 \theta d\theta = \delta_{KK'}. \quad (10)$$

For compactness, we introduce the hyperspherical harmonic functions,

$$\mathcal{Y}_\gamma^{JM}(\Omega_5, \sigma_1, \sigma_2, \boldsymbol{\xi}) = \varphi_K^{l_x l_y}(\theta) \left\{ ([Y_{l_x} \otimes Y_{l_y}]_l \otimes [X_{\sigma_1} \otimes X_{\sigma_2}]_S)_j \otimes \phi_I \right\}_{JM}, \quad (11)$$

with γ representing the set $\{KlSlJl_x l_y\}$, so that the total wave function can be written in the form:

$$\Psi^{JM} = \rho^{-5/2} \sum \chi_\gamma^J(\rho) \mathcal{Y}_\gamma^{JM}(\Omega_5, \sigma_1, \sigma_2, \boldsymbol{\xi}). \quad (12)$$

In this work, we focus on $(J, M) = (0, 0)$, corresponding to the spin of the ^{16}Be ground state.

Substituting Eqn.(12) into Eqn. (1), we are left with the following set of coupled hyper-radial equations:

$$\left(-\frac{\hbar^2}{2m} \left[\frac{d^2}{d\rho^2} - \frac{(K+3/2)(K+5/2)}{\rho^2} \right] - E_{3B} \right) + \sum_{\gamma'} V_{\gamma\gamma'}(\rho) \chi_{\gamma'}^J(\rho) = 0, \quad (13)$$

where the coupling potentials are defined as

$$V_{\gamma\gamma'}(\rho) = \langle \mathcal{Y}_{\gamma'}^{JM}(\Omega_5, \sigma_1, \sigma_2, \boldsymbol{\xi}) | \sum_{j>i=1}^3 V_{ij} | \mathcal{Y}_\gamma^{JM}(\Omega_5, \sigma_1, \sigma_2, \boldsymbol{\xi}) \rangle \quad (14)$$

Eqn. (13) must be solved under the condition that the wavefunction is regular at the origin and behaves asymptotically as,

$$\chi_{\gamma\gamma_i}^J \rightarrow \frac{i}{2} \left[\delta_{\gamma\gamma_i} H_{K+3/2}^-(\kappa\rho) - \mathbf{S}_{\gamma\gamma_i}^J H_{K+3/2}^+(\kappa\rho) \right], \quad (15)$$

when $\rho \rightarrow \infty$, where the γ_i are the components of a plane wave.

It is important also to note that the final wave function will have to be summed over γ_i , as we do not assume a specific incoming wave for our ^{16}Be system.

B. Hyperspherical R-matrix method

The set of coupled hyper-radial equations could, in principle, be solved by direct numerical integration. However, at low scattering energies, the centrifugal barrier - $(K+3/2)(K+5/2)$ - found in every channel, including $K=0$, would likely cause this method to develop numerical inaccuracies. Instead, we use the hyperspherical R-matrix method [2] (Ch. 6).

In the hyperspherical R-matrix method, we first create a basis, w_γ^n , by solving the uncoupled equations, corresponding to Eqn. (13) with all couplings set to zero except for the diagonal, in a box of size ρ_{max} ,

$$[T_\gamma(\rho) + V_{\gamma\gamma}(\rho) - \varepsilon_{n\gamma}] w_\gamma^n(\rho) = 0. \quad (16)$$

By enforcing all logarithmic derivatives,

$$\beta = \frac{d \ln(w_\gamma^n(\rho))}{d\rho}, \quad (17)$$

to be equal for $\rho = \rho_{max}$, the set of functions, w_γ^n , form a complete, orthogonal basis within the box. Then, the scattering equation inside the box can be solved by expanding in this R-matrix basis:

$$g_\gamma^p(\rho) = \sum_{n=1}^N c_\gamma^{pn} w_\gamma^n(\rho). \quad (18)$$

The corresponding coupled channel equations are:

$$[T_\gamma(\rho) + V_{\gamma\gamma}(\rho)] g_\gamma^p(\rho) + \sum_{\gamma' \neq \gamma} V_{\gamma\gamma'}(\rho) g_{\gamma'}^p(\rho) = e_p g_\gamma^p(\rho). \quad (19)$$

To find the coefficients c_γ^{np} , we insert Eqn. (18) into Eqn. (19), multiply the resulting equation by $w_{\gamma'}^{n'}$ and integrate over the box size. This results in a matrix equation:

$$\varepsilon_{n\gamma} c_\gamma^{np} + \sum_{\gamma' \neq \gamma} \sum_{n'} \langle w_\gamma^n(\rho) | V_{\gamma\gamma'}(\rho) | w_{\gamma'}^{n'}(\rho) \rangle = e_p c_\gamma^{pn}, \quad (20)$$

which, when solved, provides the coefficients c_γ^{np} of the expansion Eqn. (18). Since $g_\gamma^p(\rho)$ are only complete inside the box, and do not have the correct normalization, the full three-body scattering wavefunction is given by a superposition of these solutions which is then matched to the correct asymptotic form:

$$\chi_{\gamma\gamma_i}^J(\rho) = \sum_{p=1}^P A_{\gamma_i}^p g_\gamma^p(\rho). \quad (21)$$

The new expansion parameter p corresponds to the number of poles considered in the R-matrix. The normalization coefficients, $A_{\gamma_i}^p$, connect the inside wavefunction with the asymptotic behavior of Eqn. (15). The explicit relation is [2] (Ch. 6),

$$A_{\gamma_i}^p = \frac{\hbar^2}{2m} \frac{1}{e_p - E} \sum_{\gamma'} g_{\gamma'}^p(\rho_{max}) \left[\delta_{\gamma\gamma'} \left(H_{K+3/2}^{-'}(\kappa_{\gamma'} \rho_{max}) - \beta H_{K+3/2}^{-}(\kappa_{\gamma'} \rho_{max}) \right) - \mathbf{S}_{\gamma'\gamma_i} \left(H_{K+3/2}^{+'}(\kappa_{\gamma'} \rho_{max}) - \beta H_{K+3/2}^{+}(\kappa_{\gamma'} \rho_{max}) \right) \right].$$

From the values of the $g_{\gamma}^p(\rho)$ function at the surface, one can determine the R-matrix [2] (Ch. 6),

$$R_{\gamma\gamma'} = \frac{\hbar^2}{2m\rho_{max}} \sum_{p=1}^P \frac{g_{\gamma}^p(\rho_{max})g_{\gamma'}^p(\rho_{max})}{e_p - E_{3B}}. \quad (22)$$

Once the R-matrix is obtained, the S-matrix can be directly computed:

$$\mathbf{S} = [\mathbf{H}^+ - \rho_{max}\mathbf{R}(\mathbf{H}^{+'} - \beta\mathbf{H}^+)]^{-1} * [\mathbf{H}^- - \rho_{max}\mathbf{R}(\mathbf{H}^{-'} - \beta\mathbf{H}^-)] \quad (23)$$

along with the phase shifts for each channel, from the diagonal elements of the S-matrix, $S_{\gamma\gamma} = e^{2i\delta_{\gamma\gamma}}$ (more details in [2]). However, the S-matrix is not necessarily diagonal, and due to all of the off-diagonal terms, the diagonal does not have special significance. Instead of using only $S_{\gamma\gamma}$ directly, it is common to diagonalize the S-matrix and extract the eigenphases, δ^e [22]. The resonance energies and widths can then be extracted from these eigenphases.

C. Width calculation

If one assumes a Breit-Wigner shape, resonant properties for a single-channel calculation can be directly extracted from the phase shift through the relation:

$$\tan\delta = \frac{\Gamma/2}{E_{3B} - E_{res}}, \quad (24)$$

where Γ is the width of the resonance and E_{res} is the resonance energy. If this is valid, the width can be computed as the full width at half maximum (FWHM) from the energy derivative of the phase shift, $\Gamma = 2/(\partial\delta/\partial E_{3B})$. In the case of multiple channels with weak coupling, one can add the various partial widths to obtain the total width of the three-body resonance. For this potentially strongly coupled three-body problem, we do not expect the pure Breit-Wigner approach to be valid. Nevertheless, for completeness, we do try to identify channels for which such an approach may be applicable.

We can also construct the total three-body cross section as a function of energy which would be measured

during an elastic scattering experiment, as defined in [22],

$$\sigma_{3:3}^J(E_{3B}) \propto \frac{1}{4\kappa^5} \left[\sum_{\gamma} |1 - S_{\gamma\gamma}(E_{3B})|^2 + \sum_{\gamma\gamma'} |S_{\gamma\gamma'}(E_{3B})|^2 \right] \quad (25)$$

in which we might see a resonance as a peak. Theoretically, three-body elastic scattering could be measured if the ^{14}Be and two neutrons could be impinged upon one another simultaneously with total energy, E_{3B} . Because this quantity includes a sum over all K , this method can justify that the resonance energies extracted from a single phase shift do indeed represent the resonance energy of the total system, as all channels are included in this calculation.

III. NUMERICAL DETAILS

A. Input interactions versus data

In the three-body model, each of the two-body interactions must be constrained, typically from experimental data. However, very little is known about ^{15}Be [17], so shell model calculations are used to supplement the available data. Shell model calculations for ^{15}Be were provided [23] using the WBP interaction [24]. Since the ground state in the shell model calculation was an $l = 2$ state and was 1 MeV higher than the experimentally observed $l = 2$ state in ^{15}Be [17], the levels that were used to constrain the ^{14}Be -n interactions were the shell model levels lowered by 1 MeV, shown in Figure 2.




FIG. 2: Level scheme for ^{15}Be . The first column shows the shell model calculation provided by [23], while the second column shows the ^{15}Be levels that we used in this work; here, the shell model levels are lower by 1 MeV so the $1d_{5/2}$ state in the shell model calculation reproduces the experimental $l = 2$ energy from [17], as shown in the third column.

The ^{14}Be -n interaction for each partial wave has a Woods-Saxon shape with $a = 0.65$ fm and $R = 1.2A^{1/3}$ fm, where A is the mass number of the ^{14}Be core. The depths depend on angular momentum, and are obtained by fitting the single-particle resonances in ^{15}Be , described

in Fig. 2, using the code POLER [25]. The core deformation is taken into account by allowing an l -dependence in the potential. A spin-orbit interaction was also included with the same geometry as the central nuclear force with the depth adjusted to reproduce the split between the $1d_{5/2}$ and $1d_{3/2}$ states. We use the definition of the spin-orbit strengths of FaCE [26]. Potential depths for the various models included are as indicated in Table I.

The lowest s- and p-orbitals in ^{14}Be are assumed to be full. In order to remove the effect of these occupied states in the ^{14}Be core, the $1s_{1/2}$, $1p_{3/2}$, and $1p_{1/2}$ states were projected out through a supersymmetric transformation [26].

B. Description of models

There are four three-body models for ^{16}Be that we consider in this work. In **D3B**, the ground state of ^{15}Be is a $1d_{5/2}$ state and a three-body force is included to reproduce the experimental three-body ground state energy of ^{16}Be . This three-body force is also of Woods-Saxon form with radius of 3.02 fm and diffuseness of 0.65 fm. In **D**, the ground state of ^{15}Be is a $1d_{5/2}$ state but no three-body force is included. In **S**, the ground state of ^{15}Be is a $2s_{1/2}$ state but no three-body force is included.

All models D3B, D, and S include the GPT NN interaction [27], as in previous three-body studies [28–31]. This interaction reproduces NN observables up to 300 MeV. Although it is simpler than the AV18 [32] and Reid soft-core [33] interactions, its range is more than suitable for the energy scales used in this work. We also consider the effects of removing the NN interaction completely. This model is named **DNN**. In Table I, we provide the depths for the various terms of the interaction and the coefficient α_{NN} by which we multiply the GPT force in each of our calculations.

In Table II, we summarize the energies for the $1d_{5/2}$ and $2s_{1/2}$ states in the subsystem ^{15}Be as well as the ground state energy of ^{16}Be in the various models considered in Table I. For all of the models considered, the $1d_{3/2}$ state was placed at 6.0 MeV.

Parameter	D3B	D	DNN	S
V_s	-26.182	-26.182	-26.182	-41.182
V_p	-30.500	-30.500	-30.500	30.500
V_d	-42.73	-42.730	-42.730	-42.730
$V_{so} (l \neq 2)$	-10.000	-10.000	-10.00	-10.000
$V_{so} (l=2)$	-33.770	-33.770	-33.770	-33.770
V_{3B}	-7.190	0.000	-7.190	0.000
α_{NN}	1.000	1.000	0.000	1.000

TABLE I: Interaction parameters for the various models considered. All depths are given in MeV. Details in the text.

	D3B	D	DNN	S
$^{15}\text{Be}(1d_{5/2})$	1.80	1.80	1.80	1.80
$^{15}\text{Be}(2s_{1/2})$	~ 3	~ 3	~ 3	0.48
$^{16}\text{Be}(gs)$	1.32	1.88	3.08	1.60*

TABLE II: Energy levels, in MeV, for ^{16}Be and ^{15}Be for the various models considered. Energies are measured with respect to the ^{14}Be threshold. Details in the text. *This is an excited state.

C. Convergence

Our methods rely on basis expansions, and our model space is determined by a number of numerical parameters. In this section we demonstrate convergence for various quantities, including the ground state energy of ^{16}Be and the phase shifts. The truncation of the expansion in hyperspherical harmonics is controlled by the hyper-momentum, K . In Figure 3, we show the convergence of the lowest 0^+ three-body resonance energy of ^{16}Be as K_{max} increases. The width of ^{16}Be with respect to K_{max} shows the same trend. Our results are converged within 0.05 MeV by $K_{max} = 28$ for both observables.

Our results are very sensitive to the the number of R-matrix basis functions N (which essentially determines the hyper-radial discretization) as well as the maximum box size, ρ_{max} . In Tables III and IV, we show the convergence of the three-body resonance energy of several parameters for the $K = 0, L = 0, S = 0$ channel. The convergence with respect to the number of R-matrix basis functions is shown in Table III. Convergence is slow but results are very close to converged for $N = 95$.

We also needed to check the dependence on the box size, ρ_{max} . When increasing the box size, one also needs to increase the number of R-matrix basis functions that span the radial space for consistency. These results are given in Table IV. We summarize the minimum convergence requirements in Table V.

N	E_{3B} (MeV)
70	2.06
75	1.95
80	1.84
85	1.78
90	1.74
95	1.71
100	1.69
105	1.67

TABLE III: Convergence of E_{3B} as a function of the number of radial R-matrix functions, N , for $\rho_{max} = 60$ fm.




FIG. 3: Convergence of the three-body energy as a function of the maximum K value included in the model space.

ρ_{max} (fm)	N	E_{3B} (MeV)
50	80	1.70
60	95	1.71
70	110	1.72

TABLE IV: As the box size, ρ_{max} , increases, a greater number of R-matrix radial functions, N , are need to keep the same resonance energy, E_{3B} .

IV. RESULTS

Using model D, we calculated the eigenphases for ^{16}Be . The converged eigenphases can be found in Figure 4 (solid). As we would expect for this type of system, the resonance energy in model D is above the experimental energy observed for the ground state. We include a three-body force, as described in Table I. The eigenphases, including this three-body interaction (model D3B), are shown in Figure 4 (dashed).

One can also extract a resonance energy from the three-body total cross section, shown in Figure 5 as a function of three-body energy. This observable also contains contributions from all of the channels included in the model space. If one investigates the structure of the wavefunction of model D3B for the pole closest to the resonance energy, we conclude that the state is 37% $K = 0$, $l_x = l_y = 0$, 30% $K = 2$, $l_x = l_y = 0$, and 13% $K = 4$, $l_x = l_y = 0$.

Although the lowest experimentally observed state in ^{15}Be was an $l = 2$ state, we wanted to investigate the possibility of an s-wave ground state in ^{15}Be , below the observed state. Such a state exists in ^{10}Li and was only observed after other higher lying resonances were well known [34]. With this in mind, we developed model S, described in Table I. We use the same model space as in

Parameter	Value
K_{max}	28
$l_x(max), l_y(max)$	10
N_{Jac}	65
ρ_{max} (fm)	60
N	95

TABLE V: Minimum convergence values for the three-body wave function expansion.




FIG. 4: Eigenphases as a function of three-body energy for ^{16}Be models D (solid, black), D3B (dashed, red), DNN (dotted, green), and S (double-dash dotted, blue).

Table V. The dot-dashed line in Figure 4 shows the corresponding eigenphases for model S. The resulting cross section is also depicted in Figure 5 by the dot-dashed line. The clear evidence for the resonance seen in models D and D3B, is washed out in model S; however, there does appear to be a resonant-like shape in the eigenphase, just above the resonance in D3B at 1.60 MeV. We will come back to this in Section V.

Finally we also consider the results when the NN interaction is switched off (model DNN). A resonance is still seen in $\delta^e(E)$ (Figure 4, dotted), around 3 MeV, demonstrating the importance of the NN correlations in producing the observed state in ^{16}Be . Our results show that the configuration of the system is strongly modified by switching off the NN interaction.

V. DISCUSSION

To investigate whether or not dineutron emission is the predominant form of decay for ^{16}Be , we can study the structure of the ^{16}Be through the spatial density distribution. Two spatially correlated neutrons should be




FIG. 5: Three-body cross section as a function of three-body energy for ^{16}Be models D(solid, black), D3B (dashed, red), DNN (dotted, green), S (double-dash dotted, blue).

primarily emitted via dineutron emission - rather than a three-body decay.

In calculating the spatial probability distribution of the three-body system,

$$P(x, y) = \int |\Psi^{JM}(\mathbf{x}, \mathbf{y})|^2 d\Omega_x d\Omega_y \quad (26)$$

we can determine the location of the two neutrons with respect to the core. The wave function used here is that of Eqn. 7, calculated at the resonance energy, $E = 1.32$ MeV. The calculation includes the wave function up to ρ_{max} , and no additional binning is included to localize the resonance. Note, also, that $P(x, y)$ contains the contribution of all components, and not just a single K . From this density distribution we can determine the configuration of the two neutrons in ^{16}Be - dineutron, helicopter, or triangle (Figure 6 a, b, and c, respectively).




FIG. 6: Three-body configurations, a) dineutron (two neutrons close together and far from the core), b) helicopter (two neutrons are close to the core and far from each other), and c) three-body (the three bodies are equally spaced).

Figure 7 shows the resulting density distribution for the ^{16}Be system with the D3B model. The density distribution mainly shows a dineutron configuration, although a small component of a helicopter configuration is present. This is consistent with what was seen in [13]. Even though the three-body resonance energy shifts by about 0.5 MeV when the three-body interaction is removed, this does not change the relative strength of the dineutron component of the density distribution.




FIG. 7: Three-body density as a function of the distance between the two neutrons (r) and the distance between the nn pair and the core (s) for D3B. The scale on the right is given in fm^{-5} .

There are several quantities that we can look at to extract a width for this system. If we extract the width from the FWHM of the derivative of the three-body eigenphase shift in Figure 4 we obtain 0.17 MeV (consistent with the observed width of the nearest R-matrix pole, 0.17 MeV). Both of these are smaller than the 0.8 MeV width found by experiment [13]. This discrepancy is most likely due to the effect of experimental resolution (etc.), which has not been taken into account when comparing our calculations with experiment. Work to include these effects is currently ongoing. To further calculate the yield of ^{16}Be when produced from a proton removal of ^{17}B , as in [13], a reaction model would be needed that includes the overlap functions between the original ^{17}B and the resulting three-body continuum states in ^{16}Be .

When we switch off the NN interaction (model DNN), the density distribution shown in Figure 8 has equal contributions from the dineutron and the helicopter configurations. Increasing or decreasing the strength of the three-body interaction does not change this picture. This illustrates that it is indeed the NN interaction that is responsible for the strong dineutron character of the ^{16}Be ground state.

For comparison with all of these models, Figure 9 shows the density distribution for a ^{16}Be that has both the NN and $n\text{-}^{15}\text{Be}$ interactions removed (all K components have been summed). This system does not contain




FIG. 8: Same as Fig. 7 for the model DNN.

any resonance, so the density distribution is calculated at the ^{16}Be experimental resonance energy, 1.35 MeV. The distribution has less structure and is pushed farther away from the center of the system.




FIG. 9: Same as Fig. 7 for a plane wave solution of ^{16}Be , at $E_{3B} = 1.35$ MeV, for comparison.

Let us now turn our attention back to the hypothesis of there being a lower s-wave resonance in ^{15}Be (model S). Although $\delta^e(E)$ goes through 90° , there is no clear signature of a resonance in the total cross section. Indeed the ^{16}Be system becomes bound, also indicated by the double hump structure 4 (double-dash dotted). Only by using a much shallower s-wave potential could we regain a resonance in the low energy ^{16}Be spectrum. These results make it much less likely that ^{15}Be has an s-wave ground state.

Using three-neutron coincidences, Kuchera, et. al. proposed that there is a small chance of finding the $1d_{3/2}$ state in ^{15}Be at 2.69 MeV [35]. Including this state, keeping the $1d_{5/2}$ at 1.8 MeV, and using the same s-wave

as models D and D3B, the ground state energy of ^{16}Be produced by our model had $E_{3B} = 1.05$ MeV, without including a three-body interaction. The density distribution was nearly identical to that shown in Figure 7. In this case, the only way to reproduce the experimental ground state of ^{16}Be would be to include a repulsive three-body interaction, which is unusual.

VI. CONCLUSIONS

In summary, a three-body model for ^{16}Be was developed to investigate the properties of the system in the continuum. The hyperspherical R-matrix method was used to solve the three-body scattering problem, with the n- ^{14}Be interactions constrained by experimental data on ^{15}Be . As usual in three-body models, we included a three-body potential to reproduce the experimental ground state energy of ^{16}Be . We obtained convergence results for phase shifts, density distributions, and three-body cross sections.

We study the properties of the resulting three-body continuum around the resonant energy of ^{16}Be and conclude that it has a strong dineutron configuration, consistent with experimental observations [13]. The estimate of the width obtained from our calculations is consistent among the various methods of extraction but is smaller than the experimental value [13]. We find that the NN interaction is important in producing the strong dineutron configuration in the ground state of ^{16}Be , since the structure of the resonance is completely different when switching off the NN interaction. In contrast, the three-body force needed to shift the resonance energy to the observed experimental energy of ^{16}Be ground state, has little effect on the structure of the state. We also explore a possible s-wave ground state in ^{15}Be and find that the results are incompatible with the observed ^{16}Be ground state [13]. In fact, the structure of the ^{15}Be ground state being a $d_{5/2}$ -wave is crucial to reproducing the ^{16}Be experimental results. Only with a $1d_{5/2}$ ground state and higher lying $2s_{1/2}$ and $1d_{3/2}$ state in ^{15}Be can a resonance energy of 1.35 MeV be reproduced in ^{16}Be with a physical three-body interaction.

The ^{16}Be experiment [13] provided a variety of correlation observables that would be very interesting to compare to our model. However, our predictions need to be introduced into a full experimental simulation code that includes the appropriate three-body assumptions as well as efficiencies and acceptance of the detector setup. Work along these lines is currently underway.

ACKNOWLEDGMENTS

The authors would like to thank Artemis Spyrou, Michael Thoennessen, Simin Wang, and Anthony Kuchera for useful discussions. The authors would also like to acknowledge iCER and the High Performance Computing Center at Michigan State University for their

computational resources. This work was supported by the Stewardship Science Graduate Fellowship program under Grant No. DE-NA0002135 and by the National Science Foundation under Grant. No. PHY-1403906 and PHY-1520929. This work was performed under the aus-

pices of the U.S. Department of Energy through NNSA contract DE-FG52-08NA28552 and by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

-
- [1] I. Tanihata, H. Savajols, and R. Kanungo, *Progress in Particle and Nuclear Physics* **68**, 215 (2013), ISSN 0146-6410, URL <http://www.sciencedirect.com/science/article/pii/S0146641012001081>.
- [2] I. J. Thompson and F. M. Nunes, *Nuclear Reactions for Astrophysics* (Cambridge University Press, 2009).
- [3] V. Goldansky, *Nuclear Physics* **19**, 482 (1960), ISSN 0029-5582, URL <http://www.sciencedirect.com/science/article/pii/0029558260902583>.
- [4] J. Giovinazzo, B. Blank, M. Chartier, S. Czajkowski, A. Fleury, M. J. Lopez Jimenez, M. S. Pravikoff, J.-C. Thomas, F. de Oliveira Santos, M. Lewitowicz, et al., *Phys. Rev. Lett.* **89**, 102501 (2002), URL <http://link.aps.org/doi/10.1103/PhysRevLett.89.102501>.
- [5] M. Pftzner, E. Badura, C. Bingham, B. Blank, M. Chartier, H. Geissel, J. Giovinazzo, L.V. Grigorenko, R. Grzywacz, M. Hellstrm, et al., *Eur. Phys. J. A* **14**, 279 (2002), URL <http://dx.doi.org/10.1140/epja/i2002-10033-9>.
- [6] B. Blank, A. Bey, G. Canchel, C. Dossat, A. Fleury, J. Giovinazzo, I. Matea, N. Adimi, F. De Oliveira, I. Stefan, et al., *Phys. Rev. Lett.* **94**, 232501 (2005), URL <http://link.aps.org/doi/10.1103/PhysRevLett.94.232501>.
- [7] C. Dossat, A. Bey, B. Blank, G. Canchel, A. Fleury, J. Giovinazzo, I. Matea, F. d. O. Santos, G. Georgiev, S. Grévy, et al., *Phys. Rev. C* **72**, 054315 (2005), URL <http://link.aps.org/doi/10.1103/PhysRevC.72.054315>.
- [8] I. Mukha, K. Stümmerer, L. Acosta, M. A. G. Alvarez, E. Casarejos, A. Chatillon, D. Cortina-Gil, J. Espino, A. Fomichev, J. E. García-Ramos, et al., *Phys. Rev. Lett.* **99**, 182501 (2007), URL <http://link.aps.org/doi/10.1103/PhysRevLett.99.182501>.
- [9] J. Gómez del Campo, A. Galindo-Uribarri, J. R. Beene, C. J. Gross, J. F. Liang, M. L. Halbert, D. W. Stracener, D. Shapira, R. L. Varner, E. Chavez-Lomeli, et al., *Phys. Rev. Lett.* **86**, 43 (2001), URL <http://link.aps.org/doi/10.1103/PhysRevLett.86.43>.
- [10] L. V. Grigorenko, R. C. Johnson, I. G. Mukha, I. J. Thompson, and M. V. Zhukov, *Phys. Rev. C* **64**, 054002 (2001), URL <http://link.aps.org/doi/10.1103/PhysRevC.64.054002>.
- [11] L. V. Grigorenko and M. V. Zhukov, *Phys. Rev. C* **68**, 054005 (2003), URL <http://link.aps.org/doi/10.1103/PhysRevC.68.054005>.
- [12] L. V. Grigorenko, I. G. Mukha, C. Scheidenberger, and M. V. Zhukov, *Phys. Rev. C* **84**, 021303 (2011), URL <http://link.aps.org/doi/10.1103/PhysRevC.84.021303>.
- [13] A. Spyrou, Z. Kohley, T. Baumann, D. Bazin, B. A. Brown, G. Christian, P. A. DeYoung, J. E. Finck, N. Frank, E. Lunderberg, et al., *Phys. Rev. Lett.* **108**, 102501 (2012), URL <http://link.aps.org/doi/10.1103/PhysRevLett.108.102501>.
- [14] Z. Kohley, T. Baumann, D. Bazin, G. Christian, P. A. DeYoung, J. E. Finck, N. Frank, M. Jones, E. Lunderberg, B. Luther, et al., *Phys. Rev. Lett.* **110**, 152501 (2013), URL <http://link.aps.org/doi/10.1103/PhysRevLett.110.152501>.
- [15] M. D. Jones, N. Frank, T. Baumann, J. Brett, J. Bullaro, P. A. DeYoung, J. E. Finck, K. Hammer-ton, J. Hinnefeld, Z. Kohley, et al., *Phys. Rev. C* **92**, 051306 (2015), URL <http://link.aps.org/doi/10.1103/PhysRevC.92.051306>.
- [16] A. Spyrou, J. K. Smith, T. Baumann, B. A. Brown, J. Brown, G. Christian, P. A. DeYoung, N. Frank, S. Mosby, W. A. Peters, et al., *Phys. Rev. C* **84**, 044309 (2011), URL <http://link.aps.org/doi/10.1103/PhysRevC.84.044309>.
- [17] J. Snyder, T. Baumann, G. Christian, R. A. Haring-Kaye, P. A. DeYoung, Z. Kohley, B. Luther, M. Mosby, S. Mosby, A. Simon, et al., *Phys. Rev. C* **88**, 031303 (2013), URL <http://link.aps.org/doi/10.1103/PhysRevC.88.031303>.
- [18] F. M. Marqués, N. A. Orr, N. L. Achouri, F. Delaunay, and J. Gibelin, *Phys. Rev. Lett.* **109**, 239201 (2012), URL <http://link.aps.org/doi/10.1103/PhysRevLett.109.239201>.
- [19] A. Spyrou, Z. Kohley, T. Baumann, D. Bazin, B. A. Brown, G. Christian, P. A. DeYoung, J. E. Finck, N. Frank, E. Lunderberg, et al., *Phys. Rev. Lett.* **109**, 239202 (2012), URL <http://link.aps.org/doi/10.1103/PhysRevLett.109.239202>.
- [20] L. V. Grigorenko, I. G. Mukha, and M. V. Zhukov, *Phys. Rev. Lett.* **111**, 042501 (2013), URL <http://link.aps.org/doi/10.1103/PhysRevLett.111.042501>.
- [21] K. Hagino and H. Sagawa, *Phys. Rev. C* **89**, 014331 (2014), URL <http://link.aps.org/doi/10.1103/PhysRevC.89.014331>.
- [22] B. Danilin, I. Thompson, J. Vaagen, and M. Zhukov, *Nuclear Physics A* **632**, 383 (1998), ISSN 0375-9474, URL <http://www.sciencedirect.com/science/article/pii/S0375947498000025>.
- [23] B. Brown, private communication (2013).
- [24] E. K. Warburton and B. A. Brown, *Phys. Rev. C* **46**, 923 (1992), URL <http://link.aps.org/doi/10.1103/PhysRevC.46.923>.
- [25] I. J. Thompson (2013).
- [26] I. Thompson, F. Nunes, and B. Danilin, *Computer Physics Communications* **161**, 87 (2004), ISSN 0010-4655, URL <http://www.sciencedirect.com/science/article/pii/S0010465504002140>.
- [27] D. Gogny, P. Pires, and R. D. Tourreil, *Physics Letters B* **32**, 591 (1970), ISSN 0370-2693, URL <http://www.sciencedirect.com/science/article/pii/0370269370905526>.
- [28] F. Nunes, J. Christley, I. Thompson, R. Johnson, and

- V. Efros, Nuclear Physics A **609**, 43 (1996), ISSN 0375-9474, URL <http://www.sciencedirect.com/science/article/pii/0375947496002849>.
- [29] N. K. Timofeyuk, I. J. Thompson, and J. A. Tostevin, Journal of Physics: Conference Series **111**, 012034 (2008), URL <http://stacks.iop.org/1742-6596/111/i=1/a=012034>.
- [30] I. Brida and F. M. Nunes, International Journal of Modern Physics E **17**, 2374 (2008), <http://www.worldscientific.com/doi/pdf/10.1142/S0218301308011641>, URL <http://www.worldscientific.com/doi/abs/10.1142/S0218301308011641>.
- [31] I. Brida, F. Nunes, and B. Brown, Nuclear Physics A **775**, 23 (2006), ISSN 0375-9474, URL <http://www.sciencedirect.com/science/article/pii/S0375947406003630>.
- [32] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, Phys. Rev. C **51**, 38 (1995), URL <http://link.aps.org/doi/10.1103/PhysRevC.51.38>.
- [33] R. V. R. Jr., Annals of Physics **50**, 411 (1968), ISSN 0003-4916, URL <http://www.sciencedirect.com/science/article/pii/0003491668901267>.
- [34] M. Thoennessen, S. Yokoyama, A. Azhari, T. Baumann, J. A. Brown, A. Galonsky, P. G. Hansen, J. H. Kelley, R. A. Kryger, E. Ramakrishnan, et al., Phys. Rev. C **59**, 111 (1999), URL <http://link.aps.org/doi/10.1103/PhysRevC.59.111>.
- [35] A. N. Kuchera, A. Spyrou, J. K. Smith, T. Baumann, G. Christian, P. A. DeYoung, J. E. Finck, N. Frank, M. D. Jones, Z. Kohley, et al., Phys. Rev. C **91**, 017304 (2015), URL <http://link.aps.org/doi/10.1103/PhysRevC.91.017304>.