

This is the accepted manuscript made available via CHORUS. The article has been published as:

Bulk and isospin instabilities in hot nuclear matter

V. M. Kolomietz and S. Shlomo

Phys. Rev. C **95**, 054613 — Published 16 May 2017

DOI: [10.1103/PhysRevC.95.054613](https://doi.org/10.1103/PhysRevC.95.054613)

Bulk and isospin instabilities in hot nuclear matter

V.M. Kolomietz¹⁾ and S. Shlomo^{2,3)}

¹⁾Institute for Nuclear Research, 03680 Kiev, Ukraine

²⁾Cyclotron Institute, Texas A&M University, College Station, Texas 77843, USA

³⁾Department of Elementary Particles and Astrophysics, the Weizmann Institute of Science, Rehovot 76100, Israel

The instability of hot asymmetric nuclear matter with respect to bulk density distortions is considered. The equation of state of extended Thomas-Fermi approximation (ETFA) is used. The origin of the anomalous dispersion and the influence of the Fermi-surface distortion effects on the bulk and isospin instabilities in homogenous nuclear matter are investigated. It is shown that the development of both instabilities is significantly reduced due to the Fermi surface distortion effects. The dependence of the bulk instability on the temperature and on the multipolarity of the particle density distortions are shown for the ^{208}Pb nucleus. The dependence of the formation of the decay modes (fission or multifragmentation) of nuclei on the temperature and the Fermi-surface distortion effects are demonstrated. It is shown that in the case of low temperatures the preferable mode for the bulk instability is binary fission. For higher temperature, the preferred modes are the multifragmentations to small clusters. The number of clusters increases with increasing temperature.

Keywords: hot nuclear matter; instability growth rate; Fermi surface distortions

PACS: PACS number: 25.70-z, 25.70.Mn, 24.75.+i

I. INTRODUCTION

Diluted nuclear matter is unstable with respect to particle density fluctuations. One expects that such a kind of bulk instability plays a significant role in internal clusterization and multifragmentation in heavy ion collisions. The bulk instability of nuclear matter was studied earlier in Refs. [1,2,3,4,5,6,7,8] where the different aspects of the nuclear equation of state were taken into consideration.

We note, however, that in actual nuclear processes (heavy-ion reactions, nuclear fission, etc.), the development of instability depends not only on the equation of state, but also on dynamic effects such as the dynamic Fermi-surface distortions (FSD). In particular, the FSD effects lead to collisional relaxation on the distorted Fermi surface and to non-Markovian motion in a viscous nuclear Fermi liquid [9]. In the present paper, we mainly focus on these aspects in studying bulk instability, considering the regimes of frequent and rare internucleon collisions.

Some specific difficulties appear in the theoretical description of the instability in an arbitrary dilute many body system due to the necessity to consider the small density fluctuations around a nonequilibrium state of the system. Usually the nonequilibrium state can be fixed through the introduction of a formal constrained field [10]. Note also that in considering the development of bulk instability in a homogeneous nuclear matter one has to use an equation of state (EOS) which is extended to include the gradient terms of particle density ρ . That is because the particle density perturbations $\delta\rho(\mathbf{r}, t)$ are, in general, non-homogeneous ones. In the present paper we use the EOS which is based on the semiclassical extended Thomas-Fermi approximation for the internal kinetic energy of nucleons and the potential energy due to the Skyrme effective nucleon-nucleon interaction. The inclusion of the gradient terms in the EOS leads to a specific effect of the anomalous dispersion which plays a significant role in the description of the instability growth rate in nuclear matter.

The present paper is an extension of our previous work [7] where the general concept of the Fermi motion effects was adopted in studying the bulk instability of nuclear matter. In the present paper we generalize our approach to both the spinodal (bulk) and isospin instabilities and carry out numerical calculations employing commonly used Skyrme interactions. We pay special attention to the derivation of the stiffness coefficients, such as the incompressibility and isospin symmetry coefficients, beyond the equilibrium point by employing the variational Euler-Lagrange procedure. We analyze the occurrence of the anomalous dispersion in both the isoscalar and the isovector channels. We consider the temperature dependence of the instability growth rate for the bulk mode. We study also the dependence of the instability growth rate on the multipolarity of the

nuclear density distortions in hot nuclei with increasing temperature. The last aspect is important since an increase of the temperature strongly reduces the Fermi-surface distortion effects and the relaxation processes and thereby the growth of instabilities. We point out that the use of the Skyrme interaction with parameters adjusted to reproduce the ground state properties of nuclei within the mean-field model, is a reasonable approximation for our purposes. In fact, the effective interaction is modified only slightly (by a few percent) in a wide temperature range, $T = 0 \div 20$ MeV [11,12].

In Section II we describe the fluid dynamic approach to the instability of nuclear matter by using the extended EOS which includes the gradient corrections to the energy density functional. The approach is based on the extended Thomas-Fermi approximation and the effective Skyrme interactions. In Section III we provide numerical results for the instability growth rate and its dependence on the temperature and the multipolarity of the particle density perturbations. Our conclusions are given in Section IV.

I. BULK INSTABILITY OF ASYMMETRIC NUCLEAR MATTER

We consider the \mathbf{r} -dependent density fluctuations $\delta\rho(\mathbf{r},t)$ around an arbitrary nonequilibrium density $\rho_0 = \text{const}$ of a homogeneous nuclear matter. Following the Euler-Lagrange method, the nonequilibrium particle density ρ_0 can be fixed by introducing a formal constrained field into the relevant variational procedure for the total energy E of a many body system. Employing this method to nuclear matter and assuming the possible \mathbf{r} -dependence of the particle density ρ , we write its total energy as

$$E = \int d\mathbf{r} \epsilon_{\text{tot}}[\rho], \quad \epsilon_{\text{tot}}[\rho] = \epsilon_{\text{kin}}[\rho] + \epsilon_{\text{pot}}[\rho], \quad (1)$$

where $\epsilon_{\text{kin}}[\rho]$ and $\epsilon_{\text{pot}}[\rho]$ are the \mathbf{r} -dependent kinetic and potential energy densities of nuclear matter which are derived below. An arbitrary density ρ_0 is then given by the following variational Euler-Lagrange equation

$$\frac{\delta}{\delta\rho} \left[\int d\mathbf{r} (\epsilon_{\text{tot}}[\rho] - \xi\rho) \right] \Big|_{\rho=\rho_0} = 0, \quad (2)$$

where $-\xi\rho$ is the constrained field and ξ is the Lagrange multiplier of the Euler-Lagrange method. Below we use the density ρ_0 as an independent variable and Eq. (2) is considered as a derivation of the Lagrange multiplier ξ only. The inclusion of the constrained field in the variational procedure of Eq. (2) plays an important conceptual role because it allows us to consider

the adiabatic stiffness coefficients beyond the equilibrium point.

In evaluating the potential energy density $\epsilon_{\text{pot}}[\rho]$ in Eq. (1), we adopt the Skyrme type effective nucleon-nucleon interaction in the following standard form [13,14,15]:

$$\begin{aligned} v_{12} = & t_0(1 + x_0 P_{12}^\sigma) \delta(\mathbf{r}_1 - \mathbf{r}_2) + \frac{1}{2} t_1(1 + x_1 P_{12}^\sigma) \left[\overleftarrow{k}_{12}^2 \delta(\mathbf{r}_1 - \mathbf{r}_2) + \delta(\mathbf{r}_1 - \mathbf{r}_2) \overrightarrow{k}_{12}^2 \right] \\ & + t_2(1 + x_2 P_{12}^\sigma) \overleftarrow{k}_{12} \delta(\mathbf{r}_1 - \mathbf{r}_2) \overrightarrow{k}_{12} + \frac{1}{6} t_3(1 + x_3 P_{12}^\sigma) \rho^\nu \left(\frac{\mathbf{r}_1 + \mathbf{r}_2}{2} \right) \delta(\mathbf{r}_1 - \mathbf{r}_2) \\ & + i W_0 \overleftarrow{k}_{12} \delta(\mathbf{r}_1 - \mathbf{r}_2) (\boldsymbol{\sigma}_1 + \boldsymbol{\sigma}_2) \times \overrightarrow{k}_{12}. \end{aligned} \quad (3)$$

where t_i , x_i , ν and W_0 are the parameters of the interaction, P_{12}^σ is the spin exchange operator, $\boldsymbol{\sigma}_i$ is the Pauli spin operator, $\overrightarrow{k}_{12} = -i(\overrightarrow{\nabla}_1 - \overrightarrow{\nabla}_2)/2$, and $\overleftarrow{k}_{12} = -i(\overleftarrow{\nabla}_1 - \overleftarrow{\nabla}_2)/2$. Here, the right and left arrows indicate that the momentum operators act on the right and on the left, respectively.

In a Fermi liquid, the linearized equation of motion for small variations $\delta\rho = \rho - \rho_0$ of the particle density near arbitrary ρ_0 can be written as, see Refs. [7,9],

$$m \frac{\partial^2}{\partial t^2} \delta\rho = \nabla\rho_0 \nabla \frac{\delta E}{\delta\rho} + \nabla_\nu \nabla_\mu P'_{\nu\mu}. \quad (4)$$

The equation of motion (4) is obtained from the kinetic equation for the Wigner's distribution function $f \equiv f(\mathbf{r}, \mathbf{p}; t)$ and includes both the relaxation phenomena and the Fermi-surface distortion effects [9]. The pressure tensor $P'_{\nu\mu}$ in Eq. (4) is caused by the dynamic distortions of the Fermi-surface, see Refs. [9,16,17]. We point out that the constrained field $-\xi\rho$ does not contribute to the equation of motion (4) since $\xi = \text{const.}$

Note that the total energy E of the homogeneous nuclear matter in Eq. (4) is written beyond the point $\rho_0 = \text{const.}$, where the particle density $\rho = (\mathbf{r}, t)$ becomes \mathbf{r} -dependent. To take this fact into consideration, we use the extended Thomas-Fermi approximation (ETFA) for the kinetic energy density $\epsilon_{\text{kin}}[\rho]$ (see Refs. [18,19]) and the density-dependent Skyrme interaction for the potential energy density $\epsilon_{\text{pot}}[\rho]$ [20] which include the gradient terms $\sim \nabla\rho$. Since the values $\epsilon_{\text{kin}}[\rho]$ and $\epsilon_{\text{pot}}[\rho]$ contain the gradients of the particle density, the evaluation of the functional derivative $\delta E/\delta\rho$ in Eq. (4) must have the form

$$\frac{\delta E}{\delta\rho} = \left[\frac{\partial}{\partial\rho} - \nabla \frac{\partial}{\partial(\nabla\rho)} + \nabla^2 \frac{\partial}{\partial(\nabla^2\rho)} \right] \epsilon_{\text{tot}}[\rho]. \quad (5)$$

We consider an asymmetric nuclear matter with asymmetry parameter $X = (N - Z)/(N + Z)$, where N and Z are the number of neutrons and protons, respectively. We introduce the isoscalar density $\rho_+ = \rho_n + \rho_p$ and the neutron excess (isovector) density $\rho_- = \rho_n - \rho_p = X\rho_+$, where ρ_n is the neutron density and ρ_p is the proton density. Below we restrict ourselves to

the commonly used case of a small isospin asymmetry $X \ll 1$ which allows one, in particular, to derive the isospin symmetry energy and is sufficient for all existing nuclei. Since $\rho_- \ll \rho_+$, the total energy density $\epsilon_{\text{tot}}[\rho]$ can be reduced to the following convenient form, see Appendix A, Eq. (A3),

$$\begin{aligned} \epsilon_{\text{tot}}[\rho] = & A_1(\rho_+) + A_2(\rho_+) \rho_-^2 + [B_1(\rho_+) + C_1(\rho_+, \rho_-)] (\nabla \rho_+)^2 \\ & + B_2(\rho_+) (\nabla \rho_-)^2 \\ & + D_1(\rho_+, \rho_-) \nabla^2 \rho_+, \end{aligned} \quad (6)$$

where the density dependent functions $A_i(\rho)$, $B_i(\rho)$, $C_1(\rho)$ and $D_1(\rho)$ are given in Appendix A, Eq. (A4). The expression (6) is written up to the second order in X . The higher order correction $\sim X^4$ is negligible in all practically interesting cases.

Using Eqs. (1) and (5), taking pressure tensor $P'_{\nu\mu}$ from Ref. [9] and assuming $\delta\rho \sim \exp[i(q \cdot r - \omega t)]$, the equation of motion (4) is transformed to the following form

$$-m \omega^2 \delta\rho_{\pm} = [C_{\pm}(\rho_0) - P_{F,\pm}(\rho_0)] \nabla^2 \delta\rho_{\pm} - F_{S,\pm}(\rho_0) \nabla^2 \nabla^2 \delta\rho_{\pm}. \quad (7)$$

Here, the transport coefficients C_{\pm} and $P_{F,\pm}$ are given by

$$\begin{aligned} C_+(\rho_0) &= K(\rho_0)/9, \quad C_-(\rho_0) = 2C_{\text{sym}}(\rho_0), \quad P_{F,\pm}(\rho_0) \\ &= \frac{4}{3} \frac{i\omega\tau_{\pm}}{1 - i\omega\tau_{\pm}} (P_0/\rho_0), \end{aligned} \quad (8)$$

where P_0 is the pressure $P_0 = \rho_0 p_F^2/5m$, p_F is the Fermi momentum and τ_{\pm} is the relaxation time which can be different for isoscalar and isovector channels [21]. The transport coefficients $F_{S,\pm}$ in Eq. (7) are the coefficients of the anomalous dispersion which play a significant role in the case of short-wave-length density fluctuations.

Taking into account the finite temperature, we use the adiabatic (isentropic) [22] values of transport coefficients in Eq. (7). Using Eqs. (7), (A5) and (A6) of Appendix A, one obtains for the isoscalar mode the incompressibility coefficient,

$$\begin{aligned} K(\rho_0) = & 10 \alpha \frac{\hbar^2}{2m} \rho_0^{2/3} + \frac{27}{4} t_0 \rho_0 + \frac{9}{16} (2 + \nu)(1 + \nu) t_3 \rho_0^{1+\nu} \\ & + 10 \alpha \left(\frac{3t_1 + 5t_2}{4} + t_2 x_2 \right) \rho_0^{5/3} \\ & + X^2 \left[\frac{10}{9} \alpha \frac{\hbar^2}{2m} \rho_0^{4/3} + \frac{3}{8} \nu(\nu - 1) t_3 \left(x_3 + \frac{1}{2} \right) \rho_0^{\nu+1} \right. \\ & \left. - \frac{10}{9} \alpha \left(\frac{t_2}{2} + \frac{5t_2 x_2 - 3t_1 x_1}{8} \right) \rho_0^{5/3} \right] \end{aligned} \quad (9)$$

and the isoscalar coefficient of the anomalous dispersion

$$F_{S,+}(\rho_0) = \frac{1}{2} \frac{\hbar^2}{2m} \eta + \left[\frac{1}{6} t_1 - \frac{25}{72} t_2 \left(1 + \frac{4}{5} x_2 \right) \right] \rho_0$$

$$-X^2 \left[\frac{1}{2} \frac{\hbar^2}{2m} \eta + \frac{1}{72} \left[t_1 \left(1 + \frac{x_1}{2} \right) + t_2 \left(1 + \frac{x_2}{2} \right) \right] \rho_0 \right], \quad (10)$$

where $\alpha = (3/5) (3\pi^2/2)^{2/3}$.

Similarly, for the isovector mode one has the coefficient C_{sym} of isospin symmetry,

$$C_{\text{sym}}(\rho_0) = \frac{5}{9} \alpha \frac{\hbar^2}{2m} \rho_0^{2/3} - \frac{t_0}{4} \left(x_0 + \frac{1}{2} \right) \rho_0 - \frac{t_3}{24} \left(x_3 + \frac{1}{2} \right) \rho_0^{v+1} \\ + \frac{5}{9} \alpha \left(\frac{t_2}{2} + \frac{5t_2x_2 - 3t_1x_1}{8} \right) \rho_0^{5/3} \quad (11)$$

and the isovector coefficient $F_{S,-}$ of the anomalous dispersion,

$$F_{S,-}(\rho_0) = \frac{1}{2} \eta \frac{\hbar^2}{2m} - \left[\frac{t_1}{24} \left(1 + \frac{5x_1}{2} \right) + \frac{t_2}{18} \left(1 + \frac{19x_2}{8} \right) \right] \rho_0. \quad (12)$$

Below, we present results of numerical calculations of the incompressibility coefficient $K(\rho_0)$ and the symmetry energy coefficient $C_{\text{sym}}(\rho_0)$ for the SkM* [20], Sly203b [15] and the more modern KDE0v1 [23] Skyrme interactions. We note that the KDE0v1 interaction was determined by a fit to extensive data on binding energies, charge radii of nuclei, single particle spin-orbit splitting, and for the first time inclusion of the radii of valence single particle neutron orbits and the energies of the isoscalar giant monopole resonance (ISGMR) in nuclei. Also included in the fit are additional constraints, such as the Landau stability conditions, positive derivative of the symmetry energy, $L = 3\rho dC_{\text{sym}}/d\rho$, at large particle density, ρ , [24,25] and a minimal value for the enhancement factor, κ , of the energy weighted sum rule (EWSR) of the isovector giant dipole resonance (IVGDR). It is important to point out that 240 Skyrme interactions, published in the literature, were analyzed by an independent group [26] for their ability to pass constraints relating to experimental data on properties of nuclear matter and their derivatives, such as incompressibility coefficient K , symmetry energy coefficient, C_{sym} , and its slope, L , effective mass, m^*/m , and observational data of neutron stars. Only the KDE0v1, LNS, NRPAR, SKRA and SQMC700 interactions passed the test. These five interactions were then tested in Ref. [27] for their ability to reproduce binding energies and fission barriers of nuclei. Only KDE0v1 passed the test. We add that although Skyrme parameterizations are not expected, in general, to reproduce data to which they were not fitted, the neutron star and fission barrier data that are reproduced by the KDE0v1 interaction were not included in the fit. We note that although the LNS, NRPAR, SKRA and SQMC700 were fitted to a wide range of data for the equation of states, we have adopted the KDE0v1 interaction for its ability to reproduce properties of finite nuclei, a reasonable approximation for our purposes. In Table I we present the values of the Skyrme parameters for the SkM*, KDE0v1 and Sly230b interactions and the corresponding

physical quantities of nuclear matter at saturation density, ρ_{eq} .

Table I: Values of the Skyrme parameters and the corresponding physical quantities of nuclear matter for the SkM*, KDE0v1 and Sly230b interactions.

Parameters	<i>SkM*</i>	<i>KDE0v1</i>	<i>Sly230b</i>
t_0 ($\text{MeV} \cdot \text{fm}^3$)	-2645.00	-2553.0843	-2488.91
t_1 ($\text{MeV} \cdot \text{fm}^5$)	410.00	411.6963	486.82
t_2 ($\text{MeV} \cdot \text{fm}^5$)	-135.00	-419.8712	-546.39
t_3 ($\text{MeV} \cdot \text{fm}^{3(1+\nu)}$)	15595.00	14603.6069	13777.0
x_0	0.0900	0.6483	0.8340
x_1	0.0000	-0.3472	-0.3438
x_2	0.0000	-0.9268	-1.0
x_3	0.0000	0.9475	1.3539
W_0 ($\text{MeV} \cdot \text{fm}^5$)	130.00	124.4100	122.69
ν	0.16667	0.1673	0.166667
E/A	15.78	16.23	15.972
K (MeV)	216.7	227.54	229.90
ρ_{eq} (fm^{-3})	0.160	0.165	0.160
m^*/m	0.79	0.74	0.695
C_{sym} (MeV)	30.03	34.58	32.01
L (MeV)	45.78	54.69	45.97
κ	0.53	0.23	0.25
T_{crit} (MeV)	14.62	14.74	14.67

Note that, in general, the transport coefficients K , C_{sym} , and $F_{S,\pm}$ in the equations of motion (7) are temperature dependent. This can be seen, in a transparent way, by normalizing the density ρ_0 to the equilibrium density $\rho_{\text{eq}}(T)$ and using the dimensionless ratio $\rho_0/\rho_{\text{eq}}(T)$ as a variable. To avoid any misunderstanding, we point out that the particle density ρ_0 is an independent variable which is fixed by the Lagrange multiplier ξ in the variational Euler-Lagrange equation (2). In the case of $\xi = 0$ and zero temperature $T = 0$, the variational equation (2) provides the actual equilibrium state with the saturation density $\rho_0 = \rho_{\text{eq}}(T = 0)$. In a heated system at $T \neq 0$ and below the phase separation point, the equilibrium density $\rho_{\text{eq}}(T)$ is derived by the equilibrium condition for the pressure $P(\rho, T) = 0$, where $P(\rho, T) = \rho^2 \partial F(\rho, T) / \partial \rho$ and $F(\rho, T)$ is the free energy. For higher temperatures above the point of the phase separation, the value of $\rho_{\text{eq}}(T)$ is obtained from the interphase equilibrium condition [22]. The temperature dependence of the equilibrium density $\rho_{\text{eq}}(T)$ can be approximated as $\rho_{\text{eq}}(T) = \rho_{\text{eq}}(T = 0)(1 - 1.6 \cdot 10^{-3} T^2)$, where temperature T is taken in MeV, see Ref. [28]. In Figs. 1 and 2, we show the density dependence of the incompressibility, K , and the symmetry

energy coefficient, C_{sym} , respectively, for different temperatures T for three sets of Skyrme forces, SkM^{*}, KDE0v1 and Sly230b.




Fig. 1. The density dependence of the incompressibility coefficient $K(\rho_0)$ for different temperatures $T = 0, 8$ and 14 MeV (shown) for Skyrme interactions SkM^{*} (dotted lines), KDE0v1 (solid lines) and SLy230b (dashed lines).

As seen from Fig.1, the instability regime where $K < 0$ is shifted to higher values of the ratio $\rho_0/\rho_{\text{eq}}(T)$ with increasing temperature T . The stable mode disappears at the critical temperature $T_{\text{crit}} = 14 \div 15$ MeV where $K = 0$ at $\rho_0/\rho_{\text{eq}}(T) = 1$, see Table I.

Fig. 2 shows that the dependence of symmetry energy coefficient, $C_{\text{sym}}(\rho_0)$, on the particle density ρ_0 is strongly sensitive to the choice of the Skyrme interactions. In contrast to the behavior of the incompressibility coefficient $K(\rho_0)$ in Fig. 1, the ρ -dependence of the symmetry energy coefficient $C_{\text{sym}}(\rho_0)$ is completely different for the SkM^{*}, KDE0v1 and Sly230b interactions. This fact was noted earlier in Ref. [14]. In the case of SkM^{*} interaction, the symmetry energy coefficient $C_{\text{sym}}(\rho_0)$ is a non-monotonic function of the density and it disappears in a superdense nuclear matter. In this case, the symmetry energy coefficient $C_{\text{sym}}(T)$ and thereby the isospin stability of Fermi liquid decreases with temperature for the dilute regime at $\rho_0/\rho_{\text{eq}}(T) \lesssim$

1. This behavior is reversed for the super dense regime at $\rho_0/\rho_{\text{eq}}(T) \gtrsim 2$. For the KDE0v1 and Sly230b interactions, the symmetry energy coefficient $C_{\text{sym}}(\rho_0)$ is an increasing function of particle density and the nuclear matter does not reach the instability regime. Thus, the density dependence of the symmetry energy coefficient $C_{\text{sym}}(\rho_0)$ and the occurrence of the isospin instability are sensitive to the Skyrme interaction parametrization.




Fig. 2. The same as in Fig. 1, but for symmetry energy coefficient C_{sym} .

III. FERMI-SURFACE DISTORTION AND RELAXATION EFFECTS ON THE GROWTH OF BULK INSTABILITY

The solutions of the equations of motion (7) are significantly different for the stable and unstable modes. Focusing on the unstable modes, $K < 0$ (or $C_{\text{sym}} < 0$), we introduce a growth rate of instability $\Gamma = -i\omega$ (Γ is real, $\Gamma > 0$) and obtain from Eqs. (7) the following dispersion relations

$$\Gamma_{\pm}^2 = |u_{1,\pm}|^2 q^2 - \zeta(\Gamma_{\pm}) q^2 - \kappa_{s,\pm} q^4, \quad (13)$$

where $u_{1,+} = \sqrt{K/9m}$, $u_{1,-} = \sqrt{2C_{\text{sym}}/m}$ and $\kappa_{s,\pm} = (2/m)F_{s,\pm}$. In the case of stable modes at $K > 0$ and $C_{\text{sym}} > 0$, the values $u_{1,+}$ and $u_{1,-}$ are the first sound velocities of the isoscalar and the isovector modes, respectively. The quantity $\zeta(\Gamma_{\pm})$ in Eq. (13) occurs due to the Fermi-surface distortion effect and it is given by, see also [7],

$$\zeta(\Gamma_{\pm}) = \frac{4}{3m} \frac{\Gamma_{\pm} \tau_{\pm}}{1 + \Gamma_{\pm} \tau_{\pm}} \frac{P_0}{\rho_0}. \quad (14)$$

We have performed numerical calculations of the growth rate Γ_+ of bulk instability for the Skyrme interaction KDE0v1. The relaxation time was taken in the form, see [29],

$$\tau_{\pm} = \hbar \beta_{\pm} / T^2 \quad (15)$$

with $\beta_+ = 9.2$ MeV, $\beta_- = 4.6$ MeV [30] and we have used $\eta = 4/9$ for the Weizsäcker's correction in Eqs. (10) and (12). The bulk density ρ_0 of the unstable mode was taken as $\rho_0 = 0.3 \rho_{\text{eq}}(T)$.




Fig. 3. The dependence of the isoscalar instability growth rate $\Gamma_+(q)$ on the wave number q (in units of $k_F = p_F/\hbar$). The solid curves are for the Fermi liquid for different temperatures (shown near solid curves in MeV) with bulk density $\rho_0 = 0.3 \rho_{\text{eq}}(T)$ and the relaxation parameter $\beta = 9.2$ MeV (see Eq. (15)). The dashed line is the result for the nonviscous liquid without the Fermi surface distortion effects at temperature $T = 2$ MeV. The calculations were performed for KDE0v1 interaction.

The non-monotony behavior of the isoscalar instability growth rate $\Gamma_+(q)$ as a function of the wave number q is caused by the anomalous dispersion term in Eq. (13). This term provides the stability with respect to the short-wave-length density fluctuations with increasing q . Comparing the dashed line and solid line 2 in Fig. 3, one concludes that the presence of the Fermi surface distortion effects reduces significantly the instability. This fact was noted earlier in Ref. [7]. As seen from Fig. 3, the FSD effects are decreasing with increasing temperature T . This is due to the increase of the smearing of the Fermi surface with increasing T and thereby the role of the Fermi surface distortions becomes weaker with T . The left slopes of lines in Fig. 3 indicate preference for nuclear multifragmentation (an increase of $\Gamma_+(q)$ with increasing q) while the right slopes are for the nuclear fission (an increase of $\Gamma_+(q)$ with decreasing q). With increasing temperature, a point from the right slope of curve $\Gamma(q)$ can appear on its left slope. Due to this fact, a nucleus which is unstable with respect to the fission mode at low temperature T can become unstable to the multifragmentation mode at higher T .

Considering the isovector mode, we first note that out of the three parametrizations of Skyrme-type interaction considered in this work, an instability regime with $C_{\text{sym}} < 0$ is reached in a superdense nuclear matter for the SkM^{*} interaction (representing other interactions with negative value of C_{sym} in a superdense matter). We add that the regime of superdense matter can be of interest, in particular, for the astrophysics. The results of numerical calculations of the isovector growth rate $\Gamma_-(q)$ for the Skyrme interaction SkM^{*} are shown in Fig. 4. As seen from Fig. 4, in contrast to Fig. 3 for $\Gamma_+(q)$, the isovector growth rate $\Gamma_-(q)$ in a superdense nuclear matter is a steadily increasing function of the wave number q . This is due to the fact that the coefficient of the anomalous dispersion $F_{s,-}$ in Eq. (7) changes sign from positive to negative in a superdense nuclear matter due to the presence of the term $\sim \rho_0$ in Eq. (12). Note also that the Fermi surface distortions have little effect on the isovector growth rate $\Gamma_-(q)$ (compare the dashed and solid lines 2 in Fig. 4). This is similar to the case of zero sound modes in stable nuclear matter [8] where the Fermi-surface distortion effects do not strongly influence the characteristics of the isovector excitations.




Fig. 4. The dependence of the isovector instability growth rate $\Gamma_-(q)$ on the wave number q (in units of $k_F = p_F/\hbar$). The solid curves are for the Fermi liquid for different temperatures (shown near solid curves in MeV) with bulk density $\rho_0 = 6 \rho_{eq}(T)$ and the relaxation parameter $\beta = 4, 6$ MeV (see Eq. (15)). The dashed line is the result for the nonviscous liquid without the Fermi surface distortion effects at temperature $T = 2$ MeV. The calculations were performed for SkM* interaction.

The dispersion relation (13) determines also the critical value $q_{\pm, \text{crit}}$ of the wave number which is given by the condition $\Gamma_{\pm}(q_{\pm, \text{crit}}) = 0$. The presence of the critical value of the wave number $q_{\pm, \text{crit}}$ plays an important role in the case of finite nuclei. For a finite nucleus the growth rate $\Gamma_+(q)$ depends on the multipolarity L of the density distortions $\delta\rho_{\pm}$ because the wave number $q = q_L$ is fixed by the boundary conditions which depend on the multipolarity L of the surface distortions [31,32]. Considering a finite nucleus, the dispersion relation (13) is completed by the boundary condition which can be taken similarly to the classical form [31,32] as

$$m |u_+|^2 \rho_0 j_L(qR_0) = (L-1)(L+2) \sigma j'_L(qr)/q R_0^2|_{r=R_0}, \quad (16)$$

where R_0 is the nuclear radius, σ is the surface tension coefficient and u_+ is the zero sound velocity, see Refs. [7,9]. The secular equation (16) gives the eigenvalue of the wave number q_L and the instability growth rate $\Gamma_+(q)$ depends on the position of q_L in the interval of $q = 0 \div q_{\text{crit}}$ [33].

The eigenvalues q_L of secular equation (16) are temperature dependent because the surface tension coefficient σ in Eq. (16) is temperature dependent. In numerical calculations, we have used the following expression for $\sigma(T)$, (see Refs. [34,35]),

$$\sigma(T) = 1.1 \left(\frac{T_{crit}^2 - T^2}{T_{crit}^2 + T^2} \right)^{5/4} \text{ MeV} \cdot \text{fm}^{-2}. \quad (17)$$




Fig. 5. The dependence of the isoscalar instability growth rate $\Gamma_+(L)$ on the multipolarity L of the particle density fluctuations for the nucleus ^{208}Pb for different temperatures. 0.5, 4, 6 and 8 **MeV** which are shown near the curves. The calculations were performed for KDE0v1 interaction.

In Fig. 5 we have plotted the isoscalar instability growth rate $\Gamma_+(L)$ as function of the multipolarity L of the particle density fluctuations for the nucleus ^{208}Pb for different temperatures T . As seen from Fig. 5, the development of instability is significantly different for different temperature regimes. In the case of low temperatures (line $T = 0.5$), the preferable mode is the binary fission (low multipolarity $L = 2$ is most unstable). For higher temperature (lines $T = 4$ and 8), the more preferable unstable modes are the ones with high multipolarity L which correspond to the multifragmentation to small clusters. The number of clusters (the interval of accessible multiplicities L on the left slope of curve $\Gamma_+(L)$) increases with the further increasing

temperature (compare lines 4, 6 and 8 MeV in Fig. 5). Note also that the modes with very high multipolarity L become overdamped having, $\Gamma_+(L) < 0$. This is in agreement with the results of Fig. 3 and reflects the fact that the eigenvalues q_L are increasing with L .

IV. SUMMARY AND CONCLUSIONS

We have considered the appearance and the development of instabilities in an asymmetric nuclear matter in both the isoscalar and the isovector channels. Our analysis is based on the equations of motion for the quantum Fermi liquid in the presence of the Fermi-surface distortion effects and the relaxation processes. We point out that a realistic description of the unstable modes in a homogeneous nuclear matter requires the extension of the equation of state by taking into consideration the gradient corrections to the total energy functional. The presence of the gradient corrections leads to the anomalous dispersion term in the equation of motion and influences significantly the behavior of the instability growth rate.

Studying the appearance of the bulk and isospin instabilities, we have performed numerical calculations of the dependence of the incompressibility coefficient $K(\rho_0)$ and the symmetry energy coefficient $C_{\text{sym}}(\rho_0)$ on the nuclear density ρ_0 . To evaluate the values of $K(\rho_0)$ and $C_{\text{sym}}(\rho_0)$ at non-equilibrium density $\rho_0 \neq \rho_{\text{eq}}$ we have applied the cranking approach. The external cranking field was used in the form which excludes the direct contribution of the cranking field to the incompressibility coefficient $K(\rho_0)$ for the Skyrme interactions SkM^{*}, KDE0v1 and Sly230b for different temperatures and particle densities. We have established the critical temperature $T_{\text{crit}} = 14 \div 15$ MeV where the nuclear matter becomes unstable at the equilibrium density $\rho_{\text{eq}}(T)$. A peculiarity of the isovector mode is that the instability of nuclear matter with respect to isovector density fluctuation, i.e. regime $C_{\text{sym}}(T) < 0$, can occur in the superdense nuclear matter only and depends significantly on the choice of the Skyrme interaction parametrization, such as the case of SkM^{*}.

We have shown that the Fermi-surface distortion effects strongly hinder the development of instabilities in nuclear matter. The dependence of the instability growth rate $\Gamma_+(q)$ on the wave number q has a specific non-monotonic behavior (see Figs. 3 and 5) which is caused by the anomalous dispersion term. Different slopes of the curve $\Gamma_+(q)$ reflect different regimes (fission or multifragmentation) of instability, see also Ref. [33]. To illustrate this fact, we have considered the behavior of instability growth rate $\Gamma_+(q)$ in the finite nucleus ^{208}Pb for different multipolarities L of particle density fluctuations and different temperatures T . The results

presented in Fig. 5 show that the fission mode (low L) is preferable at low temperature. The instability with respect to multifragmentation (high L) increases with temperatures. Moreover, one can expect that the yield of small fragments, which correspond to the highest values of L , is strongly increasing for high temperatures, see corresponding lines 4, 6 and 8 MeV in Fig. 5.

V. ACKNOWLEDGMENTS

This work was supported in part by the National Academy of Sciences of Ukraine under grant # CO-2-14/2016 (V. M. K.) and by the US Department of Energy under contract # DOE-FG03-93ER40773 (S. S.). S. Shlomo also thanks the Weizmann Institute of Science for the Weston Visiting Professorship Award and the nice hospitality extended to him.

APPENDIX A. Skyrme energy functional

In this paper we use the extended Thomas-Fermi approximation (ETFA) for the kinetic energy of nucleons and the Skyrme potential energy due to internucleon interactions. Assuming $\rho_- \ll \rho_+$ and using the standard form of the Skyrme interaction, Eq. (3), we represent the total energy density $\epsilon_{\text{tot}}[\rho_n, \rho_p]$ in the following form [20,36]

$$\begin{aligned} \epsilon_{\text{tot}}[\rho_+, \rho_-] = & \frac{\hbar^2}{2m} \alpha \rho_+^{5/3} + \frac{3}{8} t_0 \rho_+^2 + \frac{1}{16} t_3 \rho_+^{2+\nu} + \frac{1}{4} \left(\frac{3t_1 + 5t_2}{4} + t_2 x_2 \right) \alpha \rho_+^{8/3} + \\ & \left[\frac{\hbar^2}{2m} \frac{5}{9} \alpha \rho_+^{2/3} - \frac{t_0}{4} \left(x_0 + \frac{1}{2} \right) \rho_+ - \frac{t_3}{24} \left(x_3 + \frac{1}{2} \right) \rho_+^{1+\nu} + \frac{5}{9} \left(\frac{t_2}{2} + \frac{5t_2 x_2 - 3t_1 x_1}{8} \right) \alpha \rho_+^{5/3} \right] \frac{\rho_-^2}{\rho_+} + \\ & \frac{1}{4} \eta \frac{\hbar^2}{2m} \frac{(\nabla \rho_+)^2}{\rho_+} + \left(\frac{t_1}{12} - \frac{25}{144} t_2 \left(1 + \frac{4}{5} x_2 \right) \right) (\nabla \rho_+)^2 + \end{aligned} \quad (\text{A1})$$

$$\begin{aligned} & \left[\frac{1}{4} \eta \frac{\hbar^2}{2m} \left(\frac{\Delta \rho_+}{\rho_+} - \frac{(\nabla \rho_+)^2}{\rho_+^2} \right) + \frac{1}{144} \left(t_1 \left(1 + \frac{x_1}{2} \right) + t_2 \left(1 + \frac{x_2}{2} \right) \right) \Delta \rho_+ \right] \frac{\rho_-^2}{\rho_+} \\ & + \frac{1}{4} \eta \frac{\hbar^2}{2m} \frac{(\nabla \rho_-)^2}{\rho_+} - \left[\frac{t_1}{48} \left(1 + \frac{5x_1}{2} \right) + \frac{t_2}{36} \left(1 + \frac{19x_2}{8} \right) \right] (\nabla \rho_-)^2. \end{aligned} \quad (\text{A2})$$

Here, $t_0, t_1, t_2, t_3, x_0, x_1, x_2, x_3$ and ν are the parameters of Skyrme interaction and η is the free Weizsäcker's parameter of the ETFA.

One can rewrite (A1), using the following notations, in the form,

$$\epsilon_{\text{tot}}[\rho_+, \rho_-] = A_1(\rho_+) + A_2(\rho_+) \rho_-^2 + [B_1(\rho_+) + C_1(\rho_+, \rho_-)] (\nabla \rho_+)^2 + B_2(\rho_+) (\nabla \rho_-)^2$$

$$+D_1(\rho_+, \rho_-)\nabla^2\rho_+, \quad (\text{A3})$$

where

$$A_1(\rho_+) = \frac{\hbar^2}{2m}\alpha\rho_+^{5/3} + \frac{3}{8}t_0\rho_+^2 + \frac{1}{16}t_3\rho_+^{2+\nu} + \frac{1}{4}\left(\frac{3t_1+5t_2}{4} + t_2x_2\right)\alpha\rho_+^{8/3},$$

$$A_2(\rho_+) = \frac{\hbar^2}{2m}\frac{5}{9}\alpha\rho_+^{-1/3} - \frac{t_0}{4}\left(x_0 + \frac{1}{2}\right) - \frac{t_3}{24}\left(x_3 + \frac{1}{2}\right)\rho_+^\nu \\ + \frac{5}{9}\left(\frac{t_2}{2} + \frac{5t_2x_2 - 3t_1x_1}{8}\right)\alpha\rho_+^{2/3},$$

$$B_1(\rho_+) = \frac{1}{4}\eta\frac{\hbar^2}{2m}\frac{1}{\rho_+} + \left[\frac{t_1}{12} - \frac{25}{144}t_2\left(1 + \frac{4}{5}x_2\right)\right], \quad (\text{A4})$$

$$B_2(\rho_+) = \frac{1}{4}\eta\frac{\hbar^2}{2m}\frac{1}{\rho_+} - \left[\frac{t_1}{48}\left(1 + \frac{5x_1}{2}\right) + \frac{t_2}{36}\left(1 + \frac{19x_2}{8}\right)\right],$$

$$D_1(\rho_+, \rho_-) = \frac{1}{4}\eta\frac{\hbar^2}{2m}\frac{\rho_-^2}{\rho_+^2} + \frac{1}{144}\left[t_1\left(1 + \frac{x_1}{2}\right) + t_2\left(1 + \frac{x_2}{2}\right)\right]\frac{\rho_-^2}{\rho_+},$$

$$C_1(\rho_+, \rho_-) = -\frac{1}{4}\eta\frac{\hbar^2}{2m}\frac{\rho_-^2}{\rho_+^3}.$$

Using $\epsilon_{\text{tot}}[\rho]$ of the form of Eq. (A3), one can evaluate the functional derivatives $\delta E/\delta\rho_\pm$ for the equations of motion (4) which are significantly different for the isoscalar and isovector modes.

(i) *Isoscalar mode:*

$$\frac{\delta E}{\delta\rho_+} = \frac{\partial}{\partial\rho_+}A_1(\rho_+) + \rho_-^2\frac{\partial}{\partial\rho_+}A_2(\rho_+) + (\nabla\rho_+)^2\frac{\partial}{\partial\rho_+}[B_1(\rho_+) + C_1(\rho_+, \rho_-)] \\ + (\nabla\rho_-)^2\frac{\partial}{\partial\rho_+}B_2(\rho_+) + (\nabla^2\rho_+)\frac{\partial}{\partial\rho_+}D_1(\rho_+, \rho_-) \\ - 2\nabla[B_1(\rho_+) + C_1(\rho_+, \rho_-)]\nabla\rho_+ + \nabla^2D_1(\rho_+, \rho_-)$$

$$\frac{\delta E}{\delta\rho_+} = \left(\frac{\delta E}{\delta\rho_+}\right)_{\rho_0} + \left(\frac{\partial^2 A_1}{\partial\rho_+^2}\right)_{\rho_0}\delta\rho_+ + \left(\rho_-^2\frac{\partial^2 A_2}{\partial\rho_+^2}\right)_{\rho_0}\delta\rho_+ + \left(\frac{\partial(B_1 + C_1)}{\partial\rho_+}\right)_{\rho_0}(\nabla\delta\rho_+)^2 \\ + \left(\frac{\partial D_1}{\partial\rho_+}\right)_{\rho_0}\nabla^2\delta\rho_+ - 2\nabla\left(\frac{\partial(B_1 + C_1)}{\partial\rho_+}\right)_{\rho_0}\delta\rho_+ \cdot (\nabla\rho_+)_0$$

$$\begin{aligned}
& -2\nabla[B_1(\rho_+) + C_1(\rho_+, \rho_-)]_{\rho_0} \cdot \nabla \delta \rho_+ - 2 \left(\frac{\partial(B_1 + C_1)}{\partial \rho_+} \right)_{\rho_0} \delta \rho_+ (\nabla^2 \rho_+)_0 \\
& -2([B_1(\rho_+) + C_1(\rho_+, \rho_-)])_{\rho_0} \nabla^2 \delta \rho_+ + \left(\frac{\partial D_1}{\partial \rho_+} \right)_{\rho_0} \nabla^2 \delta \rho_+ \\
& \frac{\delta E}{\delta \rho_+} = \left(\frac{\delta E}{\delta \rho_+} \right)_{\rho_0} + \left[\left(\frac{\partial^2 A_1}{\partial \rho_+^2} \right)_{\rho_0} + \left(\rho_-^2 \frac{\partial^2 A_2}{\partial \rho_+^2} \right)_{\rho_0} \right] \delta \rho_+ \\
& + \left[2 \left(\frac{\partial D_1}{\partial \rho_+} \right)_{\rho_0} - 2([B_1(\rho_+) + C_1(\rho_+, \rho_-)])_{\rho_0} \right] \nabla^2 \delta \rho_+
\end{aligned} \tag{A5}$$

(ii) *Isovector mode:*

$$\begin{aligned}
\frac{\delta E}{\delta \rho_-} &= 2 A_2(\rho_+) \rho_- + (\nabla \rho_+)^2 \frac{\partial}{\partial \rho_-} C_1(\rho_+, \rho_-) + (\nabla^2 \rho_+) \frac{\partial}{\partial \rho_-} D_1(\rho_+, \rho_-) \\
& - 2 \nabla B_2(\rho_+) \cdot \nabla \rho_- \\
\frac{\delta E}{\delta \rho_-} &= \left(\frac{\delta E}{\delta \rho_-} \right)_{\rho_0} + 2 [A_2(\rho_+)]_0 \delta \rho_- + \left((\nabla \rho_+)^2 \frac{\partial^2}{\partial \rho_-^2} C_1(\rho_+, \rho_-) \right)_{\rho_0} \delta \rho_- \\
& + \left((\nabla \rho_+)^2 \frac{\partial^2}{\partial \rho_-^2} D_1(\rho_+, \rho_-) \right)_{\rho_0} \delta \rho_- \\
& - 2[\nabla B_2(\rho_+)]_{\rho_0} \cdot \nabla \delta \rho_- - 2[B_2(\rho_+)]_{\rho_0} \nabla^2 \rho_- \\
\frac{\delta E}{\delta \rho_-} &= \left(\frac{\delta E}{\delta \rho_-} \right)_{\rho_0} + 2 [A_2(\rho_+)]_{\rho_0} \delta \rho_- - 2[B_2(\rho_+)]_{\rho_0} \nabla^2 \rho_-
\end{aligned} \tag{A6}$$

We note that the first term on the right hand side in Eqs. (A5) and (A6) do not contribute to the equation of motion (4) because of the variational condition (2) and that the chemical potentials $\mu_{\pm} = \text{const.}$, namely,

$$\begin{aligned}
\mu_+ &= \frac{\partial E}{\partial A} \Big|_V = \left(\frac{\delta E}{\delta \rho_+} \right)_{\rho_0} = \text{const.}, \quad \mu_- = \frac{\partial E}{\partial (N - Z)} \Big|_V = \left(\frac{\delta E}{\delta \rho_-} \right)_{\rho_0} \\
&= \text{const.}
\end{aligned} \tag{A7}$$

- [1] H. Heiselberg, C. J. Pethick and D. G. Ravenhall, Ann. Phys. **223**, 37 (1993).
- [2] M. Colonna, Ph. Chomaz and J. Randrup, Nucl. Phys. **A567**, 637 (1994).
- [3] M. Colonna and Ph. Chomaz, Phys. Rev. C **49**, 1908 (1994).
- [4] J. Konopka, H. Graf, H. Stöcker, and W. Greiner, Phys. Rev. C **50**, 2085 (1994).
- [5] H. Müller and B. D. Serot, Phys. Rev. C **52**, 2072 (1995).

- [6] V. Baran, M. Colonna, M. Di Toro and A. B. Larionov, Nucl. Phys. **A632**, 287 (1998).
- [7] V. M. Kolomietz and S. Shlomo, Phys. Rev. C **60**, 044612 (1999).
- [8] V. M. Kolomietz and S. Shlomo, Phys. Rev. C **64**, 044304 (2001).
- [9] V. M. Kolomietz and S. Shlomo, Phys. Rep. **390**, 133 (2004).
- [10] B. Jacquot, M. Colonna, S. Ayik and Ph. Chomaz, Nucl. Phys. **A617**, 356 (1998).
- [11] A. Lejeune, P. Grange, M. Martzolf and J. Cugnon, Nucl. Phys. **A453**, 189 (1986).
- [12] J. Cugnon, A. Lejeune, M. Baldo and U. Lombardo, Nucl. Phys. **A492**, 173 (1989).
- [13] T. H. R. Skyrme, Nucl. Phys. **9**, 615 (1959).
- [14] E. Chabanat, P. Bonche, P. Haensel, J. Meyer and R. Schaeffer, Nucl. Phys. **A627**, 710 (1997).
- [15] E. Chabanat, R. Bonche, R. Haensel, J. Meyer, and R. Schaeffer, Nucl. Phys. **A635**, 231 (1998).
- [16] J. R. Nix and A. J. Sierk, Phys. Rev. C **21**, 396 (1980).
- [17] D. Kiderlen, V. M. Kolomietz and S. Shlomo, Nucl. Phys. **A608**, 32 (1996).
- [18] D. A. Kirzhnits, *Field Theoretical Methods in Many Body Systems* (Pergamon, London, 1967).
- [19] P. Ring and P. Schuck, *The Nuclear Many-Body Problem*, (Springer-Verlag, New York, 1980), Ch. 13.
- [20] M. Brack, C. Guet, H.-B. Håkansson, Phys. Rep. **123**, 275 (1985).
- [21] M. Di Toro, V. M. Kolomietz and A. B. Larionov, Phys. Rev. C **59**, 3099 (1999).
- [22] L. D. Landau and E. M. Lifshitz, *Statistical Physics* (Pergamon Press, Oxford, 1958)
- [23] B. K. Agrawal, S. Shlomo, and V. K. Au, Phys. Rev. C **72**, 014310 (2005).
- [24] C. J. Horowitz and J. Piekarewicz, Phys. Rev. C **66**, 055803 (2002).
- [25] J. R. Stone, J. C. Miller, R. Koncewicz, P. D. Stevenson, and M. R. Strayer, Phys. Rev. C **68**, 034324 (2003).
- [26] M. Dutta, O. Lourenco, J. S. Sa Martins, A. Delfino, J. R. Stone, and P. D. Stevenson, Phys. Rev. C **85**, 035201 (2012).
- [27] P. D. Stevenson, P. D. Goddard, J.R. Stone, and M. Dutta, arXiv:1210.1592v1[nucl-th] 2012.
- [28] S. Shlomo and V. M. Kolomietz, Rep. Prog. Phys. **68**, 1 (2005).
- [29] A. A. Abrikosov and I. M. Khalatnikov, Rep. Prog. Phys. **22**, 329 (1959).
- [30] V. M. Kolomietz, V.A. Plujko and S. Shlomo, Phys. Rev. C **54**, 3014 (1996).
- [31] H. Lamb, *Hydrodynamics* (Dover, New York, 1945).
- [32] A. Bohr and B. R. Mottelson, *Nuclear Structure*, Vol. 2, Ch.6 (Benjamin, New York, 1975).
- [33] C. J. Pethick and D. G. Ravenhall, Nucl. Phys. A **471**, 19c (1987); C. J. Pethick and D. G.

Ravenhall, Ann. Phys. (N.Y.) **183**, 131 (1988).

[34] D. G. Ravenhall, C. J. Pethick and J. M. Lattimer, Nucl. Phys. **A407**, 571 (1983).

[35] V. M. Kolomietz, A. I. Sanzhur, S. Shlomo and S. A. Firin, Phys. Rev. C **64**, 054315 (2001).

[36] V. M. Kolomietz and A. I. Sanzhur, Eur. Phys. J. A **38**, 345 (2008).