

This is the accepted manuscript made available via CHORUS. The article has been published as:

Event-by-event correlations between soft hadrons and  $D^0$  mesons in 5.02 TeV PbPb collisions at the CERN Large Hadron Collider

Caio A. G. Prado, Jacquelyn Noronha-Hostler, Roland Katz, Alexandre A. P. Suaide, Jorge Noronha, and Marcelo G. Munhoz

Phys. Rev. C **96**, 064903 — Published 18 December 2017

DOI: [10.1103/PhysRevC.96.064903](https://doi.org/10.1103/PhysRevC.96.064903)

# Event-by-event correlations between soft hadrons and $D^0$ mesons in 5.02 TeV PbPb collisions at LHC

Caio A. G. Prado,<sup>1,2</sup> Jacquelyn Noronha-Hostler,<sup>3,4</sup> Roland Katz,<sup>1</sup>  
Alexandre A. P. Suaide,<sup>1</sup> Jorge Noronha,<sup>1</sup> and Marcelo G. Munhoz<sup>1</sup>

<sup>1</sup>*Instituto de Física, Universidade de São Paulo, C.P. 66318, 05315-970 São Paulo, SP, Brazil*

<sup>2</sup>*Institute of Particle Physics and Key Laboratory of Quark and Lepton Physics (MOE),  
Central China Normal University, Wuhan, 430079, China*

<sup>3</sup>*Department of Physics, University of Houston, Houston TX 77204, USA*

<sup>4</sup>*Department of Physics and Astronomy, Rutgers University, Piscataway, NJ 08854, USA*

In this paper heavy quark energy loss models are embedded in full event-by-event viscous hydrodynamic simulations to investigate the nuclear suppression factor and the azimuthal anisotropy of  $D^0$  mesons in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV in the  $p_T$  range 8–40 GeV. In our model calculations, the  $R_{AA}$  of  $D^0$  mesons is consistent with experimental data from the CMS experiment. We present the first calculations of heavy flavor cumulants  $v_2\{2\}$  and  $v_3\{2\}$  (and also discuss  $v_2\{4\}$ ), which is also consistent with experimental data. Event-shape engineering techniques are used to compute the event-by-event correlation between the soft hadron  $v_n$  and the heavy meson  $v_n$ . We predict a linear correlation between these observables on an event-by-event basis.

## I. INTRODUCTION

In the last decade, remarkable progress has been made towards understanding the properties of the strongly interacting Quark-Gluon Plasma (QGP) produced in ultra-relativistic heavy ion collisions [1, 2]. A defining feature of the QGP is its ability to flow as a nearly perfect liquid where the shear viscosity to entropy density ratio  $\eta/s \sim 0.1$  is an order of magnitude smaller than in ordinary fluids such as water [3].

Event-by-event relativistic viscous hydrodynamic simulations [4] have revealed that in this QCD liquid viscous effects are so small that the spatial inhomogeneities present in the initial state are efficiently converted into final state momentum space anisotropy [5, 6]. One finds that the low- $p_T$  elliptic and triangular flows of all charged particles,  $v_2$  and  $v_3$ , are linearly correlated with the corresponding eccentricities of the initial state,  $\varepsilon_2$  and  $\varepsilon_3$  [7–10], while higher order flow harmonics exhibit some degree of nonlinear response [11–14].

In contrast, the physical mechanism responsible for azimuthal anisotropies at high- $p_T$  ( $p_T \gtrsim 10$  GeV) relies not on pressure and flow gradients but rather on differences in the path length of highly energetic probes quenched by the expanding medium [15, 16]. This qualitative understanding has been recently confirmed by the first jet energy loss + event-by-event viscous hydrodynamic calculations performed in [17, 18]. A novel feature found in [17, 18] is that the approximate linear response between  $v_2$  and  $\varepsilon_2$  also holds at high  $p_T$  on an event-by-event basis. This implies that the quantum randomness in the position of the nucleons in the incident nuclei, which determines the fluctuations of the initial conditions used in the subsequent hydrodynamic evolution, significantly affects the distribution of path lengths traversed by jets in the medium.

The observation of large azimuthal anisotropy of open heavy flavor mesons [19–21] adds important new elements to the overall picture discussed above. Heavy quarks are produced by hard processes in the initial stages of the collision with a non-thermal transverse momentum spectrum, which is expected to relax towards a nearly thermal distribution within a relaxation time scale  $\tau_R \sim \frac{M}{T} \frac{\eta}{sT}$  [22] ( $M$  is the heavy quark mass). The difference between the charm and bottom quark masses suggests that, at low- $p_T$ , the  $D^0$  meson  $v_n$  should be larger than the corresponding coefficients for  $B^0$  mesons [23]. Additionally, at low- $p_T$  quark coalescence [24–26] between heavy and light flavors [27, 28] can substantially increase the elliptic flow of heavy mesons [29, 30], as well as various effects such as Langevin type behavior and hadronic rescattering.

For  $p_T \gtrsim 10$  GeV heavy quarks hadronize mostly via fragmentation<sup>1</sup> and the various low- $p_T$  effects can be neglected. This provides a simpler scenario for studying how initial state spatial anisotropies are mapped into the final state heavy flavor azimuthal anisotropy. We address this problem by exploring new techniques that connect soft physics and heavy flavor observables. A heavy quark energy loss model is embedded on top of event-by-event viscous hydrodynamic backgrounds to study the nuclear suppression factor and  $v_n$  of  $D^0$  mesons in PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV for 8–40 GeV. We do not intend to find constraints in the chosen energy loss models but rather to discuss the sensitivity of the presented observables that could be used to better understand the physics of heavy quarks on an event-by-event basis. Our event-by-event approach predicts a linear correlation between the soft hadron and the heavy meson  $v_2$  and  $v_3$ , which could be verified at LHC. This linear relationship is not an obvious feature as the anisotropies in the soft and heavy sectors emerge from two very different production, interaction, and hadronization mechanisms.

In fact, non-linearities have already been observed in a study using all charged particles at high  $p_T$  [18]. The degree of linear correlation between  $\varepsilon_n \rightarrow v_n^{hard}$  can be quantified through a Pearson coefficient,  $Q_n$ ,

$$Q_n = \frac{\langle \varepsilon_n v_n(p_T) \cos n(\phi_n - \psi_n(p_T)) \rangle}{\sqrt{\langle \varepsilon_n^2 \rangle \langle v_n(p_T)^2 \rangle}} \quad (1)$$

that demonstrates the correlation between two vectors i.e. both the magnitude  $\varepsilon_n$  and angle  $\phi_n$  of the initial condition and how that maps onto the final magnitude  $v_n$  and angle  $\psi_n$  of the azimuthal anisotropy. Then a value of  $Q_n = (-)1$  implies a perfect (anti-)linear correlation whereas  $Q_n = 0$  implies absolutely no correlation. For all charged particles at high  $p_T$ ,  $Q_2$  for  $\varepsilon_n \rightarrow v_n^{hard}$  is shown in Fig. 6 from [18] where  $Q_2 \gtrsim 0.9$ , which indicates a quite linear correlation. However, compared to the soft sector it is clear that more non-linearities appear at high  $p_T$  because  $Q_2$  is smaller at high  $p_T$ . Furthermore,  $Q_2$  demonstrates even more non-linearities for a larger  $\eta/s$  and for peripheral collisions. Finally, higher order flow harmonics demonstrate even larger non-linearities demonstrated by values of  $Q_3 \approx 0.7$ – $0.9$ . The implications of these non-linearities and how they may vary depending on the mass of the particle still remains an interesting question that we will start to explore in this paper.

---

<sup>1</sup> We neglect possible D-like states that might form in the QGP [31].

## II. DETAILS OF THE MODEL

To simulate the propagation of heavy quark jets in the medium, we developed a modular Monte Carlo code in C++, named DABMod, combined with ROOT [32] and PYTHIA8 [33] libraries, that allows for a variety of energy loss models to be implemented. Starting from the initial conditions, we sample charm quarks (c) inside the medium with initial momentum distribution given by pQCD FONLL calculations [34, 35] and random initial direction. We neglect effects of the jets on the medium [36] but the local space-time dependence of the hydrodynamic fields is considered in the energy loss calculations.

For this first study we employ the following simple parametrization for the heavy quark energy loss per unit length [37]  $\frac{dE}{dx}(T, v) = -f(T, v) \Gamma_{\text{flow}}$ , where  $T$  is the temperature experienced by the heavy quark,  $v$  is the heavy quark velocity,  $\Gamma_{\text{flow}} = \gamma [1 - v_{\text{flow}} \cos(\varphi_{\text{quark}} - \varphi_{\text{flow}})]$  with  $\gamma = 1/\sqrt{1 - v_{\text{flow}}^2}$ , takes into account the boost from the local rest frame of the fluid [38],  $\varphi_{\text{quark}}$  is the angle defined by the propagating jet in the transverse plane, and  $\varphi_{\text{flow}}$  is the flow local azimuthal angle. The jet propagates only in the transverse plane, starting from a production point  $\mathbf{x}_0$ , moving in the direction defined by  $\varphi_{\text{quark}}$ .

We investigate the cases where  $f(T, v) = \xi T^2$  and  $f(T, v) = \alpha$ , with  $\xi$  and  $\alpha$  being constants. Both forms are simplified approximations for the interactions between heavy quarks and the strongly interacting QGP. The first choice is inspired by conformal AdS/CFT calculations [39] (for non-conformal plasma see [40]). The second choice is inspired by Ref. [41] which showed that a non-decreasing drag coefficient near the phase transition is favored for a simultaneous description of heavy flavor  $R_{\text{AA}}(p_T)$  and  $v_2(p_T)$  (this is also supported by  $T$ -matrix calculations [42, 43]). These models are fairly simple to implement and they give a good description of  $R_{\text{AA}}$ .

Our calculations use hydrodynamical profiles to provide the temperature and flow fields at each time step for each event. We generate hydrodynamic profiles using the 2D + 1 event-by-event relativistic viscous hydrodynamical model, v-USPhydro [44–46], which passes standard accuracy tests [47]. Our setup is the same as [17, 46] (MCKLN initial conditions at PbPb  $\sqrt{s_{\text{NN}}} = 5.02$  TeV [48–50],  $\eta/s = 0.05$ , and initial time  $\tau_0 = 0.6$  fm), which describes experimental data in the soft sector, such that all the hydrodynamic parameters are fixed in the present study. The heavy quarks are evolved on top of hydrodynamic backgrounds until they reach the jet-medium decoupling temperature  $T_d$  below which hadronization is performed using the Peterson fragmentation function [51]. This parameter encodes the large uncertainties regarding hadronization of jets in the QGP and is set to vary between  $T_d = 120$  MeV and  $T_d = 160$  MeV, inspired by [18, 52, 53]. The parameter for the Peterson fragmentation function is fixed so that the heavy meson spectra matches FONLL calculations. Coalescence is not taken into account but it will be implemented in future work to extend the validity of our calculations to low- $p_T$ . Hadronic rescattering is not significant at high- $p_T$  [54, 55] and is neglected here.

In this work, an event-by-event analysis is possible by oversampling each individual hydro event with millions of heavy quarks. From each individual event  $R_{\text{AA}}^c(p_T, \varphi)$  for charm quarks is obtained and the corresponding azimuthal coefficients

$$v_n^c(p_T) = \frac{\frac{1}{2\pi} \int_0^{2\pi} d\varphi \cos[n\varphi - n\psi_n^c(p_T)] R_{\text{AA}}^c(p_T, \varphi)}{R_{\text{AA}}^c(p_T)}, \quad (2)$$

are calculated based on [56] with

$$\psi_n^c(p_T) = \frac{1}{n} \tan^{-1} \left( \frac{\int_0^{2\pi} d\varphi \sin(n\varphi) R_{\text{AA}}^c(p_T, \varphi)}{\int_0^{2\pi} d\varphi \cos(n\varphi) R_{\text{AA}}^c(p_T, \varphi)} \right). \quad (3)$$

In reality, there are very few heavy quarks per event and our approach gives the probability for an event with a certain  $v_n$  and  $\psi_n$  in the soft sector to produce the heavy flavor quantities  $v_n^c(p_T)$  and  $\psi_n^c(p_T)$ . To compare with experimental data the scalar product method [57, 58] is used to calculate the 2 and 4-particle cumulants with the inclusion of multiplicity weighting and centrality class re-binning as described in [18, 59]. We have at least a couple of thousand hydrodynamic events in each centrality class and we checked that our statistical error bars (computed using jackknife resampling [60]) are on the order of  $10^{-4}$ .

The free parameters  $\xi$  and  $\alpha$  that define our two energy loss scenarios are determined by matching our model calculations for  $D^0 R_{\text{AA}}$  to experimental data at  $p_T \gtrsim 10$  GeV in the 0–10% centrality class. The same procedure has been used to fix the jet-medium coupling in the light sector in [17, 18, 37]. The parameters must be fixed for every decoupling temperature  $T_d$  separately. With these parameters fixed<sup>2</sup> we can perform a full simulation across all  $p_T$  and centralities.

<sup>2</sup> We use  $\xi = (16.000, 18.500)$  and  $\alpha = (0.764, 1.087)$  MeV, where the first term in the parenthesis represents  $T_d = 120$  MeV while the second represents  $T_d = 160$  MeV.




FIG. 1. (Color online)  $R_{AA}$  for  $D^0$  mesons in 0–10% PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV and decoupling temperature  $T_d = 120$  MeV comparing two energy loss models with experimental data from CMS collaboration [61].




FIG. 2. (Color online)  $v_2\{2\}$  of  $D^0$  mesons in 30–50% PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV comparing two energy loss models with CMS data [64]. The band corresponds to the decoupling temperature interval  $120 \text{ MeV} \leq T_d \leq 160 \text{ MeV}$ .

### III. NUMERICAL RESULTS

Fig. 1 shows our results for  $D^0$   $R_{AA}$ , together with run 2 LHC CMS data [61]. In this plot we use the decoupling temperature  $T_d = 120$  MeV and compare the two energy loss scenarios. One can observe that both energy loss models give the same nuclear modification factor in the  $p_T$  range considered. These results are robust with respect to variations in  $T_d$ , by appropriately fixing  $\xi$  and  $\alpha$ . Finally, since we do not use the same  $p_T$  dependence, our results differ from previous implementations of AdS/CFT-inspired energy loss calculations such as [62, 63].

We compute the differential  $v_2\{2\}$  [17, 18] for  $D^0$  meson at 30–50% centrality and compare it to experimental data in Fig. 2. At intermediate and high- $p_T$ , the temperature dependent energy loss model clearly underestimates the data, whereas the constant model  $dE/dx \sim \alpha$  is within the uncertainties. Both energy loss models underestimate the data at low- $p_T$ , where effects such as coalescence, shadowing, and stochastic dynamics could play a significant role. The constant model gives the largest elliptic flow (as it occurred in [41]) while the bands illustrate the dependence of these observables with  $T_d$ . We find that  $v_2\{2\}$  increases when  $T_d$  is lowered from 160 MeV to 120 MeV, which is expected due to the larger time available to build up the azimuthal anisotropy. Moreover, we note that even though the chosen  $T_d$  range is large, our results are quite robust concerning this parameter.

Our calculations for the corresponding  $v_3\{2\}$  of  $D^0$  can be found in Fig. 3, together with experimental data for




FIG. 3. (Color online)  $v_3\{2\}$  of  $D^0$  in 30–50% PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV comparing two energy loss models with CMS [64]. The band corresponds to the decoupling temperature interval  $120 \text{ MeV} \leq T_d \leq 160 \text{ MeV}$ .

comparison.  $v_3\{2\}$  is a factor  $\sim 3$  smaller than the corresponding  $v_2\{2\}$  shown in Fig. 2 and falls slightly below the experimental data at low- $p_T$ . By considering event-by-event simulations, one sees that a non-decreasing drag coefficient [41] gives not only the largest  $v_2\{2\}$  but also the largest  $v_3\{2\}$ , thereby amplifying the survival of the initial state fluctuations perceived by heavy flavor. Also,  $v_3\{2\}$  is particularly sensitive to the decoupling between heavy quarks and the medium in comparison with  $v_2\{2\}$  as the bands almost overlap. We find a slightly smaller  $D^0$  triangular flow than Refs. [29, 52], which is likely due to the event-plane decorrelation effect present at high  $p_T$  [17, 18, 65]. Finally, our calculations in Fig. 1–3 show that  $v_n$  are more sensitive to the energy loss models than  $R_{AA}$ .




FIG. 4. (Color online) Event-by-event correlation between  $D^0$   $v_2$  in the  $p_T$  range 8–13 GeV and the soft hadron  $v_2$  for two energy loss models in 30–50% PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. The solid lines stem from binning the heavy meson result by the soft hadron  $v_2$ .

Azimuthal anisotropy fluctuations can be systematically investigated using multi-particle cumulants [66, 67] which, in the present context of heavy-light flavor flow correlations, should give valuable information about the spectrum of path length fluctuations of heavy quark jets in the medium. However, it is not clear whether current statistics allows for proper measurement of higher order heavy flavor cumulants at high  $p_T$  at LHC (or even  $v_3\{2\}$  with small enough error bars). Because we oversample each event we have enough statistics to compute multi-particle cumulants such as  $v_2\{4\}$  which measures the correlation between a heavy flavor candidate and 3 soft particles [18]. For the 30–50% centrality class and  $p_T$  range 8–40 GeV we find that  $v_2\{4\}/v_2\{2\} \sim 0.95$  within statistical error bars regardless of variations in the energy loss,  $T_d$ , and quark flavor. A similar value, in the case of light flavor jets, was reported in [18]. The  $v_2\{4\}/v_2\{2\}$  ratio is related to the variance of the  $v_2^2(p_T)$  distribution, which reflects the event-by-event fluctuations in the hard sector due to the initial density fluctuations within our model. Therefore, this ratio probes

the magnitude of the initial density fluctuations and the role they play in the energy loss process. A more detailed study, involving the centrality dependence of  $v_2\{4\}/v_2\{2\}$  and also the calculation of even higher order heavy flavor cumulants will be presented elsewhere.




FIG. 5. (Color online) Correlation between  $v_n^{(heavy)}\{2\}$  for  $p_T$  in 8–13 GeV and  $v_n^{(soft)}$  for  $D^0$  in 30–50% PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV using two energy loss models. The bands represent the variation of the decoupling temperature in  $120 \text{ MeV} \leq T_d \leq 160 \text{ MeV}$ .

We now propose a new observable that encodes the event-by-event fluctuations of heavy flavor  $v_n$  at high- $p_T$  that does not require the challenging high statistics needed in multi-particle cumulants analyses. For a given hydrodynamic event characterized by a soft hadron  $v_n\{2\}$ , we calculate the corresponding coefficient for  $D^0$  in each event (this would be akin to the probability that a certain soft event corresponds to this particular heavy flavor  $v_n$ ), which is shown in the scatter plot in Fig. 4 for  $v_2$ . This plot shows that the soft and the heavy elliptic flow are correlated event-by-event within a given centrality class. Experimentally, one can bin the soft  $v_2\{2\}$  and calculate the corresponding heavy meson  $v_2$  for those set of events, as was done for high  $p_T$  identified hadrons by ATLAS [68]. This type of event-shape engineering procedure [69] gives rise to the solid black lines in Fig. 4. If there were no  $v_2$  fluctuations in the heavy flavor sector one would see a flat, horizontal line. Rather, our calculations predict a clear linear correlation between the heavy meson  $v_2$  and the elliptic flow of all charged hadrons.

This correlation is investigated in detail in Fig. 5 for  $v_2$  (top panel) and  $v_3$  (bottom panel) where we binned soft vs. heavy  $v_n$  and varied the energy loss model as well as  $T_d$ . Similar separations between the energy loss models are observed for the cumulants, in Figs. 2 and 3, and the correlations, though the latter are highly dependent on the integrated  $p_T$  range and should be investigated experimentally in order to determine an optimal choice given the error bars. Also, even though the widths of the  $T_d$  bands are equivalent,  $v_3$  is more sensitive to it than  $v_2$ . These results indicate that novel event-shape engineering techniques involving the flow of soft hadrons and heavy flavor will

be instrumental in determining the collective behavior of heavy quarks in the QGP.

#### IV. LINEAR SCALING WITH ECCENTRICITIES

While there are clear qualitative differences between the two energy loss models, the understanding of how a single energy loss model leads to the development of the final azimuthal anisotropy can be further explored. It is clear from Fig. 4 that a nearly linear relationship exists between  $v_n^{soft}$  and  $v_n^{heavy}$ . Considering that in the soft sector elliptical and triangular flow are developed from a nearly linear relationship between  $\varepsilon_n \rightarrow v_n^{soft}$ , one then expects that the relationship between  $\varepsilon_n \rightarrow v_n^{heavy}$  is also nearly linear.




FIG. 6. (Color online) Pearson coefficient described in Eq. (1) for  $Q_2$  and  $Q_3$  for  $dE/dx = \alpha$  and  $dE/dx = \xi T^2$  varying the decoupling temperature  $T_d = 120$ – $160$  MeV. Integrated  $v_n^{heavy}$  is used with the cut of  $8 < p_T [\text{GeV}] < 13$ .

In order to quantify this, we return to the Pearson coefficient in Eq. (1) and study  $Q_2$  and  $Q_3$  for the two different energy loss models as well as the range of  $T_d$ . In Fig. 6 one can see that, indeed,  $Q_2$  has a nearly linear relationship between  $\varepsilon_2 \rightarrow v_2^{heavy}$ . Central collisions do experience some non-linearities, which is to be expected since they are highly fluctuations-driven. Additionally, we find that  $dE/dx = \xi T^2$  has a more linear relationship between  $\varepsilon_2 \rightarrow v_2^{heavy}$  whereas  $dE/dx = \alpha$  has more non-linearities, especially in central collisions.

Triangular flow experiences more non-linearities between  $\varepsilon_3 \rightarrow v_3^{heavy}$  than elliptical flow does. Our results appear to be equivalent to the same effect seen in [18] for the order of magnitude of  $Q_2$  and  $Q_3$ .

We note also that the longer the heavy quarks are coupled to the medium, the more linear the relationship between  $\varepsilon_n \rightarrow v_n^{heavy}$ . For  $Q_2$  this is not a large effect. However,  $Q_3$  is strongly dependent on the decoupling temperature and this plays a larger role in determining the linearity than the energy loss model itself. This implies that  $v_2$  is a better constraint for  $dE/dx$  whereas  $v_3$  could provide more information about the time that heavy quarks are coupled to the QGP medium.

#### V. CONCLUSIONS

In summary, we presented the first theoretical calculations of  $D^0$  multi-particle flow cumulants in heavy ion collisions performed using realistic event-by-event viscous hydrodynamics. Our calculations with the constant energy loss model  $dE/dx \sim \alpha$  are consistent with the published  $R_{AA}$ ,  $v_2$ , and  $v_3$  for PbPb run 2 LHC data for  $p_T = 8$ – $40$  GeV, whereas the temperature dependent energy loss model  $dE/dx \sim \xi T^2$  underestimates the  $v_2$  data. We computed for the first time the heavy flavor 4-particle cumulant  $v_2\{4\}$ , which paves the way for experimental and theoretical studies of heavy flavor elliptic flow fluctuations.

Hydrodynamic viscosity was constrained here by soft bulk flow modeling and further investigation is needed to gauge its effect on our analysis. In [18] it was observed that an increase from 0.05 to 0.12 in  $\eta/s$  change cumulants by at most 5%. Energy loss fluctuations, though not shown in this paper, have been considered following an approach similar to [37] and we observed that reasonable fluctuations affect the results for the azimuthal coefficients by a few percent. Further analysis on energy loss fluctuations will be presented elsewhere.



The linear correlation predicted here between the  $D^0$   $v_n$  and the underlying soft hadron  $v_n$  provides a novel signature of collectivity in the heavy flavor sector event-by-event. Experimental confirmation of this behavior requires extending the current cutting edge event-shape engineering techniques [68, 70] to consider soft-heavy correlations, which should be feasible during the LHC run 2 and 3. This would not only allow for a comparison between the azimuthal anisotropies of heavy quarks and charged hadrons on an event-by-event basis but also give new insight into how the ubiquitous quantum mechanical fluctuations present in the initial state affect the energy loss experienced by heavy quarks in the plasma.

## ACKNOWLEDGMENTS

The authors thank Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for support. C.A.G.P. is supported by the NSFC under grant No. 11521064, MOST of China under Project No. 2014CB845404. J.N.H. was supported by the National Science Foundation under grant № PHY-1513864 and she acknowledges the use of the Maxwell Cluster and the advanced support from the Center of Advanced Computing and Data Systems at the University of Houston to carry out the research presented here. J.N. thanks the University of Houston and Rutgers University for their hospitality.

- 
- [1] M. Gyulassy and L. McLerran, *Quark-Gluon Plasma. New Discoveries at RHIC: Case of Strongly Interacting Quark-Gluon Plasma*, Nucl. Phys. **A750**, 30 (2005), arXiv:nucl-th/0405013.
  - [2] E. V. Shuryak, *Quark-Gluon Plasma. New Discoveries at RHIC: Case of Strongly Interacting Quark-Gluon Plasma*, Nucl. Phys. **A750**, 64 (2005), arXiv:hep-ph/0405066 [hep-ph].
  - [3] L. P. Csernai, J. Kapusta, and L. D. McLerran, Phys. Rev. Lett. **97**, 152303 (2006), arXiv:nucl-th/0604032.
  - [4] U. Heinz and R. Snellings, Annu. Rev. Nucl. Part. Sci. **63**, 123 (2013), arXiv:1301.2826 [nucl-th].
  - [5] J. Takahashi, B. M. Tavares, W. L. Qian, R. Andrade, F. Grassi, Y. Hama, T. Kodama, and N. Xu, Phys. Rev. Lett. **103**, 242301 (2009), arXiv:0902.4870 [nucl-th].
  - [6] B. Alver and G. Roland, Phys. Rev. **C81**, 054905 (2010), [Erratum: *ibid.* **C82**, 039903 (2010)], arXiv:1003.0194 [nucl-th].
  - [7] D. Teaney and L. Yan, Phys. Rev. **C83**, 064904 (2011), arXiv:1010.1876 [nucl-th].
  - [8] F. G. Gardim, F. Grassi, M. Luzum, and J.-Y. Ollitrault, Phys. Rev. **C85**, 024908 (2012), arXiv:1111.6538 [nucl-th].
  - [9] H. Niemi, G. S. Denicol, H. Holopainen, and P. Huovinen, Phys. Rev. **C87**, 054901 (2013), arXiv:1212.1008 [nucl-th].
  - [10] F. G. Gardim, J. Noronha-Hostler, M. Luzum, and F. Grassi, Phys. Rev. **C91**, 034902 (2015), arXiv:1411.2574 [nucl-th].
  - [11] D. Teaney and L. Yan, Phys. Rev. **C86**, 044908 (2012), arXiv:1206.1905 [nucl-th].
  - [12] H. Niemi, K. J. Eskola, and R. Paatelainen, Phys. Rev. **C93**, 024907 (2016), arXiv:1505.02677 [hep-ph].
  - [13] J. Noronha-Hostler, L. Yan, F. G. Gardim, and J.-Y. Ollitrault, Phys. Rev. **C93**, 014909 (2016), arXiv:1511.03896 [nucl-th].
  - [14] J. Qian and U. Heinz, Phys. Rev. **C94**, 024910 (2016), arXiv:1607.01732 [nucl-th].
  - [15] X.-N. Wang, Phys. Rev. **C63**, 054902 (2001), arXiv:nucl-th/0009019.
  - [16] M. Gyulassy, I. Vitev, and X. N. Wang, Phys. Rev. Lett. **86**, 2537 (2001), arXiv:nucl-th/0012092.
  - [17] J. Noronha-Hostler, B. Betz, J. Noronha, and M. Gyulassy, Phys. Rev. Lett. **116**, 252301 (2016), arXiv:1602.03788 [nucl-th].
  - [18] B. Betz, M. Gyulassy, M. Luzum, J. Noronha, J. Noronha-Hostler, I. Portillo, and C. Ratti, Phys. Rev. **C95**, 044901 (2017), arXiv:1609.05171 [nucl-th].
  - [19] B. Abelev *et al.* (ALICE Collaboration), Phys. Rev. Lett. **111**, 102301 (2013), arXiv:1305.2707 [nucl-ex].
  - [20] B. B. Abelev *et al.* (ALICE Collaboration), Phys. Rev. **C90**, 034904 (2014), arXiv:1405.2001 [nucl-ex].
  - [21] J. Adam *et al.* (ALICE Collaboration), J. High Energy Phys. **09**, 028 (2016), arXiv:1606.00321 [nucl-ex].
  - [22] G. D. Moore and D. Teaney, Phys. Rev. **C71**, 064904 (2005), arXiv:hep-ph/0412346 [hep-ph].
  - [23] M. Nahrgang, J. Aichelin, P. B. Gossiaux, and K. Werner, J. Phys. Conf. Ser. **668**, 012024 (2016).
  - [24] V. Greco, C. M. Ko, and P. Levai, Phys. Rev. Lett. **90**, 202302 (2003), arXiv:nucl-th/0301093.
  - [25] V. Greco, C. M. Ko, and P. Levai, Phys. Rev. **C68**, 034904 (2003), arXiv:nucl-th/0305024.
  - [26] R. J. Fries, B. Muller, C. Nonaka, and S. A. Bass, Phys. Rev. Lett. **90**, 202303 (2003), arXiv:nucl-th/0301087.
  - [27] V. Greco, C. M. Ko, and R. Rapp, Phys. Lett. **B595**, 202 (2004), arXiv:nucl-th/0312100.
  - [28] Y. Oh, C. M. Ko, S. H. Lee, and S. Yasui, Phys. Rev. **C79**, 044905 (2009), arXiv:0901.1382 [nucl-th].
  - [29] M. Nahrgang, J. Aichelin, S. Bass, P. B. Gossiaux, and K. Werner, Phys. Rev. **C91**, 014904 (2015), arXiv:1410.5396 [hep-ph].
  - [30] S. Cao, T. Luo, G.-Y. Qin, and X.-N. Wang, Phys. Rev. **C94**, 014909 (2016), arXiv:1605.06447 [nucl-th].
  - [31] A. Adil and I. Vitev, Phys. Lett. **B649**, 139 (2007), arXiv:hep-ph/0611109 [hep-ph].
  - [32] R. Brun and F. Rademakers, in *New Computing Techniques in Physics Research V*, Nuclear Instruments and Methods in Physics Research, Vol. A389, edited by M. Werlen and D. Perret-Gallix (North Holland, Amsterdam, 1997) pp. 81–86.
  - [33] T. Sjostrand, S. Mrenna, and P. Z. Skands, Comput. Phys. Commun. **178**, 852 (2008), arXiv:0710.3820 [hep-ph].
  - [34] M. Cacciari, M. Greco, and P. Nason, J. High Energy Phys. **05**, 007 (1998), arXiv:hep-ph/9803400 [hep-ph].

- [35] M. Cacciari, S. Frixione, and P. Nason, J. High Energy Phys. **03**, 006 (2001), arXiv:hep-ph/0102134 [hep-ph].
- [36] R. P. G. Andrade, J. Noronha, and G. S. Denicol, Phys. Rev. **C90**, 024914 (2014), arXiv:1403.1789 [nucl-th].
- [37] B. Betz and M. Gyulassy, J. High Energy Phys. **08**, 090 (2014), [Erratum: *ibid.* **10**, 043 (2014)], arXiv:1404.6378 [hep-ph].
- [38] R. Baier, A. H. Mueller, and D. Schiff, Phys. Lett. **B649**, 147 (2007), arXiv:nucl-th/0612068.
- [39] S. S. Gubser, Phys. Rev. **D74**, 126005 (2006), arXiv:hep-th/0605182 [hep-th].
- [40] R. Rougemont, A. Ficnar, S. Finazzo, and J. Noronha, J. High Energy Phys. **04**, 102 (2016), arXiv:1507.06556 [hep-th].
- [41] S. K. Das, F. Scardina, S. Plumari, and V. Greco, Phys. Lett. **B747**, 260 (2015), arXiv:1502.03757 [nucl-th].
- [42] H. van Hees, M. Mannarelli, V. Greco, and R. Rapp, Phys. Rev. Lett. **100**, 192301 (2008), arXiv:0709.2884 [hep-ph].
- [43] F. Riek and R. Rapp, Phys. Rev. **C82**, 035201 (2010), arXiv:1005.0769 [hep-ph].
- [44] J. Noronha-Hostler, G. S. Denicol, J. Noronha, R. P. G. Andrade, and F. Grassi, Phys. Rev. **C88**, 044916 (2013), arXiv:1305.1981 [nucl-th].
- [45] J. Noronha-Hostler, J. Noronha, and F. Grassi, Phys. Rev. **C90**, 034907 (2014), arXiv:1406.3333 [nucl-th].
- [46] J. Noronha-Hostler, J. Noronha, and M. Gyulassy, Phys. Rev. **C93**, 024909 (2016), arXiv:1508.02455 [nucl-th].
- [47] H. Marrochio, J. Noronha, G. S. Denicol, M. Luzum, S. Jeon, and C. Gale, Phys. Rev. **C91**, 014903 (2015), arXiv:1307.6130 [nucl-th].
- [48] H.-J. Drescher, A. Dumitru, A. Hayashigaki, and Y. Nara, Phys. Rev. **C74**, 044905 (2006), arXiv:nucl-th/0605012.
- [49] H.-J. Drescher and Y. Nara, Phys. Rev. **C76**, 041903 (2007), arXiv:0707.0249 [nucl-th].
- [50] H. J. Drescher and Y. Nara, Phys. Rev. **C75**, 034905 (2007), arXiv:nucl-th/0611017.
- [51] C. Peterson, D. Schlatter, I. Schmitt, and P. M. Zerwas, Phys. Rev. **D27**, 105 (1983).
- [52] M. Nahrgang, J. Aichelin, P. B. Gossiaux, and K. Werner, Phys. Rev. **C93**, 044909 (2016), arXiv:1602.03544 [nucl-th].
- [53] F. Becattini, M. Bleicher, T. Kollegger, T. Schuster, J. Steinheimer, and R. Stock, Phys. Rev. Lett. **111**, 082302 (2013), arXiv:1212.2431 [nucl-th].
- [54] E. L. Bratkovskaya, T. Song, H. Berrehrah, D. Cabrera, J. M. Torres-Rincon, L. Tolos, and W. Cassing, J. Phys. Conf. Ser. **668**, 012008 (2016), arXiv:1508.03887 [nucl-th].
- [55] T. Song, H. Berrehrah, D. Cabrera, W. Cassing, and E. Bratkovskaya, Phys. Rev. **C93**, 034906 (2016), arXiv:1512.00891 [nucl-th].
- [56] A. M. Poskanzer and S. A. Voloshin, Phys. Rev. **C58**, 1671 (1998), arXiv:nucl-ex/9805001.
- [57] M. Luzum and J.-Y. Ollitrault, Phys. Rev. **C87**, 044907 (2013), arXiv:1209.2323 [nucl-ex].
- [58] M. Luzum and H. Petersen, J. Phys. **G41**, 063102 (2014), arXiv:1312.5503 [nucl-th].
- [59] F. G. Gardim, F. Grassi, M. Luzum, and J. Noronha-Hostler, Phys. Rev. **C95**, 034901 (2017), arXiv:1608.02982 [nucl-th].
- [60] R. G. Miller, Biometrika **61**, 1 (1974).
- [61] CMS Collaboration, *D<sup>0</sup> Meson Nuclear Modification Factor in PbPb Collisions at  $\sqrt{s_{NN}} = 5.02$  TeV*, Tech. Rep. CMS-PAS-HIN-16-001 (CERN, Geneva, 2016).
- [62] Y. Akamatsu, T. Hatsuda, and T. Hirano, Phys. Rev. **C79**, 054907 (2009), arXiv:0809.1499 [hep-ph].
- [63] W. A. Horowitz and M. Gyulassy, Phys. Lett. **B666**, 320 (2008), arXiv:0706.2336 [nucl-th].
- [64] CMS Collaboration, *D<sup>0</sup> Meson  $v_n$  Harmonics in PbPb Collisions at 5.02 TeV*, Tech. Rep. CMS-PAS-HIN-16-007 (CERN, Geneva, 2016).
- [65] J. Jia, Phys. Rev. **C87**, 061901 (2013), arXiv:1203.3265 [nucl-th].
- [66] A. Bilandzic, R. Snellings, and S. Voloshin, Phys. Rev. **C83**, 044913 (2011), arXiv:1010.0233 [nucl-ex].
- [67] A. Bilandzic, C. H. Christensen, K. Gulbrandsen, A. Hansen, and Y. Zhou, Phys. Rev. **C89**, 064904 (2014), arXiv:1312.3572 [nucl-ex].
- [68] G. Aad *et al.* (ATLAS Collaboration), Phys. Rev. **C92**, 034903 (2015), arXiv:1504.01289 [hep-ex].
- [69] J. Schukraft, A. Timmins, and S. A. Voloshin, Phys. Lett. **B719**, 394 (2013), arXiv:1208.4563 [nucl-ex].
- [70] J. Adam *et al.* (ALICE Collaboration), Phys. Rev. **C93**, 034916 (2016), arXiv:1507.06194 [nucl-ex].