

This is the accepted manuscript made available via CHORUS. The article has been published as:

Effects of longitudinal asymmetry in heavy-ion collisions

Rashmi Raniwala, Sudhir Raniwala, and Constantin Loizides

Phys. Rev. C **97**, 024912 — Published 23 February 2018

DOI: [10.1103/PhysRevC.97.024912](https://doi.org/10.1103/PhysRevC.97.024912)

Effects of Longitudinal Asymmetry in Heavy-Ion Collisions

Rashmi Raniwala,^{1,*} Sudhir Raniwala,^{1,†} and Constantin Loizides^{2,‡}

¹*Physics Department, University of Rajasthan, Jaipur, India*

²*Lawrence Berkeley National Laboratory, Berkeley, California, USA*

(Dated: February 5, 2018)

In collisions of identical nuclei at a given impact parameter, the number of nucleons participating in the overlap region of each nucleus can be unequal due to nuclear density fluctuations. The asymmetry due to the unequal number of participating nucleons, referred to as longitudinal asymmetry, causes a shift in the center of mass rapidity of the participant zone. The information of the event asymmetry allows us to isolate and study the effect of longitudinal asymmetry on rapidity distribution of final state particles. In a Monte Carlo Glauber model the average rapidity-shift is found to be almost linearly related to the asymmetry. Using toy models, as well as Monte Carlo data for Pb–Pb collisions at 2.76 TeV generated with HIJING, two different versions of AMPT and DPMJET models, we demonstrate that the effect of asymmetry on final state rapidity distribution can be quantitatively related to the average rapidity shift via a third-order polynomial with a dominantly linear term. The coefficients of the polynomial are proportional to the rapidity shift with the dependence being sensitive to the details of the rapidity distribution. Experimental estimates of the spectator asymmetry through the measurement of spectator nucleons in a Zero Degree Calorimeter may hence be used to further constrain the initial conditions in ultra-relativistic heavy-ion collisions.

PACS numbers:

I. INTRODUCTION

In collisions of heavy ions the geometrically overlapping region created by interacting nucleons from each nucleus is called the participant zone. Even at fixed impact parameter, the number of participating nucleons from each nucleus fluctuates around the mean due to fluctuations in the positions of the nucleons around the mean nuclear density profile. Event-by-event, the participant zone therefore has a net non-zero momentum in the nucleon-nucleon centre-of-mass (CM) frame, and hence its rapidity is shifted with respect to the CM frame, and is denoted by y_0 [1].

Experimental data and simulated data from different event generators show that the total produced particle multiplicity, measured over a wide phase space region, scales approximately with the number of participants [2, 3]. The observed N_{part} scaling indicates that the number of participants or wounded nucleons is a relevant parameter affecting the production and distribution of produced particles even at LHC energies. Fluctuations of the fireball shape in the longitudinal direction are expected to create nontrivial rapidity correlations, as explicitly demonstrated using the wounded nucleon model [4]. The different components of the fluctuating fireball shape suggested in [4], have been recently extracted from the measured two-particle rapidity correlations [5].

The distribution of charged particles averaged over a large number of events in collisions of identical nu-

clei is observed to be symmetric about the rapidity of the nucleon-nucleon CM frame. The observed forward backward asymmetry in collisions of non-identical nuclei d–Au [6] and p–Pb [7] has been argued to be due to shift of the rapidity of the participant zone. Unequal number of participants in a collision of two identical nuclei produces a shift of the participant zone. This shift can be observed by measuring asymmetry in the energy of the Zero Degree Calorimeters on either side of the interaction vertex, even though its estimate in central collisions is marred by large relative fluctuations; the measurement of shift may facilitate to separate effect of fluctuations on various observables [8]. Preliminary results of the ALICE collaboration show a difference between pseudorapidity distribution of charged particles in Pb–Pb collision events of different asymmetries, as estimated from the measurement of spectator neutrons in the neutron Zero Degree Calorimeters [9]. Simulations based on a fluid dynamical framework demonstrate the effect of longitudinal fluctuations on the azimuthal anisotropy coefficients and their rapidity dependence; a significant decrease in the values of $v_1(y)$, and a wide plateau like behaviour for $v_2(y)$, both near midrapidity, has been estimated due to the event-by-event fluctuations affecting the rapidity of the participant zone, the latter being a conserved quantity [10]. The similar transverse-momentum dependence of the rapidity-even directed flow and the corresponding estimate from two-particle correlations at mid-rapidity indicate a weak correlation between fluctuating participant and spectator symmetry planes and suggest the possibility of using the spectator nucleons to further determine and constrain the effect of initial conditions [11]. This possibility has recently been further explored by model studies using AMPT [12].

It has been argued that the vorticity arising due to ini-

*Electronic address: Rashmi.Raniwala@cern.ch

†Electronic address: raniwalasudhir@gmail.com

‡Electronic address: Constantin.Loizides@cern.ch

tial state angular momentum may survive the evolution process in a low viscosity state of quark-gluon plasma and may manifest in the rapidity dependence of directed flow, $v_1(y)$, however its observability is affected by initial state fluctuations necessitating the requirement to determine the event-by-event centre of mass [13]. The need for determination of the event-by-event centre of mass rapidity is also highlighted by the possible observation of Λ -polarisation in the CM frame, which may confirm attainment of local thermodynamical equilibrium and persistence of vorticity until freezeout of the expanding matter [14, 15]. The observation of Λ -polarisation at the Relativistic Heavy Ion Collider requires that the models describing the evolution of a heavy-ion collisions incorporate the effect of large vorticity, thereby providing a complete characterisation of the system necessary to understand the dynamics of quarks and gluons in extreme conditions [16].

In the present work, we investigate the possible effect of the net (non-zero) momentum of the participant zone on experimentally measurable distributions of produced particles by exploring possible correlations between the participant asymmetry and the distribution of particles in the kinematic phase space. Events of the same net-momentum can be selected by classifying events on the basis of measured asymmetry in spectators for any centrality. The rapidity distribution of events of any asymmetry class is studied relative to corresponding distributions of another asymmetry class. The method has the advantage that most experimental uncertainties and corrections affecting the single particle distributions are cancelled. All variables used to develop this analysis can be estimated experimentally.

The paper is organized as follows. The rapidity-shift of the participant zone due to asymmetry of the event is discussed in Section II. The effect of a rapidity-shift is estimated using a toy model on a Gaussian rapidity distribution and is discussed in Section III. Section IV discusses the effect on various charged particle rapidity distributions for variable rapidity-shifts as calculated using Glauber model. The results of the present work are summarized in Section V.

II. RAPIDITY SHIFT OF PARTICIPANT ZONE

If the number of nucleons participating from the two colliding nuclei is A and B , respectively, then the participant zone has a net momentum in the nucleon-nucleon CM frame. The net momentum corresponds to a shift in the rapidity of the participant zone, which can be approximated as

$$y_0 \cong \frac{1}{2} \ln \frac{A}{B}. \quad (1)$$

Assuming each of the A (B) nucleons has a fixed momentum p ($-p$), Eq. 1 is obtained using the sum of four-momentum vectors $(0, 0, Ap, AE)$ and $(0, 0, -Bp, BE)$,

Centrality	b_{\min} (fm)	b_{\max} (fm)	$\langle N_{\text{part}} \rangle$	$ y_0 $
0–5%	0.0	3.61	383.9	0.0144
5–10%	3.61	5.12	330.0	0.0263
10–15%	5.12	6.27	280.2	0.0352
15–20%	6.27	7.24	236.5	0.0431
20–25%	7.24	8.09	198.5	0.0512
25–30%	8.09	8.86	165.5	0.0589
30–35%	8.86	9.57	136.5	0.0678
35–40%	9.57	10.23	110.2	0.0777
40–45%	10.23	10.85	88.6	0.0887
45–50%	10.85	11.43	70.2	0.1012
50–55%	11.43	11.99	54.6	0.1155
55–60%	11.99	12.52	41.6	0.1326
60–65%	12.52	13.03	31.0	0.1528
65–70%	13.03	13.52	22.5	0.1764

TABLE I: Centrality classes defined by impact parameter as well as corresponding $\langle N_{\text{part}} \rangle$ and $\langle |y_0| \rangle$ values.

with $E^2 = m_0^2 + p^2$, and neglecting $m_0 \ll p$. Since at the LHC in the TeV scale, $m_0/p < 10^{-6}$, we replace the ' \cong ' sign by the equality sign hitherto.

Defining the asymmetry of participants for each event as $\alpha_{\text{part}} = \frac{A-B}{A+B}$, the rapidity-shift y_0 can be written as

$$y_0 = \frac{1}{2} \ln \frac{1 + \alpha_{\text{part}}}{1 - \alpha_{\text{part}}}, \quad (2)$$

and has a unique correspondence with α_{part} . For small α_{part} , the shift follows $y_0 \cong \alpha_{\text{part}}$. The unequal number of nucleons in the participant zone imply unequal number of spectators of the two colliding nuclei, $N - A$ and $N - B$, respectively, where N is the total number of nucleons in each nucleus. The spectator asymmetry $\alpha_{\text{spec}} = \frac{(N-A)-(N-B)}{(N-A)+(N-B)} = \frac{B-A}{2N-(A+B)}$ is related to the participant asymmetry via $\alpha_{\text{spec}} = -\alpha_{\text{part}} \frac{A+B}{2N-(A+B)}$. Finally, the rapidity shift y_0 is related to the spectator asymmetry as

$$y_0 = \frac{1}{2} \ln \frac{(A+B)(1 + \alpha_{\text{spec}}) - 2N\alpha_{\text{spec}}}{(A+B)(1 - \alpha_{\text{spec}}) + 2N\alpha_{\text{spec}}} \quad (3)$$

which is accessible to experiment. Unlike the unique correspondence between α_{part} and y_0 , the presence of the $(A+B)$ term in Eq. 3 leads to a distribution of y_0 for a given value of α_{spec} , even at fixed impact parameter or centrality.

The rapidity-shift y_0 (Eq. 1) as well as its dependence on α_{part} (Eq. 2) and α_{spec} (Eq. 3) can be calculated within a Monte Carlo Glauber (MCG) framework [17], as implemented in Refs. [18, 19] or in Ref. [20]. In the present work, we have generated 1.2 million minimum bias events of Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV using HIJING (v1.383) [20] with default settings. The generated impact parameter (b) is used to define the event centrality. Limits on b used for 5%-wide centrality intervals with corresponding $\langle N_{\text{part}} \rangle$ are provided in Tab. I. The mean number of participants provide an estimate of




FIG. 1: The distributions of the participant-zone rapidity shift y_0 (left panel), of the participant asymmetry α_{part} (middle panel) and of the spectator asymmetry α_{spec} for 0–5%, 20–25% and 40–45% Pb–Pb centrality classes calculated with the HIJING MCG.

the order of magnitude of the rapidity-shift. Assuming that the number of nucleons from each of the two nuclei fluctuate by their root mean square while retaining the total number to be equal to N_{part} , the resulting value of rapidity shift for the most central class of events would be ≈ 0.05 . In practice, events in any centrality class will have a distribution peaked at zero. The distributions of y_0 , α_{part} and α_{spec} calculated with the HIJING MCG are shown in Fig. 1 for three centrality classes along with a Gaussian fit for each. The width of the y_0 distribution increases with decreasing centrality, i.e. for larger impact parameters the relative fluctuations increase since




FIG. 2: Event-by-event distributions of y_0 versus α_{part} (left panel) and y_0 versus α_{spec} (right panel) for the 20–25% Pb–Pb centrality class calculated with the HIJING MCG.

Figure 2 displays event-by-event distributions of y_0 versus α_{part} and α_{spec} , respectively, obtained for the 20–25% Pb–Pb centrality class with the HIJING MCG. The unique correspondence between y_0 and α_{part} is illustrated in the left panel of Fig. 2. The lack of a unique relation

the number of participants decreases. Events can be classified according to their rapidity-shift: $y_0 < 0$ are events of negative asymmetry (*-asym*) and $y_0 > 0$ are events of positive asymmetry (*+asym*). For each centrality class, the mean value of $|y_0|$ is also reported in Tab. I. The increase of $|y_0|$ with decreasing collision centrality is in agreement with results obtained earlier [1]. The α_{part} distributions are nearly identical to the y_0 distributions, while the α_{spec} distributions are different. The widths of the α_{spec} distributions increase with increasing centrality, i.e. the relative fluctuations increase with decreasing number of spectator nucleons.

between y_0 and α_{spec} due to the presence of the $(A + B)$ term in Eq. 3 leads to a distribution of y_0 for a given value of α_{spec} , even at fixed impact parameter or centrality, as illustrated in the right panel of Fig. 2. It can be regarded as the response matrix to obtain the values of $\langle y_0 \rangle$ for a given range of α_{spec} . The mean values $\langle y_0 \rangle$ as a function of the α_{spec} asymmetry are shown in Fig. 3 for three different centralities for 0–5%, 20–25% and 40–45% Pb–Pb centrality classes calculated with the HIJING MCG.

If the experiments could measure the number of nucleons in the participant zone, A and B , the participant-zone rapidity shift y_0 could be determined for each collision. However, neither A and B , nor α_{part} is directly amenable to experimental measurement. The asymmetry α_{spec} can be estimated by measuring the number of spectator nucleons through their energy deposited in the zero degree calorimeters on either side of the interaction vertex in collider experiments [8]. Using unfolding methods and the estimated values of α_{spec} , one can obtain an estimate of $\langle y_0 \rangle$, e.g. by using the response matrix in the right panel of Fig. 2. Almost all estimates of y_0 based on Glauber like models will show similar results even if there are differences in details. The ALICE experiment has determined a response matrix using information on the number of neutrons in the spectator and also us-




FIG. 3: The mean participant-zone rapidity shift $\langle y_0 \rangle$ versus α_{spec} for 0-5%, 20-25% and 40-45% Pb-Pb centrality classes calculated with the HIJING MCG.

ing the energy deposited in the Zero Degree Calorimeter, where a Glauber Monte Carlo model has been tuned to reproduce the energy distributions in neutron Zero Degree Calorimeters [21]

III. CONSTANT RAPIDITY SHIFT AND GAUSSIAN CHARGED PARTICLE RAPIDITY DISTRIBUTION

The measured rapidity distribution of charged particles produced in collisions of identical nuclei can be described by distributions which are symmetric about the CM rapidity [2, 3]. A Gaussian form is amongst the more common distributions used to describe data. We assume that the particles produced in asymmetric collisions of identical nuclei are also distributed symmetrically in the CM frame of the participant zone. Considering that the rapidity of the participant zone is shifted by y_0 from the rapidity of the nucleon-nucleon CM system, a symmetric distribution in the participant zone will appear as a shifted distribution in the nucleon-nucleon CM frame, which is also the laboratory frame for most collider experiments. For fixed y_0 , the rapidity distributions of produced particles can be written as a Gaussian distribution of width σ .

$$\frac{dN}{dy} = N_0 \exp\left(-\frac{(y - y_0)^2}{2\sigma^2}\right). \quad (4)$$

For symmetric collisions $y_0 = 0$, while for longitudinally asymmetric collisions y_0 is finite. The positive and negative values of y_0 correspond to the net momentum of the participant zone in the positive and negative direction, respectively, causing positive and negative participant (α_{part}) asymmetries, respectively. Taking the ra-

tio of single particle rapidity distributions of different asymmetry classes will eliminate the uncertainties arising due to experimental corrections and fluctuations affecting event-by-event distribution. The ratio of the rapidity distribution of particles in collisions with positive asymmetry to the distribution in collisions of negative asymmetry yields

$$\begin{aligned} \frac{\left(\frac{dN}{dy}\right)_{+\text{asym}}}{\left(\frac{dN}{dy}\right)_{-\text{asym}}} &= \exp\left(\frac{4yy_0}{\sigma^2}\right) \\ &= \sum_{n=0}^{\infty} c_n^g(y_0, \sigma) y^n, \end{aligned} \quad (5)$$

where $c_n^g = (4y_0/\sigma^2)^n/n!$ are the coefficients of the Taylor expansion of the exponential function, and the superscript g of the coefficients stands for the Gaussian shape of the parent rapidity distribution. The coefficients de-




FIG. 4: Gaussian $\frac{dN}{dy}$ distributions for $\sigma = 3.6$ shifted by $y_0 \pm 0.1$ obtained by the toy model (top panel). Ratio of the $\frac{dN}{dy}$ distributions fitted to a third-order polynomial (bottom panel). The coefficients are 0.015, 1.1×10^{-4} and 2.2×10^{-7} , respectively with $\chi^2/\text{dof} = 125/117$.

pend upon the parameters of the parent rapidity distribution and on the rapidity shift y_0 . These parameters are effectively fixed by selecting events on the basis of centrality and asymmetry. For typical values of y_0 equal to 0.1 and σ equal to 3.6, the ratio $\frac{c_2^g}{c_1^g} \sim 0.015$ and the ratio $\frac{c_3^g}{c_1^g} \sim 0.00015$ with subsequent terms having negligible contribution to the values of the function describing the ratio of the two rapidity distributions. Hence, for a Gaussian rapidity distribution, where the relation between the shift of the participant-zone rapidity and the coefficients c_n are analytically known, the dominant contribution can be expected for the linear term. The linear

term is related to the rapidity-shift via $y_0 = \frac{c_1^g \sigma^2}{4}$.




FIG. 5: The dependences of c_1^g , c_2^g and c_3^g on y_0 for fixed $\sigma = 3.6$ obtained by the toy model. Dashed lines correspond to analytical functions of the coefficients from Eq. 5.

These results have been validated using a toy model simulation by generating Gaussian rapidity distributions which are shifted by a constant magnitude. The top panel of Fig. 4 shows two rapidity distributions for $\sigma = 3.6$ shifted by $y_0 \pm 0.1$. The unshifted distribution would obviously lie in between the two distributions and is not drawn. The bottom panel of Fig. 4 shows the ratio of the $\frac{dN}{dy}$ distributions for events with $y_0 = +0.1$ to those with $y_0 = -0.1$, and the ratio is fitted to a third-order polynomial. The coefficient of the linear term is dominant with the other coefficients smaller by 2 and 4 orders of magnitude, respectively, as expected from Eq. 5. The same calculation is repeated for different values of the shift y_0 to obtain the coefficients of the third-order polynomial fit as function of y_0 , as shown in Fig. 5. The dependence of the coefficients on the rapidity-shift known from Eq. 5 is indicated with dashed lines, and agrees very well with the numerical calculation. The figure demonstrates that the dominant contribution to the $\frac{dN}{dy}$ ratio arises from the first coefficient, which is linearly related to the rapidity shift y_0 .

IV. VARIABLE SHIFT AND VARIOUS CHARGED PARTICLE RAPIDITY DISTRIBUTIONS

In the previous section we discussed the relation of coefficients of the third-order polynomial fit to the ratio of the $\frac{dN}{dy}$ distributions, assuming that the $\frac{dN}{dy}$ distributions are Gaussian in nature, and that all events have the same value of y_0 . In the following, the values of y_0 for different events are chosen according to the distribution of y_0 from the HIJING MCG, as shown for a few centrality classes in

the left panel of Fig. 1. As before, the rapidity distribution of particles $\frac{dN}{dy}$ are generated using a toy simulation taking into account event-by-event the rapidity-shift y_0 .

In addition to a Gaussian shape for the rapidity distribution, inspired by the possibility of a double Woods-Saxon distribution [22], or a Woods-Saxon-like distribution [2], we also consider that the rapidities of produced particles can be described by a Woods-Saxon distribution.

$$\frac{dN}{dy} = N_0 \frac{1}{1 + \exp\left(\frac{|(y-y_0)|-a}{c}\right)} \quad (6)$$

where $y_0 = 0$ for symmetric events, positive for events with positive asymmetry, and negative for events with negative asymmetry. Taking the ratio of the $\frac{dN}{dy}$ for events of opposite asymmetry, and making a Taylor expansion about $y = 0$ yields a polynomial in y , which can be written as

$$\left(\frac{dN}{dy}\right)_{+\text{asym}} = \sum_{n=0}^{\infty} c_n^{\text{ws}}(y_0, a, c) y^n. \quad (7)$$

The coefficients c_n^{ws} depend on the shift in rapidity and the parameters of the Woods-Saxon distribution in a non-trivial way. For a given set of parameters, however, the dependence on the rapidity-shift y_0 can be computed numerically.

We investigate the systematic effect of the rapidity-shift on the coefficients characterising the ratio of Gaussian and Woods-Saxon rapidity distributions for different asymmetry classes. Using the parameterized form of experimental η and p_T distributions in conjunction with the relative yield of pions, kaons and protons [3, 23], toy-model simulations provide the rapidity distribution. This resulting $\frac{dN}{dy}$ distribution is fitted once to a Gaussian form and then to a Woods-Saxon form to obtain the values of the parameters. Using these values of the parameters and the y_0 distribution obtained from HIJING events (Fig. 1), the rapidity distributions are obtained separately for positive and negative values of y_0 corresponding to different centrality classes. The ratio of the two distributions is fitted to a third order polynomial as shown in Fig. 6 for four centrality classes. In each case, the χ^2/dof are the smallest in the most central class and are about 0.64 and 0.83 for Gaussian and Woods-Saxon rapidity distribution.

To investigate the possible contribution originating from the dynamics of the particle production mechanism, or in general, of any final state effects, the charged-particle rapidity distribution obtained from some commonly used event generators have also been used. We have generated about 1.2 million events of HIJING and about 1 million events each of both versions of AMPT (Default and String Melting) [24], and of DPMJET [25] for Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV. The rapidity shift y_0 is determined for each generated event from




FIG. 6: The ratio of $\frac{dN}{dy}$ distributions for events with positive ($y_0 > 0$) and with negative ($y_0 < 0$) asymmetry. The different panels are for rapidity distributions corresponding to (a) parametrised Gaussian form (b) parametrised Woods-Saxon form (c) HIJING (d) AMPT (Default) (e) AMPT (String Melting) and (f) DPMJET (default). The ratios, along with fits to third order polynomials, are shown for four centrality classes.

the number of participating nucleons from each of the two colliding nuclei. The average rapidity distributions corresponding to positive and negative values of y_0 are obtained. The ratio of these two rapidity distributions is fitted to a third order polynomial to obtain the values of the coefficients. The ratios and the fits are shown in Fig. 6, along with the results for toy model simulations for parametrised rapidity distributions. The differences in the parent rapidity distributions manifest themselves in the ratios and the values of the coefficients in the polynomial.

For all rapidity distributions, the coefficients for the quadratic and cubic terms in the polynomial fitting of the ratio are much smaller than those of the linear term. The dependence of the first three coefficients c_n on the mean rapidity shift $\langle |y_0| \rangle$ is shown in Fig. 7. The six different rapidity distributions yield different dependence of the coefficients on $\langle |y_0| \rangle$, indicating a possibility of determining the details of the rapidity distribution from the knowledge of the behaviour of the coefficients.

V. SUMMARY

In collisions of identical nuclei at a given impact parameter, the number of nucleons participating in the overlap region of each nucleus, estimated with a Monte Carlo Glauber model, can be unequal due to nuclear den-




FIG. 7: The coefficients c_1 , c_2 , and c_3 of the third-order polynomial fit to the ratio of $\frac{dN}{dy}$ distributions versus $\langle |y_0| \rangle$ for the six forms of rapidity distributions. Polynomial fits are to guide the eye. The coefficients c_n demonstrate a dependence on $\langle |y_0| \rangle$. The rapidity distributions are obtained in 14 centrality intervals up to 70% centrality with corresponding $\langle |y_0| \rangle$ as given in Tab. I.

sity fluctuations (Fig. 1). The asymmetry due to the unequal number of participating nucleons causes a rapidity-shift of the participant zone (Fig. 2), which may be experimentally accessible by measuring the energy of the spectator nucleons, as was also argued in [8]. The average rapidity-shift has been found to be almost linearly related to the asymmetry (Fig. 3). The effect of the small rapidity-shift in the rapidity distributions was estimated by taking the ratio of the distributions of events of positive and negative asymmetries. The success of the method was demonstrated by using a toy model to illustrate that such a ratio can effectively be described by a third-order polynomial (Fig. 4), where the coefficients are related to the rapidity-shift and are shown for Gaussian rapidity distributions for constant rapidity-shift (Fig. 5). The effect on the ratio of rapidity distributions for positive and negative asymmetry has been systematically studied for Gaussian and Woods-Saxon particle rapidity distributions, using a toy model simulation and taking the distribution of y_0 from Fig. 1. The ratios and the polynomial fits are shown in Fig. 6 (a) and (b). The possible effect of dynamics of particle production is investigated using the ratio of rapidity distributions for positive and negative asymmetries from different event generators. The ratios and the fitted polynomials for rapidity distributions obtained from HIJING, AMPT and DPMJET are shown in Fig. 6 (c) to (f) for four centrality classes. The ratios are sensitive to the detailed shape of the parent rapidity distribution (Fig. 6), and can be quantitatively described by a third-order polynomial with a dominantly linear term (Fig. 7). The relation between coefficients and the rapidity shift confirm that the effect of initial state longitudinal asymmetry survives

through the particle production process and the subsequent evolution to the observed final state. Experimentally, estimates of the longitudinal asymmetry via measurements of the spectator asymmetry can be used to systematically investigate the influence of the longitudinal asymmetry on various observables, and hence may further constrain the initial conditions in ultra-relativistic heavy ion collisions. Recent results from the ALICE collaboration confirm that the longitudinal fluctuations affect the pseudorapidity distributions; the effect finds a simple explanation in terms of the rapidity shift of the participant zone, and shows a sensitivity to the shape of the rapidity distribution [21]. As indicated in Sec. I, determination of an event-by-event CM may be necessary for a complete description of the evolution of heavy-ion collisions, to separate the effects of initial state fluctuations from the dynamical evolution. Further attempts to devise methods for the determination of the CM of the participant zone in each event, using experimentally measurable quantities, is under investigation.

VI. ACKNOWLEDGEMENTS

We thank Jurgen Schukraft for fruitful discussions. R. Raniwala and S. Raniwala acknowledge the financial support of the Department of Science and Technology of the Government of India. The work of C. Loizides is supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under contract number DE-AC02-05CH11231.

-
- [1] V. Vovchenko, D. Anchishkin, and L. P. Csernai, “Longitudinal fluctuations of the center of mass of the participants in heavy-ion collisions,” *Phys. Rev. C* **88** no. 1, (2013) 014901, [arXiv:1306.5208 \[nucl-th\]](#).
 - [2] **PHOBOS** Collaboration, B. Alver *et al.*, “Phobos results on charged particle multiplicity and pseudorapidity distributions in Au–Au, Cu–Cu, d–Au, and pp collisions at ultra-relativistic energies,” *Phys. Rev. C* **83** (2011) 024913, [arXiv:1011.1940 \[nucl-ex\]](#).
 - [3] **ALICE** Collaboration, E. Abbas *et al.*, “Centrality dependence of the pseudorapidity density distribution for charged particles in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **726** (2013) 610–622, [arXiv:1304.0347 \[nucl-ex\]](#).
 - [4] A. Bzdak and D. Teaney, “Longitudinal fluctuations of the fireball density in heavy-ion collisions,” *Phys. Rev. C* **87** no. 2, (2013) 024906, [arXiv:1210.1965 \[nucl-th\]](#).
 - [5] J. Jia, S. Radhakrishnan, and M. Zhou, “Forward-backward multiplicity fluctuation and longitudinal harmonics in high-energy nuclear collisions,” *Phys. Rev. C* **93** no. 4, (2016) 044905, [arXiv:1506.03496 \[nucl-th\]](#).
 - [6] P. Steinberg, “Inclusive pseudorapidity distributions in $p(d) + A$ collisions modeled with shifted rapidity distributions,” *Submitted to: Phys. Lett. B* (2007), [arXiv:nucl-ex/0703002 \[NUCL-EX\]](#).
 - [7] G. Martinez-Garcia, “Particle rapidity distribution in proton-nucleus collisions using the proton-contributor reference frame,” [arXiv:1408.3108 \[hep-ph\]](#).
 - [8] L. P. Csernai, G. Eyyubova, and V. K. Magas, “New method for measuring longitudinal fluctuations and directed flow in ultrarelativistic heavy ion reactions,” *Phys. Rev. C* **86** (2012) 024912, [arXiv:1204.5885 \[hep-ph\]](#). [Erratum: *Phys. Rev. C* **88**, 019902(2013)].
 - [9] R. Raniwala, “Longitudinal asymmetry and its measurable effects in Pb–Pb collisions at 2.76 TeV,” in *25th International Conference on Ultra-Relativistic Nucleus-Nucleus Collisions (Quark Matter 2015) Kobe, Japan, September 27-October 3, 2015*. 2015. [arXiv:1512.08177 \[nucl-ex\]](#). <http://inspirehep.net/record/1411530/files/arXiv:1512.08177.pdf>.
 - [10] L. P. Csernai, V. K. Magas, H. Stocker, and D. D. Strottman, “Fluid Dynamical Prediction of Changed v_1 -flow at LHC,” *Phys. Rev. C* **84** (2011) 024914, [arXiv:1101.3451 \[nucl-th\]](#).

- [11] **ALICE** Collaboration, B. Abelev *et al.*, “Directed flow of charged particles at midrapidity relative to the spectator plane in Pb–Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV,” *Phys. Rev. Lett.* **111** no. 23, (2013) 232302, [arXiv:1306.4145 \[nucl-ex\]](#).
- [12] V. Bairathi, S. Chatterjee, M. Rihan Haque, and B. Mohanty, “Probing Pb+Pb collisions at $\sqrt{s_{NN}} = 2760$ GeV with spectators,” *Phys. Lett.* **B754** (2016) 144–150, [arXiv:1508.02338 \[nucl-th\]](#).
- [13] L. P. Csernai, V. K. Magas, and D. J. Wang, “Flow Vorticity in Peripheral High Energy Heavy Ion Collisions,” *Phys. Rev.* **C87** no. 3, (2013) 034906, [arXiv:1302.5310 \[nucl-th\]](#).
- [14] F. Becattini, L. Csernai, and D. J. Wang, “ Λ polarization in peripheral heavy ion collisions,” *Phys. Rev.* **C88** no. 3, (2013) 034905, [arXiv:1304.4427 \[nucl-th\]](#). [Erratum: *Phys. Rev.* **C93**, no. 6, 069901 (2016)].
- [15] Y. Xie, D. Wang, and L. P. Csernai, “Global Λ polarization in high energy collisions,” *Phys. Rev.* **C95** no. 3, (2017) 031901, [arXiv:1703.03770 \[nucl-th\]](#).
- [16] **STAR** Collaboration, L. Adamczyk *et al.*, “Global Λ hyperon polarization in nuclear collisions: evidence for the most vortical fluid,” *Nature* **548** (2017) 62–65, [arXiv:1701.06657 \[nucl-ex\]](#).
- [17] M. L. Miller, K. Reygers, S. J. Sanders, and P. Steinberg, “Glauber modeling in high energy nuclear collisions,” *Ann. Rev. Nucl. Part. Sci.* **57** (2007) 205–243, [arXiv:nucl-ex/0701025 \[nucl-ex\]](#).
- [18] B. Alver, M. Baker, C. Loizides, and P. Steinberg, “The PHOBOS Glauber Monte Carlo,” [arXiv:0805.4411 \[nucl-ex\]](#).
- [19] C. Loizides, J. Nagle, and P. Steinberg, “Improved version of the PHOBOS Glauber Monte Carlo,” *SoftwareX* **1-2** (2015) 13–18, [arXiv:1408.2549 \[nucl-ex\]](#).
- [20] X.-N. Wang and M. Gyulassy, “HIJING: A Monte Carlo model for multiple jet production in pp, pA and AA collisions,” *Phys. Rev.* **D44** (1991) 3501–3516.
- [21] **ALICE** Collaboration, S. Acharya *et al.*, “Longitudinal asymmetry and its effect on pseudorapidity distributions in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” [arXiv:1710.07975 \[nucl-ex\]](#).
- [22] K. J. Eskola, K. Kajantie, P. V. Ruuskanen, and K. Tuominen, “Rapidity dependence of particle production in ultrarelativistic nuclear collisions,” *Phys. Lett.* **B543** (2002) 208–216, [arXiv:hep-ph/0204034 \[hep-ph\]](#).
- [23] **ALICE** Collaboration, B. Abelev *et al.*, “Centrality dependence of π , K, p production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev.* **C88** (2013) 044910, [arXiv:1303.0737 \[hep-ex\]](#).
- [24] Z.-W. Lin, C. M. Ko, B.-A. Li, B. Zhang, and S. Pal, “A Multi-phase transport model for relativistic heavy ion collisions,” *Phys. Rev.* **C72** (2005) 064901, [arXiv:nucl-th/0411110 \[nucl-th\]](#).
- [25] S. Roesler, R. Engel, and J. Ranft, “The Monte Carlo event generator DPMJET-III,” in *Advanced Monte Carlo for radiation physics, particle transport simulation and applications. Proceedings, Conference, MC2000, Lisbon, Portugal, October 23-26, 2000*, pp. 1033–1038. 2000. [arXiv:hep-ph/0012252 \[hep-ph\]](#). <http://www-public.slac.stanford.edu/sciDoc/docMeta.aspx?slacPubNumber=SLAC-PUB-8740>.