



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Lifetime measurements in ^{156}Gd

A. Aprahamian, R. C. de Haan, S. R. Leshner, C. Casarella, A. Stratman, H. G. Börner, H. Lehmann, M. Jentschel, and A. M. Bruce

Phys. Rev. C **98**, 034303 — Published 6 September 2018

DOI: [10.1103/PhysRevC.98.034303](https://doi.org/10.1103/PhysRevC.98.034303)

Lifetime Measurements in ^{156}Gd

A. Aprahamian,^{1,*} R. C. de Haan,¹ S. R. Lesher,^{1,2} C. Casarella,¹

A. Stratman,¹ H. G. Börner,³ H. Lehmann,³ M. Jentschel,³ and A.M. Bruce⁴

¹*Department of Physics, University of Notre Dame, Notre Dame, IN 46556, USA*

²*Department of Physics, University of Wisconsin – La Crosse, La Crosse, WI 54601, USA*

³*Institut Laue-Langevin, F-38042, Grenoble, France*

⁴*University of Brighton, Brighton, BN2 4GJ, United Kingdom*

(Dated: August 13, 2018)

Abstract

Background: The nature of low-lying oscillations or excitations around the equilibrium deformed nuclear shape remains an open question in nuclear structure. The question revolves around the possible degrees of freedom in deformed nuclei. Rotational motion is an expected feature of deformed nuclei, the open challenge is whether the “granularity” of nuclei allows single or multiple quanta of vibrational oscillations or excitations superimposed on the equilibrium deformed shape of the nucleus. Special emphasis is placed on the $K^\pi = 0^+$, β vibration whose existence is open to debate some forty years after Bohr-Mottelson-Rainwater’s Nobel prize for connecting nucleon motion to the emergence of collectivity.

Purpose: The ^{156}Gd nucleus is an excellent test case for the search of the predicted oscillations since it has one of the most developed level schemes up to 2.35 MeV and it lies in the well deformed rare-earth region of the chart of nuclides. This nucleus has previously been studied by (n, γ) , (n, e^-) , (e, e') , (p, p') , (d, p) , and (d, t) reactions with six known excited $K^\pi = 0^+$ bands. We measured level lifetimes of ^{156}Gd in order to determine the nature of the low-lying excited bands.

Method: Lifetimes of excited states in the ^{156}Gd nucleus were measured following neutron capture using the GRID technique at the Institut Laue-Langevin (ILL) in Grenoble, France.

Results: Twelve level lifetimes were measured from four excitation bands in the ^{156}Gd nucleus including the lifetimes of three of the $K^\pi = 0^+$ bands.

Conclusions: There are two $K^\pi = 0^+$ bands in this nucleus connected to the ground state band. Transitions from the $K^\pi = 0_2^+$ band at 1049.5 keV to the ground state band are more collective than the ones from the $K^\pi = 0_3^+$ band at 1168.2 keV. The moments of inertia of the various $K^\pi = 0^+$, the $K^\pi = 2^+$, and the $K^\pi = 4^+$ bands show that all the bands except the $K^\pi = 0_3^+$ band at 1168.2 keV have nearly identical moments of inertia with the ground state band pointing to the fact that all of the bands discussed here with the exception of this indicating $K^\pi = 0_3^+$ band seem to be collective excitations built on the ground state. This result is consistent with various theoretical predictions. B(E2) calculations for transitions from the $K^\pi = 2^+$ band to the ground state band supports the assignment of this band as the γ band. Also, the $K^\pi = 0_4^+$ and the $K^\pi = 4_1^+$ bands at 1715.2 keV and 1510.6 keV, respectively, are shown to be strongly connected to the $K^\pi = 2^+$ γ band presenting evidence for the observation of a second set of two-phonon $\gamma\gamma$ vibrational excitations albeit with greatly varying degrees of anharmonicity in comparison to the case of ^{166}Er .

* aapraham@nd.edu

I. INTRODUCTION

The 1975 Nobel prize in Physics was awarded to Bohr, Mottelson, and Rainwater for the discovery of the connection between nucleon motion and the emergent collective behavior. Bohr-Mottelson-Rainwater described nuclei geometrically as a shape and the oscillations of the nucleus around that shape. The lowest lying shape effecting oscillations or vibrations would be quadrupole ($\lambda = 2$) in nature, resulting in two types of vibrations in deformed nuclei: β with oscillations along the symmetry axis ($K^\pi = 0^+$) and γ breaking axial symmetry with a projection of $K^\pi = 2^+$ on the symmetry axis. The γ vibration seems to be well characterized as the first $K^\pi = 2_1^+$ (or 2_γ^+) band and exhibits a systematic behavior across the region of deformed nuclei with typical $B(E2 : 2_\gamma^+ \rightarrow 0_{g.s.}^+)$ values of a few Weisskopf units (W.u.) [1].

Today, over forty years later, the existence and characterization of the low-lying β vibration remains an open question in nuclear structure [1–35]. This is due to the lack of sufficient experimental data on the identification and characterization of 0^+ excitations in deformed nuclei and to the interpretation of what is expected of a β vibration. The absence of a β vibrational excitation in deformed nuclei will call into question why nuclei, unlike all other quantum systems, do not exhibit this mode of oscillatory motion.

In well-deformed regions of nuclei, excitations built on a deformed ground state have traditionally been described in terms of quadrupole excitations leading to the classifications of the first excited 0^+ bands as single-phonon β -vibrational excitations. However, discussions in recent years have focused on a debate about the absence, or lack of, a ($K^\pi = 0^+$) β vibration with a multitude of possible interpretations including the possibility of phase changes at the onset of deformation (for example, at $N=90$ and $Z=64$) and the application of new symmetries to describe these nuclei [4, 36–41], or the change in the expectation of a β vibration. The discussions on the existence or absence of the $K^\pi = 0^+$ β vibrational excitations in nuclei have spanned a wide spectrum of possibilities from shape co-existence, where a competing shape is not the lowest favored shape but occurs low in the excitation spectrum of a given nucleus [2], to a redefinition of what can, in fact, be interpreted as a “ β ” vibration [42]. In the IBM [43–45], the first excited 0^+ and 2^+ bands are members of the same representation and in a pure $SU(3)$ limit would not decay to the ground state (g.s.). Most deformed nuclei however are not pure $SU(3)$ and the breaking of that symmetry

gives rise to interband transitions from both of the low-lying $K^\pi = 2^+$ and $K^\pi = 0^+$ (“ γ ” and “ β ”) bands of significant strengths. Another recent development describes nuclei at the point of phase change from spherical to deformed in terms of β and γ shape parameters, or the SU(3) symmetry [17, 18, 20] or the pseudo-SU(3) [35]. There is also the possibility of Partial Dynamical Symmetries where the SU(3) symmetry is obeyed by some of the states and broken in others [46]. A systematic theoretical study of even-even deformed nuclei in the Hartree-Fock-Bogoliubov approach extended by the generator coordinate method and mapped into a five-dimensional collective Hamiltonian for even-even nuclei from $Z = 10 - 110$, provides guidelines to distinguishing between coexistence and β vibrational oscillations. These studies and others [30, 32] on the nature of $K^\pi = 0^+$ bands in deformed nuclei predict widely varying levels of collectivity depopulating the first excited 0^+ states [1].

The Gd isotopes lie in a well deformed region of the chart of nuclides with the ratio of the first two excited states $4^+/2^+$ ($R_{4/2}$) varying from 3.0 to 3.3 for ^{154}Gd to ^{160}Gd allowing a fertile testing ground for previous theories. The focus of this paper is ^{156}Gd with an $R_{4^+/2^+}$ ratio of 3.24, a well known level scheme up to an excitation energy of 2.35 MeV and six excited $K^\pi = 0^+$ bands, four of them below the pairing gap at approximately 2 MeV. The ^{156}Gd nucleus has been studied extensively and with high precision using bent crystals GAMS 2/3 for (n,γ) , the BILL electron spectrometer for (n,e^-) measurements [47], and the early tests of the GRID (GammaRay Induced Doppler broadening) technique [48–50] using the GAMS4 spectrometer at the Institut Laue Langevin in Grenoble, and by (d,t) and (d,p) reactions at the Munich Tandem Accelerator [47]. Numerous other studies included electron, proton, and photon scattering [51, 52]. The ^{156}Gd nucleus was used for comparisons with early tests of the Interacting Boson Model numerical studies for the SU(3) limit [53] as well as the later tests of Partial Dynamical Symmetry tests [46]. The focus of this paper is the excited $K^\pi = 0^+$ bands. We have measured twelve level lifetimes in these bands in an attempt to characterize the nature of these states in search of the β and other low-lying vibrational excitations.

II. EXPERIMENT

The experiment was performed at the Institut Laue-Langevin (ILL) neutron High Flux Reactor in Grenoble, France. The ^{156}Gd nucleus was populated by neutron capture on a

5.349 g Gd₂O₃ (Gadolinium Oxide) target. The GRID technique [49, 54] of lifetime measurements is based on measuring the broadening of decay gamma-ray lines using perfect crystals to measure the associated γ ray wavelength. The broadening is due to the initial recoil velocity of the nucleus where the width of a given gamma-ray transition emitted in flight results from the competition between the slowing down process and the level lifetime. Knowing the slowing down process from simulations, the nuclear level lifetime is extracted. The recoil velocities are typically $10^{-4}c$ to $10^{-6}c$, resulting in a broadening of only a few eV. The γ -ray wavelengths rely on crystal diffraction from nearly perfect flat crystals made of silicon or germanium. GAMS4 is a double-flat-crystal spectrometer with remarkable energy resolution, and high precision of a few eV [50, 55–57]. The broadened γ -ray peaks were fitted using the code GRIDDLE [58]. Fig. 1 shows the broadening of the widths of gamma-ray transitions depopulating from two different levels at 1129.44 keV and 1248.01 keV in the ¹⁵⁶Gd nucleus.

The largest uncertainties in these measurements arise from the unknown feeding of the level of interest. Therefore, in cases where the feeding of a particular nuclear level is not completely understood, rather extreme and conservative assumptions have been made in order to extract conservative *upper* and *lower* limits. The upper limit of the extracted lifetime is determined assuming that the level is totally fed by cascades of γ -ray transitions from the compound capture state at 8.536 MeV. The cascades are typically several MeV unobserved transitions via two-step cascades connecting the level of interest to the compound state. This would yield the maximum broadening for the gamma-rays depopulating a given level or the longest slowing down times resulting in the longest possible lifetime for the level of interest. Lifetimes shorter than the upper limit would yield more collective B(E2) values. The lower limit of the lifetime is extracted by assuming that the missing feeding comes from the unplaced low energy transitions measured in this nucleus. These are extreme and conservative limits to the measured lifetimes. The more realistic scenario would probably lie somewhere in the middle of the lifetimes resulting from these intentionally extreme feeding assumptions. Table I lists the percentage of known feeding for the levels of interest as well the results of the calculations for the extraction of upper and lower limits on the lifetimes. The extracted lifetime ranges are listed in Table II along with a listing for comparison of six previously measured lifetimes [47, 59, 60] for the states of interest, four were measured using the GAMS4 spectrometer in its early incarnation [47] and two were measured by coulomb




FIG. 1. Broadening curves for two γ -ray transitions 1040.474 and 1159.000 keV depopulating levels at 1129.440 and 1248.008 keV, respectively. The dotted line is the instrumental response expected without broadening and the solid black line is the fit of the data points. The lifetime for a given transition is extracted from the observed broadening.

excitation [59, 60]. In all the cases where previous measurements existed, there is agreement. A $0.6 \times \tau_{max}$ was used as a calibration factor for figures which corresponds to the approximate middle of the range of the GRID measurement to match the lifetime of the 1154.2 keV level. A more detailed explanation on calibration is found in Ref. [61].

Figure 2 presents a partial level scheme of ^{156}Gd showing in red the levels whose lifetimes were measured in this work. A value of $0.6 \times \tau_{max}$ was used to calculate the $B(E2)$ values of the transitions for all the levels of interest that were measured with GRID. The $B(E2)$ values from the $K^\pi = 4^+$, $J^\pi = 4^+$ level was previously measured by coulomb excitation.

TABLE I. The energy levels and depopulating transitions measured in ^{156}Gd , the missing feeding of each level, and the upper and lower lifetime limits, resulting in the conservative overall range for the lifetimes.

$E_x(\text{keV})$	J^π	$E_\gamma(\text{keV})$	Feeding ^a	$\tau_{upper}(\text{ps})$	$\tau_{lower}(\text{ps})$	$\tau_{range}(\text{ps})$
$K^\pi = 0_2^+$ band at 1049.479 keV						
1049.479	0^+	960.510	44.3%	$2.86^{+1.88}_{-0.82}$	$2.52^{+1.59}_{-0.71}$	$1.81 < \tau < 4.75$
1129.440	2^+	1040.474	20.4%	$2.08^{+0.50}_{-0.34}$	$1.48^{+0.30}_{-0.22}$	$1.26 < \tau < 2.58$
1297.825	4^+	1009.622	40.5%	$2.52^{+0.84}_{-0.51}$	$1.84^{+0.54}_{-0.34}$	$1.49 < \tau < 3.36$
$K^\pi = 2^+$ band at 1154.151 keV						
1154.151	2^+	1065.182	43.7%	$1.20^{+0.14}_{-0.12}$	$0.96^{+0.11}_{-0.09}$	$0.87 < \tau < 1.35$
1248.008	3^+	1159.000	52.3%	$0.92^{+0.09}_{-0.08}$	$0.74^{+0.08}_{-0.06}$	$0.68 < \tau < 1.01$
1355.425	4^+	1067.236	34.8%	$0.91^{+0.10}_{-0.09}$	$0.64^{+0.07}_{-0.06}$	$0.58 < \tau < 1.02$
1506.868	5^+	1218.710	21.8%	$1.21^{+0.61}_{-0.31}$	$0.34^{+0.18}_{-0.10}$	$0.25 < \tau < 1.82$
$K^\pi = 0_3^+$ band at 1168.190 keV						
1168.190	0^+	1079.230	19.1%	$6.34^{+8.35}_{-2.31}$	$4.61^{+5.04}_{-1.60}$	$3.00 < \tau < 14.7$
1258.075	2^+	1169.092	14.4%	$3.46^{+1.60}_{-0.84}$	$3.17^{+1.84}_{-0.86}$	$2.31 < \tau < 5.06$
$K^\pi = 0_4^+$ band at 1715.181 keV						
1715.181	0^+	472.700	11.4%	$3.69^{+3.39}_{-1.21}$	1.88 ± 1.86	$0.02 < \tau < 7.08$
1771.089	2^+	1682.184	25.7%	$0.60^{+0.13}_{-0.10}$	$0.17^{+0.06}_{-0.04}$	$0.13 < \tau < 0.73$
1893.395	4^+	1605.217	1.3%	$0.37^{+0.08}_{-0.06}$	$0.0002^{+0.0001}_{-0.00005}$	$0.00015 < \tau < 0.45$

^aPercentage of known level feeding [47, 59].

III. RESULTS

The results of the lifetime measurements are shown in Table II with the uncertainties that have led to the broad ranges of lifetimes as described in the experimental section. Table III shows the levels of interest including lifetimes and their depopulating transitions for the first three excited $K^\pi = 0^+$ bands and from the $K^\pi=2^+$ γ band. This work presents the lifetime measurements of all four excited states of the first $K^\pi = 0^+$ band, the $K^\pi=2^+$ band, the $K^\pi = 0_2^+$, as well as two additional $K^\pi = 0^+$ bands. In each case, the measurements of the

TABLE II. Level energies and lifetime ranges for all twelve states measured in this work in comparison with previous measurements. Four of the previous values were measured by the GRID technique using the GAMS4 spectrometer [47], another three were measured by coulomb excitation [59, 60].

$E_x(\text{keV})$	$\tau_{grid}(\text{ps})$	Prev. Meas.(ps)
1049.479	$1.81 < \tau < 4.75$	$1.39 < \tau < 12.97$ [47]
1129.440	$1.26 < \tau < 2.58$	2.27 ± 0.17 [60]
1154.151	$0.87 < \tau < 1.35$	$0.87 < \tau < 1.44$ [47] 0.82 ± 0.03 [59]
1168.190	$3.00 < \tau < 14.69$	
1248.008	$0.68 < \tau < 1.01$	
1258.075	$2.31 < \tau < 5.06$	2.22 ± 0.22 [59]
1297.825	$1.49 < \tau < 3.36$	
1355.425	$0.58 < \tau < 1.02$	$1.15 < \tau < 7.71$ [47]
1506.868	$0.24 < \tau < 1.82$	
1715.181	$0.02 < \tau < 7.08$	
1771.089	$0.13 < \tau < 0.73$	$0.14 < \tau < 1.44$ [47]
1893.395	$0.00015 < \tau < 0.45$	

$K^\pi = 0^+$ bands include the lifetimes of the $J^\pi = 0^+$ bandheads. In the deformed rare earth region of the chart of nuclides, there are very few lifetimes for the $J^\pi = 0^+$ levels whereas there are numerous lifetime measurements for the $J^\pi = 2^+$ and $J^\pi = 4^+$ levels of $K^\pi = 0^+$ bands. This presents a unique opportunity to test predictions regarding the characters of these excited $K^\pi = 0^+$ bands. In addition, we have included lifetimes and depopulating transition probabilities to and from the $K^\pi = 4^+$ banded state at 1510.6 keV to allow for comparison with the $K^\pi = 2^+$ γ -band. These lifetimes had been previously measured by coulomb excitation

$K^\pi = 0_2^+$ band at 1049.479 keV: The lowest energy excited band in ^{156}Gd is the $K^\pi=0^+$ band at 1049.5 keV. Lifetimes were extracted for the band head and first two excited states. The $J^\pi=0^+$ level was previously measured using GRID with a lifetime range of $1.39 < \tau < 12.97$ [47]. In the current measurement, using the same technique, the new

TABLE III. Measured level lifetimes in the ^{156}Gd nucleus and the extracted B(E2) values. All of the level lifetimes are measured in this work with the exception of the $K^\pi = 4^+$ band head at 1510.595 keV [67]. Transition intensities and conversion coefficients are from Klora *et al.* [47]. The last column is the square root of the B(E) λ divided by the $\text{CG}^2 \left(\sqrt{\frac{B(E)\lambda}{\text{CG}^2}} \right)$. The CG^2 coefficients are the standard Alaga rules $(J_i K_i 2\Delta K | J_f K_f)^2$ values. If the exact multipole admixture is unknown, 100% E2 is calculated. The level lifetime for the $K^\pi = 4^+$ band head state at 1510.6 keV was not measured in this work but it is included to allow discussion and comparison.

K_i^π, J_i^π	E_x	τ	E_γ	K_f^π, J_f^π	I_γ	α	multipolarity	B(E1) or B(E2)	B(E) λ	CG^2	Reduced Matrix Element
	(keV)	(ps)	(keV)					mW.u. or W.u.	(e^2b^2) ^a		(eb)
$0_2^+, 0^+$	1049.479	$1.81 < \tau < 4.75$	960.510	$0_{g.s.}^+, 2^+$	138	0.00259	E2	4.20 \rightarrow 11.0	0.021 \rightarrow 0.055	1.000	0.145 \rightarrow 0.234
$0_2^+, 2^+$	1129.440	$1.26 < \tau < 2.58$	1129.423	$0_{g.s.}^+, 0^+$	80	0.00197	E2	0.56 \rightarrow 1.14	0.0028 \rightarrow 0.0057	0.200	0.118 \rightarrow 0.169
			1040.474	$0_{g.s.}^+, 2^+$	294	0.011	E2+M1+E0	3.10 \rightarrow 6.28	0.016 \rightarrow 0.031	0.286	0.233 \rightarrow 0.331
			841.243	$0_{g.s.}^+, 4^+$	119	0.0032	E2	3.61 \rightarrow 7.42	0.018 \rightarrow 0.037	0.515	0.187 \rightarrow 0.268
$0_2^+, 4^+$	1297.825	$1.49 < \tau < 3.36$	1208.875	$0_{g.s.}^+, 2^+$	97	0.00159	E2	0.95 \rightarrow 2.14	0.0047 \rightarrow 0.011	0.286	0.129 \rightarrow 0.196
			1009.622	$0_{g.s.}^+, 4^+$	83	0.0168	E2+M1+E0	1.96 \rightarrow 4.43	0.0098 \rightarrow 0.022	0.260	0.194 \rightarrow 0.291
			713.104	$0_{g.s.}^+, 6^+$	11.7	0.0044	E2	1.60 \rightarrow 3.60	0.0080 \rightarrow 0.018	0.455	0.132 \rightarrow 0.199
			168.382	$0_2^+, 2^+$	1.08	0.0234	E2	200 \rightarrow 440	1.00 \rightarrow 2.19	0.286	1.87 \rightarrow 2.77
$2_1^+, 2^+$	1154.151	$0.87 < \tau < 1.35$	1154.151	$0_{g.s.}^+, 0^+$	498	0.00192	E2	2.74 \rightarrow 4.25	0.014 \rightarrow 0.021	0.200	0.265 \rightarrow 0.324
			1065.182	$0_{g.s.}^+, 2^+$	549	0.00219	89% E2	4.01 \rightarrow 6.23	0.020 \rightarrow 0.031	0.286	0.264 \rightarrow 0.329
			865.971	$0_{g.s.}^+, 4^+$	26.9	0.00272	E2	0.62 \rightarrow 0.96	0.0031 \rightarrow 0.0048	0.015	0.454 \rightarrow 0.565
$2_1^+, 3^+$	1248.008	$0.68 < \tau < 1.01$	1159.000	$0_{g.s.}^+, 2^+$	659	0.00178	E2	6.12 \rightarrow 9.10	0.031 \rightarrow 0.046	0.358	0.294 \rightarrow 0.355
			959.823	$0_{g.s.}^+, 4^+$	173	0.00265	E2	4.12 \rightarrow 6.12	0.021 \rightarrow 0.030	0.143	0.383 \rightarrow 0.458
$2_1^+, 4^+$	1355.425	$0.58 < \tau < 1.02$	1266.451	$0_{g.s.}^+, 2^+$	94	0.0014	E2	1.40 \rightarrow 2.46	0.007 \rightarrow 0.012	0.120	0.241 \rightarrow 0.316
			1067.236	$0_{g.s.}^+, 4^+$	235	0.00206	E2	8.24 \rightarrow 14.5	0.041 \rightarrow 0.072	0.351	0.342 \rightarrow 0.453
			201.269	$2_1^+, 2_1^+$	0.63	0.078	E2	86.1 \rightarrow 150	0.429 \rightarrow 0.748	0.120	1.89 \rightarrow 2.60
$2_1^+, 5^+$	1506.868	$0.25 < \tau < 1.82$	1218.710	$0_{g.s.}^+, 4^+$	81	0.00168	E2	2.41 \rightarrow 17.6	0.012 \rightarrow 0.088	0.319	0.194 \rightarrow 0.525
			922.186	$0_{g.s.}^+, 6^+$	30	0.00268	E2	3.60 \rightarrow 26.2	0.018 \rightarrow 0.131	0.182	0.314 \rightarrow 0.848
			258.860	$2_1^+, 3^+$	1.15	0.073	E2	74 \rightarrow 538	0.370 \rightarrow 2.69	0.191	1.39 \rightarrow 3.75
$0_3^+, 0^+$	1168.190	$3.00 < \tau < 14.7$	1079.230	$0_{g.s.}^+, 2^+$	111	0.00215	E2	0.76 \rightarrow 3.72	0.0038 \rightarrow 0.019	1.000	0.062 \rightarrow 0.138
$0_3^+, 2^+$	1258.075	$2.31 < \tau < 5.06$	1258.092	$0_{g.s.}^+, 0^+$	68	0.00145	E2	0.14 \rightarrow 0.31	0.00070 \rightarrow 0.0015	0.200	0.059 \rightarrow 0.088
			1169.092	$0_{g.s.}^+, 2^+$	180	0.00272	10% E2	0.053 \rightarrow 0.12	0.00026 \rightarrow 0.00060	0.286	0.030 \rightarrow 0.046
			969.868	$0_{g.s.}^+, 4^+$	250	0.0024	E2	1.90 \rightarrow 4.13	0.0095 \rightarrow 0.021	0.515	0.136 \rightarrow 0.202

^a E1 transitions calculated in units of e^2b and E2 transitions calculated in units of e^2b^2 .

K_i^π, J_i^π	E_x	τ	E_γ	K_f^π, J_f^π	I_γ	α	multipolarity	B(E1) or B(E2)	B(E λ)	CG ²	Reduced Matrix Element
	(keV)	(ps)	(keV)					mW.u. or W.u.	(e ⁿ b ⁿ) ^a		(eb)
$4_1^+, 4^+$	1510.595	274 \pm 7	1421.601	$0_{g.s.}^+, 2^+$	79	0.00117	E2	0.0020	10×10^{-6}	-	-
			1222.432	$0_{g.s.}^+, 4^+$	189	0.00191	E2	0.0102	50×10^{-6}	-	-
			925.920	$0_{g.s.}^+, 6^+$	22.6	0.00284	E2	0.0049	24×10^{-6}	-	-
			381.155	$0_2^+, 2^+$	3.47	0.0241	E2	0.062	0.00031	-	-
			356.466	$2_1^+, 2^+$	68	0.0252	E2	1.71	0.0085	0.556	0.124
			262.589	$2_1^+, 3^+$	31.9	0.069	E2	3.54	0.018	0.312	0.240
			212.771	$0_2^+, 4^+$	0.21	0.166	E2+M1	0.061	0.00030	-	-
			155.168	$2_1^+, 4^+$	9.0	0.505	20% E2	1.97	0.0098	0.110	0.298
$0_4^+, 0^+$	1715.181	0.02 < τ < 7.08	561.024	$2_1^+, 2^+$	4.81	0.0082	E2	5.4 \rightarrow 1890	0.027 \rightarrow 9.48	1.000	0.164 \rightarrow 3.07
			472.700	$1^-, 1^-$	28.6	0.0058	E1	0.35 \rightarrow 120	(1.75 \rightarrow 600) $\times 10^{-6}$	1.000	0.0013 \rightarrow 0.025
			348.726	$0^-, 1^-$	3.1	0.006	E1	0.095 \rightarrow 33	(0.48 \rightarrow 165) $\times 10^{-6}$	1.000	0.0007 \rightarrow 0.013
$0_4^+, 2^+$	1771.089	0.13 < τ < 0.73	1682.184	$0_{g.s.}^+, 2^+$	158	0.00103	50% E2	0.74 \rightarrow 4.18	0.0037 \rightarrow 0.021	0.286	0.114 \rightarrow 0.271
			528.627	$1^-, 1^-$	1.29	0.0032	E1	0.023 \rightarrow 0.13	(0.12 \rightarrow 0.65) $\times 10^{-6}$	0.100	0.001 \rightarrow 0.002
			513.021	$0_3^+, 2^+$	3.3	0.024	M1	-	-	-	-
			494.942	$1^-, 3^-$	11.9	0.0053	E1	0.26 \rightarrow 1.43	(1.3 \rightarrow 7.2) $\times 10^{-6}$	0.400	0.0018 \rightarrow 0.0042
			404.633	$0^-, 1^-$	1.78	0.02	E1	0.069 \rightarrow 0.39	(0.34 \rightarrow 1.9) $\times 10^{-6}$	0.400	0.0009 \rightarrow 0.0022
$0_4^+, 4^+$	1893.395	0.00015 < τ < 0.45	1605.217	$0_{g.s.}^+, 4^+$	59	0.0011	59% E2	1.70 \rightarrow 5109	0.0085 \rightarrow 25	0.286	0.17 \rightarrow 9.44
			617.24	$1^-, 3^-$	1.4	0.0046	E1	0.064 \rightarrow 190	(0.32 \rightarrow 950) $\times 10^{-6}$	0.351	0.0010 \rightarrow 0.052
			595.580	$0_2^+, 4^+$	0.67	0.0184	M1+E2	4.6 \rightarrow 13,800	0.023 \rightarrow 70	0.260	0.297 \rightarrow 16
			537.954	$2_1^+, 4^+$	5.6	0.0142	56% E2	36 \rightarrow 107,000	0.18 \rightarrow 540	0.351	0.715 \rightarrow 39
			485.274	$1^-, 5^-$	2.4	0.0062	E1	0.224 \rightarrow 673	(1.12 \rightarrow 3360) $\times 10^{-6}$	0.334	0.0018 \rightarrow 0.100
			431.123	$0_3^+, 4^+$	0.58	0.043	M1	-	-	-	-

^a E1 transitions calculated in units of e²b and E2 transitions calculated in units of e²b².

lifetime range of 1.81 \rightarrow 4.75 ps is in excellent agreement but with a smaller range of values yielding a B(E2: $0_{K^\pi=0_2^+}^+ \rightarrow 2_{g.s.}^+$) = 4.20 \rightarrow 11.0 W.u. The J ^{π} =2⁺ level at 1129.4 keV lifetime range was measured at 1.26 \rightarrow 2.58 ps and compares well with the previously reported value of 2.27 \pm 0.17 ps from coulomb excitation [60]. The dominant transition probability from this level is E _{γ} = 841.2 keV to the J ^{π} = 4⁺ _{$g.s.$} level with a B(E2: $2_{K^\pi=0_2^+}^+ \rightarrow 4_{g.s.}^+$) range of 3.61 \rightarrow 7.42 W.u. The J⁺ = 4⁺ level at 1297.8 keV has a measured lifetime range of 1.49 \rightarrow 3.36 ps yielding a B(E2: $4_{K^\pi=0_2^+}^+ \rightarrow 6_{g.s.}^+$) range of 1.60 \rightarrow 3.60 W.u. Also calculated is an intraband transition from the J ^{π} = 4⁺ level to the J ^{π} = 2⁺ level of the same band yielding a B(E2: $4_{K^\pi=0_2^+}^+ \rightarrow 2_{K^\pi=0_2^+}^+$) range of 200 \rightarrow 440 W.u. typical of intraband rotational transition strengths.




FIG. 2. (color online) A partial level scheme for ^{156}Gd showing levels for which lifetimes were measured using $0.6 \times \tau_{\text{max}}$ of the GRID range as described in the text. The $K^\pi = 4^+$, $J^\pi = 4^+$ level was previously measured by coulomb excitation. The width of the transition lines depopulating the levels of interest are in proportion to the transition probability in W.u. Absolute $B(E2)$ values are presented in Table III.

Reduced matrix elements are also listed in Table III and are defined as the square roots of the $B(E2)$ values divided by the Clebsch-Gordon coefficient. The reduced matrix elements for those transitions depopulating the first excited $K^\pi = 0^+$ band average between 0.163 and 0.241 eb. These numbers are deduced from the averages of the lower(upper) limits for all

the transitions depopulating this band. The measured $B(E2:0_{K^\pi=0_2^+}^+ \rightarrow 2_{g.s.}^+)$ value is in the range of $0.021 \rightarrow 0.055 e^2b^2$. This value is divided by $2J_i + 1$ to yield a $B(E2:2_{g.s.}^+ \rightarrow 0_{K^\pi=0_2^+}^+)$ range of $0.0042 - 0.011 e^2b^2$. The intraband reduced matrix element is between $1.87 - 2.77 eb$ in agreement with values from coulomb excitation [59].

$K^\pi = 2_1^+$ band at 1154.151 keV: Our results confirm previous reports of the $K^\pi=2^+$ band as the γ band. The lifetimes of the four lowest energy levels ($2^+, 3^+, 4^+, 5^+$) of this band were measured. The $J^\pi=2^+$ level was measured in this experiment to be $0.87 \rightarrow 1.35$ ps in this experiment, in agreement with previously results of $0.82(3)$ from coulomb excitation [59] and a previous GRID measurement of $0.87 < \tau < 1.44$ ps [47].

All states in this band are connected to the ground state yielding reduced matrix element average values between $0.305 - 0.465 eb$ whereas the intraband transitions are $1.64 - 3.12 eb$. The $B(E2)$ values of transitions from the measured levels in this band to the ground state band are sufficiently strong to be considered collective transitions as calculated in Table III and shown in Fig. 2. There are also several intraband transitions known and the measured lifetimes result in $B(E2)$ strengths of tens to hundreds of W.u. providing evidence that these levels are members of the same band.

$K^\pi = 0_3^+$ band at 1168.190 keV: The $J^\pi = 0^+$ band head located at 1168.190 keV and $J^\pi = 2^+$ level at 1258.075 keV resulted in lifetimes of $3.00 \rightarrow 14.7$ ps and $2.31 \rightarrow 5.06$ ps, respectively. The $B(E2)$ values from these levels to the ground state are given in Table III and illustrated in Fig. 2. The average matrix elements for the transitions depopulating the two levels of this band are smaller and range from $0.030 - 0.202 eb$. An examination of the other $K^\pi = 0^+$ bands in comparison with this one reveals somewhat weaker $B(E2)$ transition probabilities and therefore would not be chosen as the collective excitation built on the ground state band. Further evidence comes from the dynamic moments of inertia for all the $K^\pi = 0^+$ bands of interest in this nucleus. Fig. 3 shows the dynamic moments of inertia for the $K^\pi = 0^+$ bands, including the $K^\pi = 0_1^+$ ground state band, the first excited $K^\pi = 0_2^+$ band at 1049.479 keV, the $K^\pi = 0_3^+$ band at 1168.190 keV, and the $K^\pi = 0_4^+$ band at 1715.181 keV. The only band with some significant variation in the dynamic moment of inertia is the 1168.190 keV band. All others have identical dynamic moments of inertia similar to the g.s. within a 7% variation in the slopes. The $K^\pi = 0_3^+$ band at 1168.2 keV is significantly different from the other $K^\pi = 0^+$ bands. Ref. [25] reports on a study of quantum fluctuations around the equilibrium deformed shape of rare earth nuclei with

respect to the nature of collective 0^+ states. The work includes studies of the Gd isotopes where they show wave functions for 0^+ states built on the equilibrium deformed shape with one and two nodes (one and two phonon) oscillations. Their work does not report on a third excited 0^+ . The deviation of the $K^\pi = 0_3^+$ states in the dynamic moment of inertia indicates a 0^+ state of a different nature while the $K^\pi = 0_2^+$ and the $K^\pi = 0_4^+$ do indeed show themselves as oscillations or fluctuations around an equilibrium shape in this ^{156}Gd nucleus. A similar study [28] in the ^{178}Hf nucleus where there are four excited $K^\pi = 0^+$ bands below 2 MeV showed that two of the $K^\pi = 0^+$ bands that were strongly connected by collective B(E2) transitions had identical dynamic moments of inertia and indicated that they were collective oscillations built on the ground state equilibrium shape whereas the two intermediate $K^\pi = 0^+$ bands showed significantly different moments of inertia.

$K^\pi = 4_1^+$ band at 1510.595 keV: A lifetime of 274(7) ps was previously measured by coulomb excitation [59, 60] and not remeasured in this work. B(E2) transitions from this $J^\pi = 4^+$ state to the g.s. are K forbidden and show very weak transition probabilities to the g.s. band. The transitions depopulating to the 2^+ , 3^+ , and 4^+ members of the $K^\pi=2_1^+$ γ band however are much stronger yielding matrix elements on the average of 0.221 *eb*. The energy ratio of the ($K^\pi = 4_1^+$)/($K^\pi = 2_1^+$) bands is 1.31, well below the expected value of 2 and the E2 matrix elements are of the same order as the $K^\pi = 2_1^+$ to the ground state transitions.

$K^\pi = 0_4^+$ band at 1715.181 keV: The 1715.2 keV band is the third $K^\pi = 0^+$ for which lifetimes were measured including the first three levels ($0^+, 2^+, 4^+$). Low intensity transitions from these levels resulted in fewer statistics. The feeding to these transitions is not well known: the $J^\pi = 0^+$ level at 1715.181 keV, $J^\pi = 2^+$ level at 1771.089 keV and the $J^\pi = 4^+$ level at 1893.395 keV have known feeding intensities of 11%, 26%, and 1%, respectively. This resulted in the extracted lifetimes with larger ranges than other measurements in this experiment. B(E2) transition strengths were calculated for transitions depopulating the $K^\pi = 0_4^+$ and despite the less than ideal lifetime ranges some conclusions concerning the nature of this band can still be made. The E2 transitions from this band to states within the ground state band include the transitions and B(E2) values B(E2: $2_{K^\pi=0_4^+}^+ \rightarrow 2_{g.s.}^+$) and B(E2: $4_{K^\pi=0_4^+}^+ \rightarrow 4_{g.s.}^+$) range of 0.74 \rightarrow 4.18 W.u. and > 1.70 W.u., respectively. The B(E2: $0_{K^\pi=0_4^+}^+ \rightarrow 2_{K^\pi=2_1^+}^+$) range of 5.4 \rightarrow 1890 W.u. and B(E2: $4_{K^\pi=0_4^+}^+ \rightarrow 4_{K^\pi=2_1^+}^+$) range yields > 36 W.u. connecting the 0^+ and 4^+ state of this band to the $K^\pi = 2_1^+$ γ band. These




FIG. 3. (color online) Dynamic moments of inertia for the four $K^\pi = 0^+$ bands, including the $K^\pi = 0_1^+$ ground state band, the first excited $K^\pi = 0_2^+$ band at 1049.479 keV, the $K^\pi = 0_3^+$ band at 1168.190 keV, and the $K^\pi = 0_4^+$ band at 1715.181 keV. Only the 1168.190 keV band has some significant variation in the dynamic moment of inertia.

transitions show the $K^\pi = 0_4^+$ band to be collectively built on the $K^\pi = 2_1^+$ band and is evidence that the $K^\pi = 0_4^+$ band is a $K^\pi = 0^+$ $\gamma\gamma$ two-phonon vibrational band ($K^+ = 0_{\gamma\gamma}^+$). Fig. 4 shows the dynamic moments of inertia plots with identical (within 7%) slopes for the four bands considered here, the g.s. $K^\pi = 0^+$, the $K^\pi = 2^+$, the $K^\pi = 4^+$, and the $K^\pi = 0_4^+$ band at 1715.181 keV. This is an indication that all of them are excitations built on the g.s. band. A summary of the relevant energy and B(E2) ratios are shown in Table IV.




FIG. 4. (color online) Dynamic moments of inertia for the four bands considered including the g.s. band, the $K^\pi = 2^+$ gamma band, the $K^\pi = 4^+$ band and the $K^\pi = 0^+$ band at 1715.181 keV. These four bands have dynamic moments of inertia with less than 5% variation in slope. This is an indication that all of them are excitations built on the g.s. band.

TABLE IV. The experimental energy and B(E2) ratios for the proposed single and double phonon excitations.

$E(4_{\gamma\gamma}^+)/E(2_\gamma^+)$	1.31
$E(0_{\gamma\gamma}^+)/E(2_\gamma^+)$	1.49
$B(E2:4_{\gamma\gamma}^+ \rightarrow 2_\gamma^+)/B(E2:2_\gamma^+ \rightarrow 0_{g.s.}^+)$	0.39
$B(E2:0_{\gamma\gamma}^+ \rightarrow 2_\gamma^+)/B(E2:2_\gamma^+ \rightarrow 0_{g.s.}^+)$	1.96

IV. DISCUSSION

A comprehensive theoretical study of the low energy structure of well deformed nuclei was carried out for even-even nuclei from $Z = 10$ to $Z = 110$ using the Hartree-Fock Bogoliubov theory extended by the generator coordinate method and mapped onto a five dimensional collective quadrupole Hamilton (5DCH) by Delaroche *et al.* [62]. This approach is able to distinguish and to separate between excited 0^+ bands that are β vibrations from those that are in fact due to co-existing minima on the basis of relative quadrupole transition strengths. Ref. [62] has a calculated data set of over 1710 even-even nuclei and they find the shape coexistence phenomenon to be more prevalent but characterize the first excited $K^\pi = 0^+$ band in the ^{152}Sm to be a β vibration. The 5DCH calculations for the ^{156}Gd nucleus result in a β -deformation of 0.347 and a $4^+/2^+$ energy ratio of 3.27. The experimental $4^+/2^+$ energy ratio is 3.24. The calculated first excited $K^\pi = 0^+$ and $K^\pi = 2^+$ band heads are at 1274 and 1159 keV, in comparison to the experimental values at 1049.48 and 1129.40 keV, respectively. The Delaroche calculations place the first excited $K^\pi = 0^+$ band above the $K^\pi = 2_1^+$ band and predict $B(E2: 2_{K^\pi=0_2^+}^+ \rightarrow 0_{g.s.}^+) = 0.0253 e^2b^2$ in comparison with the measured range of $0.0028 \rightarrow 0.0057 e^2b^2$. The theoretical predication is 9 to 4 times larger than the experimental $B(E2)$ value range.

Perhaps one of the most important structure indicators are the intra- and interband E2 reduced transition probabilities. This work provides measured absolute transition probabilities for both types of transitions in ^{156}Gd to allow for experimental comparison. The reduced matrix elements extracted from measurements for the transitions depopulating the $K^\pi = 0_2^+$ average between 0.163 and 0.241 eb. Whereas the intraband reduced matrix element is between $1.87 \rightarrow 2.77$ eb and dividing by e yields a range of $1.28 \rightarrow 1.94$ b [63] for the matrix elements extracted from the measurements.

The ground state $B(E2: 2_{g.s.}^+ \rightarrow 0_{g.s.}^+)$ value yields a matrix element of 1.9 barn from coulomb excitation [64]. The $B(E2)$ strength of transitions from each of the levels of this $K^\pi = 0_2^+$ band to the ground state band is evidence that this lowest lying $K^\pi = 0^+$ band is a collective excitation build on the ground state and perhaps most likely, the β vibrational one-phonon band albeit with less collectivity than expected by the 5DCH approach.

The viability of the first excited $K^\pi = 0^+$ band as a β vibration is dependent on a consistent intrinsic matrix element for the interband transitions. The $B(E2)$ values for transitions between $K_1, K_2 = 0$ bands considered here are given by the equation:

$$B(E2 : J_1 \rightarrow J_2) = (J_1 0 2 0 | J_2 0)^2 \frac{5}{4\pi} e^2 Q_0^2 \quad (1)$$

The measured ranges of $B(E2)$ values presented in the results table are used to plot the extracted intrinsic Quadrupole moments shown in Fig. 5. Ideally, we would expect a complete overlap of the error bars for the values of the intrinsic quadrupole moment but the results vary slightly and yield a value of approximately 0.3 b indicating that the first excited $K^\pi = 0^+$ band is a potential beta excitation built on the ground state.

An interpretation by Leviatan *et al.* [46] has proposed the use of partial dynamical symmetry (PDS) as a selection criterion for states in a given nucleus that obey a specific symmetry while others break the symmetry strongly. They presented their calculations for ^{156}Gd and made available calculated $B(E2)$ values presuming that the first excited $K^\pi = 0^+$ band is a β vibration. These calculations, in comparison with measured values are shown in Table V. The PDS calculated $B(E2: 0_{K^\pi=0_2^+}^+ \rightarrow 2_{g.s.}^+) = 0.034 e^2 b^2$ values in comparison with a measured range of $0.021 \rightarrow 0.055 e^2 b^2$; and the $B(E2: 2_{K^\pi=0_2^+}^+ \rightarrow 0_{g.s.}^+) = 0.0055 e^2 b^2$ in comparison to the measured range of $0.0028 \rightarrow 0.0057 e^2 b^2$ are in the same order as expected for a β vibration. The transitions depopulating the $K^\pi = 0_4^+$ band at 1715.181 keV show strong connections to the the $K^\pi = 2_\gamma^+$ band and is evidence that the $K^\pi = 0_4^+$ band is a collective excitation and perhaps the $K^\pi = 0^+$ $\gamma\gamma$ two-phonon vibrational band. The dynamic moments of inertia for the various excitation bands in ^{156}Gd show nearly identical slopes with the exception of the $K^\pi = 0^+$ band at 1168.190 keV indicating that the excitations are oscillations around the equilibrium deformed shape of the nucleus. The $K^\pi = 0_2^+$, $K^\pi = 2^+$, the $K^\pi = 4^+$, and the $K^\pi = 0_4^+$ band at 1715.181 keV all have identical dynamic moments of inertia as shown in Fig. 4.

A focus in nuclear structure has been whether two phonon excitations can occur with negative anharmonicity or a ratio of less than two. The ^{166}Er nucleus is the only other case of a known case of a two phonon $K^\pi = 0_{\gamma\gamma}^+$ and $K^\pi = 4_{\gamma\gamma}^+$ set of vibrations built on the γ band where the excitation energy ratios of $K^\pi = 4_{\gamma\gamma}^+$, $K^\pi = 0_{\gamma\gamma}^+$ excitations [65] to the $K^\pi = 2_\gamma^+$ is 2.6 and 2.5, respectively, in comparison with the expected values of two. In ^{156}Gd , the energy ratio of the $(K^\pi = 0_{\gamma\gamma}^+)/ (K^\pi = 2_{g.s.}^+)$ band heads is 1.49, well below the




FIG. 5. The intrinsic quadrupole moment for all the transitions depopulating the first excited $K^\pi = 0^+$ band extracted from the upper and lower ranges of the $B(E2)$ values determined from the lifetime measurements in this work shown in Table III.

TABLE V. Comparison of measured $B(E2)_{\text{exp}}$ values and theoretical $B(E2)$ values, PDS calculated values are from Ref. [46] for the first excited 0^+ band, $K^\pi = 0_2^+$.

K_i^π, J_i^π	E_x (keV)	E_γ (keV)	K_i^π, J_i^π	$B(E2)_{\text{exp}}$ (e^2b^2)	PDS (e^2b^2)
$0_2^+, 0^+$	1049.479	960.510	$0_{g.s.}^+, 2^+$	$0.021 \rightarrow 0.055$	0.034
$0_2^+, 2^+$	1129.440	1129.423	$0_{g.s.}^+, 0^+$	$0.0028 \rightarrow 0.0057$	0.0055
		1040.474	$0_{g.s.}^+, 2^+$	$0.016 \rightarrow 0.031$	0.0084
		841.243	$0_{g.s.}^+, 4^+$	$0.018 \rightarrow 0.037$	0.02
$0_2^+, 4^+$	1297.825	1208.875	$0_{g.s.}^+, 2^+$	$0.0047 \rightarrow 0.011$	0.0067
		1009.622	$0_{g.s.}^+, 4^+$	$0.0098 \rightarrow 0.022$	0.0067
		713.104	$0_{g.s.}^+, 6^+$	$0.0080 \rightarrow 0.018$	0.021
		168.382	$0_2^+, 2^+$	$1.00 \rightarrow 2.19$	0.951

expected value of 2.0 and the ratio of $B(E2)$ values is for $B(E2:0_{\gamma\gamma}^+ \rightarrow 2_\gamma^+)/B(E2:2_\gamma^+ \rightarrow 0_{g.s.}^+)$ is 1.96 well below the pure geometric expectation value of 5. In a study [66] of ^{232}Th , a $K^\pi = 4_{\gamma\gamma}^+$ band to the γ band ratio of 1.8 was observed with a $B(E2)$ ratio of 3.1 ± 0.4 . A similar observation [28] was made in the two phonon $K^\pi = 0_{\beta\beta}^+$ to the one phonon $K^\pi = 0_\beta^+$ energy ratio for the ^{178}Hf nucleus where the energy ratio is 1.5 instead of 2. The growing number of negative anharmonicity (less than 2) for two phonon oscillations warrants further examination.

V. SUMMARY

To summarize, 12 lifetimes were measured from levels in ^{156}Gd . Six of these are new measurements while another six were previously measured lifetimes that agree with our data. The $B(E2)$ values have been calculated using the intensities and electron conversion coefficients of Klora *et al.* [47]. The results show that the $K^\pi = 0_2^+$ band is strongly connected to the ground state band with matrix elements of the same size as those of the $K^\pi = 2_1^+$ showing that the first excited $K^\pi = 0_2^+$ band is collectively built on the ground state band and may in fact be the β vibrational band. The transitions from the $K^\pi = 0_3^+$ band to the ground state band are less strong and the dynamic moment of inertia for this band varies

fairly significantly from the other excitations built on the g.s. band as shown in Fig. 3. Further evidence comes from the slopes of the dynamic moments of inertia of all the four $K^\pi = 0^+$ bands. PDS calculations indicate that the first excited 0^+ is the collective β vibrational excitation and the measured $B(E2)$ values for transitions from the first excited $K^\pi = 0_2^+$ band are in good agreement with calculated values as shown in Table V. The 5DCH calculations [62] predict greater collectivity for the β vibration as discussed in the discussion section. Quantum fluctuation studies [25] of deformed nuclei and collective 0^+ excitations indicate two excited $K^\pi = 0^+$ bands built on the ground state equilibrium shape with one and two node wave functions that point to one and two phonon excitations in ^{156}Gd . The two phonon excitation in this case is the $\gamma\gamma$ $K^\pi = 0^+$.

The $K^\pi = 4_1^+$ band is lower in excitation energy than would be expected of a $\gamma\gamma$ two phonon excitation but it does show considerable strength to the $K^\pi = 2_1^+$ band. This can of course just be a reflection of ΔK preference. The excitation energy ratio of the $K^\pi = 4_1^+$ band with the single $K^\pi = 2_1^+$ γ band is significantly less than the expected value of 2. The $K^\pi = 0_4^+$ band also does not show strong transitions to the ground state band but does have very collective transitions to the $K^\pi = 2_1^+$, γ band. Furthermore, the slopes of the dynamic moments of inertia for the $K^\pi = 4_1^+$, the $K^\pi = 2_1^+$ band, and the $K^\pi = 0_4^+$ band are quite similar indicating common origin. The $K^\pi = 0_4^+$ band is most likely the $K^\pi = 0_{\gamma\gamma}^+$ multi-phonon excitation or at the very least the $\gamma\gamma$ multi-phonon excitation is a significant element of the band's character. Figure 6 shows the systematics of the deformed Gd nuclei from $A = 154 - 160$. In ^{154}Gd , the first excited $K^\pi = 0^+$ band is much lower than the $K^\pi = 2^+$ band with strong $B(E2)$ values connecting it to the ground state. There are numerous 0^+ states identified in this nucleus but they lack lifetime measurements. The ^{156}Gd presented in this work shows a low lying first excited $K^\pi = 0^+$ band near the $K^\pi = 2^+$ band with equivalent collectivity in $B(E2)$ values connecting it to the g.s. band. In ^{158}Gd , the collective 0^+ was shown to be higher than the pairing gap and in ^{160}Gd , there are only limits for the lifetimes of the known 0^+ states. The indications point to the first excited $K^\pi = 0^+$ band in ^{156}Gd to be a β vibration of the deformed ground state.




FIG. 6. (color online) Partial level schemes for $^{154-160}\text{Gd}$ showing the lowest $K^\pi = 2^+$ bands with the known 0^+ states, transition probabilities, and the two-proton($\Delta\pi$) and two-neutron ($\Delta\nu$) pairing gaps shown as horizontal lines. The width of the transition lines depopulating the levels of interest are in proportion to the transition probability in W.u.

VI. ACKNOWLEDGMENTS

The authors gratefully acknowledge discussions with G. Bertsch and the warm hospitality at the ILL in Grenoble, France where this experiment was performed. This work was supported by the National Science Foundation under contract PHY-1419765.

-
- [1] A. Aprahamian, S. R. Leshner, C. Casarella, H. G. Börner, and M. Jentschel, *Phys. Rev. C* **95**, 024329 (2017).
 - [2] K. Heyde and J. L. Wood, *Rev. Mod. Phys.* **83**, 1467 (2011).
 - [3] P. E. Garrett, W. D. Kulp, J. L. Wood, D. Bandyopadhyay, S. Choudry, D. Dashdorj, S. R. Leshner, M. T. McEllistrem, M. Mynk, J. N. Orce, and S. W. Yates, *Phys. Rev. Lett.* **103**, 062501 (2009).
 - [4] R. F. Casten and P. von Brentano, *Phys. Rev. C* **50**, 1280(R) (1994).
 - [5] A. Aprahamian, R. C. de Haan, S. R. Leshner, J. Döring, A. M. Bruce, H. G. Börner, M. Jentschel, and H. Lehmann, *J. Phys.* **G25**, 685 (1999).
 - [6] X. Wu, A. Aprahamian, J. Castro-Ceron, and C. Baktash, *Phys. Lett. B* **316**, 235 (1993).
 - [7] X. Wu, A. Aprahamian, S. M. Fischer, W. Reviol, G. Liu, and J. X. Saladin, *Phys. Rev. C* **49**, 1837 (1994).
 - [8] S. R. Leshner, A. Aprahamian, L. Trache, A. Oros-Peusquens, S. Deyliz, A. Gollwitzer, R. Hertenberger, B. D. Valnion, and G. Graw, *Phys. Rev. C* **66**, 051305(R) (2002).
 - [9] D. A. Meyer, V. Wood, R. F. Casten, C. R. Fitzpatrick, G. Graw, D. Bucurescu, J. Jolie, P. von Brentano, R. Hertenberger, H.-F. Wirth, N. Braun, T. Faestermann, S. Heinze, J. L. Jerke, R. Krücken, M. Mahgoub, O. Möller, D. Mücher, and C. Scholl, *Phys. Rev. C* **74**, 044309 (2006).
 - [10] D. Bucurescu, G. Graw, R. Hertenberger, H.-F. Wirth, N. Lo Iudice, A. V. Sushkov, N. Y. Shirikova, Y. Sun, T. Faestermann, R. Krücken, M. Mahgoub, J. Jolie, P. von Brentano, N. Braun, S. Heinze, O. Möller, D. Mücher, C. Scholl, R. F. Casten, and D. A. Meyer, *Phys. Rev. C* **73**, 064309 (2006).
 - [11] S. R. Leshner, J. N. Orce, Z. Ammar, C. D. Hannant, M. Merrick, N. Warr, T. B. Brown, N. Boukharouba, C. Fransen, M. Scheck, M. T. McEllistrem, and S. W. Yates, *Phys. Rev. C*

- 76**, 034318 (2007).
- [12] L. Bettermann, S. Heinze, J. Jolie, D. Mücher, O. Möller, C. Scholl, R. F. Casten, D. A. Meyer, G. Graw, R. Hertenberg, H.-F. Wirth, and D. Bucurescu, *Phys. Rev. C* **80**, 044333 (2009).
 - [13] C. Bernards, R. F. Casten, V. Werner, P. von Brentano, D. Bucurescu, G. Graw, S. Heinze, R. Hertenberg, J. Jolie, S. Lalkovski, D. A. Meyer, D. Mücher, P. Pejovic, C. Scholl, and H.-F. Wirth, *Phys. Rev. C* **87**, 064321 (2013).
 - [14] C. Bernards, R. F. Casten, V. Werner, P. von Brentano, D. Bucurescu, G. Graw, S. Heinze, R. Hertenberg, J. Jolie, S. Lalkovski, D. A. Meyer, D. Mücher, P. Pejovic, C. Scholl, and H.-F. Wirth, *Phys. Rev. C* **87**, 024318 (2013).
 - [15] N. V. Zamfir, J.-y. Zhang, and R. F. Casten, *Phys. Rev. C* **66**, 057303 (2002).
 - [16] Y. Sun, A. Aprahamian, J. Zhang, and C.-T. Lee, *Phys. Rev. C* **68**, 061301(R) (2003).
 - [17] N. Pietralla and O. M. Gorbachenko, *Phys. Rev. C* **70**, 011304(R) (2004).
 - [18] K. Dusling, N. Pietralla, G. Rainovski, T. Ahn, B. Bochev, A. Costin, T. Koike, T. C. Li, A. Linnemann, S. Pontillo, and C. Vaman, *Phys. Rev. C* **73**, 014317 (2006).
 - [19] R. Fossion, C. E. Alonso, J. M. Arias, L. Fortunato, and A. Vitturi, *Phys. Rev. C* **76**, 014316 (2007).
 - [20] D. Bonatsos, E. A. McCutchan, R. F. Casten, R. J. Casperson, V. Werner, and E. Williams, *Phys. Rev. C* **80**, 034311 (2009).
 - [21] D. Bonatsos, I. E. Assimakis, N. Minkov, A. Martinou, S. Sarantopoulou, R. B. Cakirli, R. F. Casten, and K. Blaum, *Phys. Rev. C* **95**, 064326 (2017).
 - [22] D. Bonatsos, I. E. Assimakis, N. Minkov, A. Martinou, R. B. Cakirli, R. F. Casten, and K. Blaum, *Phys. Rev. C* **95**, 064325 (2017).
 - [23] R. M. Clark, R. F. Casten, L. Bettermann, and R. Winkler, *Phys. Rev. C* **80**, 011303(R) (2009).
 - [24] N. Lo Iudice, V. Yu. Ponomarev, Ch. Stoyanov, A. V. Sushkov, and V. V. Voronov, *J. Phys. G: Nucl. Part. Phys.* **39**, 043101 (2012).
 - [25] F.-Q. Chen, Y. Sun, and P. Ring, *Phys. Rev. C* **88**, 014315 (2013).
 - [26] P. E. Garrett, M. Kadi, C. A. McGrath, V. Sorokin, M. Li, M. Yeh, and S. W. Yates, *Phys. Lett. B* **400**, 250 (1997).

- [27] R. C. de Haan, A. Aprahamian, H. G. Börner, C. Doll, M. Jentschel, A. M. Bruce, and S. R. Leshner, *J. Res. Natl. Inst. Stand. Technol.* **105**, 125 (2000).
- [28] A. Aprahamian, R. C. de Haan, H. G. Börner, H. Lehmann, C. Doll, M. Jentschel, A. M. Bruce, and R. Piepenbring, *Phys. Rev. C* **65**, 031301(R) (2002).
- [29] A. Aprahamian, *Phys. Atom. Nucl.* **67**, 1750 (2004).
- [30] J. F. Sharpey-Schafer, T. E. Madiba, S. P. Bvumbi, E. A. Lawrie, J. J. Lawrie, A. Minkova, S. M. Mullins, P. Papka, D. G. Roux, and Timár, *Eur. Phys. J. A* **47**, 6 (2011).
- [31] J. F. Sharpey-Schafer, *AIP Conf. Proc.* **1377**, 205 (2011).
- [32] J. F. Sharpey-Schafer, S. M. Mullins, R. A. Bark, J. Kau, F. Komati, E. A. Lawrie, J. J. Lawrie, T. E. Madiba, P. Maine, A. Minkova, S. H. T. Murray, N. J. Ncapayi, and P. A. Vymers, *Eur. Phys. J. A* **47**, 5 (2011).
- [33] T. Papenbrock and H. A. Weidenmüller, *Phys. Scr.* **91**, 053004 (2016).
- [34] E. A. Coello Pérez and T. Papenbrock, *Phys. Rev. C* **92**, 014323 (2015).
- [35] G. Popa, J. G. Hirsch, and J. P. Draayer, *Phys. Rev. C* **62**, 064313 (2000).
- [36] A. Arima and F. Iachello, *Ann. Phys. (NY)* **99**, 253 (1976).
- [37] R. F. Casten, P. von Brentano, and N. V. Zamfir, *Phys. Rev. C* **49**, 1940 (1994).
- [38] R. F. Casten and P. von Brentano, *Phys. Rev. C* **51**, 3528 (1995).
- [39] F. Iachello, N. V. Zamfir, and R. F. Casten, *Phys. Rev. Lett.* **81**, 1191 (1998).
- [40] V. Werner, E. Williams, R. J. Casperson, R. F. Casten, C. Scholl, and P. von Brentano, *Phys. Rev. C* **78**, 051303(R) (2008).
- [41] D. Tonev, A. Dewald, T. Klug, P. Petkov, J. Jolie, A. Fitzler, O. Möller, S. Heinze, P. von Brentano, and R. F. Casten, *Phys. Rev. C* **69**, 034334 (2004).
- [42] P. E. Garrett, *J. Phys. G: Nucl. Part. Phys.* **27**, R1 (2001).
- [43] D. D. Warner and R. F. Casten, *Phys. Rev. C* **25**, 2019 (1982).
- [44] D. D. Warner and R. F. Casten, *Phys. Rev. Lett.* **48**, 1385 (1982).
- [45] R. F. Casten and D. D. Warner, *Rev. Mod. Phys.* **60**, 389 (1988).
- [46] A. Leviatan, J. E. García-Ramos, and P. Van Isacker, *Phys. Rev. C* **87**, 021302(R) (2013).
- [47] J. Klorá, H. G. Börner, T. von Egidy, R. Georgii, J. Jolie, S. Judge, V. A. Khitrov, B. Krusche, V. A. Libman, H. Lindner, L. L. Litvinsky, U. Mayerhofer, A. V. Murzin, S. J. Robinson, A. M. Sukhovej, and H. Trieb, *Nucl. Phys. A* **561**, 1 (1993).

- [48] H. G. Börner, J. Jolie, F. Hoyler, S. Robinson, M. S. Dewey, G. Greene, E. E. Kessler, Jr., and R. D. Deslattes, *Phys. Lett. B* **215**, 45 (1988).
- [49] H. Börner, *I.O.P. Conf. Ser.* **88**, 143 (1988).
- [50] M. Dewey, E. Kessler, Jr., G. Greene, R. Deslattes, H. Börner, and J. Jolie, *Nucl. Instr. and Meth.* **A284**, 151 (1989).
- [51] D. Bohle, A. Richter, W. Steffen, A. Dieperink, N. LoIudice, F. Palumbo, and O. Scholten, *Phys. Lett. B* **137**, 27 (1984).
- [52] U. E. P. Berg, C. Bläsing, J. Drexler, R. D. Heil, U. Kneissel, W. Naatz, R. Ratzek, S. Schenach, R. Stock, T. Weber, H. Wickert, B. Fischer, H. Hollick, and D. Kollewe, *Phys. Lett. B* **149**, 59 (1984).
- [53] F. Iachello, *Ann. of Phys.* **192**, 133 (1989).
- [54] J. Jolie, S. Ulbig, H. G. Börner, K. P. Lieb, S. J. Robinson, P. Schillebeeckx, E. G. Kessler, M. S. Dewey, and G. L. Greene, *Europhys. Lett.* **10**, 231 (1989).
- [55] R. D. Deslattes, E. G. Kessler, Jr., W. C. Sauder, and A. Henins, *Ann. of Phys.* **129**, 378 (1980).
- [56] E. G. Kessler Jr., G. L. Greene, R. D. Deslattes, and H. G. Börner, *Phys. Rev. C* **32**, 374 (1985).
- [57] E. G. Kessler, Jr., G. L. Greene, M. S. Dewey, R. D. Deslattes, H. Börner, and F. Hoyler, *J. Phys. G* **14**, S167 (1988).
- [58] S. Robinson and J. Jolie, *ILL Internal Report* **92Ro15T** (1992).
- [59] M. Sugawara, H. Kusakari, Y. Yoshizawa, H. Inoue, T. Morikawa, T. Shizuma, and J. Srebrny, *Phys. Rev. C* **83**, 064308 (2011).
- [60] F. K. McGowan and W. T. Milner, *Phys. Rev. C* **23**, 1926 (1981).
- [61] H. G. Börner, M. Jentschel, N. V. Zamfir, R. F. Casten, M. Krückova, and W. Andrejtscheff, *Phys. Rev. C* **59**, 2432 (1999).
- [62] J. P. Delaroche, M. Girod, J. Libert, H. Goutte, S. hilaire, S. Péru, N. Pillet, and G. F. Bertsch, *Phys. Rev. C* **81**, 014303 (2010).
- [63] B. Pritychenko, M. Birch, B. Singh, and M. Horoi, *At. Data Nucl. Data Tables* **107**, 1 (2016).
- [64] C. W. Reich, *Nucl. Data Sheets* **113**, 2537 (2012).
- [65] P. E. Garrett, M. Kadi, Min. Li, C. A. McGrath, V. Sorokin, M. Yeh, and S. W. Yates, *Phys. Rev. Lett.* **78**, 4545 (1997).

- [66] W. Korten, T. Hartlein, J. Gerl, D. Habs, and D. Schwalm, *Phys. Lett. B* **317**, 19 (1993).
- [67] A. Bäcklin, G. Hedin, B. Fogelberg, M. Saraceno, R. Greenwood, C. Reich, H. Koch, H. Baader, H. Breitig, O. Schult, K. Schreckenbach, T. Von Egidy, and W. Mampe, *Nucl. Phys.* **A380**, 189 (1982).