



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Beam-target helicity asymmetry E in $K^0\Lambda$ and $K^0\Sigma^0$ photoproduction on the neutron

D. H. Ho *et al.* (The CLAS Collaboration)

Phys. Rev. C **98**, 045205 — Published 19 October 2018

DOI: [10.1103/PhysRevC.98.045205](https://doi.org/10.1103/PhysRevC.98.045205)

Beam-Target Helicity Asymmetry E in $K^0\Lambda$ and $K^0\Sigma^0$ Photoproduction on the Neutron

D.H. Ho,^{1,*} R.A. Schumacher,^{1,†} A. D'Angelo,^{25,37} A. Deur,⁴¹ J. Fleming,⁴³ C. Hanretty,^{41,47} T. Kageya,⁴¹ F.J. Klein,⁶ E. Klempt,¹⁹ M.M. Lowry,⁴¹ H. Lu,²¹ V.A. Nikonov,¹⁶ P. Peng,⁴⁷ A.M. Sandorfi,⁴¹ A.V. Sarantsev,¹⁶ I.I. Strakovsky,¹⁸ N.K. Walford,⁶ X. Wei,⁴¹ R.L. Workman,¹⁸ K.P. Adhikari,^{35,‡} S. Adhikari,¹⁴ D. Adikaram,³⁵ Z. Akbar,¹⁵ J. Ball,⁷ L. Barion,²² M. Bashkanov,⁴³ C.D. Bass,⁴¹ M. Battaglieri,²⁴ I. Bedlinskiy,²⁸ A.S. Biselli,^{12,1} W.J. Briscoe,¹⁸ S. Boiarinov,⁴¹ V.D. Burkert,⁴¹ F. Cao,¹⁰ D.S. Carman,⁴¹ A. Celentano,²⁴ S. Chandavar,³⁴ G. Charles,²⁷ P. Chatagnon,²⁷ T. Chetry,³⁴ G. Ciullo,^{22,13} B. A. Clary,¹⁰ P. Cole,²⁰ P. Collins,⁶ M. Contalbrigo,²² V. Crede,¹⁵ R. De Vita,²⁴ M. Defurne,⁷ S. Diehl,¹⁰ C. Djalali,³⁹ R. Dupre,^{27,7,2} H. Egiyan,⁴¹ M. Ehrhart,²⁷ A. El Alaoui,⁴² L. El Fassi,³¹ P. Eugenio,¹⁵ G. Fedotov,³⁴ S. Fegan,^{44,§} R. Fersch,^{9,48} A. Filippi,²⁶ A. Fradi,²⁷ Y. Ghandilyan,⁴⁹ K.L. Giovanetti,²⁹ F.X. Girod,⁴¹ W. Gohn,^{10,¶} E. Golovatch,³⁸ R.W. Gothe,³⁹ K.A. Griffioen,⁴⁸ M. Guidal,²⁷ K. Hafidi,² H. Hakobyan,^{42,49} N. Harrison,⁴¹ M. Hattawy,² D. Heddle,^{9,41} K. Hicks,³⁴ M. Holtrop,³² Y. Ilieva,³⁹ D.G. Ireland,⁴⁴ B.S. Ishkhanov,³⁸ E.L. Isupov,³⁸ D. Jenkins,⁴⁶ H.S. Jo,³⁰ S. Johnston,² M.L. Kabir,³¹ D. Keller,⁴⁷ G. Khachatryan,⁴⁹ M. Khachatryan,³⁵ M. Khandaker,^{33,**} A. Kim,¹⁰ W. Kim,³⁰ A. Klein,³⁵ V. Kubarovsky,^{41,36} V. Laine,⁸ L. Lanza,²⁵ P. Lenisa,²² K. Livingston,⁴⁴ I. J. D. MacGregor,⁴⁴ D. Marchand,²⁷ N. Markov,¹⁰ M. Mayer,³⁵ B. McKinnon,⁴⁴ C.A. Meyer,¹ M. Mirazita,²³ V. Mokeev,^{41,38} R.A. Montgomery,⁴⁴ A. Movsisyan,²² C. Munoz Camacho,²⁷ P. Nadel-Turonski,⁴¹ S. Niccolai,²⁷ G. Niculescu,²⁹ T. O'Connell,¹⁰ M. Osipenko,²⁴ A.I. Ostrovidov,¹⁵ M. Paolone,⁴⁰ R. Paremuzyan,³² K. Park,⁴¹ E. Pasyuk,^{41,3} O. Pogorelko,²⁸ J.W. Price,⁴ Y. Prok,^{47,45} D. Protopopescu,⁴⁴ M. Rehfuss,⁴⁰ M. Ripani,²⁴ D. Riser,¹⁰ A. Rizzo,^{25,37} G. Rosner,⁴⁴ F. Sabatié,⁷ C. Salgado,^{41,33} Y.G. Sharabian,⁴¹ Iu. Skorodumina,^{39,38} G.D. Smith,⁴³ D.I. Sober,⁶ D. Sokhan,⁴⁴ N. Sparveris,⁴⁰ S. Strauch,³⁹ M. Taiuti,^{17,24} J.A. Tan,³⁰ M. Ungaro,^{41,36} H. Voskanyan,⁴⁹ E. Voutier,²⁷ R. Wang,²⁷ D.P. Watts,⁴³ L.B. Weinstein,³⁵ C.S. Whisnant,²⁹ M.H. Wood,^{5,39} N. Zachariou,⁴³ J. Zhang,⁴⁷ Z.W. Zhao,¹¹ and I. Zonta³⁷

(The CLAS Collaboration)

¹*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213*

²*Argonne National Laboratory, Argonne, Illinois 60439*

³*Arizona State University, Tempe, Arizona 85287-1504*

⁴*California State University, Dominguez Hills, Carson, CA 90747*

⁵*Canisius College, Buffalo, NY, 14208*

⁶*Catholic University of America, Washington, D.C. 20064*

⁷*IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France*

⁸*Université Blaise Pascal, Clermont-Ferrand, Aubière Cedex 63178, France*

⁹*Christopher Newport University, Newport News, Virginia 23606*

¹⁰*University of Connecticut, Storrs, Connecticut 06269*

¹¹*Duke University, Durham, North Carolina 27708-0305*

¹²*Fairfield University, Fairfield CT 06824*

¹³*Università di Ferrara, 44121 Ferrara, Italy*

¹⁴*Florida International University, Miami, Florida 33199*

¹⁵*Florida State University, Tallahassee, Florida 32306*

¹⁶*Particle and Nuclear Physics Institute, Orlova Roshka 1, 188300 Gatchina, Russia*

¹⁷*Università di Genova, Dipartimento di Fisica, 16146 Genova, Italy*

¹⁸*The George Washington University, Washington, DC 20052*

¹⁹*Helmholtz-Institute für Strahlen- und Kernphysik der Rheinischen Friedrich-Wilhelms Universität, Nussallee 14-16, 53115 Bonn, Germany*

- ²⁰ *Idaho State University, Pocatello, Idaho 83209*
- ²¹ *University of Iowa, Iowa City, IA 52242*
- ²² *INFN, Sezione di Ferrara, 44100 Ferrara, Italy*
- ²³ *INFN, Laboratori Nazionali di Frascati, 00044 Frascati, Italy*
- ²⁴ *INFN, Sezione di Genova, 16146 Genova, Italy*
- ²⁵ *INFN, Sezione di Roma Tor Vergata, 00133 Rome, Italy*
- ²⁶ *INFN, Sezione di Torino, 10125 Torino, Italy*
- ²⁷ *Institut de Physique Nucléaire, CNRS/IN2P3 and Université Paris Sud, Orsay, France*
- ²⁸ *Institute of Theoretical and Experimental Physics, Moscow, 117259, Russia*
- ²⁹ *James Madison University, Harrisonburg, Virginia 22807*
- ³⁰ *Kyungpook National University, Daegu 41566, Republic of Korea*
- ³¹ *Mississippi State University, Mississippi State, MS 39762-5167*
- ³² *University of New Hampshire, Durham, New Hampshire 03824-3568*
- ³³ *Norfolk State University, Norfolk, Virginia 23504*
- ³⁴ *Ohio University, Athens, Ohio 45701*
- ³⁵ *Old Dominion University, Norfolk, Virginia 23529*
- ³⁶ *Rensselaer Polytechnic Institute, Troy, New York 12180-3590*
- ³⁷ *Universita' di Roma Tor Vergata, 00133 Rome Italy*
- ³⁸ *Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, 119234 Moscow, Russia*
- ³⁹ *University of South Carolina, Columbia, South Carolina 29208*
- ⁴⁰ *Temple University, Philadelphia, PA 19122*
- ⁴¹ *Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606*
- ⁴² *Universidad Técnica Federico Santa María, Casilla 110-V Valparaíso, Chile*
- ⁴³ *Edinburgh University, Edinburgh EH9 3JZ, United Kingdom*
- ⁴⁴ *University of Glasgow, Glasgow G12 8QQ, United Kingdom*
- ⁴⁵ *Virginia Commonwealth University, Richmond, VA 23220*
- ⁴⁶ *Virginia Tech, Blacksburg, Virginia 24061-0435*
- ⁴⁷ *University of Virginia, Charlottesville, Virginia 22901*
- ⁴⁸ *College of William and Mary, Williamsburg, Virginia 23187-8795*
- ⁴⁹ *Yerevan Physics Institute, 375036 Yerevan, Armenia*

(Dated: September 17, 2018)

We report the first measurements of the E beam-target helicity asymmetry for the $\bar{\gamma}n \rightarrow K^0\Lambda$, and $K^0\Sigma^0$ channels in the energy range of $1.70 \leq W \leq 2.34$ GeV. The CLAS system at Jefferson Lab used a circularly polarized photon beam and a target consisting of longitudinally polarized solid molecular hydrogen deuteride (HD) with low background contamination for the measurements. The multivariate analysis method of Boosted Decision Trees was used to isolate the reactions of interest. Comparisons with predictions from the KaonMAID, SAID, and Bonn-Gatchina models are presented. These results will help separate the isospin $I = 0$ and $I = 1$ photo-coupling transition amplitudes in pseudoscalar meson photoproduction.

* Current address: GAC R&D Center Silicon Valley Inc.

† Contact: schumacher@cmu.edu

‡ Current address: Mississippi State University, Mississippi State, MS 39762-5167

§ Current address: The George Washington University, Washington, DC 20052

¶ Current address: University of Kentucky, Lexington, KY 40506

** Current address: Idaho State University, Pocatello, Idaho 83209

I. INTRODUCTION

An accurate description of excited nucleons and their interaction with probes such as photons at GeV energies has remained elusive for decades. The Standard Model [1, 2] underpins the structure of the nucleons and their excitations, but in the low-energy non-perturbative regime, competing semi-phenomenological models of specific reaction dynamics are all that are available. Present-day lattice QCD calculations [3, 4] and quark models [5–10] predict a richer baryon spectrum than experimentally observed [11–13] —the so-called *missing resonance problem*. There are theoretical approaches for the nucleon resonance spectrum that predict some quark-model states do not exist, including models with quasi-stable diquarks [14], AdS/QCD string-based models [15], and “molecular” models in which some baryon resonances are dynamically generated from the unitarized interaction among ground-state baryons and mesons [16]. But finding such missing states may in part be an experimental problem: high-mass nucleon resonances may couple weakly to πN and may thus have escaped detection in the analysis of πN elastic scattering experiments. Further, they are wide and overlapping, and partial wave analysis (PWA) of reaction data for specific final states remains difficult due to channel coupling effects and insufficient experimental constraints. The experimental results discussed here represent one step in the direction of adding constraints to the hyperon photoproduction database, which ultimately impacts models for nucleon excitations.

Cross section measurements alone are not enough to constrain PWA models of meson production amplitudes. Polarization observables related to the spins of the beam photons, target, and recoiling baryons are also needed. Photoproduction of pseudoscalar mesons is governed by four complex amplitudes that lead to an interaction cross sections and 15 spin observables [17–23]. A *mathematically complete* experiment would require data, with negligible uncertainties, on a minimum of eight well-chosen observables at each center-of mass (c.m.) energy, W , and meson polar angle, $\cos\theta_{c.m.}$. In practice, with realistically achievable uncertainties, measurements of many more are needed to select between competing partial wave solutions, and even knowledge of the

sign of an asymmetry can provide valuable discrimination [22]. Furthermore, avoiding ambiguities in PWA solutions requires measurements of observables from each spin configuration of the three combinations of beam-target, target-recoil and beam-recoil polarization [22, 23].

Furthermore, while isospin $I = 3/2$ transitions (Δ^* excitations) can be studied with proton target data alone, both proton- and neutron-target observables are necessary to study $I = 1/2$ transitions and isolate the separate $\gamma p N^*$ proton and $\gamma n N^*$ neutron photocouplings [24]. Information from neutron targets is comparatively scarce [25], particularly in the hyperon channels [26, 27], which is why the present measurement is of value. Furthermore, the hyperon photoproduction channels $\gamma N \rightarrow K\Lambda(\Sigma^0)$ are attractive for analysis for two reasons. First, the threshold for two-body hyperon final states is at $W \simeq 1.6$ GeV, above which lie numerous poorly-known resonances. Two-body strange decay modes, rather than cascading non-strange many-body decays, may be easier to interpret. Second, the hyperon channels give easy access to recoil polarization observables on account of their self-analyzing weak decays. While the present work does not involve measurement of hyperon polarizations, previous work has shown the benefit of using such information to extract properties of higher-mass nucleon resonances [28–35]. Thus, pursuing “complete” amplitude information in the hyperon photoproduction channels can be complimentary to the analogous quest in, say, pion photoproduction.

In this article, we present first-time measurements of the beam-target observable E on a longitudinally polarized neutron bound in deuterium in the quasi-free reaction $\gamma n(p) \rightarrow K^0 Y(p)$. The helicity asymmetry E is formally defined as the normalized difference in photoproduction yield between anti-parallel (σ^A) and parallel (σ^P) configurations, *i.e.*, settings where the incident photon beam polarization is anti-aligned or aligned, respectively, with the longitudinal polarization of the target. Following Refs. [18, 22] write

$$E = \frac{\sigma^A - \sigma^P}{\sigma^A + \sigma^P}. \quad (1)$$

This helicity asymmetry, E , is related to the cross section by

$$\left(\frac{d\sigma}{d\Omega}\right) = \left(\frac{d\sigma}{d\Omega}\right)_0 (1 - P_T P_\odot E), \quad (2)$$

where $(d\sigma/d\Omega)_0$ is the differential cross section averaged over initial spin states and summed over the final states, and P_T and P_\odot are the target longitudinal and beam circular polarizations, respectively.

The asymmetry results obtained will be compared with several model predictions. The first is a single-channel effective Lagrangian approach, KaonMAID [36, 37], with parameter constraints largely imposed from SU(6). Without experimental constraints on the $N^*\Lambda K^0$ and $\gamma n N^*$ vertices, the reaction of interest is difficult to model accurately. The second model giving predictions for the present results is the data description given by SAID [38, 39]. In general, SAID is more up to date than KaonMAID; for the present reaction channels the SAID predictions are a polynomial fit to all available data before 2008, assuming final state interactions for these polarization observables can be neglected [40]. The third comparison is made to the multi-channel K-matrix formalism of the Bonn-Gatchina [41] group, which is the most up to date, being constrained by recent first-time measurements [26] of the differential cross section for the reaction $\gamma n(p) \rightarrow K^0 \Lambda(p)$ (with (p) as the spectator proton).

II. EXPERIMENTAL PROCEDURES

The experiment was performed at the Thomas Jefferson National Accelerator Facility (JLab) using the CEBAF Large Acceptance Spectrometer (CLAS) [42]. This setup has been used for several studies of K^+ photoproduction of hyperonic final states on a proton target [29–31, 33, 34, 43–45] and on an effective neutron (deuteron) target [26, 27]. The present results stem from the so-called “g14” run period between December 2011 and May 2012, from which non-strange results have been previously reported [46]. The CEBAF accelerator provided longitudinally polarized electron beams with energies of $E_e = 2.281$ GeV, 2.257 GeV, and 2.541 GeV, and an *average* electron beam polarization for the present study of $P_e = 0.82 \pm 0.04$, which was measured routinely by the Hall-B Möller polarimeter [47]. The electron beam helicity was pseudo-randomly flipped between +1 and -1 with a 960 Hz flip rate. The electron beam was incident on the thin gold radiator of the Hall-B Tagger sys-

tem [48] and produced circularly polarized tagged photons. The polarization of the photons was determined using the Maximon and Olsen formula [49]

$$P_\odot = P_e \frac{4k - k^2}{4 - 4k + 3k^2}, \quad (3)$$

where P_\odot and P_e are the photon and electron polarizations, respectively, and $k = E_\gamma/E_e$ is the ratio between the photon energy and the electron beam energy.

A 5-cm-long solid target of hydrogen deuteride (HD) was used in the experiment [50, 51]. It achieved vector polarizations of 25-30% for deuterons, i.e. for *bound* neutrons in the deuteron with relaxation times of about a year. The polarized target was held at the center of CLAS using an in-beam cryostat (IBC) that produced a 0.9 T holding field and operated at 50 mK. The target polarization was monitored using nuclear magnetic resonance measurements [50]. The orientation of the target longitudinal polarization direction was inverted between periods of data taking, either parallel or anti-parallel to the direction of the incoming photon beam. Background events from the unpolarizable target wall material and aluminum cooling wires [51] were removed using empty-target data, as discussed in Sec. III A and III B.

The specific reaction channel for this discussion came from events of the type $\gamma d \rightarrow \pi^+ \pi^- \pi^- p(X)$ using a read-out trigger requiring a minimum of two charged particles in different CLAS sectors. After particle identification we required the “spectator”, X , to be an undetected low-momentum proton and possibly a photon, via the missing mass technique, as explained in the next section. In order to determine the E asymmetry experimentally, the event yields in a given kinematic bin of W and kaon center-of-mass angle were obtained by counting events with total c.m. helicity $h = 3/2$ (laboratory frame anti-parallel configuration) called N_A and $1/2$ (laboratory frame parallel configuration) called N_P , respectively. The E observable was then computed as

$$E = \frac{1}{\overline{P_T} \cdot \overline{P_\odot}} \left(\frac{N_A - N_P}{N_A + N_P} \right), \quad (4)$$

where $\overline{P_T}$ and $\overline{P_\odot}$ are the run-averaged target and beam polarizations, respectively.

III. DATA ANALYSIS

The performance of the system was extensively studied for a reaction with much higher count-rates than the present one. The non-strange reaction $\gamma d \rightarrow \pi^- p(X)$ was investigated using many of the same analysis steps and methods discussed in this article to extract the E observable for $\gamma n \rightarrow \pi^- p$ [46]. The analysis steps outlined below were all tested on that reaction. In particular, the Boosted Decision Tree (BDT) selection procedure [52, 53] used below was validated against alternative “cut-based” and kinematic fit methods, with the result that the BDT procedure resulted in $\sim 30\%$ larger yields of signal events and therefore gave better statistical precision on the final E asymmetry.

A. Particle identification

For this particular analysis, we required that every selected event consists of at least two positive tracks and two negative tracks with associated photon tagger hits [48]. The CLAS detector system determined the path length, the charge type, the momentum and the flight time for each track [54–56]. For each track of momentum \vec{p} , we compared the measured time of flight, TOF_m , to a hadron’s expected time of flight, TOF_h , for a pion and proton of identical momentum and path length. CLAS-standard cuts were placed on the difference between the measured and expected time of flight, $\Delta TOF = TOF_m - TOF_h$. We selected events for which the two positively charged particles were the proton and π^+ , and the two negatively charged were both π^- . Well-established CLAS fiducial cuts were applied to select events with good spatial reconstruction.

Events originating from unpolarized target material—aluminum cooling wires and polychlorotrifluoroethylene (pCTFE)—dilute E and must be taken into account. A period of data taking was dedicated to an *empty* target cell in which the HD material was not present. This set of data was used to study and remove the bulk of the target material background on the basis of a loose missing mass cut. Figure 1 shows the resulting reconstructed reaction vertex for 4-track data along the beam line for both a full target and for an empty target




FIG. 1. The vertex distribution of events along the beam line for a full target is the open histogram. The dashed red lines show the nominal target boundaries. The peaks at $z > 0$ are from target-independent foils in the cryostat; the positions of two are highlighted with dotted blue lines [50]. The filled histogram is the scaled target-empty background distribution.

scaled to match the counts in several downstream target foils. The full-to-empty ratio of about 3.3:1 in the target region was important in selecting the optimal BDT cut discussed below.

Figure 2 shows the resulting target-full missing mass distribution for spectator X in $\gamma d \rightarrow \pi^- \pi^+ \pi^- p(X)$, after these cuts. A clear peak corresponding to the spectator proton is seen at Point 1 for events that produced a Λ particle. A loose cut was applied to reject events with missing mass larger than $1.4 \text{ GeV}/c^2$ at Point 4 because of the presence of $\Sigma^0 \rightarrow \pi^- p(\gamma)$ events. These have a 73 MeV photon in the final state in addition to the proton, and the distribution peaks at Point 2 and has a kinematic tail to about Point 3.

B. $K^0 Y$ event selection using BDT analysis

Because of the small reaction cross section in this experiment, a method was needed to optimally isolate the events of interest with minimal statistics loss. The multivariate analysis tool called the Boosted Decision Tree (BDT) approach was used to select the exclusive events of interest in this study. Three steps were needed to achieve this result. The first BDT was created to select events from both the $\gamma d \rightarrow \pi^- \pi^+ \pi^- p(p_S)$ and




FIG. 2. The missing mass distribution, $\gamma d \rightarrow \pi^- \pi^+ \pi^- p X$ after PID cuts showing the dominant spectator proton peak at “1”. The magenta line at “4” indicates a loose event rejection for $m_X > 1.4 \text{ GeV}/c^2$. This rejects unambiguous background but keeps $\Sigma^0 \rightarrow \pi^- p(\gamma)$ events in which both a proton and a photon are missing between “2” and “3”. (See text.)

$\gamma d \rightarrow \pi^- \pi^+ \pi^- p(p_S \gamma)$ final states, consistent with quasi-free production from a deuteron. This was to reject target-material background and events with high missing momentum of the undetected spectator nucleon, p_S . The second BDT was created to remove the non-strange pionic background with the same final states, that is, to pick out events with Λ and Σ^0 intermediate-state particles. The third BDT was to separate the $K^0 \Lambda$ and $K^0 \Sigma^0$ events.

This BDT algorithm is more efficient than a simple “cut” method in both rejecting background and keeping signal events [53, 57]. The method builds a “forest” of *distinct decision trees* that are linked together by a *boosting* mechanism. Each decision tree constitutes a *disjunction* of logical conjunctions (i.e., a graphical representation of a set of *if-then-else* rules). Thus, the entire reaction phase-space is considered by every decision tree. Before employing the BDT for signal and background classification, the BDT algorithm needs to be constructed (or trained) with *training* data—wherein the category of every event is definitively known. We used the ROOT implementation of the BDT algorithm [58]. Every event processed by the constructed BDT algorithm is assigned a value between -1 and $+1$ that quantifies how likely the processed event is a background event (closer to -1) or a signal event (closer to $+1$). An optimal cut on the BDT output is chosen to maximize the $S/\sqrt{S+B}$ ratio,

where S , B are the estimations, based on training data, of the initial number of signal and background events, respectively.

The initial assignment of the π^- particles to either K^0 or Λ decay was studied with Monte Carlo simulation, and a loose selection based on invariant masses was made. Specific details of these cuts are found in Ref. [57].

The first BDT was trained using real empty-target data for the background training. A signal Monte-Carlo simulating quasi-free hyperon production on the neutron was used for signal training data. The momentum distribution of the spectator proton, p_s , followed the Hulthén potential [59, 60] for the deuteron. Based on this training, an optimal BDT cut that maximized the estimated initial $S/\sqrt{S+B}$ ratio was selected. Figure 3 shows the total (blue histogram) and rejected (black histogram) events by the first BDT cut. When comparing Figs. 1 and 3, two items should be noted. Firstly, the BDT was trained to remove target-material background events with missing momentum not consistent with a Hulthén distribution. Secondly, the BDT background-rejection efficiency was not perfect, leaving some target-material background events that was removed in a subsequent step (Sec. III C). We then rejected events with $z > -2 \text{ cm}$ on the reaction vertex to remove remaining unambiguous background events due to various cryostat foils.

The second-step BDT was trained using a 4-body phase-space $\gamma d \rightarrow \pi^- \pi^+ \pi^- p(p_S)$ simulation as background training data and the $\gamma d \rightarrow K^0 \Lambda(p_S)$ simulation as signal training data. There were two negative pions in each event: one from the decay of the K^0 and one from the decay of the hyperon. The goal of the BDT analysis was to use the available correlations among all particles to sort the pions correctly and to select events with decaying strange particles. The main training variables at this stage of the analysis included the 3-momenta of all the particles and the detached decay vertices of the K^0 s and the hyperons. After the optimized BDT cut was placed, Fig. 4 shows the total (red histogram) and rejected (black histogram) events after this second BDT analysis step. The efficiency of the second BDT was less than 100%, thus, there are remaining target background events in the selected data sample. The dips near the signal maxima in the background spectra show that the background is slightly undersubtracted. This issue is dis-




FIG. 3. The reconstructed distribution of the reaction vertex along the beam line showing target-full events in the top histogram (blue) after the loose $K^0 Y^0$ selection and the missing mass cut shown in Fig. 2. Events selected by the first BDT are the middle histogram (red), and the rejected events are the bottom histogram (black). The magenta line indicates a loose cut to reject unambiguous target-material background.

cussed and corrected below. A fit with a Breit-Wigner line shape and a polynomial was used to estimate that the strange-to-non-strange ratio of events in the data set at this stage was about 2.3:1 in the peak regions.

For the final task, separating the $K^0 \Lambda$ and $K^0 \Sigma^0$ channels, the third BDT was trained using $\gamma d \rightarrow K^0 \Sigma^0(p_S)$ simulation as “background” training data and $\gamma d \rightarrow K^0 \Lambda(p_S)$ simulation as “signal” training data. Note that the term “background” used here is just for semantic convenience, since both channels were retained after applying the third optimized BDT cut. Figure 5 shows in the left (right) histogram the classification success of the third BDT on $\gamma d \rightarrow K^0 \Lambda(p_S)$ ($\gamma d \rightarrow K^0 \Sigma^0(p_S)$) simulation data. The histograms reveal that a small number of $K^0 \Lambda$ events would be misclassified as $K^0 \Sigma^0$ events and vice-versa. In the next section, the correction for the contamination on both final data sets will be discussed. Figure 6 shows the separation result from the third BDT on real data.




FIG. 4. The invariant $\pi_{\Lambda}^- p$ mass (top) and invariant $\pi_{K^0}^- \pi^+$ mass (bottom) after target material background rejection by the first BDT cut. The black histograms show events rejected by the second BDT cut. Fits of a sum (red) of a Breit-Wigner line-shape (blue) and a 3rd order polynomial (black) are shown. The fits aid the discussion in the text but were not used in the subsequent analysis.

C. Corrections for remaining backgrounds and asymmetry calculation

The E asymmetry values for both target-material and non-strange background events were statistically consistent with zero [57]; therefore, we implemented an approximation procedure to correct for the dilution effect from the remaining background. We estimated two ratios: one for the remaining fraction of target background (TGT), R^{TGT} , and one for the fraction of remaining non-strange (NS) final-state events mixed with the hyperon events, R^{NS} . We write $R^{TGT} = N^{remain}/N^{HD}$,




FIG. 5. The distributions of missing mass from the reconstructed K^0 , $\gamma n \rightarrow \pi_{K^0}^- \pi^+ X$ for simulation data, assuming that the target is an at-rest neutron. On the left, the magenta histogram represents events with correct $K^0\Lambda$ classification, while the cyan histogram represents events with the *wrong* $K^0\Sigma^0$ classification. On the right, the cyan histogram represents events with the correct $K^0\Sigma^0$ classification, while the magenta histogram represents events with the *wrong* $K^0\Lambda$ classification.




FIG. 6. The distribution of missing mass from the reconstructed K^0 , $\gamma n \rightarrow \pi_{K^0}^- \pi^+ X$ for real data, assuming that the target is an at-rest neutron, after rejecting non-hyperon background by the second BDT cut. The magenta (cyan) histogram was classified as $K^0\Lambda$ ($K^0\Sigma^0$) using the third BDT selection step.

and $R^{NS} = Y^{remain}/Y^{K^0Y}$. N^{remain} and N^{HD} are the estimated number of remaining target-material background events and the true deuteron events after the first BDT and $z = -2$ cm vertex cuts, respectively. Y^{remain} and Y^{K^0Y} are the estimated number of remaining non-strange and true K^0Y events after the second BDT cut,

respectively. Next, let Y_{BDT} be the number of events that passed the z -vertex cut and the first two BDT selections, then Y_{BDT} can be partitioned into

$$\begin{aligned} Y_{BDT} &= (1 + R^{NS}) Y^{K^0Y} \\ &= (1 + R^{NS}) [Y_{HD}^{K^0Y} + Y_{TGT}^{K^0Y}], \end{aligned} \quad (5)$$

since Y^{K^0Y} also comprises events from the remaining target-material background and the bound signal events. If we further allow $Y_{TGT}^{K^0Y}/Y_{HD}^{K^0Y} = N^{remain}/N^{HD} = R^{TGT}$, then Y_{BDT} can finally be expressed as:

$$Y_{BDT} = (1 + R^{NS}) (1 + R^{TGT}) Y_{HD}^{K^0Y}, \quad (6)$$

or

$$Y_{HD}^{K^0Y} = (1 + R^{NS})^{-1} (1 + R^{TGT})^{-1} Y_{BDT}. \quad (7)$$

These relations should remain valid for both $Y_{BDT}^{K^0\Lambda}$ and $Y_{BDT}^{K^0\Sigma^0}$, which are the $K^0\Lambda$ and $K^0\Sigma^0$ signal events from bound neutrons, respectively. The backgrounds that leak through the BDT filters will be helicity independent and will subtract in the numerator of Eq. 4. Using Eq. 7 to correct the summed yields in the denominator gives the corrected asymmetry as

$$E_{corrected}^{K^0Y} = (1 + R^{NS}) \times (1 + R^{TGT}) E_{BDT}^{K^0Y}, \quad (8)$$

where $E_{BDT}^{K^0Y}$ is obtained from $Y_{BDT}^{K^0Y}$ (or, more exactly, Y_{BDT}^P and Y_{BDT}^A of the K^0Y parallel and anti-parallel

subsets). From the simulations we found average values of R^{TGT} and R^{NS} of 0.09 and 0.17, respectively, with some dependence on the specific run period.

Next we discuss a correction for the third BDT classification result. Recall that the third BDT selection separates the true signal $K^0 Y$ events into two subsets: one is mostly $K^0 \Lambda$ events, and the other is mostly $K^0 \Sigma^0$. If we denote N_{Λ}^{BDT} and $N_{\Sigma^0}^{BDT}$ as the number of events the third BDT identified as $K^0 \Lambda$ and $K^0 \Sigma^0$ events, respectively, then we have the expressions

$$N_{\Lambda}^{BDT} = \omega_{\Lambda} N_{\Lambda}^{true} + (1 - \omega_{\Sigma^0}) N_{\Sigma^0}^{true}, \quad (9)$$

$$N_{\Sigma^0}^{BDT} = (1 - \omega_{\Lambda}) N_{\Lambda}^{true} + \omega_{\Sigma^0} N_{\Sigma^0}^{true}, \quad (10)$$

where ω_{Λ} and ω_{Σ^0} are the fractions of events correctly identified—these values were estimated based on simulation data. After rearrangement, we arrive at the expressions

$$N_{\Lambda}^{true} = \left[\omega_{\Lambda} - \frac{(1 - \omega_{\Sigma^0})(1 - \omega_{\Lambda})}{\omega_{\Sigma^0}} \right]^{-1} \times \left[N_{\Lambda}^{BDT} - \frac{(1 - \omega_{\Sigma^0})}{\omega_{\Sigma^0}} N_{\Sigma^0}^{BDT} \right], \quad (11)$$

$$N_{\Sigma^0}^{true} = \left[\omega_{\Sigma^0} - \frac{(1 - \omega_{\Lambda})(1 - \omega_{\Sigma^0})}{\omega_{\Lambda}} \right]^{-1} \times \left[N_{\Sigma^0}^{BDT} - \frac{(1 - \omega_{\Lambda})}{\omega_{\Lambda}} N_{\Lambda}^{BDT} \right]. \quad (12)$$

The *corrected* E asymmetry was obtained using the derived N_{Λ}^{true} and $N_{\Sigma^0}^{true}$ by using Eq. 4. From the simulations we found average values of ω_Y of 0.87 and 0.91 for Λ and Σ^0 events, respectively.

The neutron polarization in the deuteron is smaller than the deuteron polarization because the deuteron wavefunction has, in addition to an S-wave component, a D-wave component in which the spin of the neutron need not be aligned with the deuteron spin. This was studied using data for the $\gamma n \rightarrow \pi^- p$ reaction and reported in our previous publication Ref. [46]. It was found that for spectator recoil momenta of less than 100 MeV/c the correction was negligible. Had we cut on recoil momentum at 200 MeV/c rather than 100 MeV/c, a measured dilution factor of $(8.6 \pm 0.1)\%$ would have been necessary for the non-strange channel. But different reaction channels

may exhibit different sensitivities to recoil momentum. For the reaction under discussion here we could not afford the statistical loss by cutting on recoil momentum, and we elected to make a conservative correction based on the general considerations of Ref. [61]. The neutron polarization can be estimated as $P_n = P_d(1 - \frac{3}{2}P_D)$, where P_n and P_d are neutron and deuteron polarizations, respectively, and P_D denotes the deuteron D-state probability. The latter is not strictly an observable and needs only to be treated consistently within a given NN potential. Following Ref. [61], we take the D-state contribution averaged over a range of NN potentials as about 5%, which implies the neutron polarization is 92.5% of the deuteron polarization, or a 7.5% dilution factor.

D. Systematic Uncertainties

We implemented four systematic studies to quantify the robustness of the trained BDT algorithms and the sensitivity of our results on the correction procedures introduced in the previous section. Two tests studied the effect of loosening the first and the second BDT cuts, respectively. One test focused on the sensitivity of the E results on the third correction—the correction procedure that was implemented to “purify” the final selected $K^0 \Sigma^0 (K^0 \Lambda)$ sample. Lastly we reduced the beam and target polarizations by one standard deviation of their respective total uncertainties (statistical and systematic) to study the changes on the E results.

Finally, we note a complication that could occur when summing Λ yields to create the E asymmetries. The relative angular distribution between the π^- and the p that are used to reconstruct a Λ carries information on the recoil polarization of the latter. When summed over azimuthal angles, this information is lost. However, limitations in detector acceptance could result in an incomplete integration, which in principle could introduce into Eq. 2 a dependence on six additional observables [22]. The gaps in CLAS acceptance are modest and, due to lower than expected production cross sections, the data below are presented in broad kinematic bins, which tends to dilute such effects. On the scale of our statistical uncertainties, such corrections are expected to be negligible and we have not attempted to correct for them.

IV. RESULTS

We present here the results for the E asymmetry in two W energy bins. The lower bin is from 1.70 GeV to 2.02 GeV and denoted as W_1 , while the higher bin is from 2.02 GeV to 2.34 GeV and referred to as W_2 . Due to small cross sections for $K^0 Y$ photoproduction, and to detector inefficiencies that are amplified by the required identification of four charged particles, our statistics are sufficient for only three bins in K^0 center-of-mass production angle. The measurements for the $\gamma n \rightarrow K^0 \Lambda$ reaction are plotted together with predictions from the KaonMAID, SAID, and Bonn-Gatchina (BnGa) models in Fig. 7. The data show that the $K^0 \Lambda$ asymmetry is largely positive below 2 GeV and mostly negative above 2 GeV, without more discernible trends. Values of E must approach $+1$ at $\cos \theta_{K^0}^{c.m.} \rightarrow \pm 1$ to conserve angular momentum. Thus, the values for E in bin W_2 must change rather rapidly near the extreme angles.

For comparison, PWA combine results from many experiments at different energies, and this results in varying degrees of sensitivity to energy and angle. This is illustrated in Fig. 7 by the SAID and BnGa PWA predictions at the limits of the energy bins. None of the models were tuned to these results; that is, the models are all predictions based on fits to previously published data on other observables. First, one observes that the data are not statistically strong enough to strongly discriminate among the models. In the lower W bin all three models can be said to agree with the data. In the higher W bin the SAID model may be slightly favored by the data among the three.

The results for the $\gamma n \rightarrow K^0 \Sigma^0$ channel are plotted in Fig. 8, together with model predictions from SAID and Kaon-MAID. In contrast to the $K^0 \Lambda$ channel at lower W , here the data hint at less positive values for E . In the bin for W above 2 GeV, the data are also consistent with zero for $K^0 \Sigma^0$, whereas the $K^0 \Lambda$ data tended to be negative. In fact, the $K^0 \Sigma^0$ asymmetry is consistent with zero in all available bins. The model comparisons show that the KaonMAID prediction for the $K^0 \Sigma^0$ channel in the higher W bin are probably not consistent with the data, while the SAID result is consistent with the data. For the $K^0 \Sigma^0$ case we do not have predictions from the Bonn-Gatchina model because the unpolarized




FIG. 7. The helicity asymmetry E for the $K^0 \Lambda$ final state (with combined statistical and systematic uncertainties) vs. $\cos \theta_{K^0}$. The asymmetries are shown with the neutron-target theoretical models KaonMaid [36] (red dashed) and SAID [38] (blue dot-dashed) and Bonn-Gatchina [32, 41] (solid black). Because of the 0.32 GeV-wide W bins, each model is represented by two curves, computed at the bin endpoint W values, as labeled.

differential cross section has not been measured yet, and without it the model does not have a prediction available.

In order to show one other comparison between data and theory, we plot some of the present results for a neutron target together with the model predictions for the $K^+ \Lambda$ reaction on a *proton* target in Fig. 9. This is intended to show the difference between the model predictions on the proton and the neutron. One sees how different the three model predictions are for protons versus neutrons. One notes that the predictions for the proton target calculations all tend to be closer to the new data we are presenting for a neutron target. This suggests that calculations of the E observable for a neutron target can be improved. Thus, we may expect these present results to have some impact on the further development of these models.

So-far unpublished CLAS results for the corresponding reaction $\gamma p \rightarrow K^+ \Lambda$ have higher statistics and finer energy bins than the present results (since the identification of this final state requires the detection of fewer particles). The present $K^0 \Lambda$ results are, within our uncertainties, similar to the $K^+ \Lambda$ asymmetries in Ref. [62].




FIG. 8. The helicity asymmetry E for the $K^0\Sigma^0$ final state (with combined statistical and systematic uncertainties) vs. $\cos\theta_{K^0}$ for two 0.32 GeV-wide energy bands in W , as labeled. The model curves are as for the previous figure.




FIG. 9. The helicity asymmetry E for the $K\Lambda$ final state vs. $\cos\theta_{K^0}$ for energy band W_2 . On the left are the data from Fig. 7 together with model predictions for a NEUTRON target. On the right are model calculations for the $K^+\Lambda$ reaction on a PROTON target, as computed using KaonMaid [36] (red dashed), SAID [38] (blue dot-dashed) and Bonn-Gatchina [32, 41] (black and black-dashed). The curves on the right are closer to the (reaction mismatched) data shown on the left.

The numerical values of the measured $K^0\Lambda$ and $K^0\Sigma^0$ E asymmetries, together with their statistical and systematic uncertainties, are reported in Table I.

V. CONCLUSIONS

We have reported the first set of the E asymmetry measurements for the reaction $\gamma d \rightarrow K^0 Y(p_s)$ for $1.70 \text{ GeV} \leq W \leq 2.34 \text{ GeV}$. In particular, we described the three-step BDT-based analysis method developed to select a clean sample of $p\pi^+\pi^-\pi^-$ with intermediate hyperons. We have plotted the E asymmetry as a function of $\cos\theta_{K^0}^{CM}$. Several systematic uncertainty tests led to the conclusion that statistical uncertainties dominated the final results. The numerical values of the measured E asymmetries and their statistical and systematic uncertainties are reported in Table I.

Evidently, this analysis is limited by the small cross sections of the channels of interest, leading to large uncertainties on the measurements of the E asymmetry. At present, comparison with several models makes no decisive selections among the model approaches. Overall, the BnGa predictions are of similar quality to the SAID predictions. The Kaon-MAID predictions for both channels seem less successful. Among all three model comparisons, the distinction between proton and neutron target predictions are differentiated by the data: The proton-target predictions compare better than the neutron-target predictions with the experimental results. In principle, this information is valuable since it hints at the necessary isospin decomposition of the hyperon photoproduction mechanism.

At present, multipole analyses for the $K^0 Y$ channels are severely limited by available data. Higher statistics data on these channels for a number of other polarization observables, from a much longer (unpolarized) target, have been collected during the $g13$ running period with CLAS and is under analysis. A greater number of different polarization observables is generally more effective than precision at determining a photoproduction amplitude [22]. When these $g13$ results become available, the present data on the beam-target E asymmetry are likely to have a larger impact.

		$\cos \theta_{K^0}$		
		-0.6	0.0	+0.6
$K^0\Lambda$	W_1	$0.834 \pm 0.499 \pm 0.287$	$-0.144 \pm 0.436 \pm 0.098$	$1.066 \pm 0.419 \pm 0.231$
	W_2	$-0.533 \pm 0.752 \pm 0.345$	$-0.263 \pm 0.618 \pm 0.101$	$-0.648 \pm 0.464 \pm 0.136$
$K^0\Sigma^0$	W_1	$-0.110 \pm 0.723 \pm 0.406$	$0.581 \pm 0.539 \pm 0.144$	$-0.319 \pm 0.541 \pm 0.460$
	W_2	$-0.471 \pm 0.446 \pm 0.391$	$0.0002 \pm 0.317 \pm 0.150$	$0.054 \pm 0.281 \pm 0.065$

TABLE I. Numerical values of the E asymmetry measurements for the $K^0\Lambda/K^0\Sigma^0$ channels. The uncertainties are statistical and systematic, respectively. The center-of-mass energy ranges are $1.70 < W_1 < 2.02$ GeV and $2.02 < W_2 < 2.34$ GeV.

ACKNOWLEDGMENTS

We acknowledge the outstanding efforts of the staff of the Accelerator and Physics Divisions at Jefferson Lab that made this experiment possible. The work of the Medium Energy Physics group at Carnegie Mellon University was supported by DOE grant DE-FG02-87ER40315. The Southeastern Universities Research Association (SURA) operated the Thomas Jefferson National Accelerator Facility for the United States Depart-

ment of Energy under contract DE-AC05-84ER40150. Further support was provided by the National Science Foundation, the Chilean Comisión Nacional de Investigación Científica y Tecnológica (CONICYT), the French Centre National de la Recherche Scientifique, the French Commissariat à l’Energie Atomique, the Italian Istituto Nazionale di Fisica Nucleare, the National Research Foundation of Korea, the Scottish Universities Physics Alliance (SUPA), the United Kingdom’s Science and Technology Facilities Council.

-
- [1] David J. Gross and Frank Wilczek, “Ultraviolet Behavior of Nonabelian Gauge Theories,” *Phys. Rev. Lett.* **30**, 1343–1346 (1973).
- [2] H. David Politzer, “Reliable Perturbative Results for Strong Interactions?” *Phys. Rev. Lett.* **30**, 1346–1349 (1973).
- [3] Robert G. Edwards, Jozef J. Dudek, David G. Richards, and Stephen J. Wallace, “Excited state baryon spectroscopy from lattice QCD,” *Phys. Rev.* **D84**, 074508 (2011).
- [4] R. G. Edwards, N. Mathur, D. G. Richards, and S. J. Wallace (Hadron Spectrum), “Flavor structure of the excited baryon spectra from lattice QCD,” *Phys. Rev.* **D87**, 054506 (2013).
- [5] Simon Capstick and W. Roberts, “Quark models of baryon masses and decays,” *Prog. Part. Nucl. Phys.* **45**, S241–S331 (2000).
- [6] Simon Capstick and W. Roberts, “Strange decays of non-strange baryons,” *Phys. Rev.* **D58**, 074011 (1998).
- [7] Simon Capstick and Nathan Isgur, “Baryons in a Relativized Quark Model with Chromodynamics,” *Proceedings, International Conference on Hadron Spectroscopy: College Park, Maryland, April 20-22, 1985*, *Phys. Rev.* **D34**, 2809 (1986), [AIP Conf. Proc. 132, 267 (1985)].
- [8] Ulrich Loring, Bernard C. Metsch, and Herbert R. Petry, “The Light baryon spectrum in a relativistic quark model with instanton induced quark forces: The Nonstrange baryon spectrum and ground states,” *Eur. Phys. J.* **A10**, 395–446 (2001).
- [9] L. Ya. Glozman, Willibald Plessas, K. Varga, and R. F. Wagenbrunn, “Unified description of light and strange baryon spectra,” *Phys. Rev.* **D58**, 094030 (1998).
- [10] M. M. Giannini, E. Santopinto, and A. Vassallo, “Hypercentral constituent quark model and isospin dependence,” *Eur. Phys. J.* **A12**, 447–452 (2001).
- [11] C. Patrignani *et al.* (Particle Data Group), “Review of Particle Physics,” *Chin. Phys.* **C40**, 100001 (2016).
- [12] Eberhard Klempt and Jean-Marc Richard, “Baryon spectroscopy,” *Rev. Mod. Phys.* **82**, 1095–1153 (2010).
- [13] Roman Koniuk and Nathan Isgur, “Where Have All the Resonances Gone? An Analysis of Baryon Couplings in a Quark Model With Chromodynamics,” *Baryon 1980:217*, *Phys. Rev. Lett.* **44**, 845 (1980).
- [14] Mauro Anselmino, Enrico Predazzi, Svante Ekelin, Sverker Fredriksson, and D. B. Lichtenberg, “Diquarks,” *Rev. Mod. Phys.* **65**, 1199–1234 (1993).
- [15] Stanley J. Brodsky, “Hadron Spectroscopy and Structure from AdS/CFT,” *Eur. Phys. J.* **A31**, 638–644 (2007).
- [16] E. E. Kolomeitsev and M. F. M. Lutz, “On baryon resonances and chiral symmetry,” *Phys. Lett.* **B585**, 243–252

- (2004).
- [17] G. F. Chew, M. L. Goldberger, F. E. Low, and Yoichiro Nambu, “Relativistic dispersion relation approach to photomeson production,” *Phys. Rev.* **106**, 1345–1355 (1957).
- [18] I. S. Barker, A. Donnachie, and J. K. Storrow, “Complete Experiments in Pseudoscalar Photoproduction,” *Nucl. Phys.* **B95**, 347–356 (1975).
- [19] C. G. Fasano, Frank Tabakin, and Bijan Saghai, “Spin observables at threshold for meson photoproduction,” *Phys. Rev.* **C46**, 2430–2455 (1992).
- [20] Wen-Tai Chiang and Frank Tabakin, “Completeness rules for spin observables in pseudoscalar meson photoproduction,” *Phys. Rev.* **C55**, 2054–2066 (1997).
- [21] Greg Keaton and Ron Workman, “Ambiguities in the partial wave analysis of pseudoscalar meson photoproduction,” *Phys. Rev.* **C54**, 1437–1440 (1996).
- [22] A. M. Sandorfi, S. Hoblit, H. Kamano, and T. S. H. Lee, “Determining pseudoscalar meson photo-production amplitudes from complete experiments,” *J. Phys.* **G38**, 053001 (2011).
- [23] J. Nys, J. Ryckebusch, D. G. Ireland, and D. I. Glazier, “Model discrimination in pseudoscalar-meson photoproduction,” *Phys. Lett.* **B759**, 260–265 (2016).
- [24] A. M. Sandorfi and S. Hoblit, “Hyperon photoproduction from polarized H and D towards a *complete* N^* experiment,” *Proceedings, 11th International Conference on Hypernuclear and Strange Particle Physics (HYP 2012): Barcelona, Spain, October 1-5, 2012*, *Nucl. Phys.* **A914**, 538–542 (2013).
- [25] A. V. Anisovich, V. Burkert, N. Compton, K. Hicks, F. J. Klein, E. Klempt, V. A. Nikonov, A. M. Sandorfi, A. V. Sarantsev, and U. Thoma, “Neutron helicity amplitudes,” *Phys. Rev.* **C96**, 055202 (2017).
- [26] N. Compton *et al.* (CLAS), “Measurement of the differential and total cross sections of the $\gamma d \rightarrow K^0 \Lambda(p)$ reaction within the resonance region,” *Phys. Rev. C* **96**, 065201 (2017).
- [27] S. Anefalos Pereira *et al.* (CLAS), “Differential cross section of $\gamma n \rightarrow K^+ \Sigma^-$ on bound neutrons with incident photons from 1.1 to 3.6 GeV,” *Phys. Lett.* **B688**, 289–293 (2010).
- [28] C. A. Paterson *et al.* (CLAS), “Photoproduction of Λ and Σ^0 hyperons using linearly polarized photons,” *Phys. Rev.* **C93**, 065201 (2016).
- [29] R. Bradford *et al.* (CLAS), “First measurement of beam-recoil observables C_x and C_z in hyperon photoproduction,” *Phys. Rev.* **C75**, 035205 (2007).
- [30] R. Bradford *et al.* (CLAS), “Differential cross sections for $\gamma + p \rightarrow K^+ + Y$ for Λ and Σ^0 hyperons,” *Phys. Rev.* **C73**, 035202 (2006).
- [31] J. W. C. McNabb *et al.* (CLAS), “Hyperon photoproduction in the nucleon resonance region,” *Phys. Rev.* **C69**, 042201 (2004).
- [32] A. V. Anisovich, V. Kleber, E. Klempt, V. A. Nikonov, A. V. Sarantsev, and U. Thoma, “Baryon resonances and polarization transfer in hyperon photoproduction,” *Eur. Phys. J.* **A34**, 243–254 (2007).
- [33] M. E. McCracken *et al.* (CLAS), “Differential cross section and recoil polarization measurements for the $\gamma p \rightarrow K^+ \Lambda$ reaction using CLAS at Jefferson Lab,” *Phys. Rev.* **C81**, 025201 (2010).
- [34] B. Dey *et al.* (CLAS), “Differential cross sections and recoil polarizations for the reaction $\gamma p \rightarrow K^+ \Sigma^0$,” *Phys. Rev.* **C82**, 025202 (2010).
- [35] A. V. Anisovich *et al.*, “ N^* resonances from $K\Lambda$ amplitudes in sliced bins in energy,” *Eur. Phys. J.* **A53**, 242 (2017).
- [36] T. Mart and C. Bennhold, “Evidence for a missing nucleon resonance in kaon photoproduction,” *Phys. Rev.* **C61**, 012201 (2000).
- [37] F. X. Lee, T. Mart, C. Bennhold, and L. E. Wright, “Quasifree kaon photoproduction on nuclei,” *Nucl. Phys.* **A695**, 237–272 (2001).
- [38] R. Arndt, W. Briscoe, I. Strakovsky, and R. Workman (George Washington Data Analysis Center), “SAID,” (2015), SAID web site <http://gwddac.phys.gwu.edu>.
- [39] R. A. Adelseck, C. Bennhold, and L. E. Wright, “Kaon Photoproduction Operator for Use in Nuclear Physics,” *Phys. Rev.* **C32**, 1681–1692 (1985).
- [40] I. Strakovsky (George Washington Data Analysis Center), “SAID,” (2017), Private Communication.
- [41] A. V. Anisovich, R. Beck, E. Klempt, V. A. Nikonov, A. V. Sarantsev, and U. Thoma, “Pion- and photo-induced transition amplitudes to ΛK , ΣK , and $N\eta$,” *Eur. Phys. J.* **A48**, 88 (2012).
- [42] B. A. Mecking *et al.*, “The CEBAF Large Acceptance Spectrometer (CLAS),” *Nucl. Instrum. Meth.* **A503**, 513–553 (2003).
- [43] K. Moriya *et al.* (CLAS), “Spin and parity measurement of the $\Lambda(1405)$ baryon,” *Phys. Rev. Lett.* **112**, 082004 (2014).
- [44] K. Moriya *et al.* (CLAS), “Differential Photoproduction Cross Sections of the $\Sigma^0(1385)$, $\Lambda(1405)$, and $\Lambda(1520)$,” *Phys. Rev.* **C88**, 045201 (2013), [Addendum: *Phys. Rev.* **C88**, no.4, 049902 (2013)].
- [45] K. Moriya *et al.* (CLAS), “Measurement of the $\Sigma\pi$ photoproduction line shapes near the $\Lambda(1405)$,” *Phys. Rev.*

- C87**, 035206 (2013).
- [46] D. Ho *et al.* (CLAS), “Beam-Target Helicity Asymmetry for $\vec{\gamma}\vec{n} \rightarrow \pi^- p$ in the N^* Resonance Region,” *Phys. Rev. Lett.* **118**, 242002 (2017).
- [47] J. M. Grames *et al.*, “Unique electron polarimeter analyzing power comparison and precision spin-based energy measurement,” *Phys. Rev. ST Accel. Beams* **7**, 042802 (2004).
- [48] D. I. Sober *et al.*, “The bremsstrahlung tagged photon beam in Hall B at JLab,” *Nucl. Instrum. Meth.* **A440**, 263–284 (2000).
- [49] Haakon Olsen and L. C. Maximon, “Photon and Electron Polarization in High-Energy Bremsstrahlung and Pair Production with Screening,” *Phys. Rev.* **114**, 887–904 (1959).
- [50] M. M. Lowry *et al.*, “A cryostat to hold frozen-spin polarized HD targets in CLAS: HDice-II,” *Nucl. Instrum. Meth.* **A815**, 31–41 (2016).
- [51] C. D. Bass *et al.*, “A portable cryostat for the cold transfer of polarized solid HD targets: HDice-I,” *Nucl. Instrum. Meth.* **A737**, 107–116 (2014).
- [52] H. Drucker and C. Cortes, “Boosting decision trees,” in *Advances in Neural Information Processing Systems*, 8, edited by D. Touretzky, M. Moser, and M. Hasselmo (MIT Press, Cambridge, Massachusetts, 1996) pp. 479–485.
- [53] B. P. Roe, H-J. Yang, J. Zhu, Y. Liu, I. Stancu, and G. McGregor, “Boosted decision trees as an alternative to artificial neural networks for particle identification,” *Nucl. Instrum. Meth.* **A543**, 577 – 584 (2005).
- [54] Y. G. Sharabian *et al.*, “A new highly segmented start counter for the CLAS detector,” *Nucl. Instrum. Meth.* **A556**, 246–258 (2006).
- [55] M. D. Mestayer *et al.*, “The CLAS drift chamber system,” *Nucl. Instrum. Meth.* **A449**, 81–111 (2000).
- [56] E. S. Smith *et al.*, “The time-of-flight system for CLAS,” *Nucl. Instrum. Meth.* **A432**, 265–298 (1999).
- [57] Dao Ho, “Measurement of the E Polarization Observable for $\gamma d \rightarrow \pi^- p(p_s)$, $\gamma d \rightarrow K^0 \Lambda(p_s)$, and $\gamma d \rightarrow \pi^+ \pi^- d(0)$ using CLAS g14 Data at Jefferson Lab,” Ph.D. Thesis, Carnegie Mellon University (2015), available online at http://www.jlab.org/Hall-B/general/clas_thesis.html.
- [58] A. Höcker, P. Speckmayer, J. Stelzer, F. Tegenfeldt, and H. Voss, “TMVA, toolkit for multivariate data analysis with ROOT,” in *Statistical issues for LHC physics. Proceedings, Workshop, PHYSTAT-LHC, Geneva, Switzerland, June 27-29, 2007*, edited by H.B. Prosper, L. Lyons, and A. DeRoeck (2007) pp. 184–187.
- [59] J. B. Cladis, W. N. Hess, and B. J. Moyer, “Nucleon momentum distributions in deuterium and carbon inferred from proton scattering,” *Phys. Rev.* **87**, 425–433 (1952).
- [60] L. Lamia, M. La Cognata, C. Spitaleri, B. Irgaziev, and R. G. Pizzone, “Influence of the d-state component of the deuteron wave function on the application of the Trojan horse method,” *Phys. Rev.* **C85**, 025805 (2012).
- [61] G. Ramachandran, R. S. Keshavamurthy, and M. V. N. Murthy, “Target Asymmetry And Effective Neutron Polarization With Polarized Deuteron Targets,” *Phys. Lett.* **87B**, 252–256 (1979).
- [62] L. Casey, “The Search for Missing Resonances in $\gamma p \rightarrow K^+ \Lambda$ Using Circularly Polarized Photons on a Longitudinally Polarized Frozen Spin Target,” Ph.D. Thesis, Catholic University of America (2011), available online at http://www.jlab.org/Hall-B/general/clas_thesis.html.