

This is the accepted manuscript made available via CHORUS. The article has been published as:

First limit on neutrinoless quadruple β decay of ^{150}Nd
to the 0_{1}^{+} state of ^{150}Gd

M. F. Kidd and W. Tornow

Phys. Rev. C **98**, 055501 — Published 9 November 2018

DOI: [10.1103/PhysRevC.98.055501](https://doi.org/10.1103/PhysRevC.98.055501)

First limit on neutrinoless quadruple beta decay of ^{150}Nd to the 0_1^+ state of ^{150}Gd

M. F. Kidd

Tennessee Technological University, Cookeville, Tennessee 38506 USA

W. Tornow

Department of Physics, Duke University, Durham, North Carolina 27708 USA

Triangle Universities Nuclear Laboratory, Durham, North Carolina 27708 USA

(Dated: October 2, 2018)

Background: Observation of lepton number violation via detection of neutrinoless double beta decay requires that the neutrino be a Majorana particle. If the neutrino is a Dirac particle, a potential lepton-number violating process is neutrinoless quadruple beta decay. Only a few nuclei can undergo neutrinoless quadruple beta decay; one of these nuclei is ^{150}Nd .

Purpose: This study yields the first half-life limit of the neutrinoless quadruple beta decay to the excited 0_1^+ state of ^{150}Gd .

Method: We searched for neutrinoless quadruple beta decay events to excited final states of ^{150}Gd by detecting the de-excitation gamma rays of the daughter nucleus in coincidence. These gamma rays have energies of 569.031 keV and 638.050 keV, and are emitted in coincidence through a $0_1^+ \rightarrow 2_1^+ \rightarrow 0_{g.s.}^+$ transition.

Results: The enriched Nd_2O_3 sample consisted of 40.33 ± 0.02 g ^{150}Nd and was observed for 642.8 days at the Kimballton Underground Research Facility. A half-life limit for the decay to the 0_1^+ state of ^{150}Gd was found to be $T_{1/2} > 1.76 \times 10^{20}$ years (90% CL).

Conclusions: We report the first search for this decay to excited final states. Though the predicted half life of this decay is many orders of magnitudes larger, constraining this value experimentally is vital to check for potential enhancements to the decay rate.

I. INTRODUCTION

One major result that would come from the detection of neutrinoless double-beta ($0\nu\beta\beta$) decay is the observation of lepton number violation (LNV). For $0\nu\beta\beta$ decay to proceed, the neutrino must be its own anti-particle, a Majorana particle. If the neutrino is a Dirac particle, $0\nu\beta\beta$ decay will not proceed. However, as stated by Heeck and Rodejohann [1] the Majorana nature of the neutrino is not required for LNV. One possible allowed decay with Dirac neutrinos that would demonstrate LNV is neutrinoless quadruple beta decay ($0\nu 4\beta$),

$$(A, Z) \rightarrow (A, Z + 4) + 4e^-, \quad (1)$$

where lepton number conservation is violated by 4 units rather than 2 units. Though $4\nu 4\beta$ decay is allowed, the Q -dependent phase space factor (Q^{23}) for eight outgoing particles severely impacts the decay rate, compared to Q^{11} for $0\nu 4\beta$ decay. The daughter ($Z+4$) nucleus must of course have a smaller mass than the parent nucleus, so not all $0\nu\beta\beta$ decay nuclei are good candidates. In fact, only three $0\nu 4\beta$ decay candidates have been identified: ^{96}Zr , ^{136}Xe , and ^{150}Nd . [1]

The Q value of $0\nu 4\beta$ decay of ^{150}Nd to ^{150}Gd is 2.079 MeV, and is the largest of the three candidates. Competing decays include the two-neutrino double-beta ($2\nu\beta\beta$) decay of ^{150}Nd , ($T_{1/2} = 8.2 \times 10^{18}$ years) and $2\nu\beta\beta$ decay to excited states ($T_{1/2} = 1.2 \times 10^{20}$ years) ([2] and references therein). Similarly, there is a possibility of the $0\nu 4\beta$ decay proceeding via a 0_1^+ excited state of ^{150}Gd , which reduces the Q value by 1207.135 keV [3].

Though [1] estimates a half-life to be inordinately large, they emphasize that constraining this value experimentally is extremely valuable to provide inputs for new physics models. To this end, the NEMO-3 collaboration has established a half-life limit for this decay to the ground state of ^{150}Gd [4]. We propose to search for the $0\nu 4\beta$ decay of ^{150}Nd to the 0_1^+ excited state of ^{150}Gd .

The daughter nucleus of the $0\nu 4\beta$ decay of ^{150}Nd is ^{150}Gd , which has a level scheme shown in Figure 1. Beta decay to the 3^- state is highly unlikely due to spin-parity considerations, so we will look for decays to the 0_1^+ state at 1207.135 keV. Decays to this state would result in a $0_1^+ \rightarrow 2_1^+ \rightarrow 0_{g.s.}^+$ decay, emitting 569.031 keV and 638.050 keV gamma rays in coincidence [3]. The rationale for concentrating on the $0\nu 4\beta$ decay to excited states is based on the fact that coincidence experiments, although less efficient, may provide greater sensitivity in searches for extremely rare events than singles measurements due to the reduction of background events.

II. EXPERIMENTAL METHOD

Our experimental setup is optimized to search for coincident gamma rays emitted from a disk-shaped source. Two high-purity germanium (HPGe) detectors surrounded by several layers of active and passive shielding sandwich the source to be studied. These detectors are operated in both “singles” mode and coincidence mode, in order to detect gamma rays emitted in subsequent decays of an excited state of a nucleus which has undergone a specific decay. The coincidence requirement




FIG. 1. Level scheme of ^{150}Nd $0\nu 4\beta$ decay to ^{150}Gd .




FIG. 2. Setup diagram for the double-beta decay apparatus. Please see text for details.

of two gamma rays of specific energies deposited simultaneously results in a significant reduction of random background events in the region of interest. Additionally, the uniqueness of the required coincident energies means there are few or no background candidates. These detectors are p-type coaxial germanium detectors with a diameter of 8.8 cm and thickness of 5.0 cm. They have nickel-plated magnesium endcaps that are 2.54 mm thick, and are connected to their cryostats in the j-type configuration. This configuration is ideal because it allows for the easy insertion of the crystals into the veto annulus. The preamplifier and high-voltage filter are located near the cryostat, away from the crystals, in order to reduce backgrounds.

Surrounding the HPGe detectors and sample is a sodium iodide (NaI) annulus with two plastic-plate scintillators on the endcaps. Three photomultiplier tubes (PMTs) on each end of the annulus and two PMTs on each plastic plate scintillator are operated in anti-coincidence with the HPGe detectors to form an active veto. Next is a layered passive shield composed of 3/4 inch thick Oxygen-Free High Conductivity (OFHC) copper plates followed by 6 inches of lead. For a full discussion of the electronics and data acquisition please see [5]. Finally, this setup was operated underground at the Kimballton Underground Research Facility (KURF) (details on KURF in [6]). A drawing of the apparatus is shown in Figure 2.

This setup was used to determine the half-life of $2\nu\beta\beta$ decay of ^{100}Mo and ^{150}Nd to final excited states ([7], [5]), and to determine a half-life limit of $0\nu\text{E}^{\text{CEC}}$ on ^{112}Sn [8]. More recently, the apparatus was used in searches for the $2\nu\beta\beta$ decay of ^{92}Zr to excited states in ^{96}Mo [9] and for the single beta decay of ^{96}Zr [10].

Here, we use data acquired from October 2008 – August 2010 for determination of $2\nu\beta\beta$ decay of ^{150}Nd to final excited states [5]. The ^{150}Nd sample used in this experiment consists of 50.00 g of Nd_2O_3 powder (43.09 ± 0.02 g Nd) enriched to 93.60% ^{150}Nd (40.33 ± 0.02 g ^{150}Nd). The mass of the ^{150}Nd content is determined via isotopic analysis. This powder is compressed into a cylindrical cavity 0.780 ± 0.005 cm thick and 2.86 cm in radius. This diameter sample makes the most of the coincidence efficiency of the detectors, which is higher near the center of the detectors. Data acquisition took place in 3-5 day-long runs over a period of 642.8 days.

III. COINCIDENCE EFFICIENCY

The coincidence efficiency for this setup was determined in [7] using the decay of ^{102}Ru , which emits two gamma rays in coincidence in a $0^+ \rightarrow 2^+ \rightarrow 0^+$ decay sequence. The energies of the gamma rays in this decay are 468.64 keV and 475.10 keV. To adapt this efficiency measurement to our purposes, corrections need to be made to account for attenuation differences and the energy dependence of the efficiency. Full details on the corrections made to the efficiency for $2\nu\beta\beta$ decay of ^{150}Nd are given in [5]. Here, the corrections were recalculated for the energies of the $0\nu 4\beta$ decay of ^{150}Nd . The final results yields a total coincidence efficiency for the $0\nu 4\beta$ decay of ^{150}Nd of $\epsilon_{\gamma\gamma} = (0.634 \pm 0.043)\%$.

IV. BACKGROUND CANDIDATES

An extremely thorough discussion of potential background sources can be found in [5]. Here, we discuss the only option that contributes in the coincidence region of interest (ROI). As discussed in [5], there is a known contamination of ^{232}Th decay products in the enriched ^{150}Nd sample. A search of the NNDC database [3] for coincident gamma rays in the ROI reveals only one potential source that could reasonably be present in our setup. This transition is located in the β decay of ^{228}Ac to ^{228}Th . In this transition, gamma rays of energy 570.91 keV (intensity, $I_\gamma = 0.182\%$), 572.14 keV ($I_\gamma = 0.150\%$), and 640.32 keV ($I_\gamma = 0.054\%$) can be emitted. Indeed, in the single-detector spectrum, a peak can be observed at 569 keV as shown in Figure 3, though no peak is obvious at 640 keV. To determine if background from this source is a concern, the level diagram of ^{228}Th must be inspected. The 640.32 keV gamma ray originates from the 1901.92 keV energy level ($J^\pi = 6^+$) with a relative intensity of 48%. This transition proceeds through the 1261.57 keV (4^+) level, and 36% of these transitions continue through the 1091.048 keV (4^+) level. From here, 2.3 % of these decays result in a 571.8 keV gamma ray. Therefore, it is unlikely there would be significant contamination from this decay in the coincidence spectrum. To check this, we analyzed 1.10 days of a ThO_2 sample inserted in the apparatus. As shown in Figure 4, no events above background are shown in the coincidence ROI.

However, as mentioned above, there is a sizable peak around 569 keV. Investigation of the coincidence spectrum reveals that this peak occurs in coincidence with 1063 keV. This level transition occurs in the $^{207}\text{Pb}(n,n'\gamma)$ interaction when the excited ^{207}Pb nucleus decays to its ground state via the 1633.36 keV state, emitting 1063.66 keV and 569.70 keV gamma rays in coincidence. However, this interaction requires a neutron flux, and is only apparent when the ^{150}Nd sample is present. We believe (α,n) reactions are taking place in the carbon in the teflon holder surrounding the ^{150}Nd sample. The α decays are




FIG. 3. The potential background peak at 569.70 keV is shown in a single-detector spectrum. Also note the lack of any excess of counts at 640 keV. See text for details of the origin of the background radiation.

due to the ^{232}Th contamination in the ^{150}Nd sample itself. Though this peak is in our single-gamma ROI, no coincidence candidate exists.

V. ANALYSIS

The coincidence data is investigated by plotting events in detector 2 versus the events in detector 1. Only the events which meet the coincidence timing requirement (about 4 μs) between the two detectors and the anti-coincidence timing requirement (about 10 μs) between the detectors and the veto are plotted in this spectrum. In the two-dimensional spectrum, the events of interest would occur at coordinates of (569.03 keV, 638.05 keV) and (638.05 keV, 569.03 keV). The significance of the counts in the two-dimensional regions of interest can best be seen by projecting them onto the x- or y-axis. To accomplish this, an energy condition (*i.e.* 569.03 ± 2.50 keV) is applied to events in one detector, and all the events which occur in the other detector in coincidence are projected into a histogram. This results in four histograms; two are in coincidence with 569.03 keV in detectors 1 and 2, and two are in coincidence with 638.05 keV in detectors 1 and 2. The corresponding histograms can then be summed so that all events in coincidence with a 569.03 keV photon are in one histogram, and vice versa, reducing the number of histograms to two (see Fig. 5). To avoid any histogram summing effects, the detectors were carefully gain matched and the bin widths are extremely close.

VI. QUADRUPLE-BETA DECAY OF ^{150}Nd TO 0_1^+ EXCITED STATE OF ^{150}Gd

Over 642.8 days of data acquisition, seven events in the 569-638 keV coincidence window are observed with an average background of 5.48 ± 0.57 counts (see Figure




FIG. 4. (a) The two-dimensional spectrum shows the ThO₂ data where no events are in coincidence above background in our region of interest (indicated by the dashed lines at 569.05 keV – 638.05 keV). (b) The singles spectrum for the ThO₂ data shows no peak at 569 keV. See text for details.

5), therefore, a half-life limit can be set. The half-life limit is given by

$$T_{1/2} > \frac{\ln(2)N_0 t f_b \epsilon_{\gamma\gamma}^{tot}}{N_d}, \quad (2)$$

Here, N_0 is the number of ^{150}Nd nuclei in the sample, t is total counting time (642.8 days or 1.76 years), $\epsilon_{\gamma\gamma}^{tot}$ is the total coincidence detection efficiency for the 569.08-638.05 keV cascade, and f_b is the branching ratio for the particular cascade. The value of N_0 is $1.620 \pm 0.003 \times 10^{23}$ nuclei of ^{150}Nd . The coincidence efficiency for the 569.08-638.05 keV cascade is $\epsilon_{\gamma\gamma} = (0.634 \pm 0.043)\%$, and f_b is 100% for this cascade. N_d is a value chosen using the TRolke class in the ROOT analysis framework [11]. This class considers a Poisson signal in the presence of an uncertain background and efficiency, and computes the confidence interval for this situation, using the method outlined in [12]. For the cal-


culation described here, we used a background estimate simultaneously measured from the sidebands of our signal, and assumed a Poisson uncertainty in this estimate, with a Gaussian uncertainty in the efficiency measurement. Here, with an observed signal of 7 events with a background of 5.48 ± 0.57 events in the region of interest, the Rolke method gives an upper limit to the profile likelihood of $N_d/\epsilon_{\gamma\gamma}^{tot} = 1123$ events, resulting in a half-life limit of $T_{1/2} > 1.76 \times 10^{20}$ years (90% CL). This limit was obtained with only 40.33 g of ^{150}Nd ; significant improvements to the limit could be achieved with a larger sample.

VII. CONCLUSIONS

We have presented the first search for $0\nu 4\beta$ decay of ^{150}Nd to excited final states of ^{150}Gd . Though this decay is predicted to have a half-life many orders of magnitude larger than this limit, it is important to experimentally constrain these quantities to check for potential enhancements to the decay rate, as discussed in [1].

VIII. ACKNOWLEDGEMENTS

This work was partially supported by the U.S. Department of Energy, Office of Nuclear Physics, under grant No. DE-FG02-97ER41033. The authors would like to thank Dr. Graham Giovanetti for his advice regarding the use of the TRolke class.



inin

FIG. 5. Coincidence data for decays to the 0_1^+ state in ^{150}Gd . Spectrum (a) is in coincidence with 569.03 keV, and spectrum (b) is in coincidence with 638.05 keV. The coincidence-energy window in spectra (a) and (b) is 2.50 keV for detector 2 and 3.00 keV for detector 1. Events within the region of interest are colored red.

-
- [1] J. Heeck and W. Rodejohann, EPL **103**, 32001 (2013).
 - [2] A. S. Barabash, Nucl. Phys. A **935**, 52 (2015).
 - [3] A. Sonzogni, Int. Conf. on Nucl. Data for Sci. and Tech. **ND 2007**, 029 (2007).
 - [4] R. Arnold. et al. (NEMO-3 Collaboration), Phys. Rev. Lett. **119**, 041801 (2017).
 - [5] M. F. Kidd, J. H. Esterline, S. W. Finch, and W. Tornow, Phys. Rev. C **90**, 055501 (2014).
 - [6] P. Finnerty, S. MacMullin, H. Back, R. Henning, A. Long, K. Macon, J. Strain, R. Lindstrom, and R. Vogelaar, Nucl. Inst. and Meth. A **642**, 65 (2011).
 - [7] M. F. Kidd, J. H. Esterline, W. Tornow, A. S. Barabash, and V. Umatov, Nucl. Phys. A **821**, 251 (2009).
 - [8] M. F. Kidd, J. H. Esterline, and W. Tornow, Phys. Rev. C **78**, 035504 (2008).
 - [9] S. W. Finch and W. Tornow, Phys. Rev. C **92**, 045501 (2015).
 - [10] S. W. Finch and W. Tornow, Nucl. Inst. and Meth. A **806**, 70 (2016).
 - [11] R. Brun and F. Rademakers, Nucl. Inst. and Meth. in Phys. Res. A **389**, 81 (1997).
 - [12] W. A. Rolke, A. M. Lopez, and J. Conrad, Nucl. Inst. and Meth. A **551**, 493 (2005).