



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Search for high-energy neutrino emission from radio-bright AGN

Bei Zhou, Marc Kamionkowski, and Yun-feng Liang
Phys. Rev. D **103**, 123018 — Published 16 June 2021
DOI: [10.1103/PhysRevD.103.123018](https://doi.org/10.1103/PhysRevD.103.123018)

Search for High-Energy Neutrino Emission from Radio-Bright AGN

Bei Zhou,^{1,*} Marc Kamionkowski,^{1,†} and Yun-feng Liang^{2,‡}

¹*Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA*

²*Laboratory for Relativistic Astrophysics, Department of Physics, Guangxi University, Nanning 530004, China*

(Dated: May 05, 2021)

We investigate the possibility that radio-bright active galactic nuclei (AGN) are responsible for the TeV–PeV neutrinos detected by IceCube. We use an unbinned maximum-likelihood-ratio method, 10 years of IceCube muon-track data, and 3388 radio-bright AGN selected from the Radio Fundamental Catalog. None of the AGN in the catalog have a large global significance. The two most significant sources have global significance of $\simeq 1.5\sigma$ and 0.8σ , though 4.1σ and 3.8σ local significance. Our stacking analyses show no significant correlation between the whole catalog and IceCube neutrinos. We infer from the null search that this catalog can account for at most 30% (95% CL) of the diffuse astrophysical neutrino flux measured by IceCube. Moreover, our results disagree with recent work that claimed a 4.1σ detection of neutrinos from the sources in this catalog, and we discuss the reasons of the difference.

I. INTRODUCTION

The TeV–PeV diffuse astrophysical neutrinos detected by IceCube [1–6] have opened a new window for astrophysics [7–21] and provide a new high-energy-particle source for elementary particle physics [22–45]. However, the origin of these neutrinos remains a mystery, albeit one that is thought to be related to the origin of high/ultrahigh-energy cosmic rays [46–51], which still remains obscure. In the past decade, significant efforts have been made to search for the sources of these astrophysical neutrinos [52–59]. A few sources were found to be very likely TeV–PeV neutrino emitters, including TXS 0506+056 [60, 61] and NGC 1068 [58]. The blazar TXS 0506+056 was found to have a 3σ (global significance) spatial and temporal association with a high-energy muon-track event induced by a ~ 300 TeV neutrino [61]. This led to the discovery of a neutrino flare between 2014 and 2015 from the location of TXS 0506+056, showing 3.5σ global significance [60]. Note that these two measurements are statistically independent. The Seyfert II galaxy NGC 1068 showed an excess of 2.9σ (global significance) in the time-integrated source searches using 10 years of IceCube data [58].

Still, the majority of the diffuse astrophysical neutrinos remain largely unexplained. Many different types of sources have been proposed and searched with data, including gamma-ray bursts [62–68], gamma-ray blazars [69–73], choked jet supernovae [74, 75], pulsar wind nebulae [76], etc. However, none of these searches found a strong correlation. For example, motivated by theoretical considerations [77–79] and TXS 0506+056 [60, 61], searches for high-energy neutrino emission from gamma-ray blazars were implemented [69, 70, 73]. However, they were found to contribute at most $\sim 20\%$ to the diffuse neutrino flux. (See also Ref. [71]

for a general discussion.) On the other hand, it has been shown that there exist high-energy neutrino sources hidden in GeV–TeV gamma rays [80–82]. Otherwise, the gamma rays from all the transparent sources would overshoot the isotropic diffuse gamma-ray flux measured by *Fermi*-LAT [83]. Therefore, other categories of sources must be investigated, including those not characterized by gamma rays.

Recently, the possibility that radio-bright AGN with strong parsec-scale cores are associated with IceCube neutrino events has been considered [84, 85]. The 3388 radio-bright AGN that were used for the analysis form a complete catalog and were selected from the Radio Fundamental Catalog [86], which collects VLBI (very-long-baseline interferometry) observations since 1980. These radio-bright AGN have very diverse gamma-ray properties. In Ref. [84], a 3.1σ significance was found when 56 muon-track events (> 200 TeV) from 2009 to 2019 were considered. In Ref. [85], a pre-trial p-value sky map published by the IceCube Collaboration [87] was found to have a 3σ correlation with radio-bright AGN. The sky map was obtained from the analysis of muon-track data between 2008 and 2015. The combination of the two analyses yields a 4.1σ -significance detection [85]. Moreover, the sources with stronger X-band ($\simeq 8$ GHz) flux densities show more significant correlations, implying that the X-band flux could be an indicator of TeV–PeV neutrino emission.

In this paper, we revisit the correlation between these radio-bright AGN and the TeV–PeV astrophysical neutrinos detected by IceCube. We use 10 years of IceCube muon-track data (April 2008 to July 2018) [88, 89], made publicly available just recently, and an unbinned maximum-likelihood-ratio method [52, 53, 90, 91]. The method, which has been extensively used for analyzing IceCube data [52, 53, 69, 70, 73, 76], takes into account the information provided by every single event. It is also widely used in other types of astroparticle measurements (e.g., Refs. [92–95]). The conclusion of our analysis is that there is no significant correlation between IceCube neutrinos and these 3388 radio-bright AGN. Our null

* beizhou@jhu.edu; orcid.org/0000-0003-1600-8835

† kamion@jhu.edu; orcid.org/0000-0001-7018-2055

‡ liangyf@gxu.edu.cn; orcid.org/0000-0002-6316-1616




FIG. 1. Distributions of the 3388 radio-bright AGN (*red*) and 1134450 neutrino events (*blue*) used in this work on the celestial sphere. The sky map is in the equatorial coordinate system, with the *horizontal* and *vertical* axes representing right ascension (RA) and declination (Dec), respectively. To IceCube, the upper half (Dec $> 0^\circ$) is the northern sky (*seen through Earth*), while the lower half (Dec $< 0^\circ$) is the southern sky. The neutrino events are shown as numbers in 360×180 of RA \times sin(Dec) bins, so that all the bins have the same solid angle. See Sec. II for details.

search suggests that no more than 30% of the IceCube neutrino flux may have come from these sources.

This paper is organized as follows. In Sec. II, we describe the radio-bright AGN (Sec. II A) and the 10 years of IceCube muon-track data (Sec. II B). Sec. III discusses the unbinned maximum-likelihood-ratio method used for our analysis. Sec. IV shows the results for both the analysis that searches for neutrino emission from every source location in the catalog (Sec. IV A) and the analysis stacking all the sources together for the correlation with all-sky neutrinos (Sec. IV B). We also compare our results with previous work (Sec. IV C). We conclude in Sec. V.

II. DATA

A. Radio-bright AGN catalog from VLBI observations

The radio-bright AGN used in this work are selected from the Radio Fundamental Catalog [86] in the Astrogéo database [96]. The Radio Fundamental Catalog collects archival VLBI flux densities at several frequencies between 2 and 22 GHz as well as precise positions with milliarcsecond accuracies for more than 10^5 compact radio sources. It uses all available VLBI (very-long-baseline interferometry) observations under absolute astrometry and geodesy programs since April 11, 1980. The catalog is regularly updated with more sources and more observations per source. In this work, we use the version

“rfc_2019c”, which comprises 16466 objects.

Moreover, as in Ref. [84], we obtain a complete catalog of radio-bright AGN by selecting the sources in the Radio Fundamental Catalog *with X-band ($\simeq 8$ GHz) flux densities larger than 0.15 Jy*. This gives us 3388 sources, with the total X-band flux density $\simeq 1518$ Jy. This is the only deep statistically complete catalog that can be obtained from the Radio Fundamental Catalog (see Ref. [84] for details). Moreover, these radio-bright AGN have very diverse gamma-ray properties. Fig. 1 shows the distribution of the 3388 sources on the celestial sphere. Their locations are nearly isotropic.

B. Neutrino data from IceCube

The IceCube Neutrino Observatory detects neutrinos by detecting the Cherenkov photons emitted by relativistic secondary charged particles from neutrino interactions in and outside the detector [97, 98]. Two basic kinds of event topologies are formed in the detector. One is an elongated track formed by muons due to their low energy-loss rate in matter. The other is a shower which looks like a round and big blob formed by electrons (electromagnetic shower) or hadrons (hadronic shower). The track events have a better angular resolution (as good as $< 1^\circ$) while worse energy resolution ($\sim 200\%$ at ~ 100 TeV) than the shower events ($\sim 10^\circ$ – 15° and $\sim 15\%$ above 100 TeV) [99]. Therefore, the track events are suited to search for point sources.

There are two kinds of track events, through-going and starting tracks. The through-going tracks are those with parent neutrinos that have interacted outside the detector, while inside for starting tracks. Overall, the through-going tracks have better angular resolution and ~ 10 times larger effective area than that of starting tracks [56].

The data released by the IceCube Collaboration span from April 2008 to July 2018 [88, 89]. The same data have been used in IceCube's 10-year time-integrated neutrino point-source search [58]. In total, there are 1134450 muon-track events. The information for each track is provided. These ten years of data can be grouped into five samples corresponding to different construction phases of IceCube, including 1) IC40, 2) IC59, 3) IC79, 4) IC86-I, and 5) IC86-II to IC86-VII. The numbers in the names represent the numbers of strings in the detector on which Digital Optical Modules are deployed.

There are two main features in the new dataset that are different from the previously released three years of data from June 2010 to April 2012 [88, 89]. First, the reconstructed directional uncertainty assumes a lower limit of 0.2° in order to avoid unaccounted-for uncertainties and to ensure no single event dominates the likelihood analysis. Second is the given uptime, which contains lists of time periods of good runs for each season.

Fig. 1 shows the distribution of the muon-track events on the celestial sphere in the equatorial coordinate system. The distribution is shown as numbers in 360×180 of RA \times sin(Dec) bins, so that all the bins have the same solid angle. More events came from the northern sky (upper half; Dec $> 0^\circ$) than the southern sky (lower half; Dec $< 0^\circ$), because the latter has a higher energy threshold to the events to reduce the contamination from atmospheric muons [88]. For the southern sky, the events are nearly evenly distributed. For the northern sky, there are more events as it is closer to Dec = 0° because of the absorption of neutrinos by Earth. For more details, see Refs. [88, 89].

III. ANALYSIS FORMALISM

Our analysis uses an unbinned maximum-likelihood-ratio method [52, 53, 90, 91]. The likelihood function is given by the product of probability density functions (PDFs) of each muon-track event (indexed by i) in the five data samples (indexed by k),

$$\mathcal{L}(n_s) = \prod_k \prod_{i \in k} \left[\frac{n_s^k}{N_k} S_i^k + \left(1 - \frac{n_s^k}{N_k}\right) B_i^k \right], \quad (1)$$

where n_s is the total number of events from the source(s) combining all five samples, which is to be fit in our analysis, and N_k is the total number of events in sample k . The S_i^k and B_i^k are the signal and background PDFs of event i in the sample k , representing the probability of the event coming from the source(s) or background, respectively.




FIG. 2. RA-integrated background PDF for different phases of IceCube, as labeled. Shown are IC40 ($\simeq 1$ -year exposure, 36900 events), IC59 ($\simeq 1$ -year exposure, 107011 events), IC79 ($\simeq 1$ -year exposure, 93133 events), IC86-I ($\simeq 1$ -year exposure, 136244 events), and IC86-II to IC86-VII ($\simeq 6$ -year exposure, 761162 events). The left half is the northern sky while right half the southern sky with respect to IceCube.

Note the k in the superscript is not an exponent but an index. Moreover, n_s^k is the number of expected signal events coming from sample k , which is determined by

$$n_s^k = f_k \times n_s, \quad (2)$$

where f_k is the expected fractional contribution from sample k , given by Eq. (6) and (7).

We calculate the background PDFs directly from data. For each sample, given the Dec δ_i of an event, the background PDF is determined by the relative number of events in $\delta_i \pm 3^\circ$ divided by the solid angle. We do not use events with $|\delta_i| > 87^\circ$ (only account for $\simeq 0.1\%$ of the sky and the total events). The calculated background PDFs have precision of less than 2% for most directions. The dependence on RA is negligible because IceCube, located at the South Pole, has a nearly uniform acceptance in RA.

Fig. 2 shows the RA-integrated background PDFs. They are consistent with the distribution of muon-track events in Fig. 1 (see also Sec. IIB).

The signal PDF is expressed as a 2D symmetric Gaussian spatial distribution,

$$S_{ij}^k \equiv S_k(\vec{x}_i, \sigma_i, \vec{x}_j) = \frac{1}{2\pi\sigma_i^2} \exp\left(-\frac{D(\vec{x}_i, \vec{x}_j)^2}{2\sigma_i^2}\right) \quad (3)$$

where \vec{x}_i and \vec{x}_j are the coordinates of the event i and source j on the celestial sphere, respectively, and $D(\vec{x}_i, \vec{x}_j)$ is the angular distance between them. Here,

σ_i is the uncertainty of the reconstructed direction of the track (much larger than the scattering angle between the incoming neutrino and the outgoing muon [54]). We do not include the energy part of the signal PDF because of lower statistics in the published smearing matrices [88, 89]. However, our analysis is sensitive to different neutrino spectrum through the energy- and zenith-dependent effective area [88, 89].

For the case of multiple sources, the signal PDF is a weighted sum of the signal PDFs (S_{ij}^k) of all the sources, indexed by j [52, 53],

$$S_i^k = \frac{\sum_j w_{j,\text{model}} w_{j,\text{acc}}^k S_{ij}^k}{\sum_j w_{j,\text{model}} w_{j,\text{acc}}^k}, \quad (4)$$

where the w_j are weighting factors which are proportional to the number of expected events from source j . The $w_{j,\text{acc}}^k$ reflects the detector's acceptance, which is proportional to the convolution of detector effective area (A_{eff}^k) and neutrino spectrum,

$$w_{j,\text{acc}}^k(\delta_j) \propto t_k \times \int A_{\text{eff}}^k(E_\nu, \delta_j) E_\nu^\Gamma dE_\nu, \quad (5)$$

where δ_j is the Dec of the source, t_k the total detector uptime of the sample k , E_ν neutrino energy, and Γ is the spectral index of the signal neutrinos, which is assumed to be the same for all the sources.

The $w_{j,\text{model}}$ describes the expected relative number of events due to the intrinsic properties of the sources, as discussed in Sec. IV B 1.

The f_k in Eq. (2) can therefore be calculated by

$$f_k = \frac{w_{j,\text{acc}}^k}{\sum_k w_{j,\text{acc}}^k}, \quad (6)$$

for one source, or

$$f_k = \frac{\sum_j w_{j,\text{model}} w_{j,\text{acc}}^k}{\sum_k \sum_j w_{j,\text{model}} w_{j,\text{acc}}^k}, \quad (7)$$

for multiple sources.

The test statistic (TS) of our analysis is,

$$\text{TS}(n_s) = 2 \ln \frac{\mathcal{L}(n_s)}{\mathcal{L}(n_s = 0)}, \quad (8)$$

where the denominator is the background/null hypothesis that all the events come from background. The best-fit number \hat{n}_s of signal events is obtained by maximizing the TS value (TS_{max}).

If the background hypothesis is true, the probability distribution for TS_{max} is approximately a χ^2 distribution, i.e.,

$$\text{PDF}(\text{TS}_{\text{max}}) \simeq \chi_1^2(\text{TS}_{\text{max}}), \quad (9)$$

when the number of observations (N_k for our case) is large (Wilks' theorem [100]). The subscript "1" is the




FIG. 3. Distribution of maximized TS values for the radio-bright AGN with $\hat{n}_s \geq 0$ from our likelihood analysis with 10 years of IceCube muon-track data. The error bar is the square root of the number in each bin. The red dashed line is the best normal distribution fit, which is proportional to the standard normal distribution.

degree of freedom of the χ^2 distribution, determined by the difference in the number of free parameters in the signal hypothesis and background hypothesis. Therefore, the $\sqrt{\text{TS}_{\text{max}}}$ is a good approximation of the significance for rejecting the background hypothesis, as it follows the standard normal distribution. We also verify in Sec. IV the use of Wilks' theorem in our analysis by comparing with simulations.

IV. RESULTS AND DISCUSSIONS

We now present our analysis with the catalog of radio-bright AGN, using the formalism in Sec. III and 10 years of IceCube data. We begin in Sec. IV A by analyzing the significance of each source location in the catalog. Then, in Sec. IV B we study the correlation of all-sky diffuse astrophysical neutrinos with the whole catalog by stacking the sources together in the likelihood analysis [Eq. (4)]. We have checked our analysis code with relevant analyses in previous work [58, 60, 61, 69, 70, 73], and we get consistent results. Finally in Sec. IV C, we compare our result with previous work [84, 85].

A. Single-source analysis

Fig. 3 shows the $\sqrt{\text{TS}_{\text{max}}}$ for radio-bright AGN in the catalog. We plot it as the number of sources in each

TABLE I. List of the five sources with highest significance.

IVS name	J2000 name	X-band flux density (Jy)	\hat{n}_s	TS _{max}	Pre-trial p-value, significance	Post-trial p-value, significance
1303-170	J1306-1718	0.208	21.6	16.6	2.28e-5, 4.1 σ	0.074, 1.5 σ
2245+029	J2247+0310	0.434	50.8	14.5	7.14e-5, 3.8 σ	0.21, 0.8 σ
0228-163	J0231-1606	0.162	15.9	9.8	8.90e-4, 3.1 σ	0.95, 0
1424+240	J1427+2348	0.187	38.1	8.9	1.42e-3, 3.0 σ	0.99, 0
0958+559	J1001+5540	0.180	27.2	8.3	2.02e-3, 2.9 σ	1.0, 0

$\sqrt{\text{TS}_{\text{max}}}$ bin, with the error bar given by the square root of the number. ~~The numbers with error bars can be very well fit by the standard normal distribution times a constant, with the reduced chi-square found to be 0.69.~~ We use a normal distribution to fit the numbers with error bars. The best fit is consistent with the standard normal distribution times the number of sources divided by the number of bins per unit x-axis and has a reduced chi-square found to be 0.69. Therefore, the probability density distribution of $\sqrt{\text{TS}_{\text{max}}}$ of the sources in this catalog follows the standard normal distribution. According to Wilks' theorem [100] (Sec. III), this means that at the positions of most sources the background hypotheses are **strongly** favored, implying no **very** strong correlation between diffuse astrophysical neutrinos and this catalog. **This is further and quantitatively supported by our stacking analysis in Sec. IV B.**

Table I lists the five sources with the highest TS_{max}, from which we calculate the pre-trial (local) p-value, p_{local} , and the corresponding significance in units of standard normal deviations. The post-trial (global) p-value can be calculated by $p_{\text{global}} = 1 - (1 - p_{\text{local}})^{N_{\text{src}}}$, from which we get the post-trial significance. Here the N_{src} is the number of sources. The source with the highest post-trial significance is J1306-1718, which has $\simeq 1.5\sigma$ ($p_{\text{global}} = 0.074$). All the other sources have post-trial significance less than 1. The X-band flux densities of the sources with highest local significance are not high compared to other sources in the catalog, implying that the X-band flux densities of radio-bright AGN might not be an indicator of high-energy emission. Moreover, we do not find anything special about these sources with high local significance compared to other sources in the catalog.

To check the pre-trial p-value (and significance) calculated from Wilks' theorem [100], we simulate $N_{\text{src}}^{\text{simu}}$ ($> 10^6$) random source locations on the celestial sphere following uniform distributions in RA and $\sin(\text{Dec})$ [= $-\cos(\text{zenith angle})$ for IceCube location]. For each location, we maximize the TS value. The p-value of a real source in the catalog is then determined by the number of the simulated locations with TS_{max} larger than that of the real source ($M_{\text{src}}^{\text{simu}}$) divided by $N_{\text{src}}^{\text{simu}}$, i.e., $M_{\text{src}}^{\text{simu}}/N_{\text{src}}^{\text{simu}}$. We find that the p-values calculated from Wilks' theorem and simulation are consistent with each other within uncertainties ($\simeq 1/\sqrt{M_{\text{src}}^{\text{simu}}}$).

B. Stacking analysis

The previous subsection shows that none of the sources in the catalog show significant neutrino emission, and the distribution of their TS_{max} strongly favors the background hypothesis. In this subsection, we do a stacking analysis which integrates possible signals from all the sources. Then, we derive their **muon** neutrino ($\nu_{\mu} + \bar{\nu}_{\mu}$) flux by

$$\hat{n}_s = 2\pi \sum_k t_k \int d\sin\delta \int A_{\text{eff}}^k(E_{\nu}, \delta) \frac{dF}{dE_{\nu}}(E_{\nu}) dE_{\nu}, \quad (10)$$

where $dF(E_{\nu})/dE_{\nu} = \Phi_0 \times (E_{\nu}/100 \text{ TeV})^{\Gamma}$ and Φ_0 is the flux normalization at 100 TeV. We also compare their flux with the diffuse astrophysical neutrino flux measured by IceCube [2, 5].

1. Weighting schemes, $w_{j,\text{model}}$

We use two weighting schemes [Eq. (4)]. For the first, we assume that the high-energy neutrino fluxes of the sources are independent of their intrinsic properties. This leads to

$$w_{j,\text{model}} = 1. \quad (11)$$

For the second scheme, we assume that the high-energy neutrino fluxes are proportional to the X-band (radio) flux densities, i.e.,

$$w_{j,\text{model}} = [\text{X-band flux density}]. \quad (12)$$

This is motivated by the findings in Refs. [84, 85] that sources with higher X-band flux densities contribute more to the correlation significance and that the significance increases when a subsample with a higher threshold of X-band flux density is used. Therefore, in this scheme, sources brighter in X-band have higher weights in their contribution to the signal PDF [Eq. (4)].

2. Results of the stacking analysis

Fig. 4 shows the results of our stacking analysis. *For both panels*, our results of all six-flavor neutrino flux combining all the sources in the catalog are shown in red




FIG. 4. Our results of all six-flavor neutrino fluxes combining all the sources in the catalog for the two weighting schemes (red and blue lines), each with three spectra indices as labeled in each panel. Also shown is all six-flavor diffuse astrophysical neutrino flux measured by IceCube [2, 5] with uncertainties (green band; the three green lines represent lower limits, center value, and upper limits). Both our results and IceCube measurements assume equal flux for all six flavors. Both panels use the same line styles. *Left*: TS values as a function of the energy fluxes at 100 TeV. The black dashed horizontal line shows where we set the upper limits at 95% confidence level on each case, corresponding to $TS \simeq -3.84$. *Right*: the energy fluxes versus neutrino energy. Note in the right panel the red and blue lines are *upper limits* at 95% confidence level.

(for equal weighting) and blue lines (for X-band flux-density weighting), each with three different spectral indices. The choice of the spectral indices is explained below. Also shown for comparison is the all six-flavor astrophysical neutrino flux measured by IceCube with uncertainty [2, 5]. Both our results and IceCube measurements assume equal flux for all six flavors.

The left panel shows the TS value as a function of the energy fluxes at 100 TeV, which is about the median energy in the IceCube measurement range (see the right panel). If other energies are chosen, the curves shift only horizontally. As shown by the plots, none of the cases show significant correlation between the catalog and IceCube neutrinos. In fact, all six cases peak at $\hat{n}_s = 0$ with $TS_{\max} = 0$.

Therefore, we set an upper limit at the 95% confidence level for each case, relative to the background hypothesis ($n_s = 0$). According to Wilks' theorem [100], this corresponds to $TS \simeq -3.84$. At 100 TeV, this catalog accounts for at most 27.8% ($\Gamma = -2.00$), 10.3% ($\Gamma = -2.53$), and 1.7% ($\Gamma = -3.0$) for equal weighting, and at most 12.6% ($\Gamma = -2.0$), 4.8% ($\Gamma = -2.53$), and 0.84% ($\Gamma = -3.0$) for an X-band flux-density weighting.

The right panel of Fig. 4 shows the energy fluxes versus neutrino energy (E_ν). Here the red and blue lines are upper limits at the 95% confidence level. The hardest spectrum that we choose, $\Gamma = -2.0$, is motivated by the Fermi acceleration mechanism as well as IceCube's measurements using 10 years of muon-track events between a few TeV and 10 PeV, which gives $-2.28^{+0.08}_{-0.09}$ [3]. The

medium spectrum, $\Gamma = -2.53$, is chosen to match IceCube's measurements using shower events from 16 TeV to 2.6 PeV [5] (-2.53 ± 0.07) as well as earlier measurements using both track and shower events in a similar energy range (-2.50 ± 0.09) [2]. The softest spectrum, $\Gamma = -3.0$, is motivated by IceCube's measurements using high-energy starting events (HESE; > 60 TeV deposit energy), which gives $-2.89^{+0.20}_{-0.19}$ [6].

For the energy flux ($E_\nu dF/dE_\nu$) integrated from 16 TeV to 2.6 PeV, at the 95% confidence level, this catalog accounts for at most 30.6% ($\Gamma = -2.00$), 10.3% ($\Gamma = -2.53$), and 2.3% ($\Gamma = -3.0$) for equal weighting, and at most 13.8% ($\Gamma = -2.0$), 4.8% ($\Gamma = -2.53$), and 1.1% ($\Gamma = -3.0$) for X-band flux-density weighting.

Overall, the X-band flux-density weighting gives less correlation significance and neutrino flux, so the X-band flux densities of radio-bright AGN may not be an indicator of high-energy emission.

In the stacking analysis above we choose two well-motivated weighting schemes. We believe other possibly well-motivated schemes would not lead to a significant correlation, because the distribution of $\sqrt{TS_{\max}}$ of all the sources closely follows the standard normal distribution and none of the sources show a strong global significance (Sec. IV A).

In Appendix A, we estimate the significance if there were signal neutrinos from these sources. We do this by injecting signal neutrinos to the current dataset (Sec. II B) and performing the same stacking analysis. Our results show that even a small contribution to the

total diffuse astrophysical neutrino flux from the sources would have shown significance in our analysis.

C. Comparison with previous work

The correlation between IceCube neutrinos and the same catalog of radio-bright AGN was previously studied in Refs. [84, 85], the first of which found a 3.1σ correlation with 56 muon-track events (> 200 TeV; 2009–2019). Our analysis differs from theirs in the following ways: We include 1) the probability [Eq. (1)], which is not negligible, that the events come from the background, 1) the background component in a more rigorous way [Eq. (1)], using the background PDF directly calculated from data, and 2) a 2D Gaussian term [Eq. (3)], that weights the events differently in terms of their distance from the sources, instead of using the same weights within certain circular regions around the source locations. Moreover, Ref. [85] found a 3σ correlation with a pre-trial p-value sky map calculated from muon-track data from 2008–2015 [87] and 4.1σ after combining the above two analyses. We, on the other hand include the information, including direction and angular error, available for each neutrino event.

Ref. [84] found that four sources—1253-055 (3C 279), 2145+067, 1741-038, and 1730-130 (NRAO 530)—that were among the brightest in the X-band fluxes drove the significance of their signal. The p-value increased from $\simeq 0.1\%$ to only $\simeq 10\%$ if the four sources were removed. Our results, however, do not show any significance from the four sources. In particular, for 3C 279, our analysis yields a local p-value of only 0.7, consistent with 0.63 obtained by IceCube collaboration also using the ten years of data [58]. We also note that three of the four sources above, including 3C 279, were within the enlarged circles (by 0.5°) claimed to take into account IceCube’s systematic uncertainties. Neither our analysis nor those done by IceCube (e.g., Refs [52–61, 63–69, 76]) use enlarged angular errors.

In the stacking analysis that correlates all the sources with all-sky muon-track events (Sec. IV B), our results do not show any significance either. Moreover, our analysis shows that this catalog of radio-bright AGN could account for *at most* 30% (95% confidence level) of the total diffuse astrophysical neutrino flux, in comparison to an estimated $\sim 25\%$ contribution found in Ref. [84].

V. CONCLUSIONS

High-energy astrophysical neutrinos provide a crucial window to study our universe [7–21] and fundamental physics [22–45]. Fully opening the window necessitates the knowledge of their origins, for which we know little at present. Encouraging results including TXS 0506+056 [60, 61] and NGC 1068 [58] have been obtained after dedicated searches by the IceCube Collaboration [52–54, 56, 58]. However, the majority of the dif-

fuse astrophysical neutrinos remain unexplained, while some types of sources are being excluded as dominant neutrino emitters [62–76, 101]. Therefore, more types of sources should be considered, including those not characterized by high-energy electromagnetic emission [80–82].

In this paper, we investigate the possibility that radio-bright AGN contribute to the high-energy astrophysical neutrinos. We use an unbinned maximum-likelihood-ratio method [52, 53, 90, 91] and recently published IceCube muon-track data for 10 years of observation [88, 89]. The 3388 radio-bright AGN form a complete catalog [84] and are selected from the Radio Fundamental Catalog [86], which collects VLBI observations since 1980 (Sec. II A). These sources show very diverse gamma-ray properties. The data comprise 1134450 muon-track events in five different construction phases of IceCube from 2008 to 2018, which have different instrumental response functions (Sec. II B). The unbinned likelihood method (Sec. III) takes into account the information of every single event and is extensively used by IceCube collaboration [52–59] and extensively used other astroparticle experiments (e.g., Refs. [92–95]).

Our results are as follows. From our analysis which investigates every source location (Sec. IV A), we do not find any source with large global significance. The two sources with the highest TS_{max} have global significance of only $\simeq 1.5\sigma$ and 0.8σ despite local significance of $\simeq 4.1\sigma$ and 3.8σ (Table I). Our calculated $\sqrt{\text{TS}_{\text{max}}}$ of all the sources follow the standard normal distribution (Fig. 3), which implies that the background hypothesis is strongly favored, implying that these radio-bright AGN *are not* might not be strong high-energy neutrino emitters. This is further and quantitatively supported by our stacking analysis in Sec. IV B.

Our stacking analysis (Sec. IV B), which seeks a correlation between the neutrinos and the sources, does not find a signal (Fig. 4 left). All the scenarios we considered have their TS values peak at zero with zero best-fit signal event. These null results allow upper limits (95% confidence level) to be placed on the contribution of these sources to the energy flux of the all-sky diffuse astrophysical neutrinos measured by IceCube [2, 5] (Fig. 4 right). In the equal-weighting scheme that assumes high-energy neutrino emission is independent of the sources’ intrinsic properties, these radio-bright AGN account for at most 30.6% between 16 TeV and 2.6 PeV. In the X-band flux-density weighting that weights to the sources by their X-band brightness, these radio-bright AGN account for at most 13.8%. We believe other possibly well-motivated schemes would not lead to a significant correlation, as the distribution of $\sqrt{\text{TS}_{\text{max}}}$ of all the sources closely follows the standard normal distribution and none of the sources show strong global significance (Sec. IV A).

Moreover, our results show that the X-band flux densities of radio-bright AGN might not be an indicator of high-energy neutrino emission because 1) the highest significant sources (some are shown in Table I) do not have higher X-band flux densities compared to the other

sources in the catalog, and 2) from the stacking analysis, the X-band flux-density weighting gives even lower upper limits on the sources' neutrino flux than the equal weighting.

The discovery prospects of high-energy neutrino sources keep increasing rapidly with more and more data to be collected by IceCube [2], KM3NeT [102], Baikal-GVD [103], P-One [104], and especially IceCube-Gen2 (about 10 times bigger than IceCube) [105, 106]. The improvements include statistics, energy range, flavor information, better reconstructions (especially angular resolution), and better sensitivities to certain parts of the sky. Meanwhile, other sources and catalogs in other wavelengths should also be tested, including those that do not show strong GeV–TeV gamma-ray emission [80–82].

ACKNOWLEDGMENTS

We are grateful for helpful discussions with John Beacom, Mauricio Bustamante, Po-Wen Chang, Dan Hooper, Xiaoyuan Huang, Ali Kheirandish, Yuri Y. Kovalev, Ranjan Laha, Tim Linden, and especially Francis Halzen, Qinrui Liu, Kohta Murase, and Alexander Plavin. This work was supported at Johns Hopkins by NSF Grant No. 1818899 and the Simons Foundation. YFL was supported by the National Natural Science Foundation of China (No. U1738136).

Appendix A: Significance of the stacking analysis with injected signal events




FIG. 5. Significance ($\simeq \sqrt{TS_{\max}}$) from our stacking analysis versus fraction of total astrophysical neutrino flux that we inject to these sources. See text for details.

Here we show that if the 3388 sources in the catalog contributed to the diffuse astrophysical neutrino flux observed by IceCube, our analysis would have returned a significant detection. We do not include the simulation uncertainties.

We start with the total diffuse astrophysical neutrino flux measured by IceCube [2, 5], which is about $dF(E_\nu)/dE_\nu = \Phi_0 \times (E_\nu/100 \text{ TeV})^{-2.53}$, where $\Phi_0 = 4.98 \times 10^{-18} \text{ GeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ is the flux normalization at 100 TeV. We then get the number of neutrinos N_ν at different energies and declinations (= zenith angles -90° for IceCube location) in each phase of IceCube by

$$N_\nu^k(E_\nu, \delta) \simeq 2\pi t_k A_{\text{eff}}^k(E_\nu, \delta) E_\nu \frac{dF}{dE_\nu}(E_\nu), \quad (\text{A1})$$

where k labels different phases or data samples of IceCube, just as in Sec. III.

Next, we assign these neutrinos to the 3388 sources with the same weight. For each neutrino with energy E_ν , we use the instrumental response functions for different phases of IceCube [PDF $^k(\sigma|E_\nu)$] provided by Ref. [89] to determine its possible angular error σ . Then, the possible angular distance between the event and its assigned source is obtained by smearing the direction of the event by a 2D symmetric Gaussian distribution with σ . We mix these fake signal events with the true events in each data sample discussed in Sec. II B.

Finally, we do a stacking analysis in the equal weighting scheme just as in Sec. IV B. We get the TS_{\max} for different fluxes of injected neutrinos, characterized by $f \times F(E_\nu)$, where f is the fraction of the astrophysical neutrino flux in Eq. (A1). The background PDFs are unchanged.

Fig. 5 shows the $\sqrt{TS_{\max}}$ (\simeq significance) versus f . The significance increases linearly with the injected signal events because of the relation, $\text{significance} \propto N_S/\sqrt{N_B}$, where N_S and N_B are numbers of signal and background events, respectively. The figure shows that even a small f would have provided a significant signal in our analysis. In fact, an $f = 15\%$ contribution to the total flux would have given a $\sim 3\sigma$ signal, and a 25% contribution would have shown $\sim 5\sigma$. Note that 2σ corresponds to $\sim 10\%$ contribution, which is consistent with our upper limit in the case of equal-weighting and $\Gamma = -2.53$ from our stacking analysis (Sec. IV B).

Appendix B: Results with previously released 2010–2012 data

This work initially used previously released three years of IceCube muon-track data [107] (June 2010 – May 2012) until the 10 years of data were released [88, 89]. Here we present the results using the former dataset as a comparison. This dataset corresponds to IC79, IC86-I, and IC86-II in the newly released dataset except for a few differences as discussed in Sec. II B.




FIG. 6. Same as the left panel of fig. 4 but using previously released three years of IceCube muon-track data.

Fig. 6 shows the result. None of the scenarios show a large significance. The highest peak only has $TS_{\max} \simeq 0.48$. When we change this dataset to the corresponding IC79, IC86-I, and IC86-II in the new dataset, the highest peak lowers to $TS_{\max} \simeq 0.15$ (not shown in the figure), possibly because the events originally have angular errors of less than 0.2° was set to be 0.2° in order to avoid unaccounted-for uncertainties and to ensure no single event dominates the likelihood analysis [89]. The constraints on the fluxes are weaker due to fewer statistics. We do not show the case of $\Gamma = -2.0$ in the equal weighting scheme because it overshoots the astrophysical neutrino spectrum measured by IceCube [2, 5] at higher energies.

-
- [1] M. G. Aartsen *et al.* (IceCube), “Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector,” *Science* **342**, 1242856 (2013), [arXiv:1311.5238 \[astro-ph.HE\]](#).
 - [2] M. G. Aartsen *et al.* (IceCube), “A combined maximum-likelihood analysis of the high-energy astrophysical neutrino flux measured with IceCube,” *Astrophys. J.* **809**, 98 (2015), [arXiv:1507.03991 \[astro-ph.HE\]](#).
 - [3] J. Stettner (IceCube), “Measurement of the Diffuse Astrophysical Muon-Neutrino Spectrum with Ten Years of IceCube Data,” *PoS ICRC2019*, 1017 (2020), [arXiv:1908.09551 \[astro-ph.HE\]](#).
 - [4] Juliana Stachurska (IceCube), “First Double Cascade Tau Neutrino Candidates in IceCube and a New Measurement of the Flavor Composition,” in *36th International Cosmic Ray Conference (ICRC 2019) Madison, Wisconsin, USA, July 24-August 1, 2019* (2019) [arXiv:1908.05506 \[astro-ph.HE\]](#).
 - [5] M. G. Aartsen *et al.* (IceCube), “Characteristics of the diffuse astrophysical electron and tau neutrino flux with six years of IceCube high energy cascade data,” *Phys. Rev. Lett.* **125**, 121104 (2020), [arXiv:2001.09520 \[astro-ph.HE\]](#).
 - [6] R. Abbasi *et al.* (IceCube), “The IceCube high-energy starting event sample: Description and flux characterization with 7.5 years of data,” (2020), [arXiv:2011.03545 \[astro-ph.HE\]](#).
 - [7] Matthew D. Kistler and John F. Beacom, “Guaranteed and Prospective Galactic TeV Neutrino Sources,” *Phys. Rev. D* **74**, 063007 (2006), [arXiv:astro-ph/0607082 \[astro-ph\]](#).
 - [8] John F. Beacom and Matthew D. Kistler, “Dissecting the Cygnus Region with TeV Gamma Rays and Neutrinos,” *Phys. Rev. D* **75**, 083001 (2007), [arXiv:astro-ph/0701751 \[astro-ph\]](#).
 - [9] Kohta Murase, Todd A. Thompson, Brian C. Lacki, and John F. Beacom, “New Class of High-Energy Transients from Crashes of Supernova Ejecta with Massive Circumstellar Material Shells,” *Phys. Rev. D* **84**, 043003 (2011), [arXiv:1012.2834 \[astro-ph.HE\]](#).
 - [10] Kohta Murase and Kunihito Ioka, “TeVPeV Neutrinos from Low-Power Gamma-Ray Burst Jets inside Stars,” *Phys. Rev. Lett.* **111**, 121102 (2013), [arXiv:1306.2274 \[astro-ph.HE\]](#).
 - [11] Kohta Murase, Markus Ahlers, and Brian C. Lacki, “Testing the Hadronuclear Origin of PeV Neutrinos Observed with IceCube,” *Phys. Rev. D* **88**, 121301 (2013), [arXiv:1306.3417 \[astro-ph.HE\]](#).
 - [12] Markus Ahlers and Francis Halzen, “Pinpointing Extragalactic Neutrino Sources in Light of Recent IceCube Observations,” *Phys. Rev. D* **90**, 043005 (2014), [arXiv:1406.2160 \[astro-ph.HE\]](#).
 - [13] Irene Tamborra, Shin’ichiro Ando, and Kohta Murase, “Star-forming galaxies as the origin of diffuse high-energy backgrounds: Gamma-ray and neutrino connections, and implications for starburst history,” *JCAP* **1409**, 043 (2014), [arXiv:1404.1189 \[astro-ph.HE\]](#).
 - [14] Kohta Murase, Yoshiyuki Inoue, and Charles D. Dermer, “Diffuse Neutrino Intensity from the Inner Jets of Active Galactic Nuclei: Impacts of External Photon Fields and the Blazar Sequence,” *Phys. Rev. D* **90**, 023007 (2014), [arXiv:1403.4089 \[astro-ph.HE\]](#).
 - [15] Keith Bechtol, Markus Ahlers, Mattia Di Mauro, Marco Ajello, and Justin Vandenbroucke, “Evidence against star-forming galaxies as the dominant source of IceCube neutrinos,” *Astrophys. J.* **836**, 47 (2017), [arXiv:1511.00688 \[astro-ph.HE\]](#).
 - [16] Matthew D. Kistler and Ranjan Laha, “Multi-PeV Sig-

- nals from a New Astrophysical Neutrino Flux Beyond the Glashow Resonance,” *Phys. Rev. Lett.* **120**, 241105 (2018), arXiv:1605.08781 [astro-ph.HE].
- [17] I. Bartos, M. Ahrens, C. Finley, and S. Marka, “Prospects of Establishing the Origin of Cosmic Neutrinos using Source Catalogs,” *Phys. Rev.* **D96**, 023003 (2017), arXiv:1611.03861 [astro-ph.HE].
- [18] Takahiro Sudoh, Tomonori Totani, and Norita Kawanaka, “High-energy gamma-ray and neutrino production in star-forming galaxies across cosmic time: Difficulties in explaining the IceCube data,” *Publ. Astron. Soc. Jap.* **70**, Publications of the Astronomical Society of Japan, Volume 70, Issue 3, 1 June 2018, 49, <https://doi.org/10.1093/pasj/psy039> (2018), arXiv:1801.09683 [astro-ph.HE].
- [19] Imre Bartos, Doga Veske, Azadeh Keivani, Zsuzsa Marka, Stefan Countryman, Erik Blaufuss, Chad Finley, and Szabolcs Marka, “Bayesian Multi-Messenger Search Method for Common Sources of Gravitational Waves and High-Energy Neutrinos,” *Phys. Rev.* **D100**, 083017 (2019), arXiv:1810.11467 [astro-ph.HE].
- [20] Mauricio Bustamante and Markus Ahlers, “Inferring the flavor of high-energy astrophysical neutrinos at their sources,” *Phys. Rev. Lett.* **122**, 241101 (2019), arXiv:1901.10087 [astro-ph.HE].
- [21] T. Hovatta *et al.*, “Association of IceCube neutrinos with radio sources observed at Owens Valley and Metsähovi Radio Observatories,” (2020), arXiv:2009.10523 [astro-ph.HE].
- [22] Dan Hooper, “Measuring high-energy neutrino nucleon cross-sections with future neutrino telescopes,” *Phys. Rev.* **D65**, 097303 (2002), arXiv:hep-ph/0203239 [hep-ph].
- [23] E. Borriello, A. Cuoco, G. Mangano, G. Miele, S. Pastor, O. Pisanti, and P. D. Serpico, “Disentangling neutrino-nucleon cross section and high energy neutrino flux with a km³ neutrino telescope,” *Phys. Rev.* **D77**, 045019 (2008), arXiv:0711.0152 [astro-ph].
- [24] Amy Connolly, Robert S. Thorne, and David Waters, “Calculation of High Energy Neutrino-Nucleon Cross Sections and Uncertainties Using the MSTW Parton Distribution Functions and Implications for Future Experiments,” *Phys. Rev.* **D83**, 113009 (2011), arXiv:1102.0691 [hep-ph].
- [25] Spencer R. Klein and Amy Connolly, “Neutrino Absorption in the Earth, Neutrino Cross-Sections, and New Physics,” in *Proceedings, 2013 Community Summer Study on the Future of U.S. Particle Physics: Snowmass on the Mississippi (CSS2013): Minneapolis, MN, USA, July 29-August 6, 2013* (2013) arXiv:1304.4891 [astro-ph.HE].
- [26] Rhorry Gauld, Juan Rojo, Luca Rottoli, Subir Sarkar, and Jim Talbert, “The prompt atmospheric neutrino flux in the light of LHCb,” *JHEP* **02**, 130 (2016), arXiv:1511.06346 [hep-ph].
- [27] M. G. Aartsen *et al.* (IceCube), “Measurement of the multi-TeV neutrino cross section with IceCube using Earth absorption,” *Nature* **551**, 596–600 (2017), arXiv:1711.08119 [hep-ex].
- [28] Mauricio Bustamante and Amy Connolly, “Extracting the Energy-Dependent Neutrino-Nucleon Cross Section above 10 TeV Using IceCube Showers,” *Phys. Rev. Lett.* **122**, 041101 (2019), arXiv:1711.11043 [astro-ph.HE].
- [29] F. Cornet, Jose I. Illana, and M. Masip, “TeV strings and the neutrino nucleon cross-section at ultrahigh-energies,” *Phys. Rev. Lett.* **86**, 4235–4238 (2001), arXiv:hep-ph/0102065 [hep-ph].
- [30] Paolo Lipari, “CP violation effects and high-energy neutrinos,” *Phys. Rev.* **D64**, 033002 (2001), arXiv:hep-ph/0102046 [hep-ph].
- [31] Jaime Alvarez-Muniz, Jonathan L. Feng, Francis Halzen, Tao Han, and Dan Hooper, “Detecting microscopic black holes with neutrino telescopes,” *Phys. Rev.* **D65**, 124015 (2002), arXiv:hep-ph/0202081 [hep-ph].
- [32] John F. Beacom, Nicole F. Bell, Dan Hooper, Sandip Pakvasa, and Thomas J. Weiler, “Decay of High-Energy Astrophysical Neutrinos,” *Phys. Rev. Lett.* **90**, 181301 (2003), arXiv:hep-ph/0211305 [hep-ph].
- [33] Tao Han and Dan Hooper, “Effects of electroweak instantons in high-energy neutrino telescopes,” *Phys. Lett.* **B582**, 21–26 (2004), arXiv:hep-ph/0307120 [hep-ph].
- [34] Tao Han and Dan Hooper, “The Particle physics reach of high-energy neutrino astronomy,” *New J. Phys.* **6**, 150 (2004), arXiv:hep-ph/0408348 [hep-ph].
- [35] Dan Hooper, Dean Morgan, and Elizabeth Winstanley, “Probing quantum decoherence with high-energy neutrinos,” *Phys. Lett.* **B609**, 206–211 (2005), arXiv:hep-ph/0410094 [hep-ph].
- [36] Dan Hooper, Dean Morgan, and Elizabeth Winstanley, “Lorentz and CPT invariance violation in high-energy neutrinos,” *Phys. Rev.* **D72**, 065009 (2005), arXiv:hep-ph/0506091 [hep-ph].
- [37] M. C. Gonzalez-Garcia, F. Halzen, and M. Maltoni, “Physics reach of high-energy and high-statistics icecube atmospheric neutrino data,” *Phys. Rev.* **D71**, 093010 (2005), arXiv:hep-ph/0502223 [hep-ph].
- [38] Kenny C. Y. Ng and John F. Beacom, “Cosmic neutrino cascades from secret neutrino interactions,” *Phys. Rev.* **D90**, 065035 (2014), [Erratum: *Phys. Rev.* **D90**, no.8, 089904 (2014)], arXiv:1404.2288 [astro-ph.HE].
- [39] Kunihto Ioka and Kohta Murase, “IceCube PeVEeV neutrinos and secret interactions of neutrinos,” *PTEP* **2014**, 061E01 (2014), arXiv:1404.2279 [astro-ph.HE].
- [40] Mauricio Bustamante, John F. Beacom, and Kohta Murase, “Testing decay of astrophysical neutrinos with incomplete information,” *Phys. Rev.* **D95**, 063013 (2017), arXiv:1610.02096 [astro-ph.HE].
- [41] Jordi Salvado, Olga Mena, Sergio Palomares-Ruiz, and Nuria Rius, “Non-standard interactions with high-energy atmospheric neutrinos at IceCube,” *JHEP* **01**, 141 (2017), arXiv:1609.03450 [hep-ph].
- [42] Pilar Coloma, Pedro A. N. Machado, Ivan Martinez-Soler, and Ian M. Shoemaker, “Double-Cascade Events from New Physics in Icecube,” *Phys. Rev. Lett.* **119**, 201804 (2017), arXiv:1707.08573 [hep-ph].
- [43] Bei Zhou and John F. Beacom, “Neutrino-nucleus cross sections for W-boson and trident production,” *Phys. Rev.* **D101**, 036011 (2020), arXiv:1910.08090 [hep-ph].
- [44] Bei Zhou and John F. Beacom, “W -boson and trident production in TeV–PeV neutrino observatories,” *Phys. Rev. D* **101**, 036010 (2020), arXiv:1910.10720 [hep-ph].
- [45] M. G. Aartsen *et al.* (IceCube), “Detection of a particle shower at the Glashow resonance with IceCube,” *Nature* **591**, 220–224 (2021).
- [46] Francis Halzen, Alexander Kappes, and Aongus O’Murchadha, “Prospects for identifying the sources

- of the Galactic cosmic rays with IceCube,” *Phys. Rev. D* **78**, 063004 (2008), arXiv:0803.0314 [astro-ph].
- [47] M. G. Aartsen *et al.* (IceCube), “Probing the origin of cosmic rays with extremely high energy neutrinos using the IceCube Observatory,” *Phys. Rev. D* **88**, 112008 (2013), arXiv:1310.5477 [astro-ph.HE].
- [48] Matthew D. Kistler, Todor Stanev, and Hasan Yüksel, “Cosmic PeV Neutrinos and the Sources of Ultrahigh Energy Protons,” *Phys. Rev. D* **90**, 123006 (2014), arXiv:1301.1703 [astro-ph.HE].
- [49] Luis A. Anchordoqui, Haim Goldberg, Morgan H. Lynch, Angela V. Olinto, Thomas C. Paul, and Thomas J. Weiler, “Pinning down the cosmic ray source mechanism with new IceCube data,” *Phys. Rev. D* **89**, 083003 (2014), arXiv:1306.5021 [astro-ph.HE].
- [50] Ke Fang, Kumiko Kotera, Kohta Murase, and Angela V. Olinto, “Testing the Newborn Pulsar Origin of Ultrahigh Energy Cosmic Rays with EeV Neutrinos,” *Phys. Rev. D* **90**, 103005 (2014), [Erratum: *Phys. Rev. D* **92**, no.12, 129901 (2015); *Phys. Rev. D* **90**, 103005 (2014)], arXiv:1311.2044 [astro-ph.HE].
- [51] M. G. Aartsen *et al.* (IceCube), “Constraints on Ultrahigh-Energy Cosmic-Ray Sources from a Search for Neutrinos above 10 PeV with IceCube,” *Phys. Rev. Lett.* **117**, 241101 (2016), [Erratum: *Phys. Rev. Lett.* **119**, no.25, 259902 (2017)], arXiv:1607.05886 [astro-ph.HE].
- [52] R. Abbasi *et al.* (IceCube), “Time-Integrated Searches for Point-like Sources of Neutrinos with the 40-String IceCube Detector,” *Astrophys. J.* **732**, 18 (2011), arXiv:1012.2137 [astro-ph.HE].
- [53] M. G. Aartsen *et al.* (IceCube), “Search for Time-independent Neutrino Emission from Astrophysical Sources with 3 yr of IceCube Data,” *Astrophys. J.* **779**, 132 (2013), arXiv:1307.6669 [astro-ph.HE].
- [54] M. G. Aartsen *et al.* (IceCube), “Searches for Extended and Point-like Neutrino Sources with Four Years of IceCube Data,” *Astrophys. J.* **796**, 109 (2014), arXiv:1406.6757 [astro-ph.HE].
- [55] M. G. Aartsen *et al.* (IceCube), “Searches for Time Dependent Neutrino Sources with IceCube Data from 2008 to 2012,” *Astrophys. J.* **807**, 46 (2015), arXiv:1503.00598 [astro-ph.HE].
- [56] M. G. Aartsen *et al.* (IceCube), “All-sky Search for Time-integrated Neutrino Emission from Astrophysical Sources with 7 yr of IceCube Data,” *Astrophys. J.* **835**, 151 (2017), arXiv:1609.04981 [astro-ph.HE].
- [57] M. G. Aartsen *et al.* (IceCube), “Search for steady point-like sources in the astrophysical muon neutrino flux with 8 years of IceCube data,” *Eur. Phys. J. C* **79**, 234 (2019), arXiv:1811.07979 [hep-ph].
- [58] M. G. Aartsen *et al.* (IceCube), “Time-Integrated Neutrino Source Searches with 10 Years of IceCube Data,” *Phys. Rev. Lett.* **124**, 051103 (2020), arXiv:1910.08488 [astro-ph.HE].
- [59] R. Abbasi *et al.* (IceCube), “A search for time-dependent astrophysical neutrino emission with IceCube data from 2012 to 2017,” (2020), arXiv:2012.01079 [astro-ph.HE].
- [60] M. G. Aartsen *et al.* (IceCube), “Neutrino emission from the direction of the blazar TXS 0506+056 prior to the IceCube-170922A alert,” *Science* **361**, 147–151 (2018), arXiv:1807.08794 [astro-ph.HE].
- [61] M. G. Aartsen *et al.* (IceCube, Fermi-LAT, MAGIC, AGILE, ASAS-SN, HAWC, H.E.S.S., INTEGRAL, Kanata, Kiso, Kapteyn, Liverpool Telescope, Subaru, Swift NuSTAR, VERITAS, VLA/17B-403), “Multimes-senger observations of a flaring blazar coincident with high-energy neutrino IceCube-170922A,” *Science* **361**, eaat1378 (2018), arXiv:1807.08816 [astro-ph.HE].
- [62] Eli Waxman and John N. Bahcall, “High-energy neutrinos from cosmological gamma-ray burst fireballs,” *Phys. Rev. Lett.* **78**, 2292–2295 (1997), arXiv:astro-ph/9701231 [astro-ph].
- [63] R. Abbasi *et al.* (IceCube), “Search for muon neutrinos from Gamma-Ray Bursts with the IceCube neutrino telescope,” *Astrophys. J.* **710**, 346–359 (2010), arXiv:0907.2227 [astro-ph.HE].
- [64] R. Abbasi *et al.* (IceCube), “Limits on Neutrino Emission from Gamma-Ray Bursts with the 40 String IceCube Detector,” *Phys. Rev. Lett.* **106**, 141101 (2011), arXiv:1101.1448 [astro-ph.HE].
- [65] R. Abbasi *et al.* (IceCube), “An absence of neutrinos associated with cosmic-ray acceleration in γ -ray bursts,” *Nature* **484**, 351–353 (2012), arXiv:1204.4219 [astro-ph.HE].
- [66] M. G. Aartsen *et al.* (IceCube), “Search for Prompt Neutrino Emission from Gamma-Ray Bursts with IceCube,” *Astrophys. J. Lett.* **805**, L5 (2015), arXiv:1412.6510 [astro-ph.HE].
- [67] M. G. Aartsen *et al.* (IceCube), “An All-Sky Search for Three Flavors of Neutrinos from Gamma-Ray Bursts with the IceCube Neutrino Observatory,” *Astrophys. J.* **824**, 115 (2016), arXiv:1601.06484 [astro-ph.HE].
- [68] M. G. Aartsen *et al.* (IceCube), “Extending the search for muon neutrinos coincident with gamma-ray bursts in IceCube data,” *Astrophys. J.* **843**, 112 (2017), arXiv:1702.06868 [astro-ph.HE].
- [69] M. G. Aartsen *et al.* (IceCube), “The contribution of Fermi-2LAC blazars to the diffuse TeV-PeV neutrino flux,” *Astrophys. J.* **835**, 45 (2017), arXiv:1611.03874 [astro-ph.HE].
- [70] Dan Hooper, Tim Linden, and Abby Vieregg, “Active Galactic Nuclei and the Origin of IceCube’s Diffuse Neutrino Flux,” *JCAP* **1902**, 012 (2019), arXiv:1810.02823 [astro-ph.HE].
- [71] Chengchao Yuan, Kohta Murase, and Peter Mszros, “Complementarity of Stacking and Multiplet Constraints on the Blazar Contribution to the Cumulative High-Energy Neutrino Intensity,” *Astrophys. J.* **890**, 25 (2020), arXiv:1904.06371 [astro-ph.HE].
- [72] Jia-Wei Luo and Bing Zhang, “Blazar - IceCube neutrino association revisited,” *Phys. Rev. D* **101**, 103015 (2020), arXiv:2004.09686 [astro-ph.HE].
- [73] Daniel Smith, Dan Hooper, and Abigail Vieregg, “Revisiting AGN as the source of IceCubes diffuse neutrino flux,” *JCAP* **2103**, 031 (2021), arXiv:2007.12706 [astro-ph.HE].
- [74] Nicholas Senno, Kohta Murase, and Peter Mszros, “Constraining high-energy neutrino emission from choked jets in stripped-envelope supernovae,” *JCAP* **1801**, 025 (2018), arXiv:1706.02175 [astro-ph.HE].
- [75] Arman Esmaili and Kohta Murase, “Constraining high-energy neutrinos from choked-jet supernovae with IceCube high-energy starting events,” *JCAP* **1812**, 008 (2018), arXiv:1809.09610 [hep-ph].
- [76] M. G. Aartsen *et al.* (IceCube), “IceCube Search for High-Energy Neutrino Emission from TeV Pul-

- sar Wind Nebulae,” *Astrophys. J.* **898**, 117 (2020), [arXiv:2003.12071 \[astro-ph.HE\]](#).
- [77] Armen Atoyan and Charles D. Dermer, “High-energy neutrinos from photomeson processes in blazars,” *Phys. Rev. Lett.* **87**, 221102 (2001), [arXiv:astro-ph/0108053 \[astro-ph\]](#).
- [78] Warren Essey, Oleg Kalashev, Alexander Kusenko, and John F. Beacom, “Role of line-of-sight cosmic ray interactions in forming the spectra of distant blazars in TeV gamma rays and high-energy neutrinos,” *Astrophys. J.* **731**, 51 (2011), [arXiv:1011.6340 \[astro-ph.HE\]](#).
- [79] R. J. Protheroe, “High-energy neutrinos from blazars,” *ASP Conf. Ser.* **121**, 585 (1997), [arXiv:astro-ph/9607165 \[astro-ph\]](#).
- [80] Kohta Murase, Dafne Guetta, and Markus Ahlers, “Hidden Cosmic-Ray Accelerators as an Origin of TeV-PeV Cosmic Neutrinos,” *Phys. Rev. Lett.* **116**, 071101 (2016), [arXiv:1509.00805 \[astro-ph.HE\]](#).
- [81] Antonio Capanema, Arman Esmaili, and Kohta Murase, “New constraints on the origin of medium-energy neutrinos observed by IceCube,” *Phys. Rev. D* **101**, 103012 (2020), [arXiv:2002.07192 \[hep-ph\]](#).
- [82] Antonio Capanema, Arman Esmaili, and Pasquale Dario Serpico, “Where do IceCube neutrinos come from? Hints from the diffuse gamma-ray flux,” *JCAP* **2102**, 037 (2021), [arXiv:2007.07911 \[hep-ph\]](#).
- [83] M. Ackermann *et al.* (Fermi-LAT), “The spectrum of isotropic diffuse gamma-ray emission between 100 MeV and 820 GeV,” *Astrophys. J.* **799**, 86 (2015), [arXiv:1410.3696 \[astro-ph.HE\]](#).
- [84] Alexander Plavin, Yuri Y. Kovalev, Yuri A. Kovalev, and Sergey Troitsky, “Observational Evidence for the Origin of High-energy Neutrinos in Parsec-scale Nuclei of Radio-bright Active Galaxies,” *Astrophys. J.* **894**, 101 (2020), [arXiv:2001.00930 \[astro-ph.HE\]](#).
- [85] A. V. Plavin, Y. Y. Kovalev, Yu. A. Kovalev, and S. V. Troitsky, “Directional Association of TeV to PeV Astrophysical Neutrinos with Radio Blazars,” *Astrophys. J.* **908**, 157 (2021), [arXiv:2009.08914 \[astro-ph.HE\]](#).
- [86] <http://astrogeo.org/rfc/>.
- [87] IceCube Collaboration (2019), “All-sky search for point sources with seven years of IceCube data. IceCube Neutrino Observatory. Dataset.” [10.21234/exm3-tm26](#).
- [88] R. Abbasi *et al.* (IceCube), “IceCube Data for Neutrino Point-Source Searches Years 2008-2018,” (2021), [10.21234/CPKQ-K003](#), [arXiv:2101.09836 \[astro-ph.HE\]](#).
- [89] IceCube Collaboration (2021), “All-sky point-source IceCube data: years 2008-2018. Dataset. ,” [10.21234/sxvs-mt83](#).
- [90] Jim Braun, Jon Dumm, Francesco De Palma, Chad Finley, Albrecht Karle, and Teresa Montaruli, “Methods for point source analysis in high energy neutrino telescopes,” *Astropart. Phys.* **29**, 299–305 (2008), [arXiv:0801.1604 \[astro-ph\]](#).
- [91] Jim Braun, Mike Baker, Jon Dumm, Chad Finley, Albrecht Karle, and Teresa Montaruli, “Time-Dependent Point Source Search Methods in High Energy Neutrino Astronomy,” *Astropart. Phys.* **33**, 175–181 (2010), [arXiv:0912.1572 \[astro-ph.IM\]](#).
- [92] K. Bays *et al.* (Super-Kamiokande), “Supernova Relic Neutrino Search at Super-Kamiokande,” *Phys. Rev. D* **85**, 052007 (2012), [arXiv:1111.5031 \[hep-ex\]](#).
- [93] M. Ackermann *et al.* (Fermi-LAT), “Updated search for spectral lines from Galactic dark matter interactions with pass 8 data from the Fermi Large Area Telescope,” *Phys. Rev. D* **91**, 122002 (2015), [arXiv:1506.00013 \[astro-ph.HE\]](#).
- [94] M. L. Ahnen *et al.* (MAGIC, Fermi-LAT), “Limits to Dark Matter Annihilation Cross-Section from a Combined Analysis of MAGIC and Fermi-LAT Observations of Dwarf Satellite Galaxies,” *JCAP* **1602**, 039 (2016), [arXiv:1601.06590 \[astro-ph.HE\]](#).
- [95] H. Abdallah *et al.* (HESS), “Search for γ -Ray Line Signals from Dark Matter Annihilations in the Inner Galactic Halo from 10 Years of Observations with H.E.S.S.” *Phys. Rev. Lett.* **120**, 201101 (2018), [arXiv:1805.05741 \[astro-ph.HE\]](#).
- [96] <http://astrogeo.org/>.
- [97] A. Achterberg *et al.* (IceCube), “First Year Performance of The IceCube Neutrino Telescope,” *Astropart. Phys.* **26**, 155–173 (2006), [arXiv:astro-ph/0604450 \[astro-ph\]](#).
- [98] <https://icecube.wisc.edu/>.
- [99] https://www.icrc2019.org/uploads/1/1/9/0/119067782/nu4b_schneider.pdf.
- [100] S. S. Wilks, “The Large-Sample Distribution of the Likelihood Ratio for Testing Composite Hypotheses,” *The Annals of Mathematical Statistics* **9**, 60 – 62 (1938).
- [101] Kohta Murase and Eli Waxman, “Constraining High-Energy Cosmic Neutrino Sources: Implications and Prospects,” *Phys. Rev. D* **94**, 103006 (2016), [arXiv:1607.01601 \[astro-ph.HE\]](#).
- [102] S. Adrian-Martinez *et al.* (KM3NeT), “Letter of intent for KM3NeT 2.0,” *J. Phys.* **G43**, 084001 (2016), [arXiv:1601.07459 \[astro-ph.IM\]](#).
- [103] A. D. Avrorin *et al.* (BAIKAL), “The prototyping/early construction phase of the BAIKAL-GVD project,” *Proceedings, 4th Roma International Conference on Astroparticle Physics (RICAP 13): Rome, Italy, May 22-24, 2013*, *Nucl. Instrum. Meth.* **A742**, 82–88 (2014), [arXiv:1308.1833 \[astro-ph.IM\]](#).
- [104] Matteo Agostini *et al.* (P-ONE), “The Pacific Ocean Neutrino Experiment,” *Nature Astron.* **4**, 913–915 (2020), [arXiv:2005.09493 \[astro-ph.HE\]](#).
- [105] Erik Blaufuss, C. Kopper, and C. Haack (IceCube-Gen2), “The IceCube-Gen2 High Energy Array,” *Proceedings, 34th International Cosmic Ray Conference (ICRC 2015): The Hague, The Netherlands, July 30-August 6, 2015*, *PoS ICRC2015*, 1146 (2016).
- [106] M. G. Aartsen *et al.* (IceCube Gen2), “IceCube-Gen2: The Window to the Extreme Universe,” (2020), [arXiv:2008.04323 \[astro-ph.HE\]](#).
- [107] IceCube Collaboration (2018), “All-sky point-source IceCube data: years 2010-2012. Dataset.” [10.21234/B4F04V](#).