



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Dark Energy Survey Year 3 Results: Three-point shear correlations and mass aperture moments

L. F. Secco et al. (DES Collaboration)

Phys. Rev. D **105**, 103537 — Published 31 May 2022

DOI: [10.1103/PhysRevD.105.103537](https://doi.org/10.1103/PhysRevD.105.103537)

Dark Energy Survey Year 3 Results: Three-Point Shear Correlations and Mass Aperture Moments

L. F. Secco,^{1,*} M. Jarvis,² B. Jain,² C. Chang,^{1,3} M. Gatti,² J. Frieman,^{1,3,4} S. Adhikari,^{1,3}
A. Alarcon,⁵ A. Amon,^{6,7} K. Bechtol,⁸ M. R. Becker,⁵ G. M. Bernstein,² J. Blazek,^{9,10} A. Campos,¹¹
A. Carnero Rosell,^{12,13,14} M. Carrasco Kind,^{15,16} A. Choi,¹⁷ J. Cordero,¹⁸ J. DeRose,¹⁹ S. Dodelson,^{11,20}
C. Doux,² A. Drlica-Wagner,^{3,4,1} S. Everett,²¹ G. Giannini,²² D. Gruen,²³ R. A. Gruendl,^{15,16} I. Harrison,^{24,18,25}
W. G. Hartley,²⁶ K. Herner,⁴ E. Krause,²⁷ N. MacCrann,²⁸ J. McCullough,²⁹ J. Myles,^{30,29,31} A.
Navarro-Alsina,³² J. Prat,^{3,1} R. P. Rollins,¹⁸ S. Samuroff,¹¹ C. Sánchez,² I. Sevilla-Noarbe,³³ E. Sheldon,³⁴
M. A. Troxel,³⁵ D. Zeurher,³⁶ M. Aguena,¹³ F. Andrade-Oliveira,³⁷ J. Annis,⁴ D. Bacon,³⁸ E. Bertin,^{39,40}
S. Bocquet,²³ D. Brooks,⁴¹ D. L. Burke,^{29,31} J. Carretero,²² F. J. Castander,^{42,43} M. Crocce,^{42,43} L. N. da
Costa,^{13,44} M. E. S. Pereira,^{37,45} J. De Vicente,³³ H. T. Diehl,⁴ P. Doel,⁴¹ K. Eckert,² I. Ferrero,⁴⁶
B. Flaugher,⁴ D. Friedel,¹⁵ J. García-Bellido,⁴⁷ G. Gutierrez,⁴ S. R. Hinton,⁴⁸ D. L. Hollowood,²¹
K. Honscheid,^{49,50} D. Huterer,³⁷ K. Kuehn,^{51,52} N. Kuropatkin,⁴ M. A. G. Maia,^{13,44} J. L. Marshall,⁵³
F. Menanteau,^{15,16} R. Miquel,^{54,22} J. J. Mohr,^{23,55} R. Morgan,⁸ J. Muir,⁵⁶ F. Paz-Chinchón,^{15,57}
A. Pieres,^{13,44} A. A. Plazas Malagón,⁵⁸ M. Rodríguez-Monroy,³³ A. Roodman,^{29,31} E. Sanchez,³³
S. Serrano,^{42,43} E. Suchyta,⁵⁹ M. E. C. Swanson,⁶⁰ G. Tarle,³⁷ D. Thomas,³⁸ C. To,⁴⁹ and J. Weller^{55,61}

(DES Collaboration)

¹Kavli Institute for Cosmological Physics, University of Chicago, Chicago, IL 60637, USA

²Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104, USA

³Department of Astronomy and Astrophysics, University of Chicago, Chicago, IL 60637, USA

⁴Fermi National Accelerator Laboratory, P. O. Box 500, Batavia, IL 60510, USA

⁵Argonne National Laboratory, 9700 South Cass Avenue, Lemont, IL 60439, USA

⁶Kavli Institute for Cosmology, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

⁷Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge, CB3 0HA

⁸Physics Department, 2320 Chamberlin Hall, University of Wisconsin-Madison,
1150 University Avenue Madison, WI 53706-1390

⁹Department of Physics, Northeastern University, Boston, MA 02115, USA

¹⁰Laboratory of Astrophysics, École Polytechnique Fédérale de Lausanne (EPFL),
Observatoire de Sauverny, 1290 Versoix, Switzerland

¹¹Department of Physics, Carnegie Mellon University, Pittsburgh, Pennsylvania 15312, USA

¹²Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain

¹³Laboratório Interinstitucional de e-Astronomia - LIneA,
Rua Gal. José Cristino 77, Rio de Janeiro, RJ - 20921-400, Brazil

¹⁴Universidad de La Laguna, Dpto. Astrofísica, E-38206 La Laguna, Tenerife, Spain

¹⁵Center for Astrophysical Surveys, National Center for Supercomputing
Applications, 1205 West Clark St., Urbana, IL 61801, USA

¹⁶Department of Astronomy, University of Illinois at Urbana-Champaign, 1002 W. Green Street, Urbana, IL 61801, USA

¹⁷California Institute of Technology, 1200 East California Blvd, MC 249-17, Pasadena, CA 91125, USA

¹⁸Jodrell Bank Center for Astrophysics, School of Physics and Astronomy,
University of Manchester, Oxford Road, Manchester, M13 9PL, UK

¹⁹Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA

²⁰NSF AI Planning Institute for Physics of the Future,
Carnegie Mellon University, Pittsburgh, PA 15213, USA

²¹Santa Cruz Institute for Particle Physics, Santa Cruz, CA 95064, USA

²²Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology,
Campus UAB, 08193 Bellaterra (Barcelona) Spain

²³Faculty of Physics, Ludwig-Maximilians-Universität, Scheinerstr. 1, 81679 Munich, Germany

²⁴Department of Physics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

²⁵School of Physics and Astronomy, Cardiff University, CF24 3AA, UK

²⁶Department of Astronomy, University of Geneva, ch. d'Écogia 16, CH-1290 Versoix, Switzerland

²⁷Department of Astronomy/Steward Observatory, University of Arizona,
933 North Cherry Avenue, Tucson, AZ 85721-0065, USA

²⁸Department of Applied Mathematics and Theoretical Physics,
University of Cambridge, Cambridge CB3 0WA, UK

²⁹Kavli Institute for Particle Astrophysics & Cosmology,
P. O. Box 2450, Stanford University, Stanford, CA 94305, USA

³⁰Department of Physics, Stanford University, 382 Via Pueblo Mall, Stanford, CA 94305, USA

³¹SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA

³²Instituto de Física Gleb Wataghin, Universidade Estadual de Campinas, 13083-859, Campinas, SP, Brazil

- ³³ *Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain*
- ³⁴ *Brookhaven National Laboratory, Bldg 510, Upton, NY 11973, USA*
- ³⁵ *Department of Physics, Duke University Durham, NC 27708, USA*
- ³⁶ *Department of Physics, ETH Zurich, Wolfgang-Pauli-Strasse 16, CH-8093 Zurich, Switzerland*
- ³⁷ *Department of Physics, University of Michigan, Ann Arbor, MI 48109, USA*
- ³⁸ *Institute of Cosmology and Gravitation, University of Portsmouth, Portsmouth, PO1 3FX, UK*
- ³⁹ *CNRS, UMR 7095, Institut d'Astrophysique de Paris, F-75014, Paris, France*
- ⁴⁰ *Sorbonne Universités, UPMC Univ Paris 06, UMR 7095, Institut d'Astrophysique de Paris, F-75014, Paris, France*
- ⁴¹ *Department of Physics & Astronomy, University College London, Gower Street, London, WC1E 6BT, UK*
- ⁴² *Institut d'Estudis Espacials de Catalunya (IEEC), 08034 Barcelona, Spain*
- ⁴³ *Institute of Space Sciences (ICE, CSIC), Campus UAB, Carrer de Can Magrans, s/n, 08193 Barcelona, Spain*
- ⁴⁴ *Observatório Nacional, Rua Gal. José Cristino 77, Rio de Janeiro, RJ - 20921-400, Brazil*
- ⁴⁵ *Hamburger Sternwarte, Universität Hamburg, Gojenbergsweg 112, 21029 Hamburg, Germany*
- ⁴⁶ *Institute of Theoretical Astrophysics, University of Oslo. P.O. Box 1029 Blindern, NO-0315 Oslo, Norway*
- ⁴⁷ *Instituto de Física Teórica UAM/CSIC, Universidad Autónoma de Madrid, 28049 Madrid, Spain*
- ⁴⁸ *School of Mathematics and Physics, University of Queensland, Brisbane, QLD 4072, Australia*
- ⁴⁹ *Center for Cosmology and Astro-Particle Physics, The Ohio State University, Columbus, OH 43210, USA*
- ⁵⁰ *Department of Physics, The Ohio State University, Columbus, OH 43210, USA*
- ⁵¹ *Australian Astronomical Optics, Macquarie University, North Ryde, NSW 2113, Australia*
- ⁵² *Lowell Observatory, 1400 Mars Hill Rd, Flagstaff, AZ 86001, USA*
- ⁵³ *George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, and Department of Physics and Astronomy, Texas A&M University, College Station, TX 77843, USA*
- ⁵⁴ *Institució Catalana de Recerca i Estudis Avançats, E-08010 Barcelona, Spain*
- ⁵⁵ *Max Planck Institute for Extraterrestrial Physics, Giessenbachstrasse, 85748 Garching, Germany*
- ⁵⁶ *Perimeter Institute for Theoretical Physics, 31 Caroline St. North, Waterloo, ON N2L 2Y5, Canada*
- ⁵⁷ *Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK*
- ⁵⁸ *Department of Astrophysical Sciences, Princeton University, Peyton Hall, Princeton, NJ 08544, USA*
- ⁵⁹ *Computer Science and Mathematics Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831*
- ⁶⁰ *National Center for Supercomputing Applications, 1205 West Clark St., Urbana, IL 61801, USA*
- ⁶¹ *Universitäts-Sternwarte, Fakultät für Physik, Ludwig-Maximilians Universität München, Scheinerstr. 1, 81679 München, Germany*

(Dated: April 26, 2022)

We present high signal-to-noise measurements of three-point shear correlations and the third moment of the mass aperture statistic using the first 3 years of data from the Dark Energy Survey. We additionally obtain the first measurements of the configuration and scale dependence of the four three-point shear correlations which carry cosmological information. With the third-order mass aperture statistic, we present tomographic measurements over angular scales of 4 to 60 arcminutes with a combined statistical significance of 15.0σ . Using the tomographic information and measuring also the second-order mass aperture, we additionally obtain a skewness parameter and its redshift evolution. We find that the amplitudes and scale-dependence of these shear 3pt functions are in qualitative agreement with measurements in a mock galaxy catalog based on N-body simulations, indicating promise for including them in future cosmological analyses. We validate our measurements by showing that B-modes, parity-violating contributions and PSF modeling uncertainties are negligible, and determine that the measured signals are likely to be of astrophysical and gravitational origin.

I. INTRODUCTION

Two-point (2pt) auto-correlation functions of the shear field (sometimes referred to as cosmic shear) have been widely used in the recent literature to constrain cosmological parameters. Current works utilize different statistical measures and exploit the shear distributions in both real (configuration) space as well as harmonic space (Asgari et al. (2021), Hamana et al. (2020), Hikage et al.

(2019), Amon et al. (2021), Secco & Samuroff et al., (2021)). One of the main products of years of effort by the community is the accurate determination of the amplitude parameter $S_8 \equiv \sigma_8 \sqrt{\Omega_m}/0.3$, where σ_8 is the root mean square amplitude of the linear-theory matter power spectrum at $z = 0$ over an 8 Mpc/h scale, and Ω_m is the matter density at $z = 0$. This amplitude is in mild tension with the value inferred from fluctuations of the Cosmic Microwave Background (Planck Collaboration 2020) by about 2σ (depending on the survey data sample used) and its origin remains unresolved.

* secco@uchicago.edu

Extracting more cosmological information from the

shear field than that encoded in 2pt statistics may help better characterize this tension and is an important goal in itself. To be useful, the additional information should have *its systematics well-understood and controlled*. The aim of this work is to address both points above: we use data from DES Y3, the first 3 years of data from the Dark Energy Survey (DES Collaboration (2022), Gatti, Sheldon et al. (2021b), Sevilla-Noarbe & Bechtol et al., (2021)) to obtain high signal-to-noise measurements of three-point (3pt) correlation functions of the shear field and show that potential contaminants in these measurements coming from observational and instrumental origins are negligible.

The benefits of utilizing higher order correlations as a cosmological probe are plenty and go far beyond simply enabling access to non-Gaussian information in the shear and matter bispectrum. Compared to 2pt functions, 3pt correlations in lensing carry different cosmological parameter degeneracies (Bernardeau et al. 2002, Kayo & Takada 2013, Takada & Jain 2003b) and when combined with 2pt functions can additionally constrain astrophysical and systematic nuisance parameters (Huterer et al. 2006, Pyne & Joachimi 2021, Semboloni et al. 2013, Troxel & Ishak 2012). The combination of 2pt and 3pt lensing data vectors is thus greater than the sum of its parts, and enables degeneracy-breaking in both the cosmological and nuisance parameter spaces.

The community has followed several approaches to extracting the information contained in higher order shear statistics. For example, non-Gaussian information can be obtained with position-dependent or integrated 2pt lensing signatures (Halder et al. 2021, Jung et al. 2021), peak statistics (Kacprzak et al. 2016, Zürcher et al. 2021), density splits of the shear field (Friedrich et al. 2018, Gruen et al. 2018) as well as with techniques borrowed from artificial intelligence and neural networks (Cheng et al. 2020, Fluri et al. 2019, Jeffrey et al. 2021, Lu et al. 2021). Another approach is to *directly measure* 3rd or higher order statistics of the shear field in the form of ellipticity correlations (Van Waerbeke et al. (2002), Benabed & Scoccimarro (2006)), mass aperture moments (Fu et al. 2014, Jarvis et al. 2004, Semboloni et al. 2011) or lensing mass maps (Gatti et al. 2021a).

In this work, we follow the latter approach and directly measure 3pt statistics of the DES Y3 data in the form of “natural” correlation functions (the three-point equivalents of ξ_{\pm}) (Schneider & Lombardi 2003) and the third moment of the mass aperture statistic (Schneider et al. 1998). We detect both statistics at high significance and additionally explore the triangle configuration dependence, tomographic signals and redshift evolution of the 3pt lensing signal, none of which have been previously measured at high significance in survey data.

We also verify that several null tests of great importance for cosmological applications (such as B-mode contamination, PSF residual errors and parity-violating contributions) are consistent with zero or otherwise negligible compared to the *E*-mode signal for these 3pt statis-

tics in DES Y3. This work, therefore, represents the first step towards a cosmological analysis with DES Y3 data using the statistics presented here, which we leave for the future.

This paper is structured as follows. In Sec. II we provide an overview of the DES Y3 weak lensing shear catalog and an N-body simulation that we utilize as a check on the rough scale dependence and amplitude of the 3pt signatures. In Sec. III we review the underlying theory of three-point lensing correlations as a probe of the matter bispectrum and describe the estimators we utilize in the data. In Sec. IV we present the main results of this paper: the measured signals of the mass aperture skewness, natural shear correlations, and some explorations of their configuration and redshift dependence, as well as a comparison with existing detections. In Sec. V we validate the measured signals and verify that their origin must be astrophysical and gravitational by checking that *B*-mode, PSF and parity-violating contaminations are negligible and that our data estimator is robust. We conclude and mention future avenues and challenges in Sec. VII.

II. DATA

We describe below the data utilized in this work, the DES Y3 shape catalog and a simulated (N-body) mock. We regard the latter as providing a simplified theory estimate, serving as a basic check of the data measurement.

A. DES Y3 Data

The first 3 years of data from the Dark Energy Survey (DES Y3) cover the full footprint of the survey’s six-year campaign. Its nominal area is over 5,000 deg², which is reduced to 4143 deg² after data selections and cuts that optimize the observed samples for weak lensing and galaxy clustering measurements, with a **baseline mask described in Sevilla-Noarbe et al. (2021)**. The DES data were collected using the 570 megapixel Dark Energy Camera (DECam; Flaugher et al. (2015)) in five photometric bands *grizY* at the Blanco telescope at Cerro Tololo Inter-American Observatory (CTIO) in Chile.

Here we are interested in the METACALIBRATION (Huff & Mandelbaum 2017, Sheldon & Huff 2017) shape catalog produced and validated in the DES Y3 analysis (Gatti & Sheldon et al., (2021c)). This is the largest shear catalog to date in number of objects and area, with over 100 million objects with a mean redshift of $z = 0.63$ and a weighted source number density $n_{\text{eff}} = 5.59 \text{ arcmin}^{-2}$. An overview of the DES Y3 weak lensing and galaxy clustering cosmological analysis is available in DES Collaboration (2022), where further specifications of the data and analysis tools are available in references contained within.

In the DES Y3 cosmological analysis, source galaxies were separated into four redshift bins each with approximately equal numbers of galaxies (Myles & Alarcon et al., (2021)). In some of the measurements presented in this work, we also separate the shear data into tomographic bins. However, since the 3pt statistics have lower signal-to-noise than the 2pt measurements, we instead divide the DES Y3 METACALIBRATION catalog into just 2 redshift bins, which we label z_1 and z_2 . The lower redshift bin, z_1 , is a combination of the galaxies assigned to bins 1 and 2 in the fiducial analysis, while bin z_2 is a combination of the galaxies originally assigned to bins 3 and 4 in that analysis. Weighting the galaxy redshifts in these two newly defined bins by their inverse-variance ellipticity and shear response, we obtain mean redshifts $\langle z_1 \rangle = 0.42$ and $\langle z_2 \rangle = 0.81$ with widths of 0.30 and 0.27 respectively.

Since the shape catalog used to derive the cosmic shear results in DES Y3 (Amon et al. (2021), Secco & Samuroff et al., (2021)) has been extensively validated, we use the same data quality cuts and sample specification in the 3pt analysis below.

B. T17 Mock Catalog

To support our findings reported in the following sections, the same 3rd order correlation measurement pipelines applied to DES Y3 data are also applied to an N-body mock galaxy catalog based on Takahashi et al. (2017) [hereafter T17].

We use full-sky lensing convergence and shear maps from T17 to create a DES Y3-like, tomographic shape catalog. In particular, we used a single one out of their 108 available sets of convergence and shear map snapshots, which span a redshift range between $z = 0.05$ and 5.3 at intervals of $150 h^{-1}$ Mpc comoving distance. The maps have been obtained via ray-tracing using the algorithm GRAYTRIX (Hamana et al. 2015), based on the output of different N-body simulations. The N-body simulations have been run using the code L-GADGET-2 (Springel 2005), assuming a flat Λ CDM WMAP 9 cosmology (Hinshaw et al. 2013) with parameters given by $(\sigma_8, n_s, h, \Omega_m, \Omega_b) = (0.82, 0.97, 0.7, 0.279, 0.046)$.

The shear and convergence maps come in the form of HEALPIX¹ (Górski et al. 2005, Zonca et al. 2019) maps with resolution NSIDE = 4096. We first produced shear maps for each of the tomographic bins by averaging the shear snapshots weighted by the redshift distributions of the bins. To this aim, we used the approximate DES Y3 redshift distributions (Myles & Alarcon et al., (2021)). Galaxy catalogs are then created by sampling the simulated shear maps at the positions of real DES Y3 galaxies, matching their number density. While, in principle,

shape noise can be added to the mock in order to closely match the real data specifications, we do not include it in our mock and instead regard simulation measurements as simple theory estimates.

III. THREE-POINT SHEAR CORRELATIONS

We now describe the basic theory of the higher order correlations we are interested in, the estimator methods that are applied to the simulated and observed data described in the previous section, as well as data covariance matrix estimates based on jackknife.

A. Theory Basics

Second order statistics (two-point correlation functions, power spectra, second moments etc.) contain only the Gaussian part of the shear field. To probe non-Gaussian information, one has to appeal to higher-order statistics. We focus here on lensing 3rd order correlations. A fundamental aspect of these correlations is that they are projections of the matter bispectrum under some lensing kernel, so we take that as our starting point.

We first define the matter bispectrum $B_\delta(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$, that is, the Fourier transform of 3-point correlations of matter overdensities $\delta(\mathbf{k})$ in wavenumbers \mathbf{k} :

$$\langle \delta(\mathbf{k}_1) \delta(\mathbf{k}_2) \delta(\mathbf{k}_3) \rangle = B_\delta(k_1, k_2, k_3) \delta_D(\mathbf{k}_1 + \mathbf{k}_2 + \mathbf{k}_3), \quad (1)$$

where the Dirac delta δ_D enforces the bispectrum definition over wavenumbers \mathbf{k}_i forming triangles, though with statistical isotropy the dependence is only on the magnitude of the modes k_1 , k_2 and k_3 of the triangle. The matter fluctuations give rise to a lensing signal that depends on the redshift distribution of the sources along a unit line-of-sight $\hat{\mathbf{n}}$. This is quantified in real space by the lensing convergence $\kappa(\hat{\mathbf{n}})$:

$$\kappa(\hat{\mathbf{n}}) = \int_0^\infty dz W(\chi) \delta(\hat{\mathbf{n}}, \chi), \quad (2)$$

where $\chi = \chi(z)$ is the comoving distance to redshift z and the lensing efficiency along the line-of-sight is

$$W(\chi) = \frac{3\Omega_m H_0^2}{2c^2} \frac{\chi}{a(\chi)} \int_\chi^\infty d\chi' n(z(\chi')) \frac{dz}{d\chi'} \frac{\chi' - \chi}{\chi'}, \quad (3)$$

where Ω_m is the matter density at redshift $z = 0$, $H_0 = 100h$ km/s/Mpc is the Hubble parameter, a is the scale factor, $n(z)$ is the normalized redshift distribution of sources, and c is the speed of light. Under this lensing kernel, the 3-dimensional matter bispectrum in eq. (1) can be projected down to the 2-dimensional harmonic space convergence bispectrum using the Limber approximation (Limber 1953, LoVerde & Afshordi 2008):

$$B_\kappa(\ell_1, \ell_2, \ell_3) = \int_0^\infty d\chi \frac{W(\chi)^3}{\chi^4} B_\delta(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3; \chi) \quad (4)$$

¹ <http://healpix.sf.net>

With a weak lensing survey, we can probe the shear field at the positions of source galaxies and quantify its statistics with the lensing bispectrum above. We can fine the spin-2 shear field along some direction (e.g., a line connecting two source galaxies) as $\gamma(\boldsymbol{\theta}) = \gamma_t(\boldsymbol{\theta}) + i\gamma_\times(\boldsymbol{\theta})$ where γ_t is the shear component oriented perpendicularly with respect to that direction, γ_\times is the 45° orientation, and $\boldsymbol{\theta}$ are vectors on the plane of the sky with magnitude θ . A natural choice for two-point correlations of the shear field is to take the direction $\boldsymbol{\theta}$ to be that of the line separating a pair of source galaxies, in which case these correlations are given by

$$\xi_\pm(\boldsymbol{\theta}) = \langle \gamma_t \gamma_t \rangle(\boldsymbol{\theta}) \pm \langle \gamma_\times \gamma_\times \rangle(\boldsymbol{\theta}) \equiv \gamma_{tt} \pm \gamma_{\times\times}, \quad (5)$$

with the angle brackets denoting averages taken over all possible pairs of galaxies, and where the right-hand equivalence introduces a shorthand notation for the multiplication of shears.

While the choice for an orientation of shear projections in the three-point case is less obvious (e.g., the orthocenter of the triangle, or the side directions, etc.), there are “natural components” of cosmic shear with rotation and invariance properties analogous to ξ_\pm that we can utilize (Schneider & Lombardi 2003) [hereafter SL03]. We follow SL03 and define:

$$\Gamma_0 \equiv \langle \gamma(\boldsymbol{\theta}_1) \gamma(\boldsymbol{\theta}_2) \gamma(\boldsymbol{\theta}_3) \rangle = \gamma_{ttt} - \gamma_{t\times\times} - \gamma_{\times t\times} - \gamma_{\times\times t} + i[\gamma_{tt\times} + \gamma_{t\times t} + \gamma_{\times tt} - \gamma_{\times\times\times}], \quad (6)$$

$$\Gamma_1 \equiv \langle \gamma^*(\boldsymbol{\theta}_1) \gamma(\boldsymbol{\theta}_2) \gamma(\boldsymbol{\theta}_3) \rangle = \gamma_{ttt} - \gamma_{t\times\times} + \gamma_{\times t\times} + \gamma_{\times\times t} + i[\gamma_{tt\times} + \gamma_{t\times t} - \gamma_{\times tt} + \gamma_{\times\times\times}], \quad (7)$$

$$\Gamma_2 \equiv \langle \gamma(\boldsymbol{\theta}_1) \gamma^*(\boldsymbol{\theta}_2) \gamma(\boldsymbol{\theta}_3) \rangle = \gamma_{ttt} + \gamma_{t\times\times} - \gamma_{\times t\times} + \gamma_{\times\times t} + i[\gamma_{tt\times} - \gamma_{t\times t} + \gamma_{\times tt} + \gamma_{\times\times\times}], \quad (8)$$

$$\Gamma_3 \equiv \langle \gamma(\boldsymbol{\theta}_1) \gamma(\boldsymbol{\theta}_2) \gamma^*(\boldsymbol{\theta}_3) \rangle = \gamma_{ttt} + \gamma_{t\times\times} + \gamma_{\times t\times} - \gamma_{\times\times t} + i[-\gamma_{tt\times} + \gamma_{t\times t} + \gamma_{\times tt} + \gamma_{\times\times\times}]. \quad (9)$$

It has been shown by SL03 as well as by Schneider et al. (2002), Takada & Jain (2003a) that, for general triangle configurations, all of the correlations above can be non-zero and their imaginary parts do not necessarily vanish. Parity invariance, however, implies that the Γ_i for equilateral configurations are purely real (all terms with an odd number of \times -components vanish) and that some, but not all, imaginary components of these statistics for isosceles configurations vanish. The correlations above thus have a complex configuration dependence and can be divided into a total of 8 data vectors (the real and imaginary part of each Γ_i), and should contain the entire 3pt information in the shear field.

The Γ_i are connected to the convergence bispectrum in eq. (4) since, in harmonic space, the shear components can be written in terms of the convergence

$\gamma(\boldsymbol{\ell}) = e^{2i\beta} \kappa(\boldsymbol{\ell})$, where β is the polar angle of $\boldsymbol{\ell}$. The exact expressions for each Γ_i in terms of the convergence bispectrum is worked out in detail in Schneider et al. (2005); for brevity, we simply quote their result for Γ_0 in simplified notation:

$$\Gamma_0(\theta_1, \theta_2, \theta_3) = (2\pi) \int_0^\infty \frac{\ell_1 d\ell_1}{(2\pi)^2} \int_0^\infty \frac{\ell_2 d\ell_2}{(2\pi)^2} \times \int_0^{2\pi} d\phi B_\kappa(\ell_1, \ell_2, \phi) \sum_{j=1}^3 e^{i\alpha_j} J_6(A_j), \quad (10)$$

where J_6 is the 6-th order Bessel function of the first kind, and $B_\kappa = B_\kappa(\ell_1, \ell_2, \phi)$ due to statistical isotropy, with ϕ the polar angle between ℓ_1 and ℓ_2 . We refer readers to Schneider et al. (2005) for the definitions of the coefficients α_j and $A_j = A_j(\theta_1, \theta_2, \theta_3)$ (see their eq. 15).

The shear field can also be decomposed into a different pair of statistics: the mass aperture statistic M_{ap} and its cross-component M_\times (Crittenden et al. 2002, Schneider et al. 1998). The mass aperture term is generally defined as a filtered version of the convergence κ :

$$M_{\text{ap}}(\boldsymbol{\theta}) = \int d^2\mathbf{r} U_\theta(r) \kappa(\mathbf{r}), \quad (11)$$

and we can also introduce it in terms of the tangential shear in circular apertures plus a cross-component shear term (expected to be null for an E -mode field) as:

$$M(\boldsymbol{\theta}) = M_{\text{ap}}(\boldsymbol{\theta}) + iM_\times(\boldsymbol{\theta}) = \int d^2\mathbf{r} Q_\theta(r) \gamma_t(\mathbf{r}) + i \int d^2\mathbf{r} Q_\theta(r) \gamma_\times(\mathbf{r}), \quad (12)$$

where again θ is the magnitude of a planar vector (an “aperture radius” over which the integrals above are computed), and \mathbf{r} is a vector on the plane of the sky.

There is some freedom in defining the filter functions $U_\theta(r)$ and $Q_\theta(r)$, but in this work we stick to the form proposed by Crittenden et al. (2002):

$$U_\theta(r) = \frac{1}{2\pi\theta^2} \left(1 - \frac{r^2}{2\theta^2} \right) \exp\left(-\frac{r^2}{2\theta^2} \right), \quad (13)$$

$$Q_\theta(r) = -U_\theta(r) + \frac{2}{r^2} \int_0^r r' dr' U_\theta(r') \quad (14)$$

$$= \frac{r^2}{4\pi\theta^4} \exp\left(-\frac{r^2}{2\theta^2} \right), \quad (15)$$

for an aperture of radius θ . The statistics defined by eqs. (12)–(15) have several interesting properties which have been explored in the literature (Crittenden et al. 2002, Kilbinger & Schneider 2005, Schneider et al. 2005). In particular, M_{ap} and M_\times cleanly separate, respectively, E - and B -modes of the shear field (Shi et al. 2014) and offer

a relatively compact weighting over angular scales (note that the filter $Q_\theta(r)$ can be significantly non-zero for radii r up to a factor of a few larger than the nominal aperture θ , a feature we will come back to later). Additionally, these forms are mathematically tractable as they mainly involve Gaussian integrals. The ease of integration means that the connection between the third-order correlation of the mass aperture and the bispectrum is straightforward. Again following [Schneider et al. \(2005\)](#), we have:

$$\langle M_{\text{ap}}^3 \rangle(\theta_1, \theta_2, \theta_3) = \frac{3}{(2\pi)^3} \int_0^\infty \ell_1 d\ell_1 \int_0^\infty \ell_2 d\ell_2 \int_0^{2\pi} d\phi \times B_\kappa(\ell_1, \ell_2, \phi) \tilde{U}(\theta_1 \ell_1) \tilde{U}(\theta_2 \ell_2) \tilde{U}(\theta_3 \ell'), \quad (16)$$

where $\tilde{U}(x) = (x^2/2)e^{-x^2/2}$ is the Fourier transform of the filter $U_\theta(r)$ in eq. (13) and $\ell' = \sqrt{\ell_1^2 + \ell_2^2 + 2\ell_1\ell_2 \cos\phi}$. The relatively compact weighting over ℓ multipoles provided by the filter and the absence of fast oscillatory functions in eq. (16) compared to eq. (10) make it a computationally tractable tool for astrophysical predictions leading to cosmology, and indeed it has been a preferred statistic in the literature for cosmological constraints employing real space shear correlations (Fu et al. 2014, Jarvis et al. 2004, Semboloni et al. 2011). As a data vector, $\langle M_{\text{ap}}^3 \rangle(\theta_1, \theta_2, \theta_3)$ is easily tractable because it contains all three-point E -mode information in the field over all triangle configurations, as opposed to the complex splitting of the signal across the 8 non-zero $\Gamma_i(\theta_1, \theta_2, \theta_3)$'s. We will also obtain measurements in the special case $\theta_1 = \theta_2 = \theta_3 = \theta$ so that $\langle M_{\text{ap}}^3 \rangle = \langle M_{\text{ap}}^3 \rangle(\theta)$, which means all aperture radii are the same (though still accounting for different triangle configurations inside the apertures, not to be confused with a strict equilateral assumption). A schematic example of the angle variables used above and in Sec. III B below is shown in Fig. 8 (Appendix B).

It is interesting to consider, additionally, that as structure in the universe becomes more non-Gaussian at lower redshifts, the third order moments of the 3-dimensional density field should increase towards $z \rightarrow 0$. For lensing fields, projection along the line of sight must also be included, and the evolution of non-Gaussian features is quantified via the reduced skewness $S(\theta; z)$ (Schneider et al. 1998), showing the amplitude of the third moment relative to the second moment:

$$S(\theta; z) = \frac{\langle M_{\text{ap}}^3(z) \rangle}{\langle M_{\text{ap}}^2(z) \rangle^2}(\theta), \quad (17)$$

which is tightly related to the usual definition of the reduced bispectrum in terms of $B(k_1, k_2, k_3)/[P(k_1)P(k_2) + \text{perm.}]$ (Cooray & Sheth 2002). This ratio encapsulates the contribution of non-Gaussian statistics to our low-redshift lensing data, arising predominantly from nonlinear structure formation at the scales considered in this work.

B. Estimating Γ_i and $\langle M_{\text{ap}}^3 \rangle$

Motivated by the connection between theory and observables in Sec. III A above, we now turn to the main objective of this work: to obtain and validate a measurement of shear correlations Γ_i and $\langle M_{\text{ap}}^3 \rangle$.

Our starting point is to measure the Γ_i 's. Their most straightforward data estimator is not conceptually different from estimating the usual 2pt statistics $\xi_\pm(\theta)$ in eq. (5). It relies on counting triplets (or pairs in the 2pt case) of galaxies in the survey, and accumulating the product of their shears in tangential and crossed orientations. So, for a catalog with ellipticities $\mathbf{e} = e_t + ie_x$ with per-galaxy weights w , the estimator $\hat{\Gamma}_0$, for example, is

$$\hat{\Gamma}_0 = \frac{\sum_{ijk} w_i w_j w_k \mathbf{e}_i \mathbf{e}_j \mathbf{e}_k}{\sum_{ijk} w_i w_j w_k}, \quad (18)$$

where the sum (ijk) runs over all galaxy triplets. In DES Y3, the weighting w is given by the inverse variance of the ellipticity estimates in METACALIBRATION (see Gatti & Sheldon et al., (2021c) Sec. 4.3), and the ellipticities \mathbf{e} are mean-subtracted and divided by the combination of shear and selection responses². Similar to the two-point ξ_\pm case, this estimator is largely unaffected by masking and geometry of the survey.

For the other statistic, $\langle M_{\text{ap}}^3 \rangle$, there are at least two conceptually different estimators. One relies on sampling apertures over the survey footprint and averaging over the tangential and cross components, directly probing integrals on the right-hand side of equation (12) as proposed by Schneider et al. (1998). One of the main benefits of this method is that the estimation runtime can be made very fast (Porth et al. 2020), and consequently it becomes feasible to obtain empirical survey covariance matrices of nearly arbitrary order in the mass aperture moments (Porth & Smith 2021). A potential drawback of this estimator, however, is that survey masks, holes, edges and other common observational issues in real data can potentially bias the mass aperture estimate.

A second method, which is our favored choice for the present work and was originally proposed by Schneider et al. (2002) and Crittenden et al. (2002), relies on estimating the n -point statistics of the aperture mass by integrating over the n -point shear correlations themselves, as estimated from data. It was shown by Jarvis et al. (2004) that, by assuming the filtering function of Crittenden et al. (2002), one obtains concise expressions for the $M(\theta)$ integration:

$$\langle M^3 \rangle(\theta) = \int \frac{s ds}{\theta^2} \int \frac{d^2 \mathbf{t}'}{2\pi\theta^2} \Gamma_0(s, \mathbf{t}') T_0\left(\frac{s}{\theta}, \frac{\mathbf{t}'}{\theta}\right), \quad (19)$$

² Example usage of the DES Y3 shear catalogs is provided in <https://github.com/des-science/DESY3Cats/>

$$\langle M^2 M^* \rangle(\theta) = \int \frac{s ds}{\theta^2} \int \frac{d^2 \mathbf{t}'}{2\pi\theta^2} \Gamma_1(s, \mathbf{t}') T_1 \left(\frac{s}{\theta}, \frac{\mathbf{t}'}{\theta} \right), \quad (20)$$

where we have used the special case $\theta = \theta_1 = \theta_2 = \theta_3$, where s and \mathbf{t}' are triangle sides as defined in eq.(B1), and the functions T_0 and T_1 are defined in eqs.(B2) and (B3) (see Appendix B). The separate tangential and cross components $\langle M_{\text{ap}}^3 \rangle$ and $\langle M_{\times}^3 \rangle$ can be written as linear combinations of the $\langle M^3 \rangle$ and $\langle M^2 M^* \rangle$ defined above. In particular, with \mathcal{R} denoting the real part of an imaginary quantity, we have

$$\langle M_{\text{ap}}^3 \rangle(\theta) = \frac{1}{4} \mathcal{R} [3 \langle M^2 M^* \rangle(\theta) + \langle M^3 \rangle(\theta)]. \quad (21)$$

We utilize TREECORR (Jarvis et al. 2004) in order to estimate the quantities in eqs. (18)-(21) above. TREECORR is an efficient tree-based algorithm for computing 2pt and 3pt correlation functions in real space data. The estimator follows closely equations (18), (19) and (20) in the sense that galaxy shears are first aggregated by their triangle configuration and side lengths, and in a post-processing step the Γ_i are integrated over with the $T_{0,1}$ functions to obtain $M_{\text{ap}/\times}$. The base algorithm itself is the same utilized for correlation function measurements in the two-point DES Y3 cosmology results (DES Collaboration 2022). We refer the reader to the source code and documentation webpage for more information³.

Even with a highly efficient tree algorithm, we find that runtime is a limiting factor when computing 3pt correlations of the spin-2 shear fields in our data (see Sec. VA further below). Therefore, for all measurements presented in this work, we divide the survey (and simulation) footprints into 100 patches of nearly equal number of galaxies. With DES Y3 data, each patch contains about $N = 1\text{M}$ galaxies. The main advantage of this approach is to significantly reduce the number of galaxies dealt with in each measurement and to better parallelize it.

We define the patch centers and assign galaxies to them using the k -means implementation in TREECORR, which yields patches of roughly similar area $\gtrsim 40 \text{ deg}^2$ (a characteristic length $\gtrsim 6 \text{ deg}$). This choice is sub-optimal, because measuring correlations in finite patches of an otherwise contiguous area necessarily neglects the signal contributions coming from triangles formed by galaxies that lie in different patches. However, since the area of the DES Y3 footprint is large compared to the relatively small angular scales over which we present our measurements in Sec. IV, this is not a significant issue. We return to this and other estimator tests in Sec. VA.

Due to the angular binning performed by TREECORR, for triangles of side lengths $d_3 \leq d_2 \leq d_1$, we define, more conveniently

$$\theta_{\text{medium}} = d_2 \quad (22)$$

as a proxy to index the Γ_i data vector, and unless explicitly noted otherwise we average over all triangles that fall within a bin around θ_{medium} . TREECORR uses internal variables u and v (defined in eqs. B4 and B5) that characterize triangles by their configuration (eg. squeezed or equilateral). We then estimate the mean 3pt signals for each natural component i of $\Gamma_i(\theta_{\text{medium}})$ via the weighted sample mean over the patches α , with $\alpha \in [1, 100]$:

$$\Gamma_i(\theta_{\text{medium}}) = \frac{\sum_{\alpha} \sum_{uv} (1/\text{Var}[\Gamma_{i,\alpha}]) \Gamma_{i,\alpha}(\theta_{\text{medium}}, u, v)}{\sum_{\alpha} \sum_{uv} (1/\text{Var}[\Gamma_{i,\alpha}])}, \quad (23)$$

where inverse-variance weights are estimated in the shape noise regime (more details in Sec. III C). Analogously, we compute the skewness of the mass aperture in each patch using eq. (21) and then combine them so the mean signal is

$$\langle M_{\text{ap}}^3 \rangle(\theta_1, \theta_2, \theta_3) = \frac{\sum_{\alpha} (1/\text{Var}[M_{\text{ap}}^3]) \langle M_{\text{ap}}^3 \rangle_{\alpha}}{\sum_{\alpha} (1/\text{Var}[M_{\text{ap}}^3])}. \quad (24)$$

C. Covariance Matrix

With the computation of the measurement over $N = 100$ patches of the DES Y3 data, we can readily obtain a jackknife estimate of the covariance matrix:

$$\text{Cov}[\zeta(\theta_i), \zeta(\theta_j)] = \frac{N-1}{N} \sum_{\alpha} \Delta\zeta_{\alpha}(\theta_i) \Delta\zeta_{\alpha}(\theta_j)^T \quad (25)$$

where ζ is the data vector of the statistic under consideration ($\langle M_{\text{ap}}^3 \rangle(\theta)$ or $\Gamma_i(\theta_{\text{medium}})$ for instance), $\langle \zeta \rangle$ is its average value over the N patches, and $\Delta\zeta_{\alpha} \equiv \zeta_{\alpha} - \langle \zeta \rangle$. When inverting the covariance matrix, we also apply a ‘‘Hartlap correction’’ factor (Dodelson & Schneider 2013, Hartlap et al. 2007, Sellentin & Heavens 2016, Taylor et al. 2013) given by $(P - N - 1)/(N - 2)$ where P is the dimension of the data vector and N is the number of patches ($P = 7$ and 55 for $\langle M_{\text{ap}}^3 \rangle(\theta)$ and $\Gamma_i(\theta_{\text{medium}})$ respectively, and $N = 100$ in both cases).

In Fig. 1, we show the normalized covariance matrices (correlation matrices) for the $\langle M_{\text{ap}}^3 \rangle(\theta)$ and $\Gamma_0(\theta_{\text{medium}})$ estimates which we present in the following Section. We additionally show, on the bottom panel of that Figure, how the diagonal $\langle M_{\text{ap}}^3 \rangle$ standard deviation compares with empirical and analytic estimates of the error in the shape noise dominated regime. We obtain an empirical estimate of the shape noise signal (light blue curve in Fig. 1) by repeating the $\langle M_{\text{ap}}^3 \rangle$ measurement over patches in which each individual galaxy shear has been randomly rotated. This effectively cancels out the cosmic signal and variance, leaving us with an estimate of the shape noise

³ <https://github.com/rmjarvis/TreeCorr>




FIG. 1. Normalized covariance matrix estimates for $\langle M_{\text{ap}}^3 \rangle$ (top panel), Γ_0 (middle panel) and $\langle M_{\text{ap}}^3 \rangle$ standard deviation (bottom panel). With the exception of the analytic shape noise estimate (dashed line in bottom panel), all other estimates are obtained from the jackknife measurements on DES Y3 data. We note that in most scales below around 20 arcmin, the shape noise contribution to the error bars is of around 50% or more, and at larger scales the errors are dominated by cosmic variance.

that preserves any masking or geometry effects of the real data. We additionally overplot in that same panel an analytic estimate of shape noise. The analytic estimate comes from the propagation of the weighted variance of Γ_i into $\langle M_{\text{ap}}^3 \rangle$, which in turn can be written as

$$\text{Var}[\mathcal{R}\{\Gamma\}] = 4\sigma_e^6 \frac{\sum_{ijk} w_i^2 w_j^2 w_k^2}{\left(\sum_{ijk} w_i w_j w_k\right)^2} \quad (26)$$

where w are weights associated to the data ellipticities, $\sigma_e^2 = \langle (e_i - \langle e_i \rangle)^2 \rangle$ is the variance of single-component ellipticities, and the sums (ijk) run over all possible triplets of galaxies. We note that this reduces to $\text{Var}[\mathcal{R}\{\Gamma\}] = 4\sigma_e^6/N_\Delta$ for equal galaxy weighting, where N_Δ is the number of triangles in a given angular bin. We find that, for values of θ less than ~ 20 arcmin, the shape noise contributes $> 50\%$ of the estimated error bars in $\langle M_{\text{ap}}^3 \rangle$.

While jackknife covariances are known to be biased on scales that approach the characteristic length of an individual patch, the covariances we utilize should be reliable for the simple S/N estimates at the relatively smaller angular scales studied in this work. Survey data covariances are generally difficult to obtain and can directly impact likelihood analyses, especially at the 3pt level (Joachimi et al. 2009, Sato & Nishimichi 2013). We therefore intend to further study the suitability of our existing jackknife matrices in a follow-up work focusing on the inference of cosmology constraints.

IV. MEASUREMENT RESULTS

We now apply the estimators defined in Sec. III B to the DES Y3 data split into 100 patches. We measure the 3pt correlations Γ_i within an angular range of $\theta_{\text{medium}} \in [1, 240]$ arcmin, approximately the same range of scales validated in DES Y3 for weak lensing applications. For Γ_i , angular bins in θ_{medium} are log-spaced (with 0.1 spacing, leading to 55 bins) and TREECORR's internal variables u and v are linearly-spaced (0.1 spacing, leading to respectively 10 and 20 bins; see Appendix B) to ensure stability of the integrals that lead to M_{ap} moments. When plotting Γ_i results and obtaining its covariance, we further average over every 5 bins in θ_{medium} for ease of visualization and to reduce noise. For the results on the M_{ap} estimation, however, we focus on a narrower range of scales and limit aperture radii to the interval $\theta \in [4, 60]$ arcmin in 7 bins, avoiding measurement biases that can arise if the aperture filtering in eq. (15) spans scales over which the Γ_i were not obtained (further details in Sec. V A).

We present the non-tomographic signal in Sec. IV A, along with splits of triangles by configuration type, and then we divide our data into two tomographic bins in Sec. IV B. In what follows, we define the signal-to-noise (S/N) of our detections as (see Appendix C, where this




FIG. 2. The non-tomographic 3rd order natural shear correlations Γ_i in DES Y3 as a function of angular scale in arcminutes of the medium-length side of triangles, θ_{medium} (eq. 22). Solid lines correspond to averaged measurements over 100 patches of the DES Y3 footprint; error bars are estimated with a jackknife method. To guide the eye, the dashed lines show corresponding measurements on a T17 N-body mock catalog (Sec. II B) with DES Y3-like redshift distributions but without shape noise and for an older set of cosmological parameters. Signal-to-noise (S/N) estimates are obtained with eq.(27). **Left panel:** Real parts of the natural 3pt shear components. The lensing signal is distributed rather evenly across the 4 components, and for most of them the null-hypothesis is clearly rejected at high confidence. **Right panel:** Imaginary parts of the natural 3pt shear components, which are expected to be zero for certain triangle configurations (e.g., equilateral) but not in general, thus leading to smaller overall S/N.

is derived)

$$S/N \equiv \begin{cases} \sqrt{\chi^2 - N_{\text{d.o.f}}} & \text{if } \chi^2 \geq N_{\text{d.o.f}} + 1 \\ \text{"Null"} & \text{otherwise} \end{cases}, \quad (27)$$

where $N_{\text{d.o.f}}$ is degrees of freedom (here the number of data points) and $\chi^2 = \mathbf{d}^T \mathbf{C}^{-1} \mathbf{d}$ with \mathbf{d} representing the measurement vector and \mathbf{C}^{-1} representing the inverse data covariance. In the low signal-to-noise regime (which is the case for many of the null tests presented later), it may be that $\chi^2 < N_{\text{d.o.f}} + 1$, in which case S/N is less than 1.0 or imaginary, which we consider a “Null” signal (consistent with no detection). Additionally, for practical purposes, we define a data vector to be *significantly* rejecting the null-hypothesis (at $X\sigma$) if $S/N = X > 2.5$, which as an equivalent p -value yields $p \lesssim 0.01$.

A. Non-tomographic 3pt Shear Signal

We first focus on the non-tomographic setting, treating all galaxies in the survey as if their line-of-sight distances belonged to a thin plane on the sky. We show the real and imaginary parts of the non-tomographic Γ_i and their S/N in Fig. 2. We report significant detections (ruling out

the null-hypothesis at 2.5σ or more) of the real parts of all natural shear components $\Gamma_i(\theta_{\text{medium}})$, and an overall lower significance for their imaginary parts. This is expected since, in specific triangle configurations, but not generally all of them, the imaginary parts vanish due to parity conservation. We overplot measurements obtained from the T17 N-body mock with dashed lines as a guide to the eye, though it should not be expected that these curves serve as a fit to the data, which we return to below.

We additionally report a strong detection (ruling out the null hypothesis at more than 11σ) of the non-tomographic lensing E -mode term $\langle M_{\text{ap}}^3 \rangle$ in the left panel of Fig. 3, in the special case of a single aperture radius $\theta_1 = \theta_2 = \theta_3$. The higher S/N of the mass aperture in comparison with individual Γ_i 's is in principle expected: the tangential projection of shears for a given triangle configuration contains a large fraction of the signal (Takada & Jain 2003b) and the $\langle M_{\text{ap}}^3 \rangle$ statistic sums over that projection across many configurations in an aperture θ , while Γ_i splits the contribution over a total of 8 independent correlations γ_{abc} with $a, b, c \in [t, \times]$.

We find that the overall amplitude of the simulated and data signals in both Γ_i and $\langle M_{\text{ap}}^3 \rangle$ closely resemble each other. A more careful assessment beyond the scope of this work would be necessary to verify whether discrepancies




FIG. 3. The 3rd order mass aperture correlations in DES Y3 for the special case of a single aperture radius ($\theta_1 = \theta_2 = \theta_3 = \theta$) in arcminutes. Solid lines correspond to averaged measurements over 100 patches of the DES Y3 footprint, and error bars are estimated with a jackknife method. As in Fig. 2, dashed lines show measurements for the T17 N-body mock (Sec. II B), and S/N estimates are obtained with eq.(27). **Left panel:** non-tomographic M_{ap} and M_{\times} moments. We find a significant detection of the pure E -mode term $\langle M_{\text{ap}}^3 \rangle$, and find the other combinations, which either violate parity or imply significant B -mode contamination, to be consistent with the null-hypothesis. **Right panel:** Tomographic mass aperture cross-correlations using DES Y3 data split into 2 wide redshift bins z_1 and z_2 . We find a significant detection of the cross-correlations that include the higher redshift bin z_2 , and the total combined data vector is detected with 15.0σ significance.

669 between solid and dashed lines in Fig 3 imply our data 695
 670 are statistically rejecting the cosmology (or gravity-only 696
 671 implementation) of the T17 simulations. 697

672 Several known factors could result in these differences: 698
 673 the difference in assumed cosmology, small scale astro-699
 674 physical systematic effects, and shear calibration. Dis-700
 675 cerning between these factors would entail obtaining 3pt-701
 676 functions in the ensemble of 108 mocks in T17 as op-702
 677 posed to the single shape-noise free mock utilized in this-703
 678 work, a computationally expensive task (see Sec. V A for-704
 679 details on the estimator performance), and carrying out-705
 680 likelihood analyses over scales where the theory model-706
 681 ing is not excessively uncertain. We do note, however, 707
 682 that based on the left panel of Fig. 3 the largest off-708
 683 sets are on small scales (roughly below $10'$) and result 709
 684 in a $\Delta\chi^2 \approx 40$ when comparing data and mock within 710
 685 $\langle M_{\text{ap}}^3 \rangle(\theta < 10')$ for the non-tomographic case. Similarly, 711
 686 the tomographic measurements $\langle M_{\text{ap}}(z_1)M_{\text{ap}}(z_2)^2 \rangle(\theta < 712$
 687 $10')$ and $\langle M_{\text{ap}}(z_1)^2M_{\text{ap}}(z_2) \rangle(\theta < 10')$ over the same 713
 688 scales show a combined $\Delta\chi^2 \approx 30$, so it may be possi-714
 689 ble that the origin of the non-tomographic discrepancy is 715
 690 driven by the redshift cross-correlations. This likely rules 716
 691 out strong baryonic feedback in the data as an explana-717
 692 tion for the discrepancy (as that would also have shown 718
 693 up strongly in the lowest-redshift $\langle M_{\text{ap}}(z_1)^3 \rangle(\theta < 10')$ 719
 694 for most feedback scenarios) as well as significant contri-720

698 butions from shear calibration bias (which would likely 919
 699 have appeared as a scale-independent offset affecting ad- 920
 700 ditionally the auto-redshift correlations). We leave fur- 921
 701 ther detailed explorations for a future work. 922

Comparing the $\langle M_{\text{ap}/\times}^3 \rangle(\theta)$ and $\Gamma_i(\theta_{\text{medium}})$ statistics 923
 presented in Fig. 2 and in the left panel of Fig. 3 we find 924
 that they separate the signal contributions in different 925
 ways. While for general triangle configurations the E 926
 and B mode signals are split rather evenly between the 927
 Γ_i , they are more concentrated in M_{ap} as opposed to M_{\times} . 928
 We will exploit this feature in more detail in Sec. V as 929
 an assessment of systematics. 930

While the mass apertures involve a sum over many 931
 triangles and effectively mix their contributions to the 932
 signal, eqs. (6)-(9) on the other hand suggest that the 933
 natural components Γ_i can be combined to separate spe- 934
 cific triangle configurations and projections. Several tri- 935
 angle geometries were used by Takada & Jain (2003b) to 936
 demonstrate that certain configurations (e.g., equilateral 937
 and isosceles triangles) have vanishing projections due to 938
 parity conservation, and that for general triangle shapes 939
 all 8 possible projections of γ_{abc} with $a, b, c \in [t, \times]$ are 940
 non-zero. 941

We can similarly explore the dependence of the signal 942
 on projection and configuration in our data by construct- 943
 ing γ_{ttt} and $\gamma_{\times\times\times}$, the components with all shears pro- 944




FIG. 4. Three-point shear signals with different triangle configuration and scale dependences. Solid lines correspond to non-tomographic DES Y3 data measurements, and dashed lines are the same measurements made on a T17 N-body mock. **Left panel:** the purely tangential γ_{ttt} component for isosceles triangles as a function of opening angle between sides d_1 and d_2 . An oscillating pattern (see text) can be seen at both the small scales (red) as well as large-scale triangles (blue, multiplied by a factor of 50 for visualization). **Right panel:** purely tangential (γ_{ttt}) and cross-projections ($\gamma_{\times\times\times}$) of equilateral triangles. A detection is clear in the tangential case and, according to expectations the equilateral, odd-parity $\gamma_{\times\times\times}$ is consistent with zero.

jected tangentially and at 45° with respect to the triangle center respectively, via

$$\gamma_{ttt} = \frac{1}{4}\mathcal{R}[\Gamma_0 + \Gamma_1 + \Gamma_2 + \Gamma_3] \quad (28)$$

$$\gamma_{\times\times\times} = \frac{1}{4}\mathcal{I}[-\Gamma_0 + \Gamma_1 + \Gamma_2 + \Gamma_3], \quad (29)$$

where \mathcal{R} and \mathcal{I} correspond to real/imaginary parts. Using the triangle side lengths (d_1, d_2, d_3) we obtain the shear signal for two types of configurations: isosceles triangles ($d_1 \approx d_2 \neq d_3$, with ϕ being the opening angle between d_1 and d_2), and equilateral triangles ($d_1 \approx d_2 \approx d_3$)⁴. Furthermore, we can separate “small scale” isosceles triangles with sides $d_1 \approx d_2$ smaller than 5 arcmin, and “large scale” isosceles with $5 < d_1 \approx d_2 < 60$ arcmin. We show our results in Fig. 4, where again dashed lines correspond to a measurement on a T17 mock. The left panel of the figure shows a characteristic oscillatory dependence on opening angle, somewhat similar to what was predicted for even-parity modes in Takada & Jain (2003b)

using a halo model approach, and in qualitative agreement with the T17 simulation result. The right panel of Fig. 4 shows the tangential and cross components of equilateral triangles as a function of angular separation $\theta_{\text{medium}} = d_2 \approx d_1 \approx d_3$. We find a significant signal in the even-parity γ_{ttt} part, while the parity-violating term $\gamma_{\times\times\times}$ is consistent with zero; both are thus consistent with expectation. While the similarity of our signals with halo model studies such as Takada & Jain (2003b), Zaldarriaga & Scoccimarro (2003) and Ho & White (2004) is visually striking, it is not exact. In particular, we find peaked signals on isosceles opening angles $\phi \rightarrow 0^\circ$ and $\phi \rightarrow 180^\circ$ that do not exactly match the expectation based on either work, but follow closely the T17 result. We believe that a quantitative comparison of these measured signals with theory and the information this could provide on gravity, nonlinear structure evolution and halo shapes certainly merits further exploration.

We further explore the general definition of the mass aperture skewness for three different aperture radii $\langle M_{\text{ap}}^3 \rangle(\theta_1, \theta_2, \theta_3)$ in equation (16), and obtain the signal in some specific setups as shown in Fig. 5. We fix the aperture radius θ_1 at 4, 15 and 60 arcmin, representing roughly the smallest, intermediate and largest scales probed with this statistic, and plot the signal as a function of the two other apertures. We find that the amplitude of the third-order mass aperture tends to be higher as we go to smaller scales. We note also that,

⁴ These relations are only approximate in the data. For these specific configuration tests, we allow for small departures from exact triangle shapes, with side ratios binned with a $\pm 15\%$ tolerance in relative side lengths.




FIG. 5. A detection of the generalized mass aperture skewness, expected to contain the entire 3pt information in the lensing field, with a dependence on three angular separations $(\theta_1, \theta_2, \theta_3)$. We fix three choices of representative scales θ_1 at 4, 15 and 60 arcmin, respectively the top, middle and bottom panels. In all cases, we find a significant detection of the generalized signal.

while the generalized $\langle M_{\text{ap}}^3 \rangle(\theta_1, \theta_2, \theta_3)$ contain the entire E -mode information of the field, they do not necessarily contain the highest signal-to-noise individually, a factor that should be taken into account in a future likelihood inference study. Nevertheless, in all cases we again find a significant detection of this particular lensing signal.

B. Tomography

Motivated by the significant detections obtained in the non-tomographic regime, we proceed to split the DES Y3 catalog into redshift bins and attempt a first tomographic measurement of the third moment of the mass aperture.

We implement the same 2-bin redshift split described in Sec. II A on the T17 mock described in Sec. II B. The original, 4-bin redshift distributions in that mock resemble the actual DES Y3 $n(z)$'s but do not reproduce their substructure exactly, so we expect that 3pt statistics obtained from the mock should provide an approximate expectation for the scale dependence and amplitude of the tomographic signal on the data.

We present our results for the cross-tomographic mass apertures in the right-hand panel of Fig. 3, in qualitative agreement with the T17 result at most scales and redshift bins. We compute the signal-to-noise ratios S/N again using eq. (27), and find significant detections of cross-correlations of $\langle M_{\text{ap}}^3 \rangle$ that include the high-redshift bin z_2 . For the complete data vector built with the 4 concatenated cross-tomographic measurements and including their cross-covariances, we find a total S/N of 15.0σ . Interestingly, this detection is non-zero on angular scales that are relatively large ($\theta \sim 1^\circ$), reaching quasi-linear and linear regimes. This implies that non-Gaussian signals may add significant information to common two-point analyses even if these mostly rely on the linear regime due to conservative scale cuts (see, for instance, Gatti et al. (2021a)).

We note several points related to this tomographic measurement. First, the signal in the higher redshift bin z_2 (red curve in the right panel of Fig. 3) is significantly larger than that for the lower bin z_1 . As with the 2pt shear measurement, this trend can be attributed to the fact that the lensing kernel for the higher redshift bin probes more large-scale structure than the kernel limited to low redshifts. Second, the signal-to-noise of $\langle M_{\text{ap}}(z_1)M_{\text{ap}}(z_2)^2 \rangle$ ($S/N = 13.3$) and $\langle M_{\text{ap}}(z_2)^3 \rangle$ ($S/N = 11.5$) are both higher than the non-tomographic case ($S/N = 11.2$). While this may seem counter-intuitive at first, it is not against expectations: there are many low-redshift galaxy triplets in the non-tomographic sample whose 3pt correlations add significant noise but insignificant signal due to the lack of depth of the lensing kernel in the lowest redshift bin, and the overall S/N goes up once these are removed. Third, it is expected that $\langle M_{\text{ap}}(z_1)M_{\text{ap}}(z_2)^2 \rangle$ should have the highest S/N : for redshift bins with approximately the same number of galaxies, a cross-correlation contains a larger number

of galaxy triplets than any auto-correlation, and additionally shot-noise contributions to the uncertainties are diagonal on the redshift bins.

In addition to the signals presented above, we measure the reduced skewness parameter in eq. (17). We again use TREECORR in order to estimate $\langle M_{\text{ap}}^2 \rangle$ in our data and mocks over the same patches where the 3pt observables were obtained. We show $S(\theta; z)$ in Fig. 6. A significant redshift evolution of the reduced skewness parameter can be seen, with the low- z bin showing more power than the high- z bin. This is in line with our expectation that the shear field should be more Gaussian at higher redshift. This is due to the larger projection distance for high redshift, which means more uncorrelated structure contributes to the lensing and a version of the central limit theorem (considering the accumulated signal as a random walk along the line-of-sight) makes the resulting shear field closer to Gaussian (Bernardeau et al. 1997, Jain & Seljak 1997). Note that there is no such expectation for the 3-dimensional density field, where the skewness is redshift-independent in leading order perturbation theory. The lensing skewness is largely independent of the power spectrum shape and normalization, and its approximate redshift evolution was given by e.g. Bernardeau et al. (1997) who obtained $S \sim z^{-1.35}$. While that scaling depends on the cosmological model and the assumptions on the source redshift distribution, we find it to be in qualitative agreement with our measurement: for a representative scale of $10'$ the ratio $S(\theta = 10'; z_1)/S(\theta = 10'; z_2)$ is about 2 to 3, with the mean of redshift bins z_1 and z_2 being at 0.42 and 0.81 (see Sec. II), roughly following the expected scaling.

V. ASSESSMENT OF SYSTEMATICS

We now turn to the validation of the signal with the aim of showing that the detection is not contaminated by systematics of observational/instrumental origin. The results of the tests detailed below indicate that the significant 3rd order lensing signals found in DES Y3 data are of astrophysical and gravitational origin.

A. Estimator Uncertainties

Potential uncertainties in the estimation of Γ_i 's and $\langle M_{\text{ap}}^3 \rangle$ have three different sources, two of them originating from approximations needed to bring the computational runtime to a reasonable level and one, much easier to mitigate, stemming from the mass aperture filtering. We begin by describing this last one, the filtering feature, which we have essentially mitigated in this work by employing angular scale cuts at the measurement level.

The filter defined in eq. (13) decays quickly as a function of angular separation, and is small (but not negligibly so) at separations of a factor λ of about $3 \times$ wider than the angular bin at which $\langle M_{\text{ap}}^3 \rangle$ is evaluated. This non-




FIG. 6. A measurement of the reduced mass aperture defined in eq. (17). Solid lines and error bars correspond to measurements on the DES Y3 data split into 2 wide redshift bins, and dashed lines correspond to measurements on an N -body mock based on the T17 simulations (see Sec. II B), which are not expected to fit the data, but serve as a guiding comparison. The redshift evolution of the skewness parameter $S(\theta; z)$ indicates, according to expectation, that more non-Gaussian structure contributes to the signal at low- z than at high- z .

localization of the filter implies that, for a measurement of $\langle M_{\text{ap}}^3 \rangle$ at an angular separation $\hat{\theta}$, the integrals over Γ_i have significant support over a range $[\hat{\theta}/\lambda, \hat{\theta}\lambda]$ where λ is a characteristic scale of filter. We employ a factor $\lambda = 4$ after empirically testing estimates of $\langle M_{\text{ap}}^3 \rangle$ over different angular ranges and finding them to stabilize very well at that chosen width. This choice is similar to previous studies (Fu et al. 2014) and justifies our choice of scales of $[1', 240']$ for $\Gamma_i(\theta_{\text{medium}})$ and $[4', 60']$ for the aperture radii in $\langle M_{\text{ap}}^3 \rangle$.

The other two potential sources of estimator uncertainties that we have explored are a decreased binning accuracy w.r.t. analogous calculations of 2pt functions, and the jackknifing method utilized. Binning accuracy in TREECORR is determined by code parameters `binslop` and `binsize`. Larger values of the former allow for larger errors when binning triangles by ratios of their side lengths (see App. B), and larger values of the latter imply coarser binning by triangle configuration. We empirically vary both on a reduced number of data patches to verify their impact on our measurement. First, we find that the recommended value of `binsize`=0.1 is sufficient for the integration over Γ_i and yields a stable $\langle M_{\text{ap}}^3 \rangle$. Second, while runtime increases prohibitively with smaller `binslop`, we find that a value of `binslop`=1.0 makes computing time feasible and does not bias the correlation functions, although it increases the diagonal covariance of the measurement by around 15%. With these choices, we

find that the computing time for 1M objects in 28 2.4GHz CPUs on the Midway2 cluster⁵ is around 300 minutes (still very expensive when compared to a timing of under 20 minutes for the 2pt $\xi_{\pm}(\theta)$ auto-correlation of 25M galaxies in one redshift bin of DES Y3 with `binslop=0.0` which approximates brute-force pair-counting).

Finally, there are the uncertainties associated with the jackknife method, which we employ for more efficient parallelization and to obtain an estimate of the covariance matrix. A source of uncertainty comes from triangles whose corners are not all located on the same patch, cause these triangles are not included in the subsequent calculations. We run a feasibly short test on the 100M-object catalog by focusing only on several angular scales of approximately equilateral triangles and find that splitting the full footprint into disjoint patches misses approximately 10% of the nearly equilateral triangles with a side length of 200 arcmin. The missing triplets enhance the shot noise contribution in those large scales but should not contribute a bias because there is no preferential shear projection that is missed due to the patch splitting.

B. B-modes and Parity

In general, a three-point signature of B -modes of astrophysical origin can come from a limited number of effects. In particular, at the 3pt level the main sources of B -modes are intrinsic alignments (Semboloni et al. 2011, Troxel & Ishak 2012, 2015) and the spatial clustering of source galaxies which are otherwise expected to randomly sample the survey footprint (Schneider et al. 2002). These effects are expected to be small compared to the lensing-induced E -mode signal, so at first a reasonable approach is to consider any significant B -mode detection as pointing to potential data systematics (PSF residuals, for instance).

Within the statistics we explore, the main correlations where B -modes could be searched for are $\langle M_{\text{ap}} M_{\times}^2 \rangle(\theta)$ which would point to B -modes correlated with E -modes. In the non-tomographic case, as shown in the left panel of Fig. 3, we find that the signal-to-noise of $\langle M_{\text{ap}} M_{\times}^2 \rangle(\theta)$ is compatible with the null-hypothesis according to the definition in eq. (27), meaning S/N is lesser than 1 or imaginary. In a similar way, we verify that $\langle M_{\text{ap}}(z_1) M_{\times}^2(z_2) \rangle$ and $\langle M_{\text{ap}}(z_2) M_{\times}^2(z_1) \rangle$, the tomographic versions of the same test which would respectively point to B -modes in the higher(lower) redshift bin correlating with E -modes in the lower(higher) redshift bin, are also consistent with the “Null” condition defined in eq. (27).

Other correlations including odd powers of the B -mode field M_{\times} such as $\langle M_{\text{ap}}^2 M_{\times} \rangle(\theta)$ are expected to vanish due to parity (Schneider 2003). A parity-violating field

would necessarily come from systematics of the data, as no astrophysical source could produce it. We indeed find the parity-violating terms $\langle M_{\text{ap}}^2 M_{\times} \rangle(\theta)$ and $\langle M_{\times}^3 \rangle(\theta)$ presented in the left panel of Fig. 3 to be consistent with the null-hypothesis. Finally, we have also shown in Fig. 4 another parity-violating correlation, $\gamma_{\times \times \times}(\theta)$ for approximately equilateral triangles, which is similarly consistent with zero.

C. PSF Residuals

We follow the approach of Rowe (2010) in order to estimate the contribution of additive PSF modeling errors to our lensing observables. We obtain the mass aperture skewness of the so-called “ ρ -statistics” (see Appendix A), which quantify the residual correlations caused by errors in the PSF modeling and deconvolution, modulated by empirically-obtained coefficients α and β .

We estimate the PSF uncertainty impact via eq. (A7) using a catalog of stars to compare them to the actual data signal. In doing so, we need input values for the coefficients α and β that multiply deconvolution errors and modeling residuals, respectively. We set $\alpha = 0.01$ and $\beta = 2$ as inputs for the additive contaminations, considering the bounds on these parameters presented in Gatti & Sheldon et al., (2021c) (respectively $\alpha = 0.001 \pm 0.005$ and $\beta = 1.09 \pm 0.07$). This choice of input values is a very conservative one, which amplifies the estimated impact of these systematics. As the additive PSF contaminations considered here have their origin in the 1-point ellipticities, we do not expect the values of those coefficients to depend on which statistics are used to measure them (apart from practical aspects such as the signal-to-noise of the chosen statistic). We therefore do not pursue a measurement of α and β based on 3pt observables, and utilize those bounds obtained in Gatti & Sheldon et al., (2021c) based on 1- and 2-point PSF correlations.

Despite the conservative choice in input coefficients, we find additive PSF systematics to be entirely negligible. We show in Fig. 7 a breakdown of the PSF contributions to individual skewness component ($\langle M_{\text{ap}}^3 \rangle$, $\langle M_{\text{ap}}^2 M_{\times} \rangle$, etc) and by PSF correlation type ($\langle e_p^3 \rangle$, $\langle e_p^2 q \rangle$, etc), where e_p is the PSF ellipticity and q the ellipticity residual error after modeling. In all cases, we find the 3rd order moments of PSF uncertainties to be negligible, well below a percent of the E -mode data signal $\langle M_{\text{ap}}^3 \rangle$.

D. Mean Shear and Other Observational Systematics

Several other features of 3pt statistics are also relevant for their robustness against systematics. In particular we consider contributions to the signal arising from a residual mean shear in ellipticities $\langle e_1 \rangle$ and $\langle e_2 \rangle$.

While a mean shear that is coherent across angular scales produces a ξ_{+} signal (eq. 5) at the 2pt level, it

⁵ <https://rcc.uchicago.edu/>

does not produce any signature on the Γ_i . This can easily be demonstrated by considering a constant shear field in cartesian coordinates, $\gamma = \gamma_1 + i\gamma_2 = c_1$, herent across some angular length scale. For 2pt functions ξ_{\pm} we project shears along the direction $\alpha + \psi$ where α is the direction of the line that connects the galaxy pair and ψ is the (random) orientation of the pair with respect to the reference of the cartesian ordinates, so $\gamma \rightarrow \gamma' = \gamma \exp[-2i(\alpha + \psi)]$. Then the natural 2pt functions of the field are $\xi_+ = \langle \gamma' \gamma'^* \rangle = c_1^2$ and $\xi_- = \langle \gamma' \gamma' \rangle = c_1^2 \langle \exp[-4i(\alpha + \psi)] \rangle = 0$ as the averaging is essentially over the multiple random orientations ψ .

For the natural 3pt functions, in comparison, the projection of each of the 3 shear components is along different direction ($\alpha + \psi$, $\beta + \psi$ or $\delta + \psi$) and many reference points are possible - the triangle incenter, the center of the side opposing a given angle, etc (Schneider & Lombardi 2003), with some projections leading to $\alpha + \beta + \delta = 0$. In the same situation of a constant shear in cartesian coordinates we have, for an example case: $\Gamma_0 = \langle c_1^3 \exp[-6i(\alpha + \beta + \delta) - 6i\psi] \rangle = 0$ due to the averaging over ψ , and similarly for all other Γ_i with $i = 1, 2, 3$ defined in eqs. (6)-(9). This insensitivity to an additive mean shear over coherent scales can be useful when compared to 2pt functions because it would not lead to the requirement of an extra correction at the data level as in (Gatti & Sheldon et al., (2021c)), and would potentially minimize the need for corrections due to additive systematics such as presented in Kitching et al. (2021).

We additionally expect that any other observational systematics that arise from statistics that are well described by Gaussian processes should have negligible contributions to 3pt functions. A potential example which we leave for a further exploration is the atmospheric contribution to PSFs. As that is well characterized by Gaussian processes with vanishing odd-order correlations, we expect it to be significantly suppressed in importance when dealing with 3pt shear correlations.

VI. COMPARISON WITH PREVIOUS WORK

Among the several types of 3pt shear statistics presented so far, some had already been detected and explored in the survey science literature while others had not. In what follows, we compare our findings with a number of previous results.

As a starting point, our 3pt S/N can be compared with the 2pt DES Y3 cosmic shear measurements. The null-hypothesis signal-to-noise defined in eq. (27) yields $S/N = 40.2$ for the joint ξ_{\pm}^{ij} data vector (eq. 5) presented in Amon et al. (2021), Secco & Samuroff et al., (2021)

⁶ Note that the definition of S/N utilized in these works is different than the one employed here.

before the removal of relatively small angular scales that are not included in the likelihood due to modeling uncertainties (a total $N_{d.o.f.} = 400$ degrees of freedom). After “fiducial” scale cuts, the DES Y3 cosmic shear data vector has $S/N = 27.5$ ($N_{d.o.f.} = 227$), and after “optimized” scale cuts we obtain $S/N = 30.1$ ($N_{d.o.f.} = 273$).

While the signal-to-noise ratio of our 3pt measurements are smaller than the corresponding 2pt S/N , it is realistic to expect that real-space 3pt shear correlations can tighten posteriors in key cosmology results because parameter degeneracies are different between two- and three-point functions. That is indeed the case with Gatti et al. (2021a), wherein an improvement of $\sim 15\%$ is seen in the lensing amplitude S_8 when combining second and third order moments of the lensing convergence.

Regarding three-point detections of cosmic shear observables, Stage-II surveys presented some of the first results: a first detection was claimed by Bernardeau et al. (2002) in the VIRMOS-DESCART 8.5deg^2 survey (Van Waerbeke et al. 2002), followed by detections of the third moment of the mass aperture by Jarvis et al. (2004) with the CTIO 75deg^2 survey data, Semboloni et al. (2011) with HST COSMOS data (Schrabback et al. 2010, Scoville et al. 2007) and, more recently, Fu et al. (2014) with CFHTLenS data (Erben et al. 2013). These first detections of lensing third moments with signal-to-noise around 3σ advanced the field. Our measurements significantly improve upon those detections and bring them up to S/N of around 15σ , a significance that enables quantitative interpretation.

To the best of our knowledge and at the time of this writing, we have reported in this work the first significant detection of the four natural 3pt cosmic shear components (Fig. 2), the first detection of tomographic 3pt mass aperture signals (right panel of Fig. 3), and the first significant detection of components split by their configuration dependence (Fig. 4). Equally important, our measurement (along with the Gatti et al. (2021a) measurement of the skewness of κ in the same data), extends to large scales approaching 1 degree, where quasi-linear theory is reliable and uncertainties due to baryonic physics can be neglected. Thus it will enable robust interpretations of cosmology and gravitational physics.

VII. CONCLUSIONS AND OUTLOOK

Using over 100M galaxies spread across the $4,143\text{deg}^2$ footprint of the first 3 years of data from the Dark Energy Survey, we presented measurements of the three-point correlations of the lensing shear field. We also combined all three point correlations into the third moment of the mass aperture statistic and verified that systematics of observational origin are negligible in our measurements. We expect this work to be a stepping stone for future applications of these 3pt statistics, in particular a joint 2pt and 3pt cosmology analyses. Our main results are summarized below:




FIG. 7. Upper bound on PSF systematics due to their potentially incorrect modelling and deconvolution, assuming coefficients $\alpha = 0.01$ and $\beta = 2.0$ in eq. (A7). Horizontal axes show angular separations and vertical axes in each panel, from top to bottom respectively, show the absolute value of PSF $\langle M_{\text{ap}}^3 \rangle$, $\langle M_{\text{ap}}^2 M_{\times} \rangle$, $\langle M_{\text{ap}} M_{\times}^2 \rangle$ and $\langle M_{\times}^3 \rangle$ correlations divided by the E-mode signal $\langle M_{\text{ap}}^3 \rangle$ of the data, which is always at the sub-percent level and significantly smaller than the measurement errors. The dashed black line shows the 1% level and blue, light blue, salmon and red lines correspond to different cross-correlation of PSF properties e_p (the PSF ellipticity) and q (the ellipticity residual error after modeling).

- 1111 • In a non-tomographic analysis, we measure the nat-1166
1112 ural cosmic shear correlations Γ_i (the 3pt functions-1167
1113 analogous to the two-point functions ξ_{\pm}) in DES-1168
1114 Y3 data at high signal-to-noise ($2.5 \lesssim S/N \lesssim 7.0$ -1169
1115 for the real part of the correlations) and also ex-1170
1116 plore the triangle configuration dependence of 3pt-1171
1117 shear projections (respectively Figs. 2 and 4); 1172
- 1118 • Also in a non-tomographic setting, we measure the-1173
1119 skewness of the mass aperture statistic $\langle M_{\text{ap}}^3 \rangle$ both-1174
1120 in 1 aperture radius and in the generalized case of-1175
1121 3 aperture radii (left panel of Fig. 3 and Fig. 5)-1176
1122 The detection significance in all cases is very high-1177
1123 ($7.0 \lesssim S/N \lesssim 11.0$); 1178
- 1124 • We detect, for the first time, a tomographic-1180
1125 $\langle M_{\text{ap}}^3 \rangle(\theta)$ signature with high significance (total to-1181
1126 mographic $S/N = 15.0$) and additionally verify an-1182
1127 expected redshift evolution of the skewness param-1183
1128 eter $S(\theta; z)$ (respectively the right panel of Fig. 3-1184
1129 and Fig. 6); 1185
- 1130 • We verify that the third-order signatures found-1186
1131 are robust against B -mode systematics, parity-1187
1132 violating contributions and PSF modeling errors-1188
1133 thus validating that our measurements are likely a-1189
1134 result of astrophysical and gravitational phenom-1190
1135 ena (left panel of Fig. 3, right panel of Fig. 4 and-1191
1136 Fig. 7); 1192
- 1137 • We reproduce the main results in an N-body mock-1194
1138 catalog and verify that overall angular scale de-1195
1139 pendences and signal amplitudes of our measure-1196
1140 ments are broadly consistent with theoretical ex-1197
1141 pectations. 1198

1142 Given the high S/N of the data vectors here presented-200
1143 and the fact that systematics of observational origin are-201
1144 well under control, carrying forward with a cosmological-202
1145 analysis is a reasonable path. It is also interesting to note-203
1146 that our detected signals are non-zero even on relatively-204
1147 large angular separations of a degree or more, implying-205
1148 that non-Gaussian information coming from quasi-linear-206
1149 and linear scales could significantly add even to a conser-207
1150 vative 2pt cosmic shear analysis. We do, however, iden-208
1151 tify below several challenges that a joint 2pt+3pt pro-209
1152 gram would face. 1210

1153 First, analytic covariances for higher order moments-211
1154 of shear are notoriously complex, and their uncertain-212
1155 ties can significantly affect parameter posteriors. While-213
1156 it remains to be tested, it is possible that the jackknife-214
1157 approach employed here for the simple S/N estimates-215
1158 might not be sufficiently accurate for the more subtle-216
1159 inference of cosmological parameters. A mock-based co-217
1160 variance would be straightforward method, but we note-218
1161 that the 3pt measurement runtime is computationally ex-219
1162 pensive and could make that approach impractical unless-220
1163 we select 3pt statistics that minimize that computational-221
1164 cost. In particular, $\langle M_{\text{ap}}^3 \rangle$ and its generalized form have-222
1165 a high signal-to-noise with a relatively small number of-223

data points, which would make the use of mock covari-
ances more feasible.

Second, the modeling of astrophysical systematics such
as intrinsic alignments and baryons is likely to preclude
the use of small angular scales presented in our measure-
ments, and therefore it might be necessary to remove part
of those data points when fitting a theory model. How-
ever, the statistical uncertainties in our measurements
are large compared to those for 2pt cosmic shear, so it is
not unreasonable to expect that in fact relatively simple
theory modeling can be used for the 3pt data vector and
still maintain an acceptable level of potential biases. We
also point out that the nonlinear dark matter bispectrum
modeling itself is a challenge, although methods based on
fitting formulas calibrated against simulations have been
employed in the literature (Lazanu et al. 2016, Takahashi
et al. 2020).

Third, redshift and shape measurement uncertainties
propagate significantly into 3pt observables. These un-
certainties are calibrated to high accuracy and precision
in 2pt analyses, and a comparably careful analysis is
needed for 3pt correlations to determine their contribu-
tion to the error budget.

Many of the challenges above have already been ad-
dressed in cosmology studies including higher order lens-
ing correlations, in particular in the convergence mo-
ments work of Gatti et al. (2021a). In detail, the covari-
ance matrix estimation was made feasible in that work
with a data compression technique retaining a number
of data points smaller than the full length of the data
vector. Also, scale cuts were determined by the impact
of baryonic physics and other astrophysical contaminants
(such as intrinsic alignments and 3rd order contributions
such as source clustering) were modeled or shown to be
negligible. Finally, the nonlinear matter bispectrum was
obtained with a fitting formula calibrated on simulations
(Scoccimarro & Couchman 2001), and its computation
was made faster with an emulator technique.

We expect to employ similar methodologies for the real
space analysis of 2pt+3pt cosmic shear, but with some
differences in the details owing to the different choice
of estimators. The resulting constraints would provide
an important consistency check to the results of Gatti
et al. (2021a), with the advantage that the real space
statistics presented in this work (specifically the general,
three-aperture radii $\langle M_{\text{ap}}^3 \rangle(\theta_1, \theta_2, \theta_3)$) are guaranteed to
contain the total E -mode content in the shear field along
with specific configuration-dependent information. Fi-
nally, our measurements of the full 3-point function of
the shear field lay the groundwork to test for primordial
non-Gaussianity in the density field, e.g. via constraints
on the f_{NL} parameter, as studied theoretically by Takada
& Jain (2003a) and Hilbert et al. (2012).

With many practical challenges overcome and a
steadily increasing level of maturity, it is realistic to ex-
pect that that lensing 2pt+3pt analyses will be among
the central probes of S_8 and the Dark Energy equation-
of-state parameter w in current and future surveys such

as the Vera C. Rubin Observatory’s Legacy Survey of Space and Time⁷ (LSST), ESA’s Euclid mission⁸ and the Roman Space Telescope⁹. That is especially important since these experiments represent a massive investment of resources, and extracting as much useful information as possible from their data is highly desirable.

ACKNOWLEDGMENTS

We would like to thank Lucas Porth for comments and Ryuichi Takahashi and collaborators for making the T17 simulations utilized in this work publicly available. This work was completed in part with resources provided by the University of Chicago Research Computing Center. MJ is supported in part by National Science Foundation Award 1907610. BJ is supported in part by the US Department of Energy grant DE-SC0007901. CC is supported by DOE grant DE-SC0021949.

Funding for the DES Projects has been provided by the U.S. Department of Energy, the U.S. National Science Foundation, the Ministry of Science and Education of Spain, the Science and Technology Facilities Council of the United Kingdom, the Higher Education Funding Council for England, the National Center for Supercomputing Applications at the University of Illinois at Urbana-Champaign, the Kavli Institute of Cosmological Physics at the University of Chicago, the Center for Cosmology and Astro-Particle Physics at the Ohio State University, the Mitchell Institute for Fundamental Physics and Astronomy at Texas A&M University, Financiadora de Estudos e Projetos, Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro, Conselho Nacional de Desenvolvimento Científico e Tecnológico and the Ministério da Ciência, Tecnologia e Inovação, the Deutsche Forschungsgemeinschaft and the Collaborating Institutions in the Dark Energy Survey.

The Collaborating Institutions are Argonne National Laboratory, the University of California at Santa Cruz, the University of Cambridge, Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas-Madrid, the University of Chicago, University College London, the DES-Brazil Consortium, the University of Edinburgh, the Eidgenössische Technische Hochschule (ETH) Zürich, Fermi National Accelerator Laboratory, the University of Illinois at Urbana-Champaign, the Institut de Ciències de l’Espai (IEEC/CSIC), the Institut de Física d’Altes Energies, Lawrence Berkeley National Laboratory, the Ludwig-Maximilians Universität München and the associated Excellence Cluster Universe, the University of Michigan, the National Optical Astronomy Observatory, the University of Nottingham, The Ohio State

University, the University of Pennsylvania, the University of Portsmouth, SLAC National Accelerator Laboratory, Stanford University, the University of Sussex, Texas A&M University, and the OzDES Membership Consortium.

The DES data management system is supported by the National Science Foundation under Grant Numbers AST-1138766 and AST-1536171. The DES participants from Spanish institutions are partially supported by MICINN under grants ESP2017-89838, PGC2018-094773, PGC2018-102021, SEV-2016-0588, SEV-2016-0597, and MDM-2015-0509, some of which include ERDF funds from the European Union. IFAE is partially funded by the CERCA program of the Generalitat de Catalunya. Research leading to these results has received funding from the European Research Council under the European Union’s Seventh Framework Program (FP7/2007-2013) including ERC grant agreements 240672, 291329, and 306478. We acknowledge support from the Brazilian Instituto Nacional de Ciência e Tecnologia (INCT) do e-Universo (CNPq grant 465376/2014-2). We acknowledge support from the Australian Research Council Centre of Excellence for All-sky Astrophysics (CAASTRO), through project number CE110001020.

This manuscript has been authored by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics. The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

Based in part on observations at Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy (AURA) under a cooperative agreement with the National Science Foundation. This work made use of Matplotlib (Hunter 2007) and NASA’s Astrophysics Data System Bibliographic Services (ADS).

Appendix A: Derivation of PSF corrections

As ellipticities e are measured from galaxy images, their PSF must be deconvolved. As in Gatti & Sheldon et al., (2021c), we define the errors in the PSF modeling as well improper deconvolution both as additive contributions to the measured ellipticities:

$$e = \gamma + \delta_e, \quad (\text{A1})$$

where the additive factor δ_e is defined by

$$\delta_e \equiv \alpha e^p + \beta q; \quad q \equiv e_* - e^p, \quad (\text{A2})$$

where e^p is the modeled PSF ellipticity (referred to as e_{model} in Gatti & Sheldon et al., (2021c)) and e_* is the

⁷ <https://www.lsst.org>

⁸ <https://www.euclid-ec.org>

⁹ <https://roman.gsfc.nasa.gov>

1327 actually measured PSF. That means the coefficients α
 1328 and β are respectively interpreted as a leakage of the
 1329 modeled PSF shape onto the galaxy ellipticity e (coming
 1330 possibly from incorrect deconvolutions) and errors in the
 1331 interpolation of the PSF shape. The shear fields e^p and q
 1332 are estimated from *reserved stars* which do not contribute
 1333 to the PSF fitting, that is, where both the modeled PSF
 1334 and the true PSF are known, otherwise we would have
 1335 $q \rightarrow 0$ by construction.

Using the same definitions in eq. (12), we can propa-
 gate the PSF correction in eq. (A1) to the 1-point quan-
 tities:

$$M_{\text{ap}}(R) = \int d^2 R Q(R) \gamma_t + \int d^2 R Q(R) [\alpha e_t^p + \beta q_t]$$

$$M_{\times}(R) = \underbrace{\int d^2 R Q(R) \gamma_{\times}}_{=0} + \int d^2 R Q(R) [\alpha e_{\times}^p + \beta q_{\times}].$$

As the cross-projections of the PSF residuals e_{\times}^p and q_{\times}
 can generally have non-zero statistical moments, we see
 that the additive PSF errors defined above can contaminate
 both E-modes and B-modes. As a stepping stone
 for the third-order case, we can again follow Jarvis et al.
 (2004) and get, for the second-order mass aperture:

$$\langle M^2 \rangle = \int d^2 R_1 d^2 R_2 Q(R_1) Q(R_2) \langle (\gamma + \delta_e) (\gamma + \delta_e) \rangle$$

$$\times \exp(-2i(\phi_1 + \phi_2))$$

$$\langle MM^* \rangle = \int d^2 R_1 d^2 R_2 Q(R_1) Q(R_2) \langle (\gamma + \delta_e) (\gamma + \delta_e)^* \rangle$$

$$\times \exp(-2i(\phi_1 - \phi_2)).$$

We can safely assume that the expected value of correla-
 tions between the gravitational shear and PSF residuals is
 zero so the cross-terms $\langle \gamma \delta_e \rangle$ vanish. Then, defining the
 PSF correlations ξ_{\pm}^{psf} analogously to how the (gravita-
 tional) shear correlations are defined, that is $\xi_+ \equiv \langle \gamma \gamma^* \rangle$,
 $\xi_- \equiv \langle \gamma \gamma \exp(-4i\theta) \rangle$, we see that the PSF corrections
 are simply additive at the mass aperture level:

$$\langle M^2 \rangle(R) = \int \frac{s ds}{R^2} \left(\xi_-^{\gamma}(s) + \xi_-^{\text{psf}}(s) \right) T_- \left(\frac{s}{R} \right) \quad (\text{A3})$$

$$\langle MM^* \rangle(R) = \int \frac{s ds}{R^2} \left(\xi_+^{\gamma}(s) + \xi_+^{\text{psf}}(s) \right) T_+ \left(\frac{s}{R} \right) \quad (\text{A4})$$

1336 where $\xi_{\pm}^{\text{psf}} \equiv \alpha^2 \langle e_p e_p \rangle_{\pm} + \alpha \beta \langle e_p q \rangle_{\pm} + \beta^2 \langle qq \rangle_{\pm}$, and
 1337 where the functions T_{\pm} are defined in Appendix B.

1338 The terms $\langle M_{\text{ap}}^2 \rangle$ and $\langle M_{\times}^2 \rangle$ can be expressed as sim-
 1339 ple linear combinations of the quantities above (Jarvis
 1340 et al. 2004). While $\langle M_{\times}^2 \rangle$ would represent B-mode sig-
 1341 nal which can generally become non-zero in the presence
 1342 of uncorrected PSF errors, the term $\langle M_{\text{ap}} M_{\times} \rangle$, if found
 1343 to be non-negligible, would additionally imply a parity
 1344 violating contribution.

The reasoning above also applies to the third-order mo-
 ments of the same observables. We define the 3pt PSF

correlations in the same way we define the natural com-
 ponents of the shear signal and write

$$\langle M^3 \rangle(R) = - \int d^2 R_1 d^2 R_2 d^2 R_3 Q(R_1) Q(R_2) Q(R_3)$$

$$\times \langle (\gamma + \delta_e)^3 \exp[-2i(\alpha + \beta + \delta)] \rangle$$

$$= - \int \frac{s ds}{R^2} \int \frac{d^2 t}{2\pi R^2} \left(\Gamma_0 + \Gamma_0^{\text{psf}} \right) T_0(s, t) \quad (\text{A5})$$

$$\langle M^2 M^* \rangle(R) = \int d^2 R_1 d^2 R_2 d^2 R_3 Q(R_1) Q(R_2) Q(R_3)$$

$$\times \langle (\gamma + \delta_e)^2 (\gamma + \delta_e)^* \exp[-2i(\alpha + \beta - \delta)] \rangle$$

$$= \int \frac{s ds}{R^2} \int \frac{d^2 t}{2\pi R^2} \left(\Gamma_1 + \Gamma_1^{\text{psf}} \right) T_1(s, t) \quad (\text{A6})$$

where we have introduced the PSF correction at the 3pt
 level as

$$\Gamma_{0,1}^{\text{psf}} = \alpha^3 \langle e_p^3 \rangle_{0,1} + 3\alpha^2 \beta \langle e_p^2 q \rangle_{0,1} + 3\alpha \beta^2 \langle e_p q^2 \rangle_{0,1} + \beta^3 \langle q^3 \rangle_{0,1}. \quad (\text{A7})$$

The derivation above assumes that cross-terms of
 the type $\langle \gamma \delta_e^2 \rangle$ or $\langle \gamma^2 \delta_e \rangle$ are null when averaged over
 large ensembles, as both of these terms boil down to
 whether the 1(2)-point gravitational shear correlates with
 the 2(1)-point PSF's, which should not be the case.
 The expressions for $\langle M_{\text{ap}}^3 \rangle$, $\langle M_{\text{ap}}^2 M_{\times} \rangle$, $\langle M_{\text{ap}} M_{\times}^2 \rangle$ and
 $\langle M_{\times}^3 \rangle$ can be obtained from the ones above as shown in
 Jarvis et al. (2004), and it remains true that $\langle M_{\text{ap}}^2 M_{\times} \rangle$
 and $\langle M_{\times}^3 \rangle$ are null in order to conserve parity, while
 $\langle M_{\text{ap}} M_{\times}^2 \rangle$ may include non-zero PSF B-modes that cor-
 relate with E-modes (in addition to astrophysical B-mode
 contributions).

Appendix B: Definition of coordinates and TreeCorr internal variables

Here we clarify some of the notation utilized in this
 draft, mainly in what refers to coordinates and definitions
 of triangle sides and their respective angles. We use the
 same conventions of Jarvis et al. (2004) and reproduce
 their Fig. 1 below in our Fig 8.

With \mathbf{q}_1 , \mathbf{q}_2 and \mathbf{q}_3 defined as the vectors from each of
 the triangle vertices to the centroid of the triangle, and s
 and \mathbf{t}' the sides of the triangle (notice that without loss
 of generality we fix s and take \mathbf{t}' to be at an angle α with
 respect to that line), we have

$$\mathbf{q}_1 = \frac{(s + \mathbf{t}')}{3}, \quad \mathbf{q}_2 = \frac{(\mathbf{t}' - 2s)}{3}, \quad \mathbf{q}_3 = \frac{(s - 2\mathbf{t}')}{3}. \quad (\text{B1})$$

The functions T_0 and T_1 that enter the mass aperture
 computations such as eq. (19) are purely geometrical and




FIG. 8. Convention for coordinate systems. **Top panel:** TREECORR internal variables and side length definitions as utilized in eqs. (B1)-(B5), reproduced from Jarvis et al. (2004). **Bottom panel:** definition of distances of interest and shear projections (t, \times) relevant in the $\Gamma_i(\theta_{\text{medium}})$ and $\langle M_{\text{ap}}(\theta_1, \theta_2, \theta_3) \rangle$ measurements. Notably, the reference scale θ_{medium} is a side length, while the scales θ_i ($i = 1, 2, 3$) are radii from the triangle center.

1375 dependent on the vectors above:

$$1376 \quad T_0(s, t) = -\frac{(\mathbf{q}_1^* \mathbf{q}_2^* \mathbf{q}_3^*)^2}{24} \exp\left(-\frac{q_1^2 + q_2^2 + q_3^2}{2}\right) \quad (\text{B2})$$

$$1377 \quad T_1(s, t) = -\left(\frac{(\mathbf{q}_1 \mathbf{q}_2^* \mathbf{q}_3^*)^2}{24} - \frac{q_1^2 \mathbf{q}_2^* \mathbf{q}_3^*}{9} + \frac{q_1^{*2} + 2\mathbf{q}_2^* \mathbf{q}_3^*}{27}\right) \\ \times \exp\left(-\frac{q_1^2 + q_2^2 + q_3^2}{2}\right), \quad (\text{B3})$$

1377 where bold symbols are vectors in complex notation with
1378 x/y on the real/imaginary direction, eg $\mathbf{v} = v_x + iv_y$.

1379 Additionally, internal TREECORR units utilized to bin
1380 triangles are such that, for triangles of side lengths $d_1 \leq$
1381 $d_2 \leq d_3$, we have

$$1382 \quad u = \frac{d_3}{d_2}, \quad (\text{B4})$$

1383

$$1384 \quad v = \pm \frac{(d_1 - d_2)}{d_3}, \quad (\text{B5})$$

1385 where the positive and negative signs of v correspond
1386 to whether side lengths are in clockwise or counterclockwise
1387 order respectively, and recall that we have named
1388 $\theta_{\text{medium}} = d_2$ to conveniently bin the Γ_i functions in
1389 eq.(22). Note that with these definitions we have $u \in$
1390 $[0, 1]$ and $v \in [-1, 1]$. In practice, selecting i.e. equilateral
1391 triangles of characteristic side length θ_{medium} within
1392 the output corresponds to sub-selecting the galaxies in
1393 bins $u \sim 0$ and $v \sim 0$.

1394

Appendix C: Signal-to-noise of a Vector

The signal-to-noise ratio of a scalar value, X , with a Gaussian uncertainty, σ , is well-defined. The signal is the expectation value of the measurement $\langle X \rangle$, and the noise is the standard deviation of the uncertainty E . Thus, the signal-to-noise is simply the ratio of these.

$$X = \langle X \rangle + E \quad (\text{C1})$$

$$E \sim \mathcal{N}(0, \sigma) \quad (\text{C2})$$

$$S/N(X) \equiv \frac{\langle X \rangle}{\sigma} \quad (\text{C3})$$

$$= \frac{\langle X \rangle}{\sqrt{\text{Var}(X)}} \quad (\text{C4})$$

However, it is less obvious what the corresponding quantity should be for a vector \mathbf{d} , where each component of the vector is itself a measurement with an uncertainty. We start by considering a data vector of independent measurements, each with its own Gaussian uncertainty.

$$\mathbf{d} = \{d_i\} \quad (\text{C5})$$

$$d_i = \langle d_i \rangle + E_i \quad (\text{C6})$$

$$E_i \sim \mathcal{N}(0, \sigma_i) \quad (\text{C7})$$

We consider all possible linear combinations of the vector elements,

$$X_{\mathbf{w}} \equiv \mathbf{w} \cdot \mathbf{d} = \sum_i w_i d_i, \quad (\text{C8})$$

for arbitrary weight vectors \mathbf{w} . For each choice of \mathbf{w} , the scalar quantity $X_{\mathbf{w}}$ of course has a well-defined signal-to-noise, given by Equation C4, but each choice may be different, depending on the specific weights being used.

Among all such possible choices, we take the one with the largest signal-to-noise to define the signal-to-noise of the vector \mathbf{d} .

$$S/N(\mathbf{d}) \equiv \max_{\mathbf{w}} \left(\frac{\langle X_{\mathbf{w}} \rangle}{\sqrt{\text{Var}(X_{\mathbf{w}})}} \right) \quad (\text{C9})$$

We therefore need to determine what choice of weights \mathbf{w} gives the largest signal-to-noise for $X_{\mathbf{w}}$. For a given choice of \mathbf{w} , we have

$$(S/N)^2 = \frac{\left(\sum_j w_j \langle d_j \rangle \right)^2}{\sum_j w_j^2 \sigma_j^2}. \quad (\text{C10})$$

As usual, we find w_i at the extremum by setting the derivative to 0.

$$0 = \frac{\partial (S/N)^2}{\partial w_i} = \frac{2 \left(\sum_j w_j \langle d_j \rangle \right) \langle d_i \rangle}{\sum_j w_j^2 \sigma_j^2} - \frac{2 w_i \sigma_i^2 \left(\sum_j w_j \langle d_j \rangle \right)^2}{\left(\sum_j w_j^2 \sigma_j^2 \right)^2}$$

$$\langle d_i \rangle \sum_j w_j^2 \sigma_j^2 = w_i \sigma_i^2 \sum_j w_j \langle d_j \rangle$$

$$w_i = \frac{\langle d_i \rangle}{\sigma_i^2} \quad (\text{C11})$$

The signal-to-noise for this choice of \mathbf{w} is then

$$S/N = \frac{\sum_i w_i \langle d_i \rangle}{\sqrt{\sum_i w_i^2 \sigma_i^2}}$$

$$= \frac{\sum_i \langle d_i \rangle^2 / \sigma_i^2}{\sqrt{\sum_i (\langle d_i \rangle / \sigma_i^2)^2 \sigma_i^2}}$$

$$= \sqrt{\sum_i \frac{\langle d_i \rangle^2}{\sigma_i^2}}. \quad (\text{C12})$$

Aside from the expectation value in the numerator, this is equivalent to $\sqrt{\chi^2}$, which is a relatively common approximation used to estimate the signal-to-noise of a vector. Calculating the expectation value of χ^2 , we find

$$\langle \chi^2 \rangle = \left\langle \sum_i \frac{d_i^2}{\sigma_i^2} \right\rangle$$

$$= \sum_i \frac{(\langle d_i \rangle + E_i)^2}{\sigma_i^2}$$

$$= \sum_i \frac{\langle d_i \rangle^2 + 2\langle d_i \rangle \langle E_i \rangle + \langle E_i^2 \rangle}{\sigma_i^2}$$

$$= \sum_i \frac{\langle d_i \rangle^2 + \sigma_i^2}{\sigma_i^2}$$

$$= (S/N)^2 + N_{\text{d.o.f.}} \quad (\text{C13})$$

Thus, we have derived the relatively simple relationship,

$$S/N = \sqrt{\langle \chi^2 \rangle - N_{\text{d.o.f.}}} \quad (\text{C14})$$

In practice, one does not have access to the expectation value $\langle \chi^2 \rangle$, so we replace it with its measured value, which is the best we can do:

$$S/N = \sqrt{\chi^2 - N_{\text{d.o.f.}}} \quad (\text{C15})$$

For high signal-to-noise vectors, the approximation $S/N = \sqrt{\chi^2}$ is not bad. But when χ^2 is only moderately larger than the number of degrees of freedom, the correction is important, and one should instead use Equation C15. And of course if the measured χ^2 is less than $N_{\text{d.o.f.}}$, there is no detection, and the signal-to-noise is essentially zero.

Finally, what if the uncertainties are correlated? That is, what if the data vector has a non-diagonal covariance matrix C ? It turns out that this case can be reduced to the same formula as above by diagonalizing C and changing to the basis where the covariance is diagonal.

$$\text{Cov}(\mathbf{d}) \equiv C = V \Lambda V^T \quad (\text{C16})$$

$$\mathbf{z} \equiv V^T \mathbf{d} \quad (\text{C17})$$

$$\text{Cov}(\mathbf{z}) = V^T \text{Cov}(\mathbf{d}) V$$

$$= V^T V \Lambda V^T V$$

$$= \Lambda \quad (\text{C18})$$

Given our definition (Equation C9), the signal-to-noise of \mathbf{z} is the same as the signal-to-noise of \mathbf{d} . Furthermore, the χ^2 for the two vectors are also equal:

$$\chi^2 = \mathbf{z}^T \Lambda^{-1} \mathbf{z} = \sum_i \frac{z_i^2}{\Lambda_{ii}}$$

$$= (V^T \mathbf{d})^T \Lambda^{-1} V^T \mathbf{d}$$

$$= \mathbf{d}^T V \Lambda^{-1} V^T \mathbf{d}$$

$$= \mathbf{d}^T (V \Lambda V^T)^{-1} \mathbf{d}$$

$$= \mathbf{d}^T C^{-1} \mathbf{d}, \quad (\text{C19})$$

where we used the fact that $V^T = V^{-1}$.

We know that the signal-to-noise of \mathbf{z} is given by Equation C15, since it has uncorrelated uncertainties. Since \mathbf{d} has the same signal-to-noise as \mathbf{z} , and it has the same χ^2 and $N_{\text{d.o.f.}}$, this must also be the correct formula for \mathbf{d} . Therefore, Equation C15 applies even to a vector with a non-trivial covariance matrix.

- 1409 Amon A., et al. 2021, arXiv e-prints, p. arXiv:2105.13543
1410 Asgari M., et al. 2021, A&A, 645, A104
1411 Benabed K., Scoccimarro R., 2006, A&A, 456, 421
1412 Bernardeau F., van Waerbeke L., Mellier Y., 1997, A&A, 322, 1
1413 1
1414 Bernardeau F., Mellier Y., van Waerbeke L., 2002, A&A, 389, L28
1415 L28
1416 Cheng S., et al. 2020, MNRAS, 499, 5902
1417 Cooray A., Sheth R., 2002, Phys. Rep., 372, 1
1418 Crittenden R. G., et al. 2002, Astrophys. J., 568, 20
1419 DES Collaboration 2022, Phys. Rev. D, 105, 023520
1420 Dodelson S., Schneider M. D., 2013, Phys. Rev. D, 88, 063537
1421 Erben T., et al. 2013, MNRAS, 433, 2545
1422 Flaughner B., et al. 2015, A.J., 150, 150
1423 Fluri J., et al. 2019, Phys. Rev. D, 100, 063514
1424 Friedrich O., et al. 2018, Phys. Rev. D, 98, 023508
1425 Fu L., et al. 2014, MNRAS, 441, 2725
1426 Gatti M., et al. 2021a, arXiv e-prints, p. arXiv:2110.10141
1427 Gatti M., Sheldon E., et al., 2021b, Mon. Not. Roy. Astron Soc., 504, 4312
1428 Soc., 504, 4312
1429 Gatti M., et al. 2021c, MNRAS, 504, 4312
1430 Górski K. M., et al. 2005, Astrophys. J., 622, 759
1431 Gruen D., et al. 2018, Phys. Rev. D, 98, 023507
1432 Halder A., et al. 2021, MNRAS, 506, 2780
1433 Hamana T., et al. 2015, PASJ, 67, 34
1434 Hamana T., et al. 2020, PASJ, 72, 16
1435 Hartlap J., Simon P., Schneider P., 2007, A&A, 464, 399
1436 Hikage C., et al. 2019, PASJ, 71, 43
1437 Hilbert S., et al. 2012, MNRAS, 426, 2870
1438 Hinshaw G., et al. 2013, ApJS, 208, 19
1439 Ho S., White M., 2004, Astrophys. J., 607, 40
1440 Huff E., Mandelbaum R., 2017, arXiv e-prints, p. arXiv:1702.02600
1441 arXiv:1702.02600
1442 Hunter J. D., 2007, Computing in Science & Engineering, 9, 90
1443 90
1444 Huterer D., et al. 2006, MNRAS, 366, 101
1445 Jain B., Seljak U., 1997, Astrophys. J., 484, 560
1446 Jarvis M., Bernstein G., Jain B., 2004, MNRAS, 352, 338
1447 Jeffrey N., Alsing J., Lanusse F., 2021, MNRAS, 501, 954
1448 Joachimi B., Shi X., Schneider P., 2009, A&A, 508, 1193
1449 Jung G., et al. 2021, JCAP, 2021, 055
1450 Kacprzak T., et al. 2016, MNRAS, 463, 3653
1451 Kayo I., Takada M., 2013, arXiv e-prints, p. arXiv:1306.4684
1452 Kilbinger M., Schneider P., 2005, A&A, 442, 69
1453 Kitching T. D., Deshpande A. C., Taylor P. L., 2021, arXiv e-prints, p. arXiv:2110.01275
1454 e-prints, p. arXiv:2110.01275
1455 Lazanu A., et al. 2016, Phys. Rev. D, 93, 083517
1456 Limber D. N., 1953, Astrophys. J., 117, 134
1457 LoVerde M., Afshordi N., 2008, Phys. Rev. D, 78, 123506
1458 Lu T., Haiman Z., Zorrilla Matilla J. M., 2021, arXiv e-prints, p. arXiv:2109.11060
1459 p. arXiv:2109.11060
1460 Myles J., et al. 2021, MNRAS, 505, 4249
1461 Planck Collaboration 2020, A&A, 641, A6
1462 Porth L., Smith R. E., 2021, MNRAS, 508, 3474
1463 Porth L., et al. 2020, MNRAS, 499, 2474
1464 Pyne S., Joachimi B., 2021, MNRAS, 503, 2300
1465 Rowe B., 2010, MNRAS, 404, 350
1466 Sato M., Nishimichi T., 2013, Phys. Rev. D, 87, 123538
1467 Schneider P., 2003, A&A, 408, 829
1468 Schneider P., Lombardi M., 2003, A&A, 397, 809
1469 Schneider P., et al. 1998, MNRAS, 296, 873
1470 Schneider P., van Waerbeke L., Mellier Y., 2002, A&A, 389, 729
1471 729
1472 Schneider P., Kilbinger M., Lombardi M., 2005, A&A, 431, 9
1473 Schrabback T., et al. 2010, A&A, 516, A63
1474 Scoccimarro R., Couchman H. M. P., 2001, MNRAS, 325, 1312
1475 1312
1476 Scoville N., et al. 2007, ApJS, 172, 38
1477 Secco L. F., et al. 2021, arXiv e-prints, p. arXiv:2105.13544
1478 Sellentin E., Heavens A. F., 2016, Monthly Notices of the Royal Astronomical Society, 464, 4658
1479 Semboloni E., et al. 2011, MNRAS, 410, 143
1480 Semboloni E., Hoekstra H., Schaye J., 2013, MNRAS, 434, 148
1481 148
1482 Sevilla-Noarbe I., et al. 2021, ApJS, 254, 24
1483 Sheldon E. S., Huff E. M., 2017, Astrophys. J., 841, 24
1484 Shi X., Joachimi B., Schneider P., 2014, A&A, 561, A68
1485 Shi X., Joachimi B., Schneider P., 2014, A&A, 561, A68
1486 Springel V., 2005, MNRAS, 364, 1105
1487 Takada M., Jain B., 2003a, MNRAS, 344, 857
1488 Takada M., Jain B., 2003b, ApJ, 583, L49
1489 Takahashi R., et al. 2017, Astrophys. J., 850, 24
1490 Takahashi R., et al. 2020, Astrophys. J., 895, 113
1491 Taylor A., Joachimi B., Kitching T., 2013, Monthly Notices of the Royal Astronomical Society, 432, 1928
1492 Troxel M. A., Ishak M., 2012, MNRAS, 423, 1663
1493 Troxel M. A., Ishak M., 2015, Phys. Rep., 558, 1
1494 Van Waerbeke L., et al. 2002, A&A, 393, 369
1495 Zaldarriaga M., Scoccimarro R., 2003, Astrophys. J., 584, 559
1496 Zonca A., et al. 2019, Journal of Open Source Software, 4, 1298
1497 Zürcher D., et al. 2021, arXiv e-prints, p. arXiv:2110.10135