

This is the accepted manuscript made available via CHORUS. The article has been published as:

Four curious supergravities

M. J. Duff and S. Ferrara

Phys. Rev. D **83**, 046007 — Published 16 February 2011

DOI: [10.1103/PhysRevD.83.046007](https://doi.org/10.1103/PhysRevD.83.046007)

Four curious supergravities

M. J. Duff^{1,*} and S. Ferrara^{2,3,4,†}

¹*Theoretical Physics, Blackett Laboratory, Imperial College London, London SW7 2AZ, United Kingdom*

²*Physics Department, Theory Unit, CERN, CH 1211, Geneva 23, Switzerland*

³*INFN - Laboratori Nazionali di Frascati, Via Enrico Fermi 40, 00044 Frascati, Italy*

⁴*Department of Physics and Astronomy, University of California, Los Angeles, CA USA*

(Dated: October 28, 2010)

We consider four supergravities with 16+16, 32+32, 64+64, 128+128 degrees of freedom displaying some curious properties: (1) They exhibit minimal supersymmetry ($\mathcal{N} = 1, 2, 2, 1$) but maximal rank ($r = 7, 6, 4, 0$) of the scalar coset in $D = 4, 5, 7, 11$. (2) They couple naturally to supermembranes and admit these membranes as solutions. (3) Although the $D = 4, 5, 7$ supergravities follow from truncating the maximally supersymmetric ones, there nevertheless exist M-theory compactifications with G_2 , $SU(3)$, $SU(2)$ holonomy having these supergravities as their massless sectors. (4) They reduce to $\mathcal{N} = 1, 2, 4, 8$ theories all with maximum rank 7 in $D = 4$ which (5) correspond to 0, 1, 3, 7 lines of the Fano plane and hence admit a division algebra ($\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$) interpretation consistent with the black-hole/qubit correspondence, (6) are generalized self-mirror and hence (7) have vanishing on-shell trace anomaly.

PACS numbers: 11.25.-w

Keywords: keywords

Contents

I. Introduction	2
II. Minimal supergravities in $D = 4, 5, 7, 11$	5
A. Compactifications	5
B. Supermultiplets	6
C. Lagrangians	7
D. Membrane solutions	8
E. Cosets	8
III. $\mathcal{N} = 1, 2, 4, 8$ in $D = 4$	9
A. Betti numbers	9
B. Self-mirror with vanishing trace anomaly	9
C. Cosets	11
D. Fano plane and $\mathbb{O}, \mathbb{C}, \mathbb{H}, \mathbb{R}$	12
IV. $\mathcal{N} = 2, 4, 8, 16$ in $D = 3$	14
A. Compactifications	14
B. Cosets	15
V. Acknowledgments	16
References	17

*m.duff@imperial.ac.uk

†sergio.ferrara@cern.ch

- $D = 7$: $\mathcal{N}=2$ graviton + 3 vector, $f = 40 + 40 + 3(8 + 8) = 64 + 64$

$$(g_{\mu\nu}, \mathcal{A}; 2\psi_\mu, 2\chi; A_{\mu\nu\rho}, 3A_\mu) + 3(3\mathcal{A}; 2\chi; A_\mu)$$

with rank 4 scalar coset

$$\frac{G}{H} = SO(1,1) \times \frac{SL(4,R)}{SO(4)}$$

- $D = 5$: $\mathcal{N}=2$ graviton + 2 vector + 3 hyper + 1 3-form, $f = 8 + 8 + 2(4 + 4) + 3(4 + 4) + (4 + 4) = 32 + 32$

$$(g_{\mu\nu}; 2\psi_\mu; A_\mu) + 2(\mathcal{A}; 2\chi; A_\mu) + 3(2\mathcal{A}; 2\chi; 2A) + (\mathcal{A}; 2\chi; A_{\mu\nu\rho}, 2A)$$

with rank 6 scalar coset

$$\frac{G}{H} = SO(1,1)^3 \times \frac{SO(3,4)}{SO(3) \times SO(4)} \ltimes R^2$$

- $D = 4$: $\mathcal{N}=1$ graviton + 7 WZ, $f = 2 + 2 + 7(2 + 2) = 16 + 16$

$$(g_{\mu\nu}; \psi_\mu; A_{\mu\nu\rho}) + 7(\mathcal{A}; \chi; A)$$

with rank 7 scalar coset

$$\frac{G}{H} = \frac{SL(2)^7}{SO(2)^7}$$

The U-duality is only $SL(2)^6 \times SO(1,1) \ltimes R$, however, because of the coupling of the scalars to $A_{\mu\nu\rho}$.

These theories may also be derived by compactifying M-theory on T^4 , T^6 and T^7 and truncating the massless sectors so as to obtain minimal supersymmetry ($\mathcal{N} = 2, 2$ and 1 respectively) while preserving maximum rank of the scalar coset ($r = 4, 6$ and 7 respectively). They exhibit several other remarkable properties. For example, they admit membranes as elementary (electric) solutions, by virtue of the universal presence of a 3-form $A_{\mu\nu\rho}$ and by virtue of the correct dilaton exponent which follows from the maximum rank condition. We therefore expect that they will be compatible with the superspace constraints enforced by kappa symmetry on the worldvolume of the Green-Schwarz membranes [6], but we do not address this problem here.

It will be important for our purposes to distinguish between the Lagrangians obtained directly in these two ways and the conventional supergravity Lagrangians obtained after dualization of the p -forms. For example, the latter has no $A_{\mu\nu\rho}$ field in $D = 4$ and so not only has a different symmetry, namely $SL(2)^7$ as opposed to the $SL(2)^6 \times SO(1,1) \ltimes R$ of the former, but also admits no electric membrane (domain wall) solution.

As described in section III, these four theories may be further reduced to $\mathcal{N} = 1, 2, 4$ and 8 theories all with maximum rank $r = 7$ in $D = 4$, corresponding to compactification on X^7 with independent betti numbers

$$(b_0, b_1, b_2, b_3) = (1, \mathcal{N} - 1, 3\mathcal{N} - 3, 4\mathcal{N} + 3) \quad (2)$$

Compactifications with such betti numbers may indeed be found in [12] for $\mathcal{N} = 8$, in [13, 14] for $\mathcal{N} = 4$ and in [15–17] for $\mathcal{N} = 2$ and $\mathcal{N} = 1$. We show that the corresponding supergravities before dualization are:

- $T^7 : (1, 7, 21, 35)$

$\mathcal{N}=8$ graviton, $f = 128 + 128$,

$$(g_{\mu\nu}, 7\mathcal{A}_\mu, 28\mathcal{A}; 8\psi_\mu, 56\chi; A_{\mu\nu\rho}, 7A_{\mu\nu}, 21A_\mu, 35A)$$

with rank 7 scalar coset

$$\frac{G}{H} = SO(1,1) \times \frac{SL(7,R)}{SO(7)} \ltimes R^{35}$$

- $X^4 \times T^3 : (1, 3, 9, 19)$

$\mathcal{N}=4$ graviton + 3 vector + 3 2-form, $f = 16 + 16 + 3(8 + 8) + 3(8 + 8) = 64 + 64$,

$$(g_{\mu\nu}, 3\mathcal{A}_\mu, \mathcal{A}; 4\psi_\mu, 4\chi; A_{\mu\nu\rho}, 3A_\mu, A) + 3(3\mathcal{A}; 4\chi; A_\mu, 3A) + 3(2\mathcal{A}; 4\chi; A_{\mu\nu}, A_\mu, 3A)$$

with rank 7 scalar coset

$$\frac{G}{H} = \frac{SL(2,R)}{SO(2)} \times \frac{SO(3,6)}{SO(3) \times SO(6)} \times SO(1,1) \times \frac{SL(3)}{SO(3)} \ltimes R^9$$

- $X^6 \times S^1 : (1, 1, 3, 11)$

$\mathcal{N}=2$ graviton + 3 vector + 3 hyper +1 linear, $f = 4 + 4 + 3(4 + 4) + 3(4 + 4) + (4 + 4) = 32 + 32$,

$$(g_{\mu\nu}, \mathcal{A}_\mu; 2\psi_\mu; A_{\mu\nu\rho}) + 3(\mathcal{A}; 2\chi; A_\mu, A) + 3(2\mathcal{A}; 2\chi; 2A) + (\mathcal{A}; 2\chi; A_{\mu\nu}, 2A)$$

with rank 7 scalar coset

$$\frac{G}{H} = SO(1, 1) \times \frac{SL(2, R)^3}{SO(2)^3} \times \frac{SO(3, 4)}{SO(3) \times SO(4)} \ltimes R^2$$

- $X^7 : (1, 0, 0, 7)$

$\mathcal{N}=1$ graviton +7 WZ, $f = 2 + 2 + 7(2 + 2) = 16 + 16$,

$$(g_{\mu\nu}; \psi_\mu; A_{\mu\nu\rho}) + 7(\mathcal{A}; \chi; A)$$

with rank 7 scalar coset

$$\frac{G}{H} = \frac{SL(2)^7}{SO(2)^7}$$

Interestingly enough, the cases $\mathcal{N} = 8$, $\mathcal{N} = 4$ and $\mathcal{N} = 2$ (albeit without the three hyper and one linear multiplet [18]) have already made an appearance in the context of the *black-hole/qubit correspondence* [19–21], where their 56, 24, 8 black hole charges correspond to 7, 3, 1 lines of the Fano plane [22–24]. In particular, the $\mathcal{N} = 2$ supergravity is just the STU model whose black holes have a Bekenstein-Hawking entropy given by Cayley’s hyperdeterminant, the same quantity that describes the entanglement of three qubits. The 7, 3, 1 lines of the Fano plane in turn provide the multiplication table of the imaginary octonion, quaternion, and complex numbers respectively. The fourth $\mathcal{N} = 1$ supergravity completes the set with 0 lines, corresponding to the reals.

Earlier work on the branscan gave an $\mathbb{O}, \mathbb{H}, \mathbb{C}, \mathbb{R}$ division algebra interpretation to the four sequences appearing in Table I, which have $8 + 8, 4 + 4, 2 + 2, 1 + 1$ worldvolume degrees of freedom. See [25–27] and references therein. Since our supergravities are obtained by compactification, however, the corresponding membranes all have $8 + 8$.

Furthermore, we recall that in [28] we defined a generalized mirror symmetry

$$(b_0, b_1, b_2, b_3) \rightarrow (b_0, b_1, b_2 - \rho/2, b_3 + \rho/2) \quad (3)$$

under which

$$\rho \equiv 7b_0 - 5b_1 + 3b_2 + b_3 \quad (4)$$

changes sign

$$\rho \rightarrow -\rho \quad (5)$$

Generalized self-mirror theories are defined to be those for which ρ vanishes. In the case of G_2 manifolds with $b_1 = 0$, Joyce [15, 16] refers to $\rho = 0$ as an “axis of symmetry”. For related work on mirror symmetry and Joyce-manifolds, see [17, 29, 30].

Moreover the quantity ρ also shows up in the on-shell Weyl anomaly [31, 32], before dualization [33], which is given by

$$g_{\mu\nu} \langle T^{\mu\nu} \rangle = A \frac{1}{32\pi^2} R^{*\mu\nu\rho\sigma} R^*_{\mu\nu\rho\sigma} \quad (6)$$

where

$$A = -\frac{1}{24}\rho \quad (7)$$

Since our four curious supergravities all have $\rho = 0$, they are self-mirror in the above sense and hence have vanishing Weyl anomaly.

Finally we note that a spacelike reduction gives the four Type IIA supergravities that couple to superstrings in $D = 3, 4, 6, 10$. They yield $\mathcal{N} = 16, 8, 4, 2$ supergravities in $D = 3$. While a timelike reduction from $D = 4$ to $D = 3$ yields after dualization the four cosets that play a role in the four-way entanglement of eight qubits [24, 34–37], namely $E_{8(8)}/SO^*(16)$, $SO(8, 8)/SO(4, 4)^2$, $SO(4, 4)^2/SO(2, 2)^4$, $SO(2, 2)^4/SO(1, 1)^8$.

<i>Field</i>	<i>f</i>
g_{MN}	44
ψ_M	128
A_{MNP}	84
<i>total f</i>	256

TABLE II: D=11 fields

II. MINIMAL SUPERGRAVITIES IN $D = 4, 5, 7, 11$

A. Compactifications

To derive the $D = 4, 5, 7, 11$ theories we begin with compactification on generic manifolds, tori and manifolds of special holonomy as shown in Table II, Table III, Table IV and Table V

	<i>Field</i>	<i>f</i>	<i>generic</i>	<i>torus</i>	<i>special</i>
g_{MN}	$g_{\mu\nu}$	14	d_0	1	1
	\mathcal{A}_μ	5	d_1	4	0
	\mathcal{A}	1	$-8d_0 + 3d_2$	10	10
ψ_M	ψ_μ	16	$2d_0 + d_1/2$	4	2
	χ	4	$-4d_0 + 2d_1 + 2d_2$	16	8
A_{MNP}	$A_{\mu\nu\rho}$	10	d_0	1	1
	$A_{\mu\nu}$	10	d_1	4	0
	A_μ	5	d_2	6	6
	A	1	d_1	4	0
<i>total f</i>		$16(2d_0 + 2d_1 + d_2)$		256	128
$\chi(X^4)$		$2d_0 - 2d_1 + d_2$		0	8

TABLE III: Compactify to $D = 7$ on X^4 with betti numbers: generic ($d_0 = d_4 = 1; d_1 = d_3; d_2$); torus (1; 4; 6) and $SU(2)$ holonomy (1; 0; 6).

	<i>Field</i>	<i>f</i>	<i>generic</i>	<i>torus</i>	<i>special</i>
g_{MN}	$g_{\mu\nu}$	5	c_0	1	1
	\mathcal{A}_μ	3	c_1	6	0
	\mathcal{A}	1	$-2c_0 - 2c_1 + c_2 + c_3$	21	9
ψ_M	ψ_μ	4	$2c_0 + c_1$	8	2
	χ	2	$-2c_0 + 2c_2 + c_3$	48	12
A_{MNP}	$A_{\mu\nu\rho}$	1	c_0	1	1
	$A_{\mu\nu}$	3	c_1	6	0
	A_μ	3	c_2	15	3
	A	1	c_3	20	8
$total\ f$		$4(2c_0 + 2c_1 + 2c_2 + c_3)$		256	64
$\chi(X^6)$		$2c_0 - 2c_1 + 2c_2 - c_3$		0	0

TABLE IV: Compactify to $D = 5$ on X^6 with betti numbers: generic ($c_0 = c_6 = 1; c_1 = c_5; c_2 = c_4; c_3$), torus (1; 6; 15; 20) and $SU(3)$ holonomy (1; 0; 3; 8)

Field f			generic torus special		
g_{MN}	$g_{\mu\nu}$	2	b_0	1	1
	\mathcal{A}_μ	2	b_1	7	0
	\mathcal{A}	1	$-b_1 + b_3$	28	7
ψ_M	ψ_μ	2	$b_0 + b_1$	8	1
	χ	2	$b_2 + b_3$	56	7
A_{MNP}	$A_{\mu\nu\rho}$	0	b_0	1	1
	$A_{\mu\nu}$	1	b_1	7	0
	A_μ	2	b_2	21	0
	A	1	b_3	35	7
$total\ f$		$4(b_0 + b_1 + b_2 + b_3)$	256	32	
$\rho(X^7)$		$7b_0 - 5b_1 + 3b_2 - b_3$	0	0	

TABLE V: Compactify to $D = 4$ on X^7 with betti numbers: generic ($b_0 = b_7 = 1; b_1 = b_6; b_2 = b_5, b_3 = b_4$), torus (1; 7; 21; 35) and G_2 holonomy (1; 0; 0; 7)

B. Supermultiplets

Here we group the individual fields into supermultiplets:

$$N = 1 \quad multiplet \quad f$$

$$graviton \ (g_{MN}; \psi_M; A_{MNP}) \ 128 + 128$$

TABLE VI: The $D = 11$ multiplet in the minimal $\mathcal{N} = 1$ basis

$N = 2$	$multiplet$	f	$N = 2d_0 + d_1/2$	$N = 4$	$N = 2$
$graviton$	$(g_{\mu\nu}, \mathcal{A}; 2\psi_\mu, 2\chi; A_{\mu\nu\rho}, 3A_\mu)$	$40 + 40$	d_0	1	1
$gravitino$	$(4A_\mu; 2\psi_\mu, 8\chi; 4A_{\mu\nu}, 4A)$	$64 + 64$	$d_1/4$	1	0
$vector$	$(3\mathcal{A}; 2\chi; A_\mu)$	$8 + 8$	$-3d_0 + d_2$	3	3

TABLE VII: The $D = 7$ multiplets in the minimal $\mathcal{N} = 2$ basis

$N = 2$	<i>multiplet</i>	f	$N = 2c_0 + c_1$	$N = 8$	$N = 2$
<i>graviton</i>	$(g_{\mu\nu}; 2\psi_\mu; A_\mu)$	$8 + 8$	c_0	1	1
<i>gravitino</i>	$(2\mathcal{A}_\mu; 2\psi_\mu, 2\chi; 2A_\mu)$	$12 + 12$	$c_1/2$	3	0
<i>vector</i>	$(\mathcal{A}; 2\chi; A_\mu)$	$4 + 4$	$-c_0 - c_1 + c_2$	8	2
<i>hyper</i>	$(2\mathcal{A}; 2\chi; 2A)$	$4 + 4$	$-c_0 - c_1/2 + c_3/2$	6	3
<i>2-form</i>	$(2\chi; A_{\mu\nu}, A)$	$4 + 4$	c_1	6	0
<i>3-form</i>	$(\mathcal{A}; 2\chi; A_{\mu\nu\rho}, 2A)$	$4 + 4$	c_0	1	1

TABLE VIII: The $D = 5$ multiplets in the minimal $\mathcal{N} = 2$ basis

$N = 1$	<i>multiplet</i>	f	$N = b_0 + b_1$	$N = 8$	$N = 1$
<i>graviton</i>	$(g_{\mu\nu}; \psi_\mu; A_{\mu\nu\rho})$	$2 + 2$	b_0	1	1
<i>gravitino</i>	$(\mathcal{A}_\mu; \psi_\mu)$	$2 + 2$	b_1	7	0
<i>vector</i>	$(\chi; A_\mu)$	$2 + 2$	b_2	21	7
<i>WZ</i>	$(\mathcal{A}; \chi; A)$	$2 + 2$	$-b_1 + b_3$	28	7
<i>linear</i>	$(\chi; A_{\mu\nu}, A)$	$2 + 2$	b_1	1	1

TABLE IX: The $D = 4$ multiplets in the minimal $\mathcal{N} = 1$ basis

C. Lagrangians

The bosonic sector of the toroidally compactified $D = 11$ supergravity prior to dualization may be found in [38, 39]. It will be useful to split the metric scalars \mathcal{A} into $\vec{\phi}$, the $(11 - D)$ -vector of dilatonic scalar fields coming from the diagonal components of the internal metric, and the rest, which we continue to describe by the letter \mathcal{A} . The original eleven-dimensional fields g_{MN} and A_{MNP} will give then rise to the following fields in D dimensions,

$$\begin{aligned} g_{MN} &\rightarrow g_{\mu\nu}, & \vec{\phi}, & \mathcal{A}_\mu^i, & \mathcal{A}^i_j \\ A_{MNP} &\rightarrow A_{\mu\nu\rho}, & A_{\mu\nu k}, & A_{\mu j k}, & A_{i j k} \end{aligned} \quad (8)$$

where the indices i, j, k run over the $(11 - D)$ internal toroidally-compactified dimensions. If we denote the rank $p + 1$ field strengths of the rank p potentials by a subscript $(p + 1)$, the Lagrangian is

$$\begin{aligned} \frac{\mathcal{L}}{\sqrt{-g}} = & R - \frac{1}{2} (\partial\vec{\phi})^2 - \frac{1}{48} e^{\vec{a}\cdot\vec{\phi}} F_{(4)}^2 - \frac{1}{12} \sum_i e^{\vec{a}_i\cdot\vec{\phi}} (F_{(3)i})^2 - \frac{1}{4} \sum_{i < j} e^{\vec{a}_{ij}\cdot\vec{\phi}} (F_{(2)ij})^2 \\ & - \frac{1}{4} \sum_i e^{\vec{b}_i\cdot\vec{\phi}} (\mathcal{F}_{(2)}^i)^2 - \frac{1}{2} \sum_{i < j < k} e^{\vec{a}_{ijk}\cdot\vec{\phi}} (F_{(1)ijk})^2 - \frac{1}{2} \sum_{i < j} e^{\vec{b}_{ij}\cdot\vec{\phi}} (\mathcal{F}_{(1)j}^i)^2 + \mathcal{L}_{FFA} \end{aligned} \quad (9)$$

where the “dilaton vectors” \vec{a} , \vec{a}_i , \vec{a}_{ij} , \vec{a}_{ijk} , \vec{b}_i , \vec{b}_{ij} are constants that characterise the couplings of the dilatonic scalars $\vec{\phi}$ to the various gauge fields [40]

$$F_4 : \quad \vec{a} = -\vec{g} \quad (10)$$

$$F_3 : \quad \vec{a}_i = \vec{f}_i - \vec{g} \quad (11)$$

$$F_2 : \quad \vec{a}_{ij} = \vec{f}_i + \vec{f}_j - \vec{g} \quad (12)$$

$$F_1 : \quad \vec{a}_{ijk} = \vec{f}_i + \vec{f}_j + \vec{f}_k - \vec{g} \quad (13)$$

$$\mathcal{F}_2 : \quad \vec{b}_i = -\vec{f}_i \quad (14)$$

$$\mathcal{F}_1 : \quad \vec{b}_{ij} = -\vec{f}_i + \vec{f}_j \quad (15)$$

$$(16)$$

where the vectors \vec{g} and \vec{f}_i have $(11 - D)$ components in D dimensions, and are given by

$$\begin{aligned}\vec{g} &= 3(s_1, s_2, \dots, s_{11-D}), \\ \vec{f}_i &= \left(\underbrace{0, 0, \dots, 0}_{(10-i)}, (10-i)s_i, s_{i+1}, s_{i+2}, \dots, s_{11-D} \right)\end{aligned}\quad (17)$$

where $s_i = \sqrt{2/((10-i)(9-i))}$. Note that the 4-dimensional metric is related to the eleven-dimensional one by

$$ds_{11}^2 = e^{\frac{1}{3}\vec{g}\cdot\vec{\phi}} ds_4^2 + \sum_i e^{2\vec{\gamma}_i\cdot\vec{\phi}} (h^i)^2 \quad (18)$$

where

$$\vec{\gamma}_i = \frac{1}{6}\vec{g} - \frac{1}{2}\vec{f}_i \quad (19)$$

and

$$h^i = dz^i + \mathcal{A}^i + \mathcal{A}^i_j dz^j \quad (20)$$

In general, the field strengths appearing in the kinetic terms are not simply the exterior derivatives of their associated potentials, but have non-linear Kaluza-Klein modifications as well. On the other hand the terms included in \mathcal{L}_{FFA} , which denotes the dimensional reduction of the $F_{(4)} \wedge F_{(4)} \wedge A_{(3)}$ term in $D = 11$, are best expressed purely in terms of the potentials and their exterior derivatives. The complete details may be found in [40], where it is shown that the symmetry of the Lagrangian is

$$GL(11 - D, R) \ltimes R^q \quad (21)$$

with

$$q = \frac{1}{6}(11 - D)(10 - D)(9 - D) \quad (22)$$

Our *minimal* Lagrangians in $D = 7, 5, 4$ follow by appropriate truncations that nevertheless keep all the $\vec{\phi}$. We shall not show these explicitly.

D. Membrane solutions

According to [4, 40, 41], the existence of an elementary membrane solution in D dimensions requires a metric, 3-form potential and dilaton described by the action

$$\frac{\mathcal{L}}{\sqrt{-g}} = R - \frac{1}{2}(\partial\phi)^2 - \frac{1}{48}e^{a\phi}F_{(4)}^2 \quad (23)$$

Moreover the dilaton coupling must be such that

$$a^2 = \frac{2(11 - D)}{(D - 2)} \quad (24)$$

or $a = 0, 8/5, 4, 7$ in $D = 11, 7, 5, 4$. But if we start with

$$\frac{\mathcal{L}}{\sqrt{-g}} = R - \frac{1}{2}(\partial\vec{\phi})^2 - \frac{1}{48}e^{\vec{a}\cdot\vec{\phi}}F_{(4)}^2 \quad (25)$$

and make the ansatz

$$a\vec{\phi} = \vec{a}\phi \quad (26)$$

noting that

$$\vec{a}\cdot\vec{a} = \frac{2(11 - D)}{(D - 2)} = a^2 \quad (27)$$

then the two Lagrangians coincide. Note that this ansatz would not have worked had we failed to implement the maximum rank condition by omitting some of the components of $\vec{\phi}$.

E. Cosets

The scalar cosets, before and after dualization are shown in Table X and XI

<i>theory</i>	<i>charges</i>	<i>G/H</i>	<i>dim</i>	<i>rank</i>	<i>max G/H</i>	<i>dim</i>	<i>rank</i>
$D = 11$	32	0	0	0	$\subset 0$	0	0
$D = 7$	16	$SO(1,1) \times SL(4,R)/SO(4)$	10	4	$\subset SL(5,R)/SO(5)$	14	4
$D = 5$	8	$SO(1,1)^3 \times SO(4,3)/[SO(4) \times SO(3)] \ltimes R^2$	17	6	$\subset SO(1,1) \times SL(6,R)/SO(6) \ltimes R^{20}$	41	6
$D = 4$	4	$SL(2,R)^7/SO(2)^7$	14	7	$\subset SO(1,1) \times SL(7,R)/SO(7) \ltimes R^{35}$	63	7

TABLE X: $D = 4, 5, 7, 11$ cosets before dualization

<i>theory</i>	<i>charges</i>	<i>G/H</i>	<i>dim</i>	<i>rank</i>	<i>max G/H</i>	<i>dim</i>	<i>rank</i>
$D = 11$	32	0	0	0	$\subset 0$	0	0
$D = 7$	16	$SO(1,1) \times SL(4,R)/SO(4)$	10	4	$\subset SL(5,R)/SO(5)$	14	4
$D = 5$	8	$SO(1,1)^2 \times SO(4,4)/SO(4)^2$	18	6	$\subset E_{6(6)}/Usp(8)$	42	6
$D = 4$	4	$SL(2,R)^7/SO(2)^7$	14	7	$\subset E_{7(7)}/SU(8)$	70	7

TABLE XI: $D = 4, 5, 7, 11$ cosets after dualization

III. $\mathcal{N} = 1, 2, 4, 8$ IN $D = 4$

A. Betti numbers

These four theories may be further reduced to $\mathcal{N} = 1, 2, 4$ and 8 theories all with maximum rank $r = 7$ in $D = 4$, corresponding to compactification on $X^{(8-\mathcal{N})} \times T^{(\mathcal{N}-1)}$. Denote the betti numbers of X^7, X^6, X^4 by b, c, d , respectively. The betti numbers of S^1 are $(1, 1)$, of T^3 are $(1, 3, 3, 1)$, of T^4 are $(1, 4, 6, 4, 1)$, of T^7 are $(1, 7, 21, 35, 21, 7, 1)$, so we have

$$\begin{aligned}
X^7 &: (b_0, b_1, b_2, b_3) \\
X^6 \times S^1 &: (c_0, c_0 + c_1, c_1 + c_2, c_2 + c_3) \\
X^4 \times T^3 &: (d_0, 3d_0 + d_1, 3d_0 + 3d_1 + d_2, d_0 + 4d_1 + 3d_2)
\end{aligned} \tag{28}$$

The number of fields in $D = 4$ is given by Table XII.

<i>Field</i>	<i>f</i>	360A	X^7	$X^6 \times S^1$	$X^4 \times T^3$	T^7
g_{MN}	$g_{\mu\nu}$	2 848	b_0	c_0	d_0	1
	\mathcal{A}_μ	2 -52	b_1	$c_0 + c_1$	$3d_0 + d_1$	7
	\mathcal{A}	1 4	$-b_1 + b_3$	$-c_0 - c_1 + c_2 + c_3$	$-2d_0 + 3d_1 + 3d_2$	28
ψ_M	ψ_μ	2 -233	$b_0 + b_1$	$2c_0 + c_1$	$4d_0 + d_1$	8
	χ	2 7	$b_2 + b_3$	$c_1 + 2c_2 + c_3$	$4d_0 + 7d_1 + 4d_2$	56
A_{MNP}	$A_{\mu\nu\rho}$	0 -720	b_0	d_0	c_0	1
	$A_{\mu\nu}$	2 364	b_1	$c_0 + c_1$	$3d_0 + d_1$	7
	A_μ	2 -52	b_2	$c_1 + c_2$	$3d_0 + 3d_1 + d_2$	21
	A	1 4	b_3	$c_2 + c_3$	$d_0 + 4d_1 + 3d_2$	35

$$A = -\rho/24 \quad A = -\chi/24 \quad A = 0 \quad A = 0$$

TABLE XII: $X^7, X^6 \times S^1, X^4 \times T^3, T^7$ compactification of D=11 supergravity.

B. Self-mirror with vanishing trace anomaly

Finally, we note that in [28] we defined a generalized mirror symmetry

$$(b_0, b_1, b_2, b_3) \rightarrow (b_0, b_1, b_2 - \rho/2, b_3 + \rho/2) \tag{29}$$

under which

$$\rho \equiv 7b_0 - 5b_1 + 3b_2 + b_3 \quad (30)$$

changes sign

$$\rho \rightarrow -\rho \quad (31)$$

Moreover the quantity ρ also shows up in the on-shell trace anomaly (before dualization), which is given by

$$g_{\mu\nu} \langle T^{\mu\nu} \rangle = A \frac{1}{32\pi^2} R^{*\mu\nu\rho\sigma} R^*_{\mu\nu\rho\sigma} \quad (32)$$

The value of the A coefficients for each field is given in Table XII that shows compactification on X^7 , $X^6 \times S^1$, $X^4 \times T^3$ and T^7 . We adopt the interpretation of [33] that assigns different anomalies to $A_{\mu\nu}$ and \mathcal{A} even though they are naively dual to one another and nonzero anomaly to $A_{\mu\nu\rho}$. Remarkably, we find that the total anomaly depends on ρ

$$A = -\frac{1}{24}\rho \quad (33)$$

So the anomaly flips sign under generalized mirror symmetry and vanishes for generalized self-mirror theories.

In the case of ($\mathcal{N} = 1, D = 11$) on $X^6 \times S^1$, or equivalently (Type IIA, D=10) on X^6 ,

$$A = -\frac{1}{24}\chi \quad (34)$$

where χ is the Euler number of X^6 .

Here we group the individual fields into supermultiplets as shown in Tables XIII to XVI.

$\mathcal{N} = 1$	<i>multiplet</i>	f	$360A$	$\mathcal{N} = b_0 + b_1$	$\mathcal{N} = 8$	$\mathcal{N} = 1$
<i>graviton</i>	$(g_{\mu\nu}; \psi_\mu; A_{\mu\nu\rho})$	$2 + 2$	-105	b_0	1	1
<i>gravitino</i>	$(\mathcal{A}_\mu; \psi_\mu)$	$2 + 2$	-285	b_1	7	0
<i>vector</i>	$(\chi; A_\mu)$	$2 + 2$	-45	b_2	21	0
<i>WZ</i>	$(\mathcal{A}; \chi; A)$	$2 + 2$	15	$-b_1 + b_3$	28	7
<i>linear</i>	$(\chi; A_{\mu\nu}, A)$	$2 + 2$	375	b_1	7	0
<i>total f</i>				$4(b_0 + b_1 + b_2 + b_3)$	256	32
<i>total A</i>				$-(7b_0 - 5b_1 + 3b_2 - b_3)/24$	0	0

TABLE XIII: The $D = 4$ multiplets in an $\mathcal{N}=1$ basis.

$\mathcal{N} = 2$	<i>multiplet</i>	f	$360A$	$\mathcal{N} = 2c_0 + c_1$	$\mathcal{N} = 8$	$\mathcal{N} = 2$
<i>graviton</i>	$(g_{\mu\nu}, \mathcal{A}_\mu; 2\psi_\mu, A_{\mu\nu\rho})$	$4 + 4$	-390	c_0	1	1
<i>gravitino</i>	$(\mathcal{A}_\mu; \psi_\mu, \chi; A_\mu)$	$4 + 4$	-330	c_1	6	0
<i>vector</i>	$(\mathcal{A}, 2\chi; A_\mu, A)$	$4 + 4$	-30	c_2	15	3
<i>hyper</i>	$(2\mathcal{A}; 2\chi; 2A)$	$4 + 4$	30	$-c_0 - c_1 + c_3/2$	3	3
<i>linear</i>	$(\mathcal{A}; 2\chi; A_{\mu\nu}, 2A)$	$4 + 4$	390	$c_0 + c_1$	7	1
<i>total f</i>				$4(2c_0 + 2c_1 + 2c_2 + c_3)$	256	64
<i>total A</i>				$-(2c_0 - 2c_1 + 2c_2 - c_3)/24$	0	0

TABLE XIV: The $D = 4$ multiplets in an $\mathcal{N}=2$ basis.

$\mathcal{N} = 4$	<i>multiplet</i>	f	$360A$	$\mathcal{N} = 4d_0 + d_1$	$\mathcal{N} = 8$	$\mathcal{N} = 4$
<i>graviton</i>	$(g_{\mu\nu}, 3\mathcal{A}_\mu, \mathcal{A}, 4\psi_\mu, 4\chi, A_{\mu\nu\rho}, 3A_\mu, A)$	$16 + 16$	-1080	d_0	1	1
<i>gravitino</i>	$(\mathcal{A}_\mu, 3\mathcal{A}, \psi_\mu, 7\chi, A_{\mu\nu}, 3A_\mu, 4A)$	$16 + 16$	0	d_1	4	0
<i>vector</i>	$(3\mathcal{A}; 4\chi; A_\mu, 3A)$	$8 + 8$	0	$-3d_0 + d_2$	3	3
<i>2-form</i>	$(2\mathcal{A}; 4\chi; A_{\mu\nu}, A_\mu, 3A)$	$8 + 8$	360	$3d_0$	3	3
<i>total f</i>				$16(2d_0 + 2d_1 + d_2)$	256	128
<i>total A</i>				0	0	0

TABLE XV: The $D = 4$ multiplets in an $\mathcal{N}=4$ basis.

$\mathcal{N} = 8$	<i>multiplet</i>	f	$360A$	$\mathcal{N} = 8$
<i>graviton</i>	$(g_{\mu\nu}, 7\mathcal{A}_\mu, 28\mathcal{A}; 8\psi_\mu, 56\chi; A_{\mu\nu\rho}, 7A_{\mu\nu}, 21A_\mu, 35A)$	256	0	1
<i>total f</i>				256
<i>total A</i>				0

TABLE XVI: The $D = 4$ multiplets in an $\mathcal{N}=8$ basis.

C. Cosets

The $D = 4$ scalar cosets, before and after dualization are given in Tables XVII and XVIII

<i>theory</i>	<i>charges</i>	G/H	<i>dim</i>	<i>rank</i>
$N = 8$	32	$SO(1, 1) \times SL(7, R)/SO(7) \ltimes R^{35}$	63	7
$N = 4$	16	$SL(2)/SO(2) \times SO(6, 3)/[SO(6) \times SO(3)] \times SL(3, R)/SO(3) \ltimes R^9$	35	7
$N = 2$	8	$SO(1, 1) \times SL(2)^3/SO(2)^3 \times SO(4, 3)/[SO(4) \times SO(3)] \ltimes R^2$	21	7
$N = 1$	4	$SL(2, R)^7/SO(2)^7$	14	7

TABLE XVII: $D = 4$ cosets before dualization

<i>theory</i>	<i>charges</i>	<i>G/H</i>	<i>dim</i>	<i>rank</i>
$N = 8$	32	$E_7/SU(8)$	70	7
$N = 4$	16	$SL(2)/SO(2) \times SO(6,6)/SO(6)^2$	38	7
$N = 2$	8	$SL(2)^3/SO(2)^3 \times SO(4,4)/SO(4)^2$	22	7
$N = 1$	4	$SL(2, R)^7/SO(2)^7$	14	7

TABLE XVIII: $D=4$ cosets after dualization

D. Fano plane and $\mathbb{O}, \mathbb{C}, \mathbb{H}, \mathbb{R}$




FIG. 1: The Fano plane has seven points and seven lines (the circle counts as a line) with three points on every line and three lines through every point. The points A, B, C, D, E, F, G represent the seven qubits and the seven lines $ABD, BCE, CDF, DEG, EFA, FGB, GAC$ represent the tripartite entanglement.

Next we turn to the black-hole/qubit correspondence [19–24]. The number of electric and magnetic black hole charges of these $\mathcal{N} = 8, 4, 2, 1$ theories are 56, 24, 8, 0, respectively. These correspond to 7, 3, 1, 0 lines of the Fano plane of Fig 1, which in turn admit an interpretation in terms of entangled qubits, as may be seen by writing them all in an $SL(2)^7$ basis:

- $\mathcal{N} = 8$

$$E_{7(7)} \supset SL(2)_A \times SL(2)_B \times SL(2)_C \times SL(2)_D \times SL(2)_E \times SL(2)_F \times SL(2)_G, \quad (35)$$

and the **56** decomposes as

$$\begin{aligned}
\mathbf{56} \rightarrow & (\mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}) \\
& + (\mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}) \\
& + (\mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}) \\
& + (\mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}) \\
& + (\mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}) \\
& + (\mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{2}) \\
& + (\mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}).
\end{aligned} \quad (36)$$

corresponding to the seven lines of the Fano plane describing a tripartite entanglement of seven qubits (Alice, Bob, Charlie, Daisy, Emma, Fred and George):

$$\begin{aligned}
|\psi\rangle_{56} = & a_{ABD}|ABD\rangle \\
& + b_{BCE}|BCE\rangle \\
& + c_{CDF}|CDF\rangle \\
& + d_{DEG}|DEG\rangle \\
& + e_{EFA}|EFA\rangle \\
& + f_{FGB}|FGB\rangle \\
& + g_{GAC}|GAC\rangle
\end{aligned} \tag{37}$$

• $\mathcal{N} = 4$

$$SL(2)_A \times SO(6,6) \supset SL(2)_A \times SL(2)_B \times SL(2)_C \times SL(2)_D \times SL(2)_E \times SL(2)_F \times SL(2)_G, \tag{38}$$

and the $(\mathbf{2}, \mathbf{12})$ decomposes as

$$\begin{aligned}
(\mathbf{2}, \mathbf{12}) \rightarrow & (\mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}) \\
& + (\mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{2}, \mathbf{1}) \\
& + (\mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{2}).
\end{aligned} \tag{39}$$

corresponding to the three lines of the Fano plane describing a tripartite entanglement of Alice with Bob and Daisy, Alice with Emma and Fred, Alice with Charlie and George:

$$\begin{aligned}
|\psi\rangle_{24} = & a_{ABD}|ABD\rangle \\
& + e_{EFA}|EFA\rangle \\
& + g_{GAC}|GAC\rangle
\end{aligned} \tag{40}$$

• $\mathcal{N} = 2$

$$SL(2)_A \times SL(2)_B \times SL(2)_D \times SO(4,4) \supset SL(2)_A \times SL(2)_B \times SL(2)_C \times SL(2)_D \times SL(2)_E \times SL(2)_F \times SL(2)_G, \tag{41}$$

and the $(\mathbf{2}, \mathbf{2}, \mathbf{2}, \mathbf{1})$ decomposes as

$$(\mathbf{2}, \mathbf{2}, \mathbf{2}, \mathbf{1}) \rightarrow (\mathbf{2}, \mathbf{2}, \mathbf{1}, \mathbf{2}, \mathbf{1}, \mathbf{1}, \mathbf{1}) \tag{42}$$

corresponding to the one line of the Fano plane describing a tripartite entanglement of three qubits, Alice, Bob and Daisy:

$$|\psi\rangle_8 = a_{ABD}|ABD\rangle \tag{43}$$

• $\mathcal{N} = 1$

$$\begin{aligned}
& SL(2)_A \times SL(2)_B \times SL(2)_C \times SL(2)_D \times SL(2)_E \times SL(2)_F \times SL(2)_G \\
& \supset SL(2)_A \times SL(2)_B \times SL(2)_C \times SL(2)_D \times SL(2)_E \times SL(2)_F \times SL(2)_G,
\end{aligned} \tag{44}$$

$$(\mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}) \rightarrow (\mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}) \tag{45}$$

corresponding to no lines.

The black hole entropies are given by the qubit entanglement measures which are quartic polynomials in the a, b, c, d, e, f, g coefficients, namely Cartan's E_7 invariant, the analogous $SL(2) \times SO(6,6)$ invariant and Cayley's $SL(2)^3$ hyperdeterminant.

Since the 7, 3, 1, 0 lines of the Fano plane also describe the multiplication table of the octonions, quaternions, complex and real, it was conjectured in [24] that there is an $\mathbb{O}, \mathbb{H}, \mathbb{C}, \mathbb{R}$ interpretation not just for the charges but for the entire theories. Their field content and trace anomalies are given in Tables XIX and XX.

<i>Field</i>	<i>f</i>	360 <i>A</i>	\mathbb{A}	\mathbb{O}	\mathbb{H}	\mathbb{C}	\mathbb{R}
$g_{\mu\nu}$	2	848	1	1	1	1	1
\mathcal{A}_μ	2	-52	$\mathcal{N}-1$	7	3	1	0
$\vec{\phi}$	1	4	7	7	7	7	7
\mathcal{A}	1	4	$3(\mathcal{N}-1)$	21	9	3	0
ψ_μ	2	-233	\mathcal{N}	8	4	2	1
χ	2	7	$7\mathcal{N}$	56	28	14	7
$A_{\mu\nu\rho}$	0	-720	1	1	1	1	1
$A_{\mu\nu}$	1	364	$\mathcal{N}-1$	7	3	1	0
A_μ	2	-52	$3(\mathcal{N}-1)$	21	9	3	0
A	1	4	$4\mathcal{N}+3$	35	19	11	7
<i>total f</i>			$32\mathcal{N}$	256	128	64	32
<i>total A</i>			0	0	0	0	0

TABLE XIX: Vanishing anomaly in \mathbb{O} , \mathbb{H} , \mathbb{C} \mathbb{R} theories.

$\mathcal{N}=1$	<i>multiplet</i>	<i>f</i>	360 <i>A</i>	\mathbb{A}	\mathbb{O}	\mathbb{H}	\mathbb{C}	\mathbb{R}
<i>graviton</i>	$(g_{\mu\nu}; \psi_\mu; A_{\mu\nu\rho})$	2+2	-105	1	1	1	1	1
<i>gravitino</i>	$(\mathcal{A}_\mu; \psi_\mu)$	2+2	-285	$\mathcal{N}-1$	7	3	1	0
<i>vector</i>	$(\chi; A_\mu)$	2+2	-45	$3(\mathcal{N}-1)$	21	9	3	0
WZ_ϕ	$(\vec{\phi}; \chi; A)$	2+2	15	7	7	7	7	7
WZ_A	$(\mathcal{A}; \chi; A)$	2+2	15	$3(\mathcal{N}-1)$	21	9	3	0
<i>linear</i>	$(\chi; A_{\mu\nu}, A)$	2+2	375	$\mathcal{N}-1$	7	3	1	0
<i>total f</i>				$32\mathcal{N}$	256	128	64	32
<i>total A</i>				0	0	0	0	0

TABLE XX: The $D=4$ multiplets in an $\mathcal{N}=1$ basis from X^7 with $(b_0, b_1, b_2, b_3) = (1, \mathcal{N}-1, 3\mathcal{N}-3, 4\mathcal{N}+3)$

IV. $\mathcal{N}=2, 4, 8, 16$ IN $D=3$

A. Compactifications

Consider Type IIA in $D=10$. In the NS sector we have the fields $(g_{MN}, \Phi; \psi_M, \chi; A_{MN})$ with $f=64+64$; in the R-R we have the fields $(\mathcal{A}_M; \psi_M, \chi; A_{MNP})$ also with $f=64+64$. We compactify on generic X^7 with independent betti numbers (b_0, b_1, b_2, b_3) , $X^6 \times S^1$ with independent X^6 betti numbers (c_0, c_1, c_2, c_3) , $X^4 \times S^3$ with independent X^4 betti numbers (d_0, d_1, d_2) and on T^7 with $(1, 7, 21, 35)$. The results for NS and RR combined are shown in Table XXI. In Table XXII, we group into $\mathcal{N}=2$ multiplets.

Field f			X^7	$X^6 \times S^1$	$X^4 \times T^3$	T^7
g_{MN}	$g_{\mu\nu}$	0	b_0	c_0	d_0	1
	\mathcal{A}_μ	1	$b_0 + b_1$	$2c_0 + c_1$	$4d_0 + d_1$	8
	\mathcal{A}	1	b_3	$c_2 + c_3$	$d_0 + 4d_1 + 3d_2$	35
Φ	Φ	1	b_0	c_0	d_0	1
ψ_M	ψ_μ	0	$2b_0 + 2b_1$	$4c_0 + 2c_1$	$8d_0 + 2d_1$	16
	χ	1	$2b_0 + 2b_1 + 2b_2 + 2b_3$	$4c_0 + 4c_1 + 4c_2 + 2c_3$	$16d_0 + 16d_1 + 8d_2$	128
A_{MNP}	$A_{\mu\nu\rho}$	0	b_0	c_0	d_0	1
	$A_{\mu\nu}$	0	$b_0 + b_1$	$2c_0 + c_1$	$3d_0 + d_1$	8
	A_μ	1	$b_1 + b_2$	$c_0 + 2c_1 + c_2$	$6d_0 + 4d_1 + d_2$	28
	A	1	$b_2 + b_3$	$c_1 + 2c_2 + c_3$	$4d_0 + 7d_1 + 4d_2$	56
<i>total f</i>			$4(b_0 + b_1 + b_2 + b_3)$	$4(2c_0 + 2c_1 + 2c_2 + c_3)$	$16(2d_0 + 2d_1 + d_2)$	256

TABLE XXI: $X^7, X^6 \times S^1, X^4 \times T^3, T^7$ compactification of Type IIA

$\mathcal{N} = 2$ multiplet	content	f	$\mathcal{N} = 2b_0 + 2b_1$	$\mathcal{N} = 16$	$\mathcal{N} = 8$	$\mathcal{N} = 4$	$\mathcal{N} = 2$
graviton	$(g_{\mu\nu}, \mathcal{A}_\mu, \mathcal{A}; 2\psi_\mu, 2\chi; A_{\mu\nu\rho}, A_{\mu\nu})$	$2 + 2$	b_0	1	1	1	1
gravitino	$(\mathcal{A}_\mu, \mathcal{A}; 2\psi_\mu, 2\chi)$	$2 + 2$	b_1	7	3	1	0
vector	$(2\chi; A_\mu, A)$	$2 + 2$	b_2	21	9	3	0
hyper	$(\mathcal{A}; 2\chi; A)$	$2 + 2$	$-b_1 + b_3$	28	16	10	7
linear	$(2\chi; A_{\mu\nu}, A_\mu, A)$	$2 + 2$	b_1	7	3	1	0
<i>total f</i>			$4(b_0 + b_1 + b_2 + b_3)$	256	128	64	32

TABLE XXII: The $D = 3$ multiplets in an $\mathcal{N} = 2$ basis

B. Cosets

The $D = 3$ scalar cosets after dualization for spacelike and timelike reductions are given in Tables XXIII and XXIV

theory	G/H	dim	rank
$\mathcal{N} = 16$	$E_8/SO(16)$	128	8
$\mathcal{N} = 8$	$SO(8, 8)/SO(8)^2$	64	8
$\mathcal{N} = 4$	$SO(4, 4)^2/SO(4)^4$	32	8
$\mathcal{N} = 2$	$SL(2, R)^8/SO(2)^8$	16	8

TABLE XXIII: $D = 3$ cosets after dualization

theory	G/H	dim	rank
$\mathcal{N} = 16$	$E_8/SO^*(16)$	128	8
$\mathcal{N} = 8$	$SO(8, 8)/SO(4, 4)^2$	64	8
$\mathcal{N} = 4$	$SO(4, 4)^2/SO(2, 2)^4$	32	8
$\mathcal{N} = 2$	$SL(2, R)^8/SO(1, 1)^8$	16	8

TABLE XXIV: $D = 3$ cosets from timelike reduction

V. ACKNOWLEDGMENTS

Conversations and correspondence with Leron Borsten, Duminda Dahanayake, Phil Gibbs, Alessio Marrani, Chris Pope, William Rubens, Ashoke Sen, Ergin Sezgin, Samson Shatashvili, John Schwarz and Edward Witten are much appreciated. MJD is supported in part by the STFC under rolling Grant No. ST/G000743/1. S. F is supported by the ERC Advanced Grant no 226455, “Supersymmetry, Quantum Gravity and Gauge Fields” and in part by DOE Grant DE-FG03-91ER40662. MJD is grateful for hospitality at the CERN theory division, where he was supported by the above ERC Advanced Grant, and at the Mitchell Institute for Fundamental Physics and the Institute for Quantum Studies, Texas A&M University.

-
- [1] M. B. Green and J. H. Schwarz, Phys. Lett. **B136**, 367 (1984).
 - [2] A. Achucarro, J. M. Evans, P. K. Townsend, and D. L. Wiltshire, Phys. Lett. **B198**, 441 (1987).
 - [3] M. J. Duff and J. X. Lu, Nucl. Phys. **B390**, 276 (1993), hep-th/9207060.
 - [4] M. J. Duff, R. R. Khuri, and J. X. Lu, Phys. Rept. **259**, 213 (1995), hep-th/9412184.
 - [5] M. J. Duff, P. S. Howe, T. Inami, and K. S. Stelle, Phys. Lett. **B191**, 70 (1987).
 - [6] E. Bergshoeff, E. Sezgin, and P. K. Townsend, Phys. Lett. **B189**, 75 (1987).
 - [7] M. J. Duff and K. S. Stelle, Phys. Lett. **B253**, 113 (1991).
 - [8] E. Bergshoeff, E. Sezgin, and P. K. Townsend, Phys. Lett. **B169**, 191 (1986).
 - [9] B. A. Ovrut and D. Waldram, Nucl. Phys. **B506**, 236 (1997), hep-th/9704045.
 - [10] J. C. Baez and J. Huerta (2009), 0909.0551.
 - [11] J. C. Baez and J. Huerta (2010), 1003.3436.
 - [12] E. Cremmer, B. Julia, and J. Scherk, Phys. Lett. **B76**, 409 (1978).
 - [13] S. Ferrara and C. Kounnas, Nucl. Phys. **B328**, 406 (1989).
 - [14] A. Sen and C. Vafa, Nucl. Phys. **B455**, 165 (1995), hep-th/9508064.
 - [15] D. Joyce, J. Differential Geometry **43** (1996).
 - [16] D. Joyce, J. Differential Geometry **43**, 329 (1996).
 - [17] M. R. Gaberdiel and P. Kaste, JHEP **08**, 001 (2004), hep-th/0401125.
 - [18] R. D'Auria, L. Sommovigo, and S. Vaula, JHEP **11**, 028 (2004), hep-th/0409097.
 - [19] M. J. Duff, Phys. Rev. **D76**, 025017 (2007), hep-th/0601134.
 - [20] R. Kallosh and A. Linde, Phys. Rev. **D73**, 104033 (2006), hep-th/0602061.
 - [21] P. Lévy, Phys. Rev. **D74**, 024030 (2006), hep-th/0603136.
 - [22] M. J. Duff and S. Ferrara, Phys. Rev. **D76**, 025018 (2007), quant-ph/0609227.
 - [23] P. Lévy, Phys. Rev. **D75**, 024024 (2007), hep-th/0610314.
 - [24] L. Borsten, D. Dahanayake, M. J. Duff, H. Ebrahim, and W. Rubens, Phys. Rep. **471**, 113 (2009), 0809.4685.
 - [25] T. Kugo and P. K. Townsend, Nucl. Phys. **B221**, 357 (1983).
 - [26] J. M. Evans, Nucl. Phys. **B298**, 92 (1988).
 - [27] M. J. Duff, Class. Quant. Grav. **5**, 189 (1988).
 - [28] M. J. Duff and S. Ferrara (2010), 1009.4439.
 - [29] S. L. Shatashvili and C. Vafa, Selecta Math. **1**, 347 (1995), hep-th/9407025.
 - [30] B. S. Acharya, Nucl. Phys. **B524**, 269 (1998), hep-th/9707186.
 - [31] M. J. Duff, Nucl. Phys. **B125**, 334 (1977).
 - [32] M. J. Duff, Class. Quant. Grav. **11**, 1387 (1994), hep-th/9308075.
 - [33] M. J. Duff and P. van Nieuwenhuizen, Phys. Lett. **B94**, 179 (1980).
 - [34] P. Levay (2010), 1004.3639.
 - [35] L. Borsten, D. Dahanayake, M. J. Duff, A. Marrani, and W. Rubens (2010), 1005.4915.
 - [36] L. Borsten et al. (2010).
 - [37] P. Gibbs (2010), 1009.0076.
 - [38] E. Cremmer, B. Julia, H. Lu, and C. N. Pope, Nucl. Phys. **B523**, 73 (1998), hep-th/9710119.
 - [39] E. Cremmer, B. Julia, H. Lu, and C. N. Pope, Nucl. Phys. **B535**, 242 (1998), hep-th/9806106.
 - [40] H. Lu and C. N. Pope, Nucl. Phys. **B465**, 127 (1996), hep-th/9512012.
 - [41] I. V. Lavrinenko, H. Lu, and C. N. Pope, Nucl. Phys. **B492**, 278 (1997), hep-th/9611134.