

This is the accepted manuscript made available via CHORUS. The article has been published as:

Constraining the mixing matrix for the standard model with four generations: Time-dependent and semileptonic CP asymmetries in $B_{\{d\}}^{\{0\}}$, $B_{\{s\}}$, and $D^{\{0\}}$

Soumitra Nandi and Amarjit Soni

Phys. Rev. D **83**, 114510 — Published 16 June 2011

DOI: [10.1103/PhysRevD.83.114510](https://doi.org/10.1103/PhysRevD.83.114510)

Constraining the mixing matrix for Standard Model with four generations: time dependent and semi-leptonic CP asymmetries in B_d^0 , B_s and D^0

Soumitra Nandi¹ and Amarjit Soni²

¹*Physique des Particules, Université de Montréal,*

C.P. 6128, succ. centre-ville, Montréal, QC, Canada H3C 3J7

²*Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

Using existing experimental information from K, B and D decays as well as electroweak precision tests and oblique parameters, we provide constraints and correlations on the parameters of the 4X4 mixing matrix for the Standard Model (SM) with four generations (SM4). We emphasize that some correlations amongst the parameters have important repercussions for key observables. We work with a particular representation of this matrix which is highly suited for extracting information from B-decays. In particular, in our parameterization for SM4, we extend the hierarchical structure seen in SM with three generations (SM3) as an expansion in powers of λ , the sine of the Cabbibo angle. Implications of the resulting constraints for time dependent and semileptonic CP asymmetries for D^0 , B^0 and for B_s are also given. While we show that the semi-leptonic asymmetries may be significantly enhanced in SM4 over the SM, there are important constraints and correlations with other observables. In particular, we find that despite significant enhancement in the semi-leptonic asymmetry a_{sl}^s for B_s over the SM, it is very difficult for SM4 to account for the central value of the recent D0 result, though given the large experimental error and other considerations, we do not regard this as a problem for SM4. Regarding the gold-plated measurement of $\sin 2\beta$ via $S_{\psi K_s}$, while SM4 can remove the tension that SM shows, as a consequence of one of the important correlations we find that the semi-leptonic asymmetry a_{sl}^d for B_d gets appreciably restricted in SM4. In this context we suggest that existing data from B-factories taken on $\Upsilon(4S)$ and $\Upsilon(5S)$, and in the relevant continuum be used to constrain the semi-leptonic asymmetries for B_d , B_s as well as their linear combination. Of course, the data from the Tevatron and LHCb experiments can provide non-trivial tests of SM4 as well.

I. INTRODUCTION

In the past few years a number of tensions in the CKM fits for the Standard Model (SM) with 3 generations have been revealed [1–5]. There are quite serious indications that the “predicted” value of $\sin 2\beta$ is larger compared to the value measured directly via the “gold-plated” ψK_S mode by as much as $\approx 3.3\sigma$ [6]. Of course, the value of $\sin 2\beta$ determined from the penguin dominated modes tends to be even smaller compared to that from the ψK_s mode and therefore that constitutes even

a larger deviation from the SM predicted value [2]. There are other anomalies as well that appear related. The difference in the partial rate asymmetries between $B^0 \rightarrow K^+\pi^-$ and $B^+ \rightarrow K^+\pi^0$ is also too large [7] to understand [2], though QCD complications do not allow us to draw compelling conclusions in this regard [8]. But with the backdrop of the hint of presence of a new CP-odd phase in the $\Delta S = 1$ penguin dominated modes, it is highly suggestive that the direct CP problem in $K \pi$ modes is receiving, at least in part, contribution from the same new physics source.

There are also some indications from the CDF and DO experiments at the Tevatron [9]. While the earlier indication of possible non-standard effects in $B_s \rightarrow \psi\phi$ seem to have weakened somewhat at the higher luminosity around 6/fb now being used [10], D0 has announced a surprisingly large CP-asymmetry in the same sign dimuons which they attribute primarily to originate from $B_s \rightarrow X_s\mu\nu$ [11, 12]. From a theoretical standpoint if new physics exists in $\Delta S = 1$ B-decays, then it becomes highly unnatural for it not to exist in $\Delta S = 2$, B_s mixings as well.

A simple extension of the SM with four generations (SM4) can readily account for such anomalies [13–18]. Of course, even without these anomalies, SM4 is an interesting extension of the SM worth study. The two extra phases that it possesses can give rise to a host of non-standard CP asymmetries and in fact SM4 can significantly ameliorate the difficulties with regard to baryogenesis that SM3 has [19, 20]. Besides, the heavier quarks and leptons of the 4th generation may well lead to dynamical electroweak symmetry breaking and thereby become useful in addressing the hierarchy problem without the need for supersymmetry at the weak scale [21–25]. Motivated by these considerations we will continue our investigations of the physical implications of SM4. In particular we will use all the known experimental constraints such as $B \rightarrow X_s\gamma$, $B \rightarrow X_sl^+l^-$, ΔM_{B_s} , ΔM_{B_d} , $K^+ \rightarrow \pi\nu\nu$, $|\epsilon_k|$ and electroweak precision constraints from $Z \rightarrow b\bar{b}$ as well as oblique corrections [26, 27] as in our previous work [13, 14]. However, we will now use an explicit representation of the 4X4 CKM matrix of [28] given long ago. Although our basic idea to keep the elements of the fourth row as simple as possible is the same as in [28], but we are extending the hierarchical structure of SM3 [29] to SM4 in our parameterization. We make this particular choice as it is very well designed to extract constraints from B decays since it was shown in a series of papers [28, 30, 31] that SM4 is highly susceptible to those decays. While physical results should not depend on the parameterization used, we believe in practice this representation would lead to a better determination of the underlying SM4 parameters.

We will provide constraints and many correlations amongst the 6 real parameters and the 3 phases that enter the SM4. We will then apply this framework to study mixing induced CP asymmetries $S(B_d \rightarrow \psi K_s)$, $S(B_s \rightarrow \psi\phi)$, $S(D^0 \rightarrow f)$ (where f may be any self conjugate final

state such as $K_s\pi^0$, $K_s\omega$, $K_s\rho^0$, $\pi^0\pi^0$, K^+K^- , $\pi^+\pi^-$ etc.) and semi-leptonic asymmetries in D^0 , B^0 , and in B_s .

In obtaining these constraints and implications we will allow $m_{t'}$ to range from 375 to 575 GeV as suggested by current hints from study of B-decays [13, 14]. An interesting aspect of SM4 is that it is rather well constrained already. Thus, for example, while the semi-leptonic asymmetry in B_s (a_{sl}^s) can be enhanced by as much as a factor of few hundreds over SM3 it still cannot account for the central value of the recent D0 result [11]¹. Of course, that observation has only about 2- σ significance and therefore rather large errors but improved experimental results could certainly rule out or confirm SM4, since the predicted range² in SM4 for a_{sl}^s is between about (-0.005) to (0.005). Note also that in SM4 sign of a_{sl}^s has to be the same as $S_{\psi\phi}$. Furthermore, for B_d , a_{sl}^d can only be larger by around a factor of three over SM3. These semi-leptonic asymmetries also have interesting correlations with $S(B_s \rightarrow \psi\phi)$ and $S(B_d \rightarrow \psi K_s)$ respectively that should be testable.

As mentioned above one of the key difficulty for the CKM-paradigm of SM3 uncovered in recent years is that the predicted value of $\sin 2\beta$ is too large compared to the measured one [1, 6]. We will show here that SM4 tends to alleviate this tension appreciably but at the same time then it allows to place an important bound on a_{sl}^d through the correlation mentioned in the previous paragraph.

B-factories placed a bound on a_{sl}^d [10] some years ago but by now they have considerable more data. So an improved bound would be extremely worthwhile. In the past couple of years BELLE also took substantial data on Υ 5S [33]. In fact that data could provide a very clean study of a_{sl}^s as well as on A_{sl}^b , which is defined as the linear combination of a_{sl}^s and a_{sl}^d [12], since that sample provides a valuable source of this combination as well as an enriched sample of B_s . CDF, D0 and LHCb should be able to provide very useful results on these semi-leptonic asymmetries. In fact whereas the Tevatron $p\bar{p}$ collider allowed D0 to yield the sum of a_{sl}^d and a_{sl}^s , with the pp collider at LHC, the LHCb collaboration plans to study the difference of these two asymmetries [34].

We should emphasize that in this series of studies on the 4th generation [13, 14], for simplicity, and for definiteness, we have been making a tacit assumption that a heavy charge 2/3 and -1/3

¹ Actually the D0 experiment [11, 12] reports a measurement of the linear combination, A_{sl}^b , of the semi-leptonic asymmetry in B_d , a_{sl}^d and the one for B_s , a_{sl}^s . They find $A_{sl}^b = -0.00957 \pm 0.00251(stat) \pm 0.00146(syst)$ which is about 3.2σ away from the SM prediction, $A_{sl}^b(SM) = (-2.3_{-0.6}^{+0.5}) \times 10^{-4}$ [32]. Using the existing upper bound from B-factories [10], $a_{sl}^d = -0.0047 \pm 0.0046$, D0 provides $a_{sl}^s = -0.0146 \pm 0.0075$ which is just short of 2σ away from 0 and so also from the vanishingly small SM predicted value, $(2.1 \pm 0.6) \times 10^{-5}$ [32]. While the D0 result is extremely exciting it is a very challenging experiment, in particular because the B -mesons are not being identified or tagged. It is clearly extremely important to confirm their observation. For now we prefer to use the D0 results on these asymmetries with caution till confirmed and therefore we confine our comparison to their result on a_{sl}^s , wherein it is considerably diluted due to their use of the B -factory bound on a_{sl}^d .

² In arriving at this range, conservatively we have enlarged the theory error in $\frac{|\Gamma_{12}^s|}{|M_{12}^{s,SM}|}$ to 2σ .

quark doublet has weak interaction just like the previous three families allowing us to incorporate these readily into a 4X4 mixing-matrix resulting from an immediate generalization of the 3X3 case. Clearly if and when such a doublet of quarks is observed we will need to make detail tests on the weak interaction properties of the new quarks to verify that this assumption is correct.

The paper is arranged as follows. After the introduction, in Sec. IIA and IIB we provide information regarding the parametrisation and the constraints on the 4x4 CKM matrix by incorporating oblique corrections along with experimental data from important observables involving Z, B and K decays as well as B_d and B_s mixings etc. In Sec. IIC, we present the estimates of many useful observables in the SM4. Finally in Sec. III, we present our summary.

II. NUMERICAL ANALYSIS

A. Parametrisation of V_{CKM4}

We follow the idea of [28] in parameterizing the SM4 mixing matrix, then the elements of fourth row such as $V_{t'd}$, $V_{t's}$ and $V_{t'b}$, which are more relevant for the discussion of b physics, will be rather simple. However, our approach in parameterizing the elements of the SM4 mixing matrix is different from that one used in [28]. We parameterized the SM4 mixing matrix in terms of the nine parameters, λ , A , C , P , Q , r , δ_{ub} , $\delta_{t'd}$ and $\delta_{t's}$, by expanding each element of the matrix in powers of λ , the sine of the Cabbibo angle. In analogy with the Wolfenstein representation [29] for SM3 we assume a hierarchical structure for SM4 and define

$$\begin{aligned} V_{us} &= \lambda, & V_{cb} &= A\lambda^2, & V_{ub} &= A\lambda^3 C e^{-i\delta_{ub}}, \\ V_{t's} &= -Q\lambda^2 e^{i\delta_{t's}}, & V_{t'd} &= -P\lambda^3 e^{i\delta_{t'd}}, & V_{t'b} &= -r\lambda \end{aligned} \quad (1)$$

with the generalized 4×4 mixing matrix V_{SM4} as given in eq. 3. With the inputs $|V_{ub}| = (32.8 \pm 3.9) \times 10^{-4}$ and $|V_{cb}| = (40.86 \pm 1.0) \times 10^{-3}$ taken at 1σ , constraints obtained on A and C are given by

$$0.825 \leq A \leq 0.865, \quad 0.32 \leq C \leq 0.42, \quad (2)$$

while $\lambda = 0.2205 \pm 0.0018$. The phase of V_{ub} i.e δ_{ub} can be taken as the CKM angle γ of SM3. We do not have sufficient data to exclude the ranges like $P < 1$, $Q < 1$ and $r < 1$, therefore we treat them as free parameters and then constrain them.

$$\begin{pmatrix}
1 - \frac{\lambda^2}{2} + \mathcal{O}(\lambda^4) & \lambda & A\lambda^3 C e^{-i\delta_{ub}} & P\lambda^3 e^{-i\delta_{t'd}} \\
& & & + Q\lambda^3 e^{-i\delta_{t's}} + ACr\lambda^4 e^{-i\delta_{ub}} \\
& & & - P\frac{\lambda^5}{2} e^{-i\delta_{t'd}} + \mathcal{O}(\lambda^7) \\
-\lambda + \mathcal{O}(\lambda^5) & 1 - \frac{\lambda^2}{2} & A\lambda^2 & Q\lambda^2 e^{-i\delta_{t's}} \\
& & & + A\lambda^3 r - P\lambda^4 e^{-i\delta_{t'd}} \\
& & & - \frac{Q}{2}\lambda^4 e^{-i\delta_{t's}} + \mathcal{O}(\lambda^6) \\
A\lambda^3(1 - C e^{i\delta_{ub}}) & -A\lambda^2 - Qr\lambda^3 e^{i\delta_{t's}} & 1 - \frac{r^2\lambda^2}{2} + \mathcal{O}(\lambda^4) & r\lambda + \mathcal{O}(\lambda^4) \\
-P r\lambda^4 e^{i\delta_{t'd}} & + A\lambda^4(\frac{1}{2} - C e^{i\delta_{ub}}) & & \\
+ \frac{1}{2} A C \lambda^5 e^{i\delta_{ub}} + \mathcal{O}(\lambda^7) & + \mathcal{O}(\lambda^6) & & \\
-P\lambda^3 e^{i\delta_{t'd}} & -Q\lambda^2 e^{i\delta_{t's}} & -r\lambda & 1 - \frac{r^2\lambda^2}{2} + \mathcal{O}(\lambda^4)
\end{pmatrix}. \quad (3)$$

B. Inputs

In our earlier papers [13, 14], to find the limits on some of the V_{CKM4} elements, we concentrated mainly on the constraints that will come from non-decoupling oblique corrections, vertex correction to $Z \rightarrow b\bar{b}$, $\mathcal{BR}(B \rightarrow X_s \gamma)$, $\mathcal{BR}(B \rightarrow X_s l^+ l^-)$, $B_d - \bar{B}_d$ and $B_s - \bar{B}_s$ mixing, $\mathcal{BR}(K^+ \rightarrow \pi^+ \nu \nu)$ and the indirect CP violation in $K_L \rightarrow \pi\pi$ described by ϵ_k ; we did not consider ϵ'/ϵ as a constraint because of the large hadronic uncertainties, in the evaluation of its matrix elements. With the inputs given in Table I we have made the scan over the entire parameter space by a flat random number generator “RAN1(J)” where “J” is the number of iterations. As mentioned in the previous section we constrain the parameters A and C from the data on $|V_{cb}|$ and $|V_{ub}|$. $\Gamma(Z \rightarrow b\bar{b})$ and the non-decoupling oblique corrections such as T_4 are functions of $|V_{t'b}|^2$ and $|V_{tb}|^2$ or equivalently functions of r or higher powers of r , thus we get direct constraint on r from these data. All the other observables in Table I are functions of either $\lambda_{bs}^{t'} = V_{t'b} V_{t's}^*$ and $\lambda_{bs}^t = V_{tb} V_{ts}^*$ or $\lambda_{bd}^{t'} = V_{t'b} V_{t'd}^*$ and $\lambda_{bd}^t = V_{tb} V_{td}^*$ or both. Furthermore we can use unitarity relation, like $\lambda_{bq}^u + \lambda_{bq}^c + \lambda_{bq}^t + \lambda_{bq}^{t'} = 0$ with $q = d, s$, to reduce the number of independent elements, therefore it is possible to constrain the magnitudes and phases of these elements directly from the data as we did in our previous studies [13, 14]. However, in this paper, we express all these elements in terms of new SM4 parameters such as, $P, Q, r, \delta_{t'd}$ and $\delta_{t's}$ of the 4×4 mixing matrix eq. 3 or equivalently we express all the observables as functions of these parameters. We allow these parameters to vary randomly over a broad range and finally constrain them from the data. After satisfying all the constraints we get $\approx 8 \times 10^3$ data points in a multi dimensional parameter space for $J = 2 \times 10^9$.

$B_K = 0.740 \pm 0.025$ [41–43]	$R_{bb} = 0.216 \pm 0.001$
$f_{bd}\sqrt{B_{bd}} = 0.224 \pm 0.015$ GeV [44, 45]	$ V_{ub} = (32.8 \pm 2.6) \times 10^{-4}$ ^a
$\xi = 1.232 \pm 0.042$ [44, 45]	$ V_{cb} = (40.86 \pm 1.0) \times 10^{-3}$
$\eta_c = 1.51 \pm 0.24$ [46]	$\gamma = (73.0 \pm 13.0)^\circ$
$\eta_t = 0.5765 \pm 0.0065$ [47]	$\mathcal{BR}(B \rightarrow X_s \gamma) = (3.55 \pm 0.25) \times 10^{-4}$
$\eta_{ct} = 0.494 \pm 0.046$ [48]	$\mathcal{BR}(B \rightarrow X_s \ell^+ \ell^-) = (0.44 \pm 0.12) \times 10^{-6}$
$\Delta M_s = (17.77 \pm 0.12) ps^{-1}$	$\mathcal{BR}(K^+ \rightarrow \pi^+ \nu \nu) = (0.147_{-0.089}^{+0.130}) \times 10^{-9}$
$\Delta M_d = (0.507 \pm 0.005) ps^{-1}$	$\mathcal{BR}(B \rightarrow X_c \ell \nu) = (10.61 \pm 0.17) \times 10^{-2}$
$ \epsilon_k \times 10^3 = 2.32 \pm 0.007$	$T_4 = 0.11 \pm 0.14$
$\kappa_\epsilon = 0.94 \pm 0.02$ [49] ^b	$m_t(m_t) = (163.5 \pm 1.7) GeV$

^aIt is the weighted average of $V_{ub}^{inl} = (40.1 \pm 2.7 \pm 4.0) \times 10^{-4}$ and $V_{ub}^{exl} = (29.7 \pm 3.1) \times 10^{-4}$. In our numerical work to follow, we increase the error on $|V_{ub}|$ by 50% and take the total error to be around 12% because of the appreciable disagreement between the two determinations.

^bWe tacitly assume that κ_ϵ in SM4 is approximately the same as in SM3.

TABLE I: Inputs that we use in order to constrain the SM4 parameter space, when not explicitly stated, we take the inputs from Particle Data Group [7]; for the lattice inputs see also [6].

From direct searches at the Tevatron, it follows that $m_{t'} > 335 GeV$ [35]. Taking into account the limits from electroweak precision tests [36–39], perturbativity [40] and indications from our studies [13, 14], plausible ranges for $m_{t'}$ and $m_{b'}$ can be taken as,

$$375 GeV < m_{t'} < 575 GeV, \quad m_{t'} - m_{b'} \approx \left(1 + \frac{1}{5} \ln \frac{m_H}{115 GeV}\right) \times 55 GeV. \quad (4)$$

Here the mass splitting depends on the higgs mass, however for simplicity we do not consider the variation of higgs mass and take mass splitting to be $\approx 50 GeV$.

Detailed formulas for the above mentioned observables (Table I) can be seen from one of our earlier papers [14]. In this paper, we do not impose $S_{\psi K_s} = \sin 2\beta_{eff}$ as a constraint, we show SM4 prediction for $S_{\psi K_s}$ and its correlation with the semileptonic asymmetry a_{sl}^d . The expression for the semileptonic asymmetry is given by

$$a_{sl}^q = \frac{|\Gamma_{12}^q|}{|M_{12}^q|} \sin \phi_q = \frac{|\Gamma_{12}^q|}{|M_{12}^{q,SM}|} \frac{\sin \phi_q}{|\Delta_q|}, \quad (q = d, s), \quad (5)$$

where $|\Gamma_{12}^q|$, $|M_{12}^q|$ are the width and mass differences between heavy and light mass eigenstates of B_q mesons, and the CP violating phase ϕ_q ($q = d, s$) is defined as

$$\phi_q \equiv Arg \left[-\frac{M_{12}^q}{\Gamma_{12}^q} \right]. \quad (6)$$

In eq. 5, Δ_q parameterized the new physics effects in mixing and defined as

$$M_{12}^q = M_{12}^{q,SM} \left(1 + \frac{M_{12}^{q,NP}}{M_{12}^{q,SM}}\right) = M_{12}^{q,SM} \Delta_q. \quad (7)$$

In the SM the CP phases are small, $\phi_d \approx -4.3^\circ$ and $\phi_s \approx 0.22^\circ$ [32], new physics can affect the magnitude and phase of M_{12}^q , hence ΔM_q and ϕ_q can deviate substantially from their SM predictions. The semileptonic asymmetry is the function of the CP phase ϕ_q , $|\Delta_q|$ and the ratio $\frac{|\Gamma_{12}^q|}{|M_{12}^{q,SM}|}$ ($q = d, s$). The SM predictions for Γ_{12}^q suffers from large theoretical uncertainties which are $\approx 40\%$ and 30% for B_s and B_d system respectively [32], however, in the ratio $\frac{|\Gamma_{12}^q|}{|M_{12}^{q,SM}|}$ hadronic uncertainties due to the decay constants f_{B_q} cancel to a large extent. In the B_d and B_s system one has [32]

$$\begin{aligned} \frac{|\Gamma_{12}^d|}{|M_{12}^{d,SM}|} &= (52.6_{-12.8}^{+11.5}) \times 10^{-4}, \\ \frac{|\Gamma_{12}^s|}{|M_{12}^{s,SM}|} &= (4.97 \pm 0.94) \times 10^{-3}, \end{aligned} \quad (8)$$

with the theoretical error in both the cases $\approx 20\%$. However, to predict a_{sl}^q in any new physics (NP) scenario one needs to constrain $|\Delta_q|$ from the data on ΔM_q eq. 7, therefore this gets limited by the precision of the lattice calculations.

We also study $D^0 - \bar{D}^0$ mixing in the presence of a fourth generation of quarks [16]. In particular, we calculate the size of the allowed CP violation, which could be large compared to the SM, and show its parametric dependence on CKM4 elements.

Within the SM, $D^0 - \bar{D}^0$ mixing proceeds to an excellent approximation only through the box diagrams with internal b and s quark exchanges. In the case of four generations there is an additional important contribution to $D^0 - \bar{D}^0$ mixing coming from the virtual exchange of the fourth generation down quark b' .

The short distance (SD) contributions to the matrix element of the $\Delta C = 2$ effective Hamiltonian can be written as

$$\langle \bar{D}^0 | \mathcal{H}_{\text{eff}}^{\Delta C=2} | D^0 \rangle_{\text{SD}} \equiv |M_{12}^D| e^{2i\phi_D} = (M_{12}^D)^*, \quad (9)$$

where

$$M_{12}^D = \frac{G_F^2}{12\pi^2} f_D^2 \hat{B}_D m_D M_W^2 \overline{M}_{12}^D, \quad (10)$$

with

$$\begin{aligned} \overline{M}_{12}^D &= \lambda_s^{(D)*2} \eta_{cc}^{(K)} S_0(x_s) + \lambda_b^{(D)*2} \eta_{cc}^{(K)} S_0(x_b) + \lambda_{b'}^{(D)*2} \eta_{tt}^{(K)} S_0(x_{b'}) \\ &+ 2\lambda_b^{(D)*} \lambda_s^{(D)*} \eta_{cc}^{(K)} S_0(x_b, x_s) + 2\lambda_{b'}^{(D)*} \lambda_s^{(D)*} \eta_{ct}^{(K)} S_0(x_{b'}, x_s) + 2\lambda_{b'}^{(D)*} \lambda_b^{(D)*} \eta_{ct}^{(K)} S_0(x_{b'}, x_b), \end{aligned} \quad (11)$$

where

$$\lambda_i^{(D)} = V_{ci}^* V_{ui} \quad (i = s, b, b'). \quad (12)$$

For the QCD corrections we will use the approximate relations

$$\eta_{b'b'}^{(D)} \approx \eta_{tt}^{(K)}, \quad \eta_{b'b}^{(D)} \approx \eta_{b's}^{(D)} \approx \eta_{ct}^{(K)}, \quad \eta_{ss}^{(D)} \approx \eta_{bb}^{(D)} \approx \eta_{bs}^{(D)} \approx \eta_{cc}^{(K)}. \quad (13)$$

Including the long distance part the full matrix elements are given by,

$$\langle \bar{D}^0 | \mathcal{H}_{\text{eff}}^{\Delta C=2} | D^0 \rangle = (M_{12}^D + M_{12}^{\text{LD}})^* - \frac{i}{2} \Gamma_{12}^{\text{LD}*}, \quad (14)$$

$$\langle D^0 | \mathcal{H}_{\text{eff}}^{\Delta C=2} | \bar{D}^0 \rangle = (M_{12}^D + M_{12}^{\text{LD}}) - \frac{i}{2} \Gamma_{12}^{\text{LD}}. \quad (15)$$

Here Γ_{12}^{LD} and M_{12}^{LD} stand for long distance (LD) contributions with the former arising exclusively from SM3 dynamics. These contributions are very difficult to estimate reliably. In our work we scan flatly over the intervals [50, 51].

$$-0.02 \text{ ps}^{-1} \leq M_{12}^{\text{LD}} \leq 0.02 \text{ ps}^{-1}, \quad (16)$$

$$-0.04 \text{ ps}^{-1} \leq \Gamma_{12}^{\text{LD}} \leq 0.04 \text{ ps}^{-1}. \quad (17)$$

$D^0 - \bar{D}^0$ oscillations can be characterized by the normalized mass and width differences

$$x_D \equiv \frac{\Delta M_D}{\bar{\Gamma}}, \quad y_D \equiv \frac{\Delta \Gamma_D}{2\bar{\Gamma}}, \quad \bar{\Gamma} = \frac{1}{2}(\Gamma_1 + \Gamma_2), \quad (18)$$

with

$$\begin{aligned} \Delta M_D &= M_1 - M_2 = 2\text{Re} \left[\frac{q}{p} (M_{12}^D - \frac{i}{2} \Gamma_{12}^D) \right] \\ &= 2\text{Re} \sqrt{|M_{12}^D|^2 - \frac{1}{4} |\Gamma_{12}^D|^2 - i\text{Re}(\Gamma_{12}^D M_{12}^{D*})}, \end{aligned} \quad (19)$$

$$\begin{aligned} \Delta \Gamma_D &= \Gamma_1 - \Gamma_2 = -4\text{Im} \left[\frac{q}{p} (M_{12}^D - \frac{i}{2} \Gamma_{12}^D) \right] \\ &= -4\text{Im} \sqrt{|M_{12}^D|^2 - \frac{1}{4} |\Gamma_{12}^D|^2 - i\text{Re}(\Gamma_{12}^D M_{12}^{D*})}, \end{aligned} \quad (20)$$

where

$$\frac{q}{p} \equiv \sqrt{\frac{M_{12}^{D*} - \frac{i}{2} \Gamma_{12}^{D*}}{M_{12}^D - \frac{i}{2} \Gamma_{12}^D}}. \quad (21)$$

For practical purposes it is sufficient to consider the time-dependent CP asymmetry S_f as [50]

$$\frac{\Gamma(D^0(t) \rightarrow f) - \Gamma(\bar{D}^0(t) \rightarrow f)}{\Gamma(D^0(t) \rightarrow f) + \Gamma(\bar{D}^0(t) \rightarrow f)} \equiv S_f(D) \frac{t}{2\bar{\tau}_D}, \quad (22)$$

parameter	value	parameter	value
m_D	$(1.86484 \pm 0.00017)\text{GeV}$	$\bar{\tau}_D$	$(0.4101 \pm 0.0015)\text{ps}$
f_D	$(0.212 \pm 0.014)\text{GeV}$ [55]	$m_c(m_c)$	$(1.268 \pm 0.009)\text{GeV}$ [57, 58]
\hat{B}_D	$1.18^{+0.07}_{-0.05}$ [16, 56]	$m_b(m_b)$	$(4.20^{+0.17}_{-0.07})\text{GeV}$ [52]

TABLE II: Values of the input parameters for D mesons used in our analysis.

which is given by

$$\eta_f S_f(D) \simeq - \left[y_D \left(\left| \frac{q}{p} \right| - \left| \frac{p}{q} \right| \right) \cos \varphi - x_D \left(\left| \frac{q}{p} \right| + \left| \frac{p}{q} \right| \right) \sin \varphi \right], \quad (23)$$

where $\eta_f = \pm 1$ is the CP parity of the final state f . The SM3 prediction for $\eta_f S_f(D)$ is [16, 50]

$$[\eta_f S_f(D)]_{\text{SM3}} \approx -2 \cdot 10^{-6}. \quad (24)$$

Finally, the semileptonic asymmetry is defined as

$$a_{\text{SL}}(D) \equiv \frac{\Gamma(D^0(t) \rightarrow \ell^- \bar{\nu} K^{+(*)}) - \Gamma(\bar{D}^0 \rightarrow \ell^+ \nu K^{-(*)})}{\Gamma(D^0(t) \rightarrow \ell^- \bar{\nu} K^{+(*)}) + \Gamma(\bar{D}^0 \rightarrow \ell^+ \nu K^{-(*)})} = \frac{|q|^4 - |p|^4}{|q|^4 + |p|^4} \approx 2 \left(\left| \frac{q}{p} \right| - 1 \right). \quad (25)$$

The world averages based on data from BaBar, Belle and CDF are given by [52–54]

$$\begin{aligned} x_D &= (0.98^{+0.24}_{-0.26}) \%, & y_D &= (0.83 \pm 0.16) \%, \\ |q/p| &= (0.87^{+0.17}_{-0.15}), & \phi &= (-8.5^{+7.4}_{-7.0})^\circ, \\ \eta_f S_f(D) &= (-0.248 \pm 0.496) \%, \end{aligned} \quad (26)$$

with ϕ being the phase of q/p and the asymmetry $\eta_f S_f(D)$ defined in (22).

In addition to Table I the relevant input parameters for $D^0 - \bar{D}^0$ mixing are given in Table II.

C. Results

Allowed ranges for different CKM4 parameters are given in Table III. Constraint on $V_{t'b}$ or equivalently on the new parameter r (i.e $V_{t'b} = -r\lambda$) is obtained from non-decoupling oblique corrections (T_4) and vertex corrections to $Z \rightarrow b\bar{b}$. We also note the allowed ranges for the product of the different CKM4 elements, $|\lambda_{db}^{t'}| = |V_{t'd}^* V_{tb}|$, $|\lambda_{sb}^{t'}| = |V_{t's}^* V_{tb}|$ and $|\lambda_{uc}^{b'}| = |V_{ub}^* V_{cb'}|$, obtained from our analysis; these are relevant to B_d^0 , B_s and D^0 oscillations. Allowed ranges for the corresponding phases and their correlations with the magnitude of the product couplings are shown in Fig. 1. We note that values of $|\lambda_{sb}^{t'}|$ larger than 0.002 correspond to very narrow regions of the phase $\delta_{t's}$ (left panel, Fig. 1) close to 90° or 270° , whereas that for $\delta_{uc}^{b'} = \delta_{cb'} - \delta_{ub'}$ (right

parameter	allowed range	parameter	allowed range
λ	0.2205 ± 0.0018	$ V_{t'b} $	< 0.12
C	$0.32 \rightarrow 0.42$	$ V_{t'd} $	< 0.05
A	$0.825 \rightarrow 0.865$	$ V_{t's} $	< 0.11
γ	$(73 \pm 13)^\circ$	$ V_{ub'} $	< 0.05
r	< 0.5	$ V_{cb'} $	< 0.11
P	< 5.0	$ \lambda_{db'}^{t'} $	< 0.002
Q	< 2.5	$ \lambda_{sb'}^{t'} $	< 0.01
		$ \lambda_{uc'}^{b'} $	< 0.0025

TABLE III: Allowed ranges of the CKM4 parameters obtained from our analysis.

panel) is close to zero when $|\lambda_{uc'}^{b'}| \gtrsim 0.0008$. $B_d - \bar{B}_d$ and $B_s - \bar{B}_s$ mixing are sensitive to the new parameters $[P, \delta_{t'd}]$ and $[Q, \delta_{t's}]$ respectively whereas $K^0 - \bar{K}^0$ and $D^0 - \bar{D}^0$ mixing are sensitive to all these four new parameters and their parametric dependencies are given by,

$$\begin{aligned}\lambda_{ds}^{t'} &= V_{t's}^* V_{t'd} = P Q \lambda^5 e^{i(\delta_{t'd} - \delta_{t's})} \\ \lambda_{uc}^{b'} &= V_{ub'}^* V_{cb'} = Q \lambda^5 \left(Q + P e^{i(\delta_{t'd} - \delta_{t's})} \right)\end{aligned}\quad (27)$$

respectively. In this framework it is quite natural to expect that there is a strong correlation between $K^0 - \bar{K}^0$ and $D^0 - \bar{D}^0$ mixing, as pointed out in the case of purely left-handed currents [50, 59], $D^0 - \bar{D}^0$ mixing is also correlated with the observables from B_d and B_s mixing and decays. So the constraints obtained on the new parameters from the inputs given in Table I, especially ϵ_K and $Br(K^+ \rightarrow \pi^+ \nu \bar{\nu})$, are helpful to find the allowed parameter space for $\lambda_{uc}^{b'}$ and the corresponding phase difference $\delta_{uc}^{b'}$.

In Fig. 2 (upper-left panel) we show the correlation between P and Q , larger values of P corresponds to lower value of Q and vice versa. We obtain such a correlation mainly due to the constraints from ϵ_K and $Br(K^+ \rightarrow \pi^+ \nu \bar{\nu})$, although the upper bound on P and Q is coming from the B_d and B_s data (see Table I). The expressions for ϵ_K and $Br(K^+ \rightarrow \pi^+ \nu \bar{\nu})$ are sensitive to $\lambda_{ds}^{t'}$ i.e using these inputs we will get direct constraint on $\lambda_{ds}^{t'}$; as indicated in eq. 27 $\lambda_{ds}^{t'}$ is proportional to the product of P and Q . Therefore we will get direct constraint on the product not on individual P or Q and this is the reason why they follow the correlations shown. Similar correlation is possible between $V_{t'd}$ and $V_{t's}$ since they are proportional to P and Q respectively. We




FIG. 1: Fourth generation parameter space; left panel shows the variation of $|\lambda_{sb}^{t'}| = |V_{t's}^* V_{t'b}|$ with the phase $\delta_{t's}$ of $V_{t's}$, whereas right panel shows it for $|\lambda_{uc}^{b'}| = |V_{ub'}^* V_{cb'}|$ with $\delta_{uc}^{b'}$, the phase difference between the phase of $V_{ub'}^*$ and $V_{cb'}$.

also show the correlations between some other CKM4 elements³; the plot of $|V_{t's}|$ as a function of $|V_{cb'}|$ (lower-left panel) shows that $|V_{t's}| \approx |V_{cb'}|$ since leading order contribution to both the terms is proportional to Q . However, the plot of $|V_{t'd}|$ as a function of $|V_{ub'}|$ (upper-right panel) shows such a relationship only when $P \gg Q$ since the leading order contribution to $|V_{t'd}|$ is proportional to P whereas that for $|V_{ub'}|$ is proportional to a linear combination of P and Q , see eq. 3. For relatively smaller values of $|V_{t'd}|$, of $\mathcal{O}(10^{-3})$, $|V_{ub'}|$ could be as high as 0.02; this is possible when $Q > P$ or alternatively when $|V_{t's}| \gtrsim 0.025$ (middle-right panel, Fig. 2). It also shows that larger values of $|V_{ub'}|$ are still possible which correspond to $P \gg Q$ i.e for smaller values of $|V_{t's}|$. Similar feature can be observed in the correlation between $|V_{ub'}|$ and $|V_{cb'}|$ (lower-right panel). Middle-left panel of Fig. 2 shows the correlation between $|V_{t'd}|$ and $|V_{cb'}|$ which is similar to the correlation between P and Q since leading order contribution in $|V_{cb'}|$ is $\propto Q$ and that for $|V_{t'd}|$ is $\propto P$.

The expressions for the product of CKM4 elements $|\lambda_{db}^{t'}|$ and $|\lambda_{sb}^{t'}|$ are given by

$$\begin{aligned} |\lambda_{db}^{t'}| &= |V_{t'd}^* V_{t'b}| = Pr\lambda^4 \\ |\lambda_{sb}^{t'}| &= |V_{t's}^* V_{t'b}| = Qr\lambda^3, \end{aligned} \quad (28)$$

whereas that for $|\lambda_{uc}^{b'}|$ can be obtained from eq. 27 by taking its modulus, and we see that when $P \ll Q$ it is $\approx Q^2\lambda^5$. In Fig. 3 we show the correlations between the products of CKM4 elements; upper-left panel shows the correlation between $|\lambda_{db}^{t'}|$ and $|\lambda_{sb}^{t'}|$ which is similar to the correlation

³ Flavour data allows us to get direct constraints on various products of CKM4 elements. The bounds on individual CKM4 element are obtained using the constraints on the product couplings. Data for non-decoupling oblique corrections helps to get tighter constrain on $|V_{t'b}|$ which helps to constrain $|V_{t's/d}|$ from the bound on $|\lambda_{s/db}^{t'}|$.




FIG. 2: Correlations between different new CKM4 elements.

between P and Q (upper-left panel Fig. 2), as expected since the slope of the curve is given by $\frac{Q}{P\lambda}$. In the upper-right panel of Fig. 3 we show the correlation between $|\lambda_{db}'|$ and $|\lambda_{uc}'|$ and note that $|\lambda_{uc}'|$ could be as large as 0.0025 when $|\lambda_{db}'|$ is very small (say < 0.0005) i.e when $P \ll Q$ and vice versa. The most interesting one is the correlation between $|\lambda_{sb}'|$ and $|\lambda_{uc}'|$ (lower-panel Fig. 3); it shows an almost linear relationship between them which is prominent for larger values of $|\lambda_{sb}'|$ i.e for larger values of Q due to strong Q^2 dependence of $|\lambda_{uc}'|$. It plays an important role in understanding the correlations between the CP asymmetries in B_s and D system; below we will




FIG. 3: Correlations between different CKM4 product couplings, $|\lambda_{db}^{t'}| = |V_{t'd}^* V_{t'b}|$ and $|\lambda_{sb}^{t'}| = |V_{t's}^* V_{t'b}|$ (upper left panel), $|\lambda_{db}^{t'}|$ and $|\lambda_{uc}^{b'}| = |V_{ub'}^* V_{cb'}|$ (upper right panel), $|\lambda_{sb}^{t'}|$ and $|\lambda_{uc}^{b'}|$ (lower panel).

discuss it in detail. The final remark from these discussions is that the allowed parameter space for the new CKM4 parameter space are highly correlated; random choices of the CKM4 parameters should not be done, in fact chosen values should be consistent with the appropriate correlations.

Let us move to next part of our discussion where we show the effect of the fourth generation on different observables related to B_d , B_s and D system. As mentioned previously SM4 is quite different from most extensions of the SM in the sense that it is highly constrained. This motivates us to search for observables that can be used to confirm or rule out SM4. With this in mind we study the CP asymmetries $S_{\psi K_s}$, a_{sl}^d , $S_{\psi\phi}$ and a_{sl}^s and correlations among them.

In Fig. 4 (upper-left panel) we show CP asymmetry $S_{\psi K_s}$ as a function of $\lambda_{db}^{t'}$ and note that $S_{\psi K_s}$ can go down to ≈ 0.4 or can reach around 0.9 for large values of the product coupling $|\lambda_{db}^{t'}|$; so agreement with or appreciable deviation from the present experimental measurement are, in principle, both possible. We do not get any noticeable correlation between $S_{\psi K_s}$ with the phase $\delta_{t'd}$ of $\lambda_{db}^{t'}$. Finally, we note that though SM4 allows $|\Delta_d|$ to have values within the range $0.65 \lesssim |\Delta_d| \lesssim 1.25$ (eq. 7), it does not have any noticeable correlation with $S_{\psi K_s}$. In the upper-right

panel of Fig. 4 we show the semileptonic asymmetry a_{sl}^d (eqs. 5 and 8) as a function of $S_{\psi K_s}$. We see that while, in principle, in SM4, $-0.002 \lesssim a_{sl}^d \lesssim 0.002$, the experimental bound on $S_{\psi K_s}$ allows only $a_{sl}^d \gtrsim -0.001$. Recall that SM has a bound, $(-4.8_{-1.2}^{+1.0}) \times 10^{-4}$ as shown by the black band in the Fig. 4 (upper-right panel). Thus in SM4, a_{sl}^d can be large by a factor of about 2 or 3; even more importantly the correlation between a_{sl}^d and $S_{\psi K_s}$ can be very useful for testing the SM4.




FIG. 4: Various correlations in SM4 are shown. Variation of CP asymmetries with the magnitude of the product couplings; $S_{\psi K_s}$ as a function of $|\lambda_{db}^{t'}|$ (upper-left panel), $S_{\psi\phi}$ as a function of $|\lambda_{sb}^{t'}|$ (lower-left panel). Correlations between the time dependent mixing induced CP asymmetries and semileptonic CP asymmetries for B_d and B_s are shown in upper and lower right panel respectively. Blue horizontal and vertical bands are the corresponding experimental ranges. In the upper-right panel SM allowed band (thick dark) is shown in $S_{\psi K_s}$ vs a_{sl}^d plane. In the lower-right panel the red and blue regions correspond to $\frac{|\Gamma_{12}^s|}{|M_{12}^{s,SM}|}$ with the uncertainties taken at 1σ and 2σ respectively and grey horizontal band corresponds to the experimental range for a_{sl}^s and the vertical band is that for CP asymmetry.

Similarly, in the lower-left panel of Fig. 4 we are showing the allowed regions for the CP asymmetry $S_{\psi\phi}$ in $B_s \rightarrow \psi\phi$ as a function of $|\lambda_{sb}^{t'}|$, for $375 \text{ GeV} < m_{t'} < 575 \text{ GeV}$, $S_{\psi\phi}$ is bounded

by $-0.60 < S_{\psi\phi} < 0.60$ ⁴, the explicit dependence on $m_{t'}$ has been shown in our earlier papers [13, 14]. It is also interesting to note that its magnitude increases with $|\lambda'_{sb}|$; precise measurements of $S_{\psi\phi}$ will be helpful to put tighter constraints on $|\lambda'_{sb}|$ and the corresponding phase. Recently CDF and DO have updated their measurement of the CP-violating phase with data sample corresponding to an integrated luminosity of 5.2 fb^{-1} and 6.1 fb^{-1} respectively. The allowed 68% C.L ranges are [60, 61]

$$\begin{aligned} \phi_s^{\psi\phi} &\in [-0.04, -1.04] \cup [-2.16, -3.10], & CDF \\ &\in -0.76^{+0.38}_{-0.36} (stat) \pm 0.02 (syst) & DO. \end{aligned} \quad (29)$$

The corresponding 1σ ranges for $S_{\psi\phi} = \sin \phi_s^{\psi\phi}$ are given in Table IV.




FIG. 5: Correlation between $S_{\psi\phi}$ and $|\Delta_s|$ (eq. 7).

In the lower-right panel of Fig. 4 we show the correlation ⁵ between $S_{\psi\phi}$ and a_{sl}^s (eq. 5) with $\frac{|\Gamma_{12}^s|}{|M_{12}^{s,SM}|}$ (eq. 8) taken at 1σ (red) and 2σ (blue) of the theory error. We note that its magnitude increases with $S_{\psi\phi}$ as well as with the ratio $\frac{|\Gamma_{12}^s|}{|M_{12}^{s,SM}|}$, as expected from eq. 5. It is also important to note that a_{sl}^s is inversely proportional to $|\Delta_s|$. As shown in Fig. 5, $|\Delta_s|$ has a strong correlation with $S_{\psi\phi}$, we note that the values like $|\Delta_s| \approx 0.7$ are allowed for larger values of $|S_{\psi\phi}|$. Therefore a_{sl}^s is expected to get an additional boost for lower values of $|\Delta_s|$ when $|S_{\psi\phi}|$ reaches its maximum

⁴ Actually the limit is $|S_{\psi\phi}| \lesssim 0.56$, it is rounded up to 0.6.

⁵ The plot corresponds to negative solution for $S_{\psi\phi}$, we do not show the points corresponding to the positive solution of $S_{\psi\phi}$ for which one should get a region symmetric to that shown in the figure.

value (≈ 0.5). The maximum allowed ranges in SM4 are thus given by

$$\begin{aligned} |a_{sl}^s| &\lesssim 0.004, & \frac{|\Gamma_{12}^s|}{|M_{12}^{s,SM}|} @1\sigma, \\ &\lesssim 0.005, & \frac{|\Gamma_{12}^s|}{|M_{12}^{s,SM}|} @2\sigma. \end{aligned} \quad (30)$$

Perhaps by taking the theory error on $\frac{|\Gamma_{12}^s|}{|M_{12}^{s,SM}|}$ at 2σ we are exhibiting overabundance of caution. Be that as it may, it is clear that although SM4 can increase a_{sl}^s over SM3 by over two order of magnitude, SM4 has difficulty actually reaching the central value of the D0 result. Given that the D0 results on a_{sl}^s is only $\sim 2\sigma$, and considering the fact that this is an extremely difficult measurement we do not believe at this point it is a concern for SM4. In our opinion fragility of the experimental result suggests independent verification is essential ⁶.

In Table IV we summarise the allowed ranges for different CP observables in SM4, it includes time dependent CP asymmetries in $B_d \rightarrow \psi K_s$, $B_s \rightarrow \psi \phi$ as well as the semileptonic asymmetries associated with B_d and B_s system (eq. 5). We also mention the corresponding experimental ranges and SM3 predictions obtained with the inputs given in Table I.

CP observable	SM3	Exp	SM4 ranges
$S_{\psi K_s} = \sin 2\beta_d$	0.739 ± 0.049	0.67 ± 0.02	$0.40 \rightarrow 0.90$
$S_{\psi \phi} = \sin \phi_s^{\psi \phi}$	-0.04 ± 0.002	$[-0.04, -0.86]$ CDF $[-0.37, -0.90]$ DO	$-0.60 \rightarrow 0.60$
a_{sl}^d	$(-4.8_{-1.2}^{+1.0}) \times 10^{-4}$	-0.0047 ± 0.0046	> -0.002
a_{sl}^s	$(2.1 \pm 0.6) \times 10^{-5}$	-0.0146 ± 0.0075	$-0.005 \rightarrow 0.005$ (eq. 30)
$\eta_f S_{CP}(D)$	$\approx -2 \cdot 10^{-6}$	$-0.248 \pm 0.496\%$	$-0.01 \rightarrow 0.01$
$a_{sl}(D)$	$\approx 1 \cdot 10^{-4}$		$-0.6 \rightarrow 0.6$

TABLE IV: Allowed ranges of different CP observables related to B_d , B_s and D^0 systems in SM3 and SM4; current experimental status is also given.

In Fig. 6 we show the correlation between the real and imaginary parts of the short distance (SD) contribution to $D^0 - \bar{D}^0$ mixing. Note that the magnitude of $Im(M_{12}^D)$ could be as high as 0.6%, which could be negative or positive; very small number of points are allowed for $Re(M_{12}^D) < 0$, however, it could be as high as 0.032. These findings are in good agreement with Ref. [16].

In Fig. 7 we plot real (left panel) and imaginary (right panel) parts of the SD contribution to M_{12}^D as a function of $|\lambda_{uc}'|$ and note that in both the cases its magnitude increases with the product

⁶ Lack of identification of B mesons is a very serious concern for this experimental result, see also footnote 1




FIG. 6: Correlation between the real and imaginary part of the SD contribution to M_{12}^D .




FIG. 7: Real (left panel) and imaginary (right panel) parts of the SD contribution to M_{12}^D as a function $|\lambda_{uc}^{b'}|$.

coupling. In the case of the real part almost all the allowed points are for $Re(M_{12}^D) > 0$, however, in case of imaginary part we have both positive and negative solutions. As noticed before (Fig. 3), $|\lambda_{sb}^{t'}|$ has a linear relationship with $|\lambda_{uc}^{b'}|$; a tighter constraint on $|\lambda_{sb}^{t'}|$, which is possible to get by reducing the errors in the measurements of B_d or B_s observables, will be helpful to put tighter constrain on $D^0 - \bar{D}^0$ mixing.

In Fig. 8 we plot the time dependent CP asymmetry $\eta_f S_{CP}(D)$ (eq. 23) and the semileptonic asymmetry $a_{sl}(D)$ (eq. 25) in the D system as a function of the phase of $\frac{q}{p}$ (eq. 21) and $|\frac{q}{p}|$ respectively; it could be directly compared with the correlations shown in [16]. We note that with




FIG. 8: Time dependent CP asymmetry for D system, $\eta_f S_{CP}(D)$, as a function of the phase ϕ of $\frac{q}{p}$ (left panel); semileptonic CP asymmetry, $a_{sl}(D)$, as a function of $|\frac{q}{p}|$ (right panel). The corresponding SM value for $a_{sl}(D)$ is $\mathcal{O}(10^{-4})$ [16]. The grey horizontal and vertical bands represent the corresponding experimental ranges.

the present experimental bound on the phase of $\frac{q}{p}$ (eq. 26), the magnitude of $\eta_f S_{CP}(D)$ could be enhanced up to the present experimental bound. On the other hand with the present constraint on $|\frac{q}{p}|$, $a_{sl}(D)$ could be reduced to -0.6 ; again these results are also in agreement with Buras et. al [16].




FIG. 9: $\eta_f S_{CP}(D)$ as a function of $|\lambda'_{uc}|$ (left panel), $a_{sl}(D)$ as a function $|\lambda'_{uc}|$ (right panel).

In Fig. 9 we plot $S_{CP}(D)$ and $a_{sl}(D)$ as a function of $|\lambda'_{uc}|$ and note that the magnitude of both may increase with $|\lambda'_{uc}|$; SM3 predictions and the allowed ranges in SM4 for the corresponding observables are summarised in Table IV. As discussed before (Fig. 4), the magnitude of $S_{\psi\phi}$ increases with the corresponding product coupling, we also noticed that $|\lambda'_{uc}|$ increases with $|\lambda'_{sb}|$ which indicates a definite correlation between $S_{\psi\phi}$ and $\eta_f S_{CP}(D)$ [16]. In the near future if we are

able to put tighter constraints on $|\lambda_{sb}^{t'}|$, we will be able to get strong limit on $\eta_f S_{CP}(D)$ and $a_{sl}(D)$ due to fourth generation effects.

III. CONCLUSION

This paper represents a continuation of our study of some of the properties of SM4, Standard Model with four generations. One feature of SM4 that distinguishes it from practically all other beyond the Standard Model scenarios is that it is highly constrained. Recognizing this, we have few objectives in mind for this work. Firstly we want to quantitatively ascertain how well SM4 is able to address a key anomaly for the Standard Model, i.e. $S_{\psi K_s}$. We also want to see how well SM4 can address the semi-leptonic asymmetries that have been much in the news recently. Simple intuitive arguments suggest that the CP asymmetries in $S_{\psi K_s}$ and a_{sl}^d should be strongly correlated and similarly, $S_{\psi\phi}$ and a_{sl}^s .

Furthermore, we want to see how best experimental information can be used to extract the parameters of SM4 as efficiently and accurately as possible. With this in mind we are using a particular representation of the 4×4 mixing matrix which is specifically designed to be very effective in extracting information on the parameters of SM4 from B-decays, supplemented by extending the hierarchical nature as for SM3, since many B-decay modes are highly sensitive to SM4. Of course, we understand that physics will not depend on which representation of 4×4 mixing matrix one uses. Recall, though, that even for the simpler case of the SM, a number of different representations have been studied over the past few decades.

Using the specific representation for the 4×4 mixing matrix mentioned above we obtain constraints and correlations on its elements using available data from K, B and D decays as well as electroweak precision tests and oblique corrections and allowing the $m_{t'}$ mass to range from 375 to 575 GeV. Constraints obtained are then used to study the mixing induced and semi-leptonic CP asymmetries in B_d , B_s and in D^0 . Although SM4 allows $S(B_d \rightarrow \psi K_s)$ to be closer to experiment thus alleviating a key difficulty for SM3 that has been found in recent years and SM4 also allows a_{sl}^d to be bigger by a factor of $O(3)$, these two observables are strongly correlated. Thus while $S_{\psi K_s}$ can be $\approx 0.70^7$, simultaneously SM4 restricts $|a_{sl}^d| \lesssim 0.0015$. The B-factories have a lot more data

⁷ $S_{\psi K_s}$ provides a clean measurement of $\sin 2\beta$ and it is well known that in SM the contamination due to QCD or electroweak penguins is small (less than a few percent [62]) in this decay. Although the effects are sub-dominant, SM4 can contribute to the decay $b \rightarrow c\bar{s}$ through the penguin diagrams and the overall effects are expected to be below 5%. Due to the weak mass dependences of the Wilson coefficients the contributions from the QCD penguins (P_{QCD}^{SM4}) in SM4 are expected to be small compare to P_{QCD}^{SM3} since there is an additional suppression of order λ^2

since they studied this asymmetry some years ago [10]; it would be very worthwhile to update this bound as it could provide a very useful non-trivial test of SM4.

In contrast to a_{sl}^d , a_{sl}^s can be a lot bigger in SM4, and of opposite sign, than in SM3 where it is essentially negligible. Therefore, it would also be very useful to constrain this asymmetry as well as the linear combination (A_{sl}^b) [11] of the two. Interestingly, the large same sign dimuon asymmetry recently discovered by D0 [11] implies a rather large central value for a_{sl}^s . This has the same sign as in SM4 though the central value of the D0 result is somewhat larger than the expected range in SM4; however, the significance of the D0 result is only about 2σ on a_{sl}^s . We therefore do not believe it is wise to pay too much attention to the central value given the size of the experimental error ($\approx 50\%$). These asymmetries should be a high priority target for experiments at the Tevatron as well as at LHCb. In recent years Belle also has taken appreciable data at the $\Upsilon(5S)$ which should be used for placing bounds on these asymmetries. In the future, these asymmetries should also be a very useful target at the Super-B factories.

Note Added: Very recently Ref [63] presented constraints on SM4 using a completely different representation of the 4X4 mixing matrix [64].

Acknowledgments

SN thanks Theory Division of Saha Institute of Nuclear Physics (SINP) for Hospitality. SN's work is financially supported by NSERC of Canada. The work of AS is supported in part by the US DOE grant # DE-AC02-98CH10886(BNL).

-
- [1] E. Lunghi and A. Soni, Phys. Lett. B **666**, 162 (2008) [arXiv:0803.4340 [hep-ph]].
 - [2] E. Lunghi and A. Soni, JHEP **0908**, 051 (2009) [arXiv:0903.5059 [hep-ph]].
 - [3] A. Lenz *et al.* [CKMfitter Group], arXiv:1008.1593 [hep-ph].
 - [4] M. Bona *et al.* [UTfit Collaboration], Phys. Lett. B **687**, 61 (2010) [arXiv:0908.3470 [hep-ph]].
 - [5] E. Lunghi and A. Soni, Phys. Rev. Lett. **104**, 251802 (2010) [arXiv:0912.0002 [hep-ph]].
 - [6] E. Lunghi and A. Soni, arXiv:1010.6069 [hep-ph].

due to the CKM4 product coupling $\lambda_{bs}^{t'}$. Although the electroweak penguin (P_{EW}^{SM4}) in SM4 get enhanced by $\approx 20\%$ with respect to P_{EW}^{SM3} for heavy $m_{t'}$ (≈ 500 GeV), its effect on $\sin 2\beta$ measurements are expected to be of similar size as that of P_{CD}^{SM} .

- [7] K. Nakamura *et al.* [Particle Data Group], J. Phys. G **37**, 075021 (2010).
- [8] see for example, M. Gronau and J. L. Rosner, Phys. Lett. B **644**, 237 (2007) [arXiv:hep-ph/0610227]; H. Y. Cheng and C. K. Chua, Phys. Rev. D **80**, 114008 (2009) [arXiv:0909.5229 [hep-ph]].
- [9] T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **100**, 161802 (2008); [arXiv:0712.2397 [hep-ex]]; V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **101**, 241801 (2008) [arXiv:0802.2255 [hep-ex]].
- [10] The Heavy Flavor Averaging Group *et al.*, arXiv:1010.1589 [hep-ex].
- [11] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. D **82**, 032001 (2010) [arXiv:1005.2757 [hep-ex]].
- [12] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **105**, 081801 (2010) [arXiv:1007.0395 [hep-ex]].
- [13] A. Soni, A. K. Alok, A. Gird, R. Mohanta and S. Nandi, Phys. Lett. B **683**, 302 (2010) [arXiv:0807.1971 [hep-ph]].
- [14] A. Soni, A. K. Alok, A. Giri, R. Mohanta and S. Nandi, Phys. Rev. D **82**, 033009 (2010) [arXiv:1002.0595 [hep-ph]].
- [15] A. J. Buras, B. Duling, T. Feldmann, T. Heidsieck, C. Promberger and S. Recksiegel, JHEP **1009**, 106 (2010) [arXiv:1002.2126 [hep-ph]].
- [16] A. J. Buras, B. Duling, T. Feldmann, T. Heidsieck, C. Promberger and S. Recksiegel, JHEP **1007**, 094 (2010) [arXiv:1004.4565 [hep-ph]].
- [17] W. S. Hou and C. Y. Ma, Phys. Rev. D **82**, 036002 (2010) [arXiv:1004.2186 [hep-ph]].
- [18] See also, M. Bobrowski, A. Lenz, J. Riedl and J. Rohrwild, Phys. Rev. D **79**, 113006 (2009) [arXiv:0902.4883 [hep-ph]]; O. Eberhardt, A. Lenz and J. Rohrwild, Phys. Rev. D **82**, 095006 (2010) [arXiv:1005.3505 [hep-ph]].
- [19] G. W. S. Hou, arXiv:0810.3396 [hep-ph].
- [20] For earlier related works see, C. Jarlskog and R. Stora, Phys. Lett. **B208**, 288 (1988); F. del Aguila and J. A. Aguilar-Saavedra, Phys. Lett. **B386**, 241 (1996); F. del Aguila and J. A. Aguilar-Saavedra and G. C. Branco, Nucl. Phys. **B510**, 39, 1998.
- [21] J. Carpenter, R. Norton, S. Siegemund-Broka and A. Soni, Phys. Rev. Lett. **65**, 153 (1990). In passing, we note that the quark mass needed for dynamical electroweak symmetry breaking in this work, translated to the fourth family quasi-degenerate doublet, gives $m_{t'} \sim 500 \text{ GeV}$ and $m_H \approx \sqrt{2} m_{t'} \sim 700 \text{ GeV}$.
- [22] See the proceedings of the First International Symposium on the fourth family of quarks and leptons, Santa Monica, CA, Feb 1987, published by the NY Academy of Sciences; eds D. Cline and A. Soni; see also the proceedings of the Second International Symposium on the fourth family of quarks and leptons, Santa Monica, CA, Feb 1989, published by the NY Academy of Sciences; eds D. Cline and A. Soni.
- [23] B. Holdom, Phys. Rev. Lett. **57**, 2496 (1986) [Erratum-ibid. **58**, 177 (1987)]; W. A. Bardeen, C. T. Hill and M. Lindner, Phys. Rev. D **41**, 1647 (1990); C. T. Hill, M. A. Luty and E. A. Paschos, Phys. Rev. D **43**, 3011 (1991); P. Q. Hung and G. Isidori, Phys. Lett. B **402**, 122 (1997) [arXiv:hep-ph/9609518].

- [24] P. Q. Hung and C. Xiong, arXiv:0911.3892 .
- [25] M. Hashimoto and V. A. Miransky, arXiv:0912.4453 .
- [26] M. S. Chanowitz, Phys. Rev. D **79**, 113008 (2009) [arXiv:0904.3570 [hep-ph]].
- [27] M. S. Chanowitz, Phys. Rev. D , 035018 (2010) [arXiv:1007.0043 [hep-ph]].
- [28] W. S. Hou, A. Soni and H. Steger, Phys. Lett. B **192**, 441 (1987).
- [29] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983).
- [30] W. S. Hou, R. S. Willey and A. Soni, Phys. Rev. Lett. **58**, 1608 (1987) [Erratum-ibid. **60**, 2337 (1988)].
- [31] W. S. Hou, A. Soni and H. Steger, Phys. Rev. Lett. **59**, 1521 (1987).
- [32] A. Lenz and U. Nierste, JHEP **0706**, 072 (2007) [arXiv:hep-ph/0612167].
- [33] R. Louvot *et al.* [Belle Collaboration], Phys. Rev. Lett. **102**, 021801 (2009) [arXiv:0809.2526 [hep-ex]]; A. Drutskoy, arXiv:hep-ex/0605110.
- [34] Ulrich Uwer, talk at the conference “Flavor Physics in the LHC Era”, November 8-12 2010 , Singapore, <http://www.ntu.edu.sg/ias/upcomingevents/FlavorPhysics/Pages/default.aspx>
- [35] CDF collaboration, CDF note 10110, <http://www-cdf.fnal.gov/physics/S10CDFResults.html>
- [36] H. J. He, N. Polonsky and S. f. Su, Phys. Rev. D **64**, 053004 (2001) [arXiv:hep-ph/0102144].
- [37] V. A. Novikov, L. B. Okun, A. N. Rozanov and M. I. Vysotsky, Phys. Lett. B **529**, 111 (2002) [arXiv:hep-ph/0111028].
- [38] G. D. Kribs, T. Plehn, M. Spannowsky and T. M. P. Tait, Phys. Rev. D **76**, 075016 (2007) [arXiv:0706.3718 [hep-ph]].
- [39] J. Erler and P. Langacker, arXiv:1003.3211 [hep-ph].
- [40] P. Q. Hung and M. Sher, Phys. Rev. D **77**, 037302 (2008) [arXiv:0711.4353 [hep-ph]].
- [41] D. J. Antonio *et al.* [RBC Collaboration and UKQCD Collaboration], Phys. Rev. Lett. **100**, 032001 (2008) [arXiv:hep-ph/0702042]; C. Kelly, talk at Lattice 2010, June 14-19 2010, Sar- dinia, Italy, <http://www.infn.it/Lattice2010>; Y. Aoki *et al.*, arXiv:1012.4178 [hep-lat].
- [42] T. Bae *et al.*, arXiv:1008.5179 [hep-lat].
- [43] C. Aubin, J. Laiho and R. S. Van de Water, Phys. Rev. D **81**, 014507 (2010) [arXiv:0905.3947 [hep-lat]].
- [44] E. Gamiz, C. T. H. Davies, G. P. Lepage, J. Shigemitsu and M. Wingate [HPQCD Collaboration], Phys. Rev. D **80**, 014503 (2009) [arXiv:0902.1815 [hep-lat]].
- [45] E. Gamiz, private communication.
- [46] S. Herrlich and U. Nierste, Nucl. Phys. **B419**, 292 (1994) [hep-ph/9310311].
- [47] A. J. Buras, M. Jamin and P. H. Weisz, Nucl. Phys. B **347**, 491 (1990).
- [48] S. Herrlich and U. Nierste, Phys. Rev. D **52**, 6505 (1995) [hep-ph/9507262].
- [49] A. J. Buras, D. Guadagnoli and G. Isidori, Phys. Lett. B **688**, 309 (2010) [arXiv:1002.3612 [hep-ph]].
- [50] I. I. Bigi, M. Blanke, A. J. Buras, and S. Recksiegel, JHEP **07** (2009) 097, {\tt0904.1545} .
- [51] W. Altmannshofer, A. J. Buras and P. Paradisi, Phys. Lett. B **688**, 202 (2010) [arXiv:1001.3835 [hep-ph]].
- [52] Heavy Flavor Averaging Group (HFAG) Collaboration, E. Barberio *et al.*, Aver-

- ages of b -hadron properties at the end of 2006, <http://www.slac.stanford.edu/xorg/hfag>.
- [53] Heavy Flavor Averaging Group Collaboration, E. Barberio *et al.*, *Averages of b -hadron and c -hadron Properties at the End of 2007*, [\tt0808.1297](#).
 - [54] A. J. Schwartz, [\tt0911.1464](#).
 - [55] V. Lubicz and C. Tarantino, *Nuovo Cim.* **123B** (2008) 674–688, [\tt0807.4605](#).
 - [56] C. W. Bernard, T. Draper, G. Hockney and A. Soni, *Phys. Rev. D* **38**, 3540 (1988); R. Gupta, T. Bhattacharya and S. R. Sharpe, *Phys. Rev. D* **55**, 4036 (1997) [[arXiv:hep-lat/9611023](#)]; L. Lellouch and C. J. D. Lin [UKQCD Collaboration], *Phys. Rev. D* **64**, 094501 (2001) [[arXiv:hep-ph/0011086](#)]; H. W. Lin, S. Ohta, A. Soni and N. Yamada, *Phys. Rev. D* **74**, 114506 (2006) [[arXiv:hep-lat/0607035](#)]. Although these results are obtained in the quenched approximation, we know now that for B-parameters quenching effects are rather small; see Ref. 39
 - [57] J. Laiho, R. S. Van de Water, and E. Lunghi, [\tt0910.2928](#).
 - [58] **HPQCD** Collaboration, I. Allison *et al.*, *Phys. Rev.* **D78** (2008) 054513, [\tt0805.2999](#).
 - [59] K. Blum, Y. Grossman, Y. Nir and G. Perez, *Phys. Rev. Lett.* **102**, 211802 (2009) [[arXiv:0903.2118 \[hep-ph\]](#)].
 - [60] CDF collaboration, CDF note 10206, <http://www-cdf.fnal.gov/physics/new/bottom/bottom.html>
 - [61] DØcollaboration, Conference note 6098-CONF, www-d0.fnal.gov/Run2Physics/WWW/results/prelim/B/B60/
 - [62] M. Ciuchini, M. Pierini and L. Silvestrini, *Phys. Rev. Lett.* **95**, 221804 (2005) [[arXiv:hep-ph/0507290](#)]; M. Gronau and J. L. Rosner, *Phys. Lett. B* **672**, 349 (2009) [[arXiv:0812.4796 \[hep-ph\]](#)].
 - [63] A. K. Alok, A. Dighe and D. London, [arXiv:1011.2634 \[hep-ph\]](#).
 - [64] C. S. Kim and A. S. Dighe, *Int. J. Mod. Phys. E* **16**, 1445 (2007) [[arXiv:0710.1681 \[hep-ph\]](#)].