This is the accepted manuscript made available via CHORUS. The article has been published as:

Probing the Ricci dark energy model with perturbations by
using WMAP seven-year cosmic microwave background
measurements, baryon acoustic oscillations, and type Ia
supernovae

Yuting Wang, Lixin Xu, and Yuanxing Gui

Phys. Rev. D **84**, 063513 — Published 14 September 2011

DOI: [10.1103/PhysRevD.84.063513](https://doi.org/10.1103/PhysRevD.84.063513)

Probing Ricci dark energy model with perturbations by using WMAP seven-year cosmic microwave background measurements, BAO and Type Ia supernovae

Yuting Wang^a, Lixin Xu^{b,a,c*}, and Yuanxing Gui^a

^a*School of Physics and Optoelectronic Technology, Dalian University of Technology,
Dalian, Liaoning 116024, People's Republic of China*

^b*International Center for Astrophysics, Korea Astronomy and Space Science Institute,
Yuseong Daedeokdaero 776, Daejeon 305-348, R. Korea*

^c*College of Advanced Science & Technology, Dalian University of Technology,
Dalian, Liaoning 116024, People's Republic of China*

In this paper, we investigate the Ricci dark energy model with perturbations through the joint constraints of current cosmological datasets from dynamical and geometrical perspectives. We use the full cosmic microwave background information from WMAP seven-year data, the baryon acoustic oscillations from Sloan Digital Sky Survey and the Two Degree Galaxy Redshift Survey, and Type Ia supernovae from Union2 compilation of Supernova Cosmology Project collaboration. A global constraint is performed by employing the Markov chain Monte Carlo method. With the best-fitting results, we show the differences of cosmic microwave background power spectra and background evolutions for the cosmological constant model and Ricci dark energy model with perturbations.

PACS numbers: 98.80.-k, 98.80.Es

Keywords: dark energy; constraint

I. INTRODUCTION

The present observations from WMAP seven-year cosmic microwave background (CMB) radiation measurements [1], large scale structure (LSS) surveys [2, 3] and updated type Ia supernovae (SNIa) [4, 5] have confirmed at a high confidence level the first implication from SNIa in 1998 [6, 7] that our universe is undergoing an accelerated expansion. It has been an important issue of modern cosmology to unveil the mysterious face of the driving force of current accelerated expansion. One of the theoretical viewpoints is that dark energy with negative pressure drives current expansion. A flood of dark energy models have been proposed, such as a natural cosmological constant (Λ CDM model), dynamical dark energy models with scalar fields [8] or with exotic equation of state [9, 10] and so on. Please see [11] for up-to-date review on dark energy.

Although the cosmological constant model keeps a good fit to the current observations, it suffers from fine-tuning problem and cosmic coincidence problem. In the final analysis, both are related to the energy density of vacuum energy. Therefore, the cosmological constant problem is in essence an issue of quantum gravity. Before the complete theory of quantum gravity is established, the dark energy model on the basis of holographic principle of quantum gravity theory has been built in Ref. [10] by applying the energy bound proposed by Cohen et al [12] to cosmology. Namely, for a system with size L and UV cut-off Λ without decaying into a black hole, it is required that the total energy in a region of size L should not exceed the mass of a black hole of the same size, thus $L^3 \rho_\Lambda \leq LM_{pl}^2$. It reveals a duality between UV cut-off and IR cut-off. In cosmology, the UV cut-off is related to the dark energy density, and IR cut-off is related to the large scale of the universe. The holographic dark energy models with different IR cut-offs have been studied and the detailed discussions can be seen in Refs. [10, 13–16].

In Ref. [13], Gao *et. al.* firstly proposed that dark energy density is in direct proportion to Ricci scalar curvature, R , and referred to as Ricci dark energy (RDE) model. The distinct characteristic of RDE model is that the cosmic coincidence problem and causality problem can be naturally solved. However, the physical motivation for such a model was vague in Ref. [13]. Subsequently, Cai *et. al.* [17] were inspired by the motivation of the causal entropy bound in the framework of holography, found the causal connection scale, R_{CC} , and put forward an appropriate physical mechanism for this model. In [17], it was pointed out that the scale on which the black hole can be formed must be within R_{CC} , which is set by the "Jeans" length of the perturbations to $R_{CC}^{-2} = \text{Max}(\dot{H} + 2H^2, -\dot{H})$ while one assumes black hole in the universe is formed by gravitational collapse of perturbations of cosmological spacetime. Only one takes the causal connection scale with $R_{CC}^{-2} = \dot{H} + 2H^2$ as IR cut-off, can the dark energy model be consistent with current observations [17]. It is found that $R_{CC}^{-2} \propto R$. Therefore, the RDE is stressed to be the holographic dark

* lxxu@dlut.edu.cn

energy with IR cut-off of the causal connection scale. The RDE density reads

$$\rho_{de} = 3M_{pl}^2 \alpha \left(\dot{H} + 2H^2 \right), \quad (1)$$

where M_{pl} is the reduced Planck Mass $M_{pl} \equiv 1/\sqrt{8\pi G}$, α is the dimensionless parameter and has been constrained by using the distance measurements from different joint observations [18, 19]. Especially, in these studies, the compressed CMB information is used, including the "shift parameter", the "acoustic scale" and the photon decoupling epoch, the values of which are obtained on a basis of a given model in advance [20]. That is to say, before these distance parameters are used to constrain a model, one should renew to get their values basing on the constrained model. In this paper, we revisit the cosmological parameters in RDE model by using the full CMB power spectrum data from WMAP7 and additional distance measurements from baryon acoustic oscillations (BAO) and Union2 SNIa including 557 data points. Compared with the previous works, we not only focus on the constraint of background evolution, but also take dark energy perturbations into account. For there will be dark energy perturbations according to the perturbed conservation equations when the dark energy is not a cosmological constant. What's more, the perturbations of dark energy play an important role when the dark energy model is confronted with the current observations [21]. The inclusion of dark energy perturbations leaves an imprint on CMB power spectrum [22, 23] and leads to the changes of parameters' spaces in the complete fitting to observational data [23–28]. In particular, there are the significant changes on the large-scale CMB power spectra by the late integrated Sachs-Wolfe (ISW) effect [29], which arises from a time-dependent gravitational potential when dark energy dominates in the universe at late time for a flat dark energy model. The magnitude of ISW-power spectrum is dependent on dark energy models and dark energy perturbations [22, 23, 30–35]. Therefore, a complete constraint on RDE model with perturbations is performed by employing the Markov chain Monte Carlo (MCMC) method.

The paper is organized as follows. In next section, we review the background equations for RDE model and the perturbation equations in the conformal Newtonian gauge. In section III, we perform a global fitting to observational data by using MCMC method and discuss the constraint results. The last section is the conclusion.

II. REVIEW THE DYNAMICS OF RICCI DARK ENERGY: BACKGROUND AND PERTURBATION EVOLUTION

First of all, we give a brief review on the background evolution of RDE model in the framework of unperturbed and spatially flat Friedmann-Robertson-Walker (FRW) metric. The Einstein field equation is written as

$$H^2 = \frac{1}{3M_{pl}^2}(\rho_m + \rho_{de}), \quad (2)$$

where the energy density of RDE, ρ_{de} is rewritten with the variable $x = \ln a$ as

$$\rho_{de} = 3\alpha M_{pl}^2 \left(\frac{1}{2} \frac{dH^2}{dx} + 2H^2 \right). \quad (3)$$

After using the dimensionless definitions $E = \frac{H}{H_0}$ and $\Omega_{m0} = \frac{\rho_{m0}}{3M_{pl}^2 H_0^2}$, we can rewrite Eq. (2) as

$$E^2 = \Omega_{m0} e^{-3x} + \alpha \left(\frac{1}{2} \frac{dE^2}{dx} + 2E^2 \right). \quad (4)$$

The Solution of this first order differential equation with the initial condition $E^2(x=0) = 1$ is obtained

$$E^2 = \frac{2}{2-\alpha} \Omega_{m0} e^{-3x} + \left(1 - \frac{2}{2-\alpha} \Omega_{m0} \right) e^{-(4-\frac{2}{\alpha})x}. \quad (5)$$

From Eqs. (3,5), we can get the dimensionless energy density of RDE

$$\begin{aligned} \Omega_{de}(x) = \frac{\rho_{de}}{3M_{pl}^2 H_0^2} &= \frac{\alpha}{2-\alpha} \Omega_{m0} e^{-3x} + \left(1 - \frac{2}{2-\alpha} \Omega_{m0} \right) e^{-(4-\frac{2}{\alpha})x} \\ &= \frac{\alpha}{2-\alpha} \Omega_{m0} e^{-3x} + \left(\Omega_{de0} - \frac{\alpha}{2-\alpha} \Omega_{m0} \right) e^{-(4-\frac{2}{\alpha})x}, \end{aligned} \quad (6)$$

where Ω_{de0} is the present value of the dimensionless energy density of RDE. Then the corresponding equation of state (EOS) of RDE is obtained by the conservation equation

$$\begin{aligned} w(x) &= -1 - \frac{1}{3} \frac{d \ln \Omega_{de}(x)}{dx} \\ &= \frac{(\Omega_{de0} - \frac{\alpha}{2-\alpha} \Omega_{m0}) \frac{\alpha-2}{\alpha} e^{-(4-\frac{2}{\alpha})x}}{3\Omega_{de}(x)}. \end{aligned} \quad (7)$$

When EOS of dark energy is not constantly equal to -1, it is necessary to consider dark energy perturbations. We works in the conformal Newtonian gauge ¹, the perturbed metric being

$$ds^2 = -a(\eta)^2 \{ (1 + 2\psi) d\eta^2 + (1 - 2\phi) \delta_{ij} dx^i dx^j \}. \quad (8)$$

The perturbed conservation equations of dark energy perturbations [38] read

$$\delta' = -(1+w)(\theta - 3\phi') - 3\mathcal{H}(c_s^2 - w)\delta, \quad (9)$$

$$\theta' = -\mathcal{H}(1-3w)\theta - \frac{w'}{1+w}\theta + \frac{c_s^2}{1+w}k^2\delta + k^2\psi, \quad (10)$$

where δ is the energy density perturbation, θ is the velocity perturbation and is related to the velocity by $\theta = ik^j v_j$ [38]. Where the primes denote the derivatives with respect to conformal time η , $\mathcal{H}$ is the conformal Hubble function and c_s^2 is the general sound speed, being defined

$$c_s^2 \equiv \frac{\delta p}{\delta \rho}. \quad (11)$$

From the intrinsic entropy perturbation Γ [22, 37],

$$w\Gamma \equiv (c_s^2 - c_a^2)\delta = \frac{p'}{\rho} \left(\frac{\delta p}{p'} - \frac{\delta \rho}{\rho'} \right), \quad (12)$$

where the adiabatic speed of sound, c_a^2 , is purely determined by EOS,

$$c_a^2 \equiv \frac{p'}{\rho'} = w - \frac{w'}{3\mathcal{H}(1+w)}, \quad (13)$$

it is found that only in the rest frame where δ keeps gauge invariant can the general sound speed be a gauge invariant quantity since Γ and c_a^2 are gauge and scale independent.

The following transformation gives the useful relation between the gauge invariant, rest frame density perturbation, $\hat{\delta}$, and the density and velocity perturbations in a general frame, δ and θ [22, 37],

$$\hat{\delta} = \delta + 3\mathcal{H}(1+w)\frac{\theta}{k^2}. \quad (14)$$

Thus we can express the pressure perturbation in a general frame, δp , in terms of the general sound speed in the rest frame, $\hat{c}_s^2$, by using Eqs. (12,14)

$$\delta p = \hat{c}_s^2 \delta \rho + 3\mathcal{H}(1+w)(\hat{c}_s^2 - c_a^2)\rho \frac{\theta}{k^2}. \quad (15)$$

Using the above equation and Eq. (13), we have

$$\delta' = -(1+w)(\theta - 3\phi') - 3\mathcal{H}(\hat{c}_s^2 - w)\delta - 3\mathcal{H}(w' + 3\mathcal{H}(1+w)(\hat{c}_s^2 - w))\frac{\theta}{k^2}, \quad (16)$$

$$\theta' = -\mathcal{H}(1-3\hat{c}_s^2)\theta + k^2 \left(\frac{\hat{c}_s^2 \delta}{1+w} + \psi \right). \quad (17)$$

¹ Refer to the reviews of cosmological perturbation theory [36–40] for the detailed gauge transformation.

III. CONSTRAINT METHOD AND RESULTS FROM DATASETS: WMAP, BAO AND SNIA

A. Constraint Method and Data

In the process of performing the observational constraints on cosmological parameters, we adopt the MCMC method by working with the publicly available **CosmoMC**² package [41], including the **CAMB**³ code of calculating the theoretical CMB power spectrum. We modified the code for RDE model with dark energy perturbations. In the first instance, the works of including perturbations of dark energy with EOS on either side of the cosmological constant boundary, i.e. $w > -1$ or $w < -1$, were studied [42]. In these cases, the perturbed equations of dark energy can be solved well. However, when one encounters the perturbations of dark energy with EOS crossing the cosmological constant boundary -1, there inevitably exists a divergent problem due to the term $1+w$ in the denominator in Eq.(17). As pointed out in [19], RDE behaves like a "quintom" when $\alpha < \frac{1}{2}$. Here, we use the method proposed in [24], dividing the whole range of w into three regions with a small positive parameter ϵ : (1) $w > -1 + \epsilon$; (2) $-1 + \epsilon \geq w \geq -1 - \epsilon$; and (3) $w < -1 - \epsilon$. The Eqs. (16,17) are well satisfied in the regions (1) and (3). The divergence happens in the region (2). In order to deal with the divergence and keep dark energy perturbations continuous, the perturbations in the region (2) are set as follows [24]

$$\delta' = 0, \quad \theta' = 0. \quad (18)$$

While in the process of running the program, we set $\epsilon = 10^{-7}$, which is within the limited range, $\epsilon < 10^{-5}$ [24]. By using the method, Refs. [25–28] gave the global fittings on the quintom-like dark energy with parameterized EOS when the dark energy perturbations were included.

We adopt the following 7 dimensional cosmological spaces:

$$P \equiv (\omega_b, \omega_c, \Theta_S, \tau, \alpha, n_s, \log[10^{10} A_s]) \quad (19)$$

where $\omega_b = \Omega_b h^2$ and $\omega_c = \Omega_c h^2$ are the physical baryon and cold dark matter densities, Θ_S is the ratio (multiplied by 100) of the sound horizon and angular diameter distance, τ is the optical depth, α is the newly added Ricci parameter, n_s is the scalar spectral index and A_s is defined as the amplitude of initial power spectrum. The pivot scale of initial scalar power spectrum we have used is $k_{s0} = 0.05 \text{Mpc}^{-1}$. The corresponding priors are taken as follows: $\Omega_b h^2 \in [0.005, 0.1]$, $\Omega_c h^2 \in [0.01, 0.99]$, $\Theta_S \in [0.5, 10]$, $\tau \in [0.01, 0.8]$, $\alpha \in [0.1, 0.8]$, $n_s \in [0.5, 1.5]$ and $\log[10^{10} A_s] \in [2.7, 4]$. In addition, we use a top-hat prior of the cosmic age i.e. $10 \text{Gyr} < t_0 < 20 \text{Gyr}$, impose a weak Gaussian prior on the physical baryon density $\Omega_b h^2 = 0.022 \pm 0.002$ from Big Bang nucleosynthesis [43] and use the new value of Hubble constant $H_0 = 74.2 \pm 3.6 \text{km s}^{-1} \text{Mpc}^{-1}$ [44] by a Gaussian likelihood function.

In our calculations, we have taken the total likelihood $L \propto e^{-\chi^2/2}$ to be the products of the separate likelihoods of CMB, BAO and SNIa. Thus we have

$$\chi^2 = \chi_{CMB}^2 + \chi_{BAO}^2 + \chi_{SNIa}^2. \quad (20)$$

The full CMB data are used, including the new temperature and polarization power spectra from WMAP seven-year data⁴. We combine the additional distance measurements from BAO and SNIa. For BAO information, we use the values of $[r_s(z_d)/D_V(0.2), r_s(z_d)/D_V(0.35)]$ and their inverse covariance matrix [45]. For SNIa, we use the 557 Union2 SNIa data with systematic errors [5]. In order to assess the goodness of fit between RDE model and Λ CDM model on the basis of the same observational data, we also perform the best fitting to the WMAP+BAO+SNIa datasets in the Λ CDM model.

B. Fitting Results and Discussions

In this section, we firstly present the constraint results from observational data. Then using the fitting results, we investigate the imprints on CMB temperature anisotropy, background evolution, and evolutions of perturbations for RDE model with perturbations.

² <http://cosmologist.info/cosmomc/>.

³ <http://camb.info/>.

⁴ <http://lambda.gsfc.nasa.gov/product/map/current/>

Model&Data	RDE&WMAP alone		RDE&WMAP+BAO+SN Ia		Λ CDM &WMAP+BAO+SN Ia	
parameters	Mean	Best Fit	Mean	Best Fit	Mean	Best Fit
$\Omega_b h^2$	$0.0222^{+0.0006+0.0012}_{-0.0006-0.0011}$	0.0222	$0.0241^{+0.0007+0.0014}_{-0.0007-0.0013}$	0.0241	$0.0225^{+0.0005+0.0010}_{-0.0005-0.0010}$	0.0226
$\Omega_c h^2$	$0.1289^{+0.0064+0.0121}_{-0.0063-0.0125}$	0.1313	$0.1087^{+0.0052+0.0104}_{-0.0055-0.0103}$	0.1086	$0.1124^{+0.0033+0.0065}_{-0.0033-0.0063}$	0.1123
Θ_S	$1.0296^{+0.0024+0.0047}_{-0.0023-0.0048}$	1.0293	$1.0302^{+0.0024+0.0048}_{-0.0025-0.0048}$	1.0303	$1.0392^{+0.0026+0.0051}_{-0.0025-0.0050}$	1.0393
τ	$0.0986^{+0.0077+0.0309}_{-0.0092-0.0285}$	0.0931	$0.1374^{+0.0107+0.0417}_{-0.0122-0.0385}$	0.1382	$0.0877^{+0.0063+0.0245}_{-0.0070-0.0230}$	0.0877
α	$0.2310^{+0.0252+0.0526}_{-0.0257-0.0470}$	0.2180	$0.3484^{+0.0217+0.0433}_{-0.0215-0.0407}$	0.3452	-	-
n_s	$1.0159^{+0.0186+0.0385}_{-0.0186-0.0355}$	1.0080	$1.0887^{+0.0181+0.0374}_{-0.0185-0.0351}$	1.0871	$0.9680^{+0.0119+0.0238}_{-0.0119-0.0235}$	0.9691
$\log[10^{10} A_s]$	$3.113^{+0.036+0.073}_{-0.037-0.072}$	3.109	$3.120^{+0.048+0.100}_{-0.047-0.090}$	3.122	$3.187^{+0.036+0.072}_{-0.037-0.071}$	3.180
Ω_{de0}	$0.744^{+0.026+0.047}_{-0.025-0.056}$	0.745	$0.741^{+0.017+0.031}_{-0.017-0.034}$	0.746	$0.727^{+0.014+0.027}_{-0.015-0.03}$	0.728
Age/Gyr	$14.00^{+0.12+0.23}_{-0.11-0.22}$	14.03	$13.83^{+0.12+0.23}_{-0.12-0.23}$	13.82	$13.76^{+0.011+0.021}_{-0.011-0.021}$	13.75
Ω_{m0}	$0.256^{+0.025+0.056}_{-0.026-0.047}$	0.255	$0.259^{+0.017+0.035}_{-0.017-0.031}$	0.254	$0.273^{+0.015+0.03}_{-0.014-0.027}$	0.272
z_{re}	$12.36^{+1.50+2.99}_{-1.50-2.96}$	12.02	$14.49^{+1.68+3.28}_{-1.69-3.41}$	14.61	$10.51^{+1.18+2.34}_{-1.17-2.35}$	10.32
H_0	$77.13^{+3.35+6.56}_{-3.31-6.57}$	77.54	$71.63^{+1.66+3.30}_{-1.63-3.21}$	72.26	$70.32^{+1.28+2.52}_{-1.27-2.47}$	70.38

TABLE I: The mean values with 1σ , 2σ regions and best fit values for RDE model from WMAP alone and the combination of WMAP, BAO and SN Ia, respectively, and the mean values with 1σ , 2σ regions and best fit values for Λ CDM model from the combination of WMAP, BAO and SN Ia.

The constraint results of full basic parameters and derived parameters are showed in Table I, where we list the mean values with 1σ , 2σ regions and best fit values from WMAP alone in the RDE model, WMAP+BAO+SN Ia in the RDE model and WMAP+BAO+SN Ia in the Λ CDM model, respectively. It is seen that the inclusion of distance information from BAO and SN Ia leads to a significant change for best fit value of parameter α . Both the best fit values of the scalar spectral index, n_s , in the RDE model are greater than 1. That is to say, the scalar spectrum is "blue" tilted. This result is different from that in the Λ CDM model, where within 2σ regions the initial power spectrum with a "blue" tilt is not favored. For the combined constraint, the total best-fit χ^2 of RDE model is 8056.1, while $\chi^2 = 8002.6$ for the best-fit Λ CDM model. Therefore, the Λ CDM model still shows a better fit to the current data than the RDE model. In Fig. 1, we give the 1-D marginalized distributions of parameters and 2-D contours with confidence level (C.L.) from the combination of WMAP, BAO, and SN Ia. The respective contributions of WMAP, BAO, and SN Ia to cosmological parameters can be seen in Fig. 2, which shows that the 2-D contours of parameters (α , Ω_{m0}) from WMAP alone, BAO alone, SN Ia alone, and their combinations. The stringent results can be obtained from the joint constraints of dynamical and geometrical aspects.

In Fig. 3, we show CMB temperature power spectra for the RDE model by using the best-fitting results in Table I and for the Λ CDM model with the best fit values from WMAP alone constraint [1] and WMAP+BAO+SN Ia in Table I. It is found that there are significant discrepancies of CMB temperature power spectra in the two models on large angular scales, where the contribution to secondary CMB temperature power spectrum is mainly from the late ISW effect. Both the curves of CMB temperature power spectra for the RDE model on large angular scales almost tend plat. A large-scale ($l < 10$) plateau happens in the RDE model. In order to understand the imprint on CMB temperature power spectrum from parameter α alone, we illustrate the CMB temperature power spectra for different α in Fig. 4. It is shown that CMB temperature power spectra on all the scales decrease as the value of α becomes smaller. Because α has an effect on the expansion rate, the angular diameter distance to recombination becomes larger when α decreases. So there is the rightward shift of the positions of peaks. What's more, the usual dimensionless densities of RDE and matter component are changed for different parameters, α , as shown in Fig. 5. That is to say, the change in the expansion rate brings about the differences of usual dimensionless densities when there is the same matter density, Ω_{mo} . We can see at early times the usual dimensionless matter density decreases when α become larger. We try to look into the role of the change of α by taking another view of point. We consider that this effect can be viewed as coming from the different effective matter densities based on the same expansion rate, i.e. $\Omega_{mo}^{eff}(1+z)^3/E_s^2$. So there is a lower effective matter density with the increase in α . The lower effective matter density causes the delay of epoch of matter-radiation equality, i.e. the smaller z_{eq} . This change in redshift of equality lead to the enhanced late ISW effect on large scales and power spectra on small scales. Furthermore, the delay of the time of dark energy domination for smaller α also leads to the suppressed late ISW effect on large scales. In Fig. 6, we show the evolutions of cold dark matter and dark energy perturbations on the large scale, i.e. $k = 0.001\text{Mpc}^{-1}$.

The differences of background evolutions between the Λ CDM model and RDE model can be seen through the evolutions of the distance moduli and relative distance moduli in Fig. 7. Then following the method of the propagation

FIG. 1: The 1-D marginalized distributions on individual parameters and 2-D contours with 68%C.L., 95 %C.L., and 99.7%C.L. between each other using the combination of the observational data from WMAP, BAO and SNIa for RDE model.

of errors by the Fisher matrix analysis [46–48]⁵, we plot the evolution of EOS with 1σ errors in Fig. 8 by using the mean values from WMAP+BAO+SNIa. The current value of EOS is -1.0853, less than -1. What’s more, it is seen that the EOS of RDE crosses the cosmological constant boundary at $z \sim 0.05$.

IV. SUMMARY

In summary, we perform a global fitting on the RDE model with perturbations to the combined constraints from the full CMB data, BAO, and SNIa by using the MCMC method. All the cosmological parameters for the RDE model are well determined, as shown in Table I. It is worth to notice that the initial power spectrum with a ”blue” tilt is favored for RDE model. Making use of the constraint results, we investigate the CMB temperature power spectra. It is found that the differences of CMB temperature power spectra on large angular scales are significant between the Λ CDM model and the RDE model with perturbations. The main feature of CMB temperature power spectra on large angular scales for the RDE model with perturbations is that there exists a flat plateau. In view of the differences of

⁵ We refer to [49] to marginalize the cosmological parameters which are not associated with EOS and obtain the error covariance matrix of EOS parameters from the sub-Fisher matrix.

FIG. 2: The 2-D contours with 68%C.L., 95 %C.L., and 99.7%C.L. between α and Ω_{m0} for RDE model from WMAP alone (red dashed line), BAO alone (magenta dot-dashed line), SNIa alone (blue dotted line), and the combination of WMAP, BAO and SNIa (solid black line).

large-scale power spectra, on which the ISW effect play an important role, the effects on RDE parameters made by the ISW data from the cross-correlations of CMB and LSS will be worth investigating in the future work.

FIG. 3: CMB temperature power spectra vs. multipole moment l , where the black dots with black error bars denote the observed data with its corresponding uncertainty from WMAP, the red solid line is for Λ CDM model with the best fit values from WMAP alone constraint [1], the green dot-dashed line is for Λ CDM model with the best fit values from WMAP+BAO+SNIa joint constraint. the blue dashed line is for RDE with the best fit values from WMAP alone constraint, and the magenta dotted line is for RDE model with the best fit values from WMAP+BAO+SNIa joint constraint.

FIG. 4: CMB temperature power spectra vs. multipole moment l for RDE model with different parameters, α , where the blue dotted line is power spectrum with $\alpha = 0.44$, the black solid line is power spectrum with the best fit value of α from WMAP+BAO+SNIa joint constraint, and the magenta dashed line is power spectrum with $\alpha = 0.24$.

FIG. 5: The usual dimensionless energy densities of RDE (thin), $\left(\frac{\Omega_{de}}{E^2}\right)$, and matter component (thick), $\left(\frac{\Omega_{m0}(1+z)^3}{E^2}\right)$, vs. redshift z for RDE model with different parameters, α , where the blue dotted lines are dimensionless energy densities with $\alpha = 0.44$, the black solid lines are dimensionless energy densities with the best fit value of α from WMAP+BAO+SNIa joint constraint, and the magenta dashed lines are dimensionless energy densities with $\alpha = 0.24$.

FIG. 6: The evolutions of cold dark matter density perturbation, δ_c (dot-dashed line), dark energy density perturbation, $|\delta_{de}|$ (solid line) and its velocity perturbation, v_{de} (dashed line).

FIG. 7: The distance moduli μ vs. redshift z (left panel), where the grey dots with grey error bars denote the observed data with its corresponding uncertainty from SNIa, the red solid line is for Λ CDM model with the best fit values from WMAP alone constraint [1], the green dot-dashed line is for Λ CDM model with the best fit values from WMAP+BAO+SN Ia joint constraint, the blue dashed line is for RDE model with the best fit values from WMAP alone constraint, and the magenta dotted line is for RDE model with the best fit values from WMAP+BAO+SN Ia joint constraint. The relative distance moduli $\Delta\mu = \mu_{RDE} - \mu_{\Lambda CDM}$ vs. redshift z (right panel), where the magenta solid line is for the best fit values from WMAP alone constraint, and the blue dashed line is for the best fit values from WMAP+BAO+SN Ia joint constraint.

FIG. 8: The evolution of the equation of state with 1σ errors, where the mean values with 1σ errors for RDE model from WMAP+BAO+SN Ia joint constraint are used.

Acknowledgments

This work is supported by NSF (10703001) of P.R. China and the Fundamental Research Funds for the Central Universities (DUT10LK31).

-
- [1] E. Komatsu *et al.*, *Astrophys. J. Suppl.* **192**, 18 (2011), arXiv:1001.4538.
 - [2] M. Tegmark *et al.*, *Phys. Rev. D* **69**, 103501 (2004), astro-ph/0310723.
 - [3] M. Tegmark *et al.*, *Astrophys. J.* **606**, 702 (2004), astro-ph/0310725.
 - [4] A. Clocchiatti *et al.*, *Astrophys. J.* **642**, 1 (2006).
 - [5] R. Amanullah *et al.*, [Supernova Cosmology Project Collaboration], *Astrophys. J.* **716**, 712 (2010), arXiv:1004.1711.
 - [6] A. G. Riess *et al.*, *Astron. J.* **116**, 1009 (1998), astro-ph/9805201.
 - [7] S. Perlmutter *et al.*, *Astrophys. J.* **517**, 565 (1999), astro-ph/9812133.
 - [8] B. Ratra and P. J. E. Peebles, *Phys. Rev. D.* **37**, 3406 (1988);
M. S. Turner and M. White, *Phys. Rev. D* **56**, 4439 (1997);
R. R. Caldwell, R. Dave, and P. J. Steinhardt, *Phys. Rev. Lett.* **80**, 1582 (1998);
P. J. Steinhardt, M. L. Wang, and I. Zlatev, *Phys. Rev. D* **59**, 123504 (1999);
V. Sahni and A. A. Starobinsky, *Int. J. Mod. Phys. D* **9**, 373 (2000);
R. R. Caldwell, *Phys. Lett. B* **545**, 23 (2002);
N. N. Weinberg, *Phys. Rev. Lett.* **91**, 071301 (2003);
S. Nojiri and S. D. Odintsov, *Phys. Rev. D* **72**, 023003 (2005);
B. Feng, X. L. Wang, and X. M. Zhang, *Phys. Lett. B* **607**, 35 (2005);
Z. K. Guo, Y. S. Piao, X. N. Zhang, and Y. Z. Zhang, *Phys. Lett. B* **608**, 177 (2005).
 - [9] A. Y. Kamenshchik, U. Moschella, and V. Pasquier, *Phys. Lett. B* **511**, 265 (2001);
M. C. Bento, O. Bertolami, and A. A. Sen, *Phys. Rev. D* **66**, 043507 (2002);
M. C. Bento, O. Bertolami, M. J. Reboucas, and P. T. Silva, *Phys. Rev. D* **73**, 043504 (2006).
 - [10] M. Li, *Phys. Lett. B* **603**, 1 (2004).
 - [11] M. Li, X. D. Li, S. Wang, and Y. Wang, arXiv:1103.5870.
 - [12] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, *Phys. Rev. Lett.* **82**, 4971 (1999), hep-th/9803132.
 - [13] C. Gao, X. Chen, and Y. G. Shen, *Phys. Rev. D* **79**, 043511 (2009), arXiv:0712.1394.
 - [14] L. N. Granda and A. Oliveros, *Phys. Lett. B* **669**, 275 (2008), arXiv:0810.3149;
Y. T. Wang and L. X. Xu, arXiv:1004.3340.
 - [15] L. X. Xu, J. B. Lu, and W. B. Li, *Eur. Phys. J. C* **64**, 89 (2009).
 - [16] R.-J. Yang, Z.-H. Zhu, and F. Wu, *Int. J. Mod. Phys. A* **26**, 317 (2011).
 - [17] R. G. Cai, B. Hu and Y. Zhang, *Commun. Theor. Phys.* **51**, 954 (2009).
 - [18] L. Xu, W. Li, J. Lu, and B. Chang, *Mod. Phys. Lett. A* **24**, 1355 (2009);
M. Li, X. Li, and X. Zhang, arXiv:0912.3988;
M. Li, X. D. Li, S. Wang, and X. Zhang, *JCAP* **0906**, 036 (2009);
L. X. Xu and Y. T. Wang, *JCAP* **06**, 002 (2010).
 - [19] X. Zhang, *Phys. Rev. D* **79**, 103509 (2009);
 - [20] H. Li, J. Q. Xia, G. B. Zhao, Z. H. Fan, and X. M. Zhang, *Astrophys. J.* **683**, L1-L4 (2008).
 - [21] C. G. Park, J. C. Hwang, J. H. Lee, and H. Noh, *Phys. Rev. Lett.* **103**, 151303 (2009).
 - [22] R. Bean and O. Dore, *Phys. Rev. D* **69**, 083503 (2004), astro-ph/0307100.
 - [23] J. Weller and A. M. Lewis, *Mon. Not. Roy. Astron. Soc.* **346**, 987 (2003).
 - [24] G. B. Zhao, J. Q. Xia, M. Li, B. Feng, and Xinmin Zhang, *Phys. Rev. D* **72**, 123515 (2005), astro-ph/0507482.
 - [25] J. Q. Xia, G. B. Zhao, B. Feng, H. Li, and Xinmin Zhang, *Phys. Rev. D* **73**, 063521 (2006), astro-ph/0511625.
 - [26] G. B. Zhao, J. Q. Xia, B. Feng, and Xinmin Zhang, *Int. J. Mod. Phys. D* **16**, 1229 (2007), astro-ph/0603621.
 - [27] J. Q. Xia, Y. F. Cai, T. T. Qiu, G. B. Zhao, and Xinmin Zhang, *Int. J. Mod. Phys. D* **17**, 1229 (2008), astro-ph/0703202.
 - [28] J. Q. Xia, H. Li, G. B. Zhao, and Xinmin Zhang, *Phys. Rev. D* **78**, 083524 (2008), arXiv:0807.3878;
 - [29] R. K. Sachs and A. M. Wolfe, *Astrophys. J.* **147**, 73 (1967).
 - [30] B. M. Schaefer, *Mon. Not. Roy. Astron. Soc.* **388**, 1403 (2008), arXiv:0803.2239.
 - [31] Y. T. Wang, Y. X. Gui, L. X. Xu, and J. B. Lu, *Phys. Rev. D* **81**, 083514 (2010).
 - [32] G. Olivares, F. A. Brandela, and D. Pavon, *Phys. Rev. D* **77**, 103520 (2008).
 - [33] J. B. Dent, S. Dutta, and T. J. Weiler, *Phys. Rev. D* **79**, 023502 (2009).
 - [34] D. Sapone and M. Kunz, *Phys. Rev. D* **80**, 083519 (2009).
 - [35] Y. T. Wang, L. X. Xu, and Y. X. Gui, *Phys. Rev. D* **82**, 083522 (2010).
 - [36] J. M. Bardeen, *Phys. Rev. D* **22**, 1882 (1980).
 - [37] H. Kodama, and M. Sasaki, *Prog. Theo. Phys. Supp.* **78**, 1 (1984).
 - [38] C. P. Ma and E. Bertschinger, *Astrophys. J.* **455**, 7 (1995).
 - [39] W. Hu and D. J. Eisenstein, *Phys. Rev. D* **59**, 083509 (1999).
 - [40] K. A. Malik and D. Wands, *Phys. Rept.* **475**, 1 (2009).

- [41] A. Lewis and S. Bridle, Phys. Rev. D **66**, 103511 (2002).
- [42] Ch. Yèche, A. Ealet, A. Refregier, C. Tao, A. Tilquin, J.-M. Virey, and D. Yvon, astro-ph/0507170;
P. S. Corasaniti, M. Kunz, D. Parkinson, E. J. Copeland, and B. A. Bassett, Phys. Rev. D **70**, 083006 (2004).
- [43] S. Burles, K. M. Nollett, and M. S. Turner, Astrophys. J. **552**, L1 (2001).
- [44] A. G. Riess *et al.*, Astrophys. J. **699**, 539 (2009), arXiv:0905.0695.
- [45] W. J. Percival *et al.*, Mon. Not. Roy. Astron. Soc. **401**, 2148 (2010), arXiv:0907.1660.
- [46] W. H. Press *et al.*, Numerical Recipes, Cambridge University Press (1994).
- [47] U. Alam, V. Sahni, T. D. Saini, and A. A. Starobinsky, astro-ph/0406672.
- [48] S. Nesseris and L. Perivolaropoulos, Phys. Rev. D **72**, 123519 (2005), astro-ph/0511040.
- [49] T. D. Kitching and A. Amara, arXiv:0905.3383.