



# CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Inferring core-collapse supernova physics with gravitational waves

J. Logue, C. D. Ott, I. S. Heng, P. Kalmus, and J. H. C. Scargill

Phys. Rev. D **86**, 044023 — Published 17 August 2012

DOI: [10.1103/PhysRevD.86.044023](https://doi.org/10.1103/PhysRevD.86.044023)

# Inferring Core-Collapse Supernova Physics with Gravitational Waves

J. Logue,<sup>1,2,\*</sup> C. D. Ott,<sup>3,4,5,†</sup> I. S. Heng,<sup>1,‡</sup> P. Kalmus,<sup>2,3,§</sup> and J. H. C. Scargill<sup>6,2,¶</sup>

<sup>1</sup>*SUPA, Institute for Gravitational Research, School of Physics and Astronomy,  
University of Glasgow, Glasgow G12 8QQ Scotland, United Kingdom*

<sup>2</sup>*LIGO Laboratory, California Institute of Technology, Pasadena, CA 91125, USA*

<sup>3</sup>*TAPIR, MC 350-17, California Institute of Technology, Pasadena, CA 91125, USA*

<sup>4</sup>*Institute for the Physics and Mathematics of the Universe (IPMU), The University of Tokyo, Kashiwa, Japan*

<sup>5</sup>*Center for Computation and Technology, Louisiana State University, Baton Rouge, LA, USA*

<sup>6</sup>*New College, Oxford, OX1 3BN, UK*

Stellar collapse and the subsequent development of a core-collapse supernova explosion emit bursts of gravitational waves (GWs) that might be detected by the advanced generation of laser interferometer gravitational-wave observatories such as Advanced LIGO, Advanced Virgo, and LCGT. GW bursts from core-collapse supernovae encode information on the intricate multi-dimensional dynamics at work at the core of a dying massive star and may provide direct evidence for the yet uncertain mechanism driving supernovae in massive stars. Recent multi-dimensional simulations of core-collapse supernovae exploding via the neutrino, magnetorotational, and acoustic explosion mechanisms have predicted GW signals which have distinct structure in both the time and frequency domains. Motivated by this, we describe a promising method for determining the most likely explosion mechanism underlying a hypothetical GW signal, based on Principal Component Analysis and Bayesian model selection. Using simulated Advanced LIGO noise and assuming a single detector and linear waveform polarization for simplicity, we demonstrate that our method can distinguish magnetorotational explosions throughout the Milky Way ( $D \lesssim 10$  kpc) and explosions driven by the neutrino and acoustic mechanisms to  $D \lesssim 2$  kpc. Furthermore, we show that we can differentiate between models for rotating accretion-induced collapse of massive white dwarfs and models of rotating iron core collapse with high reliability out to several kpc.

PACS numbers: 04.30.Tv, 04.80.Nn, 05.45.Tp, 97.60.Bw

## I. INTRODUCTION

Almost eighty years after the proposal by Baade & Zwicky that (core-collapse) supernovae represent the transition of an ordinary massive star into a neutron star [1], we still lack a complete understanding of this phenomenon. In particular, we do not know with certitude how the *supernova mechanism* operates and converts the necessary fraction of gravitational energy of collapse into kinetic energy and light of the explosive outflow.

The basic story line of core collapse goes as follows (see [2, 3] for detailed reviews): At the end of a massive star's ( $8 - 10 M_{\odot} \lesssim M \lesssim 130 M_{\odot}$  at zero-age main sequence [ZAMS]) life, nuclear burning has ceased in its core, which is then composed primarily of iron-group nuclei (or O-Ne nuclei at the lower end of the mass range) and supported against gravity by the pressure of relativistically degenerate electrons. Eventually, the core exceeds its effective Chandrasekhar mass and dynamical collapse sets in. The collapsing core separates into subsonically infalling homologous ( $v \propto r$ ) inner core and supersonically collapsing outer core [4, 5]. When the inner

core reaches nuclear density, the repulsive core of the nuclear force leads to a stiffening of the nuclear equation of state (EOS). The inner core, suddenly supported by the stiff supernuclear EOS, overshoots its new equilibrium, then bounces back into the still infalling outer core. A shock wave forms at the sonic point between inner and outer core at an enclosed baryonic mass of  $\sim 0.5 M_{\odot}$ . It quickly moves out in radius and mass, but must do work in breaking up infalling iron-group nuclei. This and neutrino losses from electron capture in the region behind the shock sap its might. The shock succumbs to the ram pressure of the outer core, stalls, and turns into an accretion shock.

The shock must be re-energized to drive a core-collapse supernova explosion and, in the canonical scenario, leave behind a neutrino-cooling and contracting protoneutron star. This shock revival must, depending on progenitor star structure, occur within  $\sim 0.5 - 3$  s, otherwise accretion will push the protoneutron star over its maximum mass, leading to collapse and black hole formation [6]. Understanding the supernova mechanism, which must robustly revive the stalled shock in supernovae from massive stars that are observed on a daily basis, is the principle current challenge of core-collapse supernova theory.

Observational clues for the supernova mechanism are few. Electromagnetic waves are emitted in optically thin regions far from the core and thus yield only second-hand information about the supernova mechanism. Yet, observations of ejecta morphology, spatial distributions of nucleosynthetic yields and pulsar kicks are indicative

---

\*Electronic address: j.logue@physics.gla.ac.uk

†Electronic address: cott@tapir.caltech.edu

‡Electronic address: I.k.Heng@glasgow.ac.uk

§Electronic address: kalmus@caltech.edu

¶Electronic address: james.scargill@new.ox.ac.uk

of aspherical (i.e., multi-dimensional) processes bearing relevance in the explosion dynamics (e.g., [7, 8] and references therein). Neutrinos, on the other hand, are emitted deep inside the core and can provide crucial thermodynamic, structural, and, to some extent, dynamical information on what occurs in the core [2, 9, 10]. The few neutrinos captured from supernova 1987A [2, 11] have impressively confirmed the very basic picture of core collapse outlined in the above.

Gravitational waves (GWs), like neutrinos emitted from dense regions impenetrable by photons, carry dynamical information about their source. Since their emission occurs at lowest order by accelerated quadrupole motions, GWs are direct probes of multi-dimensional dynamics in the core that may play a crucial role in the supernova mechanism [3, 12].

Stellar collapse has long been considered a promising source of GWs for detectors on Earth (see the historical overview in [12]) and much effort has gone into understanding the GW signature of stellar collapse and the subsequent evolution towards a core-collapse supernova explosion. This has led to the identification of a range of emission processes, including rotating collapse and core bounce, nonaxisymmetric rotational instabilities, aspherical outflows, convection/turbulence in the protoneutron star and in the region immediately behind the shock, instabilities of the standing accretion shock, pulsations of the protoneutron star, asymmetric emission of neutrinos, and magnetic stresses (see [12, 13] for recent reviews). The most recent set of simulations [12, 14–19] suggest that GWs from the average core-collapse supernova may be visible throughout the Milky Way for the second generation of laser interferometer GW observatories, including Advanced LIGO, Advanced Virgo, and LCGT [20, 21]. Extreme emission scenarios may allow detection throughout the local group of galaxies, including the Andromeda galaxy [12, 22–24], but third-generation detectors such as the Einstein Telescope [25] might hope to observe more distant events.

If it was possible to associate an explosion mechanism with particular multi-dimensional dynamics that leads to a characteristic GW signal, then the detection or non-detection of such a signal from the next galactic core-collapse supernova could confirm or rule out this mechanism. To realize such an GW-observational test of the explosion mechanism, one must separate the signal from detector noise and determine its parameters, e.g., by matching, in some way, to signal predictions from simulations. The most straightforward method for signal extraction and parameter estimation is matched filtering (e.g., [26]), which looks for a match of detector data with waveform templates from simulations. Matched filtering requires exact knowledge of the expected signal, which is possible, e.g., for the inspiral phase of compact binaries, since the parameter space of binary systems is limited, all relevant physics is understood (at least in the black-hole – black-hole binary case) and numerical relativity simulations can predict essentially exact wave-

form templates (i.e., limited only by numerical error). However, building waveform catalogs with exact predictions for matched filtering is impossible for GWs from core-collapse supernovae. On the one hand, there is unknown physics (e.g., the nuclear EOS) and many unconstrained parameters (e.g., the details of the precollapse configuration are poorly known) in the stellar collapse problem. This alone would require extensive parameter studies to build up template databases covering the poorly constrained parameter space. On the other hand, all expected GW emission processes in core-collapse supernovae are influenced or dominated by turbulent flow. Hence, their GW signals have a stochastic component that is impossible to predict, even if all initial conditions and physics were known exactly. Matched filtering is not applicable to such GW bursts. To extract the GW signal from the next galactic core collapse event and determine source physics such as the explosion mechanism, an approach to signal extraction/reconstruction, model selection, and parameter estimation is needed that can handle the stochastic nature of the expected GW signals.

The reconstruction of both polarizations of a GW signal requires coincident observations of two detectors; linearly polarized signals can be reconstructed from data of just one detector and adding a third detector overdetermines the problem, permitting the source position on the sky to be determined [27]. GW signal reconstruction was pioneered by Gürsel & Tinto [28] with a maximum likelihood approach, variants [29, 30] of which have been incorporated into search pipelines for GW bursts [31].

Summerscales *et al.* [32] were the first to study signal reconstruction and parameter estimation for the GW burst from rotating core collapse and bounce based on waveforms of Ott *et al.* [33]. They injected signals into real detector noise of early LIGO science runs and used a maximum entropy approach to reconstruct the signal using data from two detectors without any a priori knowledge of the signal shape. Cross-correlation of the reconstructed signal with signal predictions of [33] was then used to determine source parameters.

Incorporating GW signal information from core collapse simulations into detection and signal reconstruction approaches was first considered by Brady & Ray-Majumder [34], who realized that the GW burst from rotating collapse and bounce, while being unpredictable in detail, has robust features that can be isolated mathematically. They created an orthonormal vectorspace of waveforms from [33, 35] using Gram-Schmidt orthonormalization and isolated a subspace of essential features most common to all waveforms. Heng [36] also considered waveforms from rotating core collapse and bounce and utilized the more recent waveform catalog of Dimmelmeier *et al.* [17]. He performed Principal Component Analysis (PCA; e.g., [37]), which transforms a correlated, multi-dimensional data set into a set of orthogonal components by determining the eigenvectors and eigenvalues of the covariance matrix of the data set. The principal component (PC) vectors are the eigenvectors ranked ac-

ording to their corresponding eigenvalue, with the first PC being the eigenvector with the largest eigenvalue. Cannon *et al.* [38] have also utilised this method for GW signals from compact binary coalescence.

Röver *et al.* [39] went a step further and combined PCA with Bayesian inference using the Markov-Chain Monte Carlo technique for computing marginalization integrals (see, e.g., [40] for a pedagogical introduction to Bayesian methods). They considered linearly-polarized waveforms from rotating core collapse and bounce and were able to reconstruct signals from modeled noise in a single detector and infer key parameters, e.g., the nuclear EOS used in the simulation that led to a given trial waveform.

In this paper, we present a proof-of-principle study to demonstrate that the core-collapse supernova explosion mechanism can be inferred from the GW signal of a galactic core-collapse supernova observed with second-generation GW observatories such as Advanced LIGO, Advanced Virgo, and LCGT. We consider the neutrino mechanism (e.g., [3]), the magnetorotational mechanism (e.g., [41]), and the acoustic mechanism [42, 43], discuss their essentials, and argue that they bear distinct GW signatures as first pointed out by Ott [12, 44], which is a prerequisite for our study. We follow the approach of Röver *et al.* [39] and, for simplicity, restrict ourselves to a single detector, linearly polarized signals, and a Gaussian noise model at the noise level of Advanced LIGO in broadband mode. Like Röver *et al.*, we adopt a Bayesian approach and use PCA, but, for the first time, apply it to multiple waveform catalogs. We associate each waveform catalog with one of the three mechanisms and calculate Bayes evidence ratios using the nested sampling algorithm [40, 45] to determine what mechanism's PCs match best with a given injected signal. We demonstrate that this approach can identify any of the considered explosion mechanisms with high confidence for core collapse events occurring at distances of up to  $\sim 2$  kpc. The magnetorotational explosion mechanism can even be inferred throughout the Milky Way ( $D \gtrsim 10$  kpc). In addition to studying the explosion mechanism, we also consider the problem of determining whether a core collapse event is a rapidly rotating ordinary iron core collapse or an accretion-induced collapse (AIC) of a massive white dwarf. These two processes are governed by the same physics and differences in their waveforms are subtle [46], but our approach is still capable of telling them apart.

This article is structured as follows: Section II reviews the set of considered candidate core-collapse supernova mechanisms, their individual GW signatures, and the GW signal catalogs that we draw model waveforms from. In Section III, we introduce our method for model selection via PCA and nested sampling. The results of our study are presented and discussed in detail in Section IV. We summarize and conclude in Section V.

## II. SUPERNOVA MECHANISMS AND THEIR GRAVITATIONAL WAVE SIGNATURES

In this study, we consider the neutrino mechanism, the magnetorotational mechanism, and the acoustic mechanism for core-collapse supernova explosions and describe them and their characteristic GW signal features in the following sections. Variations of these mechanisms and alternatives have been discussed elsewhere (e.g., [3, 47]).

### A. Neutrino Mechanism

The gravitational collapse of the iron core and the subsequent evolution of the nascent hot puffed-up protoneutron star to a cold compact neutron star release of order 300 B (1 Bethe =  $10^{51}$  erg) of energy,  $\sim 99\%$  of which is emitted in the form of neutrinos of all flavors [2]. If only a small fraction of the energy released in neutrinos is reabsorbed behind the stalled shock, leading to net heating, an explosion could be launched and endowed with the energy to account for the observed range of asymptotic explosion energies of 0.1-1 B of garden-variety core-collapse supernovae [48]. This is the gist of the neutrino mechanism of core-collapse supernovae, which, in its early form was proposed by Arnett [49] and Colgate & White [50], and in its modern form by Bethe & Wilson [51].

Despite its appealing simplicity, the neutrino mechanism, in its purest, spherically-symmetric (1D) form, fails to revive the shock in all but the lowest-mass massive stars with O-Ne cores [52–55]. There is now strong evidence from axisymmetric (2D) [14, 19, 56–60] and first 3D [15, 16, 61–66] simulations that the breaking of spherical symmetry is key to the success of the neutrino mechanism. In 2D and 3D, neutrino-driven convection in the region of net heating behind the shock, and the standing-accretion-shock instability (SASI) [67–70] increase the efficiency of the neutrino mechanism [60, 62, 63, 65].

Leaving rapid rotation aside for a moment, the dominant multi-dimensional GW-emitting dynamics in neutrino-driven core-collapse supernovae are convection in the protoneutron star (e.g., [72, 73]) and SASI-modulated convection in the region behind the stalled shock. GW emission from convection and SASI has been extensively studied in simulations in 2D [58, 71–74] and to some extent in 3D [15, 16, 75, 76]. The top panel in Fig. 1 shows a typical example waveform drawn from the catalog of Murphy *et al.* [71]. Right after core bounce, an initial burst of GWs is emitted by strong, so-called prompt convection [12], driven by the negative entropy gradient left behind by the stalling shock. Subsequently, the GW signal settles at lower amplitudes, then picks up again as the SASI reaches its non-linear phase and high-velocity accretion downstreams penetrate deep into the region behind the shock, where they are decelerated, leading to pronounced spikes in the wave signal [71]. The secular rise in the signal amplitude towards the end of the waveform is due to the onset of an aspherical explosion




FIG. 1: Linearly polarized GW signal predictions for a core collapse event located at 10 kpc from matter dynamics in axisymmetric simulations that can be associated with the neutrino mechanism (top panel, taken from [71]), the magnetorotational mechanism (center panel, taken from [17]), and the acoustic mechanism (bottom panel, taken from [12]). Note the varying ranges of the time and strain axes. Also note that the simulations of [17] did not include magnetic fields, since the GW signal from core bounce in magnetorotational explosions is due to rapid rotation and not influenced by magnetic fields. This, however, did not allow them to capture the expected secular rise of the waveform expected to occur once an explosion sets in [18]. See text for further discussion.

[15, 58, 71], but occurs at too low characteristic frequencies to be visible to Advanced LIGO-class detectors. Not included in the top panel of Fig. 1 is the contribution to the GW signal from anisotropic neutrino emission [77–79], which can dominate in amplitude, but, like the contribution from aspherical outflow, occurs on timescales too long to lead to emission at frequencies detectable by the upcoming generation of ground-based detectors [15, 16, 58, 72, 74, 76].

Overall, the detectable GW signal from a neutrino-driven nonrotating or slowly rotating core-collapse super-

nova will have random polarization, a broadband spectrum from  $\sim 100 - 1000$  Hz and typical strain amplitudes  $|h|$  of order  $10^{-22} (D/10 \text{ kpc})^{-1}$ , with individual peaks reaching  $10^{-21} (D/10 \text{ kpc})^{-1}$  [12, 15]. The typical duration of emission is the time from core bounce to the launch of the explosion,  $0.3 - 1$  s, but convection inside the cooling proton-neutron star can continue to emit GWs at lower amplitudes and higher frequencies for many seconds afterwards [12, 72]. Typical total emitted GW energies are in the range  $10^{-11} - 10^{-9} M_{\odot} c^2$  [12, 58, 71].

The effects of rotation on the neutrino mechanism and its GW signature are not yet fully understood (see, e.g., [14, 19, 44, 57, 59, 80, 81]) and it cannot be excluded that contributions from rotational dynamics may modify the GW signal of neutrino-driven core-collapse supernovae. However, results from the systematic rotating core-collapse studies of [17, 18, 41, 81] suggest that once rotation rates become sufficiently high to alter the dynamics, the explosion is actually more likely to occur via the magnetorotational mechanism discussed in §II B. This, however, is under the provision that the magnetorotational instability (e.g., [82, 83]) works robustly and builds up the required strong magnetic fields to drive an explosion.

Keeping the above caveats in mind, for the purpose of this study, we make the assumption that the GW signature of neutrino-driven core-collapse supernovae is unaffected by rotational effects.

### 1. GW Signal Catalogs

In this study, we use the catalog of Murphy *et al.* [71], which is available for download from [84]. The Murphy *et al.* catalog (in the following, we refer to waveforms from this catalog as Mur waveforms) encompasses 16 waveforms that were extracted via the quadrupole formula (e.g., [85]) from Newtonian axisymmetric core-collapse simulations that used a parameterized scheme for electron capture and neutrino heating/cooling and included only the monopole component of the gravitational potential as described in [60, 71]. The Murphy *et al.* simulations are nonrotating and the parameter space covered is spanned by progenitor ZAMS mass ( $\{12, 15, 20, \text{and } 40\} M_{\odot}$ ) and by the dialed-in total electron and anti-electron neutrino luminosity.

Yakunin *et al.* [58] performed self-consistent axisymmetric Newtonian (with an approximate-GR monopole term of the gravitational potential [86]) radiation-hydrodynamics simulations of neutrino-driven core-collapse supernovae. They provide three waveforms at [87], obtained from simulations using progenitors of (12, 15, and 25)  $M_{\odot}$ . We use the Yakunin waveforms (denoted, in the following, as Yak waveforms) to test the robustness of our supernova mechanism determination algorithm, which uses the PCs of the Mur waveforms.

Since we are limiting ourselves to one detector in this proof-of-principle study, we are considering only linearly polarized signals. Gravitational waveforms with + and

× polarizations from 3D simulations of neutrino-driven core-collapse supernovae [15, 16, 76] will be considered in future work.

## B. Magnetorotational Mechanism

The conservation of angular momentum in core collapse to a protoneutron star leads to a spin-up by a factor of  $\sim 1000$  [88]. Starting from a precollapse angular velocity distribution that may be expected to be more or less uniform in the inner core (e.g., [89]), homologous collapse preserves the uniform rotation of the inner core while the supersonic collapse of the outer core leads to strong differential rotation in the outer protoneutron star and in the region between protoneutron star and shock [88].

A rapidly spinning precollapse core with a period of order 1 s results in a ms-period protoneutron star, with a rotational energy of order 10 B, which is about ten times greater than the typical core-collapse supernova explosion energy. If only a fraction of this energy was tapped, a strong explosion could be triggered.

Theory and simulations (e.g., [18, 41, 90–96]) have shown that magnetorotational processes are efficient at extracting spin energy and can drive collimated outflows, leading to energetic bipolar jet-like explosions. Recent work [18, 41, 94–96] suggests that magnetic fields of the order of  $10^{15}$  G with strong toroidal components are required to yield the necessary magnetic stresses to drive a strong bipolar explosion. If  $10^{15}$  G fields were to arise from flux compression in collapse alone, precollapse core fields would have to be of order  $10^{12}$  G [41, 94], which is about 3 orders of magnitude larger than predicted by stellar evolution models (e.g., [89, 97]). It is more likely that the most significant amplification occurs after core bounce via rotational winding of poloidal into toroidal field (a linear process), the non-linear magnetorotational instability (MRI, which is not yet fully understood in the core collapse context [82, 83]). Both processes operate on the free energy stored in differential rotation, which is abundant in the outer core.

For the magnetorotational mechanism to work, precollapse spin periods  $\lesssim 4 - 5$  s appear to be required [41]. Such rapid rotation leads to a strongly centrifugally-deformed inner core with a large quadrupole moment ( $\ell = 2$ ; due to its oblateness), which rapidly changes during core bounce, leading to a strong burst of GWs. The GW signal from rotating collapse and bounce has been studied extensively and the most recent general-relativistic simulations have shown it to be of rather generic morphology with a single strong peak at bounce and a subsequent ringdown as the protoneutron star core settles into its new equilibrium [17, 98, 99]. A typical example GW signal taken from the catalog of Dimmelmeier *et al.* [17, 100] is shown in the center panel of Fig. 1. The core collapse and bounce phase proceeds essentially axisymmetrically even in very rapidly spinning

cores [99, 101, 102] and its GW signal is linearly polarized with vanishing amplitude seen by an observer located along the symmetry axis and maximum amplitude for an equatorial observer. Typical emission durations for the linearly polarized GWs from core bounce are of order 10 ms and peak GW amplitudes for rapidly spinning cores that may lead to magnetorotational explosions are of order  $10^{-21} - 10^{-20}$  at 10 kpc with most of the energy being emitted around 500 – 800 Hz in cores that reach nuclear density and bounce due to the stiffening of the nuclear EOS. Cores with initial spin periods shorter than  $\sim 0.5 - 1$  s experience a slow bounce at sub-nuclear densities strongly influenced or dominated by the centrifugal force. They emit most of the GW energy at frequencies below  $\sim 200$  Hz [12, 17]. Typical emitted GW energies are in the range  $10^{-10} - 10^{-8} M_{\odot} c^2$ . The GW signal from rotating collapse and core bounce is unlikely to be affected by MHD effects, since the build up to dynamically relevant field strengths occurs only after bounce [18, 41, 92, 93].

Due to the strong rotational deformation of the protoneutron star, neutrinos decouple from the matter at smaller radii and hotter temperatures in polar regions than near the equator. This leads to the emission of a larger neutrino flux with a harder neutrino spectrum in polar regions (e.g., [59]). This globally asymmetric neutrino emission results in a secularly rising low-frequency GW signal [12]. Similar low-frequency contributions will come from the bipolar outflow characteristic for a magnetorotational explosion and from magnetic stresses [18, 92, 93]. The low-frequency waveform components are not shown in the center panel of Fig. 1 and are not detectable by the upcoming second-generation earthbound GW observatories.

Also associated with rapid rotation and the magnetorotational mechanism are rotational instabilities that may lead to nonaxisymmetric deformations of the protoneutron star whose “bar-mode” ( $m = 2$ ) components may emit elliptically polarized GWs for tens to hundreds of milliseconds [99, 101–104]. However, these instabilities, and in particular their interplay with magnetic fields and the MRI (see, e.g., [105]), are not yet fully understood. Since we are considering only linearly polarized signals and are limited to one detector, we do not include GW signals from these nonaxisymmetric instabilities in this study.

### 1. GW Signal Catalogs

We employ the large (128 waveforms) GW signal catalog of Dimmelmeier *et al.* [17, 100] (Dim in the following), who performed 2D GR simulations of rotating iron core collapse for (11.2, 15, 20, and 40)  $M_{\odot}$  progenitors and two different nuclear EOS, varying initial rotation rate and degree of differential rotation. They approximated the effects of electron capture during collapse by parametrizing the electron fraction  $Y_e$  as a function of

density, which yields inner core sizes that are very close to those obtained with full neutrino transport [106]. The inner core size determines the amount of mass and angular momentum that can be dynamically relevant during core bounce and, hence, is a determining factor in the GW signal [98]. The `Dim` catalog was also used by the previous parameter estimation work of Röver *et al.* [39]. For testing, we use the three additional `Dim` waveforms computed for [39] that are not part of the original `Dim` catalog and were used to test their algorithm. We label this set of extra waveforms as `DimExtra`.

For studying the robustness of our mechanism-determination approach, we draw gravitational waveforms of rotating models from the catalog of Scheidegger *et al.* [102, 107], (`Sch` in the following) who performed 3D Newtonian-MHD rotating iron core collapse calculations with a spherical approximate-GR gravitational potential and employed the same EOS and electron capture treatment as Dimmelmeier *et al.* [17], but used different progenitor models.

Furthermore, we use the GW signal catalog of Abdikamalov *et al.* [46, 84] (`Abd` in the following) who used the same numerical code as Dimmelmeier *et al.* [17], but studied the rapidly spinning accretion-induced collapse (AIC) of massive white dwarfs to neutron stars. This process yields a GW signal very similar to rotating iron core collapse and explosions in AIC may occur also via the magnetorotational mechanisms [108]. We include this catalog of 106 waveforms to see if our algorithm can differentiate between rotating iron core collapse and rotating AIC assuming the `Dim` and `Abd` catalogs correctly predict the respective GW signals.

### C. Acoustic Mechanism

The core-collapse supernova evolution in the *acoustic mechanism* proposed by Burrows *et al.* [22, 42, 43] is initially identical to the one expected for the neutrino mechanism. Neutrino heating, convection and the SASI set the stage, but no explosion is triggered for  $\gtrsim 500$  ms after bounce. At this point, the SASI is in its highly non-linear phase and modulates high-velocity accretion downflows that impact on the protoneutron star and excite core pulsations (primarily  $\ell = \{1, 2\}$   $g$ -modes). Over hundreds of milliseconds, these pulsations reach large amplitudes and damp via the emission of strong sound waves. Traveling down the steep density gradient in the region behind the shock, the sound waves steepen to shocks and dissipate their energy behind and in the shock. This mechanism is robust in the simulations by Burrows *et al.* [22, 42, 43], but requires  $\gtrsim 1$  s to develop, thus leads to massive NSs, and tends to yield explosion energies on the lower side of what is observed.

The GW signature of the acoustic mechanism is dominated by the strong emission from the quadrupole components of the protoneutron star core pulsations that are quasi-periodic (their frequency shifts secularly along with

the changing protoneutron star structure) and become very strong  $\gtrsim 800 - 1000$  ms after core bounce [12, 22]. The lower panel of Fig. 1 depicts a typical example waveform from Ott *et al.* [12, 22], who studied the GW signature of the acoustic mechanism based on the simulations of Burrows *et al.* [42, 43]. At early times, the GW signal is essentially the same as expected for the neutrino mechanism, but once the protoneutron star core pulsations grow strong, they are hard to miss. The simulations of Burrows *et al.* [22, 42, 43] were axisymmetric and the resulting GW signals are linearly polarized, though in 3D, one would expect oscillation power also in non-axisymmetric components. Typical maximum strain amplitudes are of order  $\text{few} \times 10^{-21} - 10^{-20}$  and multiple modes with frequencies between  $\sim 600 - 1000$  Hz contribute to the emission. Since the pulsations last for many cycles, the emitted GW energies may be large and are predicted to be of order  $10^{-8} - 10^{-7} M_{\odot} c^2$  and extreme models reach  $\text{few} \times 10^{-5} M_{\odot} c^2$  [12, 22].

There are multiple caveats associated with the acoustic mechanism that must be mentioned. Most importantly, the acoustic mechanism has been found in simulations of only one group with a single simulation code, but others have not yet ruled out the possibility of strong protoneutron star pulsations at late times (e.g., [19]). In a non-linear perturbation study, Weinberg & Quataert [109] found that the protoneutron star pulsation amplitudes may be limited by a parametric instability involving high-order modes that damp efficiently via neutrino emission and are not presently resolved in numerical simulations. This would limit the protoneutron star pulsations to dynamically insignificant amplitudes. Moreover, the simulations of Burrows *et al.* were axisymmetric and nonrotating or only very slowly rotating. It is not clear to what amplitudes individual protoneutron star pulsation modes would grow in 3D. Rapid rotation, due to its stabilizing effect on convection and SASI [44, 59], may likely inhibit the growth of pulsations. Both 3D and rotational effects remain to be explored.

#### 1. GW Signal Catalogs

We employ the set of 7 waveforms from the models of [43] analyzed by Ott [12] and available at [84]. We refer to this set as the `Ott` catalog in the following and use them to compute PCs for the acoustic mechanism's GW signature. All waveforms were computed on the basis of the Burrows *et al.* [22, 42, 43] simulations and differ only in the employed progenitor model, covering a range in ZAMS mass from 11.2 to 25  $M_{\odot}$ .

Three additional waveforms of an earlier study of Ott *et al.* [22] are available [84]. We label this small set `OttExtra` and use them for testing our method's capability of correctly identifying them as coming from stars exploding via the acoustic mechanism.

### III. DATA ANALYSIS

#### A. Strategy

The three example gravitational waveforms shown in Fig. 1 that are associated with the three supernova mechanisms are clearly different. Provided the assumptions made in associating these signals with the various mechanisms are correct, a GW signal detected from a core-collapse supernova should, in principle, allow to determine the explosion mechanism. To do so in practice, two problems must be overcome: (i) The exact waveform of an incident signal is impossible to predict in advance. (ii) Real GW detectors are noisy instruments (see, e.g., [110] for a discussion of detectors and noise sources) and any GW signal will be contaminated by detector noise. In other words, it is necessary to develop a data analysis algorithm that is capable of distinguishing between underlying physical models (e.g., supernova mechanisms) on the basis of a noisy signal whose detailed shape cannot be predicted exactly.

In the following subsections, we describe the components of a Bayesian data analysis algorithm which classifies detected GW signals from core-collapse supernovae as belonging to one of a set of signal catalogs, representing, e.g., different explosion mechanisms. A block diagram of the analysis algorithm, which we call the Supernova Model Evidence Extractor (SMEE), is shown in Fig. 2. SMEE is implemented in MATLAB<sup>1</sup>.

In a first step, SMEE performs Principal Component Analysis (PCA) via singular value decomposition (SVD) on the waveforms in each catalog to create sets of orthogonal basis vectors, the Principal Components (PCs), which are ordered according to their prevalence in their catalog. In other words, the first PC represents the most common feature of all signals in the catalog, the second PC represents the second most common feature, and so on. Using a complete set of PCs, each waveform can be reconstructed as a linear combination of PCs for the corresponding catalog, allowing each waveform to be simply parameterized by the PC coefficients in the linear combination. However, since PCs are expected to span the parameter space defined by each catalog of waveforms efficiently, catalog waveforms may be reconstructed with good accuracy already with a set of PCs that is significantly smaller than the number of waveforms in the catalog. Moreover, non-catalog waveforms (i.e., real signals) may be identified as belonging to the same class of signals as catalog waveforms if they can be approximately matched with the first few PCs of a catalog.

SMEE then uses Bayesian model selection and computes the logarithm of the Bayes factor to distinguish between GW signal classes. The Bayes factor is the ratio of




FIG. 2: Block diagram of the Supernova Model Evidence Extractor (SMEE). A desired core-collapse supernova gravitational wave signal is injected into noise, and the algorithm compares it to the principal components (PCs) of a given waveform catalog representing a particular model. The PCs are constructed via singular value decomposition (SVD). The sign of the log Bayes factor between two PC sets indicates which model is favored by the data.

the evidences for two competing hypotheses and, for the purpose of our analysis, we weigh the evidence that the observed data supports the presence of a GW signal consistent with signals from one of any two competing catalogs. This requires summing up the likelihood function times the prior across all possible signal parameters (in our case, values of PC coefficients) to determine the evidence (also called the marginal likelihood) for two different signal models to be tested. SMEE accomplishes this efficiently via the Nested Sampling algorithm [40, 45].

#### B. Bayesian Model Selection

In our analysis, we employ Bayesian Model Selection, similar to that described in [111]. Specifically, we use the

<sup>1</sup> The MathWorks Inc., Natick, MA 01760, USA. <http://www.mathworks.com/products/matlab/>.

Bayes factor to compare the probabilities of two competing models. In general terms, the Bayes factor  $B_{ij}$  can be written as the ratio of the evidences  $p(D|M)$ ,

$$B_{ij} = \frac{p(D|M_i)}{p(D|M_j)}, \quad (1)$$

where  $M_i$  and  $M_j$  are two competing models tested using the data  $D$ . The evidence for each model is obtained by integrating the product of its likelihood function,  $p(D|\theta, M)$ , and prior,  $p(\theta|M)$ , across all model parameter values  $\theta$ , such that,

$$p(D|M) = \int_{\theta} p(\theta|M)p(D|\theta, M) d\theta. \quad (2)$$

This effectively calculates the likelihood averaged over all parameter values. The distribution of the prior reflects our bias for or against particular parameter values. For the analysis discussed in this article, we adopt a flat, uniform prior over all parameter values which means that we do not favour any particular portion of the parameter space.

The evidence will be greater for a model that is supported by the data. Therefore, the Bayes factor indicates which of the two competing models is preferred by the data. It is often more convenient to compare models by using the natural logarithm of the Bayes factor,

$$\log B_{ij} = \log p(D|M_i) - \log p(D|M_j). \quad (3)$$

In this case,  $\log B_{ij} > 0$  means  $M_i$  is the preferred model whereas  $\log B_{ij} < 0$  will point to  $M_j$  being favored.

### C. Nested Sampling Algorithm

For evaluating  $\log B_{ij}$ , we first need to calculate the evidences  $p(D|M_i)$  and  $p(D|M_j)$  for the two models  $M_i$  and  $M_j$ . From Eq. 2 we see that the evidence is the sum of the likelihood times the prior determined for all possible parameter values of the desired model. An exhaustive, brute-force approach to computing the evidence by calculating the likelihood values for every choice of parameter values is computationally prohibitive. It is also an inefficient way of determining the evidence since the likelihood values will be most significant, and therefore contribute most to the evidence, for a small subset of parameter values which constructs a waveform that closely resembles the data. For most other combinations of the model's parameters, the likelihood will be insignificant and do not contribute to the evidence.

Therefore, we choose to follow the approach of Veitch *et al.* [112] and employ Nested Sampling [40, 45] to efficiently calculate the evidence integral. The Nested Sampling algorithm determines the evidence integral by calculating the likelihood for sample values of the desired

model's parameters. Initially, the model's parameter values are randomly selected. The algorithm then iterates over different sets of the model's parameter values, calculating the likelihood for each set of parameter values. With each iteration, the smallest likelihood value obtained is compared with that calculated for a randomly chosen set of new parameter values. Only parameter values which lead to a larger likelihood value than the smallest obtained in the previous iteration are kept. Therefore, as the algorithm stochastically samples the parameter space, it iteratively converges on the set of parameter values that produce the most significant likelihood values.

The nested sampling algorithm is similar to the Markov-Chain Monte Carlo approach (e.g., [40]) except that the primary output of the Nested Sampling algorithm is the evidence, which can be immediately obtained by summation whereas the posterior distribution is only found as a by-product. To find the evidence, a set of "live points" are found through creating a stochastic sampling of the prior distribution to generate a set of  $N$  samples which are denoted as  $\theta_i$ , where  $i = 1 \dots N$ . In context of the analysis presented in later in this article (see section III E),  $\theta_i$  corresponds to  $\beta_i$  which is the  $i$ -th sample of the principal component coefficient for which we calculate the signal likelihood. The evidence integral (Eq. 2) is then written as

$$\begin{aligned} p(D|M) &= \int_{\theta} p(\theta|M)p(D|\theta, M) d\theta, \\ &\approx \sum_{i=1}^N p(D|\theta_i, M)w_i, \\ &\approx \sum_{i=1}^N L_iw_i, \end{aligned} \quad (4)$$

where the weight,

$$w_i = p(\theta_i|M)d\theta_i, \quad (5)$$

is the fraction of the prior distribution represented by the  $i$ -th sample and  $L_i$  is its likelihood. It is this weighted likelihood that is calculated by the Nested Sampling algorithm and subsequently used to obtain the evidence. More details on the Nested Sampling algorithm can be found in [45, 112].

### D. Principal Component Analysis via Singular Value Decomposition

Each core-collapse supernova waveform catalog consists of a number of GW signals obtained for different initial conditions and simulation parameters (e.g., progenitor star mass, EOS, rotational configuration etc.). While individual waveforms of one catalog are different in detail, they generally exhibit strong common general

features. This can be exploited by principal component analysis (PCA) [37], which isolates the most common features of waveforms in linearly independent principal components (PCs) ordered by their relevance. The first few PCs may already be sufficient to efficiently span their entire catalog, as was shown in [36] and [39] for the Dim catalog (see Sect. II B 1).

The PCs are obtained via singular value decomposition (SVD) (e.g., [37]) of time-domain waveforms from each catalog. To perform SVD on a catalog with  $m$  waveforms, a matrix  $\mathbf{A}$  is created such that each of its columns corresponds to a waveform of uniform length  $n$  from the catalog.

The  $n \times m$  waveform matrix  $\mathbf{A}$  is factored so that

$$\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T, \quad (6)$$

where  $\mathbf{U}$  is an  $n \times n$  matrix whose columns correspond to the eigenvectors of  $\mathbf{A}\mathbf{A}^T$ . Similarly, the columns of the  $m \times m$  matrix  $\mathbf{V}$  correspond to the eigenvectors of  $\mathbf{A}^T\mathbf{A}$  and  $\mathbf{\Sigma}$  is an  $n \times m$  diagonal matrix whose elements correspond to the square root of the corresponding eigenvalues.

Since  $\mathbf{A}\mathbf{A}^T$  is the covariance matrix of  $\mathbf{A}$ , the eigenvectors in  $\mathbf{U}$  are effectively an orthonormal basis which span the  $m$ -dimensional parameter space defined by the catalog of waveforms used to construct  $\mathbf{A}$ . Note that, in practice,  $n \gg m$  and it is impractical to determine the eigenvectors in  $\mathbf{U}$  directly. Instead, the smaller  $\mathbf{V}$  and its corresponding eigenvalues in  $\mathbf{\Sigma}$  are first determined which are subsequently used to derive  $\mathbf{U}$ .

The orthonormal eigenvectors of  $\mathbf{U}$  are the PCs and are ranked by their corresponding eigenvalues. The PC with the largest corresponding eigenvalue is referred to as the first PC and consists of the most significant common features of all waveforms in the catalog. It follows that the PC with the second largest corresponding eigenvalue is the second PC and consists of the second most significant common features and so on.

The waveforms in  $\mathbf{A}$  can be reconstructed by taking a linear combination of PCs,

$$h_i \approx \sum_{j=1}^k U_j \beta_j, \quad (7)$$

where  $h_i$  is the desired waveform from the catalog,  $U_j$  is the  $j$ th PC from the  $\mathbf{U}$  matrix and  $\beta_j$  is the corresponding PC coefficient, which can be obtained by projection of  $h_i$  onto  $U_j$ . The sum of  $k$  PCs produces an approximation of the desired waveform since  $k \leq m$ .

## E. Signal and Noise Models

For the analysis described here, two types of models are considered. The signal model  $M_s$  tests the presence

of a signal waveform  $h(\beta)$  in the data. Here, PCA is performed for each catalog (using SVD; see section III D) and each waveform is parameterized by its PC coefficients ( $\beta$ ). The Gaussian likelihood function for the signal model is

$$p(D|\beta, M_s) = \prod_{i=1}^N \frac{1}{\sigma_i \sqrt{2\pi}} \exp \left[ -\frac{(D_i - h_i(\beta))^2}{2\sigma_i^2} \right], \quad (8)$$

where  $\sigma_i$  is the standard deviation of the noise,  $h_i(\beta)$  is the desired waveform reconstructed from the PCs and  $N$  is the length of the data with a corresponding index  $i$ . The evidence for the signal model is determined by performing the integral in Eq. 2 numerically, using uniformly-distributed priors over a chosen range for  $\beta$ .

On the other hand, the noise model  $M_n$  tests the data's consistency with Gaussian noise. The likelihood function for the noise model is the same as that in Eq. 8, but with  $h(\beta) = 0$ . From this, it is straightforward to perform the integration in Eq. 2 and obtain an analytic form for the noise evidence function,

$$p(D|M_n) = \prod_{i=1}^N \frac{1}{\sigma_i \sqrt{2\pi}} \exp \left[ -\frac{D_i^2}{2\sigma_i^2} \right]. \quad (9)$$

In both Eq. 8 and 9, the standard deviation of the noise is a function of each sample in the data, because the simulated noise is designed to correspond to the expected sensitivity of Advanced LIGO, which varies as a function of frequency (see Sec. III F). To handle the frequency-dependent noise, the signal and noise evidences are calculated in the frequency domain with  $D_i$ ,  $h_i(\beta)$  and  $\sigma_i$  corresponding to data, reconstructed waveform, and noise in the  $i$ -th frequency bin respectively. In particular, each reconstructed waveform is obtained by taking a linear combination of the Fourier transforms of its corresponding PCs.

The natural logarithm of the Bayes factor used to compare the signal model to the noise model is then simply

$$\log B_{SN} = \log[p(D|M_s)] - \log[p(D|M_n)]. \quad (10)$$

## F. Generation of Simulated Noise

We generate Gaussian colored noise, assuming a single Advanced LIGO detector in the proposed broadband configuration (the so-called ‘‘zero detuning, high-power’’ mode). We employ the data file `ZERO_DET_high.P.txt` provided by [113], which contains  $\sqrt{S(f)}$ , the square root of the one-sided detector noise power spectral density in units of  $(\text{Hz})^{-1/2}$ . An open-source implementation for MATLAB of what we describe in the following can be found in [114].

The real discrete time-domain noise  $n(t_j)$ , where  $t_j$  denotes the  $j$ -th discrete time interval of size  $\Delta t$ , is obtained




FIG. 3: The first three Principal Components (PCs) of the waveforms from the Dimmelmeier (**Dim**) [17, 100], Murphy (**Mur**) [71, 84], and Ott [12, 84] catalogs, which we take to be representative of the magnetorotational, neutrino and acoustic mechanisms, respectively (see Secs. II A, II B and II C). In creating the PCs, the waveforms of each catalog are placed in a systematic way in a 3 s interval and padded left and right by zeros. The vertical axis is a dimensionless scale which represents the amplitude, which we have normalized by the maximum amplitude over all 3 PCs of each catalog shown here.

by inverse discrete Fourier transform from the complex frequency-domain noise  $\tilde{n}(f_k)$ , where  $f_k$  denotes the  $k$ -th discrete frequency interval of size  $\Delta f = 1/(N_t \Delta t)$ , where  $N_t$  is the number of intervals in the time domain. Since the time-domain noise is real and has zero mean, the frequency-domain noise must obey

$$\begin{aligned} \tilde{n}(-f) &= \tilde{n}^*(f), \\ \tilde{n}(f=0) &= \tilde{n}(f=f_{\text{Nyq}}) = \tilde{n}(f=-f_{\text{Nyq}}) = 0. \end{aligned} \quad (11)$$

Here,  $f_{\text{Nyq}} = 1/(2\Delta t)$  is the Nyquist frequency. If  $N_t$  is the even number of equally spaced bins in time of width  $\Delta t$ , then  $N_f = N_t/2 - 1$  is the number of independent frequency bins  $f_k$  in the frequency domain of width  $\Delta f$ . The frequency variable  $f_k$  assumes values from  $-f_{\text{Nyq}}$  to  $f_{\text{Nyq}}$ .

$|\tilde{n}(f_k)| = \sqrt{\tilde{n}(f_k)\tilde{n}^*(f_k)}$  is a two-sided amplitude spectral density. We generate  $\tilde{n}(f_k)$  by sampling the standard

normal distribution (zero mean, variance one) weighted by the noise transfer function  $T(f_k) = \sqrt{S(f_k)}/\sqrt{2}$ . The real and imaginary parts of  $\tilde{n}(f_k)$  are then given for each  $f_k \in (0, f_{\text{Nyq}})$  by

$$\Re(\tilde{n}(f_k)) = \frac{1}{\sqrt{2}} T(f) \text{RANDN}, \quad (13)$$

$$\Im(\tilde{n}(f_k)) = \frac{1}{\sqrt{2}} T(f) \text{RANDN}, \quad (14)$$

where **RANDN** is a random number sampled from the standard normal distribution. The remaining  $\tilde{n}(f_k)$  are then obtained via Eqs. 11 and 12.

The inverse discrete Fourier transform of  $\tilde{n}(f_k)$  to time domain noise  $n(t_j)$  will preserve its Gaussian character in the time domain in the limit of small sampling interval [39, 112]. Specifically, we use the following definition of the discrete Fourier transform (DFT) for transform-

ing noise from the frequency to the time domain when needed:

$$\tilde{n}(f_k) = \sum_{j=0}^{N-1} n(t_j) \exp(-2\pi i j k / N), \quad (15)$$

$$n(t_j) = \frac{1}{N} \sum_{k=0}^{N-1} \tilde{n}(f_k) \exp(2\pi i j k / N). \quad (16)$$

For convenience, we define the matched filter signal-to-noise ratio (SNR) of a GW signal  $h$  as

$$\text{SNR}^2 = 4 \int_0^\infty \frac{|\tilde{h}(f)|^2}{S(f)} df \quad (17)$$

$$= 4 \Delta t^2 \Delta f \sum_{k=1}^{N_f} \frac{|\tilde{h}(f_k)|^2}{S(f_k)}, \quad (18)$$

where  $S(f)$  is the one-sided noise power spectral density. The factor  $\Delta t^2$  is applied to correct the dimensions of  $\tilde{h}(f)$ , which we obtain via the DFT defined by Eq. 15.

## G. Application in SMEE

### 1. GW Signal Preparation and PCA

Before carrying out PCA and injecting signals into noise, all waveforms are buffered with zeros to be of length  $n$ , which we choose to correspond to 3 s at a sampling rate of 4096 Hz, allowing us to comfortably accommodate the longest available core-collapse supernova GW signals. The Advanced LIGO sampling rate is 16 kHz. The reduced sampling rate we choose saves computation time and is sufficient to capture the frequency content of the core-collapse supernova waveforms considered here, which have most of their power at  $\sim 50 - 1000$  Hz.

We align waveforms from the `Dim` catalog at their maximum (the spike at core bounce). Waveforms from the `Mur` and `Ott` catalogs are aligned so that the onsets of emission coincide. All waveforms are shifted so that they are aligned to the 4000-th point in the SMEE input data file, corresponding to about the 1 s mark in the 3 s interval, to leave ample space left and right of the waveform.

In Fig. 3, we present the first three PCs computed for the `Dim` (magnetorotational mechanism; left panel), `Mur` (neutrino mechanism; center panel), and `Ott` (acoustic mechanism; right panel) catalogs. Before generating PCs for the `Mur` catalog we filter out the secular low-frequency drifts present in the `Mur` waveforms (see Fig. 1) by high-passing the signal above 30 Hz. Since the low-frequency components are hidden in detector noise even when the source is nearby, dropping them improves the efficiency of our subsequent Bayesian analysis and signal reconstruction. We apply the same high-passing to trial waveforms for the neutrino mechanism before injecting them into noise.

### 2. Signal Injection and Model Selection

We inject trial GW signals into simulated Gaussian Advanced LIGO noise and use SMEE to determine which signal model (e.g., what core-collapse supernova explosion mechanism) a given injected signal belongs to via the evaluation of the logarithmic Bayes factors  $\log B_{SN}$  (Eq. 10) for an injected signal for each signal model  $S$  and the noise model  $N$ . Comparing two signal models  $i$  and  $j$  is then accomplished by computing  $\log B_{ij} = \log B_{iN} - \log B_{jN}$ .

SMEE's model selection operates in the frequency domain. Trial GW signal and the PCs belonging to the signal model under consideration are transformed into the frequency domain via DFT and the trial GW signal is added to the complex frequency-domain noise, retaining phase information. The Nested Sampling algorithm is then invoked to marginalize the PC coefficients  $\beta_k$ . The prior for each coefficient is flat and uniform. The prior range for each  $\beta_k$  is determined by first reprojecting all waveforms of a given catalog back onto the PCs to compute  $\hat{\beta}_{kl}$  for each PC  $k$  and waveform  $l$  of the catalog. The range of expected possible values of  $\beta_k$  is then found by taking the minimum and maximum of  $\hat{\beta}_{kl}$  over all  $l$  and adjusting these numbers by 10% down and up to add a margin of error to account for uncertainty due to the noise, motivated by the findings of [39].

Keeping the noise model fixed, the results of SMEE's computations will depend on the SNR of the signal, i.e. the distance to the core collapse event, and on the amount of information we can provide to SMEE about expected signals in the form of PCs.

The maximum number of PCs at SMEE's disposal is limited by the number of waveforms used to determine the set of PCs. While each catalog used in this study has a different number of waveforms, we choose to simplify our analysis by using the same number of PCs for all catalogs. Hence, the maximum number of PCs we use here is 7 and is set by the number of waveform in the `Ott` catalog (see §II C 1). Using 7 PCs gives SMEE complete information about signals belonging to the `Ott` catalog and significant, but incomplete information about waveforms from the `Dim`, `Mur`, and `Abd` catalogs. We also carry out SMEE runs with less than 7 PCs to study the dependence on the number of PCs employed. Using only a small subset of a catalog's PCs limits SMEE's ability to precisely reconstruct injected catalog waveforms, but it represents the real-life situation that the a priori information about a detected signal is severely limited. Our goal here is not to ideally reconstruct signals but to show that determining the underlying physical model of an observed signal is possible with limited advance knowledge.

## IV. RESULTS

### A. Response to Gaussian Noise

For interpreting the results of SMEE’s Bayesian model selection on the basis of Eq. 3, it is necessary to quantify and understand SMEE’s response to pure Gaussian detector noise without a signal being present. To this end, we run SMEE on 10,000 randomized instances of Advanced LIGO detector noise (generated as described in §III F) without injecting signals and compute  $\log B_{SN}$  (Eq. 10) in the absence of a signal for each signal model  $S$ . The results, shown in Fig. 4, follow a Gaussian distribution with a mean corresponding to the expected value  $-\sum_{i=1}^N \frac{h_i(\beta)^2}{2\sigma_i^2}$ , where  $N$  is the number of PCs employed (see §III E). The average logarithmic Bayes factors obtained for 10,000 instances of noise indicate that noise, or any signal fully consistent with noise, is most likely to have a logarithmic Bayes factor of  $-54.0$  when SMEE is run with 7 PCs of the `Dim` catalog. For the `Ott`, `Mur`, and `Abd` catalogs, the expected logarithmic Bayes factors for pure Gaussian noise and 7 PCs are  $-52.1$ ,  $-52.3$ , and  $-53.0$ , respectively. The observed expectation values are very comparable to those calculated for the `Dim` ( $-53.9$ ), `Ott` ( $-52.2$ ), `Mur` ( $-52.3$ ), and `Abd` ( $-52.9$ ) catalogs, respectively, verifying that SMEE is operating as expected. We have repeated this experiment for the case when only 3 PCs are used and also in this case find that SMEE closely reproduces the predicted expectation values, which are near  $-26$  in the 3-PC case.

Since the logarithmic Bayes factors follow a Gaussian distribution, we can set a threshold using the standard deviations as an indicator for the expected false alarm rate. Ideally, for the `Dim` catalog, a 1% false alarm rate would correspond to a threshold that is  $\sim 2.6$  times the standard deviation, corresponding to  $\sim 0.44$  above the mean. However, we note that the expected logarithmic Bayes factor value varies between different catalogs and, for a fixed false alarm rate, we would require a different threshold for each catalog. This variation can be addressed by re-normalising all Bayes factors so that they are the same for all catalogs when there is only noise. But, since the focus of our work here is to distinguish between different waveforms and not to perform a study on the detection efficiency of GW signals, we choose to take the more conservative approach of simply setting a higher threshold. Therefore, we conservatively choose to identify a signal as being distinct from noise if its  $\log B_{SN}$  is greater than  $-47$  (in cases in which we use 3 PCs, this number is  $-21$ ). When comparing two signal models  $M_i$  and  $M_j$ , we conservatively identify model  $M_i$  as favored if  $\log B_{ij} \geq 5$  (and vice versa).




FIG. 4: Results from running SMEE with 7 PCs of the `Dim` catalog and without an injected SMEE signal on 10,000 randomized instances of Gaussian Advanced LIGO noise, generated as described in Sec. III F. A signal consistent with noise is most likely to have a logarithmic Bayes factor of  $\sim -54.0$ . The red line plots a Gaussian distribution with a mean of  $-53.96$  and a standard deviation  $\sigma = 0.17$ .

TABLE I: Minimum SNR at which the injected waveform will give a  $\log B_{SN}$  above the threshold defined in §IV A. This is shown for 5 representative waveforms from the `Dim`, `Mur`, and `Ott` catalogs, which we take to be representative of the magnetorotational, neutrino, and acoustic mechanisms, respectively. We provide results from SMEE runs with 3 and 7 PCs.

Waveform Name	Minimum SNR	
	3 PCs	7 PCs
<b>Dim</b> [17]		
s11a1o01_Shen	13	13
s11a3o09_Shen	4	4
s20a3o05_LS	5	5
s40a3o07_LS	4	4
s40a3o13_Shen	4	4
<b>Mur</b> [71]		
20_3.8	9	4
40_10.0	4	4
40_13.0	21	15
15_3.2	10	9
15_4.0	4	4
<b>Ott</b> [12]		
nomoto13	143	4
nomoto15	45	4
s15.0WHW02	4	4
s20.0WHW02	4	4
s25.0WHW02	4	4

### B. Signal vs. Noise

The minimal GW signal strength required for SMEE to be able to select the core-collapse supernova mechanism is an important question. The primary prerequisite for an incident GW signal to be useful for model selection is that SMEE can distinguish it from detector noise, i.e.,

TABLE II:  $\log B_{SN}$  with respect to the **Dim**, **Mur**, and **0tt** PCs computed for representative injected waveforms from the **Dim**, **Mur**, and **0tt** catalogs, which we take to be representative of the magnetorotational, neutrino, and acoustic explosion mechanism, respectively. Results for source distances of 0.2, 2, and 10 kpc are given. Each table entry shows the results for 3 PCs (to the left of the vertical divider |) and the 7-PC result (to the right of the divider).  $\log B_{SN} < -47$  when 7 PCs are used and  $\log B_{SN} < -21$  when 3 PCs are used indicates that the injected signal is likely consistent with noise while larger values suggests that the signal belongs to the signal model whose PCs were used in the analysis.

Waveform Name	$\log B_{SN}$ Dim PCs			$\log B_{SN}$ Mur PCs			$\log B_{SN}$ 0tt PCs		
	0.2 kpc	2 kpc	10 kpc	0.2 kpc	2 kpc	10 kpc	0.2 kpc	2 kpc	10 kpc
<b>Dim</b> [17]									
s11a1o01_Shen	38982 48686	364 430	-10 -34	9178 25647	65 203	-22 -42	-25 222	-27 -51	-26 -52
s11a3o09_Shen	5 $\times$ 10 <sup>6</sup>  5 $\times$ 10 <sup>6</sup>	52403 54505	2071 2129	36170 58531	335 532	-11 -29	23 9191	-26 39	-26 -49
s20a3o05_LS	3 $\times$ 10 <sup>6</sup>  3 $\times$ 10 <sup>6</sup>	25738 26859	1005 1023	4139 29285	15 240	-24 -40	1 2242	-26 -31	-26 -51
s40a3o07_LS	1 $\times$ 10 <sup>7</sup>  1 $\times$ 10 <sup>7</sup>	1 $\times$ 10 <sup>5</sup>  1 $\times$ 10 <sup>5</sup>	4737 5411	47029 1 $\times$ 10 <sup>5</sup>	444 951	-7 -12	519 56978	-21 517	-26 -29
s40a3o13_Shen	2 $\times$ 10 <sup>7</sup>  2 $\times$ 10 <sup>8</sup>	2 $\times$ 10 <sup>5</sup>  2 $\times$ 10 <sup>5</sup>	8174 8302	2 $\times$ 10 <sup>5</sup>  4 $\times$ 10 <sup>5</sup>	1830 3968	48 109	160 19781	-25 145	-26 -44
<b>Mur</b> [71]									
20_3.8	4205 10631	16 50	-24 -50	59264 4 $\times$ 10 <sup>5</sup>	566 3775	-2 101	-22 -5	-27 -53	-26 -52
40_10.0	280 2362	-23 -33	-26 -53	4 $\times$ 10 <sup>5</sup>  5 $\times$ 10 <sup>5</sup>	4198 4981	143 149	-26 -29	-27 -54	-26 -52
40_13.0	365 819	-23 -48	-26 -54	3874 7591	12 23	-24 -49	-27 -42	-27 -54	-26 -52
15_3.2	7450 10699	48 51	-23 -50	23744 29439	211 241	-16 -41	-25 -12	-27 -53	-26 -52
15_4.0	2030 7658	-6 20	-25 -51	8 $\times$ 10 <sup>5</sup>  8 $\times$ 10 <sup>5</sup>	7672 7812	282 262	-12 -22	-26 -53	-26 -52
<b>0tt</b> [12]									
nomoto13	3731 6155	11 4	-24 -51	132 313	-25 -50	-26 -52	544 1 $\times$ 10 <sup>6</sup>	-21 11893	-26 426
nomoto15	1520 2428	-11 -32	-25 -53	54 146	-26 -52	-26 -52	32145 7 $\times$ 10 <sup>6</sup>	295 65635	-13 2575
s15.0WHW02	578 2417	-20 -33	-26 -53	1931 2601	-7 -27	-25 -51	6 $\times$ 10 <sup>7</sup>  6 $\times$ 10 <sup>7</sup>	6 $\times$ 10 <sup>5</sup>  6 $\times$ 10 <sup>5</sup>	24653 24655
s20.0WHW02	622 2895	-20 -28	-26 -53	-19 587	-27 -47	-26 -52	4 $\times$ 10 <sup>7</sup>  4 $\times$ 10 <sup>7</sup>	4 $\times$ 10 <sup>5</sup>  4 $\times$ 10 <sup>5</sup>	17975 17957
s25.0WHW02	2343 5963	-3 3	-25 -51	1918 2969	-7 -24	-25 -51	1 $\times$ 10 <sup>8</sup>  1 $\times$ 10 <sup>8</sup>	1 $\times$ 10 <sup>6</sup>  1 $\times$ 10 <sup>6</sup>	45853 45829

we must find the minimum signal strength (i.e., SNR) so that  $\log B_{SN} > -47$  (when 7 PCs are used; Eq. 10 and §IV A).

In order to determine the range of minimum SNR required across and within core-collapse supernova GW signal types, we draw 5 representative waveforms from the **Dim**, **Mur**, and **0tt** catalogs and run them through SMEE at varying SNR, using 7 PCs generated from the catalog to which each injected waveform belongs. We choose the waveforms in such a way to capture the full range of variation within each catalog. The result of this exercise is shown in Fig. 5 and summarized in Tab. I. Generally, an SNR  $\gtrsim 4 - 5$  is required for SMEE to find  $\log B_{SN} > -47$  in the idealized setting that we consider here. In a real fully blind search, unknown arrival times and non-Gaussianity of real detector noise will generally require an SNR in excess of 8 for a detection statement (e.g., [115]).

In Fig. 5, the waveforms associated with the acoustic mechanism (**0tt** catalog) require the smallest SNR, followed by those of the neutrino mechanism (**Mur** catalog) and the magnetorotational mechanism (**Dim** catalog). This hierarchy in minimum SNR, however, is not fundamental, but a consequence of the fact that we have chosen to carry out this test using 7 PCs for each waveform catalog. Since the **0tt** catalog comprises only 7 waveforms, the set of 7 PCs completely spans it and allows perfect reconstruction, maximizing  $p(D|M_s)$  (Eq. 8). In the case

of less than perfect knowledge of the signal, the minimum SNR will always be greater. This is why the **Dim** and **Mur** catalogs, which have many more than 7 waveforms, require larger minimum SNR than the **0tt** waveforms.

The situation is somewhat different, when we recalculate  $\log B_{SN}$  using only 3 PCs. The **0tt** catalog is very small and rather diverse in the time domain. Its first few PCs do not efficiently span the catalog and, when only the first 3 PCs are used, the minimum SNRs for waveforms poorly reconstructed with these PCs increases dramatically, as shown for the **0tt** waveforms *nomoto13* and *nomoto15* in Tab. I. Some of the **Mur** waveforms also exhibit increased minimum SNRs, indicating that there is significant time-domain variation that is not captured by the first 3 PCs. The large **Dim** catalog, on the other hand, is very efficiently spanned already by its first few PCs, as previously pointed out by [36, 39], and the minimum SNR for waveforms from this catalog remains practically unchanged when going from 7 to 3 PCs in our analysis.

### C. Determining the Core-Collapse Supernova Explosion Mechanism

The basic assumption of this study is that the neutrino, magnetorotational, and acoustic core-collapse supernova explosion mechanism have robustly distinct GW signatures. In this section, we test this assumption by inject-




FIG. 5: Mean  $\log B_{SN}$  as a function of signal-to-noise ratio (SNR; Eq. 18) for 5 representative waveforms from the **Mur**, **0tt** and **Dim** catalogs that encapsulate the variations within each catalog using 7 Principal Components (PC). The shaded areas represent the standard error in the mean value of  $\log B_{SN}$  for each waveform catalog computed as  $\sigma = \pm N^{-1}(\sum_i(\bar{x} - x_i)^2)^{1/2}$ , where  $\bar{x}$  is the mean and  $x_i$  are the individual SNRs and  $N$  is the number of waveforms. Values of  $\log B_{SN}$  below  $-47$  in the 7-PC case and below  $-21$  in the 3-PC case indicate that the algorithm considers it more likely that there is no signal detectable in the noise. Table I summarizes numerical results for the minimum SNR for which  $\log B_{SN} \geq -47$ .

ing waveforms into simulated noise and running SMEE on the data using PCs of waveform catalogs representative of the neutrino, magnetorotational, and acoustic mechanisms. If our assumption is correct and the GW signatures of these mechanisms are truly distinct, then SMEE should (i) yield the largest value of  $\log B_{SN}$  when the set of PCs is used that corresponds to the mechanism the waveform is representative of, and, (ii)  $\log B_{ij}$  (Eq. 3) should be positive (and larger than  $\sim 5$ ; see §IV A) if the injected waveform is most consistent with mechanism  $i$ , negative if it is most consistent with mechanism  $j$ , and near zero if the result is inconclusive.

We carry out our SMEE calculations for events located at 0.2 kpc, 2 kpc, and 10 kpc and with 3 and 7 PCs. Betelgeuse ( $\alpha$  Orionis), a red supergiant star of  $15 - 20 M_{\odot}$  [116], is located  $197 \pm 45$  pc from Earth [117] and will eventually explode as a Type II-P core-collapse supernova. Hence, studying a potential event at 0.2 kpc will tell us what we can learn from the observation of GWs from Betelgeuse’s collapse and supernova. 2 kpc is still nearby on the galactic scale, but the Galactic volume out to this radius already contains hundreds of supergiants,

one of which may make the next galactic supernova [118]. Finally, 10 kpc is the fiducial Galactic distance scale and we consider it to state what could be inferred throughout the Milky Way. As in Sec. IV B, we carry out SMEE runs with 3 and 7 PCs to study the sensitivity of the results on the amount of knowledge about the injected waveform we grant SMEE.

Table II summarizes the results from  $\log B_{SN}$  calculations for five representative waveforms from the **Mur** (neutrino mechanism), **Dim** (magnetorotational mechanism), and **0tt** (acoustic mechanism) catalogs. The larger the value of  $\log B_{SN}$ , the greater the confidence that the data contain a signal consistent with the employed set of PCs.  $\log B_{SN} \leq -47$  (for the 7-PC case;  $-21$  in the 3-PC case) indicates that the signal is more consistent with noise. The results show that it is indeed possible to clearly associate any injected waveform with its catalog and, thus, select the explosion mechanism for a core-collapse supernova out to at least 2 kpc. However, at the galactic scale (10 kpc), a significant fraction of waveforms representative of the neutrino mechanism, due to their intrinsically low GW amplitudes, can be told apart from noise only marginally or are even most consistent with noise. Typical magnetorotational explosions and explosions driven by the acoustic mechanism are still clearly identifiable at 10 kpc. However, when the reduced set of 3 PCs is used, the **0tt** catalog, which is not well spanned by only 3 PCs, suffers most and two out of the five representative **0tt** waveforms are not or only marginally identifiable with the 3 first **0tt** PCs at 10 kpc.

Provided that there is confidence that a signal has been detected, we can compute  $\log B_{ij}$  (Eq. 3) to study if the signal is more likely to be consistent with mechanism  $i$  or mechanism  $j$ . Since we know  $\log B_{SN}$  for all signal models, we can simply compute  $\log B_{ij} = \log B_{iN} - \log B_{jN}$  from the data for representative waveforms provided in Tab. II. We use 7 PCs for these calculations.

In Fig. 6, we show results of injection studies of *all* waveforms from the **Dim**, **Mur**, and **0tt** catalogs run through SMEE and analyzed with the **Dim**, **Mur**, and **0tt** PCs at a source distance of 10 kpc. The full numerical results on whose basis Fig. 6 was generated are available online [119]. The left panel depicts the  $\log B_{\text{DimMur}}$  result for injected waveforms from the **Dim** and **Mur** catalogs, that we take to be representative of the magnetorotational and neutrino mechanism, respectively. Even at 10 kpc the vast majority of waveforms characteristic for magnetorotational explosions are clearly identified as belonging to this mechanism. For the neutrino mechanism, the evidence is generally significantly weaker and only  $\sim 44\%$  of the **Mur** waveforms are identified with  $\log B_{\text{DimMur}} < -100$  and none have  $\log B_{\text{DimMur}} < -1000$ , while  $\sim 19\%$  are in the inconclusive regime of  $-5 < \log B_{\text{DimMur}} < 5$ .

In the center panel of Fig. 6, we show results for  $\log B_{\text{Dim0tt}}$  for injected waveforms corresponding to the magnetorotational (**Dim**) and the acoustic (**0tt**) mechanism. The case is clear cut and most waveforms are correctly identified as most likely belonging to their re-

spective catalog/mechanism. Finally, the right panel of Fig. 6 presents  $\log B_{\text{MurOtt}}$  for waveforms representative of the neutrino (**Mur**) and acoustic (**Ott**) mechanism. As in the previous panel, SMEE associates the waveforms corresponding to the acoustic mechanism with high confidence to the **Ott** catalog. The evidence suggesting correct association of the neutrino mechanism waveforms is considerably less strong, but  $\log B_{\text{MurOtt}}$  is still conclusive for  $\sim 88\%$  of the **Mur** waveforms.

Figure 7 shows the results for  $\log B_{\text{DimMur}}$ ,  $\log B_{\text{DimOtt}}$ , and  $\log B_{\text{MurOtt}}$  obtained by SMEE with 7 PCs at a source distance of 2 kpc. Here, all acoustic mechanism waveforms (**Ott** catalog), all magnetorotational mechanism waveforms (**Dim** call), and all neutrino mechanism waveforms (**Mur** catalog) are correctly identified as belonging to their respective catalog and explosion mechanism.

#### D. Deciding Between Rotating Accretion-Induced Collapse and Rotating Iron Core Collapse

The waveforms of the **Dim** catalog are representative of the GW signal emitted by rotating collapse and bounce of iron cores of massive stars with ZAMS masses  $\gtrsim 8 - 10 M_{\odot}$ . In the accretion-induced collapse (AIC) of rapidly rotating O-Ne white dwarfs, very similar dynamics occurs and the corresponding GW signals, as predicted by Abdikamalov *et al.* [46], share many of the basic features of the rotating iron core collapse and bounce waveforms of, e.g., the **Dim** catalog (see the discussion in Sec. IV.C. of [46]). Hence, it is interesting to see if our SMEE model selection algorithm can tell them apart.

We compute the PCs for the **Abd** catalog in the same fashion as done previously for the **Dim**, **Mur**, and **Ott** catalogs and inject all **Abd** and **Dim** waveforms into simulated Advanced LIGO noise. Since the **Abd** and **Dim** catalogs are very similar, the results found for **Dim** waveforms in Sec. IV B carry over directly to **Abd** waveforms. SMEE is then run with 7 PCs to calculate  $\log B_{\text{AbdDim}}$ . The result is shown in Fig. 8 for source distances of 10 kpc and 2 kpc. Full numerical results are available on-line [119].

In spite of the strong general similarity of rotating iron core collapse and rotating AIC waveforms, SMEE correctly identifies the vast majority of injected waveforms as most likely being emitted by a rotating iron core collapse or by rotating AIC. However, for a source at 10 kpc (left panel of Fig. 8),  $\sim 6\%$  of the **Dim** and  $\sim 5\%$  of the **Abd** are incorrectly identified as belonging to the respective other catalog. For an additional 2% of the **Dim** waveforms and 14% of the **Abd** waveforms, the evidence is inconclusive.

At a source distance of 2 kpc (right panel of Fig. 8), 88% of the AIC (**Abd**) and 93% of the rotating core collapse (**Dim**) waveforms are correctly identified.

If one placed trust in the reliability of less dominant and more particular features of waveforms in the underlying catalogs, one could use a larger number of PCs in the analysis. In order to study the effect of using an increased

number of PCs, we re-run the **Abd** vs. **Dim** comparison with 14 PCs and find that the result is significantly worse than with 7 PCs:  $\sim 61\%$  of the **Abd** waveforms and  $\sim 23\%$  of the **Dim** catalog are now incorrectly attributed to the respective other catalog at 10 kpc. This counter intuitive and at first surprising result is readily explained by the overall great similarity of the AIC and iron core collapse waveforms and the nature of PCA and SMEE's Bayesian model selection. The most robust features of each waveform catalog are encapsulated in its first few PCs. The first **Dim** and **Abd** PCs are indeed significantly different, but subsequent **Abd** and **Dim** PCs exhibit rather similar secondary features. Since each PC carries the same weight in SMEE's evidence calculation, including a larger number of PCs dilutes SMEE's judgment in this case and leads to the observed false identifications.

#### E. Testing Robustness with Unknown Waveforms

In the previous sections, we have demonstrated SMEE's ability to identify an injected trial GW signal as belonging to a particular physical model (i.e., emission mechanism and/or explosion mechanism). In this, however, we have drawn the injected waveforms directly from the catalogs used to generate the PCs. In other words, we have given SMEE (limited) advance knowledge about the injected waveforms.

Here, we carry out a much more stringent test of SMEE's ability to select between models of the core-collapse supernova mechanism by injecting waveforms that were not employed in the initial PC generation and/or stem from completely independent catalogs.

##### 1. Magnetorotational Mechanism

For the magnetorotational mechanism, we employ three additional **Dim** waveforms (**DimExtra**, Sec. II B 1) that were not included in the calculation of the **Dim** PCs. Furthermore, we inject waveforms from rotating models of the **Sch** catalog of Scheidegger *et al.* [102, 107] (see Sec. II B 1). The results of the  $\log B_{SN}$  calculation for the magnetorotational, neutrino, and acoustic mechanism signal models are summarized in Tab. III. **DimExtra** waveforms are identified as being most consistent with the **Dim** catalog and, hence, the magnetorotational mechanism. This is true with high confidence when 3 or 7 PCs are used and for all **DimExtra** signals out to distances  $\gtrsim 10$  kpc.

The **Sch** waveforms were generated with a completely different numerical code and thus allow for a truly independent test of our approach in SMEE. Also, unlike the **Dim** waveforms, the **Sch** waveforms are based on 3D simulations. Hence, they are not linearly polarized. For consistency with our current approach, we neglect  $h_{\times}$  and inject only  $h_{+}$  as seen by an equatorial observer. Results of SMEE  $\log B_{SN}$  calculations for all injected **Sch**




FIG. 6: Histograms describing the outcome of signal model comparisons by means of the Bayes factors  $\log B_{ij} = \log p(D|M_i) - \log p(D|M_j)$ , where  $i \neq j$  and  $M_i$  and  $M_j$  are signal models described by the **Dim** (magnetorotational mechanism), **Mur** (neutrino mechanism), and **Ott** (acoustic mechanism) waveform catalogs. The Bayes factors are computed with 7 PCs and for a source distance of 10 kpc. A positive value  $\log B_{ij}$  indicates that the injected waveform most likely belongs to model  $M_i$ , while a negative value suggest that model  $M_j$  is the more probable explanation. The bars are color-coded according to the type of injected waveform. The results are binned into ranges of varying size from  $< -10000$  to  $> 10000$  and the height of the bars indicates what fraction of the waveforms of a given catalog falls into a given bin of  $\log B_{ij}$ . We consider the range of  $(-5, 5)$  of  $\log B_{ij}$  as inconclusive evidence (see §IV A).




FIG. 7: Same as Fig. 6, but computed for a source distance of 2 kpc.

waveforms are summarized in Tab. III. SMEE correctly identifies *all* injected **Sch** waveforms as indicative of magnetorotational explosions at a source distance of 2 kpc. At 10 kpc, still 91% of the injected **Sch** waveforms are attributed to the magnetorotational mechanism, which is an indication of the robustness of the GW signal associated with rapid rotation and magnetorotational explosions. The very few **Sch** waveforms that SMEE is not able to clearly associate with the magnetorotational mechanism have such weak SNRs that they are more consistent with noise than with any of the catalogs at 10 kpc.

## 2. Acoustic Mechanism

We test SMEE’s ability to identify core-collapse supernovae exploding via the acoustic mechanism by injecting the three **OttExtra** waveforms (see Sec. II C 1). The re-

sults of this test are again summarized in Tab. III. They suggest that the a-priori unknown **OttExtra** waveforms can be identified as belonging to the acoustic mechanism out to 2 kpc with great confidence when 7 PCs are used in the analysis. At 10 kpc, the waveforms are still correctly attributed to the acoustic mechanism, but the evidence is much weaker in the 7-PC case while the waveforms are more consistent with noise when the analysis is performed with only 3 PCs. The **OttExtra** 3 waveform, which is clearly identified at 10 kpc, has an extreme SNR of  $\sim 2530$  at this distance, while the two other waveforms have SNRs of  $\sim 50$ . SMEE’s difficulty is illustrated in the right panel of Fig. 9, which indicates that the **OttExtra** waveforms reach the threshold of  $\log B_{SN} \geq -47$  only for SNRs  $\gtrsim 35$ , whereas **Ott** waveforms are identified already at SNRs  $\gtrsim 4$ , if the full set of 7 PCs is used. This is a strong indication that the range of possible waveform features associated with the acoustic mechanism is




FIG. 8: Outcome of the SMEE analysis of injected rotating iron core collapse (Dim catalog) and rotating accretion-induced collapse (AIC, Abd catalog) waveforms. The left panel shows results for a source distance of 10 kpc and the right panel depicts the results for a distance of 2 kpc. The Bayes factors  $\log B_{\text{AbdDim}}$  are computed on the basis of 7 PCs from the Abd and Dim catalog. A positive value of  $\log B_{\text{AbdDim}}$  indicates that an injected waveform is most likely associated with rotating AIC and a negative value suggests it to be more consistent with rotating iron core collapse. The results are binned into ranges of varying size from  $< -10000$  to  $> 10000$  and the height of the color-coded bars indicates what fraction of the waveforms of a given catalog falls into a given bin of  $\log B_{\text{AbdDim}}$ . We consider the range of  $(-5, 5)$  of  $\log B_{ij}$  as inconclusive evidence (see §IV A).




FIG. 9: Mean and  $1\text{-}\sigma$  range of  $\log B_{SN}$  as a function of signal-to-noise ratio SNR comparing signal with noise evidence. The horizontal lines mark the threshold values of  $\log B_{SN}$  above which we consider an injected waveform to be distinct from Gaussian noise. **Left panel:** Results for the Sch and DimExtra. These two were both reconstructed with 7 Dim PCs. **Right panel:** Results for the Yak, MurRem and OttExtra waveforms as reconstructed with 7 Mur for the first two and 7 Ott PCs for the latter. The Dim PCs very efficiently reconstruct the Sch and DimExtra waveforms for moderate SNRs while the Yak and OttExtra require very high SNRs to be distinguished from noise by the Mur and Ott PCs, respectively.

not efficiently covered by the 7 PCs generated from the Ott catalog. This could simply be attributed to the very small number of waveforms in this catalog. However, when studying the Ott and OttExtra waveforms, one immediately notes that the time between the first peak (associated with core bounce) and the second peak (the global maximum, associated with the non-linear phase

of the protoneutron star pulsations) varies significantly between waveforms. Since we compute PCs in the time domain, such large-scale features are imprinted onto the PCs and make it difficult to identify waveforms whose two peaks are separated by significantly different intervals. An alternative method that may work much better for waveforms of this kind is to compute PCs based on

TABLE III:  $\log B_{SN}$  for gravitational waveforms that were not included in the catalogs used for PC computation. The **DimExtra**, **Sch**, **OttExtra**, and **Yak** waveforms are discussed in §II. The **MurRem** waveforms are three randomly selected waveforms from the **Mur** catalog that were removed before re-computation of the **Mur** PCs. Results are shown for source distances of 0.2 kpc, 2 kpc, and 10 kpc and for evaluations using 3 PCs (to the left of the vertical divider |) and 7 PCs (to the right of the divider). Larger values indicate stronger evidence that the waveform is matched to the model catalog from which the PCs were constructed.  $\log B_{SN} < -47$  when 7 PCs are used and  $\log B_{SN} < -21$  when 3 PCs are used indicates that the injected signal is likely consistent with noise while larger values suggests that the signal belongs to the signal model whose PCs were used in the analysis.

Waveform Name	$\log B_{SN}$ Dim PCs						$\log B_{SN}$ Mur PCs						$\log B_{SN}$ Ott PCs					
	0.2 kpc		2 kpc		10 kpc		0.2 kpc		2 kpc		10 kpc		0.2 kpc		2 kpc		10 kpc	
<b>DimExtra</b> [39]																		
s20a1o05_shen	3×10 <sup>6</sup>	3×10 <sup>6</sup>	28269	31142	1106	1194	1×10 <sup>5</sup>	2×10 <sup>5</sup>	1429	2020	32	31	352	6515	-23	12	-26	-49
s15a1o03_LS	8×10 <sup>6</sup>	9×10 <sup>6</sup>	89606	95070	3560	3751	2×10 <sup>5</sup>	3×10 <sup>5</sup>	1966	2593	54	54	384	40334	-22	350	-26	-36
s40a1o10_LS	2×10 <sup>7</sup>	2×10 <sup>7</sup>	2×10 <sup>5</sup>	2×10 <sup>5</sup>	6832	7564	3×10 <sup>5</sup>	1×10 <sup>6</sup>	2624	14076	80	513	142	13089	-25	78	-26	-47
<b>Sch</b> [102]																		
R1E1CA	20861	32817	182	271	-17	-41	68	482	-26	-48	-26	-52	-27	68	-27	-52	-26	-52
R1E1CA.L	11631	13694	90	80	-21	-48	-25	-48	-27	-54	-26	-52	-27	19	-27	-53	-26	-52
R1E1DB	19285	25307	167	196	-18	-44	63	438	-26	-49	-26	-52	-26	89	-27	-52	-26	-52
R1E3CA	27629	46426	250	408	-15	-35	1	-6	-26	-53	-26	-52	-23	160	-27	-52	-26	-52
R1STCA	8304	10006	57	44	-23	-50	114	127	-25	-52	-26	-52	-27	-33	-27	-54	-26	-52
R2E1AC	4×10 <sup>5</sup>	4×10 <sup>5</sup>	3923	4273	132	120	1133	1602	-15	-37	-25	-52	-19	991	-26	-43	-26	-52
R2E3AC	4×10 <sup>5</sup>	4×10 <sup>5</sup>	3566	3902	118	104	1151	1567	-15	-38	-25	-52	-18	578	-26	-47	-26	-52
R2STAC	8×10 <sup>5</sup>	8×10 <sup>5</sup>	7504	7785	275	260	405	1301	-22	-40	-26	-52	-25	1463	-27	-39	-26	-52
R3E1AC	3×10 <sup>6</sup>	4×10 <sup>6</sup>	28195	37420	1103	1445	7418	13743	48	84	-23	-47	126	10015	-25	47	-26	-48
R3E1AC.L	2×10 <sup>6</sup>	3×10 <sup>6</sup>	19227	25641	744	974	10361	11138	77	58	-22	-48	103	5089	-25	-2	-26	-50
R3E1CA	2×10 <sup>6</sup>	3×10 <sup>6</sup>	20652	30427	801	1165	7590	13598	50	83	-23	-47	183	7814	-24	25	-26	-49
R3E1DB	2×10 <sup>6</sup>	3×10 <sup>6</sup>	20722	30537	804	1170	10102	18050	75	127	-22	-45	176	7438	-25	21	-26	-49
R3E2AC	2×10 <sup>6</sup>	3×10 <sup>6</sup>	24203	27271	943	1039	4575	12516	19	72	-24	-47	32	7135	-26	18	-26	-49
R3E3AC	3×10 <sup>6</sup>	4×10 <sup>6</sup>	33975	39403	1334	1524	5493	8915	29	36	-24	-49	107	13629	-25	83	-26	-47
R3STAC	5×10 <sup>6</sup>	5×10 <sup>6</sup>	47277	50486	1866	1968	13231	17583	106	122	-21	-45	70	10361	-26	50	-26	-48
R4E1AC	9×10 <sup>6</sup>	1×10 <sup>7</sup>	87917	1×10 <sup>5</sup>	3492	4121	15672	26725	130	214	-20	-42	584	39541	-20	342	-26	-36
R4E1CF	5×10 <sup>7</sup>	6×10 <sup>7</sup>	5×10 <sup>5</sup>	6×10 <sup>5</sup>	21092	22361	8×10 <sup>5</sup>	3×10 <sup>6</sup>	7543	31753	277	1220	655	1×10 <sup>5</sup>	-20	1434	-26	7
R4E1EC	8×10 <sup>6</sup>	8×10 <sup>7</sup>	75553	82557	2997	3251	15653	31961	130	266	-20	-39	315	34696	-23	294	-26	-38
R4E1FC	4×10 <sup>7</sup>	4×10 <sup>7</sup>	4×10 <sup>4</sup>	4×10 <sup>5</sup>	16017	17730	3×10 <sup>5</sup>	5×10 <sup>5</sup>	3140	5586	101	173	643	1×10 <sup>5</sup>	-20	1253	-26	0
R4E1FC.L	8×10 <sup>6</sup>	1×10 <sup>7</sup>	83536	97750	3317	3859	16402	21202	138	159	-19	-44	301	33112	-23	278	-26	-39
R4STAC	9×10 <sup>6</sup>	1×10 <sup>7</sup>	94188	1×10 <sup>5</sup>	3743	5146	45510	63122	429	578	-8	-27	1310	37004	-13	317	-25	-37
R5E1AC	7×10 <sup>6</sup>	8×10 <sup>6</sup>	70290	78739	2787	3098	36378	45202	337	399	-11	-34	232	34696	-24	294	-26	-38
<b>OttExtra</b> [22]																		
m15b6	1663	2165	-10	-35	-25	-53	247	1212	-24	-41	-26	-52	427	27296	-22	220	-26	-41
s11WW	1450	7562	19	-25	-23	-51	272	1594	-24	-37	-26	-52	1056	39024	-16	337	-25	-36
s25WW	7455	-50221	51	63	-23	-49	2×10 <sup>5</sup>	4×10 <sup>5</sup>	2279	3801	66	102	1×10 <sup>5</sup>	1×10 <sup>7</sup>	1075	1×10 <sup>5</sup>	18	5236
<b>MurRem</b> [71]																		
20_3.4	4253	9073	16	34	-24	-50	15668	22046	130	167	-20	-43	-21	-32	-27	-53	-26	-52
12_3.2	1461	7459	-12	18	-25	-51	7905	14088	53	88	-23	-47	-25	-47	-27	-54	-26	-52
15_3.2	7450	10699	48	51	-23	-50	10862	17337	82	120	-22	-45	-25	-12	-27	-53	-26	-52
<b>Yak</b> [58]																		
s12_matter	212	349	-24	-53	-26	-53	777	1257	-19	-41	-26	-52	-24	-31	-27	-53	-26	-52
s15_matter	194	504	-24	-52	-26	-54	902	2071	-17	-33	-26	-51	-25	5	-27	-53	-26	-52
s25_matter	1099	1381	-15	-43	-25	-53	726	2080	-19	-32	-26	-52	-16	-25	-26	-53	-26	-52

waveform power spectra, which would remove any potentially problematic phase information.

### 3. Neutrino Mechanism

We test SMEE's ability to identify GW signals emitted by core-collapse supernovae exploding via the neutrino mechanism in two ways. First, we remove three

randomly selected **Mur** waveforms (**Mur** waveforms 20\_3.4, 12\_3.4, and 15\_3.2, labeling this set as **MurRem**) from the **Mur** catalog, recompute the PCs without these 3 waveforms, then run SMEE to compute  $\log B_{SN}$  for the three **MurRem** waveforms. The results, listed in Tab. III, show that SMEE is able to correctly identify **MurRem** waveforms as GW signals consistent with the **Mur** catalog with strong evidence out to a distance of  $\sim 2$  kpc and even at 10 kpc two out of three **MurRem** waveforms are picked

out of the noise (though with relatively weak evidence). This is consistent with the overall results for waveforms belonging to the **Mur** catalog discussed in Sec. IV C.

However, a comparison of the right panel of Fig. 9 with Fig. 5 shows that the **MurRem** waveforms require an SNR that is more than twice as high to reach values of  $\log B_{SN}$  at which we can consider them to be distinct from Gaussian noise. This is most likely due to the rather large diversity of **Mur** waveforms. Components of relevance to the **MurRem** waveforms are apparently not captured in the first 7 PCs when these waveforms are not included in the PCA.

A yet more stringent test is enabled by the waveforms of the **Yak** catalog (see Sec. II A 1) that were obtained with a completely different numerical code. We inject the three available **Yak** waveforms into Advanced LIGO noise and run SMEE on them to compute  $\log B_{SN}$ . We list the results in Tab. III. SMEE correctly and clearly associates the **Yak** waveforms with the **Mur** PCs at 0.2 kpc. At 2 kpc, the association is still possible, but at 10 kpc the **Yak** waveforms appear to be most consistent with noise for SMEE. The right panel of Fig. 9 shows that the **Yak** waveforms require an SNR to be clearly associated with the neutrino mechanism that is  $\gtrsim 7$  times higher than for **MurRem** waveforms and more than  $\sim 17$  times higher than for **Mur** waveforms. This rather disappointing result can be explained as follows: While the **Yak** waveforms are qualitatively very similar to the **Mur** waveforms, they differ significantly in quantitative aspects. The **Yak** waveforms are generally only half as long ( $\sim 1$  s for **Mur** and 0.5 s for **Yak**, whose models explode much earlier than the **Mur** models). Furthermore, the **Yak** waveforms have considerably more power at frequencies above  $\sim 800$  Hz and their energy spectra peak at  $\sim 1000$  Hz while most of the emission in the **Mur** waveforms occurs at or below  $\sim 400$  Hz. This may be due to the more simplified treatment of gravity and neutrino microphysics and transport in the study of Murphy *et al.* [71] underlying the **Mur** catalog compared to the work of Yakunin *et al.* [58] that led to the **Yak** catalog.

## V. SUMMARY AND DISCUSSION

In this article, we have described the Supernova Model Evidence Extractor (SMEE), a novel Bayesian approach to inferring physical information from observations of GW bursts emitted in stellar collapse and core-collapse supernovae. SMEE decomposes catalogs of simulated GW signals into their principle components and employs the Nested Sampling algorithm to compute the evidence that a trial signal injected into GW detector noise belongs to a given catalog.

It is evident that core-collapse supernovae are powered by the release of gravitational energy in gravitational collapse, but the central unsolved problem of core-collapse supernova theory is by what mechanism this energy is transferred from the collapsed core to the stel-

lar mantle to drive the explosion. Simulations show that multi-dimensional GW-emitting dynamics is a crucial ingredient to all potential explosion mechanisms. Hence, GWs could be ideal probes for the core-collapse supernova mechanism, provided that different mechanism models have clearly distinct GW signatures. In this paper, we have considered (i) the neutrino mechanism, (ii) the magnetorotational mechanism, and (iii) the acoustic mechanism for core-collapse supernova explosions and have treated mechanism as equally probable (i.e., having the same prior probability). The primary and dominant multi-dimensional dynamics and GW emission processes of these mechanisms are, ordered in the above order of mechanisms, (i) convection/turbulence and accretion downstreams, (ii) rapid rotation, and (iii) proton-neutron star core pulsations, respectively.

Using GW signal catalogs based on simulations representative of these three mechanisms, we have applied SMEE to infer the explosion mechanism underlying trial waveforms injected into simulated Advanced LIGO noise. Our results show that our Bayesian approach is capable of identifying any of the considered explosion mechanisms with high confidence for core collapse events occurring out to a distance of  $\sim 2$  kpc even with only rudimentary knowledge of the precise shape of the expected signal.

Magnetorotational explosions, leading to particularly strong GW signals with very robust common features, can be clearly identified throughout the Milky Way ( $D \gtrsim 10$  kpc). Our results also suggest that it is possible to further distinguish between rapidly rotating accretion-induced collapse of massive white dwarfs and rapidly rotating iron core collapse for events occurring in the galaxy, provided that the differences between the GW signals predicted by current simulations of rotating accretion-induced collapse and rotating iron core collapse are reliable.

GW signals emitted by neutrino-driven explosions have systematically lower amplitudes and, hence, are harder to distinguish from detector noise. Moreover, GW emission from convection/turbulence and accretion downstreams has not been as extensively studied, the available number of model waveforms is an order of magnitude smaller, and the currently predicted GW signals are not as reliable as in the case of rapidly rotating collapse. This reduces the efficacy of the principal component decomposition and, combined with the overall weakness of the signals, limits SMEE's robust reach for the neutrino mechanism to  $\lesssim 2$  kpc for the currently available set of model waveforms and Advanced LIGO in broadband configuration.

The GW signals from core-collapse supernovae driven by the acoustic mechanism are strong and would likely be detectable by Advanced LIGO throughout the Milky Way. However, the set of available model waveforms for this signal type is very limited and individual waveforms differ significantly at large scales in the time domain while having similar frequency content. The inclusion of phase information (by computing PCs in the time domain and using their full complex Fourier transforms) in

the current incarnation of SMEE is sub-optimal for such signals and the PCs only inefficiently span the space of possible waveforms. This makes it difficult for SMEE to clearly identify the acoustic mechanism for core collapse events occurring at distances significantly greater than  $\sim 2$  kpc.

This study is the first systematic attempt at inferring core-collapse supernova physics from observations of GW bursts emitted by multiple different underlying mechanisms. While this is a significant step beyond previous work that focused only on GWs from rotating core collapse [32, 39], our present study still suffers from a number of simplifying assumptions: We have considered only a single detector, Gaussian noise, and linearly polarized, optimally oriented GW emission. Real core-collapse supernovae will emit in both GW polarizations and will have arbitrary, generally non-optimal orientation with respect to observatories on Earth. Advanced LIGO class GW detectors will operate as networks and observations will be coincident between three or even four detectors, and the noise backgrounds of these detectors will not be stationary and Gaussian, but non-stationary and glitchy. We have also been agnostic with regard to the prior probability of the three considered core-collapse supernova mechanisms. Input from astronomical observations and theory could be used to generate (approximate) prior probabilities for each mechanism, which could then be included in a future analysis.

Near future work on SMEE will be directed towards incorporating detector networks, variations in detector configurations and sensitivities, both GW polarizations, non-Gaussian noise, arbitrary source-detector orientations, and improved principal component analysis with and without reliance on the signal phase. However, even with these improvements, successful extraction of physics

from core-collapse supernova GW signals will crucially depend on the availability of extensive catalogs of reliable predictions for both  $h_+$  and  $h_\times$ . These must be provided by the core-collapse supernova modeling community on the basis of full 3D simulations.

## Acknowledgments

We thank the Basel, MPA Garching, ORNL, and Princeton core-collapse supernova modeling groups for making their gravitational waveforms publicly available. We are happy to acknowledge helpful exchanges with E. Abdikamalov, A. Burrows, Y. Chen, D. Chernoff, N. Christensen, T. Loredo, J. Murphy, J. Nordhaus, L. Santamaria, B. Schutz, M. Vallisneri, A. Weinstein, S. Wesolowski and the core-collapse supernova working group of the LIGO Scientific Collaboration and the Virgo collaboration.

CDO is supported in part by the Sherman Fairchild Foundation and by the National Science Foundation under grant number PHY-0904015. Some of the results presented in this article were obtained through computations on the Caltech compute cluster “Zwicky” (NSF MRI award No. PHY-0960291), on the NSF Teragrid under grant TG-PHY100033, on machines of the Louisiana Optical Network Initiative under grant Ioni\_numrel07, and at the National Energy Research Scientific Computing Center (NERSC), which is supported by the Office of Science of the US Department of Energy under contract DE-AC02-05CH11231.

ISH and JL gratefully acknowledge the support of the UK Science and Technology Facilities Council and the Scottish Universities Physics Alliance (SUPA).

- 
- [1] W. Baade and F. Zwicky. *Proc. Nat. Acad. Sci.*, 20:259, 1934.
  - [2] H. A. Bethe. *Rev. Mod. Phys.*, 62:801, 1990.
  - [3] H.-T. Janka, K. Langanke, A. Marek, G. Martínez-Pinedo, and B. Müller. *Phys. Rep.*, 442:38, 2007.
  - [4] P. Goldreich and S. V. Weber. *Astrophys. J.*, 238:991, 1980.
  - [5] A. Yahil. *Astrophys. J.*, 265:1047, 1983.
  - [6] E. O’Connor and C. D. Ott. *Astrophys. J.*, 730:70, 2011.
  - [7] L. Wang and J. C. Wheeler. *Ann. Rev. Astron. Astroph.*, 46:433, 2008.
  - [8] N. Smith, S. B. Cenko, N. Butler, J. S. Bloom, M. M. Kasliwal, A. Horesh, S. R. Kulkarni, N. M. Law, P. E. Nugent, E. O. Ofek, D. Poznanski, R. M. Quimby, B. Sesar, S. Ben-Ami, I. Arcavi, A. Gal-Yam, D. Polishook, D. Xu, O. Yaron, D. A. Frail, and M. Sullivan. *Mon. Not. Roy. Astron. Soc.*, 420:1135, 2012.
  - [9] T. Lund, A. Marek, C. Lunardini, H.-T. Janka, and G. Raffelt. *Phys. Rev. D*, 82:063007, 2010.
  - [10] T. D. Brandt, A. Burrows, C. D. Ott, and E. Livne. *Astrophys. J.*, 728:8, 2011.
  - [11] K. Hirata, T. Kajita, M. Koshiba, M. Nakahata, and Y. Oyama. *Phys. Rev. Lett.*, 58:1490, 1987.
  - [12] C. D. Ott. *Class. Quantum Grav.*, 26:063001, 2009.
  - [13] K. Kotake. *submitted to a special issue of Comptes Rendus Physique “Gravitational Waves (from detectors to astrophysics)”*, *ArXiv:1110.5107*, 2011.
  - [14] B. Müller, H.-T. Janka, and A. Marek. *Submitted to the Astrophys. J. ArXiv:1202.0815*, 2012.
  - [15] E. Müller, H.-T. Janka, and A. Wongwathanarat. *Astron. Astrophys.*, 537:A63, 2012.
  - [16] K. Kotake, W. Iwakami Nakano, and N. Ohnishi. *Astrophys. J. in press; arXiv:1106.0544*, 736:124, 2011.
  - [17] H. Dimmelmeier, C. D. Ott, A. Marek, and H.-T. Janka. *Phys. Rev. D*, 78:064056, 2008.
  - [18] T. Takiwaki and K. Kotake. *Astrophys. J.*, 743:30, 2011.
  - [19] A. Marek and H.-T. Janka. *Astrophys. J.*, 694:664, 2009.
  - [20] B. P. Abbott et al. (LIGO Scientific Collaboration). *Rep. Prog. Phys.*, 72:076901, 2009.
  - [21] K. Kuroda and LCGT Collaboration. *Class. Quantum Grav.*, 27:084004, 2010.
  - [22] C. D. Ott, A. Burrows, L. Dessart, and E. Livne. *Phys.*

- Rev. Lett.*, 96:201102, 2006.
- [23] C. D. Ott, C. Reisswig, E. Schnetter, E. O'Connor, U. Sperhake, F. Löffler, P. Diener, E. Abdikamalov, I. Hawke, and A. Burrows. *Phys. Rev. Lett.*, 106:161103, 2011.
- [24] C. Fryer and K. C. B. New. *Liv. Rev. Rel.*, 14:1, 2011.
- [25] M. Punturo et al. *Class. Quantum Grav.*, 27:194002, 2010.
- [26] B. J. Owen and B. S. Sathyaprakash. *Phys. Rev. D*, 60:022002, 1999.
- [27] A. C. Searle, P. J. Sutton, and M. Tinto. *Class. Quantum Grav.*, 26:155017, 2009.
- [28] Y. Gürsel and M. Tinto. *Phys. Rev. D*, 40:3884, 1989.
- [29] É. É. Flanagan and S. A. Hughes. *Phys. Rev. D*, 57:4566, 1998.
- [30] S. Klimentenko, S. Mohanty, M. Rakhmanov, and G. Mitselmakher. *Phys. Rev. D*, 72:122002, 2005.
- [31] S. Klimentenko, I. Yakushin, A. Mercer, and G. Mitselmakher. *Class. Quantum Grav.*, 25:114029, 2008.
- [32] T. Z. Summerscales, A. Burrows, L. S. Finn, and C. D. Ott. *Astrophys. J.*, 678:1142, 2008.
- [33] C. D. Ott, A. Burrows, E. Livne, and R. Walder. *Astrophys. J.*, 600:834, 2004.
- [34] P. R. Brady and S. Ray-Majumder. *Class. Quantum Grav.*, 21:1839, 2004.
- [35] T. Zwerger and E. Müller. *Astron. Astrophys.*, 320:209, 1997.
- [36] I. S. Heng. *Class. Quantum Grav.*, 26:105005, 2009.
- [37] K. V. Mardia, J. T. Kent, and J. M. Bibby. *Multivariate analysis*. Academic Press, London, UK, 1979.
- [38] K. Cannon, A. Chapman, C. Hanna, D. Keppel, A. C. Searle, and A. J. Weinstein. *Phys. Rev. D*, 82:044025, 2010.
- [39] C. Röver, M.-A. Bizouard, N. Christensen, H. Dimmelmeier, I. S. Heng, and R. Meyer. *Phys. Rev. D*, 80:102004, 2009.
- [40] D. S. Sivia and J. Skilling. *Data Analysis: A Bayesian Tutorial*. Oxford University Press, Oxford, UK, second edition, 2006.
- [41] A. Burrows, L. Dessart, E. Livne, C. D. Ott, and J. Murphy. *Astrophys. J.*, 664:416, 2007.
- [42] A. Burrows, E. Livne, L. Dessart, C. D. Ott, and J. Murphy. *Astrophys. J.*, 640:878, 2006.
- [43] A. Burrows, E. Livne, L. Dessart, C. D. Ott, and J. Murphy. *Astrophys. J.*, 655:416, 2007.
- [44] C. D. Ott. *Class. Quantum Grav.*, 26:204015, 2009.
- [45] J. Skilling. Nested Sampling. In R. Fischer, R. Preuss, & U. V. Toussaint, editor, *AIP Conference Proceedings: 24th International Workshop on Bayesian Inference and Maximum Entropy Methods in Science and Engineering*, volume 735, page 395, 2004.
- [46] E. B. Abdikamalov, C. D. Ott, L. Rezzolla, L. Dessart, H. Dimmelmeier, A. Marek, and H.-T. Janka. *Phys. Rev. D*, 81:044012, 2010.
- [47] C. D. Ott, E. P. O'Connor, and B. Dasgupta. *Proceedings of the HAVSE 2011 conference*. *ArXiv:1111.6282*, 2011.
- [48] M. Hamuy. *Astrophys. J.*, 582:905, 2003.
- [49] W. D. Arnett. *Canadian Journal of Physics*, 44:2553, 1966.
- [50] S. A. Colgate and R. H. White. *Astrophys. J.*, 143:626, 1966.
- [51] H. A. Bethe and J. R. Wilson. *Astrophys. J.*, 295:14, 1985.
- [52] M. Liebendörfer, M. Rampp, H.-T. Janka, and A. Mezzacappa. *Astrophys. J.*, 620:840, 2005.
- [53] M. Rampp and H.-T. Janka. *Astron. Astrophys.*, 396:361, 2002.
- [54] T. A. Thompson, A. Burrows, and P. A. Pinto. *Astrophys. J.*, 592:434, 2003.
- [55] F. S. Kitaura, H.-T. Janka, and W. Hillebrandt. *Astron. Astrophys.*, 450:345, 2006.
- [56] R. Buras, H.-T. Janka, M. Rampp, and K. Kifonidis. *Astron. Astrophys.*, 457:281, 2006.
- [57] Y. Suwa, K. Kotake, T. Takiwaki, S. C. Whitehouse, M. Liebendörfer, and K. Sato. *Pub. Astr. Soc. Jap.*, 62:L49, 2010.
- [58] K. N. Yakunin, P. Marronetti, A. Mezzacappa, S. W. Bruenn, C.-T. Lee, M. A. Chertkow, W. R. Hix, J. M. Blondin, E. J. Lentz, O. E. Bronson Messer, and S. Yoshida. *Class. Quantum Grav.*, 27:194005, 2010.
- [59] C. D. Ott, A. Burrows, L. Dessart, and E. Livne. *Astrophys. J.*, 685:1069, 2008.
- [60] J. W. Murphy and A. Burrows. *Astrophys. J.*, 688:1159, 2008.
- [61] C. L. Fryer and M. S. Warren. *Astrophys. J.*, 601:391, 2004.
- [62] J. Nordhaus, A. Burrows, A. Almgren, and J. Bell. *Astrophys. J.*, 720:694, 2010.
- [63] T. Kuroda, K. Kotake, and T. Takiwaki. *Submitted to the Astrophys. J.*; *arXiv:1202.2487*, 2012.
- [64] W. Iwakami, K. Kotake, N. Ohnishi, S. Yamada, and K. Sawada. *Astrophys. J.*, 700:232, 2009.
- [65] T. Takiwaki, K. Kotake, and Y. Suwa. *Submitted to the Astrophys. J.*, *arXiv:1108.3989*, 2011.
- [66] F. Hanke, A. Marek, B. Müller, and H.-T. Janka. *Submitted to the Astrophys. J.*, *arXiv:1108.4355*, 2011.
- [67] J. M. Blondin, A. Mezzacappa, and C. DeMarino. *Astrophys. J.*, 584:971, 2003.
- [68] T. Foglizzo, P. Galletti, L. Scheck, and H.-T. Janka. *Astrophys. J.*, 654:1006, 2007.
- [69] T. Foglizzo, L. Scheck, and H.-T. Janka. *Astrophys. J.*, 652:1436, 2006.
- [70] L. Scheck, H.-T. Janka, T. Foglizzo, and K. Kifonidis. *Astron. Astrophys.*, 477:931, 2008.
- [71] J. W. Murphy, C. D. Ott, and A. Burrows. *Astrophys. J.*, 707:1173, 2009.
- [72] E. Müller, M. Rampp, R. Buras, H.-T. Janka, and D. H. Shoemaker. *Astrophys. J.*, 603:221, 2004.
- [73] A. Marek, H.-T. Janka, and E. Müller. *Astron. Astrophys.*, 496:475, 2009.
- [74] K. Kotake, N. Ohnishi, and S. Yamada. *Astrophys. J.*, 655:406, 2007.
- [75] C. L. Fryer, D. E. Holz, and S. A. Hughes. *Astrophys. J.*, 609:288, 2004.
- [76] K. Kotake, W. Iwakami, N. Ohnishi, and S. Yamada. *Astrophys. J. Lett.*, 697:L133, 2009.
- [77] R. Epstein. *Astrophys. J.*, 223:1037, 1978.
- [78] A. Burrows and J. Hayes. *Phys. Rev. Lett.*, 76:352, 1996.
- [79] E. Müller and H.-T. Janka. *Astron. Astrophys.*, 317:140, 1997.
- [80] C. L. Fryer and A. Heger. *Astrophys. J.*, 541:1033, 2000.
- [81] R. Walder, A. Burrows, C. D. Ott, E. Livne, I. Lichtenstadt, and M. Jarrar. *Astrophys. J.*, 626:317, 2005.
- [82] S. A. Balbus and J. F. Hawley. *Astrophys. J.*, 376:214, 1991.
- [83] M. Obergaulinger, P. Cerdá-Durán, E. Müller, and

- M. A. Aloy. *Astron. Astrophys.*, 498:241, 2009.
- [84] URL <http://www.stellarcollapse.org/gwcatalog>. Ott et al. Gravitational Wave Catalog.
- [85] K. S. Thorne. In S. W. Hawking and W. Israel, editors, *300 Years of Gravitation*, Cambridge, UK, 1987. Cambridge University Press.
- [86] A. Marek, H. Dimmelmeier, H.-T. Janka, E. Müller, and R. Buras. *Astron. Astrophys.*, 445:273, 2006.
- [87] URL <http://astrodev.phys.utk.edu/chimera/>. ORNL/FAU Gravitational Wave Signal Catalog.
- [88] C. D. Ott, A. Burrows, T. A. Thompson, E. Livne, and R. Walder. *Astrophys. J. Suppl. Ser.*, 164:130, 2006.
- [89] A. Heger, S. E. Woosley, and H. C. Spruit. *Astrophys. J.*, 626:350, 2005.
- [90] J. C. Wheeler, I. Yi, P. Höflich, and L. Wang. *Astrophys. J.*, 537:810, 2000.
- [91] G. S. Bisnovatyi-Kogan. *Astron. Zh.*, 47:813, 1970.
- [92] K. Kotake, S. Yamada, K. Sato, K. Sumiyoshi, H. Ono, and H. Suzuki. *Phys. Rev. D*, 69:124004, 2004.
- [93] M. Obergaulinger, M. A. Aloy, H. Dimmelmeier, and E. Müller. *Astron. Astrophys.*, 457:209, 2006.
- [94] M. Shibata, Y. T. Liu, S. L. Shapiro, and B. C. Stephens. *Phys. Rev. D*, 74:104026, 2006.
- [95] L. Dessart, A. Burrows, E. Livne, and C. D. Ott. *Astrophys. J. Lett.*, 673:L43, 2008.
- [96] P. Cerdá-Durán, J. A. Font, L. Antón, and E. Müller. *Astron. Astrophys.*, 492:937, 2008.
- [97] S. E. Woosley and A. Heger. *Astrophys. J.*, 637:914, 2006.
- [98] H. Dimmelmeier, C. D. Ott, H.-T. Janka, A. Marek, and E. Müller. *Phys. Rev. Lett.*, 98:251101, 2007.
- [99] C. D. Ott, H. Dimmelmeier, A. Marek, H.-T. Janka, I. Hawke, B. Zink, and E. Schnetter. *Phys. Rev. Lett.*, 98:261101, 2007.
- [100] URL [http://www.mpa-garching.mpg.de/rel\\_hydro/wave\\_catalogue.shtml](http://www.mpa-garching.mpg.de/rel_hydro/wave_catalogue.shtml). MPA Garching Gravitational Wave Signal Catalog.
- [101] S. Scheidegger, T. Fischer, S. C. Whitehouse, and M. Liebendörfer. *Astron. Astrophys.*, 490:231, 2008.
- [102] S. Scheidegger, R. Käppeli, S. C. Whitehouse, T. Fischer, and M. Liebendörfer. *Astron. Astrophys.*, 514:A51, 2010.
- [103] M. Shibata and Y.-I. Sekiguchi. *Phys. Rev. D*, 71:024014, 2005.
- [104] M. Rampp, E. Müller, and M. Ruffert. *Astron. Astrophys.*, 332:969, 1998.
- [105] W. Fu and D. Lai. *Mon. Not. Roy. Astron. Soc.*, 413:2207, 2011.
- [106] M. Liebendörfer. *Astrophys. J.*, 633:1042, 2005.
- [107] URL [http://phys-merger.physik.unibas.ch/~simon/gw\\_data](http://phys-merger.physik.unibas.ch/~simon/gw_data). Basel GW Signal Catalog.
- [108] L. Dessart, A. Burrows, E. Livne, and C. D. Ott. *Astrophys. J.*, 669:585, 2007.
- [109] N. N. Weinberg and E. Quataert. *Mon. Not. Roy. Astron. Soc.*, 387:L64, 2008.
- [110] M. Pitkin, S. Reid, S. Rowan, and J. Hough. *Liv. Rev. Rel.*, 14(5), 2011.
- [111] W. Del Pozzo, J. Veitch, and A. Vecchio. *Phys. Rev. D*, 83:082002, 2011.
- [112] J. Veitch and A. Vecchio. *Phys. Rev. D*, 81:062003, 2010.
- [113] David Shoemaker. Advanced ligo anticipated sensitivity curves. Technical Report LIGO-T0900288-v3, LIGO Scientific Collaboration, 2010.
- [114] URL <http://ligo-vcs.phys.uwm.edu/wsvn/MatApps/packages/stochastic/trunk/PreProcessing/detectorNoise.m>. LIGO Scientific Collaboration MatApps Package – noise generation code.
- [115] J. Abadie et al. (LIGO and VIRGO Scientific Collaborations). *Class. Quantum Grav.*, 27:173001, 2010.
- [116] N. Smith, K. H. Hinkle, and N. Ryde. *Astron. J.*, 137:3558, 2009.
- [117] G. M. Harper, A. Brown, and E. F. Guinan. *Astron. J.*, 135:1430, 2008.
- [118] M. M. Hohle, R. Neuhäuser, and B. F. Schutz. *Astron. Nachr.*, 331:349, 2010.
- [119] URL <http://www.stellarcollapse.org/SMEE>. Supernova Model Evidence Extractor (SMEE) Supplemental Data.