



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Transverse momentum cross section of e^+e^- pairs in the Z-boson region from $pp[\overline{}]$ collisions at $\sqrt{s}=1.96$ TeV

T. Aaltonen *et al.* (CDF Collaboration)

Phys. Rev. D **86**, 052010 — Published 26 September 2012

DOI: [10.1103/PhysRevD.86.052010](https://doi.org/10.1103/PhysRevD.86.052010)

1 **Transverse momentum cross section of e^+e^- pairs in the Z -boson region from $p\bar{p}$**
2 **collisions at $\sqrt{s} = 1.96$ TeV**

3 T. Aaltonen,²¹ B. Álvarez González^{z,9} S. Amerio,⁴⁰ D. Amidei,³² A. Anastassov^{x,15} A. Annovi,¹⁷ J. Antos,¹²
4 G. Apollinari,¹⁵ J.A. Appel,¹⁵ T. Arisawa,⁵⁴ A. Artikov,¹³ J. Asaadi,⁴⁹ W. Ashmanskas,¹⁵ B. Auerbach,⁵⁷
5 A. Aurisano,⁴⁹ F. Azfar,³⁹ W. Badgett,¹⁵ T. Bae,²⁵ A. Barbaro-Galtieri,²⁶ V.E. Barnes,⁴⁴ B.A. Barnett,²³
6 P. Barria^{hh,42} P. Bartos,¹² M. Baucé^{ff,40} F. Bedeschi,⁴² S. Behari,²³ G. Bellettini^{gg,42} J. Bellinger,⁵⁶
7 D. Benjamin,¹⁴ A. Beretvas,¹⁵ A. Bhatti,⁴⁶ D. Bisello^{ff,40} I. Bizjak,²⁸ K.R. Bland,⁵ B. Blumenfeld,²³ A. Bocci,¹⁴
8 A. Bodek,⁴⁵ D. Bortoletto,⁴⁴ J. Boudreau,⁴³ A. Boveia,¹¹ L. Brigliadori^{ee,6} C. Bromberg,³³ E. Brucken,²¹
9 J. Budagov,¹³ H.S. Budd,⁴⁵ K. Burkett,¹⁵ G. Busetto^{ff,40} P. Bussey,¹⁹ A. Buzatu,³¹ A. Calamba,¹⁰ C. Calancha,²⁹
10 S. Camarda,⁴ M. Campanelli,²⁸ M. Campbell,³² F. Canelli,^{11,15} B. Carls,²² D. Carlsmith,⁵⁶ R. Carosi,⁴²
11 S. Carrillo^{m,16} S. Carron,¹⁵ B. Casal^{k,9} M. Casarsa,⁵⁰ A. Castro^{ee,6} P. Catastini,²⁰ D. Cauz,⁵⁰ V. Cavaliere,²²
12 M. Cavalli-Sforza,⁴ A. Cerri^{f,26} L. Cerrito^{s,28} Y.C. Chen,¹ M. Chertok,⁷ G. Chiarelli,⁴² G. Chlachidze,¹⁵
13 F. Chlebana,¹⁵ K. Cho,²⁵ D. Chokheli,¹³ W.H. Chung,⁵⁶ Y.S. Chung,⁴⁵ M.A. Ciocci^{hh,42} A. Clark,¹⁸ C. Clarke,⁵⁵
14 G. Compostella^{ff,40} M.E. Convery,¹⁵ J. Conway,⁷ M. Corbo,¹⁵ M. Cordelli,¹⁷ C.A. Cox,⁷ D.J. Cox,⁷ F. Crescioli^{gg,42}
15 J. Cuevas^{z,9} R. Culbertson,¹⁵ D. Dagenhart,¹⁵ N. d'Ascenzo^{w,15} M. Datta,¹⁵ P. de Barbaro,⁴⁵ M. Dell'Orso^{gg,42}
16 L. Demortier,⁴⁶ M. Deninno,⁶ F. Devoto,²¹ M. d'Errico^{ff,40} A. Di Canto^{gg,42} B. Di Ruzza,¹⁵ J.R. Dittmann,⁵
17 M. D'Onofrio,²⁷ S. Donati^{gg,42} P. Dong,¹⁵ M. Dorigo,⁵⁰ T. Dorigo,⁴⁰ K. Ebina,⁵⁴ A. Elagin,⁴⁹ A. Eppig,³²
18 R. Erbacher,⁷ S. Errede,²² N. Ershaidat^{dd,15} R. Eusebi,⁴⁹ S. Farrington,³⁹ M. Feindt,²⁴ J.P. Fernandez,²⁹
19 R. Field,¹⁶ G. Flanagan^{u,15} R. Forrest,⁷ M.J. Frank,⁵ M. Franklin,²⁰ J.C. Freeman,¹⁵ Y. Funakoshi,⁵⁴ I. Furic,¹⁶
20 M. Gallinaro,⁴⁶ J.E. Garcia,¹⁸ A.F. Garfinkel,⁴⁴ P. Garosi^{hh,42} H. Gerberich,²² E. Gerchtein,¹⁵ S. Giagu,⁴⁷
21 V. Giakoumopoulou,³ P. Giannetti,⁴² K. Gibson,⁴³ C.M. Ginsburg,¹⁵ N. Giokaris,³ P. Giromini,¹⁷ G. Giurgiu,²³
22 V. Glagolev,¹³ D. Glenzinski,¹⁵ M. Gold,³⁵ D. Goldin,⁴⁹ N. Goldschmidt,¹⁶ A. Golossanov,¹⁵ G. Gomez,⁹
23 G. Gomez-Ceballos,³⁰ M. Goncharov,³⁰ O. González,²⁹ I. Gorelov,³⁵ A.T. Goshaw,¹⁴ K. Goulianos,⁴⁶ S. Grinstein,⁴
24 C. Grosso-Pilcher,¹¹ R.C. Group^{53,15} J. Guimaraes da Costa,²⁰ S.R. Hahn,¹⁵ E. Halkiadakis,⁴⁸ A. Hamaguchi,³⁸
25 J.Y. Han,⁴⁵ F. Happacher,¹⁷ K. Hara,⁵¹ D. Hare,⁴⁸ M. Hare,⁵² R.F. Harr,⁵⁵ K. Hatakeyama,⁵ C. Hays,³⁹ M. Heck,²⁴
26 J. Heinrich,⁴¹ M. Herndon,⁵⁶ S. Hewamanage,⁵ A. Hocker,¹⁵ W. Hopkins^{g,15} D. Horn,²⁴ S. Hou,¹ R.E. Hughes,³⁶
27 M. Hurwitz,¹¹ U. Husemann,⁵⁷ N. Hussain,³¹ M. Hussein,³³ J. Huston,³³ G. Introzzi,⁴² M. Iori^{jj,47} A. Ivanov^{p,7}
28 E. James,¹⁵ D. Jang,¹⁰ B. Jayatilaka,¹⁴ E.J. Jeon,²⁵ S. Jindariani,¹⁵ M. Jones,⁴⁴ K.K. Joo,²⁵ S.Y. Jun,¹⁰
29 T.R. Junk,¹⁵ T. Kamon^{25,49} P.E. Karchin,⁵⁵ A. Kashi,⁵ Y. Kato^{o,38} W. Ketchum,¹¹ J. Keung,⁴¹ V. Khotilovich,⁴⁹
30 B. Kilminster,¹⁵ D.H. Kim,²⁵ H.S. Kim,²⁵ J.E. Kim,²⁵ M.J. Kim,¹⁷ S.B. Kim,²⁵ S.H. Kim,⁵¹ Y.K. Kim,¹¹
31 Y.J. Kim,²⁵ N. Kimura,⁵⁴ M. Kirby,¹⁵ S. Klimentenko,¹⁶ K. Knoepfel,¹⁵ K. Kondo,^{54,*} D.J. Kong,²⁵ J. Konigsberg,¹⁶
32 A.V. Kotwal,¹⁴ M. Kreps,²⁴ J. Kroll,⁴¹ D. Krop,¹¹ M. Kruse,¹⁴ V. Krutelyov^{c,49} T. Kuhr,²⁴ M. Kurata,⁵¹
33 S. Kwang,¹¹ A.T. Laasanen,⁴⁴ S. Lami,⁴² S. Lammel,¹⁵ M. Lancaster,²⁸ R.L. Lander,⁷ K. Lannon^{y,36} A. Lath,⁴⁸
34 G. Latino^{hh,42} T. LeCompte,² E. Lee,⁴⁹ H.S. Lee^{q,11} J.S. Lee,²⁵ S.W. Lee^{bb,49} S. Leo^{gg,42} S. Leone,⁴² J.D. Lewis,¹⁵
35 A. Limosani^{t,14} C.-J. Lin,²⁶ M. Lindgren,¹⁵ E. Lipeles,⁴¹ A. Lister,¹⁸ D.O. Litvintsev,¹⁵ C. Liu,⁴³ H. Liu,⁵³ Q. Liu,⁴⁴
36 T. Liu,¹⁵ S. Lockwitz,⁵⁷ A. Loginov,⁵⁷ D. Lucchesi^{ff,40} J. Lueck,²⁴ P. Lujan,²⁶ P. Lukens,¹⁵ G. Lungu,⁴⁶ J. Lys,²⁶
37 R. Lysak^{e,12} R. Madrak,¹⁵ K. Maeshima,¹⁵ P. Maestro^{hh,42} S. Malik,⁴⁶ G. Manca^{a,27} A. Manousakis-Katsikakis,³
38 F. Margaroli,⁴⁷ C. Marino,²⁴ M. Martínez,⁴ P. Mastrandrea,⁴⁷ K. Matera,²² M.E. Mattson,⁵⁵ A. Mazzacane,¹⁵
39 P. Mazzanti,⁶ K.S. McFarland,⁴⁵ P. McIntyre,⁴⁹ R. McNulty^{j,27} A. Mehta,²⁷ P. Mehtala,²¹ C. Mesropian,⁴⁶
40 T. Miao,¹⁵ D. Mietlicki,³² A. Mitra,¹ H. Miyake,⁵¹ S. Moed,¹⁵ N. Moggi,⁶ M.N. Mondragon^{m,15} C.S. Moon,²⁵
41 R. Moore,¹⁵ M.J. Morello^{ii,42} J. Morlock,²⁴ P. Movilla Fernandez,¹⁵ A. Mukherjee,¹⁵ Th. Muller,²⁴ P. Murat,¹⁵
42 M. Mussini^{ee,6} J. Nachtman^{n,15} Y. Nagai,⁵¹ J. Naganoma,⁵⁴ I. Nakano,³⁷ A. Napier,⁵² J. Nett,⁴⁹ C. Neu,⁵³
43 M.S. Neubauer,²² J. Nielsen^{d,26} L. Nodulman,² S.Y. Noh,²⁵ O. Norriella,²² L. Oakes,³⁹ S.H. Oh,¹⁴ Y.D. Oh,²⁵
44 I. Oksuzian,⁵³ T. Okusawa,³⁸ R. Orava,²¹ L. Ortolan,⁴ S. Pagan Griso^{ff,40} C. Pagliarone,⁵⁰ E. Palencia^{f,9}
45 V. Papadimitriou,¹⁵ A.A. Paramonov,² J. Patrick,¹⁵ G. Pauletta^{kk,50} M. Paulini,¹⁰ C. Paus,³⁰ D.E. Pellett,⁷
46 A. Penzo,⁵⁰ T.J. Phillips,¹⁴ G. Piacentino,⁴² E. Pianori,⁴¹ J. Pilot,³⁶ K. Pitts,²² C. Plager,⁸ L. Pondrom,⁵⁶
47 S. Poprocki^{g,15} K. Potamianos,⁴⁴ F. Prokoshin^{cc,13} A. Pranko,²⁶ F. Ptohos^{h,17} G. Punzi^{gg,42} A. Rahaman,⁴³
48 V. Ramakrishnan,⁵⁶ N. Ranjan,⁴⁴ I. Redondo,²⁹ P. Renton,³⁹ M. Rescigno,⁴⁷ T. Riddick,²⁸ F. Rimondi^{ee,6}
49 L. Ristori^{42,15} A. Robson,¹⁹ T. Rodrigo,⁹ T. Rodriguez,⁴¹ E. Rogers,²² S. Rolli^{i,52} R. Roser,¹⁵ F. Ruffini^{hh,42}
50 A. Ruiz,⁹ J. Russ,¹⁰ V. Rusu,¹⁵ A. Safonov,⁴⁹ W.K. Sakumoto,⁴⁵ Y. Sakurai,⁵⁴ L. Santi^{kk,50} K. Sato,⁵¹
51 V. Savelyev^{w,15} A. Savoy-Navarro^{aa,15} P. Schlabach,¹⁵ A. Schmidt,²⁴ E.E. Schmidt,¹⁵ T. Schwarz,¹⁵ L. Scodellaro,⁹
52 A. Scribano^{hh,42} F. Scuri,⁴² S. Seidel,³⁵ Y. Seiya,³⁸ A. Semenov,¹³ F. Sforza^{hh,42} S.Z. Shalhout,⁷ T. Shears,²⁷

53 P.F. Shepard,⁴³ M. Shimojima^v,⁵¹ M. Shochet,¹¹ I. Shreyber-Tecker,³⁴ A. Simonenko,¹³ P. Sinervo,³¹ K. Sliwa,⁵²
 54 J.R. Smith,⁷ F.D. Snider,¹⁵ A. Soha,¹⁵ V. Sorin,⁴ H. Song,⁴³ P. Squillacioti^{hh},⁴² M. Stancari,¹⁵ R. St. Denis,¹⁹
 55 B. Stelzer,³¹ O. Stelzer-Chilton,³¹ D. Stentz^x,¹⁵ J. Strologas,³⁵ G.L. Strycker,³² Y. Sudo,⁵¹ A. Sukhanov,¹⁵
 56 I. Suslov,¹³ K. Takemasa,⁵¹ Y. Takeuchi,⁵¹ J. Tang,¹¹ M. Tecchio,³² P.K. Teng,¹ J. Thom^g,¹⁵ J. Thome,¹⁰
 57 G.A. Thompson,²² E. Thomson,⁴¹ D. Toback,⁴⁹ S. Tokar,¹² K. Tollefson,³³ T. Tomura,⁵¹ D. Tonelli,¹⁵ S. Torre,¹⁷
 58 D. Torretta,¹⁵ P. Totaro,⁴⁰ M. Trovatoⁱⁱ,⁴² F. Ukegawa,⁵¹ S. Uozumi,²⁵ A. Varganov,³² F. Vázquez^m,¹⁶ G. Veley,¹⁵
 59 C. Vellidis,¹⁵ M. Vidal,⁴⁴ I. Vila,⁹ R. Vilar,⁹ J. Vizán,⁹ M. Vogel,³⁵ G. Volpi,¹⁷ P. Wagner,⁴¹ R.L. Wagner,¹⁵
 60 T. Wakisaka,³⁸ R. Wallny,⁸ S.M. Wang,¹ A. Warburton,³¹ D. Waters,²⁸ W.C. Wester III,¹⁵ D. Whiteson^b,⁴¹
 61 A.B. Wicklund,² E. Wicklund,¹⁵ S. Wilbur,¹¹ F. Wick,²⁴ H.H. Williams,⁴¹ J.S. Wilson,³⁶ P. Wilson,¹⁵
 62 B.L. Winer,³⁶ P. Wittich^g,¹⁵ S. Wolbers,¹⁵ H. Wolfe,³⁶ T. Wright,³² X. Wu,¹⁸ Z. Wu,⁵ K. Yamamoto,³⁸
 63 D. Yamato,³⁸ T. Yang,¹⁵ U.K. Yang^r,¹¹ Y.C. Yang,²⁵ W.-M. Yao,²⁶ G.P. Yeh,¹⁵ K. Yiⁿ,¹⁵ J. Yoh,¹⁵ K. Yorita,⁵⁴
 64 T. Yoshida^l,³⁸ G.B. Yu,¹⁴ I. Yu,²⁵ S.S. Yu,¹⁵ J.C. Yun,¹⁵ A. Zanetti,⁵⁰ Y. Zeng,¹⁴ C. Zhou,¹⁴ and S. Zucchelli^{ee6}

(CDF Collaboration)[†]

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

²*Argonne National Laboratory, Argonne, Illinois 60439, USA*

³*University of Athens, 157 71 Athens, Greece*

⁴*Institut de Física d'Altes Energies, ICREA, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

⁵*Baylor University, Waco, Texas 76798, USA*

⁶*Istituto Nazionale di Fisica Nucleare Bologna, ^{ee}University of Bologna, I-40127 Bologna, Italy*

⁷*University of California, Davis, Davis, California 95616, USA*

⁸*University of California, Los Angeles, Los Angeles, California 90024, USA*

⁹*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

¹⁰*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

¹¹*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*

¹²*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

¹³*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

¹⁴*Duke University, Durham, North Carolina 27708, USA*

¹⁵*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*

¹⁶*University of Florida, Gainesville, Florida 32611, USA*

¹⁷*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

¹⁸*University of Geneva, CH-1211 Geneva 4, Switzerland*

¹⁹*Glasgow University, Glasgow G12 8QQ, United Kingdom*

²⁰*Harvard University, Cambridge, Massachusetts 02138, USA*

²¹*Division of High Energy Physics, Department of Physics,*

University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland

²²*University of Illinois, Urbana, Illinois 61801, USA*

²³*The Johns Hopkins University, Baltimore, Maryland 21218, USA*

²⁴*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*

²⁵*Center for High Energy Physics: Kyungpook National University,*

Daegu 702-701, Korea; Seoul National University, Seoul 151-742,

Korea; Sungkyunkwan University, Suwon 440-746,

Korea; Korea Institute of Science and Technology Information,

Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757,

Korea; Chonbuk National University, Jeonju 561-756, Korea

²⁶*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

²⁷*University of Liverpool, Liverpool L69 7ZE, United Kingdom*

²⁸*University College London, London WC1E 6BT, United Kingdom*

²⁹*Centro de Investigaciones Energeticas Medioambientales y Tecnológicas, E-28040 Madrid, Spain*

³⁰*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

³¹*Institute of Particle Physics: McGill University, Montréal, Québec,*

Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia,

Canada V5A 1S6; University of Toronto, Toronto, Ontario,

Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3

³²*University of Michigan, Ann Arbor, Michigan 48109, USA*

³³*Michigan State University, East Lansing, Michigan 48824, USA*

³⁴*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*

³⁵*University of New Mexico, Albuquerque, New Mexico 87131, USA*

³⁶*The Ohio State University, Columbus, Ohio 43210, USA*

³⁷*Okayama University, Okayama 700-8530, Japan*

³⁸*Osaka City University, Osaka 588, Japan*

³⁹*University of Oxford, Oxford OX1 3RH, United Kingdom*

113

114 ⁴⁰*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, ^{ff}University of Padova, I-35131 Padova, Italy*
 115 ⁴¹*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
 116 ⁴²*Istituto Nazionale di Fisica Nucleare Pisa, ^{gg}University of Pisa,*
 117 ^{hh}*University of Siena and ⁱⁱScuola Normale Superiore, I-56127 Pisa, Italy*
 118 ⁴³*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
 119 ⁴⁴*Purdue University, West Lafayette, Indiana 47907, USA*
 120 ⁴⁵*University of Rochester, Rochester, New York 14627, USA*
 121 ⁴⁶*The Rockefeller University, New York, New York 10065, USA*
 122 ⁴⁷*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1,*
 123 ^{jj}*Sapienza Università di Roma, I-00185 Roma, Italy*
 124 ⁴⁸*Rutgers University, Piscataway, New Jersey 08855, USA*
 125 ⁴⁹*Texas A&M University, College Station, Texas 77843, USA*
 126 ⁵⁰*Istituto Nazionale di Fisica Nucleare Trieste/Udine,*
 127 *I-34100 Trieste, ^{kk}University of Udine, I-33100 Udine, Italy*
 128 ⁵¹*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
 129 ⁵²*Tufts University, Medford, Massachusetts 02155, USA*
 130 ⁵³*University of Virginia, Charlottesville, Virginia 22906, USA*
 131 ⁵⁴*Waseda University, Tokyo 169, Japan*
 132 ⁵⁵*Wayne State University, Detroit, Michigan 48201, USA*
 133 ⁵⁶*University of Wisconsin, Madison, Wisconsin 53706, USA*
 134 ⁵⁷*Yale University, New Haven, Connecticut 06520, USA*
 135 (Dated: August 24, 2012)

The transverse momentum cross section of e^+e^- pairs in the Z-boson mass region of 66–116 GeV/ c^2 is precisely measured using Run II data corresponding to 2.1 fb⁻¹ of integrated luminosity recorded by the Collider Detector at Fermilab. The cross section is compared with two quantum chromodynamic calculations. One is a fixed-order perturbative calculation at $\mathcal{O}(\alpha_s^2)$, and the other combines perturbative predictions at high transverse momentum with the gluon resummation formalism at low transverse momentum. Comparisons of the measurement with calculations show reasonable agreement. The measurement is of sufficient precision to allow refinements in the understanding of the transverse momentum distribution.

136 PACS numbers: 13.85.Qk, 12.38.Qk

* Deceased

† With visitors from: ^aIstituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, ^bUniversity of CA Irvine, Irvine, CA 92697, USA, ^cUniversity of CA Santa Barbara, Santa Barbara, CA 93106, USA, ^dUniversity of CA Santa Cruz, Santa Cruz, CA 95064, USA, ^eInstitute of Physics, Academy of Sciences of the Czech Republic, Czech Republic, ^fCERN, CH-1211 Geneva, Switzerland, ^gCornell University, Ithaca, NY 14853, USA, ^hUniversity of Cyprus, Nicosia CY-1678, Cyprus, ⁱOffice of Science, U.S. Department of Energy, Washington, DC 20585, USA, ^jUniversity College Dublin, Dublin 4, Ireland, ^kETH, 8092 Zurich, Switzerland, ^lUniversity of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017, ^mUniversidad Iberoamericana, Mexico D.F., Mexico, ⁿUniversity of Iowa, Iowa City, IA 52242, USA, ^oKinki University, Higashi-Osaka City, Japan 577-8502, ^pKansas State University, Manhattan, KS 66506, USA, ^qKorea University, Seoul, 136-713, Korea, ^rUniversity of Manchester, Manchester M13 9PL, United Kingdom, ^sQueen Mary, University of London, London, E1 4NS, United Kingdom, ^tUniversity of Melbourne, Victoria 3010, Australia, ^uMuons, Inc., Batavia, IL 60510, USA, ^vNagasaki Institute of Applied Science, Nagasaki, Japan, ^wNational Research Nuclear University, Moscow, Russia, ^xNorthwestern University, Evanston, IL 60208, USA, ^yUniversity of Notre Dame, Notre Dame, IN 46556, USA, ^zUniversidad de Oviedo, E-33007 Oviedo, Spain, ^{aa}CNRS-IN2P3, Paris, F-75205 France, ^{bb}Texas Tech University, Lubbock, TX 79609, USA, ^{cc}Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile, ^{dd}Yarmouk University, Irbid 211-63, Jordan,

I. INTRODUCTION

In hadron-hadron collisions at high energies, massive lepton pairs are produced via the Drell–Yan process [1]. In the standard model, colliding partons from the hadrons can interact to form an intermediate W or γ^*/Z vector boson that subsequently decays into a lepton pair. Initial state quantum chromodynamic (QCD) radiation from the colliding partons imparts transverse momentum (P_T) to the boson and produces an accompanying final state jet or jets.

A recent advance in QCD fixed-order perturbative calculations at $\mathcal{O}(\alpha_s^2)$ is the evaluation of the Drell–Yan cross section that is fully exclusive and differential [2]. The exclusive cross section includes both the lepton pair produced via the W or γ^*/Z boson intermediate state, and the associated final state partons. It includes finite boson widths, boson-lepton spin correlations, and $\gamma - Z$ interference for the γ^*/Z intermediate state.

The QCD calculation of the Drell–Yan-process cross section that is differential in transverse momentum for all values of P_T employs a resummation formalism [3] that merges fixed-order calculations with an all-orders sum of large terms from soft and collinear gluon emissions. The dynamics at low P_T is factorized into a calculable perturbative form factor and a hadron-level, non-perturbative one that must be measured. The non-perturbative form factor also includes the effect of the intrinsic P_T of partons in the hadron. Refinement of the phenomenology needs precise measurements of the transverse momentum differential cross section at low P_T from hadron-hadron collisions at various center-of-momentum energies, \sqrt{s} .

Previous $p\bar{p}$ measurements at $\sqrt{s} = 0.63$ TeV [4, 5] support the resummation formalism, but with limited statistics. The next $p\bar{p}$ measurements at $\sqrt{s} = 1.8$ TeV [6–9] contributed to the phenomenology at low P_T [10]. Recent $p\bar{p}$ measurements at $\sqrt{s} = 1.96$ TeV [11] are precise enough to constrain phenomenological calculations of the Drell–Yan lepton pair P_T distribution. Early Large Hadron Collider pp results [12, 13] at $\sqrt{s} = 7$ TeV show agreement with calculations.

In this article, a new and precise measurement of the differential cross section in P_T for Drell–Yan lepton pairs from $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV is presented. The specific Drell–Yan process is $p\bar{p} \rightarrow e^+e^- + X$, where the e^+e^- pair is produced through an intermediate γ^*/Z boson, and X is the hadronic final state associated with the production of the boson. The measurement of the differential cross section is restricted to dielectron pairs within the 66–116 GeV/ c^2 mass range and is fully corrected to include all boson rapidities, electron phase space, and detector effects. Within this mass range, the dielectron pairs originate mostly from the resonant production and decay of Z bosons.

The cross section, measured using 2.1 fb $^{-1}$ of collisions recorded by the Collider Detector at Fermilab (CDF) during 2002–2007, covers $0 < P_T < 350$ GeV/ c . This range is subdivided into variable-width P_T bins. For $P_T < 25$ GeV/ c , the bin width is 0.5 GeV/ c . The cross section presented for each P_T bin is the average bin cross section, $\Delta\sigma/\Delta P_T$, where $\Delta\sigma$ is the cross section in a P_T bin, and ΔP_T its width.

The $\Delta\sigma/\Delta P_T$ measurement depends on the correct modeling of the physics and detector to unfold the effects of the detector acceptance and resolution for the $p\bar{p}$ production of Drell–Yan e^+e^- pairs. The modeling of the physics and detector is data driven. This measurement is an extension of the CDF measurements of the Drell–Yan e^+e^- pair rapidity differential cross section [14], and of the decay-electron angular-distribution coefficients [15] that reflect the polarization state of the intermediate γ^*/Z boson produced in $p\bar{p} \rightarrow \gamma^*/Z + X$. The $\Delta\sigma/\Delta P_T$ measurement uses the same 2.1 fb $^{-1}$ data set and analysis methods developed in those measurements, where both the data and the modeling of the physics and detector are well studied and understood.

Section II provides a brief overview of the QCD calculations of $\Delta\sigma/\Delta P_T$ used for comparison with this measurement. Section III provides a summary of CDF and the Tevatron collider at Fermi National Accelerator Laboratory. Section IV reports the selection of electrons and dielectrons for the $\Delta\sigma/\Delta P_T$ measurement. Section V details the simulation of the data. Section VI describes the cross section and its measurement. Section VII is the summary.

II. QCD CALCULATIONS

For the Drell–Yan process, QCD radiation from the colliding partons of the hadrons in the initial state imparts transverse momentum to the lepton pairs. Fixed-order perturbative calculations are expected to become increasingly reliable with larger transverse momentum. However, the Drell–Yan process has two energy scales: the lepton-pair invariant mass and transverse momentum. Difficulties arise in the perturbative calculation when these two scales differ significantly. This is a QCD multi-scale problem. Simpler perturbative QCD calculations usually have one scale, and this scale is often used as the scale in the strong coupling, α_s , to control accuracy. In addition, all perturbative QCD calculations have an arbitrary mass factorization scale that separates the hard parton scattering from the soft parton distribution functions (PDFs) of the hadrons. With multiple scales, scale issues can be harder to control and quantify.

At the opposite end corresponding to low transverse momentum, large contributions from soft and collinear gluon emissions begin to dominate and limit the applicability of standard perturbative calculations. The QCD resummation

190 methods are used to overcome this limitation [3]. These resummation methods may be viewed as techniques to control
 191 large and unreliable contributions from multiple QCD scales in the low transverse momentum kinematic region.

192 As neither calculation is expected to be accurate over the entire range of P_T , it is useful to compare them with
 193 measurements. Of interest is the low P_T region where the bulk of events is produced. The understanding and
 194 proper modeling of QCD at low P_T is important for many physics measurements. The Drell–Yan process can be
 195 used as a benchmark. The measurement presented here is compared with a recent QCD resummation calculation,
 196 RESBOS [10, 16–18], and a state-of-the-art QCD fixed-order $\mathcal{O}(\alpha_s^2)$ calculation (NNLO) of $\Delta\sigma/\Delta P_T$, FEWZ2 [2, 19].

197 The FEWZ2 NNLO calculation is fully exclusive and differential for the final-state leptons and partons, and includes
 198 γ^*/Z finite decay width and lepton correlation effects. For calculations, the MSTW2008 [20] NNLO nucleon PDFs
 199 with their 90% C.L. uncertainties and the default FEWZ2 electroweak parameters of the Fermi coupling constant
 200 (G_μ) scheme and fine-structure constant at the Z -boson mass ($\alpha_{\text{em}}^{-1} = 128$) are used. The QCD factorization and
 201 renormalization scales are both set to the Z -boson mass. As no significant phase-space restrictions are applied on the
 202 final state, except for the 66–116 GeV/ c^2 dilepton mass range limit, FEWZ2 is used here as an inclusive calculation.
 203 The numerical integration accuracy is set to the 1% level.

204 The RESBOS calculation utilizes the Collins, Soper, and Sterman (CSS) resummation formalism that combines
 205 fixed-order perturbative QCD calculations with an all-orders summation of large terms from gluon emissions [3]. The
 206 CSS cross section consists of two terms: a W function, which contains the large terms from gluon emissions; and a Y
 207 function, which is the fixed-order cross section minus its asymptotic (large gluon emission) terms already in W . The
 208 Y function becomes important as the magnitude of the P_T approaches the lepton-pair invariant mass. After a Fourier
 209 transformation from transverse momentum to its conjugate impact-parameter space (b), the resummation in the W
 210 function is expressed as renormalization group equations [21]. With this formalism, the lepton-pair mass and impact
 211 parameter scales are connected by the renormalization group evolution, through which large perturbative terms are
 212 reliably controlled. At small b , W is evaluated to arbitrary order in the renormalized coupling. At large b , hadron
 213 level, non-perturbative terms that must be measured become dominant. The methodologies at small and large impact
 214 parameters are joined by factorizing W into a perturbative and a non-perturbative form factor. The perturbative
 215 form factor uses the impact parameter, $b_* \equiv b/\sqrt{1 + (b/b_{\text{max}})^2}$, so that it becomes constant in the non-perturbative
 216 region.

217 The CSS gluon resummation W and Y functions should be evaluated to all orders of α_s and then combined to fully
 218 describe the physics at all P_T . However, practical implementations of the CSS gluon resummation formalism evaluate
 219 the perturbative Y function and the perturbative part of the resummed W function term to a finite order in α_s . Even
 220 with a finite order expansion, the CSS gluon resummation formalism provides a good description of the physics at
 221 low lepton-pair P_T . Above a P_T value of about the boson mass, the resummed cross section is dominated by the Y
 222 function and is close to the pure fixed-order calculation. However, in an intermediate P_T zone starting from about
 223 half the boson mass, the cancellation between the W and Y functions evaluated at finite order becomes inadequate
 224 because of an order mismatch. The W perturbative expansion terms are intrinsically all-orders from the underlying
 225 resummation formalism, but the Y terms are strictly finite-order. Within this intermediate P_T zone, $W + Y$ loses
 226 accuracy and requires compensation in practical implementations of the resummation formalism.

227 The RESBOS implementations of the W and Y functions are calculated using CTEQ6.6 PDFs [22], and are provided
 228 within RESBOS as cross-section tables on a grid of the boson mass, transverse momentum, and rapidity. The RESBOS
 229 non-perturbative form factor [10] of the W function for the Drell–Yan process is

$$\exp \left\{ \left[-g_1 - g_2 \ln \frac{Q}{2Q_0} - g_1 g_3 \ln(100x_1 x_2) \right] b^2 \right\},$$

230 where $g_1 = 0.21 \text{ GeV}^2$, $g_2 = 0.68 \text{ GeV}^2$, $g_3 = -0.6$, Q is the lepton pair mass, $Q_0 = 1.6 \text{ GeV}/c^2$ (with $b_{\text{max}} =$
 231 0.5 GeV^{-1} for b_* in the perturbative form factor), and $x_1 x_2 = Q^2/s$. The g_{1-3} are parameters derived from mea-
 232 surements. This form factor describes both low- and high-mass data at various \sqrt{s} from fixed target to colliders.
 233 The specific W and Y function cross-section tables used are W_{321} and Y_k , respectively, and the numerical integration
 234 uncertainties of RESBOS are under 1% and negligible.

235 The CSS gluon resummation W function has three separate perturbative functions: A , B , and C . In the RESBOS
 236 implementation [17] of the W function, W_{321} , those functions are evaluated to $\mathcal{O}(\alpha_s^3)$, $\mathcal{O}(\alpha_s^2)$, and $\mathcal{O}(\alpha_s)$, respectively.
 237 Its Y function is $\mathcal{O}(\alpha_s^2)$. At large P_T , RESBOS utilizes both the resummed cross section, $W + Y$, and the $\mathcal{O}(\alpha_s^2)$ fixed-
 238 order cross section. The resummed cross section becomes inaccurate in the intermediate transverse momentum region
 239 starting from about half of the boson mass because of the intrinsic order mismatch described previously. Therefore,
 240 as the P_T increases, a matching procedure between the resummed and fixed-order cross section is implemented by
 241 RESBOS to provide a reliable prediction over all transverse momentum. This matching is implemented in the Y_k

242 cross-section table¹, and is a non-trivial, phenomenological part of RESBOS. On the other hand, in the transverse
 243 momentum region above the order of the boson mass, the RESBOS calculation and its accuracy are similar to the
 244 FEWZ2 NNLO inclusive calculation considered here. The RESBOS calculation also includes the full γ^*/Z interference
 245 effects with a finite decay-width for the Z boson and with lepton correlations. The dominant electroweak corrections
 246 are included in the calculation using the effective Born approximation, as done in the LEP electroweak precision
 247 measurements.

248 III. THE EXPERIMENTAL APPARATUS

249 The CDF II [23] is a general purpose detector at the Fermilab Tevatron Run II $p\bar{p}$ collider whose center-of-momentum
 250 energy is 1.96 TeV. The CDF positive z -axis is along the proton direction. For particle trajectories, the polar angle θ
 251 is relative to the proton direction and the azimuthal angle ϕ is about the beamline axis. The energy and momentum
 252 of a particle are denoted as E and P , respectively. Their components transverse to the beamline are defined as
 253 $E_T = E \sin \theta$ and $P_T = P \sin \theta$, respectively. The particle rapidity, y , is $y = \frac{1}{2} \ln[(E + P_z c)/(E - P_z c)]$, where P_z
 254 is the component of momentum along the z -axis. The pseudorapidity of a particle trajectory is $\eta = -\ln \tan(\theta/2)$.
 255 Fixed detector coordinates are specified as $(\eta_{\text{det}}, \phi)$, where η_{det} is the pseudorapidity from the detector center ($z = 0$).
 256 Portions of the detector relevant to this analysis are briefly described next.

257 The central tracker (COT) is a 3.1 m long, open cell drift chamber that extends radially from 0.4 m to 1.3 m. The
 258 2.1 m long silicon tracker surrounds the Tevatron beam pipe and is within the inner radius of the COT. Combined,
 259 these two trackers provide efficient, high resolution tracking over $|\eta_{\text{det}}| < 1.3$. Both trackers are immersed in a 1.4 T
 260 axial magnetic field produced by a superconducting solenoid just beyond the outer radius of the COT.

261 Outside the solenoid are the central calorimeters, covering $|\eta_{\text{det}}| < 1.1$. The forward regions, $1.1 < |\eta_{\text{det}}| < 3.6$,
 262 are covered by the end-plug calorimeters. All calorimeters are scintillator-based sampling calorimeters read out with
 263 phototubes. Both calorimeters are segmented along their depth into electromagnetic (EM) and hadronic (HAD)
 264 sections and transversely into projective towers. The EM calorimeter energy resolutions measured in test beams with
 265 electrons are $\sigma/E = 14\%/\sqrt{E_T}$ for the central calorimeter, and $\sigma/E = 16\%/\sqrt{E} \oplus 1\%$ for the plug calorimeter,
 266 where the symbol \oplus is a quadrature sum, and E_T and E are in units of GeV. Both the central and plug EM
 267 calorimeters have preshower and shower-maximum detectors for electromagnetic shower identification and shower
 268 centroid measurements. The combination of the plug shower-maximum detector and silicon tracker provides enhanced
 269 tracking coverage to $|\eta_{\text{det}}| = 2.8$.

270 The Fermilab Tevatron collides bunches of protons and anti-protons at a nominal crossing frequency of 2.5 MHz.
 271 Over 2002–2007 operations, the instantaneous $p\bar{p}$ collision luminosities at the start of collisions increased over an order
 272 of magnitude to $280 \times 10^{30} \text{ cm}^{-2}\text{s}^{-1}$. Collision luminosities are continuously measured by the gas Cherenkov counters
 273 which are just outside the Tevatron beam pipe and are in the region $3.7 < |\eta_{\text{det}}| < 4.7$ [24].

274 The CDF event trigger system has three tiers, L1, L2, and L3. The L1 trigger is entirely implemented in hardware,
 275 is based on trigger primitives, and is synchronous and deadtime-less. Trigger primitives are quantities from the
 276 front-end readout used for trigger decisions. The L2 trigger, which processes events selected by the L1 trigger, is
 277 asynchronous and is a combination of hardware and software that uses L1 primitives along with additional front-end
 278 data. The L3 trigger processes events selected by the L2 trigger and is a speed-optimized version of the CDF offline
 279 reconstruction. Track- and EM-objects, which are available at all trigger levels and are refined at each level, form the
 280 basis of very efficient trigger paths for the electrons used in this measurement.

281 IV. DATA SELECTION

282 The data set consists of 2.1 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ collected during 2002–2007. Collisions producing
 283 massive Drell–Yan dielectron pairs have the following experimental signatures:

- 284 • A large fraction of the electrons have high E_T .
- 285 • There are two well-separated electrons of opposite charge.
- 286 • The electrons tend to be separated from jets and other particles from the interaction.

287 These features are used in the selection of events both at the trigger and analysis levels. Electrons in both the central
 288 and plug calorimeters are selected.

¹ To reduce the time needed to compute the Y_k cross section table to $\mathcal{O}(\alpha_s^2)$, the computation is implemented by an $\mathcal{O}(\alpha_s)$ calculation with boson mass, rapidity, and transverse momentum dependent NNLO-to-NLO K-factors.

A. Triggers

The high E_T electrons are selected from generic $p\bar{p}$ collisions by two non-attenuated (full-rate) triggers: the CENTRAL-18, and Z-NO-TRACK. Each has well-defined L1, L2, and L3 trigger paths for both physics and trigger efficiency measurements. Independent and dedicated trigger paths are used for the efficiency measurements.

The CENTRAL-18 trigger is the inclusive electron trigger for electrons with $E_T > 18$ GeV in the central calorimeter region [23]. A track is required at all trigger levels. Loose criteria applied at each level select candidates that are consistent with an electron showering in the calorimeter, including EM-shower-like lateral shower profile in the EM compartment, EM-shower-like energy leakage in the HAD compartment, and matching between the track and the shower centroid in the EM shower-maximum detector. There is no equivalent inclusive plug electron trigger because the L1 and L2 tracking and the plug calorimeter acceptance do not overlap.

The Z-NO-TRACK trigger identifies dielectrons using solely calorimeter information. No tracking information is used. Electron candidates can be in either the central or plug calorimeter region. Both candidates are required to have $E_T > 18$ GeV. The only other requirement is that shower energy sharing in the EM and HAD compartments be electron-like. While this trigger is specifically for dielectron candidates that are both in the plug calorimeter region, it accepts the small fraction of dielectron events that fail the CENTRAL-18 trigger.

B. Electron Selection

To improve the purity of the sample, CDF standard central and plug [23] electron identification requirements are applied. Fiducial requirements are always applied to ensure that the electrons are in well-instrumented regions of CDF where their reconstruction is well understood and predictable. Each electron candidate is required to have an associated track. Having track matching on both electron candidates significantly reduces backgrounds.

The track vertex position along the beamline (Z_{vtx}) is restricted to the inner region of CDF: $|Z_{\text{vtx}}| < 60$ cm. For 2002–2007 Tevatron operations, 4% of the $p\bar{p}$ luminous region along the beamline is outside this fiducial region. The $p\bar{p}$ collision profile along the beamline is measured by vertexing multiple tracks in minimum-bias events. The multiple track vertexing acceptance is relatively flat within $|Z_{\text{vtx}}| \sim 100$ cm.

As electrons in both the central and plug calorimeter regions are used, there are three exclusive Drell–Yan dielectron topologies: CC, CP, and PP, where the C (P) refers to an electron in the central (plug) calorimeter. In the measurement of the ee -pair P_T distribution, the kinematic region of the ee -pair extends over all rapidities, but is restricted to the 66–116 GeV/ c^2 pair mass range. The kinematic and fiducial regions of acceptance for electrons in the three dielectron topologies are listed below.

1. Central–Central (CC)

- $E_T > 25$ (15) GeV for electron 1 (2)
- $0.05 < |\eta_{\text{det}}| < 1.05$

2. Central–Plug (CP)

- $E_T > 20$ GeV for both electrons
- Central region: $0.05 < |\eta_{\text{det}}| < 1.05$
- Plug region: $1.2 < |\eta_{\text{det}}| < 2.8$

3. Plug–Plug (PP)

- $E_T > 25$ GeV for both electrons
- $1.2 < |\eta_{\text{det}}| < 2.8$

The CC electron E_T selection is asymmetric. Electron 1 has the highest E_T . The asymmetric selection is the result of an optimization based on the decay electron angular distribution measurement [15]. It improves the acceptance in the electron phase space. The PP electron candidates are both required to be in the same end-plug calorimeter, and these pairs extend the rapidity coverage to $|y| \sim 2.9$. At the Tevatron, the kinematic limit for $|y|$ of the ee -pair at the Z-boson mass is 3.1. Drell–Yan dielectrons in opposing end plug calorimeters have little acceptance, tend to be at low ee -pair rapidities, and are overwhelmed by QCD di-jet backgrounds.

As Drell–Yan high- E_T leptons are typically produced in isolation, the electron candidates are required to be isolated from other calorimetric activity. The isolation requirement is that the sum of E_T over towers within a 0.4 isolation cone in (η, ϕ) surrounding the electron cluster be under 4 GeV ($E_{\text{iso}} < 4$ GeV). The towers of the electron cluster

are not included in the sum. While this is a topological selection rather than an electron identification selection, it is included in the electron identification efficiencies.

Electron identification in the central calorimeter region is optimized for electrons of $P_T > 10$ GeV/ c . It utilizes the COT and silicon trackers, the longitudinal and lateral (tower) segmentation of the EM and HAD calorimeter compartments, and the shower-maximum strip detector (CES) within the EM calorimeter. The most discriminating information is provided by the trackers in combination with the CES. An electron candidate must have shower clusters within the EM calorimeter towers and CES that have EM-like lateral shower profiles. A candidate must also have an associated track that extrapolates to the three-dimensional position of the CES shower centroid. The track transverse momentum, P_T , must be consistent with the associated electron shower E_T via an E/P selection when $P_T < 50$ GeV/ c . For both the track matching in the CES and E/P selection, allowances are included for bremsstrahlung energy loss in the tracking volume, which on average is about 20% of a radiation length. The fraction of shower energy in the HAD calorimeter towers behind the EM tower cluster must be consistent with that for electrons ($E_{\text{HAD}}/E_{\text{EM}}$ requirement). These selections are more restrictive than the ones used in the trigger.

The central electron selection as described has high purity and is called the tight central electron (TCE) selection. Its average selection efficiency is 84%. The track-finding efficiency on the associated tracks is 99%. To improve the selection of central dielectrons, a looser selection, called the loose central electron (LCE) selection, is used on the second electron. The LCE selection does not use transverse shower shape constraints, the E/P constraint, nor track matching in the CES. For track associations, the track need only project into the largest-energy calorimeter tower within the cluster of towers associated with the EM shower. For electron candidates that fail the TCE selection, the LCE selection has an average exclusive efficiency of 76%.

Electron identification in the forward plug calorimeter region also utilizes the COT and silicon trackers, the longitudinal and lateral (tower) segmentation of the EM and HAD calorimeter compartments, and the shower-maximum strip detector (PES) within the EM calorimeter. However, as the plug calorimeter geometry is completely different from the central geometry, the details of the identification requirements differ.

The plate-geometry, end-plug calorimeters have projective towers, but these towers are physically much smaller than the central calorimetry towers. EM showers in the plug calorimeter are clustered into “rectangular” 3×3 tower clusters in (η, ϕ) space, with the highest-energy tower in the center. The EM calorimeter energy resolution and lateral shower shapes measured in an electron test beam use 3×3 shower clustering [25]. The EM preshower detector is the first layer of the EM calorimeter and it is instrumented and read out separately. As there are ~ 0.7 radiation lengths of material in front of it, its energy is always included in the EM-cluster shower energy.

An electron in the plug calorimeter, like those in the central region, must also have shower clusters within the EM calorimeter towers and PES that have EM-like lateral shower profiles. The longitudinal $E_{\text{HAD}}/E_{\text{EM}}$ leakage requirement is more restrictive because of the deeper depth of the EM section and the differing collision conditions in the forward region. The plug selection efficiency without the tracking requirement averages about 84%.

Tracks going into the plug calorimeters have limited geometrical acceptance in the COT for $|\eta_{\text{det}}| > 1.3$. The forward tracking coverage of the silicon tracker is exploited with a calorimetry-seeded tracking algorithm called “Phoenix”. It is similar to central tracking, where tracks found in the COT are projected into the silicon tracker and hits within a narrow road of the trajectory seed silicon track reconstruction. With the Phoenix algorithm, the track helix in the magnetic field is specified by the position of the $p\bar{p}$ collision vertex, the three-dimensional exit position of the electron into the PES, and a helix curvature. The curvature is derived from the E_T of the shower in the EM calorimeter. As the E_T provides no information on the particle charge, there are two potential helices, one for each charge. The algorithm projects each helix into the silicon tracker and seeds the silicon track reconstruction. If both projections yield tracks, the higher quality one is selected. The COT is not directly used, but tracks found by the trackers are used to reconstruct the location of the $p\bar{p}$ collision vertex.

The radial extent of the PES, relative to the beamline, is 12–129 cm. Depending on the track vertex location along the beamline (Z_{vtx}), a track traverses from 0 to 8 layers of silicon. A Phoenix track is required to have at least three silicon hits. Only plug electrons associated to tracks that traversed at least three silicon layers are accepted. Eighty percent of the tracks traverse four or more silicon layers. Within the plug region, the average Phoenix track acceptance is 94% and within this acceptance zone, the track-finding efficiency is 91%.

The Phoenix algorithm is efficient and results in low background. While the pointing resolution of a Phoenix track is good (1 mrad or better), its path length in the magnetic field at large $|\eta_{\text{det}}|$ is small and the helix curvature resolution is poor. Consequently, there is neither a P_T nor E/P requirement for plug electron identification.

The central region tracking algorithm utilizes hits in the silicon tracker if available. However, the plug Phoenix tracking algorithm requires a fully functional silicon tracker. This silicon requirement reduces the effective integrated luminosity of CP and PP topology dielectrons relative to CC dielectrons by 6%.




FIG. 1. The raw ee -pair P_T distribution for all dielectron topologies combined. No corrections or background subtractions are applied. The highest P_T is 327 GeV/ c .

392

C. Dielectron Selection

Events are required to have a reconstructed dielectron pair mass $66 < M_{ee} < 116$ GeV/ c^2 . For dielectrons of the CC topology, the two tracks are required to have opposite charge. However, for CP and PP topology dielectrons, there is no opposite charge requirement because of the significant charge misidentification on Phoenix tracks at large $|\eta_{\text{det}}|$.

The efficiency for the trigger to select events is typically over 99% for dielectrons that pass offline event selections. The CENTRAL-18 trigger has an inefficiency of 3% per single central electron due to track association requirements. The Z -NO-TRACK trigger is on average less than 0.5% inefficient for all topologies, and complements the CENTRAL-18 trigger.

401

D. Measurement Event Sample

The numbers of events passing all previously described selections in the CC, CP, and PP dielectron topologies are 51 951, 63 752, and 22 469, respectively. Figure 1 shows the raw ee -pair P_T distribution for these events. The backgrounds are small, and are from QCD or from WW , WZ , ZZ , $t\bar{t}$, W + jets, and $Z \rightarrow \tau^+\tau^-$ sources with real high- E_T electrons. The QCD background is primarily from dijets where a track in a jet fakes an electron or is an electron from a photon conversion. The high- E_T electron sources have at least one real electron. The second electron is either a real second electron or a fake one such as in W + jets.

Overall, the background from QCD and non-Drell-Yan high- E_T electrons is 0.5%. It is negligible at low pair P_T , and for $P_T > 100$ GeV/ c , it reaches the 5% level. These backgrounds are subtracted from the P_T distribution shown in Fig. 1 for the measurement of $\Delta\sigma/\Delta P_T$. Backgrounds are significantly reduced, particularly at large P_T , by requiring each electron candidate to have an associated track.

The overall QCD background level is 0.3%, and it is under 1% at all P_T . It is estimated with the data used for the P_T measurement using an “isolation extrapolation” procedure. All selection criteria are applied to both electron candidates except the isolation energy (E_{iso}) requirement on one electron candidate. Its E_{iso} distribution has a sharp peak at low E_{iso} from Drell-Yan electrons (the signal) and a broad, flat distribution extending to very large E_{iso} from QCD sources (the background). The E_{iso} distribution is fit to a signal plus a background component over the full E_{iso} range, and the background component is extrapolated into the signal region for the QCD background estimate. The signal and background shapes are derived from the unbiased data set used in the measurement, and with selections close to the electron selections to avoid biases. For the background shape event selection, two electron-like candidates are required, but one is selected to be “jet-like” by reversing the selection requirement on its E_{iso} and $E_{\text{HAD}}/E_{\text{EM}}$ parameters. The other, whose E_{iso} distribution is the background shape, has all electron selection requirements except E_{iso} applied.

The high E_T electron backgrounds from WW , WZ , ZZ , $t\bar{t}$, W + jets, and $Z \rightarrow \tau^+\tau^-$ are derived from the simulated samples. The overall background level from these sources is 0.2%, but they are the source of the 5% backgrounds for $P_T > 100$ GeV/ c .

427 Above the P_T of 150 GeV/c, there are 55 events. The ee -pair mass distribution has a clear Z -boson mass peak,
 428 and within the 66–116 GeV/ c^2 mass range, there is no indication of unexpected backgrounds. The peak location and
 429 width are consistent with expectations.

430 V. DATA SIMULATION

431 The acceptance for Drell–Yan dilepton pairs is obtained using the Monte Carlo physics event generator, PYTHIA
 432 6.214 [26], and the CDF event and detector simulations. PYTHIA generates the hard, leading order (LO) QCD
 433 interaction, $q + \bar{q} \rightarrow \gamma^*/Z$, simulates initial state QCD radiation via its parton shower algorithms, and generates
 434 the decay, $\gamma^*/Z \rightarrow l^+l^-$. The CTEQ5L [27] nucleon parton distribution functions (PDFs) are used in the QCD
 435 calculations. The underlying event and γ^*/Z boson P_T parameters are PYTHIA tune AW (i.e., PYTUNE 101, which is
 436 a tuning to previous CDF data) [26, 28].

437 Generated events are processed by the CDF event and detector simulation. The event simulation includes PHOTOS
 438 2.0 [29] which adds final-state quantum electrodynamics (QED) radiation to decay vertices with charged particles, e.g.
 439 $\gamma^*/Z \rightarrow e^+e^-$. The time-dependent beam and detector conditions for data runs recorded and used for physics analyses
 440 are simulated. The beam conditions simulated are the p and \bar{p} beamline parameters, the $p\bar{p}$ luminous region profile,
 441 and the instantaneous and integrated luminosities per run. The detector conditions simulated are detector component
 442 calibrations, which include channel gains and malfunctions. Thus, the simulated events parallel the recorded data,
 443 and are reconstructed, selected, and analyzed as the data.

444 The $\Delta\sigma/\Delta P_T$ measurement is data driven and depends on the correct modeling of both the physics and the detector.
 445 The procedure involves the measurement and tuning of the underlying kinematics and detector parameters that make
 446 the simulated, reconstructed event distributions match the actual data as precisely as possible. This is a bootstrap
 447 process that iterates if necessary for the required precision. The default simulation does not reproduce the data at
 448 the precision required. The following subsections describe the model tunings.

449 A. Physics Simulation

The Drell–Yan dilepton production is described by

$$\frac{d^4\sigma}{dM^2 dy dP_T d\Omega} = \frac{d^3\sigma}{dM^2 dy dP_T} \frac{dN}{d\Omega},$$

450 where $d^3\sigma/dM^2 dy dP_T$ is the unpolarized γ^*/Z boson production cross section at the resonance mass M with subse-
 451 quent decay to e^+e^- , and $dN/d\Omega$ the electron angular distribution of the $\gamma^*/Z \rightarrow e^+e^-$ decay in a boson rest frame.
 452 For this measurement, the single differential distributions $d\sigma/dy$ and $d\sigma/dP_T$, and the electron angular distribution,
 453 are tuned to the data. The y distribution tuning for γ^*/Z production is from the $d\sigma/dy$ measurement [14]. The
 454 tuning of the electron angular distribution is briefly reviewed next. The tuning of $d\sigma/dP_T$ is specific to this analysis,
 455 and is presented last.

456 The PYTHIA parton showering starts with the $q\bar{q} \rightarrow \gamma^*/Z$ annihilation vertex at the end of the shower chain then
 457 evolves the shower backwards in time to an initiating $q\bar{q}$ or qg state. The Compton production process cannot be
 458 fully simulated. While its gluon splitting subprocess is simulated, the gluon fusion subprocess, $qg \rightarrow q^* \rightarrow q + \gamma^*/Z$,
 459 cannot be simulated from the annihilation vertex at the end of the shower chain. The gluon fusion production rate is
 460 compensated in the shower, but there is no compensation to the boson polarization states affected by this subprocess.
 461 The boson polarization affects the decay electron angular distribution.

462 The decay electron angular distribution is analyzed in the Collins–Soper (CS) rest frame [30] of the e^+e^- pair.
 463 The CS frame is reached from the laboratory frame via a Lorentz boost along the lab z -axis into a frame where the
 464 z -component of the pair momentum is zero, followed by a boost along the P_T of the pair. At $P_T = 0$, the CS and
 465 laboratory coordinate frames are the same. Within the CS frame, the z -axis for the polar angle is the angular bisector
 466 between the proton direction and the negative of the anti-proton direction. The x -axis is the direction of the P_T . The
 467 polar and azimuthal angles of the e^- in the rest frame are denoted as ϑ and φ , respectively.

468 The general structure of the Drell–Yan decay lepton angular distribution in a boson rest frame consists of nine




FIG. 2. The P_T correction function applied to the generator level $\Delta N/\Delta P_T$ distribution that makes flat the ratio of the observed data to the simulated data. The points are at the center of the P_T bins. For the low-statistics $P_T > 120$ GeV/c region, an average correction is used.

469 helicity cross sections governed by the polarization state of the vector boson [31],

$$\begin{aligned} \frac{16\pi}{3} \frac{dN}{d\Omega} = & (1 + \cos^2 \vartheta) + \\ & A_0 \frac{1}{2} (1 - 3 \cos^2 \vartheta) + \\ & A_1 \sin 2\vartheta \cos \varphi + \\ & A_2 \frac{1}{2} \sin^2 \vartheta \cos 2\varphi + \\ & A_3 \sin \vartheta \cos \varphi + \\ & A_4 \cos \vartheta + \\ & A_5 \sin^2 \vartheta \sin 2\varphi + \\ & A_6 \sin 2\vartheta \sin \varphi + \\ & A_7 \sin \vartheta \sin \varphi . \end{aligned}$$

470 The A_{0-7} coefficients are cross section ratios, and are functions of the boson kinematics. They are zero at $P_T = 0$,
 471 except for the electroweak part of A_4 responsible for the forward-backward e^- asymmetry in $\cos \vartheta$. The A_{5-7}
 472 coefficients appear at $\mathcal{O}(\alpha_s^2)$ and are small in the CS frame. The decay-electron angular-distribution analysis [15] in
 473 the CS frame measures the large and accessible decay electron angular coefficients, A_0 , A_2 , A_3 , and A_4 , as functions
 474 of P_T . These measurements are incorporated into the modeling of $\gamma^*/Z \rightarrow e^+e^-$ decays.

475 The generator-level P_T distribution is adjusted, bin by bin, so that the shape of the reconstruction-level, simulated
 476 P_T distribution is the same as in the data. The method uses the data-to-simulation ratio of the number of reconstructed
 477 events in P_T bins as an iterative adjustment estimator for the generator level P_T bins. Successive iterations unfold
 478 the smearing of events across P_T bins. Figure 2 is the generator-level P_T correction function that makes the data-
 480 to-simulation ratio uniform. Statistical fluctuations in the ratio are smoothed out. The $\Delta N/\Delta P_T$ correction is the
 481 measurement of the shape of $d\sigma/dP_T$ that is used in the physics model.

482 B. Detector Simulation

483 The simulation is used to calculate the combined detector acceptance (A) and selection efficiency (ϵ) as a function
 484 of kinematic variables for Drell–Yan dielectrons. The combined acceptance and efficiency convolution is denoted as
 485 $A \otimes \epsilon$. Single-electron selection efficiencies are measured and incorporated into the simulation as event-weight scale
 486 factors. The scale factors are ratios of the measured efficiencies of the data to the simulated data.




FIG. 3. The overall CC topology central electron E_T distribution. The crosses are the data and the histogram is the simulated data.

487 The electron-trigger efficiencies have an E_T (calorimetry) and η_{det} (tracking) dependence that are measured and
 488 incorporated into the simulation. The electron-identification efficiencies are measured as a function of η_{det} for both
 489 central and plug region electrons. Plug region efficiencies are measured separately for CP and PP topology dielec-
 490 trons due to their different environments. Plug-electron efficiencies have a clear time dependence due to the increasing
 491 instantaneous luminosities delivered by the Tevatron. This dependence is incorporated into the simulation. Lumi-
 492 nosity effects are measured using the number of $p\bar{p}$ vertices reconstructed by the trackers per event.

A precise model of the calorimeter response in the simulation is important for the calculation of $A \otimes \epsilon$. Electron kinematics are derived from a three-momentum that uses the electron energy measured in the calorimeters for the momentum magnitude and the associated track for the direction. The simulated electron energy scale calibration and resolution versus η_{det} are tuned using the electron E_T distribution. The default scale and resolution per η_{det} bin are adjusted so that the electron E_T distribution reconstructed in simulation matches that of the data. Only the constant term in the energy resolution is adjusted. Since the default simulation parametrization of the energy resolution can already have a constant term, the resolution adjustment is done with an additional constant term c_2 ,

$$\frac{\sigma}{E} = \frac{c_0}{\sqrt{E}} \oplus c_1 \oplus c_2,$$

493 where σ is the energy resolution, E is the energy, c_0 and c_1 are the default parameters of Section III, and the \oplus denotes
 494 combination in quadrature. The tuned values of c_2 on average are 0.9% and 2.3% for the central and plug calorimeters,
 495 respectively. The steeply rising and falling parts of the electron E_T distribution dominate the constraints. The three
 496 dielectron topologies, CC, CP, and PP, provide multiple and independent central and plug electron E_T samples. The
 497 η_{det} -dependent E_T distributions of each topology are calibrated independently. After the η_{det} -dependent parameters
 498 are determined, the separate CC, CP, and PP dielectron mass distributions are used to set an overall global scale and
 499 resolution adjustment for central and plug electrons.

500 The simulation is compared to data using histogrammed electron E_T and ee -pair mass distributions. Since the
 501 backgrounds are small, they are ignored. The comparison statistic is the χ^2 between the simulation and data. The
 502 event count of the simulated data is normalized to that of the data, and only statistical uncertainties are used in the
 503 calculation.

504 The η_{det} -dependent calorimeter response tunings provide a good match between the simulated-data and data.
 505 Figure 3 shows the E_T distribution of CC-topology central electrons. The corresponding plot for PP-topology plug
 506 electrons is similar in shape except that the width of the E_T “peak” is slightly narrower. Figure 4 shows the E_T
 507 distribution of CP-topology plug electrons. The corresponding plot for CP-topology central electrons is very similar.
 508 A χ^2 test is used to evaluate the compatibility between the simulation and data. For CC-central, CP-central, CP-plug,
 509 and PP-plug electrons, the χ^2 values are 117, 100, 87, and 135, respectively, for 100 bins (90 bins for PP).

513 Figure 5 shows the CC-topology ee -pair mass distribution. The ee -pair mass distributions for the CP and PP
 514 topologies are similar. The simulated-data to data χ^2 for the CC-, CP-, and PP-topology ee -pair mass distributions
 515 are 107, 123, and 114, respectively, for 100 bins. The sharp and narrow Z -peaks provide significant constraints on the
 516 the global energy scale and resolution parameters.




FIG. 4. The overall CP-topology plug electron E_T distribution. The crosses are the data and the histogram is the simulated data.




FIG. 5. The overall CC topology ee -pair mass distribution. The crosses are the data and the histogram is the simulated data.

517

VI. THE CROSS SECTION

The differential cross section in P_T is the average cross section in a P_T bin, or $\Delta\sigma/\Delta P_T$, where $\Delta\sigma$ is the integrated cross section in a bin. The $\Delta\sigma$ is defined as:

$$\Delta\sigma = \frac{N}{\mathcal{L} A \otimes \epsilon}$$

518 where N is the background subtracted event count, \mathcal{L} is the effective integrated luminosity, and $A \otimes \epsilon$ is the combined
 519 acceptance and efficiency. The effective luminosity, \mathcal{L} , is 2057 pb^{-1} , and it includes the acceptance of the $|Z_{\text{vtx}}| <$
 520 60 cm fiducial restriction. The details of the measurement and its uncertainties are presented next.

521

A. Acceptance and Efficiency Unfolding

522 The combined acceptance and efficiency, $A \otimes \epsilon$, is calculated using the simulation to convolve individual electron
 523 η_{det} acceptances and efficiencies into an ee -pair P_T quantity. The value of $A \otimes \epsilon$ ranges from 0.22 at $P_T \simeq 0.2 \text{ GeV}/c$
 524 to 0.30 at $P_T \simeq 200 \text{ GeV}/c$. As P_T increases, the ee -pair rapidity becomes more central, the electron E_T becomes
 525 larger, and the acceptance slowly increases.

526 The smearing of the observed P_T away from the generator (γ^*/Z) level value is significant relative to the bin size
 527 at low P_T : It has an rms width of about $2.2 \text{ GeV}/c$ and is non-Gaussian. Detector resolution and QED radiation
 528 from the $\gamma^*/Z \rightarrow e^+e^-$ vertex induce distortions to the reconstructed ee -pair mass and P_T distributions. In addition,
 529 they induce a broad enhancement in the $A \otimes \epsilon$ function. It rises from 0.22 at $P_T \simeq 0.2 \text{ GeV}/c$ to a broad maximum

530 of 0.28 around a P_T of 8 GeV/c, then decreases to 0.24 at $P_T \sim 30$ GeV/c before increasing again at larger P_T due
 531 to the increased acceptance.

532 When $A \otimes \epsilon$ is used to calculate cross sections, it unfolds the effects of smearing. The $(A \otimes \epsilon)^{-1}$ correction is applied
 533 bin-by-bin and consists logically of two steps. The first step is a scaling correction on the number of reconstructed and
 534 selected events. This scales (unfolds) the number of events reconstructed in a P_T bin into the number of reconstructed
 535 events produced in the bin. The simulation provides an average scaling factor. The second step is a standard detector
 536 acceptance correction on this scaled (unfolded) event count.

537 For the cross section uncertainty evaluation, more information on event production and migration among the P_T
 538 bins is required. The number of events produced in each bin have statistical fluctuations. With smearing, there
 539 is event migration among the bins, and this migration is also subject to statistical fluctuations. At low P_T , event
 540 migration between bins is large. If these migrations are unaccounted, the cross section uncertainty will be significantly
 541 underestimated. As the event migration between bins is not measured, these migrations are estimated with the
 542 simulation.

543

B. Unfolding Uncertainty Model

544 Comparisons of fully-corrected cross section measurements with theoretical cross sections are not straightforward.
 545 Where detector smearing is significant, there are significant uncertainty correlations among the P_T bins due to the
 546 event migrations among the bins. The simulation behind the scaling correction accounts for these migrations. The
 547 scaling correction uncertainty has both statistical and systematic biases. The systematic bias is from the residual
 548 simulation model bias on the bin scaling factor. This bias has been mitigated by the model tuning described in
 549 Section V B. The sources of statistical uncertainty from event migration for the scaling correction are discussed,
 550 and a model of per-measurement (per-single-experiment) fluctuations for the uncertainty that uses the simulation is
 551 specified.

552 Within the context of the simulation, information about the event migration of reconstructed events among P_T
 553 bins is in its transfer matrix, \bar{n}_{lk} , where \bar{n}_{lk} is the expectation value of the number of events produced in bin k that
 554 migrate into bin l . The expectation value of a quantity is denoted with an overbar, e.g. \bar{n} . All expectation values
 555 are normalized to the integrated luminosity of the data. The number of events that do not migrate out of a bin is
 556 denoted by \bar{n}_g . The number of events that migrate out and in are denoted by \bar{n}_o and \bar{n}_i , respectively. In terms of the
 557 transfer matrix, \bar{n}_{lk} , the \bar{n}_g , \bar{n}_o , and \bar{n}_i for P_T bin m are, respectively, \bar{n}_{mm} , the sum of \bar{n}_{lm} over the migration index
 558 l excluding bin m , and the sum of \bar{n}_{mk} over the production index k excluding bin m . The per-measurement statistical
 559 fluctuation of a quantity from its expectation value is denoted by δ followed by the quantity, e.g., $\delta n = n - \bar{n}$. An
 560 ensemble variance is denoted by δ^2 , e.g., for Poisson statistics, $\delta^2 n = \bar{n}$, and if c is a constant, $\delta^2 cn = c^2 \bar{n}$.

561 The scaling correction factor is $\bar{\rho} \equiv \bar{N}_g / \bar{N}_r$, where $\bar{N}_g = \bar{n}_g + \bar{n}_o$ is the expectation on the number of events
 562 produced in a bin, and $\bar{N}_r = \bar{n}_g + \bar{n}_i$ is the number of events reconstructed in a bin. Any residual model systematic
 563 bias is in $\bar{\rho}$. For a given measurement, the number of events produced and reconstructed in a bin are $N_g = n_g + n_o$
 564 and $N_r = n_g + n_i$, respectively. The scaling correction estimate for N_g is $\bar{\rho} N_r$. The difference between the scaling
 565 correction estimator $\bar{\rho} N_r$ and its target N_g gives a bias between them, $B = \bar{\rho} N_r - N_g$. If there are no target
 566 fluctuations ($N_g = \bar{N}_g$), B is the statistical bias of the estimator. With target fluctuations, there are two statistical
 567 biases, $\bar{\rho} N_r - \bar{\rho} \bar{N}_r (= \delta \bar{\rho} N_r)$ and $N_g - \bar{N}_g (= \delta N_g)$, and B is their difference.

568 With no smearing, the estimator and target, along with their fluctuations, are identical, so $B = 0$ and the statistical
 569 uncertainty of the scaling correction is just that of the estimator. With smearing, the estimator and target fluctuations
 570 are not fully correlated, so $B \neq 0$ and the scaling correction statistical uncertainty is from a combination of estimator
 571 and target statistical fluctuations. The estimator ($\bar{\rho} N_r$) and target (N_g) have three statistically independent elements:
 572 n_g , n_o , and n_i . As n_g is part of both the estimator and target, the common overlap must be removed to avoid double
 573 counting. The total per-measurement fluctuation for the scaling correction, denoted as $\delta N'_g$, is defined as the sum of
 574 fluctuations ($\delta n = n - \bar{n}$) from the estimator and the target minus their common term, δn_g :

$$\begin{aligned} \delta N'_g &= \delta \bar{\rho} N_r + \delta N_g - \delta n_g \\ &= \delta \bar{\rho} (n_g + n_i) + \delta (n_g + n_o) - \delta n_g \\ &= \delta \bar{\rho} (n_g + n_i) + \delta n_o . \end{aligned}$$

575 For Poisson statistics, the P_T bin ensemble variance is:

$$\begin{aligned} \delta^2 N'_g &= \bar{\rho}^2 (\bar{n}_g + \bar{n}_i) + \bar{n}_o \\ &= \bar{\rho} \bar{N}_g + \bar{n}_o , \end{aligned}$$




FIG. 6. $R_g(P_T)$ and $\bar{\rho}(P_T)$. The solid histogram is R_g , the bin variance of the uncertainty model relative to the variance of the produced events. The abrupt drops are where the bin size changes. The lower, dashed histogram is $\bar{\rho}$.

576 where $\bar{N}_g = \bar{\rho}\bar{N}_r = \bar{\rho}(\bar{n}_g + \bar{n}_i)$ is used. The covariance from the δn_o and $\delta \bar{\rho} n_i$ terms between bins k and l is
 577 $\bar{\rho}_k \bar{n}_{kl} + \bar{\rho}_l \bar{n}_{lk}$.

578 The ratio, R_g , of $\delta^2 N'_g$ to $\delta^2 N_g (= \bar{N}_g)$ is the variance of the model relative to the variance of only the produced
 580 events. Figure 6 shows both the ratio and the scaling correction factor as functions of P_T . In the low P_T bins, \bar{n}_o and
 581 \bar{n}_i are separately much larger than \bar{n}_g . Their effects are significant as $R_g = \bar{\rho} + \bar{n}_o/\bar{N}_g$.

582 For the uncertainty evaluations, the cross section is rewritten as $\Delta\sigma = \bar{\rho}N_r/(\mathcal{L} A')$, where $A' \equiv \bar{\rho} A \otimes \epsilon$. The
 583 uncertainty on \mathcal{L} is systematic and is considered separately. Thus, the fractional uncertainty on $\Delta\sigma$ is a combination
 584 of the fractional uncertainty of $\bar{\rho}N_r$ and A' . The fractional uncertainty of $\bar{\rho}N_r$ is defined as the uncertainty of $\bar{\rho}N_r$
 585 from the model ($\delta N'_g$) divided by $\bar{\rho}N_r (= \bar{N}_g)$. The correlation of these fractional uncertainties between P_T bins l
 586 and k is given by the fractional covariance matrix: $\bar{V}_{lk}/(\bar{N}_{gl} \bar{N}_{gk})$, where \bar{V}_{lk} is the covariance matrix of the model,
 587 and \bar{N}_{gl} and \bar{N}_{gk} are the \bar{N}_g of bin l and k , respectively. The small acceptance fractional uncertainties are added in
 588 quadrature to the diagonal part of the fractional covariance matrix. The measured cross sections are used to convert
 589 the unitless fractional matrix into units of cross section squared, and this matrix is used to propagate uncertainties
 590 for the total cross section measurement and for the comparison of a prediction with the measured cross section.

591

C. Systematic Uncertainties

592 The largest source of uncertainty is the effective integrated luminosity, \mathcal{L} . It has an overall uncertainty of 5.8% that
 593 consists of a 4% uncertainty of the acceptance of the gas Cherenkov luminosity detector [24] to $p\bar{p}$ inelastic collisions and
 594 a 4.2% measurement uncertainty. It is common to all P_T bins and not explicitly included. The acceptance uncertainty
 595 is primarily from the uncertainty in the beamline and detector geometry (material), and from the uncertainty in the
 596 model of the inelastic cross section. The inelastic cross section model contributes 2% to the acceptance uncertainty.
 597 The measurement uncertainty contains the uncertainty of the absolute $p\bar{p}$ inelastic cross section.

598 The uncertainty on $A \otimes \epsilon$ has a component from the input electron efficiency measurements which depend on η_{det}
 599 and instantaneous luminosity. The simulation is used to propagate these electron measurement uncertainties into an
 600 uncertainty for the ee -pair P_T and to include correlations of the same measurements. The calculated uncertainty is
 601 uniform and amounts to about 1% over $0 < P_T < 20$ GeV/ c . It slowly decreases at higher P_T . A large fraction of the
 602 uncertainty is due to plug electron measurement uncertainties. The fractional uncertainty decreases with P_T because
 603 the fraction of plug events decreases. Because the same measurements are used on all P_T bins, the uncertainty is
 604 treated as fully correlated across bins.

605 The calorimeter response modeling uncertainty analysis is limited by the statistical precision of the simulated
 606 data. At the peak of the P_T distribution, the statistical uncertainty is 0.3%. The variations on the central and plug
 607 calorimeter global energy scale and resolutions tunings allowed by the data propagate into changes of $A \otimes \epsilon$ that are
 608 no larger than its statistical uncertainty. These changes are not independent.




FIG. 7. The $\Delta\sigma/\Delta P_T$ cross section versus P_T . Cross section values are plotted at the bin center. The horizontal bars represent the bin extent and the vertical bars are the cross section uncertainties. The solid (black) crosses are the data and all uncertainties except the integrated luminosity uncertainty are combined and plotted. The solid (red) histogram is the RESBOS calculation. The dot-dash (blue) bars of the $P_T > 25$ GeV/c region are the FEWZ2 calculation. For the calculations, only numerical uncertainties are included but they are too small to be visible. The inset is the $P_T < 25$ GeV/c region with a linear ordinate scale.

609

D. Results

The Drell-Yan $\Delta\sigma/\Delta P_T$ for e^+e^- pairs in the Z -boson mass region of $66 - 116$ GeV/ c^2 is shown in Fig. 7 and tabulated in Table I. The total cross section from the numerical integration of the cross section in each P_T bin is $256.1 \pm 1.3 \pm 2.6$ pb, where the first uncertainty is statistical and the second is the systematic uncertainty due to electron efficiency measurements. The 5.8% integrated luminosity uncertainty of 14.9 pb is not included.

Figure 7 shows that the RESBOS prediction has a general agreement with the data over the full range of P_T . The RESBOS total cross section from the numerical integration of its cross section in each P_T bin is 254 pb. Figure 8 shows the ratio of the measured cross section to the RESBOS prediction in the lower P_T region.

The detector smearing correlates neighboring P_T bin uncertainties that are estimated with the model specified in Section VI A. For the low P_T bins, the correlations spread across many bins but for $P_T > 40$ GeV/ c , the correlations are predominantly between nearest neighbors. The cross section covariance matrix eigenvalues and eigenvectors are used for the χ^2 comparison between the data and RESBOS. The eigenvalues are the measurement uncertainties (variances) of the associated eigenvector. Measurement uncertainties between eigenvectors are uncorrelated. As an eigenvector corresponds to many P_T bins because of smearing, its most probable P_T bin is used for its association to a P_T bin. The mapping of eigenvectors to P_T bins is described next.

The P_T bins are numbered consecutively, 0–81 (lowest to highest P_T), and the bin number is denoted by n . The bin-number expectation values of the eigenvectors are used for their assignment to P_T bins. The eigenvector with the lowest expectation value is assigned to P_T bin 0, the next lowest to P_T bin 1, and so on. For $P_T < 25$ GeV/ c , the rms width of the expectation value is about 4 bins, and above it, about 1 bin or less. In the 13–18 GeV/ c region, the

TABLE I. The $\Delta\sigma/\Delta P_T$ cross section versus P_T . The first uncertainty is the statistical uncertainty. The second uncertainty is the efficiency measurement systematic uncertainty, which is 100% correlated across all bins. The 5.8% luminosity uncertainty applies to all bins but is not included.

P_T bin GeV/c	$\Delta\sigma/\Delta P_T$ pb / GeV/c	P_T bin GeV/c	$\Delta\sigma/\Delta P_T$ pb / GeV/c
0.0–0.5	$(3.613 \pm 0.168 \pm 0.035) \times 10^0$	20.5–21.0	$(2.923 \pm 0.143 \pm 0.030) \times 10^0$
0.5–1.0	$(1.008 \pm 0.027 \pm 0.010) \times 10^1$	21.0–21.5	$(2.877 \pm 0.144 \pm 0.030) \times 10^0$
1.0–1.5	$(1.551 \pm 0.033 \pm 0.015) \times 10^1$	21.5–22.0	$(2.603 \pm 0.134 \pm 0.027) \times 10^0$
1.5–2.0	$(1.947 \pm 0.037 \pm 0.019) \times 10^1$	22.0–22.5	$(2.624 \pm 0.139 \pm 0.027) \times 10^0$
2.0–2.5	$(2.158 \pm 0.039 \pm 0.021) \times 10^1$	22.5–23.0	$(2.590 \pm 0.140 \pm 0.026) \times 10^0$
2.5–3.0	$(2.295 \pm 0.040 \pm 0.023) \times 10^1$	23.0–23.5	$(2.516 \pm 0.139 \pm 0.026) \times 10^0$
3.0–3.5	$(2.258 \pm 0.039 \pm 0.022) \times 10^1$	23.5–24.0	$(2.200 \pm 0.124 \pm 0.022) \times 10^0$
3.5–4.0	$(2.235 \pm 0.039 \pm 0.022) \times 10^1$	24.0–24.5	$(1.948 \pm 0.113 \pm 0.020) \times 10^0$
4.0–4.5	$(2.061 \pm 0.037 \pm 0.021) \times 10^1$	24.5–25.0	$(2.179 \pm 0.129 \pm 0.022) \times 10^0$
4.5–5.0	$(1.987 \pm 0.036 \pm 0.020) \times 10^1$	25.0–26.0	$(2.032 \pm 0.085 \pm 0.021) \times 10^0$
5.0–5.5	$(1.876 \pm 0.035 \pm 0.019) \times 10^1$	26.0–27.0	$(1.736 \pm 0.076 \pm 0.018) \times 10^0$
5.5–6.0	$(1.729 \pm 0.034 \pm 0.017) \times 10^1$	27.0–28.0	$(1.633 \pm 0.075 \pm 0.016) \times 10^0$
6.0–6.5	$(1.563 \pm 0.032 \pm 0.016) \times 10^1$	28.0–29.0	$(1.616 \pm 0.077 \pm 0.016) \times 10^0$
6.5–7.0	$(1.468 \pm 0.031 \pm 0.015) \times 10^1$	29.0–30.0	$(1.381 \pm 0.069 \pm 0.014) \times 10^0$
7.0–7.5	$(1.374 \pm 0.030 \pm 0.014) \times 10^1$	30.0–32.0	$(1.284 \pm 0.045 \pm 0.013) \times 10^0$
7.5–8.0	$(1.307 \pm 0.030 \pm 0.013) \times 10^1$	32.0–34.0	$(1.005 \pm 0.039 \pm 0.010) \times 10^0$
8.0–8.5	$(1.183 \pm 0.028 \pm 0.012) \times 10^1$	34.0–36.0	$(8.769 \pm 0.361 \pm 0.088) \times 10^{-1}$
8.5–9.0	$(1.112 \pm 0.027 \pm 0.011) \times 10^1$	36.0–38.0	$(7.959 \pm 0.352 \pm 0.079) \times 10^{-1}$
9.0–9.5	$(1.033 \pm 0.026 \pm 0.011) \times 10^1$	38.0–40.0	$(7.068 \pm 0.336 \pm 0.070) \times 10^{-1}$
9.5–10.0	$(1.024 \pm 0.027 \pm 0.011) \times 10^1$	40.0–44.0	$(5.605 \pm 0.193 \pm 0.055) \times 10^{-1}$
10.0–10.5	$(9.043 \pm 0.244 \pm 0.094) \times 10^0$	44.0–48.0	$(4.600 \pm 0.179 \pm 0.044) \times 10^{-1}$
10.5–11.0	$(8.295 \pm 0.231 \pm 0.084) \times 10^0$	48.0–52.0	$(3.552 \pm 0.156 \pm 0.033) \times 10^{-1}$
11.0–11.5	$(8.319 \pm 0.239 \pm 0.085) \times 10^0$	52.0–56.0	$(2.760 \pm 0.136 \pm 0.025) \times 10^{-1}$
11.5–12.0	$(7.780 \pm 0.229 \pm 0.079) \times 10^0$	56.0–60.0	$(2.311 \pm 0.128 \pm 0.020) \times 10^{-1}$
12.0–12.5	$(7.465 \pm 0.227 \pm 0.076) \times 10^0$	60.0–65.0	$(1.618 \pm 0.089 \pm 0.014) \times 10^{-1}$
12.5–13.0	$(6.839 \pm 0.215 \pm 0.069) \times 10^0$	65.0–70.0	$(1.343 \pm 0.084 \pm 0.011) \times 10^{-1}$
13.0–13.5	$(6.411 \pm 0.208 \pm 0.065) \times 10^0$	70.0–75.0	$(1.094 \pm 0.078 \pm 0.009) \times 10^{-1}$
13.5–14.0	$(6.220 \pm 0.208 \pm 0.064) \times 10^0$	75.0–80.0	$(8.415 \pm 0.678 \pm 0.068) \times 10^{-2}$
14.0–14.5	$(5.890 \pm 0.204 \pm 0.060) \times 10^0$	80.0–85.0	$(6.347 \pm 0.565 \pm 0.049) \times 10^{-2}$
14.5–15.0	$(5.363 \pm 0.190 \pm 0.055) \times 10^0$	85.0–90.0	$(4.982 \pm 0.504 \pm 0.038) \times 10^{-2}$
15.0–15.5	$(5.186 \pm 0.190 \pm 0.053) \times 10^0$	90.0–95.0	$(3.786 \pm 0.422 \pm 0.028) \times 10^{-2}$
15.5–16.0	$(4.792 \pm 0.181 \pm 0.049) \times 10^0$	95.0–100.0	$(2.988 \pm 0.389 \pm 0.023) \times 10^{-2}$
16.0–16.5	$(4.431 \pm 0.172 \pm 0.045) \times 10^0$	100.0–110.0	$(2.298 \pm 0.227 \pm 0.016) \times 10^{-2}$
16.5–17.0	$(4.149 \pm 0.165 \pm 0.042) \times 10^0$	110.0–120.0	$(1.449 \pm 0.178 \pm 0.010) \times 10^{-2}$
17.0–17.5	$(4.346 \pm 0.179 \pm 0.044) \times 10^0$	120.0–130.0	$(9.369 \pm 1.389 \pm 0.064) \times 10^{-3}$
17.5–18.0	$(3.931 \pm 0.166 \pm 0.040) \times 10^0$	130.0–140.0	$(8.395 \pm 1.496 \pm 0.055) \times 10^{-3}$
18.0–18.5	$(3.757 \pm 0.163 \pm 0.038) \times 10^0$	140.0–150.0	$(5.304 \pm 1.174 \pm 0.034) \times 10^{-3}$
18.5–19.0	$(3.753 \pm 0.167 \pm 0.038) \times 10^0$	150.0–175.0	$(1.861 \pm 0.331 \pm 0.012) \times 10^{-3}$
19.0–19.5	$(3.586 \pm 0.163 \pm 0.036) \times 10^0$	175.0–200.0	$(5.283 \pm 1.478 \pm 0.031) \times 10^{-4}$
19.5–20.0	$(3.303 \pm 0.154 \pm 0.034) \times 10^0$	200.0–250.0	$(2.838 \pm 1.019 \pm 0.019) \times 10^{-4}$
20.0–20.5	$(2.952 \pm 0.142 \pm 0.030) \times 10^0$	250.0–350.0	$(1.489 \pm 1.162 \pm 0.009) \times 10^{-4}$

⁶³¹ rms width is the largest, 5–6 bins.

⁶³² The χ^2 is calculated for the eigenvector associated with the P_T bin n . For reference, the uncorrelated χ^2 is also
⁶³³ calculated for the bin. The cumulative χ^2 from bin 0 to n inclusive is denoted as $\chi^2(n)$. The number of degrees of
⁶³⁴ freedom of $\chi^2(n)$ is n . A useful measure is $\chi^2(n) - n$: it is typically constant when the prediction is compatible with
⁶³⁵ the data and increases over regions with discrepancies.

⁶³⁶ Figure 9 shows the $\chi^2(n) - n$ of the the RESBOS prediction. For the correlated χ^2 , changes in $\chi^2(n) - n$ can only be
⁶³⁸ associated with a P_T region because of smearing. In the $P_T < 25$ GeV/c region (bins 0–49), there are small differences
⁶³⁹ but the data may allow further tuning of the RESBOS non-perturbative form factor that is important in this region.

⁶⁴⁰ In the $44 < P_T < 90$ GeV/c region of Fig. 8 (bins 61–70 of Fig. 9), the RESBOS prediction is systematically lower
⁶⁴¹ than the data. This region is where the resummed calculation must be matched to the fixed-order perturbative
⁶⁴² calculation. This region is where the data can also contribute to the RESBOS resummation phenomenology of the




FIG. 8. The ratio of the measured cross section to the RESBOS prediction in the $P_T < 130$ GeV/ c region. The RESBOS total cross section is normalized to the data. The inset is an expansion of the low P_T region.




FIG. 9. $\chi^2(n) - n$ versus P_T bin number of the RESBOS prediction. The solid (red) histogram includes bin correlations and the dashed (blue) histogram does not. Bins 0–49 cover the 0–25 GeV/ c region, bin 60 is 40 GeV/ c , bin 70 is 85 GeV/ c , and bin 80 is 200 GeV/ c .

643 Drell–Yan lepton pair P_T distribution at the Tevatron.

648 Figure 10 shows the ratio of the measured cross section to the FEWZ2 prediction. There is reasonable agreement
 646 with the data in the high P_T region where the RESBOS and FEWZ2 calculations are in agreement with each other. In
 647 P_T bins where the deviation of the FEWZ2 prediction from the measurement is significant, the difference provides a
 648 measure of the importance of higher order contributions above $\mathcal{O}(\alpha_s^2)$. The PDF uncertainties provided by FEWZ2
 649 are at the 3% to 4% level. The uncertainties from variations of the QCD factorization and renormalization scales
 650 (from the Z -boson mass) in the P_T regions of 25–30, 100–110, and 200–250 GeV/ c , are at the 7%, 5%, and 6%
 651 level, respectively. However, the accuracy of these scale uncertainties is unclear because of the two different scales
 652 (lepton-pair mass and transverse momentum) inherent in this QCD calculation.

653

VII. SUMMARY

654 The transverse momentum cross section of e^+e^- pairs in the Z -boson mass region of 66–116 GeV/ c^2 produced in $p\bar{p}$
 655 collisions is measured using 2.1 fb $^{-1}$ of Run II data collected by the Collider Detector at Fermilab. The measurement
 656 is data driven and corrected for the detector acceptance and smearing. The physics and detector models of the
 657 simulation used for the correction are tuned so that the simulation matches the data. The precision of the data
 658 and the measurement method require both the data and simulation to be well calibrated and understood. The
 659 measurement uncertainties are from a simulation-based model that quantifies the effects of event migration between
 660 measurement bins due to detector smearing.




FIG. 10. The ratio of the measured cross section to the FEWZ2 prediction in the $25 < P_T < 250$ GeV/ c region. The FEWZ2 cross section is not normalized to the data.

661 Comparisons of this measurement with current quantum chromodynamic $\mathcal{O}(\alpha_s^2)$ perturbative and all-orders gluon
 662 resummation calculations show reasonable agreement. The data is of sufficient precision for further refinements in
 663 the phenomenology of the Drell–Yan lepton pair transverse momentum distribution.

664

ACKNOWLEDGMENTS

665 We thank C.-P. Yuan for useful discussions and help on the CSS QCD resummation formalism and the RESBOS
 666 calculation. We thank F. Petriello for useful discussions and help on the FEWZ2 calculation. We thank the Fermilab
 667 staff and the technical staffs of the participating institutions for their vital contributions. This work was supported
 668 by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare;
 669 the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering
 670 Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science
 671 Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean
 672 World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities
 673 Council and the Royal Society, UK; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación,
 674 and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; the Academy of Finland; and the Australian
 675 Research Council (ARC).

-
- 676 [1] S. D. Drell and T.-M. Yan, Phys. Rev. Lett. **25**, 316 (1970).
677 [2] K. Melnikov and F. Petriello, Phys. Rev. D **74**, 114017 (2006).
678 [3] J. C. Collins, D. E. Soper, and G. Sterman, Nucl. Phys. **B250**, 199 (1985); J. C. Collins and D. E. Soper, *ibid.* **B193**,
679 381 (1981); **B197**, 446(E) (1982); **B213**, 545(E) (1983).
680 [4] C. Albajar *et al.* (UA1 Collaboration), Z. Phys. C **44**, 15 (1989).
681 [5] J. Alitti *et al.* (UA2 Collaboration), Z. Phys. C **47**, 523 (1990).
682 [6] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **66**, 2951 (1991).
683 [7] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **67**, 2937 (1991).
684 [8] B. Abbott *et al.* (D0 Collaboration), Phys. Rev. Lett. **80**, 5498 (1998).
685 [9] T. Affolder *et al.* (CDF Collaboration), Phys. Rev. Lett. **84**, 845 (2000).
686 [10] F. Landry, R. Brock, P. M. Nadolsky, and C.-P. Yuan, Phys. Rev. D **67**, 073016 (2003).
687 [11] V. M. Abazov *et al.* (D0 Collaboration), Phys. Lett. B **693**, 522 (2010); Phys. Rev. Lett. **106**, 122001 (2010).
688 [12] G. Aad *et al.* (ATLAS Collaboration), Phys. Lett. B **705**, 415 (2011).
689 [13] S. Chatrchyan *et al.* (CMS Collaboration), Phys. Rev. D **85**, 032002 (2012).
690 [14] T. Aaltonen *et al.* (CDF Collaboration), Phys. Lett. B **692**, 232 (2010).
691 [15] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **106**, 241801 (2011).
692 [16] G. A. Ladinsky and C.-P. Yuan, Phys. Rev. D **50**, 4239 (1994).
693 [17] C. Balázs and C.-P. Yuan, Phys. Rev. D **56**, 5558 (1997).
694 [18] A. Konychev and P. Nadolsky, Phys. Lett. B **633**, 710 (2006).
695 [19] R. Gavin, Y. Li, F. Petriello, and S. Quackenbush, Comput. Phys. Commun. **182**, 2388 (2011).
696 [20] A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt, Eur. Phys. J. C **63**, 189 (2009).
697 [21] G. 't Hooft, Nucl. Phys. **B61**, 455 (1973).
698 [22] P. M. Nadolsky *et al.* (CTEQ Collaboration), Phys. Rev. D **78**, 103004 (2008).
699 [23] A. Abulencia *et al.* (CDF Collaboration), J. Phys. G: Nucl. Part. Phys. **34**, 2457 (2007).
700 [24] D. Acosta *et al.*, Nucl. Instrum. Methods A **494**, 57 (2002).
701 [25] M. Albrow *et al.*, Nucl. Instrum. Methods A **480**, 524 (2002).
702 [26] T. Sjöstrand, P. Edén, L. Lönnblad, G. Miu, S. Mrenna, and E. Norrbin, Comput. Phys. Commun. **135**, 238 (2001).
703 [27] H. L. Lai *et al.* (CTEQ Collaboration), Eur. Phys. J. C **12**, 375 (2000).
704 [28] M. Albrow *et al.* (Tev4LHC QCD Working Group), arXiv:0610012v1 [hep-ph].
705 [29] E. Barberio and Z. Was, Computer Phys. Comm. **79**, 291 (1994); E. Barberio, B. van Eijk, and Z. Was, **66**, 115 (1991).
706 [30] J. C. Collins and D. E. Soper, Phys. Rev. D **16**, 2219 (1977).
707 [31] E. Mirkes, Nucl. Phys. **B387**, 3 (1992); E. Mirkes and J. Ohnemus, Phys. Rev. D **50**, 5692 (1994).