This is the accepted manuscript made available via CHORUS. The article has been published as:

G-flux, M5 instantons, and U(1) symmetries in F-theory

Joseph Marsano, Natalia Saulina, and Sakura Schäfer-Nameki

Phys. Rev. D **87**, 066007 — Published 22 March 2013

DOI: [10.1103/PhysRevD.87.066007](https://doi.org/10.1103/PhysRevD.87.066007)

On G -flux, M5 instantons, and $U(1)$ s in F-theory

Joseph Marsano¹, Natalia Saulina² and Sakura Schäfer-Nameki³

¹ *Enrico Fermi Institute, University of Chicago*
5640 S Ellis Avenue, Chicago, IL 60637 USA
marsano uchicago.edu

² *Perimeter Institute for Theoretical Physics*
31 Caroline St N., Waterloo, Ontario N2L 2Y5, Canada
saulina theory.caltech.edu

³ *Department of Mathematics, King's College, University of London*
The Strand, WC2R 2LS, London, UK
ss299 theory.caltech.edu

Abstract

Local aspects of singular F-theory compactifications for SUSY GUT model-building are fairly well understood in terms of Higgs bundles and their spectral data. Several global issues remain, however, including a description of G -fluxes, which are key to constructing chiral matter and stabilizing moduli, and the global realization of $U(1)$ symmetries that can forbid phenomenologically unfavorable couplings. In this paper, we sharpen our earlier proposal for describing G -fluxes and $U(1)$'s through a distinguished 'Tate' divisor. As an application, we give a general discussion of M5-instanton contributions in the presence of G -flux and exemplify this in a concrete example, where we comment on the ability of instanton induced superpotential couplings to stabilize Kähler moduli.

1 Introduction and Summary

While the rules for local¹ model-building in F-theory GUTs are becoming fairly well understood [1–6], any such model still carries intrinsic assumptions about the global completion. For starters, local models describe only a local notion of the fluxes required to generate chiral matter. Further, local models often rely on $U(1)$ s which have become massive through a Stückelberg mechanism [2, 6, 7] to solve important phenomenological problems. These $U(1)$ s are intrinsically global in nature, though, so their presence or absence depends on details of the global completion [8].

These two points have received some attention in the past [8–10] but they do not represent the only bulk physics to which local models must appeal. In many models [6, 7, 11–14] the presence of GUT-singlets must be assumed for a variety of reasons, whether it be to generate neutrino mass, break supersymmetry, or lift unwanted exotic particles. Crucial in all such models are the superpotential couplings involving the singlets, which are typically assumed to be generated by some bulk physics. Here, bulk physics usually means one or more M5-instantons. Of course M5-instantons are important for more than just singlet physics; the issue of moduli stabilization, which will involve M5-instantons in a crucial way, is lurking behind everything that local model-builders do.

Our objective in this paper is to make some progress toward understanding these global issues in order to build continually improving global completions of local models. We begin with a discussion of the two important constructs that underlie all global models: G -flux and $U(1)$ symmetries. This discussion extends and clarifies that of [10], where we introduced a formalism for dealing with these based on ‘spectral divisors’. While [10] focused on the connection to Heterotic and the ability to write chirality computations on a resolved Calabi-Yau directly in terms of intersections on the singular one, our aim here is to develop more properly the connection between the global data of $U(1)$ ’s and fluxes in the ‘spectral divisor’ formalism and the corresponding data of the local model. Since in [10] we only discussed $U(1)$ ’s in an example, we make them central to the present discussion. Our approach to $U(1)$ ’s is complementary to previous work on this topic [9] and gives a uniform way to describe G -fluxes that can be interpreted as $U(1)$ fluxes and those that cannot. The ‘spectral divisor’ formalism is motivated by Heterotic/F-theory duality [15–20] and is compatible with the spectral cover in that context whenever such a dual Heterotic description exists [10]. In a sharp sense that we describe, a ‘spectral divisor’ is a global extension of the Higgs bundle spectral cover that mimics the ‘cylinder’ of the Heterotic/F-theory ‘cylinder map’ [16]. The relation of Calabi-Yau data to spectral data of local models was first discussed in [1] and global applicability of certain spectral cover properties, such as the preservation of $U(1)$ ’s, was investigated in [9] from a different perspective using mirror symmetry.

We then turn to a discussion of M5-instantons where our aim is to study the conditions for which the couplings that they generate are nonzero. There is a vast literature on M5-instantons

¹We refer to “local” in this context as the complete description of the 8d SYM theory on 7-branes in terms of Higgs bundle data, which does include information on the local geometry and monodromies (also sometimes referred to as “semi-local”).

detailing many of these issues, including recent work focused on the connection to F-theory and the relation to D3-brane instantons in type IIB [21–28]. We pay particular attention to recasting the fermi zero mode computation in terms of cohomologies on divisors in the base B_3 of Y_4 and the interplay of M5-instantons with the G -fluxes that we construct with spectral divisors as in [10]. With this knowledge, we turn to geometries based on the threefold of [29], specify which divisors support M5s that generate nonzero couplings, and comment on the implications for moduli stabilization in those models. We also take this opportunity to present a cleaner description of the threefold in [29]. Before getting to the details, let us summarize our general approach to these issues as well as some of the results.

1.1 Approach to G -flux and $U(1)$ symmetries

The constructs of G -flux and $U(1)$ symmetries are not completely unrelated because each admits a similar geometric description. Consider, for instance, an elliptically fibered Calabi-Yau fourfold Y_4 with a surface of $SU(5)_{\text{GUT}}$ singularities. The G -fluxes that generate chiral matter are $(2, 2)$ -forms in Y_4 that have exactly 1 leg on the torus, which is to say that they can be thought of as (linear combinations of) holomorphic surfaces that are effectively orthogonal to all horizontal and vertical surfaces. If Y_4 were smooth, this condition would tell us that G has vanishing intersection with all simple surfaces obtained as pairwise intersections of divisors in Y_4 . Constructing G 's of this type is already rather difficult. Things become even more complicated, however, when the Calabi-Yau is singular since we are most interested in G 's that probe the singularity structure. The resolution $\tilde{Y}_4$ will contain holomorphic surfaces that do not sit inside the preimage of horizontal or vertical divisors under the blow-down map $P : \tilde{Y}_4 \rightarrow Y_4$. These include the matter surfaces that one obtains when a curve of singularities is resolved². Curves of singularities support matter fields that descend from wrapped M2-branes so we want to engineer G -fluxes that have nonzero integral over at least some of them.

The issue with $U(1)$ symmetries is similar. $U(1)$ s that couple to our charged matter fields, which themselves originate from wrapped M2-branes, come from the reduction of the M-theory 3-form C_3 on harmonic $(1, 1)$ -forms ω

$$C_3 = A_1 \wedge \omega. \quad (1.1)$$

When we compactify M-theory on Y_4 , this gives a 3-dimensional gauge field on $\mathbb{R}^{2,1}$. To ensure that this carries over to a 4-dimensional gauge field in the F-theory limit, where the volume of the elliptic fiber is scaled to zero, the $(1, 1)$ -form ω must be chosen to have 1 leg on the elliptic fiber. If we wish to think of ω as a divisor in Y_4 , then, this condition means that ω cannot be horizontal or vertical. If Y_4 were smooth, this would force ω to be trivial so once any nontriviality in ω is intrinsically tied to the singularities. This is to be expected because we anticipate that ω has nonzero integral over degenerate cycles. This is how 4-dimensional states from wrapped M2-branes manage to couple to $U(1)$ vector bosons in the first place.

²These also include surfaces that sit inside the divisors that one obtains from the resolution of surfaces of singularities.

In each of these cases we seek a simple geometric description, whether it be a holomorphic surface to specify G or a divisor to specify the $(1,1)$ -form ω that we need to get a $U(1)$. Unfortunately, these concepts must be refined when the geometry is singular and all of the nontrivial structure that we need is contained in that refinement. In this paper, as in previous work [10], we adopt the standard approach that the physics on a singular Calabi-Yau fourfold Y_4 should be specified by starting with a particular smooth resolution $\tilde{Y}_4$ and taking a singular limit. This is essentially what we do when assigning gauge groups and charged matter to ADE singularities anyway. We specify the physics by saying that we start with a resolution $\tilde{Y}_4$ in which the ADE singularities are resolved by small resolutions, i.e. growing 2-cycles (as opposed to, say, deforming). We can wrap M2-branes on the finite volume 2-cycles in $\tilde{Y}_4$ where their physics is clear and then we see that these states become massless in the singular limit. The physical D3-brane charge should also include the geometric contribution from the resolved geometry $\tilde{Y}_4$ as computed in several examples over the past couple of years [30, 31]. In the same way, we should understand backgrounds for C_3 , including G -fluxes and reductions of C_3 that yield $U(1)$ s, in the resolved geometry $\tilde{Y}_4$. G -flux of the type that we need to induce chirality should be describable as a holomorphic surface in $\tilde{Y}_4$ and $U(1)$ s should be describable as ordinary divisors in $\tilde{Y}_4$.

To describe G -flux and $U(1)$ symmetries, then, it is crucial to specify a resolution $\tilde{Y}_4$. We actually do not describe the resolution explicitly but rather, following ideas from Heterotic/F-theory duality, we use dP_9 -fibrations to do the job. The basic strategy is to take advantage of the fact that we are interested in small resolutions which should be describable entirely within the local geometry near a surface of singularities, S_{GUT} , which we typically assume to be of A_4 type. Because the resolution does not care about what happens beyond the local geometry, we can excise it from Y_4 and insert it into a different fourfold $\hat{Y}_4$ that has the structure of a dP_9 -fibration. This allows us to describe the vanishing cycles in terms of the homology of dP_9 and, in particular, to think about the singularity structure in terms of the Mori cone of the dP_9 fiber as one moves along the base S_{GUT} . The geometry of dP_9 tells us about the structure of small resolutions, which simply grow back the zero volume homology cycles that sit in the Mori cone, and allows us to specify holomorphic surfaces and divisors in $\hat{Y}_4$ that interact in the right way with resolved cycles. We can then restrict these to the local geometry and, in so doing, obtain noncompact surfaces and divisors that we can try to globally extend after putting everything back into Y_4 .

A procedure based on dP_9 is a natural way to extend the ideas of F-theory/Heterotic duality beyond the realm of K3-fibered Calabi-Yau fourfolds and is at the heart of the approach in [10]. In that work, we introduced the notion of a ‘spectral divisor’ as an object that generalizes the ‘cylinder’ of the cylinder map [16] to geometries without a Heterotic dual. A ‘spectral divisor’ is nothing more than a tool for understanding holomorphic divisors and surfaces in the resolved geometry in terms of objects that can be described directly in the singular one. We used ‘spectral divisors’ in [10] to rewrite the integrals of G over matter surfaces in terms of intersection theory on the singular 4-fold Y_4 ³. We further argued that $U(1)$ ’s are generated

³In order to do this, we introduced a notion of ‘dual matter surface’ which was particularly useful for understanding how the local worldvolume gauge theory computations are related to Heterotic ones when a

in 4-folds where a distinguished ‘spectral divisor’ splits into components and later gave this distinguished object the name ‘Tate divisor’ in [6]. In this work, we develop the spectral divisor formalism more thoroughly in a way that extends the relationship of local and global data beyond intersection computations. We see how the ‘Tate divisor’ leads to a natural emergence of the Higgs bundle spectral cover $\mathcal{C}_{Higgs}$, provides a direct way to map G -flux to bundle data of the local model, and clarifies the special role played by matter curves in $\mathcal{C}_{Higgs}$ in determining the sheaves to which charged matter fields couple. Along the way, we describe how $U(1)$ fluxes are captured by our formalism, comment on the D3-brane tadpole and counting of singlets, and formulate very precisely how the quantization condition of the local model is connected to Witten’s criterion for G -flux quantization in the bulk Calabi-Yau 4-fold $\tilde{Y}_4$ [39]. We also describe how ‘spectral divisors’ can be used to perform useful manipulations and computations involving objects in the resolved Calabi-Yau that are more general than in [10]. Using our methods, we provide a simple check on the quantization condition that doesn’t require explicit computation of the second Chern class of $\tilde{Y}_4$.

As we were finalizing this work [32] appeared, where small resolutions of singular Calabi-Yau fourfolds with an $SU(5)$ singularity are studied in general. There are several small resolutions, which are birationally equivalent, and it would be interesting to put our analysis of G -fluxes and $U(1)$ s in this context.

1.2 M5-instantons

In the second part of this paper we turn to the study of M5-instantons, by which we really mean those Euclidean M5s in M-theory compactified on Y_4 that descend to D3-brane instantons in the F-theory limit. M5s of this type wrap vertical divisors of the form π^*S_2 with S_2 a divisor in the base B_3 of Y_4 and $\pi : Y_4 \rightarrow B_3$ is the elliptic fibration. In this setting, we focus on the two things that can destroy an M5s ability to contribute a superpotential coupling: fermi zero modes and G -flux. Because our M5s are really describing type IIB D3s wrapping S_2^4 , one expects that the computation of fermi zero modes should be related to how “moveable” S_2 is inside B_3 , that is to cohomologies of the normal bundle $N_{S_2|B_3}$. We work out this relation using the Leray spectral sequence and then use those results to determine all surfaces S_2 in the threefold of [29] that can support an instanton with the right fermi zero mode structure.

We then discuss the interplay of G -flux with M5-instantons. If the restriction of G to an M5 is nontrivial, its effect is to introduce a source term for the chiral 2-form b_2 . This source can sometimes be effectively canceled by introducing wrapped M2-branes ending on the M5 and, in those cases, the M5 generates a charged coupling involving the wrapped M2-brane states. One can see on general grounds that such a coupling will not only be invariant under any $U(1)$ s that were engineered but more generally that it will be invariant under bulk gauge transformations of the M-theory 3-form C_3 . We believe this should incorporate the selection rules alluded to in [28] involving “massive $U(1)$ s” at the KK scale. When the source cannot be canceled by

Heterotic dual exists. In this paper, we crucially avoid any reference to ‘dual matter surfaces’ as they are rather clumsy.

⁴For a review of D3-instantons in type IIB orientifolds see [33].

wrapped M2-branes, we believe the superpotential coupling is forced to vanish⁵.

Using the spectral divisor description of G -fluxes, we are able to argue that fluxes used to generate chirality in F-theory GUT models generically have trivial restriction to M5-instantons so they generate “uncharged” couplings that can play a role in moduli stabilization when the fermi zero mode structure is correct. This is actually easy to see without a detailed computation. The G -fluxes for chirality are engineered so that they only have nonzero integral over matter surfaces obtained by combining a curve of singularities $\Sigma_{sing,i}$ with the curve C_i that degenerates there. In order for G to have nontrivial restriction to the M5, then, it must contain a full curve of singularities in Y_4 and hence contain a full matter surface in its entirety in $\tilde{Y}_4$; otherwise, G would integrate to zero over all divisors in the M5. Any curve of singularities, though, will generically meet an M5 in isolated points in Y_4 ⁶. The restriction of G to such an M5 will be trivial and the coupling generated, if any, will be “uncharged” in the sense that it will involve only moduli and not any fields that arise from wrapped M2s.

The upshot of this analysis is that vertical divisors of elliptically-fibered fourfolds based on the threefold geometry of [29] that we identify as having the right zero mode structure will have no problem generating uncharged couplings if the only G -fluxes we introduce are spectral divisor fluxes for inducing a chiral spectrum. This gives an indication of whether we have any right to expect Kähler moduli stabilization to be possible in geometries constructed from the threefold of [29]. Note that to combine Kähler moduli stabilization with complex structure moduli stabilization we must assume that the latter does not require any G -fluxes that restrict nontrivially to the M5 instantons that are used for the former. This is not a new problem and it is somewhat canonical to assume that complex structure moduli stabilization does not interfere with Kähler moduli stabilization in this way. Such an assumption should not be made cavalierly, though; it is an important one whose validity cannot be taken for granted. Any claims to actually achieve moduli stabilization in an F-theory GUT model must justify it. To be sure, we make no such claims in the present paper.

1.3 Outline

The remainder of this paper is organized as follows. In section 2 we extend the spectral divisor formalism of [10] to describe $U(1)$ s as well as G -fluxes. We also comment on the connection to Higgs bundles, the D3-brane tadpole, and flux quantization. In section 3, we turn to the study of M5-instantons and discuss the two ways that their contribution to the superpotential

⁵The absence of wrapped branes that can cancel the source amounts to the statement that there are no classical solutions to the equations of motion for b_2 in the presence of the source that are properly quantized. There is a factor of $\frac{1}{2}$ that we discuss which leads to a scenario where wrapped branes cannot cancel the source. It is possible that some object exists that can do the job but we do not see a candidate at the moment.

⁶Of course, the M5 will contain curves of singularities Σ'_{sing} along which it intersects surfaces of singularities like S_{GUT} . The only G -flux that would integrate nonzero under the surface obtained by resolving the singularity over Σ'_{sing} will be one that corresponds to a worldvolume flux. In F-theory GUT models we typically do not turn on any worldvolume fluxes except for hypercharge flux, which is globally trivial. As the intersection of the M5 with a surface S_{GUT} of singularities in Y_4 is isomorphic to a curve in S_{GUT} that is nontrivial in the base B_3 of Y_4 , the hypercharge flux will integrate to zero over it.

can be ruined: fermi zero modes and G -flux. Finally, in section 4 we apply these ideas in some generality to geometries based on the threefold of [29]. We identify divisors in the threefold that form the base of vertical divisors in an elliptically fibered Calabi-Yau fourfold that can lead to nontrivial M5 instanton corrections. We also verify that the G -flux needed to generate chirality will restrict trivially to the worldvolumes of those M5s. Appendix A contains a cleaner description of the threefold geometry that we use than the one contained in [29]. Because this description does not utilize any flop transitions we believe it to be more transparent.

2 $U(1)$ s, G -flux, and the Tate divisor

In this section, we review and extend the formalism described in [10] for describing $(2, 2)$ G -fluxes in F-theory GUTs that generate chiral spectra. We would like to emphasize a few things that were not addressed in [10]. The first is a description of how the spectral divisor formalism can be used to take the apparent $U(1)$ s of a local F-theory GUT and explicitly engineer these gauge bosons in a global extension⁷. To explicitly realize $U(1)$ s we will be lead to consider a distinguished object that we term the Tate divisor, which is a special example of a type of object referred to as a spectral divisor in [10]⁸. In addition to allowing an explicit description of $U(1)$ s, the Tate divisor also helps to obtain an explicit description of $U(1)$ flux following ideas of [10]. After this, we turn to a discussion of chiral spectra, whose determination we formulate in a language that is more amenable to the study of GUT singlets that localize on curves that do not sit inside the GUT surface (or any other surface above which the elliptic fibration exhibits non-Abelian singularities). Finally, we make some comments about the D3-brane tadpole including a general argument that the contribution from our G -fluxes arises only from surfaces of singularities in accord with a conjecture of [31]⁹. The extension of ideas that we describe here to surfaces of exceptional type singularities should be straightforward and we have some comments on this. Further generalizations are beyond the scope of this paper but would be an interesting topic of future study.

⁷That this issue is subtle was emphasized in [8] while a focused study of $U(1)$ s in global F-theory models, with which we believe our discussion is consistent, can be found in [9].

⁸As described in [10], a spectral divisor in our singular Calabi-Yau fourfold is one that contains important singularities and whose limiting behavior near those singularities causes it to behave in a favorable way upon resolution. The Tate divisor is a distinguished example of such a divisor that is suitable for describing not only G -fluxes but also $U(1)$ s, as we shall see.

⁹More specifically, [31] checks that the difference in the Euler character of several resolved fourfolds and the naive computation of the Euler character based on data of the singular fourfold only seems to get contributions from surfaces of singularities. As authors of [31] have suggested, this is indicative of flux-induced tadpoles getting contributions only from surfaces of singularities as well.

2.1 Geometric Setup and $U(1)$ s

Our basic setup is an elliptically fibered Calabi-Yau fourfold, Y_4 , that exhibits a surface of $SU(5)_{\text{GUT}}$ singularities¹⁰. We let π denote the elliptic fibration

$$\pi : Y_4 \rightarrow B_3 \quad (2.1)$$

and write the defining equation for Y_4 in the ‘‘Tate form’’

$$y^2 = x^3 + a_0 z^5 + a_2 z^3 x + a_3 z^2 y + a_4 z x^2 + a_5 x y. \quad (2.2)$$

Here, z is the holomorphic section on B_3 whose vanishing defines the surface S_{GUT} and the a_m ’s are sections of the bundles $\mathcal{O}((m-6)K_{B_3} + (m-5)S_{\text{GUT}})$. This geometry exhibits an $SU(5)_{\text{GUT}}$ singularity along $z=0$ [34] with enhancements in singularity type to $SO(10)$ where $z = a_5 = 0$ and $SU(6)$ where $z = a_0 a_5^2 - a_2 a_3 a_4 + a_3^2 a_4 = 0$. The divisor $z=0$ supports the degrees of freedom of an $\mathcal{N} = 1$ supersymmetric gauge theory with gauge group $SU(5)$ while the curves of $SO(10)$ and $SU(6)$ enhancement support charged chiral matter. The string theoretic origin of that matter is well understood: matter fields arise from M2-branes wrapping the extra vanishing cycles or, equivalently, the (p, q) strings that become massless there in the type IIB language.

In this setting, both $U(1)$ symmetries and the G -fluxes responsible for chiral spectra are somewhat subtle to describe. This is because both are sensitive to the monodromic structure of the geometry near $z=0$ (among other things), where the $SU(5)_{\text{GUT}}$ degrees of freedom live. We focused on G -fluxes in [10] so we will place a greater emphasis on $U(1)$ s in the present paper.

One of the nice features of $U(1)$ s from a phenomenological perspective is that they can distinguish different types of 4-dimensional fields that sit in the same $SU(5)_{\text{GUT}}$ representation. We are usually interested in distinguishing $\mathbf{\bar{5}}$ ’s (so that left-handed leptons do not participate in the same couplings as down type Higgs doublets) but, for simplicity of presentation, we focus instead on $\mathbf{10}$ ’s that localize on the curve $z = a_5 = 0$. This curve, $\Sigma_{\mathbf{10}}$, is the locus of $SO(10)$ singularities and, if a_5 is chosen appropriately, may split into multiple components. The type of $U(1)$ we are interested in should couple differently to $\mathbf{10}$ ’s associated to the different components so, to think about $U(1)$ s geometrically, we ask a related question: is there a way to distinguish (or tell if it is possible to distinguish) between the 2-cycles that degenerate along different components of $\Sigma_{\mathbf{10}}$ when it is reducible? In trying to distinguish these 2-cycles geometrically we will end up building the $U(1)$ that we want.

As reviewed in the introduction, $U(1)$ s in F-theory originate from harmonic 2-forms, which integrate to zero over the fiber class, F , and any curve that sits inside the section of the elliptic fibration. Given such a 2-form in our Calabi-Yau fourfold, we can reduce the M-theory 3-form C_3 on it to obtain a 3-dimensional gauge field that becomes a 4-dimensional gauge field in the F-theory limit (where the elliptic fiber is shrunk to zero volume while holding τ fixed). The

¹⁰There may be hidden sectors consisting of additional surfaces of singularities but we ignore this possibility for now, and will return to this in the context of M5 instantons.

$U(1)$ charge of chiral fields that descend from M2-branes wrapped on a cycle C , then, is simply the integral of ω over C

$$q_C \sim \int_C \omega. \quad (2.3)$$

The cycles C that house chiral matter are holomorphic so we will want ω to be a $(1,1)$ -form. Alternatively, we specify ω by a dual divisor D_ω in the resolved geometry, so that

$$q_C \sim D_\omega \cdot C. \quad (2.4)$$

If two curves C_1 and C_2 that support wrapped M2s are homologically distinct, it should be possible to construct a divisor D_ω such that

$$D_\omega \cdot C_1 \neq D_\omega \cdot C_2. \quad (2.5)$$

This can be used to construct a $U(1)$ with respect to which the fields from C_1 and C_2 carry distinct charges.

2.2 Local geometry and dP_9

Because $U(1)$ charges involve integrations over vanishing 2-cycles, the existence of a $U(1)$ that distinguishes charged fields is closely tied to how singularities are resolved. This can be quite complicated in general so our strategy will be to take a somewhat indirect route motivated by the cylinder map of Heterotic/F-theory duality [15, 16, 18–20]. Rather than explicitly resolving all singularities, we will embed our local geometry into a different fourfold $\hat{Y}_4$ in which the singularity structure can be related to the geometry of a family of dP_9 surfaces. The advantage of embedding our local geometry into a family of dP_9 's is that we can describe the singularity structure in terms of dP_9 homology cycles that degenerate above various loci in S_{GUT} ¹¹.

To proceed, then, consider the local geometry

$$y^2 = x^3 + b_0 z^5 + b_2 z^3 x + b_3 z^2 y + b_4 z x^2 + b_5 x y \quad (2.6)$$

where the b_m 's are just the restrictions of the global holomorphic sections a_m in (2.2) to S_{GUT}

$$b_m = a_m|_{S_{\text{GUT}}}. \quad (2.7)$$

This is the local model for the geometry near S_{GUT} and its defining data admits a well-known mapping to Higgs bundle data of the 7-brane worldvolume gauge theory [1, 18–20, 35]. It is a trivial matter to embed this local geometry into a K3-fibered Calabi-Yau fourfold $\hat{Y}_4$ by embedding S_{GUT} into the threefold $\hat{B}_3 = \mathbb{P}(\mathcal{O} \oplus N)$ with N the normal bundle of S_{GUT} inside B_3 . By construction, our surface S_{GUT} has an identical normal bundle in $\hat{B}_3$ as in B_3 and all

¹¹Of course this limits the type of cycles we can consider but there is no problem for phenomenologically relevant examples in which all singularities that we deal with embed into E_8 .

of the holomorphic sections b_m can be trivially extended to global ones in $\hat{B}_3$ that can be used to define $\hat{Y}_4$ ¹².

With this K3-fibered Calabi-Yau we can utilize Heterotic/F duality. Indeed, the spectral divisor formalism for G -fluxes [10] is a strategy to apply ideas of the cylinder map in a more general setting [1, 15, 16, 18, 19, 36]. We are not interested in Heterotic/F duality here per se, but rather in the stable degeneration limit [37, 38] of $\hat{Y}_4$ so that our local geometry becomes embedded into a dP_9 fibration, which we think of as a family of dP_9 's parametrized by S_{GUT} . We can explicitly present the family as

$$y^2 = x^3 + x f_4 z^4 + g_6 z^6 + w (b_0 z^5 + b_2 z^3 x + b_3 z^2 y + b_4 z x^2 + b_5 x y) , \quad (2.8)$$

where w and z are homogeneous coordinates on the $\mathbb{P}^1$ fiber of $\hat{B}_3$ ¹³.

In presenting the dP_9 as an elliptic fibration over $\mathbb{P}^1$ with section, we have distinguished two curve classes. These are the base, e_9 , and the elliptic fiber, F , which is a representative of the anti-canonical class. Additional cycles that degenerate correspond to -2 curves that fail to intersect either e_9 or F . Elements of $H_2(dP_9, \mathbb{Z})$ with this property are referred to as roots and, following [36], we label this collection by R_8

$$R_8 = \{C \in H_2(dP_9, \mathbb{Z}) | C \cdot F = C \cdot e_9 = 0, C^2 = -2\} . \quad (2.9)$$

The intersection matrix on R_8 is equivalent to (-1) times the Cartan matrix of E_8 so the roots C are naturally identified with elements of the E_8 root lattice. One inconvenient feature of the C 's is that they do not have effective holomorphic representatives in generic dP_9 surfaces (i.e. they do not sit in the Mori cone). Of course, these dP_9 's are somewhat special in that all of them exhibit $SU(5)_{\text{GUT}}$ singularities. The 4 roots of $SU(5)_\perp$ are therefore in the Mori cone throughout the family¹⁴ so the monodromy group as we move in the family must sit inside the Weyl group of the $SU(5)_\perp$ commutant of $SU(5)_{\text{GUT}}$ inside E_8 , which is the symmetric group on five objects.

2.3 From dP_9 to $U(1)$ s

The dP_9 fibers over matter curves like Σ_{10} exhibit an additional holomorphic curve corresponding to the new vanishing cycle. We can think of this as coming from a new root that enters the

¹²It may be that $\hat{Y}_4$ has some rather sick singularities far from S_{GUT} . This will not have any affect on what we say here, though, as our interest is only in small resolutions of singularities along S_{GUT} .

¹³Because the power of w out front is only 1 this does not define a Calabi-Yau fourfold. Rather, it is only 'half' of the Calabi-Yau fourfold $\hat{Y}_4$ that we can split off in the stable degeneration limit. It has the advantage that if the local geometry of Y_4 is well behaved then the dP_9 -fibration is also well-behaved in that there are no 'sick' singularities at $w = 0$. Even though the dP_9 -fibration is not itself Calabi-Yau, it is a fine place to study the small (crepant) resolutions we need to desingularize Y_4 . As described in the introduction, we could skip the K3-fibration altogether and simply start by embedding our local geometry into the dP_9 -fibration (2.8).

¹⁴This can be achieved in the standard picture of dP_9 as $\mathbb{P}^2$ blown up at 9 points by taking the limit as several of the blown up points move to nongeneric locations, such as sitting on top of one another or sitting on a common line.

Mori cone. Our task of distinguishing different kinds of **10**'s boils down to determining whether the new effective root on one component of $\Sigma_{\mathbf{10}}$ is related by monodromy to the new effective root on another component as we move in the family. The easiest way to do this would be to simply follow one of the roots through the family. This cannot be done easily because the root exits the Mori cone as soon as we move off of $\Sigma_{\mathbf{10}}$. Fortunately, however, the roots are in 1-1 correspondence with elements of a second distinguished set

$$I_8 = \{\ell \in H_2(dP_9, \mathbb{Z}) | \ell \cdot F = 1, \ell \cdot e_9 = 0, \ell^2 = -1\} \quad (2.10)$$

according to the identification

$$\ell = (F + e_9) - C. \quad (2.11)$$

Elements of I_8 are precisely the exceptional lines of dP_9 that miss the section e_9 . They are in the Mori cone of generic dP_9 's so they have nice effective holomorphic representatives throughout the family. If we want to follow the behavior of a particular root as we move along S_{GUT} , then, it is often easier to look at what happens to the effective line from I_8 that is dual to that root in the sense of (2.11).

We are interested in a particular subset of the roots that correspond to **10**'s of $SU(5)_{\text{GUT}}$. The group theoretic decomposition of the E_8 adjoint under

$$\begin{aligned} E_8 &\rightarrow SU(5)_{\text{GUT}} \times SU(5)_{\perp} \\ \mathbf{248} &\rightarrow (\mathbf{24}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{24}) \oplus [(\mathbf{10}, \mathbf{5}) \oplus \text{cc}] \oplus [(\overline{\mathbf{5}}, \mathbf{10}) \oplus \text{cc}] \end{aligned} \quad (2.12)$$

tells us that, modulo the action of $SU(5)_{\text{GUT}}$ roots, there are five such roots, transforming as a fundamental of $SU(5)_{\perp}$, that can move into the Mori cone above a given component of $\Sigma_{\mathbf{10}}$. To see how their homological classes mix in the family, we study instead the behavior of the 5 exceptional lines ℓ_i dual to those roots in the sense of (2.11). Given the form of the dP_9 fibration (2.8), we can describe the union of these five exceptional lines (fibered over S_{GUT}) by the divisor $\mathcal{C}$ [36]¹⁵

$$\mathcal{C} : b_0 z^5 + b_2 z^3 x + b_3 z^2 y + b_4 z x^2 + b_5 x y = 0. \quad (2.13)$$

To ensure that some **10**'s are distinct from others, what we need is for the divisor (2.13) to split into multiple components in our dP_9 -fibration. If this happens, some exceptional lines will not mix with others in the fibration. We will have two distinct sets of lines $\{\ell_i\}$ and $\{\ell_a\}$ and, correspondingly, two distinct sets of roots, $\{C_i\}$ and $\{C_a\}$.

Along with guaranteeing that the $\{C_i\}$ and $\{C_a\}$ do not mix, the splitting of our exceptional lines into the sets $\{\ell_i\}$ and $\{\ell_a\}$ also provides a means for distinguishing them. For concreteness, let us suppose that there are n roots in the group $\{C_i\}$ and m in the group $\{C_a\}$. In that case, consider the following element of $H_2(dP_9, \mathbb{Z})$

$$w = m \sum_{i=1}^n \ell_i - n \sum_{a=1}^m \ell_a. \quad (2.14)$$

¹⁵When restricted to any single dP_9 of the family, $\mathcal{C}$ describes a union of five exceptional lines. Because it is irreducible for a generic family with $SU(5)_{\text{GUT}}$ singularity at $z = 0$, these five exceptional lines are mixed by the generic S_5 monodromy group and hence are precisely the ℓ_i 's that are dual to the **10** roots in the sense of (2.11) provided we take a proper transform in the dP_9 .

This is a nontrivial element of $H_2(dP_9, \mathbb{Z})$ that distinguishes roots in the sets $\{C_i\}$ and $\{C_a\}$ via the intersection relations

$$w \cdot C_i = m, \quad w \cdot C_a = -n, \quad (2.15)$$

which follow from the local intersection data between the exceptional lines ℓ_i and roots C_i ($i = 1, \dots, 5$) in dP_9

$$\ell_i \cdot C_j = 1 + \delta_{ij}. \quad (2.16)$$

The action of w therefore reduces to that of a Cartan generator of $SU(5)_\perp$

$$w \leftrightarrow \text{diag}(m, m, \dots, m, -n, -n, \dots, -n). \quad (2.17)$$

When the divisor (2.13) splits into two components $\mathcal{C}^{(m)}$ and $\mathcal{C}^{(n)}$ with m and n sheets, respectively, the combination w is invariant under the monodromy action of our family so that it fibers over S_{GUT} to yield a nontrivial divisor given by the linear combination $n\mathcal{C}^{(m)} - m\mathcal{C}^{(n)}$. This divisor, in turn, provides a $(1, 1)$ -form that can be used to obtain a $U(1)$ gauge field that couples differently to M2-branes wrapping the roots $\{C_i\}$ and $\{C_a\}$ according to the group theoretic intersection (2.15).

While the preceding discussion made use of dP_9 geometry to describe the nature of the singularity, the prescriptions for distinguishing roots and computing $U(1)$ charges relied only on local intersection data near S_{GUT} . To be sure, even though the object $\mathcal{C}$ was defined globally, its intersections with the roots $\{C_i\}$ and $\{C_a\}$ all occur in the neighborhood of S_{GUT} . In that sense, the collection of exceptional lines provides us with the construction of a noncompact divisor in the local geometry near S_{GUT} that yields a $U(1)$ symmetry capable of distinguishing the $\mathbf{10}$'s on different components of $\Sigma_{\mathbf{10}}$ ¹⁶. Engineering an honest $U(1)$ in a compact model just requires us to provide a global extension of this noncompact divisor in a more general setting without the global structure of a dP_9 or K3.

2.4 The Tate divisor

Given a Tate model of the form (2.2), we propose that a suitable object that extends the noncompact divisor of our local model is the Tate divisor defined by

$$\mathcal{C}_{\text{Tate}} : \quad a_0 z^5 + a_2 z^3 x + a_3 z^2 y + a_4 z x^2 + a_5 x y, \quad (2.18)$$

where it is understood that we take a suitable proper transform when passing to the resolution of Y_4 ¹⁷. In the neighborhood of S_{GUT} this behaves in precisely the same way as the object $\mathcal{C}$ above so that we can directly apply the local intersection data (2.16) without having to work through the explicit resolution of singularities for each Y_4 .

¹⁶Factoring of this noncompact divisor is of course trivially equivalent to factoring of the Higgs bundle spectral cover.

¹⁷This is pretty important since $\mathcal{C}_{\text{Tate}}$ is singular along the surface of $SU(5)_{\text{GUT}}$ singularities where all 5 sheets come together. Resolution of the $SU(5)_{\text{GUT}}$ singularities separates the sheets and removes this singularity.

The divisor $\mathcal{C}_{\text{Tate}}$ is an example of a spectral divisor, an object that we introduced in [10] for defining global G -fluxes in a way that exploited the connection to the Heterotic string using essentially the same type of reasoning as above. Spectral divisors are singled out with respect to other divisors by the fact that their intersection with π^*S_2 contains the Higgs bundle spectral cover. For the Tate divisor we can see this by noting that the meromorphic section t defined as

$$t = \frac{y}{x} \quad (2.19)$$

is actually holomorphic when restricted to $\mathcal{C}_{\text{Tate}}$ so that we can write the defining equation (2.18) as¹⁸

$$a_0 z^5 + a_2 z^3 t^2 + a_3 z^2 t^3 + a_4 z t^4 + a_5 t^5. \quad (2.20)$$

Among the spectral divisors, there is one, the Tate divisor $\mathcal{C}_{\text{Tate}}$, that is further distinguished in that there is a simple definition for it that can be canonically associated to any Y_4 (i.e. without making any reference to the sections a_m that appear in the Tate form) and is furthermore effective at determining the presence of additional divisors for $U(1)$ symmetries. In particular, this is a divisor that can be easily chosen to extend factorization structures of the Higgs bundle spectral cover in order to build additional $U(1)$ symmetries¹⁹.

When the Tate divisor becomes reducible into components $\mathcal{C}^{(m)}$ and $\mathcal{C}^{(n)}$ with m and n sheets, respectively, it reflects the fact that some of the **10** roots that degenerate along curves in S_{GUT} are distinguished from others and also provides us with a $U(1)$ that can make the distinction. Explicitly, the $U(1)$ that does the job is a traceless combination

$$\omega = n\mathcal{C}^{(m)} - m\mathcal{C}^{(n)} - \pi^*\delta, \quad (2.21)$$

where δ is chosen to ensure that ω is orthogonal to the horizontal and vertical divisors of Y_4 ²⁰.

As a homogeneous polynomial of degree 5 in z and t it is fairly easy to choose the a_m 's so that this object splits. The algebra is completely equivalent to the splitting of Higgs bundle spectral covers in local model building [2, 7, 30, 31].

2.5 G -Fluxes, Matter Surfaces, and Chiral Spectrum

The Tate divisor can also be used to construct G -fluxes for chirality as described in [10]. We do not repeat that discussion here but simply remind the reader of the prescription to describe these G -fluxes as $(1,1)$ -forms inside $\mathcal{C}_{\text{Tate}}$. Such a $(1,1)$ -form effectively defines a $(2,2)$ -form in the fourfold obtained by resolving the singularities of Y_4 that, as an object inside $\mathcal{C}_{\text{Tate}}$, is sensitive to the details of that resolution. It is this last fact that allows G to know about the degenerate roots and integrate nontrivially over the matter surfaces, which we define as the

¹⁸We use the fact that $y = t^3$ and $x = t^2$ on $\mathcal{C}_{\text{Tate}}$.

¹⁹We expect that $\mathcal{C}_{\text{Tate}}$ splits whenever a new $U(1)$ divisor is present with nonzero intersection with some degenerate 2-cycles above S_2 . It would be interesting to show this in generality.

²⁰We have to take a traceless combination because something with pure trace, like say $\mathcal{C}^{(m)}$, will have nontrivial intersection with the fiber class by virtue of the fact that $\ell_i \cdot F = 1$ (2.10).

surface in $\tilde{Y}_4$ that maps to a curve of singularities in Y_4 under the blow-down map. In general, these surfaces should have the structure of a resolved root fibered over a curve.

The G -fluxes needed to engineer chiral matter can be defined when $\mathcal{C}_{\text{Tate}}$ is irreducible and, in fact, we can even use a different “spectral divisor” to construct G -flux whose local defining equation near the surface of $SU(5)_{\text{GUT}}$ singularities is equivalent to that of $\mathcal{C}_{\text{Tate}}$ despite differing globally. Nevertheless, to keep things simple we will always use $\mathcal{C}_{\text{Tate}}$ to study G -fluxes in this paper. In that case, we get a particularly nice type of G -flux when $\mathcal{C}_{\text{Tate}}$ is reducible and can be used to construct $U(1)$ s. We simply take a $(1, 1)$ -form ρ in B_3 , pull it back to each component of $\mathcal{C}_{\text{Tate}}$, and take a traceless combination of the two

$$G = (n\mathcal{C}^{(m)} - m\mathcal{C}^{(n)}) \cdot \pi^* \rho - G_0, \quad (2.22)$$

where G_0 is a subtraction term that we must include to ensure that G is orthogonal to horizontal and vertical divisors in Y_4 . The form (2.22) is very much in the spirit of how G -fluxes in F-theory should heuristically be related to $U(1)$ flux as

$$G \sim \omega_i \wedge F_i \quad (2.23)$$

with ω_i the $(1, 1)$ -form specifying a $U(1)$ and F_i the flux associated to that $U(1)$.

Let us turn now to determining the chiral spectrum, which involves integrating G over various matter surfaces. In [10] we studied ways to do this with a focus on rephrasing the computation as one that can be done in terms of divisor classes and intersections in Y_4 . Rather than executing gymnastics like that, we just review here how the Higgs bundle picture emerges.

The G -flux that we build will have the generic form

$$G = \mathcal{G} - G_0, \quad (2.24)$$

where $\mathcal{G}$ is obtained from the Tate divisor (as a holomorphic surface inside $\mathcal{C}_{\text{Tate}}$) and G_0 is some holomorphic surface in Y_4 whose contribution we must add to ensure that G is orthogonal to horizontal and vertical divisors in Y_4 . Chiral matter localizes on curves of singularities Σ where 2-cycles C degenerate and the flux that they couple to is obtained by restricting G to the matter surface ‘ $\Sigma \times C$ ’²¹ and integrating it over C . The net chirality follows from integrating G over the entire matter surface and this is the computation we ultimately want to describe.

Intersecting G with any holomorphic surface will require a special type of calculation. From the form (2.24) we see that intersections of $\mathcal{G}$ with generic holomorphic surfaces will generally be cancelled by intersections with G_0 . The types of intersections for which this does not happen are those that occur at the singular locus, to which $\mathcal{G}$ is sensitive but G_0 is not. By construction, then, G only has nonzero integral over surfaces that include degenerate cycles so for any computation we can focus our attention on the neighborhood of the curve of singularities in question. For $SU(5)_{\text{GUT}}$ -charged matter, it is enough to focus on the limiting behavior of $\mathcal{C}_{\text{Tate}}$ and $\mathcal{G}$ near the surface of $SU(5)_{\text{GUT}}$ singularities, which we do by taking $y/x = t \rightarrow 0$ and $z \rightarrow 0$ with $s = z/t$ fixed. Recall that $\mathcal{C}_{\text{Tate}}$ is described by

$$a_0 z^5 + a_2 z^3 x + a_3 z^2 y + a_4 z x^2 + a_5 x y. \quad (2.25)$$

²¹We put $\Sigma \times C$ in quotes because the surface does not have to be a direct product.

In the limit, then, $\mathcal{C}_{\text{Tate}}$ becomes

$$\mathcal{C}_{\text{Tate}} \rightarrow t^5 (b_0 s^5 + b_2 s^3 + b_3 s^2 + b_4 s + b_5) \quad (2.26)$$

with the b_m 's now restricted to sections on S_{GUT} and s a section of the canonical bundle on S_{GUT} . The term in $()$'s here is nothing other than the Higgs bundle spectral cover, $\mathcal{C}_{\text{Higgs}}$ [1]. The limiting behavior of $\mathcal{G}$ also produces something familiar; it gives us a divisor γ inside $\mathcal{C}_{\text{Higgs}}$ that we would like to interpret as the corresponding object in the local model.

It is interesting to see in some detail how the approach of $\mathcal{C}_{\text{Tate}}$ to the surface of $SU(5)_{\text{GUT}}$ singularities takes place. For starters, the t^5 factor tells us that every (local) sheet of $\mathcal{C}_{\text{Tate}}$ meets the surface of $SU(5)_{\text{GUT}}$ singularities once above each point on S_{GUT} . This reflects the fact that each exceptional line ℓ_i in the dP_9 picture transforms in a **10** of $SU(5)_{\text{GUT}}$ and must therefore intersect some of the $SU(5)_{\text{GUT}}$ roots at each point on the surface of singularities. This accounts for the ‘1’ in the local intersection data (2.16) and has no effect on any of our computations when the G -flux is chosen to be traceless²².

The rest of the approach is captured by the behavior of the Higgs bundle spectral cover. Consider first the curve of $SO(10)$ singularities, Σ_{10} , which sits at $y = b_5 = 0$. Within the second piece of (2.26) there is a distinguished curve, $t = b_5 = 0$, that coincides with Σ_{10} . It captures part of the Tate divisor that lands exactly on Σ_{10} as we approach the surface of $SU(5)_{\text{GUT}}$ singularities. The curve $t = b_5 = 0$ is familiar from local model building: it is the “local **10** matter curve” by which we mean the **10** matter curve of the local model that sits inside the Higgs bundle spectral cover $\mathcal{C}_{\text{Higgs}}$. Here we call this curve $\Sigma_{10, \text{Higgs}}$ and it appears naturally in the Calabi-Yau fourfold as describing the limiting behavior of the specific (local) sheet of $\mathcal{C}_{\text{Tate}}$ that lands directly on the curve of $SO(10)$ singularities. This is just another way of saying that $\Sigma_{10, \text{Higgs}}$ describes a cross section of the exceptional line ℓ_i that is dual to the root C_i whose degeneration causes the $SO(10)$ enhancement.

From this point of view, it is clear that restricting G to the matter surface ‘ $\Sigma_{10} \times C$ ’ is equivalent to projecting $\mathcal{G}$ onto $\Sigma_{10, \text{Higgs}}$, which is isomorphic to the **10** matter curve in the copy of S_{GUT} that sits in the section²³. This reproduces the standard result from Higgs bundles that charged fields are sections of

$$H^m(\Sigma_{10}, K_{\Sigma_{10}}^{1/2} \otimes i_* \gamma) \quad m = 0, 1 \quad (2.27)$$

with Σ_{10} the curve of $SO(10)$ enhancements in S_{GUT} . Recall that γ is a divisor in $\mathcal{C}_{\text{Higgs}}$ that represents the limiting behavior of $\mathcal{G}$ in our global approach and specifies the “local flux” in a local model. The map i is just the usual embedding $i : \Sigma_{10, \text{Higgs}} \rightarrow \Sigma_{10}$.

²²Indeed, from this point of view, the traceless condition of the G -flux that we imposed in [10] is really just the statement that we require our G -flux to be orthogonal to all $SU(5)_{\text{GUT}}$ roots. Trace parts may have an important role to play, however. It has been known for some time [19] that what is commonly referred to as hypercharge flux must include some component from the bulk in the sense that it is not a flux in the pure $U(1)_Y$ direction. A G -flux constructed within the Tate divisor that has a pure trace piece is exactly the sort of thing we need to give a proper definition of this hypercharge flux.

²³Note that the singularities of Y_4 , including the surface of singularities above S_{GUT} , do not lie in the section.

There is a similar story for the $\bar{\mathbf{5}}$ curve. To see this, recall that the surface of $SU(5)_{\text{GUT}}$ singularities is concretely given by the intersection of the “Tate form” (2.2)²⁴

$$y - \frac{1}{2}(b_3x + b_5z^2) = z = 0. \quad (2.28)$$

With our current variables, the first of these corresponds to²⁵

$$t^3 - \frac{z^2}{2}(b_3s^2 + b_5) = 0. \quad (2.29)$$

Now, the $\bar{\mathbf{5}}$ curve $\Sigma_{\bar{\mathbf{5}}, \text{Higgs}}$ of the local model with Higgs bundle spectral cover $\mathcal{C}_{\text{Higgs}}$ is described by

$$b_3s^2 + b_5 = b_0s^4 + b_2s^2 + b_4 = 0, \quad (2.30)$$

which we can essentially think of as the intersection of $b_3s^2 + b_5 = 0$ with $\mathcal{C}_{\text{Higgs}}$. As we send $t \rightarrow 0$, then, the curve $\Sigma_{\bar{\mathbf{5}}, \text{Higgs}}$ lands directly on the curve of $SU(6)$ singularities.

As in the case of $\mathbf{10}$ ’s, this suggests that restricting G to the matter surface ‘ $\Sigma_{\bar{\mathbf{5}}} \times C$ ’ is equivalent to projecting $\mathcal{G}$ onto the curve of $SU(6)$ singularities. This projection is now 2-1 for the usual reasons and leads to the familiar result from Higgs bundles that $\bar{\mathbf{5}}$ ’s and $\mathbf{5}$ ’s are sections of

$$H^m(\Sigma_{\bar{\mathbf{5}}}, K_{\Sigma_{\bar{\mathbf{5}}}}^{1/2} \otimes \nu_*\gamma) \quad m = 0, 1, \quad (2.31)$$

where ν is the 2-1 covering map from the curve $\Sigma_{\bar{\mathbf{5}}, \text{Higgs}}$ in $\mathcal{C}_{\text{Higgs}}$ to the curve $\Sigma_{\bar{\mathbf{5}}}$ of $SU(6)$ singularities inside S_{GUT} . We know from the seminal work of [36] that we should actually take $\Sigma_{\bar{\mathbf{5}}}$ to be the normalization of the curve of $SU(6)$ singularities. We understand this statement here in the following way. Singularities of $\Sigma_{\bar{\mathbf{5}}}$ come from nodes where the singularity type enhances. The two branches of $\Sigma_{\bar{\mathbf{5}}}$ that meet at such a node are places where homologically distinct cycles are degenerating so those branches are separated in the lift to $\mathcal{C}_{\text{Higgs}}$ (which we view as the limiting behavior of $\mathcal{C}_{\text{Tate}}$) upon resolution.

Note that this picture makes clear why matter fields are associated with sections on matter curves inside the Higgs bundle spectral cover. Those curves are capturing the actual curves of singularities, as seen by the G -flux, in the fourfold. It is along the curves of singularities, not their projection to the section (which does not meet the singularities), that the wrapped branes are found.

Even though we can say something about the actual cohomology groups associated to various charged matter fields, let us say a few more things about the simpler question of net chirality. On general grounds we expect that the net chirality on a matter curve Σ is obtained by integrating G over the corresponding matter surface ‘ $\Sigma \times C$ ’ and, from our discussion above, this amounts to integrating γ over the “local matter curves” $\Sigma_{R, \text{Higgs}}$ inside the Higgs bundle spectral cover. The only reason for the appearance of $\mathcal{C}_{\text{Higgs}}$, though, was our insistence on studying the chiral

²⁴This corresponds to $\hat{y} = z = 0$ in the corresponding Weierstrass model. The equation for y reflects the shift that is needed to go from Tate to Weierstrass form.

²⁵This relation is just setting the Y coordinate of the Weierstrass form to zero.

spectrum on matter curves that sit inside the surface of $SU(5)_{\text{GUT}}$ singularities²⁶. We can give a description of the computation of chiral matter that does not make reference to $\mathcal{C}_{\text{Higgs}}$ as follows. First, we embed the curve of singularities Σ associated to a matter surface ‘ $\Sigma \times C$ ’ into $\mathcal{C}_{\text{Tate}}$. Then, we determine the intersection of $\mathcal{G}$ with that surface.

The nice thing about this prescription is that it works equally well for matter curves that do not sit inside the surface of $SU(5)_{\text{GUT}}$ singularities. In principle, then, we should be able to determine the net chirality of $SU(5)_{\text{GUT}}$ -singlet fields that carry $U(1)$ charge in $SU(5)_{\text{GUT}}$ models that engineer an extra $U(1)$. We hope to report soon on models in which this computation is explicitly carried out.

2.6 D3-brane tadpole

Let us now make some remarks about the D3-brane tadpole. To compute the D3-brane tadpole induced by a G -flux, what we need to evaluate is a self-intersection

$$\int_{Y_4} G \wedge G, \quad (2.32)$$

which we can think of in two pieces

$$\mathcal{G} \cdot (\mathcal{G} - G_0) - G_0 \cdot (\mathcal{G} - G_0) \quad (2.33)$$

with $\mathcal{G}$ and G_0 defined as in (2.24). The second term here vanishes by construction because G_0 is an honest surface in Y_4 that is constructed as a linear combination of intersections of horizontal and vertical divisors. What we are after, then, is the first term. We can think of this as integrating G over the surface $S_{\mathcal{G}}$ in the resolution of Y_4 that is determined by $\mathcal{G}$. As usual, any integral of G over a surface will receive contributions only from places where $\mathcal{G}$ meets singularities that have to be resolved. So, what we want is to look at the full locus where $\mathcal{G}$ meets singularities and restrict $\mathcal{G}$ to that locus. If we consider, for instance, the surface of $SU(5)_{\text{GUT}}$ singularities then the restriction of $\mathcal{G}$ is the curve γ that sits inside the Higgs bundle spectral cover $\mathcal{C}_{\text{Higgs}}$. From the surface of $SU(5)_{\text{GUT}}$ singularities, then, we get a contribution to (2.32) from each point in the self-intersection $\gamma \cdot_{\mathcal{C}_{\text{Higgs}}} \gamma$. The setup is in fact almost identical to the computation of chiral matter except that we have replaced the matter curve by the curve γ in order to reflect the fact that we are integrating over the curve $S_{\mathcal{G}}$ determined by $\mathcal{G}$. When we do this, though, we must be a little careful because $\mathcal{G}$ is defined on a union of $\ell_i \sim (x_9 - e_9) - C_i$ ’s and the contribution to (2.32) that comes from a singular point arises from the C_i . As the sign is opposite from what we have when we integrate G over a matter surface we find that the contribution to (2.32) is in fact $-\gamma \cdot_{\mathcal{C}_{\text{Higgs}}} \gamma$. Note that this type of reasoning is not at all new and should be familiar from studies of Heterotic/F-theory duality [15, 16, 18–20].

More generally, we expect (2.32) to be computed by the restriction of $\mathcal{G} \cdot_{\mathcal{C}_{\text{Tate}}} \mathcal{G}$ to the locus of singularities with the appropriate signs. Note that $\mathcal{G} \cdot_{\mathcal{C}_{\text{Tate}}} \mathcal{G}$ is a curve inside $\mathcal{C}_{\text{Tate}}$ so it

²⁶The presence of the ambient surface of singularities was also responsible for the gymnastics associated with imposing tracelessness and dropping the t^5

will generically miss curves of singularities or isolated point singularities. This means that (2.32) should get contributions only from surfaces of singularities in accord with a conjecture of [31]. The discussion for a surface of $SU(5)_{\text{GUT}}$ singularities should generalize to something like the following for the D3-brane tadpole induced by G -flux when there are multiple surfaces of singularities that do not intersect one another

$$Q_{D3,\text{induced}} = -\frac{1}{2} \int_{Y_4} G \wedge G = \frac{1}{2} \sum_{\text{surfaces of singularities}, i} \int_{\mathcal{C}_i} \gamma_i^2, \quad (2.34)$$

where $\mathcal{C}_i$ is essentially the Higgs bundle spectral cover for the i th surface and γ_i the limit of $\mathcal{G}$. It is actually not hard to see that a spectral cover for an $SU(n)$ Higgs bundle emerges from the limiting behavior of $\mathcal{C}_{\text{Tate}}$ near any surface of singularities whose commutant inside E_8 is an $SU(n)$ group or a product of such groups. Looking at the behavior of $\mathcal{C}_{\text{Tate}}$ for more general types of singularities and interpreting the result would be very interesting but is beyond the scope of this paper.

2.7 Remarks on Flux Quantization

We now turn to flux quantization, an issue that we somewhat neglected in [10]. Rather than simply rewriting Witten's famous condition [39], let us recall one of the original observations that points to the need for modifying the G -flux quantization rule. This will motivate a consistency check that we perform later.

The D3-brane tadpole in Calabi-Yau fourfold compactifications of F-theory receives both a flux contribution, which we have described, and a geometric contribution from the Euler character

$$n_{D3} = \frac{\chi}{24} - \frac{1}{2} G^2. \quad (2.35)$$

For smooth Calabi-Yau fourfolds, it was shown in [40] that χ is always divisible by 6 but not necessarily by 12. When $\chi/24$ fails to be an integer, the quantization of G must be modified in order to account for this fact and ensure an integral induced D3-brane charge. In [40], it is proven that χ is always divisible by 24 when the fourfold admits a smooth Weierstrass description. More recent work [29, 41] uses similar reasoning to argue directly that G -fluxes are integrally quantized in such cases²⁷. When our Y_4 is singular, though, we need to ask questions about χ and G -flux quantization on the smooth resolution $\tilde{Y}_4$ which is not of this type. We can think of $\tilde{Y}_4$ as a smooth Calabi-Yau fourfold but cannot use any special properties attributed to fourfolds that can be realized as smooth Weierstrass models.

The connection between χ and G -flux quantization arises from a few simple facts about Calabi-Yau fourfolds. First of all, the Todd genus of a smooth Calabi-Yau fourfold $\tilde{Y}_4$ is 2, which implies that

$$\lambda^2 = 480 + \frac{\chi}{3}, \quad (2.36)$$

²⁷ [29] showed that $c_2(\tilde{Y}_4)$ is even whenever the class $c_1(B_3)^2 - c_2(B_3)$ is even in B_3 and argued that compactifications in which $c_1(B_3)^2 - c_2(B_3)$ is odd are necessarily Lorentz-violating. The work [41] actually proved that $c_1(B_3)^2 - c_2(B_3)$ is always even.

where λ denotes the second Chern class

$$\lambda = c_2(\tilde{Y}_4). \quad (2.37)$$

Using this, we can write the induced D3-brane charge as

$$\begin{aligned} n_{D3} &= 60 - \frac{1}{2} \left[G^2 - \left(\frac{\lambda}{2} \right)^2 \right] \\ &= 60 - \frac{1}{2} \left[\left(\alpha - \frac{\lambda}{2} \right)^2 - \left(\frac{\lambda}{2} \right)^2 \right] \\ &= 60 - \frac{1}{2} [\alpha^2 - \alpha \cdot \lambda], \end{aligned} \quad (2.38)$$

where we implicitly defined

$$\alpha = G + \frac{\lambda}{2}. \quad (2.39)$$

Writing things in this way is helpful because the intersection of any holomorphic surface $\tilde{S}$ inside $\tilde{Y}_4$ with the class λ satisfies [39]

$$\tilde{S} \cdot \lambda = \tilde{S}^2 \bmod 2 \quad (2.40)$$

so that the induced D3-brane charge is guaranteed to be an integer whenever α is an integral class

$$\alpha = G + \frac{\lambda}{2} \in H^4(\tilde{Y}_4, \mathbb{Z}). \quad (2.41)$$

This is just the quantization law derived by Witten [39]. We emphasize here that the joint geometric and flux contributions to the D3-brane tadpole reflect a connection between χ and the G -flux quantization law that is embodied by (2.36). This will allow a useful consistency check on any proposed quantization rule.

2.7.1 New surfaces in $\tilde{Y}_4$

We now turn to the quantization of G -flux directly, which is to say a characterization of the odd part of λ . Because λ is always even in smooth Weierstrass models, such as the fourfold we get by deforming away the singularities of Y_4 , we expect that any odd piece of $c_2(\tilde{Y}_4)$ comes from holomorphic surfaces in $\tilde{Y}_4$ that do not survive the blow-down map $\tilde{Y}_4 \rightarrow Y_4$. Surfaces of this type include matter surfaces, which have the form ‘ $C \times \Sigma$ ’ for some degenerating root C , as well as the surfaces (or rather formal linear combinations of surfaces) that can be described in the spectral divisor formalism [10]²⁸. The latter are of the form $\mathcal{S} - S_0$ for $\mathcal{S}$ a divisor inside $\mathcal{C}_{\text{Tate}}$ and S_0 the usual subtraction piece that ensures $\mathcal{S} - S_0$ is trivial after the blow-down map.

²⁸We do not claim to have identified a distinguished linearly independent basis for the set of surfaces under consideration.

Given $\tilde{S}$, we will use the notation $\tilde{S}_{\mathcal{S}}$ to denote a surface class in $\tilde{Y}_4$ that is constructed via the $\mathcal{S} - S_0$ procedure from a divisor $\mathcal{S}$ in $\mathcal{C}_{\text{Tate}}$.

We will later be interested in computing intersections of these new surfaces $\tilde{S}_{\mathcal{S}} \cdot_{\tilde{Y}_4} \tilde{S}_{\mathcal{S}}$. In principle, we can use our knowledge of the local geometry near singularities of Y_4 to do this. For simplicity, let us suppose that our Y_4 has a single surface of $SU(5)_{\text{GUT}}$ singularities and no others. In that case, the local intersection data (2.16) from the dP_9 picture tells us that the computation should reduce to one within the Higgs bundle spectral cover. Suppose that a divisor $\mathcal{S}$ in $\mathcal{C}_{\text{Tate}}$ restricts to a divisor s inside $\mathcal{C}_{\text{Higgs}}$. In the same way that we computed the D3-brane tadpole contribution (2.34), we can use the local intersection data (2.16) to obtain

$$\tilde{S}_{\mathcal{S}_1} \cdot_{\tilde{Y}_4} \tilde{S}_{\mathcal{S}_2} = -s_1 \cdot_{\mathcal{C}_{\text{Higgs}}} s_2 - (p_* s_1) \cdot_{S_{\text{GUT}}} (p_* s_2), \quad (2.42)$$

where p is the projection map

$$p : \mathcal{C}_{\text{Higgs}} \rightarrow S_{\text{GUT}}. \quad (2.43)$$

The first term is familiar from the discussion preceding (2.34). The new term involving $p_* s_i$'s accounts for the fact that $\mathcal{S}$ need not be traceless. It effectively computes the contribution from the '1' in (2.16).

We can tabulate a few interesting results here. When $\mathcal{C}_{\text{Higgs}}$ is generic, which means among other things that it *does not split*, the divisors of $\mathcal{C}_{\text{Higgs}}$ are of the form

$$\sigma \cdot_X \mathcal{C}_{\text{Higgs}} \quad \text{and} \quad p^* \Sigma, \quad (2.44)$$

where Σ is a curve in S_{GUT} , $X = \mathbb{P}(\mathcal{O} \oplus K_{S_{\text{GUT}}})$ is the usual ambient space in which $\mathcal{C}_{\text{Higgs}}$ is embedded for ease of study, and σ is a section of the $\mathbb{P}^1$ -fibration X that transforms trivially as we move along S_{GUT} ²⁹. Given this, we know that any divisor $\mathcal{S}$ must restrict to a combination

$$\mathcal{S} \rightarrow n(\sigma \cdot_X \mathcal{C}_{\text{Higgs}}) + p^* \Sigma. \quad (2.45)$$

This allows us to give general formulas. Writing c_1 as shorthand for $c_1(S_{\text{GUT}})$ as usual and denoting the class of $\mathcal{C}_{\text{Higgs}}$ inside X as

$$\mathcal{C}_{\text{Higgs}} = 5\sigma + \pi_X^* \eta, \quad \pi_X : X \rightarrow S_{\text{GUT}} \quad (2.46)$$

we have that

$$\begin{aligned} \left(\tilde{S}_{\sigma \cdot \mathcal{C}_{\text{Higgs}}} \right)^2 &= -30c_1^2 + 11c_1\eta - \eta^2 \\ \left(\tilde{S}_{\sigma \cdot \mathcal{C}_{\text{Higgs}}} \right) \cdot_{\tilde{Y}_4} \left(\tilde{S}_{p^* \Sigma} \right) &= -6\Sigma \cdot_{S_{\text{GUT}}} (\eta - 5c_1) \\ \left(\tilde{S}_{p^* \Sigma_1} \right) \cdot_{\tilde{Y}_4} \left(\tilde{S}_{p^* \Sigma_2} \right) &= -30\Sigma_1 \cdot_{S_{\text{GUT}}} \Sigma_2. \end{aligned} \quad (2.47)$$

We have abused notation somewhat here. By $\tilde{S}_{\sigma \cdot_X \mathcal{C}_{\text{Higgs}}}$ we mean $\tilde{S}_{\mathcal{S}}$ for a divisor $\mathcal{S}$ that restricts to $\sigma \cdot_X \mathcal{C}_{\text{Higgs}}$ in the Higgs bundle spectral cover.

²⁹This is the conventional notation introduced to the F-theory literature by [1].

2.7.2 G -flux quantization

A rule for G -flux quantization is naturally motivated from the study of Higgs bundles. In the latter setting, the object γ that represents the restriction of $\mathcal{G}$ to $\mathcal{C}_{\text{Higgs}}$ satisfies the rule

$$\gamma + \frac{r}{2} \in H^2(\mathcal{C}_{\text{Higgs}}, \mathbb{Z}), \quad (2.48)$$

where r is the ramification divisor of the covering

$$p : \mathcal{C}_{\text{Higgs}} \rightarrow S_{\text{GUT}}. \quad (2.49)$$

The divisor r naturally descends from a divisor class $\tilde{r}$ inside $\mathcal{C}_{\text{Tate}}$ obtained by taking the ramification divisor of the covering $\mathcal{C}_{\text{Tate}} \rightarrow B_3$ and removing the component along the surface of $SU(5)_{\text{GUT}}$ singularities. The proposed quantization rule is now that $\mathcal{G}$ should be chosen so that

$$\mathcal{G} + \frac{\tilde{r}}{2} \in H^4(\mathcal{C}_{\text{Tate}}, \mathbb{Z}). \quad (2.50)$$

This is equivalent to the claim that the combination $\tilde{r} - r_0$ captures the odd part of $c_2(\tilde{Y}_4)$ where r_0 represents our usual subtraction term. In equations, the conjecture is

$$\frac{1}{2} [\lambda(\tilde{Y}_4) - (\tilde{r} - r_0)] \in H^4(\tilde{Y}_4, \mathbb{Z}). \quad (2.51)$$

As usual, this is something that has been conjectured, at least in some form, in the setting of Heterotic/F-theory duality. In addition to describing everything intrinsically in terms of F-theory and the geometry of $\tilde{Y}_4$, we would like to perform a consistency check that we do not believe has been studied previously using duality or otherwise. This check has to do with the connection between λ and the Euler character (2.36), which tells us that the odd part of the former is related to the part of the latter that fails to be divisible by 24. More specifically, suppose we compute χ and λ in a smooth deformation of Y_4 and follow the resulting classes as we move back to the singular Y_4 . The result of this is the naive χ and λ that we would obtain by blindly applying formulae for smooth Weierstrass models without properly accounting for the singularity structure. If we pull these back to the resolved geometry $\tilde{Y}_4$, they will differ from the actual $\chi(\tilde{Y}_4)$ and $\lambda(\tilde{Y}_4)$ there by shifts

$$\delta\chi = \chi(\tilde{Y}_4) - \chi(Y_4), \quad \delta\lambda = \lambda(\tilde{Y}_4) - \lambda(Y_4). \quad (2.52)$$

The general relation (2.36) tells us that³⁰

$$\frac{\delta\chi}{12} + \frac{1}{4}(\delta\lambda)^2 \in \mathbb{Z}. \quad (2.53)$$

The odd part of $\delta\lambda$ is thus directly connected to the failure of χ to be divisible by 12. In our $SU(5)_{\text{GUT}}$ example, we conjecture that the odd part of $\delta\lambda$ is given by $(\tilde{r} - r_0)$ so the non-integer part of $(\delta\lambda)^2/4$ can be computed using the rules of section 2.7.1

$$\tilde{S}_{(\tilde{r}-r_0)} \cdot_{\tilde{Y}_4} \tilde{S}_{(\tilde{r}-r_0)} = -15(32c_1^2 - 25\eta c_1 + 5\eta^2). \quad (2.54)$$

³⁰We use the fact that $\lambda(Y_4)$ is an even class.

We can now compare this to the formulae conjectured in [30] for the shift in Euler character due to the $SU(5)_{\text{GUT}}$ singularity. This leads to the result

$$\delta\chi = -15(488c_1^2 - 211\eta \cdot c_1 + 23\eta^2). \quad (2.55)$$

From this, it is easy to verify that

$$\frac{\delta\chi}{12} + \frac{1}{4}(\delta\lambda)^2 \in \mathbb{Z}. \quad (2.56)$$

We believe this represents nice evidence in favor of the conjectures of [30] relating to the Euler character and the conjectured odd part of λ .

Note that the conjecture for $\delta\lambda$ will take a slightly different form when $U(1)$ s are engineered. This is because a split Tate divisor $\mathcal{C}_{\text{Tate}}$ will lead to a split Higgs bundle and what enters the quantization condition for γ in that case are the ramification divisors r_n of the different components of $\mathcal{C}_{\text{Higgs}}$. We therefore expect that the shift in Euler character $\delta\chi$ will be similarly affected.

The above story also provides more evidence that surfaces of singularities (and likely the local behavior of the Tate divisor) play a dominant role in determining the Euler character. If we have multiple non-intersecting surfaces of singularities in Y_4 , we expect a local Higgs bundle description (or something like it) near each one and each will carry with it a new potentially odd contribution to λ . Shifts in λ go hand in hand, according to (2.36), with shifts in the Euler character.

3 M5 instantons, G -flux, and charged couplings

We now move to a simple application of our description of G -flux: the influence of flux on M5 instantons. More generally, we would like to know, for a given choice of Calabi-Yau fourfold and G -flux, which divisors can support M5-instantons that generate nonzero couplings and whether any of those couplings involve charged fields, by which we mean states that arise from M2-branes that wrap degenerate cycles.

For starters, recall that in F-theory we are really studying backgrounds of the type $\mathbb{R}^{3,1} \times B_3$ in IIB string theory with axio-dilaton that varies over B_3 . Nonperturbative superpotential corrections can arise not just from D3-instantons [42–47] wrapped on divisors in B_3 but also from gaugino condensation [44, 48, 49] which may occur along divisors in B_3 that support non-Abelian gauge groups. For applications to F-theory GUT model-building the visible sector will not undergo gaugino condensation but, as we will review in the next section, hidden sector gauge groups can often appear and the corresponding gaugino condensates can play an important role in stabilizing Kähler moduli.

In this paper, we adopt the perspective of M-theory to describe D3-instantons as a limit of M5-instantons that are wrapped on vertical divisors of the elliptically fibered Calabi-Yau fourfold Y_4 . These vertical divisors are complex threefolds D of the form π^*S_2 where S_2 is a

holomorphic divisor in B_3 . In the F-theory limit, where the volume of the elliptic fiber of Y_4 is taken to zero, these M5-instantons descend to D3-instantons wrapped on S_2 .

In general, there are two reasons that an M5-instanton wrapping a vertical divisor fails to contribute a superpotential coupling W_{inst}

- $W_{inst} = 0$ if there are not precisely two fermion zero modes on the M5-brane worldvolume
- W_{inst} may be zero if the restriction of the G -flux to the M5 worldvolume is nontrivial

Before proceeding, let us say a few more words about the second point. The influence of G -flux is evident already from the fact that the complexified Kähler modulus³¹ $T = \int_D (\text{vol}_D + iC_6)$ shifts under gauge transformations of the M-theory 3-form C_3 when the restriction of G_4 to the M5 is nontrivial. To see why, recall that under $C_3 \rightarrow C_3 + d\Lambda_2$ the 6-form C_6 shifts as

$$C_6 \rightarrow C_6 + \frac{1}{2}\Lambda_2 \wedge G_4. \quad (3.1)$$

This is a general consequence of the gauge invariance of $G_7 = *G_4$ and the Bianchi identities

$$dG_7 = -\frac{1}{2}G_4 \wedge G_4, \quad (3.2)$$

which implies that

$$G_7 = dC_6 - \frac{1}{2}C_3 \wedge G_4. \quad (3.3)$$

As a result, the complexified Kähler modulus $T = \int_D (\text{vol}_D + iC_6)$ and hence the classical action of the M5 shifts under gauge transformations of the M-theory 3-form $C_3 \rightarrow C_3 + d\Lambda_2$ according to

$$\delta T = \frac{i}{2} \int_D \Lambda_2 \wedge G_4. \quad (3.4)$$

An interesting special case of this arises when we have a $U(1)$ symmetry of the type considered in section 2.3, by which we mean a harmonic $(1,1)$ -form ω on which we can reduce C_3 as $C_3 = A_1 \wedge \omega$.³² A gauge transformation with respect to this $U(1)$ corresponds to the special choice $\Lambda_2 = \phi \wedge \omega$ with ϕ the product of a constant 0-form on the Calabi-Yau fourfold and a nontrivial function of the Minkowski directions. In that case, the classical M5-instanton action e^{-T} transforms as though it carries charge

$$q_{M5} = \frac{1}{2} \int_D \omega \wedge G_4. \quad (3.5)$$

Any coupling generated by the M5-instanton must be gauge invariant under all potential gauge transformations of C_3 . When G_4 restricts trivially to the M5 worldvolume this is not a problem

³¹ C_6 is the magnetic dual of the 3-form gauge potential C_3 in M-theory.

³²Here A_1 gives a 3-dimensional gauge field on $\mathbb{R}^{2,1}$ in M-theory that descends to a 4-dimensional gauge field on $\mathbb{R}^{3,1}$ in the F-theory limit when ω has exactly 1 leg on the elliptic fiber.

and, in that case, the fermi zero mode structure should represent the only obstacle to obtaining a nonzero coupling. When G_4 has nontrivial restriction, we will argue that this can be accomplished in some cases by inserting wrapped M2-brane states³³. In those cases the M5 generates a coupling, again provided the fermi zero mode structure is appropriate. When this is not possible, the M5 should not generate any coupling.

Note that invariance under bulk gauge transformations of C_3 is a stronger statement than invariance under $U(1)$ symmetries of the type described in section 2.3. The full condition of bulk gauge invariance should incorporate the selection rules alluded to in [28]³⁴.

In the rest of this section we discuss each of the two obstacles to obtaining a nonzero coupling in turn. In section 3.1, we review the counting of fermion zero modes applying those results to M5-instantons in the specific geometry of [29] in section 4. Then in section 3.2 we discuss the effect of G -flux in a bit more detail, describe the impact of the G -fluxes that we discussed in section 2.5, clarify how they could in principle generate charged couplings, and demonstrate that, for the G -fluxes described in section 2.5, the restriction to M5 worldvolumes is generically trivial. This means that vertical divisors with the right number of fermi zero modes should, in the absence of additional G -fluxes with nontrivial restriction, generate uncharged couplings that can play a role in the stabilization of Kähler moduli.

3.1 Counting fermion zero modes

3.1.1 Review of fermions living on M5-instanton

To count fermion zero modes one recasts the equations of motion for fermions living on the M5-instanton as a set of equations on differential forms on the divisor D . For this purpose we recall below how world-volume fermions transform under the rotations of the normal and tangent directions. The normal bundle to the M5-brane has a product form $\mathbb{R}^3 \times \mathcal{N}$, where $\mathbb{R}^3$ stands for external space³⁵ and $\mathcal{N}$ is the line bundle describing one complex normal direction inside Y_4 . Let us introduce complex coordinates z^i , $i = 1, 2, 3$ along the divisor D and the complex coordinate w normal to D inside Y_4 .

The fermions $\theta = \begin{pmatrix} \theta_\alpha^{A+} \\ \theta_\alpha^{A-} \end{pmatrix}$ living on the M5-brane transform in the representation $\mathbf{4} \otimes \mathbf{2} \otimes \mathbf{2}$ under $Spin(6) \times SO(3) \times SO(2)$. Here $A = 1, 2$ is a spinor under external $SO(3)$, the $+$ ($-$) stands for a chiral(anti-chiral) spinor of $SO(2)$ and $\alpha = 1, \dots, 4$ is a chiral spinor of $Spin(6)$. Now we use the fact (see for example [42]) that the bundle S^+ of chiral spinors on a Kähler manifold of complex dimension three is isomorphic to the bundle

$$\left(\Omega^{(0,0)} \otimes K^{\frac{1}{2}} \right) \oplus \left(\Omega^{(0,2)} \otimes K^{\frac{1}{2}} \right). \quad (3.6)$$

³³What is really happening is that the flux induces a source term for the chiral 2-form b_2 so that the vacuum solutions about which we expand are solitons that describe suitable M2s ending on the M5.

³⁴As we understand it, gauge transformations corresponding to the massive $U(1)$ s of [28] correspond to choosing Λ_2 to be of the form $\phi \wedge \tilde{\omega}$ with ϕ as above and $\tilde{\omega}$ a 2-form on Y_4 that is not necessarily harmonic.

³⁵For computation of instanton generated superpotential we work in Euclidean signature in external 3D space.

Here $\Omega^{(0,p)}$ stands for the bundle of $(0,p)$ forms. We will further use that the normal bundle on the divisor in Y_4 is isomorphic to the canonical bundle K . Recalling that θ is a section of the bundle³⁶ $S^+ \otimes K^{\frac{1}{2}} \oplus S^+ \otimes K^{-\frac{1}{2}}$, we find the following degrees of freedom. A $(0,2)$ form $a_{(2)}^w$ taking values in the canonical bundle K , a section of K , $a_{(0)}^w$, as well as a $(0,2)$ form $b_{(2)}$ and a scalar $b_{(0)}$.

Locally we write θ in terms of these degrees of freedom as follows

$$\theta = (a_{(0)}^w + a_{i\bar{j}}^w \gamma^{\bar{i}} \tilde{\gamma}^{\bar{j}}) T_w \epsilon + (b_{(0)} + b_{i\bar{j}}^w \gamma^{\bar{i}} \tilde{\gamma}^{\bar{j}}) \epsilon, \quad (3.7)$$

where the chiral spinor ϵ satisfies

$$\tilde{\gamma}^i \epsilon = 0, \quad i = 1, 2, 3, \quad T_{\bar{w}} \epsilon = 0. \quad (3.8)$$

Here $T_w, T_{\bar{w}}$ are $SO(2)$ Dirac matrices:

$$T_w T_{\bar{w}} + T_{\bar{w}} T_w = 2g_{w\bar{w}}. \quad (3.9)$$

Meanwhile, the six dimensional chiral(anti-chiral) gamma matrices $\tilde{\gamma}_i, \tilde{\gamma}_{\bar{j}}(\gamma_i, \gamma_{\bar{j}})$ have the properties

$$\gamma_{\bar{j}} \tilde{\gamma}_i + \gamma_i \tilde{\gamma}_{\bar{j}} = 2g_{i\bar{j}}, \quad (3.10)$$

where $g_{i\bar{j}}$ is Kähler metric on the divisor D .

The equation of motion for fermions reads [50]

$$\tilde{\gamma}^i \nabla_i \theta + \tilde{\gamma}^{\bar{i}} \nabla_{\bar{i}} \theta - \frac{1}{8} T_{\bar{w}} \tilde{\gamma}^{\bar{i}j\bar{k}} G_{i\bar{j}k}^{\bar{w}} \theta - \frac{1}{8} T_w \tilde{\gamma}^{ij\bar{k}} G_{ij\bar{k}}^w \theta = 0. \quad (3.11)$$

Here the covariant derivatives $\nabla_j, \nabla_{\bar{j}}$ include the connection on the bundle of chiral $\text{Spin}(6)$ spinors as well as connection on the spin bundle derived from the normal bundle $\mathcal{N}$.

Using (3.7) we find from (3.11) the following set of equations³⁷:

$$\begin{aligned} \partial_{[\bar{i}} b_{\bar{j}\bar{k}]} &= 0 \\ 4\partial^{\bar{j}} b_{\bar{j}\bar{k}} + \partial_{\bar{k}} b_{(0)} &= 0 \\ D_{[\bar{m}_1} a_{\bar{m}_2 \bar{m}_3]}^w &= 0 \\ 4D^{\bar{j}} a_{\bar{j}\bar{k}}^w + D_{\bar{k}} a_{(0)}^w &= -G^{\bar{i}\bar{j}}{}_{\bar{k}}{}^w b_{\bar{i}\bar{j}}. \end{aligned} \quad (3.12)$$

In the equations (3.12) the covariant differentials include connection on the canonical bundle and we used the primitivity condition on G -flux.

There are always universal modes counted by $2 \times h^{(0,0)}(D) = 2$, where the factor of 2 accounts for two spinor indices in R^3 . Non-universal modes without G -flux are counted by

$$2 \times [h^{(0,1)}(D) + h^{(0,2)}(D) + h^{(0,3)}(D)]. \quad (3.13)$$

It is further known [45, 46, 51, 52] that modes counted by $h^{(0,2)}(D)$ may be lifted by G -flux since the G -flux couples to them directly in (3.12). An especially simple case of such lifting occurs [45] if $h^{(0,1)}(D) = h^{(0,3)}(D) = 0$ and $h^{(0,2)}(D) = 1$.

³⁶Here we are not writing explicitly spinor index in $\mathbb{R}^3$.

³⁷ $X_{[\bar{i}_1 \dots \bar{i}_p]} = \frac{1}{p!} (X_{\bar{i}_1 \dots \bar{i}_p} \pm \text{permutations})$

3.1.2 $h^{(0,k)}(D)$ for $D = p^*S_2$

To study the zero mode structure of an M5 then we must be able to compute the Hodge numbers $h^{(0,k)}(D)$ for a vertical divisor $D = p^*S_2$. From the connection to D3-instantons, one intuitively expects that these should be related to cohomologies of the normal bundle of S_2 inside B_3 . We describe this relation for the special case where S_2 is a Hirzebruch surface $\mathbb{F}_n$ or del Pezzo surface dP_m in the following.

Let $P : D \rightarrow S_2$ be the elliptic fibration arising as a restriction of $\pi : Y_4 \rightarrow B_3$. We now use the Leray spectral sequence to relate the cohomology of the divisor $D = \pi^*S_2$ to the cohomology of S_2 . We start from E_2^{ij} with $i = 0, 1, 2$ and $j = 0, 1$, where each entry E_2^{ij} is given by $H^i(S_2, R^j P_* \mathcal{O}_D)$

$$E_2^{ij} : \begin{array}{ccc} H^0(S_2, R^1 P_* \mathcal{O}_D) & H^1(S_2, R^1 P_* \mathcal{O}_D) & H^2(S_2, R^1 P_* \mathcal{O}_D) \\ H^0(S_2, \mathcal{O}_{S_2}) & H^1(S_2, \mathcal{O}_{S_2}) & H^2(S_2, \mathcal{O}_{S_2}) \end{array} \quad (3.14)$$

We will always take S_2 to be Fano $\mathbb{F}_n$ or del Pezzo dP_m so

$$H^1(S_2, \mathcal{O}_{S_2}) = H^2(S_2, \mathcal{O}_{S_2}) = 0. \quad (3.15)$$

Then $E_\infty^{i,j} = E_2^{i,j}$ and

$$h^{(0,0)}(D) = 1, \quad h^{(0,k)}(D) = h^{k-1}(S_2, R^1 P_* \mathcal{O}_D), \quad k = 1, 2, 3. \quad (3.16)$$

Now the key step is to relate $R^1 P_* \mathcal{O}_D$ with the normal bundle of S_2 in B_3 :

$$R^1 P_* \mathcal{O}_D = N_{S_2|D} = K_{B_3}|_{S_2} = K_{S_2} \otimes N_{S_2|B_3}^{-1}. \quad (3.17)$$

So that using Serre duality

$$\begin{aligned} h^{(0,1)}(D) &= h^2(S_2, N_{S_2|B_3}) \\ h^{(0,2)}(D) &= h^1(S_2, N_{S_2|B_3}) \\ h^{(0,3)}(D) &= h^0(S_2, N_{S_2|B_3}). \end{aligned} \quad (3.18)$$

The absence of non-universal fermi zero modes therefore corresponds to the vanishing of suitable cohomologies of the normal bundle of S_2 inside B_3 .

3.2 G -flux, the M5 path-integral, and charged couplings

Let us now turn to the effect of G -flux on the M5-instanton partition function and the corresponding generation of charged couplings. For this, we only really need to consider the topological couplings:

$$S_{M5} \sim i\tau_{M5} \int_{M5} \left(C_6 - \frac{1}{2} b_2 \wedge G_4 + \dots \right). \quad (3.19)$$

Here, b_2 is the chiral 2-form while C_6 and G_4 are restrictions of the corresponding 11d fields to the M5 worldvolume. The form of the second coupling is essentially determined by the fact that b_2 and C_6 shift under the bulk gauge transformation $C_3 \rightarrow C_3 + d\Lambda_2$ according to

$$b_2 \rightarrow b_2 + \Lambda_2, \quad C_6 \rightarrow C_6 + \frac{1}{2}\Lambda_2 \wedge G_4. \quad (3.20)$$

Once the second coupling in (3.19) is included the sum is nicely gauge invariant. One useful feature of (3.19) is that it makes clear how the physics of our M5-instanton is impacted by a nontrivial G -flux: the G -flux introduces a source for the 2-form b_2 ³⁸. This, in turn, means that classical solutions for b_2 correspond to solitons that we should interpret as M2 branes ending on the M5 [53]. In the case at hand, our M5 is Euclidean so the M2s will have two spatial directions parallel and a timelike direction transverse to the M5.

In principle, if we know the G -flux then we can determine exactly what M2-brane configuration will be described by the solitonic solutions to the equations of motion for b_2 . It is easier to think not in terms of solitons of the M5 worldvolume theory, however, but to instead introduce M2s directly into the game in order to cancel the flux-induced tadpole. Each M2 has a worldvolume coupling to C_3 along with a boundary coupling to the 2-form b_2 [54, 55]

$$S_{M2} \sim i\tau_2 \left(\int_{M2} C_3 - \int_{M2 \cap M5} b_2 + \dots \right). \quad (3.21)$$

The idea now is to include M2-branes in such a way that the coupling to b_2 in (3.21) exactly cancels the one induced by the G -flux in (3.19). The resulting action for b_2 will be the ordinary one that we would have if no G -flux were present at all. Note that the individual gauge invariance of (3.19) and (3.21) guarantees that the sum is gauge invariant as well. When the source terms for b_2 in (3.19) cancel off the boundary couplings in (3.21), then this guarantees that the net charge of the individual M2-branes under a bulk gauge transformation, as determined by the shifts in $i\tau_2 \int_{M2} C_3$, is precisely canceled by the charge of the M5-instanton, as determined by the shift in $i\tau_5 \int_{M5} C_6$. Invariance under bulk gauge transformations is thus automatic and hence invariance under any $U(1)$ s of the type considered in section 2.3 is automatic as well.

To see which M2-branes we have to add, let us rewrite the G_4 -dependent part of (3.19) as

$$-\frac{i}{2}\tau_5 \int_{M5} b_2 \wedge G_4 = -\frac{i}{2}\tau_5 \int_{PD_{M5}[G_4]} b_2, \quad (3.22)$$

where $PD_{M5}[G_4]$ is the Poincare dual of G_4 inside the M5. A collection of M2-branes that cancels this is one for which the curves $\Sigma_i = M5 \cap M2_i$ satisfy

$$\sum_i \tau_2 \Sigma_i = -\frac{1}{2}\tau_5 PD_{M5}[G_4]. \quad (3.23)$$

³⁸We are grateful to S. Sethi for emphasizing to us the important role played by the chiral 2-form and for helpful discussions about this point.

The M2s that participate in our instanton-generated coupling are determined by the Poincare dual of the G -flux inside the M5 worldvolume. Note that the gymnastics we have just described is nothing more than a procedure for describing how to associate a collection of Wilson surface observables of the type considered in [26] with a charged coupling involving 4-dimensional fields associated to a particular collection of wrapped M2's. Our description in terms of (3.23) is particularly convenient because our construction of G -flux is one in which we essentially define it by the Poincare dual. Before going too far, though, we have to be somewhat careful about normalization. When we build G -flux as in section 2.5 by specifying an integer combination of holomorphic surfaces $\mathcal{G}$ inside the Tate divisor, that flux is quantized according to

$$G_{\text{Tate}} + \frac{1}{2}c_2(Y_4) \in H^4(Y_4, \mathbb{Z}) . \quad (3.24)$$

In terms of the 3-form C_3 appearing in the action above, this G_{Tate} is then really³⁹

$$G_{\text{Tate}} = \frac{2\pi}{\tau_2} dC_3 , \quad (3.25)$$

so that, using the M2 and M5 tensions $\tau_2 = (2\pi)^{-2}$ and $\tau_5 = (2\pi)^{-5}$, we have that

$$\sum_i \Sigma_i = -\frac{1}{2} PD_{M5}[G_{\text{Tate}}] . \quad (3.26)$$

Up to a factor $(-\frac{1}{2})$, then, the surface that we use to describe the G -flux tells us precisely which wrapped M2-brane states participate in the instanton-generated coupling.

We still have to discuss the nature of $PD_{M5}[G_{\text{Tate}}]$ to get our prescription for determining the charged coupling generated by an M5 instanton. Before getting to that, though, we should say something about the puzzling factor of $-\frac{1}{2}$. This seems to be the statement that G -flux can induce a half-integral M2 charge on the M5 worldvolume. We are not aware of any reason for this charge failing to be integer quantized and, correspondingly, do not know of any ways to suitably cancel it. It seems, then, that a vertical divisor π^*S_2 to which the restriction of G_{Tate} is an odd class simply cannot house an M5 instanton. We would be very interested in a better understanding of this issue but unfortunately do not have anything substantive to say about it at the moment.

3.2.1 The nature of $PD_{M5}[G_{\text{Tate}}]$

Let us now consider how the 2-cycles specified by $PD_{M5}[G_{\text{Tate}}]$ can be determined from the geometric construction of G -fluxes in section 2.5. The G -flux there was actually constructed as a difference

$$G_{\text{Tate}} = \mathcal{G} - G_0 , \quad (3.27)$$

³⁹This can be determined in the usual way by looking at the phase of an M2 as it moves around in a flux background [39].

where $\mathcal{G}$ is a surface inside the Tate divisor, $\mathcal{C}_{\text{Tate}}$, and G_0 is a class in Y_4 that we must subtract to ensure that G_{Tate} is orthogonal to horizontal and vertical divisors. The restriction of $\mathcal{C}_{\text{Tate}}$ to an M5 worldvolume will be a spectral surface $\mathcal{C}_{\text{Tate}}|_{M5}$ in that worldvolume while $\mathcal{G}$ will be a curve $\Sigma_{\mathcal{G}}$ that sits inside it. The restriction $G_0|_{M5}$ is also a curve Σ_{G_0} inside the M5 but will not lie in the spectral surface $\mathcal{C}_{\text{Tate}}|_{M5}$. The object $PD_{M5}[G_{\text{Tate}}]$ is then given by the difference

$$PD_{M5}[G_{\text{Tate}}] = \Sigma_{\mathcal{G}} - \Sigma_{G_0} \quad (3.28)$$

in the resolved geometry.

To get a better feel for $PD_{M5}[G_{\text{Tate}}]$, let us look at its intersections with divisors inside the M5. Any intersection of $\Sigma_{\mathcal{G}}$ with a horizontal or vertical divisor will be canceled by a corresponding contribution from Σ_{G_0} . The M5 may however contain additional divisors upon resolution, if it contains a curve of singularities $\Sigma_{\text{sing},i}$. Associated with that curve of singularities is a matter surface ‘ $\Sigma_{\text{sing},i} \times C_i$ ’ that will represent a divisor in the M5 where the intersection with $\Sigma_{\mathcal{G}}$ is not canceled by a contribution from Σ_{G_0} ⁴⁰. Divisors of this type represent the only ones in M5 over which G_{Tate} can give a nonzero integral. If the M5 does not contain any curves of singularities, then, G_{Tate} will have a trivial restriction to its worldvolume. Given our geometric description of G_{Tate} , the curve $PD_{M5}[G_{\text{Tate}}]$ is actually easy to describe. We have

$$PD_{M5}[G_{\text{Tate}}] = \sum_i n_i C_i, \quad (3.29)$$

where C_i is a degenerate cycle and n_i is the intersection number $\Sigma_{\mathcal{G}} \cdot \mathcal{C}_{\text{Tate}}|_{M5} \cdot \Sigma_{\text{sing},i}$ with $\Sigma_{\text{sing},i}$ the curve of singularities inside $\mathcal{C}_{\text{Tate}}|_{M5}$ along which C_i degenerates.

We can say all of this in a much simpler way. We constructed G_{Tate} so that it would correspond to a (2,2)-form in the resolved fourfold that has nonzero integral only over matter surfaces, that is to say surfaces comprised of curves inside Y_4 and 2-cycles that degenerate along those curves. The restriction of G_{Tate} to an M5 instanton will therefore only be nontrivial if the instanton contains at least one of these matter surfaces in its entirety since otherwise there will not be surfaces over which we can integrate G_{Tate} to get a nonzero answer. Now, the M5 wraps a divisor in Y_4 while the singularities of interest lie along curves in Y_4 ⁴¹. The generic situation, then, is for the M5 to intersect a curve of singularities only in isolated points. This means

⁴⁰Note that $\Sigma_{\mathcal{G}}$ will generically intersect $\Sigma_{\text{sing},i}$ inside the M5 even though they are both codimension 2. This is because both curves actually sit inside the spectral surface $\mathcal{C}_{\text{Tate}}|_{M5}$, with respect to which each is codimension 1. Let us also emphasize that it is not enough for the M5 to contain isolated points of singularities. Even though $\Sigma_{\mathcal{G}}$ and an isolated point singularity both sit inside $\mathcal{C}_{\text{Tate}}$, they will generically not intersect. Said differently, an isolated point singularity inside the M5 does not give rise to a new divisor in the M5 upon resolution over which we can integrate G_{Tate} .

⁴¹It is true that the M5 intersects surfaces of singularities (like the one above S_{GUT}) in curves. G -fluxes that can have nonzero integral over these singularities are precisely fluxes that break the corresponding non-Abelian gauge group, i.e. worldvolume flux associated to the stack of 7-branes. Hypercharge flux is an example of such a flux so this will be an important issue for model-building applications (though trivializable hypercharge fluxes should not pose a problem because the curves in question are homologically nontrivial in B_3 as the intersection of a surface $S_{\text{GUT}} \subset B_3$ with the divisor $S_2 \subset B_3$ that is the base of the M5-instanton). For simplicity, we focus only on bulk G -fluxes in this paper.

that bulk G -fluxes of the type described in section 2.5 generically have trivial restriction to M5 instantons. Based on our previous discussions, this means that the topological term is absent from the M5-instanton action and there is no source term for b_2 . This removes one obstacle to obtaining a nonzero coupling, leaving only the structure of fermi zero modes to be checked. The classical action of such an M5-instanton is invariant under all bulk gauge transformations so this is consistent with all of the usual $U(1)$ selection rules.

4 Instanton induced superpotential: an Example

In this paper so far we have developed the general framework for studies of global G -fluxes and $U(1)$ symmetries, and gave criteria when instanton contributions to the superpotential arise. We will next apply these in a concrete example, with the base threefold that we constructed in [29]. The construction of B_3 in our earlier work [29] relied on a chain of blowups along a nodal curve in $\mathbb{P}^3$, and the final threefold was obtained after a flop transition. There is a slightly more elegant way to reach this geometry, which does not require a flop, that effectively reverses the order of the blow-ups. We only give a heuristic description of this approach in the main text below but a detailed description can be found in appendix A.

4.1 Divisors that contribute superpotential couplings

The first obstruction to obtaining superpotential couplings from an M5-instanton is the fermi zero mode structure. As we discussed in section 3.1, obtaining the right structure for a vertical divisor of the form π^*S_2 requires the cohomology groups $H^p(S_2, N_{S_2/B_3})$ to be trivial for $p = 0, 1, 2$. As a start, then, we should look for divisors S_2 in B_3 that are not moveable so that at least the $p = 0$ condition is satisfied. More precisely, the relevant divisors $D = \pi^*S_2$ have to satisfy $h^{(0,3)}(D) = h^0(S_2, N_{S_2/B_3}) = 0$. Before listing candidate divisors in the threefold of [29] that satisfy this criterion, though, let us quickly review its basic structure.

We can obtain the threefold of [29] by starting with $\mathbb{P}^3$ and picking out a distinguished plane cubic curve C_{nodal} with a node. Because of the node, we can think of C_{nodal} as a pinched torus obtained by gluing together the north and south poles of a $\mathbb{P}^1$. The construction of B_3 now proceeds in two steps:

- First, blow up $\mathbb{P}^3$ at the node

This step separates the two branches of C_{nodal} so that its proper transform, C'_{nodal} , is a smooth $\mathbb{P}^1$. The exceptional divisor of this blow-up, E , is a $\mathbb{P}^2$ that meets C'_{nodal} in exactly two points. Now, onto the second step

- Second, blow up along C'_{nodal}

Because C'_{nodal} is a $\mathbb{P}^1$, the exceptional divisor D' of this second blow-up is a Hirzebruch surface $\mathbb{F}_n$ where n is determined by the normal bundle of C'_{nodal} . It is easy to show, as we do

Figure 1: Cartoon of the threefold B_3 from [29]. When D' and E' are blown down we are left with a nodal cubic that lies in the plane $H - D' - E'$. ℓ and G' label curve classes in the notation of [29].

in Appendix A, that D' is in fact an $\mathbb{F}_4$. Note that since E met C'_{nodal} in exactly two points, the preimage E' of E under the second blow-up is a dP_2 surface. Further, D' meets E' in a sum of two curves. From the perspective of E' , these are the two exceptional curves of the dP_2 . From the perspective of D' , these curves are both in the fiber class of the Hirzebruch. As a result, the two exceptional curves of E' are in fact homologous to one another in B_3 by a 3-chain that sits inside D' . This construction can easily be generalized to engineer threefolds where E' is replaced by a dP_n surfaces provided we replace C'_{nodal} with a curve that has a higher degree singularity in place of the node.

We reproduce the cartoon of B_3 from [29] in figure 4.1. From its construction, we see that there are two obvious divisors for which $h^0(S_2, N_{S_2/B_3}) = 0$

$$\begin{aligned} E' &\cong dP_2 \\ D' &\cong \mathbb{F}_4. \end{aligned} \tag{4.1}$$

All other divisors descend from polynomials in our original $\mathbb{P}^3$ that have specified intersections with the original nodal curve, C_{nodal} , or its node. Almost all of these are obviously deformable but there is one more candidate that is not: the specific hyperplane of $\mathbb{P}^3$ in which our plane curve C_{nodal} sat. The proper transform of this hyperplane is the divisor

$$H - D' - E' \cong dP_1. \tag{4.2}$$

Indeed, the hyperplane becomes a dP_1 after the first blow-up that is unaffected by the second blow-up.

So, in total, we have three candidate divisors in B_3 that can serve as the base of M5 instantons with the right zero mode structure. In our discussion above, though, we have only

argued that $H^0(S_2, N_{S_2/B_3})$ is trivial for each candidate. It remains to check $H^p(S_2, N_{S_2/B_3})$ for $p = 1, 2$. We do this in Appendix A with the following results. All of the required cohomologies vanish for E' , $H - D' - E'$, and D' so that the fermi zero mode structure is exactly right in each of these cases.

Let us turn now to the restriction of G -flux. In general, D' and $H - D' - E'$ will not contain any entire curves of singularities. As discussed in section 3.2, this means that the G -flux used to engineer a chiral spectrum restricts trivially. Provided no other G -fluxes have nontrivial restriction, both π^*D' and $\pi^*(H - D' - E')$ can be expected to yield superpotential couplings that can be used for Kähler moduli stabilization.

What about E' ? This divisor is special because it is exactly the GUT divisor on which the 7-branes are wrapped in $SU(5)_{\text{GUT}}$ models based on the geometry of [29]. Any M5 wrapping π^*E' will therefore contain several curves of singularities. The restricted G -flux will be nontrivial in general so the M5 wrapping π^*E' (which is really just capturing the contribution of a gauge theory instanton of the 7-brane worldvolume theory) will not play any role in Kähler moduli stabilization⁴².

4.1.1 Any hope for Kähler moduli stabilization?

Since only two divisors in B_3 can lead to superpotential corrections from M5-instantons, it may be difficult to stabilize Kähler moduli. This is because we have 3 Kähler moduli, which can be parametrized by writing the divisor corresponding to the Kähler form as

$$J = mH - aE' - bD'. \quad (4.3)$$

Explicitly, the Kähler cone is easy to determine and corresponds to m , a , and b satisfying

$$a > 2b > 0, \quad m > a + b. \quad (4.4)$$

While it is not always true that we need as many nonperturbative terms in the superpotential as we have Kähler moduli [56], we believe that instanton corrections from D' and $H - D' - E'$ in the threefold B_3 are not enough by themselves to stabilize everything. Fortunately, it might be possible to engineer a little help from an unexpected place. The geometry of [29] admits a nice divisor class that does not intersect the GUT divisor $S_{\text{GUT}} = E'$. This class, which we call H because it descends from a hyperplane in $\mathbb{P}^3$ that does not contain the node of C_{nodal} , is somewhat like the divisor that supports the other E_8 in models with Heterotic dual. If we can manage to engineer a hidden sector gauge group on H that undergoes gaugino condensation, the resulting superpotential correction can be used as the third nonperturbative term that we need to stabilize all of the Kähler moduli.

4.1.2 Hidden Sector

We therefore propose that models based on the geometry of [29] should, in the future, realize a hidden sector on H and study the interplay of global fluxes with that sector to determine the

⁴²In three generation models, in fact, the Poincare dual of the G -flux inside π^*E' should be odd. In this case, we are led to expect that a single M5 wrapping π^*E' does not contribute any superpotential coupling at all.

presence or absence of gaugino condensation. Note that even the realization of a hidden sector is nontrivial; it is relatively easy to see that a generic $SU(5)_{\text{GUT}}$ model based on [29] will not have surfaces of singularities anywhere away from S_{GUT} . When trying to engineer a surface of singularities along the divisor H , one must take some care to make sure that non-minimal singularities are not introduced. We now proceed to describe one example that implements a hidden sector. In the future, we hope to report on model-building efforts that engineer hidden sectors in more realistic models while implementing the approach to $U(1)$ s and global fluxes described earlier in this paper.

To get a simple example of a hidden sector on H , let us start with the global Weierstrass models studied in [29]. There we engineered an $SU(5)_{\text{GUT}}$ singularity at $z = 0$ (on $S_{\text{GUT}} = E'$) by specifying a rather general Tate form. Expressed in the notation of [29, 57] this is

$$y^2 = x^3 + z^5 g_5 + \frac{1}{48} f_3 x z^3 + h x y - \frac{1}{4} H x^2 z + \frac{1}{12} q z^2 y. \quad (4.5)$$

Here, the global sections g_5, f_3, h, H, q are rescalings of a_m in the global Tate model. We use $P = -3Hq^2 - 3hf_3q + 2g_5h^2$ to denote the section that gives rise to the codimension two enhancement to $SU(6)$. The holomorphic sections on B_3 that we use to build the sections in (4.5) are

Holomorphic Section	Divisor class
Z_4	H
$Z_{1,2}$	$(H - E') + E' = H$
Z_3	$(H - D' - E') + (D' + E') = H$
$W_{1,2,3}$	$H - E'$
W_4	$3H - D' - 2E'$
V_1	$(3H - D' - 2E') + E' = 3H - D' - E'$
V_0	$H - D' - E'$
z	E'

(4.6)

We actually never use $Z_{1,2,3}$ or V_1 explicitly. The divisors defined by each of these are reducible which indicates that each is the product of more elementary sections. Consider the ansatz for the section h of (4.5) which is in the class $4H - D' - 2E'$

$$h = (Z_4 + zP_1(W_1, W_2, W_3))(W_4 - (Z_4 + zP_1(W_1, W_2, W_3))V_0A_1(W_1, W_2, W_3)), \quad (4.7)$$

where A_1 and P_1 are linear polynomials in W_i , $i = 1, 2, 3$ ⁴³. Making the same choices otherwise for P, H, q and $f_3 = 0$ and $g_5(P, H, q)$ determined in terms of the sections P, H, q as in example 1 of appendix E in [29], we arrive at an elliptic fibration, which automatically realizes the $SU(5)_{\text{GUT}}$ singularity at $z = 0$ with matter curves and Yukawas.

To determine whether there is any nonabelian gauge enhancement along the divisor H , we expand the discriminant around $Z_4 = 0$. For $P_1 = 0$ the discriminant has vanishing order

⁴³Note that in the examples in appendix E of [29] we considered a slightly less general ansatz, where $P_1 = 0$.

$\Delta \sim Z_4^{12}$ and in fact, can be brought into non-minimal Tate form, which is disfavored. We therefore consider P_1 a generic polynomial in $W_{1,2,3}$, which yields

$$\Delta = Z_4^6 \tilde{\Delta} + O(Z_4^7), \quad (4.8)$$

where $\tilde{\Delta}$ has vanishing order in the other sections of the order z^{12} , V_0^6 , W_i^6 and W_4^0 ⁴⁴. After shifting x, y by Z_4 , the Weierstrass form can be brought globally into Tate form for I_6 without monodromy, which is $SU(6)$ [34, 58], i.e.

$$y^2 + a_1 xy + a_3 y = x^3 + a_2 x^2 + a_4 x + a_6, \quad (4.9)$$

where

$$a_1 = O(Z_4^0), \quad a_2 = O(Z_4^1), \quad a_3 = O(Z_4^3), \quad a_4 = O(Z_4^3), \quad a_6 = O(Z_4^6). \quad (4.10)$$

So in summary this exemplifies that a non-abelian gauge group can reside on the divisor H .

4.2 Superpotential and SUSY vacua

Thus far, we have argued that geometries based on the threefold of [29] will exhibit superpotential couplings from M5-instantons wrapping $\pi^* D'$ and $\pi^*(H - D' - E')$. Provided we get help from a gaugino condensate on H , this should be enough to stabilize Kähler moduli. In this subsection, we investigate the potential that would result and show that it indeed exhibits a supersymmetric minimum.

Concretely, the superpotential of interest is

$$W_{non-pert} = \sum_{D \in \{D', H-D'-E', H\}} a_D e^{-T_D}. \quad (4.11)$$

To proceed, we should write the real parts of the T_D 's of interest in terms of the Kähler moduli. Recall that we parametrize the Kähler form as

$$J = mH - aE' - bD' \quad (4.12)$$

with the Kähler cone specified by

$$a > 2b > 0, \quad m > a + b. \quad (4.13)$$

The divisors of interest that give contributions from wrapped instantons are (4.1) and (4.2) and have volumes given by

$$\begin{aligned} \text{Re } T_{D'} &= 2b(3m - 2a - 3b) \\ \text{Re } T_{H-D'-E'} &= (m - a - b)(m + a - 5b) \\ \text{Re } T_H &= m^2 - 3b^2. \end{aligned} \quad (4.14)$$

⁴⁴We do not worry about the high degree of z in $\tilde{\Delta}$ since $z = Z_4 = 0$ admits no solutions.

Furthermore, the Kähler potential is

$$K = -2 \ln J^3 = -2 \ln (m^3 - a^3 + 6b^3 - 9mb^2 + 6ab^2) . \quad (4.15)$$

The superpotential contribution from M5 instantons is then

$$W = a_1 e^{-T_{H-D'-E'}} + a_2 e^{-T_{D'}} + a_3 e^{-T_H} + W_0 . \quad (4.16)$$

We will in the following set $a_1 = 1$ without loss of generality. The scalar potential is

$$V = e^K \left(K^{a\bar{b}} (D_a W) (D_{\bar{b}} W) - 3|W|^2 \right) , \quad (4.17)$$

where D_a is the usual Kähler covariant derivative

$$D_a f = \partial_a f + (\partial_a K) f . \quad (4.18)$$

To find SUSY vacua inside the Kähler cone, consider the three equations

$$D_{a,b,m} W = 0 . \quad (4.19)$$

In these equations, we follow convention and absorb the axion vevs into the coefficients a_D . These vevs are ultimately determined by the phases of the a_D . Instead of looking for solutions for m , a , and b in terms of the superpotential parameters, we instead solve (4.19) for W_0 , a_2 , and a_3 as functions of the Kähler moduli. That way, given a particular place in the Kähler cone, we can determine the precise superpotential couplings that give rise to a supersymmetric vacuum there. Doing this, we find

$$W_0 = - \frac{e^{(a+b-m)(a-5b+m)} (3(a-b+m) + \text{Vol})}{3a} \quad (4.20)$$

where

$$\text{Vol} = -a^3 + 6ab^2 + 6b^3 - 9b^2m + m^3 , \quad (4.21)$$

as well as

$$\begin{aligned} a_2 &= \frac{(a-b)e^{a^2-8ab-11b^2+12bm-m^2}}{a} \\ a_3 &= \frac{(m-a)e^{a^2-4ab-8b^2+6bm}}{a} . \end{aligned} \quad (4.22)$$

From these expressions it is clear that we can achieve supersymmetric solutions inside the Kähler cone with a_2, a_3 (and a_1) all positive and $W_0 < 0$.

Acknowledgments

We would like to thank I. García-Etxebarria, J. Halverson, W. Taylor, and especially S. Sethi for valuable discussions. We also thank M. Wijnholt for helpful discussions on the first draft of this paper and M. Esole for pointing out an error in the computation of bundle cohomologies on $\mathbb{F}_4$ in Appendix A.1.2 in v1. We are all very grateful to the Caltech theory group for their generous hospitality at a crucial stage of this work. JM would also like to thank the Perimeter Institute for Theoretical Physics, the theoretical physics and algebraic geometry groups at The Ohio State University, the theory group at the University of Pennsylvania, the organizers of the Spring 2011 String Vacuum Project meeting, and the Center for Theoretical Physics at the Massachusetts Institute of Technology for their hospitality at various stages of this work. JM and NS are grateful to the organizers of the Fall 2010 String Vacuum Project at The Ohio State University during the initial stages of this work. NS is grateful to the University of Chicago theory group for hospitality. SSN thanks the organizers of StringMath 2011 for hospitality at UPenn and a stimulating conference. The work of JM is supported by DOE grant DE-FG02-90ER-40560 and NSF grant PHY-0855039. We thank Kolya Gromov for communication assistance during the final stages of this work.

A Three-fold

In this appendix we explain an alternative construction of the threefold of [29], which was already outlined in section 4.1. We start with $\mathbb{P}^3$ parametrized by homogeneous coordinates $[Z_1, Z_2, Z_3, Z_4]$ and blow up the point $p_0 = [0, 0, 0, 1]$. We let H denote the class that descends from the hyperplane of $\mathbb{P}^3$ and E denote the exceptional divisor. There are correspondingly two curve classes. We use ℓ_0 for the descendent of the intersection of hyperplanes in $\mathbb{P}^3$ and ℓ the nontrivial $\mathbb{P}^1$ inside E . The intersection data in this blown-up space, which we refer to as X , is

$$\begin{array}{c|cc} & H & E \\ \hline H & \ell_0 & 0 \\ E & 0 & -\ell \end{array} \quad (\text{A.1})$$

$$\begin{array}{c|cc} & H & E \\ \hline \ell_0 & 1 & 0 \\ \ell & 0 & -1 \end{array} \quad (\text{A.2})$$

In these computations, we used the fact that $(H - E) \cdot E = \ell$ and $(H - E) \cdot \ell = 1$. Nonvanishing triple intersections are

$$H^3 = 1 \quad E^3 = 1. \quad (\text{A.3})$$

Within this blown-up space, we consider the proper transform of the nodal cubic curve

$$\mathcal{C} : \quad Z_3 = 0, \quad Z_4 Z_1 Z_2 + (Z_1 + Z_2)^3 = 0. \quad (\text{A.4})$$

A smooth cubic curve inside the $\mathbb{P}^2$ defined by $Z_3 = 0$ is a torus. The nodal cubic here is a pinched torus which is topologically equivalent to a $\mathbb{P}^1$ glued to itself by identifying the north and south poles. When we take the proper transform of $\mathcal{C}$, we separate the north and south poles and get an honest $\mathbb{P}^1$. If we want to describe the proper transform in equations, we can do it as follows. Each of the hyperplanes $Z_i = 0$ for $i = 1, 2, 3$ becomes irreducible, containing one component in the class $H - E$ and another in the class E . Accordingly, we can write

$$Z_i = \zeta W_i, \quad i = 1, 2, 3 \quad (\text{A.5})$$

for W_i sections of the bundle $\mathcal{O}(H - E)$ and ζ a section of $\mathcal{O}(E)$ whose vanishing defines the exceptional divisor. The equations for $\mathcal{C}$ now become

$$\zeta W_3 = 0, \quad \zeta^2 (Z_4 W_1 W_2 + \zeta (W_1 + W_2)^3) = 0. \quad (\text{A.6})$$

The proper transform is obtained by dropping the ζ prefactors. Note that there are no solutions to $W_1 = W_2 = W_3 = 0$ since they are really just (global extensions of) projective coordinates on the exceptional $\mathbb{P}^2$. This is what ensures that the curve

$$\mathcal{C}' : \quad W_3 = 0, \quad Z_4 W_1 W_2 + \zeta (W_1 + W_2)^3 = 0 \quad (\text{A.7})$$

is smooth. As we said $\mathcal{C}'$ is a $\mathbb{P}^1$ so, when we blow up along $\mathcal{C}'$ to get our B_3 , the exceptional divisor of that blow-up will be a Hirzebruch surface. To see which one, note that $\mathcal{C}'$ is in the class

$$[\mathcal{C}'] = (H - E) \cdot (3H - 2E). \quad (\text{A.8})$$

The normal bundle is a sum of line bundles and, because $\mathcal{C}'$ is a $\mathbb{P}^1$, these are determined by their degrees. The normal bundle inside $(H - E)$ is computed by $(H - E)(3H - 2E)^2 = 5$ while the normal bundle inside $3H - 2E$ is computed by $(H - E)^2(3H - 2E) = 1$. The normal bundle is therefore $\mathcal{O}(5) \oplus \mathcal{O}(1)$ so when we blow up along $\mathcal{C}'$ the exceptional divisor D' is an $\mathbb{F}_4$.

The final step, as we said, is blowing up along $\mathcal{C}'$ to get the threefold B_3 of [29]. We get an exceptional divisor D' and let E' denote the proper transform of E , which is a dP_2 surface. We get a new curve class as well which is the $\mathbb{P}^1$ fiber of D' . For our curve classes, we take ℓ_0 to be the descendent of the curve of the same name in both $\mathbb{P}^3$ and X . By ℓ we mean the proper transform of the curve of the same name (ℓ) in X . Finally, let us take the third curve class to be the fiber of our $\mathbb{F}_4$. Because it is the fiber of a Hirzebruch surface, let us call this curve f ⁴⁵

The intersection of divisors with curves is easy to compute

	H	E'	D'
ℓ_0	1	0	0
ℓ	0	-1	1
f	0	0	-1

(A.9)

The intersection of the Hirzebruch surface D' with ℓ is 1 because ℓ is the proper transform of a hyperplane in E' (in the class $h - e_i$) while the Hirzebruch surface is a fibration of f over the curve $\mathcal{C}'$. The curve f reduces to an exceptional curve in E' which intersects $h - e_i$ exactly once. The intersection of D' with f is computed by noting that $H - D' - E'$ meets f exactly once.

We now turn to intersections of divisors. Part of the intersection table is easy to compute

	H	E'	$H - D' - E'$
H	ℓ_0	0	$\ell_0 - 3f$
E'	0	*	*
$H - D' - E'$	$\ell_0 - 3f$	*	*

(A.10)

We can fill in the rest by noting some useful facts. First, the intersection of $H - E$ with E is just ℓ so the intersection of $H - E'$ with E' is the total transform $\ell + f$. The reason for this is that $H - E$ does not contain the curve $\mathcal{C}'$. From this we conclude that

$$(H - E') \cdot E' = \ell + f \quad \implies \quad E'^2 = -(\ell + f). \quad (\text{A.11})$$

Now, we know that E' meets D' in twice the curve class f so we have that

$$E' \cdot D' = 2f \quad \implies \quad (H - D' - E') \cdot E' = \ell - f. \quad (\text{A.12})$$

⁴⁵Note that f is the curve that we called $\ell - G'$ in [29].

Finally, we need $(H - D' - E')^2$. For this, it will be enough to get D'^2 . We can discern this, however, from

$$(H - D' - E') \cdot (3H - D' - 2E') = 0. \quad (\text{A.13})$$

This leads to

$$(H - D' - E')^2 = \ell + 5f - 2\ell_0 \quad (\text{A.14})$$

and we complete the intersection table as

	H	E'	$H - D' - E'$	
H	ℓ_0	0	$\ell_0 - 3f$	(A.15)
E'	0	$-\ell - f$	$\ell - f$	
$H - D' - E'$	$\ell_0 - 3f$	$\ell - f$	$\ell + 5f - 2\ell_0$	
D'	$3f$	$2f$	$3\ell_0 - 7f - 2\ell$	

Now, for instantons we are particularly interested in divisors S_2 with trivial $H^0(S_2, N_{S_2/B_3})$. These will be divisors that do not move in families inside B_3 . The candidate divisors are

$$\begin{aligned} E' &\cong dP_2 \\ D' &\cong \mathbb{F}_4 \\ H - D' - E' &\cong dP_1. \end{aligned} \quad (\text{A.16})$$

A.1 Divisors and Normal Bundles

To study whether a divisor S_2 plays a role in the generation of nonperturbative couplings via M5 instantons, it is not enough to know that $h^0(S_2, N_{S_2/B_3}) = 0$. We would also like to know $h^p(S_2, N_{S_2/B_3})$ for $p = 1, 2$. In this subsection, we compute these cohomologies for the divisors in (A.16).

A.1.1 $E' = dP_2$

First we start with $E' = dP_2$. Denote by h , e_1 , and e_2 the standard generators of $H_2(dP_2, \mathbb{Z})$. We have

$$e_1 \sim e_2 = f, \quad h = \ell + f. \quad (\text{A.17})$$

Now, the normal bundle is given by

$$E'^2 = -\ell - f = -h. \quad (\text{A.18})$$

We can compute the relevant cohomologies as follows

$$\begin{aligned} H^0(dP_2, \mathcal{O}(-h)) &= 0 \\ H^2(dP_2, \mathcal{O}(-h)) &= H^0(dP_2, \mathcal{O}(-2h + e_1 + e_2)) = 0. \end{aligned} \quad (\text{A.19})$$

These vanishings mean that

$$\begin{aligned} h^1(dP_2, \mathcal{O}(-h)) &= -\chi(dP_2, \mathcal{O}(-h)) \\ &= 0. \end{aligned} \quad (\text{A.20})$$

To do this computation, we use the fact that

$$\begin{aligned}\chi(S, L) &= \int_S \text{ch}(V) \wedge \text{Td}(S) \\ &= \int_S \left[\frac{1}{12} (c_1(TS)^2 + c_2(TS)) + \frac{1}{2} c_1(L)^2 + \frac{1}{2} c_1(L) c_1(TS) \right]\end{aligned}\tag{A.21}$$

along with

$$c_1(T_{dP_n}) = \mathcal{O}(3h - \sum_{i=1}^n e_i) \quad \int_{dP_n} [c_2(T_{dP_n}) + c_1(T_{dP_n})^2] = 12.\tag{A.22}$$

To summarize, we find

$$\boxed{h^{0,p}(E', N_{E'/B_3}) = 0}\tag{A.23}$$

Of course we needed this just to ensure that the 7-brane GUT theory did not contain chiral adjoints.

A.1.2 $D' = \mathbb{F}_4$

We now turn to $D' = \mathbb{F}_4$. We usually use b for the base of the Hirzebruch surface and f for the fiber. We know that the curve f in B_3 is just the fiber of $\mathbb{F}_4$ by construction. On the other hand, we expect that b is given by the intersection of D' with $(H - D' - E')$. This is because $H - D' - E'$ is the proper transform of the hyperplane containing the $\mathbb{P}^1$ along which we blew up to get D' . Intersecting D' with this hyperplane should give the base of the Hirzebruch surface. This means we have

$$f \sim f \quad b \sim (H - D' - E') \cdot D' = 3\ell_0 - 2\ell - 7f.\tag{A.24}$$

We can check this identification by computing some intersections inside D' . Using the relations

$$D' \cdot E' = 2f \quad D' \cdot (H - D' - E') = b\tag{A.25}$$

we find

$$\begin{aligned}2f \cdot_{D'} f &= E' \cdot f \\ &= 0 \\ 2f \cdot_{D'} b &= E' \cdot (3\ell_0 - 2\ell - 7f) \\ &= 2 \\ b \cdot_{D'} b &= (H - D' - E') \cdot (3\ell_0 - 2\ell - 7f) \\ &= -4.\end{aligned}\tag{A.26}$$

From this, we learn that

$$f \cdot_{D'} f = 0 \quad f \cdot_{D'} b = 1 \quad b \cdot_{D'} b = -4,\tag{A.27}$$

which is precisely the intersection table of $\mathbb{F}_4$.

To move further, it is useful to know the Chern classes of $\mathbb{F}_4$. Viewing $\mathbb{F}_n$ properly as a $\mathbb{P}^1$ fibration over $\mathbb{P}^1$ we have (recalling that $c(\mathbb{P}^1) = 1 + 2f$)

$$c(\mathbb{F}_n) = c(\mathbb{P}^1)(1 + b)(1 + b + nf) \quad (\text{A.28})$$

so that

$$c_1(\mathbb{F}_n) = 2b + (n + 2)f \quad c_2(\mathbb{F}_n) = 4 \quad (\text{A.29})$$

Note that

$$c_1(\mathbb{F}_n)^2 = 8 \quad (\text{A.30})$$

Now, the normal bundle of D' is given by

$$(D')^2 = 8f + 2\ell - 3\ell_0 = f - b \quad (\text{A.31})$$

so what we want to compute is $h^{0,p}(\mathbb{F}_4, \mathcal{O}(f - b))$. It is rather clear that

$$H^0(\mathbb{F}_4, \mathcal{O}(f - b)) = 0. \quad (\text{A.32})$$

We also have

$$H^2(\mathbb{F}_4, \mathcal{O}(f - b)) = H^0(\mathbb{F}_4, \mathcal{O}(b - f - (2b + 6f))) = 0. \quad (\text{A.33})$$

Now, we can evaluate

$$\chi(\mathbb{F}_4, \mathcal{O}(f - b)) = \frac{1}{2}(f - b) \cdot (b + 7f) = 0. \quad (\text{A.34})$$

This means that

$$H^1(\mathbb{F}_4, \mathcal{O}(f - b)) = 0. \quad (\text{A.35})$$

To summarize, we find that

$$\boxed{h^{0,p}(D', N_{D'/B_3}) = 0} \quad (\text{A.36})$$

A.1.3 $H - D' - E' = dP_1$

Let us start by identifying the hyperplane $\tilde{h}$ and exceptional curve $\tilde{e}$ of $H - D' - E'$. We have

$$\tilde{h} = (H - D' - E') \cdot H = \ell_0 - 3f \quad \tilde{e} = (H - D' - E') \cdot E' = \ell - f \quad (\text{A.37})$$

To check this, we can compute

$$\begin{aligned} \tilde{h} \cdot_{H-D'-E'} \tilde{h} &= H \cdot (\ell_0 - 3f) \\ &= 1 \\ \tilde{h} \cdot_{H-D'-E'} \tilde{e} &= H \cdot (\ell - f) \\ &= 0 \\ &= E' \cdot (\ell_0 - 3f) \\ &= 0 \\ \tilde{e} \cdot_{H-D'-E'} \tilde{e} &= E' \cdot (\ell - f) \\ &= -1. \end{aligned} \quad (\text{A.38})$$

The normal bundle of $H - D' - E'$ is given by

$$(H - D' - E')^2 = \ell + 5f - 2\ell_0 = -2\tilde{h} + \tilde{e}. \quad (\text{A.39})$$

Now it immediately follows that

$$\begin{aligned} H^0(dP_1, \mathcal{O}(-2\tilde{h} + \tilde{e})) &= 0 \\ H^2(dP_1, \mathcal{O}(-2\tilde{h} + \tilde{e})) &= H^0(dP_1, \mathcal{O}(-h)) \\ &= 0. \end{aligned} \quad (\text{A.40})$$

Now we compute

$$\chi(dP_1, \mathcal{O}(-2\tilde{h} + \tilde{e})) = 1 + \frac{1}{2}(-2\tilde{h} + \tilde{e}) \cdot (\tilde{h}) = 0 \quad (\text{A.41})$$

to conclude that

$$H^1(dP_1, \mathcal{O}(-2\tilde{h} + \tilde{e})) = 0. \quad (\text{A.42})$$

So, to summarize, we have that

$$\boxed{h^{0,p}(H - D' - E', N_{(H-D'-E')/B_3}) = 0} \quad (\text{A.43})$$

References

- [1] R. Donagi and M. Wijnholt, *Higgs Bundles and UV Completion in F-Theory*, 0904.1218.
- [2] J. Marsano, N. Saulina, and S. Schafer-Nameki, *Monodromies, Fluxes, and Compact Three-Generation F-theory GUTs*, *JHEP* **08** (2009) 046, [0906.4672].
- [3] E. Dudas and E. Palti, *On hypercharge flux and exotics in F-theory GUTs*, *JHEP* **09** (2010) 013, [1007.1297].
- [4] J. Marsano, *Hypercharge Flux, Exotics, and Anomaly Cancellation in F-theory GUTs*, *Phys.Rev.Lett.* **106** (2011) 081601, [1011.2212].
- [5] C. Ludeling, H. P. Nilles, and C. C. Stephan, *The Potential Fate of Local Model Building*, *Phys.Rev.* **D83** (2011) 086008, [1101.3346].
- [6] M. J. Dolan, J. Marsano, N. Saulina, and S. Schafer-Nameki, *F-theory GUTs with U(1) Symmetries: Generalities and Survey*, 1102.0290.
- [7] J. Marsano, N. Saulina, and S. Schafer-Nameki, *Compact F-theory GUTs with U(1)_{PQ}*, *JHEP* **04** (2010) 095, [0912.0272].
- [8] H. Hayashi, T. Kawano, Y. Tsuchiya, and T. Watari, *More on Dimension-4 Proton Decay Problem in F-theory – Spectral Surface, Discriminant Locus and Monodromy*, *Nucl.Phys.* **B840** (2010) 304–348, [1004.3870].
- [9] T. W. Grimm and T. Weigand, *On Abelian Gauge Symmetries and Proton Decay in Global F-theory GUTs*, *Phys.Rev.* **D82** (2010) 086009, [1006.0226].
- [10] J. Marsano, N. Saulina, and S. Schafer-Nameki, *A Note on G-Fluxes for F-theory Model Building*, *JHEP* **11** (2010) 088, [1006.0483].
- [11] J. Marsano, N. Saulina, and S. Schafer-Nameki, *Gauge Mediation in F-Theory GUT Models*, *Phys. Rev.* **D80** (2009) 046006, [0808.1571].
- [12] J. J. Heckman and C. Vafa, *F-theory, GUTs, and the Weak Scale*, *JHEP* **0909** (2009) 079, [0809.1098]. * Temporary entry *.
- [13] R. Tatar, Y. Tsuchiya, and T. Watari, *Right-handed Neutrinos in F-theory Compactifications*, *Nucl.Phys.* **B823** (2009) 1–46, [0905.2289].
- [14] E. Dudas and E. Palti, *Froggatt-Nielsen models from E(8) in F-theory GUTs*, *JHEP* **1001** (2010) 127, [0912.0853].
- [15] R. Friedman, J. Morgan, and E. Witten, *Vector bundles and F theory*, *Commun.Math.Phys.* **187** (1997) 679–743, [hep-th/9701162].

- [16] G. Curio and R. Y. Donagi, *Moduli in $N=1$ heterotic / F theory duality*, *Nucl.Phys.* **B518** (1998) 603–631, [[hep-th/9801057](#)].
- [17] B. Andreas, G. Curio, “On discrete twist and four flux in $N=1$ heterotic / F theory compactifications,” *Adv. Theor. Math. Phys.* **3**, 1325–1413 (1999). [[hep-th/9908193](#)].
- [18] R. Donagi and M. Wijnholt, *Model Building with F -Theory*, [0802.2969](#).
- [19] R. Donagi and M. Wijnholt, *Breaking GUT Groups in F -Theory*, [0808.2223](#).
- [20] H. Hayashi, T. Kawano, R. Tatar, and T. Watari, *Codimension-3 Singularities and Yukawa Couplings in F -theory*, *Nucl.Phys.* **B823** (2009) 47–115, [[0901.4941](#)]. *
Temporary entry *.
- [21] J. J. Heckman, J. Marsano, N. Saulina, S. Schafer-Nameki, and C. Vafa, *Instantons and SUSY breaking in F -theory*, [0808.1286](#).
- [22] J. Marsano, N. Saulina, and S. Schafer-Nameki, *An Instanton Toolbox for F -Theory Model Building*, *JHEP* **01** (2010) 128, [[0808.2450](#)].
- [23] M. Cvetič, I. García-Etxebarria, and R. Richter, *Branes and instantons at angles and the F -theory lift of $O(1)$ instantons*, *AIP Conf.Proc.* **1200** (2010) 246–260, [[0911.0012](#)].
- [24] R. Blumenhagen, A. Collinucci, and B. Jurke, *On Instanton Effects in F -theory*, *JHEP* **1008** (2010) 079, [[1002.1894](#)].
- [25] M. Cvetič, I. García-Etxebarria, and J. Halverson, *Global F -theory Models: Instantons and Gauge Dynamics*, *JHEP* **1101** (2011) 073, [[1003.5337](#)].
- [26] R. Donagi and M. Wijnholt, *MSW Instantons*, [1005.5391](#).
- [27] M. Cvetič, I. García-Etxebarria, and J. Halverson, *On the computation of non-perturbative effective potentials in the string theory landscape: IIB/ F -theory perspective*, [1009.5386](#).
- [28] T. W. Grimm, M. Kerstan, E. Palti, and T. Weigand, *On Fluxed Instantons and Moduli Stabilisation in IIB Orientifolds and F -theory*, [1105.3193](#).
- [29] J. Marsano, N. Saulina, and S. Schafer-Nameki, *F -theory Compactifications for Supersymmetric GUTs*, *JHEP* **08** (2009) 030, [[0904.3932](#)].
- [30] R. Blumenhagen, T. W. Grimm, B. Jurke, and T. Weigand, *Global F -theory GUTs*, *Nucl.Phys.* **B829** (2010) 325–369, [[0908.1784](#)].
- [31] T. W. Grimm, S. Krause, and T. Weigand, *F -Theory GUT Vacua on Compact Calabi-Yau Fourfolds*, *JHEP* **1007** (2010) 037, [[0912.3524](#)].

- [32] M. Esole and S.-T. Yau, *Small resolutions of $SU(5)$ -models in F -theory*, 1107.0733.
- [33] R. Blumenhagen, M. Cvetič, S. Kachru, and T. Weigand, *D-Brane Instantons in Type II Orientifolds*, *Ann.Rev.Nucl.Part.Sci.* **59** (2009) 269–296, [0902.3251].
- [34] M. Bershadsky, K. A. Intriligator, S. Kachru, D. R. Morrison, V. Sadov, *et. al.*, *Geometric singularities and enhanced gauge symmetries*, *Nucl.Phys.* **B481** (1996) 215–252, [hep-th/9605200].
- [35] R. Donagi and M. Wijnholt, *Gluing Branes, I*, 1104.2610.
- [36] H. Hayashi, R. Tatar, Y. Toda, T. Watari, and M. Yamazaki, *New Aspects of Heterotic– F Theory Duality*, *Nucl. Phys.* **B806** (2009) 224–299, [0805.1057].
- [37] D. R. Morrison and C. Vafa, *Compactifications of F -Theory on Calabi–Yau Threefolds – I*, *Nucl. Phys.* **B473** (1996) 74–92, [hep-th/9602114].
- [38] D. R. Morrison and C. Vafa, *Compactifications of F -Theory on Calabi–Yau Threefolds – II*, *Nucl. Phys.* **B476** (1996) 437–469, [hep-th/9603161].
- [39] E. Witten, *On flux quantization in M -theory and the effective action*, *J. Geom. Phys.* **22** (1997) 1–13, [hep-th/9609122].
- [40] S. Sethi, C. Vafa, and E. Witten, *Constraints on low-dimensional string compactifications*, *Nucl. Phys.* **B480** (1996) 213–224, [hep-th/9606122].
- [41] A. Collinucci and R. Savelli, *On Flux Quantization in F -Theory*, 1011.6388.
- [42] E. Witten, *Nonperturbative superpotentials in string theory*, *Nucl.Phys.* **B474** (1996) 343–360, [hep-th/9604030].
- [43] O. J. Ganor, *A Note on zeros of superpotentials in F theory*, *Nucl.Phys.* **B499** (1997) 55–66, [hep-th/9612077].
- [44] F. Denef, M. R. Douglas, and B. Florea, *Building a better racetrack*, *JHEP* **0406** (2004) 034, [hep-th/0404257].
- [45] N. Saulina, *Topological constraints on stabilized flux vacua*, *Nucl.Phys.* **B720** (2005) 203–210, [hep-th/0503125].
- [46] R. Kallosh, A.-K. Kashani-Poor, and A. Tomasiello, *Counting fermionic zero modes on $M5$ with fluxes*, *JHEP* **0506** (2005) 069, [hep-th/0503138].
- [47] F. Denef, M. R. Douglas, B. Florea, A. Grassi, and S. Kachru, *Fixing all moduli in a simple f -theory compactification*, *Adv.Theor.Math.Phys.* **9** (2005) 861–929, [hep-th/0503124].

- [48] P. K. Tripathy and S. P. Trivedi, *Compactification with flux on K3 and tori*, *JHEP* **0303** (2003) 028, [[hep-th/0301139](#)].
- [49] L. Gorlich, S. Kachru, P. K. Tripathy, and S. P. Trivedi, *Gaugino condensation and nonperturbative superpotentials in flux compactifications*, *JHEP* **0412** (2004) 074, [[hep-th/0407130](#)].
- [50] R. Kallosh and D. Sorokin, *Dirac action on M5 and M2 branes with bulk fluxes*, *JHEP* **0505** (2005) 005, [[hep-th/0501081](#)].
- [51] D. Lust, S. Reffert, W. Schulgin, and P. K. Tripathy, *Fermion zero modes in the presence of fluxes and a non-perturbative superpotential*, *JHEP* **0608** (2006) 071, [[hep-th/0509082](#)].
- [52] D. Tsimpis, *Fivebrane instantons and Calabi-Yau fourfolds with flux*, *JHEP* **0703** (2007) 099, [[hep-th/0701287](#)].
- [53] P. S. Howe, N. Lambert, and P. C. West, *The Selfdual string soliton*, *Nucl.Phys.* **B515** (1998) 203–216, [[hep-th/9709014](#)].
- [54] E. Bergshoeff, E. Sezgin, and P. Townsend, *Supermembranes and Eleven-Dimensional Supergravity*, *Phys.Lett.* **B189** (1987) 75–78.
- [55] E. Bergshoeff, E. Sezgin, and P. Townsend, *Properties of the Eleven-Dimensional Super Membrane Theory*, *Annals Phys.* **185** (1988) 330.
- [56] K. Bobkov, V. Braun, P. Kumar, and S. Raby, *Stabilizing All Kahler Moduli in Type IIB Orientifolds*, *JHEP* **1012** (2010) 056, [[1003.1982](#)].
- [57] B. Andreas and G. Curio, *From Local to Global in F-Theory Model Building*, *J.Geom.Phys.* **60** (2010) 1089–1102, [[0902.4143](#)].
- [58] S. Katz, D. R. Morrison, S. Schafer-Nameki, and J. Sully, *Tate’s algorithm and F-theory*, [1106.3854](#).