

CHCRUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Non-Abelian quasigapless modes localized on mass vortices in superfluid ^{3}He-B

Muneto Nitta, Mikhail Shifman, and Walter Vinci Phys. Rev. D 87, 081702 — Published 25 April 2013 DOI: 10.1103/PhysRevD.87.081702

On Non-Abelian Quasi-Gapless Modes Localized on Mass Vortices in Superfluid ³He-B

MUNETO NITTA¹, MIKHAIL SHIFMAN² AND WALTER VINCI^{3,4,5}

¹Department of Physics, and Research and Education Center for Natural Sciences, Keio University, 4-1-1 Hiyoshi, Yokohama, Kanagawa 223-8521, Japan

² William I. Fine Theoretical Physics Institute, University of Minnesota, Minneapolis, MN 55455, USA

 ³University of Pisa, Department of Physics "E. Fermi", INFN, Largo Bruno Pontecorvo 7, 56127, Italy
 ⁴ London Centre for Nanotechnology, University College London, London, WC1H 0AH, UK.
 ⁵ Department of Computer Science, University College London, London WC1E 6BT, UK.

nitta(at)phys-h.keio.ac.jp
shifman(at)physics.umn.edu
w.vinci(at)ucl.ac.uk

Abstract

Kelvin waves, or Kelvons, have been known for a long time as gapless excitations propagating along superfluid vortices. These modes can be interpreted as the Nambu-Goldstone excitations arising from the spontaneous breaking of the translational symmetry. Recently a different type of gapless excitations localized on strings – the so-called non-Abelian modes – attracted much attention in high-energy physics. We discuss their relevance in condensed matter physics. Non-Abelian rotational quasi-gapless excitations could appear on the mass vortices in the B phase of the superfluid ³He, due to the fact that the order parameter in ³He-B is tensorial. While the U(1) rotational excitations are well established in vortices with asymmetric cores, the non-Abelian rotational excitations, belonging to the same family, were not considered.

1 Introduction

In this paper we argue that vortices in certain superfluids with a tensorial order parameter, such as the B-phase of superfluid ³He, can have non-Abelian quasi-gapless modes. The existence of such modes depends on the values of a number of phenomenological parameters (see Eq. (2.2) below). Therefore, strictly speaking, for the time being we can only say that our assertion refers to a superfluid with the symmetry of ³He-B, characterized by relatively small values of parameters γ_2 and γ_3 , which could be arranged in ultra cold Fermi gases with p-wave pairings.

By non-Abelian we mean that the moduli (collective coordinates) that give rise to these modes are described by non-Abelian sigma models, say, the O(3) model. The very notion of non-Abelian gapless (or quasi-gapless) modes came from the study of topological defects in high-energy physics, especially in supersymmetric non-Abelian field theories, where they attracted much attention recently [1], see also [2] for reviews. These non-Abelian modes are called *orientational*; they play an important role in the theory of confinement.

Non-Abelian Nambu-Goldstone (NG) excitations localized on the vortices occur when a non-Abelian global symmetry of the ground state is spontaneously broken by the vortex. An interesting case that can be realized in Nature is quark matter at high densities, such as in the neutron star cores. In this case nucleons are melt into the quark matter in which color SU(3) and flavor SU(3) symmetries are spontaneously broken down to a diagonal SU(3) color-flavor locked symmetry [3], resembling the B phase of ³He. The vortices in this medium spontaneously break the SU(3) color-flavor locked symmetry of the ground state. As a result, non-Abelian NG gapless modes appear, localized on and propagating along the non-Abelian vortices [4].¹

Motivated by these developments in high-energy physics we would like to study whether the results obtained there are relevant in condensed matter applications.

Superfluid Helium-3 is one of the most interesting states of matter thoroughly studied both experimentally [6] and theoretically [7–10]. Unlike conventional superfluids, the fermionic ³He atoms condense after forming bosonic Cooper pairs [7]. The attractive interaction between ³He atoms has a strong short-range repulsive core that favors a *P*-wave pairing and bound states with unit angular momentum and spin. The consequence of this fact is that the order parameter describing the condensate has a tensorial structure: it is a 3 by 3 matrix $e_{\mu i}$, where μ and *i* are the spin and orbital indices, respectively [8,11,12].

Low-energy physics of superfluids can be described by gapless excitations, the NG modes associated with spontaneously broken global symmetries. The most important NG modes are: (i) phonons associated with the spontaneously broken phase symmetry $U(1)_p$ (here the subscript p stands for phase), and (ii) magnons associated with spontaneously broken SO(3)spin symmetry [13].

In conventional superfluids the breaking of the Abelian phase symmetry $U(1)_p \rightarrow 1$ leads to the existence of phonon excitations in the bulk and of topologically stable superfluid vortices [14] when the superfluid sample is under rotation [15–17]. Long ago Lord Kelvin predicted that vortices support vibrating modes (called Kelvons) which correspond to fluctuations of the vortex line [18, 16, 19]. Kelvons have been directly observed recently [20].

¹The above modes are gapped, however, after taking into account quantum corrections [5].

In unconventional superfluids, such as ³He, the order parameter is tensorial and non-Abelian symmetries related to spatial rotations are usually broken in the vacuum. If we neglect the spin-orbit interaction, rotations of spinorial and orbital indices can be performed independently; the full symmetry of ³He is $G = U(1)_p \times SO(3)_S \times SO(3)_L$, where $SO(3)_S$ and $SO(3)_L$ are spin and angular rotations. Two phases were theoretically predicted and experimentally (indirectly) observed (in the absence of external magnetic fields). We focus on the so-called B phase, where the ground state preserves a locked SO(3) symmetry,

$$G = U(1)_p \times SO(3)_S \times SO(3)_L \to H_{\mathrm{B}} = SO(3)_{S+L}.$$

The locked $SO(3)_{S+L}$ symmetry can be identified with the overall spatial rotations $SO(3)_R$. The expression above imply the existence of bulk "non-Abelian" NG bosons, called magnons in the condensed matter terminology, due to spontaneous breaking of non-Abelian global symmetries. The B phase supports topologically stable vortices:

$$\pi_1 \left(U(1)_p \times SO(3)_S \times SO(3)_L / SO(3)_{S+L} \right) = \pi_1 \left(U(1)_p \times SO(3)_{S-L} \right) = \mathbb{Z} \oplus \mathbb{Z}_2.$$

The first \mathbb{Z} factor corresponds to the breaking of the Abelian $U(1)_p$ symmetry and supports the so-called mass vortices. These vortices are created and stabilized in a lattice once the sample is rotated, and are characterized by a non-vanishing superfluid current and angular momentum, much in the same way as vortices in conventional superfluids.² Both the bulk magnons and the Kelvin excitations were recently observed [21, 20] in experiments with superfluid ³He and ⁴He.

In this paper we will argue that a hitherto unexplored type of quasi-gapless modes localized on the mass vortices in the superfluid ³He-B exist. While Kelvons can be interpreted as the NG modes arising from the breaking of translational symmetry, the excitations we discuss arise from the breaking of the spatial rotation symmetry $H_{\rm B} = SO(3)_{S+L}$ by the vortex solution. Because of their origin we will refer to them as non-Abelian.

It is known that the B phase is divided into two sub-phases classified according to the core structure of the mass vortex under rotations around the vortex axis. The core can be either axially symmetric or axially asymmetric [22, 11]. A possible signal of the axial symmetry breaking in the core of the mass vortices has been already observed [23]. The axial symmetry gives rise to a U(1) NG mode localized on the given mass vortex. This is the conventional U(1) NG excitation, while our task is to focus on non-Abelian excitations.

Considering the whole non-Abelian symmetry, there can exist up to two or three additional modes, due to breaking of the bulk symmetry $H_{\rm B} = SO(3)_{S+L}$ on the vortex, in accordance with the fact that

$$SO(3)_{S+L}/U(1)_z \simeq S^2$$
 or $SO(3)_{S+L}/1 \simeq S^3/\mathbb{Z}_2$

relevant respectively for the axially symmetric core (left) and for the asymmetric core (right).

As far as we know, these excitations were not discussed in the literature, nor observed in experiments. This is the first example of spatially localized non-Abelian modes in condensed

²The second \mathbb{Z}_2 factor stabilizes a more exotic type of vortices, the spin vortices. They are not directly created by the superfluid rotation; they were detected, however, as bound states with the mass vortices

matter physics. It is important to stress here that these modes can be quasi-gapless, or become gapped upon inclusion of quantum effects.

The arguments that lead us to this conclusion are explained in more detail in the next Sections. They can be applied to other unconventional superfluids with tensorial order parameters. The crucial points in our considerations are (i) the presence (in the B phase) of an unbroken non-Abelian (global rotational) $SO(3)_{S+L}$ symmetry, which may be spontaneously broken by the vortex, and (ii) tensorial nature of the order parameter. The rotational symmetry is broken by the vortex solution at two levels. As in the case of conventional superfluids, it is broken by the physical shape of the vortex itself – a line defect in three spatial dimensions. (iii) It is known that the moduli coming from the latter breaking are not independent of those originated by translations, in the case of scalar order parameter, while we argue that independent (quasi)moduli are generated in the case of tensorial order parameters.

From a more formal point of view, if the ground state of the theory is symmetric under H and the vortex solution preserves only a subgroup H' of the group H, then a particular vortex gives rise to a family of (degenerate) vortices which are characterized by a number $\nu = \dim H - \dim H'$ of continuous parameters (called moduli) of the solutions and ν_{NG} NG excitation modes. In relativistic field theories, the NG theorem implies the existence of a massless particle per each broken generator ($\nu_{NG} = \nu$). In the non-relativistic case [24] the NG theorem is subtler [25]: the number of the gapless excitations can be smaller than or equal to the number of the broken generators (moduli).³

Our proposal is that the rotational symmetry $H = SO(3)_{S+L}$ broken by a non-trivial configuration of the tensorial order parameter on the vortex solution, generates rotational non-Abelian (quasi)moduli ω^{ℓ} ($\ell = 1, 2, 3$) which are coupled, in the general case, to the conventional Kelvin excitations. The Kelvin excitation presents an example of the nonrelativistic type II NG mode. With two broken translations, there is only one independent circularly polarized fluctuation of the vortex. The number of the gapless rotational excitations, given a certain number of rotational moduli, must be established on the case-by-case basis in non-relativistic theories, through the quantization procedure.

2 The Ginzburg-Landau description of ³He

The starting point of our analysis is the Ginzburg-Landau description of the superfluid phase of ³He near the critical temperature. The order parameter is a 3×3 matrix $e_{\mu i}$ with spin indices μ (where $\mu = 1, 2, 3$) and orbital indices i (i = 1, 2, 3) that transform under the vector representations of $SO(3)_S$ and $SO(3)_L$;

$$e_{\mu i} \to e^{i\psi} S_{\mu\nu} L_{ij} e_{\nu j} \,, \tag{2.1}$$

where S and L are $SO(3)_S$ and $SO(3)_L$ orthogonal rotation matrices. ψ is a phase parameter corresponding to $U(1)_p$ rotations.

³One NG mode with a *linear* dispersion relation (called type-I) corresponds to one broken generator (as in the relativistic case), and, at the same time, one NG mode with a *quadratic* dispersion relation (called type-II) corresponds to *two* broken generators.

The most general time dependent expression for the free energy, invariant under the $U(1) \times SO(3)_S \times SO(3)_L$ symmetry of the theory takes the form

$$F_{GL} \equiv \hbar i \, e_{\mu i} \partial_t e^*_{\mu i} + \gamma_1 \partial_i e_{\mu j} \partial_i e^*_{\mu j} + \gamma_2 \partial_i e_{\mu i} \partial_j e^*_{\mu j} + \gamma_3 \partial_i e_{\mu j} \partial_j e^*_{\mu i} - \alpha \, e^*_{\mu i} e_{\mu i} \\ + \beta_1 e^*_{\mu i} e^*_{\mu i} e_{\nu j} e_{\nu j} + \beta_2 e^*_{\mu i} e_{\mu i} e^*_{\nu j} e_{\nu j} + \beta_3 e^*_{\mu i} e^*_{\nu i} e_{\mu j} e_{\nu j} + \beta_4 e^*_{\mu i} e_{\nu i} e^*_{\nu j} e_{\mu j} + \beta_5 e^*_{\mu i} e_{\nu i} e_{\nu j} e^*_{\mu j} .$$
(2.2)

The coefficients γ and β are phenomenological parameters depending both on temperature and pressure. Experimental determination of all these coefficients is very difficult [26]. One can roughly evaluate them from BCS-like calculations [12]. Note that the γ_2 and γ_3 terms in the second line differ only by a total derivative. Hence, in deriving the equations of motion in the bulk they can be identified. BCS-like calculations are valid at weak coupling, and predict the B-phase, were they are more reliable. Strong coupling effects have to be taken into account in general, especially to predict the appearance of the A phase. However, for the considerations presented in this paper the precise values of the coefficients γ , α and β are not important.

Minimizing the potential term in Eq. (2.2) with respect to the order parameter gives the B-phase ground state [10]

$$(e_0)_{\mu i} = e^{i\psi} \Delta (R_0)_{\mu i} , \qquad \Delta^2 \equiv \frac{\alpha}{6(\beta_1 + \beta_2) + 2(\beta_3 + \beta_4 + \beta_5)} ,$$
 (2.3)

where the ground state solution is parametrized (in the most general form) in terms of an element $(R_0)_{\mu i}$ of the SO(3) group. The configuration $(e_0)_{\mu i} = e^{i\psi}\Delta (R_0)_{\mu i}$ contains 4 parameters, representing the NG excitations in the bulk, or magnons, discussed in the introduction.

The unbroken symmetry corresponds to a combined action of spinorial and orbital transformations, namely,

$$S_{\mu\tilde{\mu}}L_{i\tilde{i}} (R_0)_{\mu\tilde{i}} = (R_0)_{\mu i} .$$
(2.4)

Without loss of generality we can set $(R_0)_{\mu i} = \delta_{\mu i}$ (i.e. $\theta = 0$ and $\psi = 0$) in the ground state. Then Eq. (2.4) becomes obvious provided S = L.

If the parameters $\gamma_{2,3}$ in (2.2) are small, the symmetry of the free energy (2.2) is enhanced. Indeed, in the limit $\gamma_2, \gamma_3 \to 0$, the spatial $SO(3)_R$ rotations of coordinates can be done independently from the spinorial $SO(3)_S$ and orbital rotation $SO(3)_L$ without changing the free energy. In this case, the $SO(3)_S$ and $SO(3)_L$ symmetries, together with the locked $SO(3)_{S+L}$ symmetry effectively act as internal symmetries. Then if we switch on $\gamma_{2,3} \neq 0$ these two terms in the free energy explicitly break $SO(3)_S \times SO(3)_L \times SO(3)_R$ down to a "locked" $SO(3)_{S+L} = SO(3)_R$. This observations will be important below.

Vortices in ³He-B

The simplest topologically stable mass vortex in the B phase of ${}^{3}\text{He}$ has the following asymptotic behavior

$$e_{\mu i}^{\text{vort}} = \Delta e^{i\alpha} \delta_{\mu i} \text{ at } r \to \infty,$$
 (2.5)

where α is the polar angle in the perpendicular plane. We identify ψ with the polar angle, so that the phase of the order parameter winds by 2π . From the topological point of view, a mass vortex is stable because $\pi_1(U(1)_p)$ is nontrivial. The asymptotic behavior (2.5) of the vortex solution does not break the $SO(3)_{L+S}$ symmetry of the bulk. Thus, no modulus associated with the phase winding appears. At generic distances, the ansatz (2.5) is generalized as follows,

$$e_{\mu i}^{\text{vort}}(x,y) = \Delta e^{i\alpha} \,\tilde{e}_{\mu i}(x,y) \,, \quad \tilde{e}_{\mu i} \equiv \begin{pmatrix} g_1(x,y) & f_{12}(x,y) & f_{13}(x,y) \\ f_{21}(x,y) & g_2(x,y) & f_{23}(x,y) \\ f_{31}(x,y) & f_{32}(x,y) & g_3(x,y) \end{pmatrix}, \quad (2.6)$$

with boundary conditions $\tilde{e}_{\mu i} \to \delta_{\mu i}$ at $r \to \infty$.

As already mentioned in the Introduction, vortices with an asymmetric core and a rotational modulus can exist. They are characterized by a profile matrix $\tilde{e}_{\mu i}(x, y)$ that break SO(2) rotations around the vortex axis, for example if all f's vanish, but $g_1 \neq g_2$ in the core. This is a well-known property that distinguishes the mass vortices in ³He-B from those in conventional superfluids [22]. Thus, the ³He vortices cannot be considered as diagonal embeddings of more conventional U(1) superfluid vortices.

The mentioned rotational gapless mode is Abelian. In the following, we want to consider its non-Abelian counterpart.

3 Scalar vs. nonscalar order parameters

To begin with, let us consider the usual case of a superfluid vortex in a theory with scalar order parameter (e.g. in ⁴He). Such vortices are characterized by two translational moduli, corresponding to translations in the x and y directions (assuming that the vortex axis is oriented along the z axis). Let us examine the physical reason behind the existence of these translational moduli (and, hence, the Kelvin mode). The underlying theory has a number invariances in the ground state. In the case of the scalar order parameter these symmetries are (a) translational (three generators); (b) rotational (three generators). Some of the above symmetries are obviously spontaneously broken by the vortex defect, namely, translations in the x and y direction $(T_{x,y})$ and rotations outside the x, y plane $(L_{zx} \text{ and } L_{zy})$.

It turns out that the two translational modes are one the conjugate momentum of the other, thus giving a single mode known as the Kelvin mode (see e.g. [24]). What is the fate of the rotational symmetry which is also spontaneously broken by the vortex? Two of the rotations are broken, while the SO(2) rotation around the z axis remains unbroken. At first sight the SO(3)/SO(2) pattern of the symmetry breaking must lead to two extra moduli, corresponding to broken L_{zx} and L_{zy} . However, these extra moduli are absent because the above rotations are already represented by the translational moduli,⁴ which we will call ξ_I (where the capital Latin subscript runs over I = 1, 2). In the quantization procedure the moduli ξ_I become the fields $\xi_I(t, z)$. Thus, in the case at hand, no extra moduli are needed to represent the spontaneously broken O(3) symmetry since local translations are equivalent to local rotations [27].

Dynamics of the ξ_I on the vortex is described by the gradient terms $\sim (\partial_z \xi_I)^2$, plus a time derivative. The string oriented along the z axis corresponds to the ground state:

$$\xi_I = 0. \tag{3.1}$$

 $^{{}^{4}}$ In the context of high energy physics this observation is sometimes referred to as the inverse Higgs mechanism [27, 28]

A rotated string is described by another ground state,

$$\xi_I(z) = a_I z$$
, a_I arbitrary real numbers. (3.2)

This is also a solution of the equation of motion. An apparent increase in the ground state energy for (3.2) compared to (3.1) is an illusion: for the rotated vortex the element of distance along the vortex changes too. The overall tension remains the same.

Let us pass to the problem of interest in which the relevant order parameter carries spatial indices, see Eq. (2.2). We will start our discussion from the simplest limit $\gamma_{2,3} = 0$ (or very small). If $\gamma_{2,3} = 0$ the symmetry of the ground state is enhanced: in addition to the translational symmetry and that of spatial rotations R, it has "internal" symmetries $SO(3)_S \times SO(3)_L$ spontaneously broken in the ground state down to $SO(3)_{L+S}$.

The vortex solution itself breaks T and R spontaneously, by the orientation of the vortex axis. As was explained above, this breaking generates only two (not four) moduli, ξ_I . The breaking of the vacuum $SO(3)_{L+S}$ symmetry (which in the above limit is "internal") necessarily produces either three rotational moduli in the vortex solution (if $SO(3)_{L+S}$ is completely broken on the given solution) or two such moduli (if $SO(3)_{L+S}$ is broken down to SO(2)). For definiteness let us focus on the first case. If the axially asymmetric solution exists in the limit $\gamma_{2,3} = 0$, it can be automatically elevated to the status of the full-fledged non-Abelian vortex. Indeed, since it breaks SO(2) by definition, one can "smear" this SO(2)over the $SO(3)_{L+S}$ by considering all possible embeddings of the broken SO(2) in $SO(3)_{L+S}$. In this way, we acquire three (non-Abelian) rotational moduli. The axially asymmetric solution exists provided that either $g_1 \neq g_2$, or one or more f's in Eq. (2.6) do not vanish, or both.

We did not attempt to systematically explore the parameter space $\beta_{1,...,5}$ in search of all possible global vortex solutions at $\gamma_{2,3} = 0$. This will be the subject of a special investigation. However, for some values of $\beta_{1,...,5}$ particular relevant solutions can be found in a qualitative way in the following *ansatz*:

$$e_{\mu i}^{\text{vort}} = \Delta(r_{\perp}) \, e^{i\alpha} \delta_{\mu i} + \varepsilon_{\mu i k} \, \chi^k(r_{\perp}) \tag{3.3}$$

where $\vec{r}_{\perp} = \{x, y\}$ and χ^i is a vector with regards to the unbroken $SO(3)_{L+S}$. In addition to (2.5), $\chi^k(r_{\perp}) \to 0$ at $r_{\perp} \to \infty$. In the core of the vortex, at small r_{\perp} , an instability for χ^i develops, $\chi^i \neq 0$, and we obtain two rotational moduli (corresponding to $SO(3) \to SO(2)$).

Now let us consider a general situation and switch on $\gamma_{2,3} \neq 0$. Applying $SO(3)_{L+S} = SO(3)_R$ to the vortex profile we get, generally speaking, three moduli ω^{ℓ} in the matrix structure of $e_{\mu i}^{\text{vort}}(x, y)$. Simultaneously the center of the solution gets shifted too by ξ_I , see Eq. (3.2), due to the entanglement of rotations and translations. The shift parameters a_I are linearly dependent on $\omega_{1,2}$.

There are two consequences: a mass gap proportional to $\gamma_{2,3}$ is generated for those moduli that are not associated with the U(1) rotation and, in addition, a derivative coupling connecting the translational and rotational moduli appears too.

4 Low-energy effective theories on the vortex

In this section we will briefly outline a general construction of low-energy effective theories on the vortex which is not very widely used in condensed matter (albeit is absolutely standard in high-energy physics, e.g. [29]). This theory allows one to describe dynamics of gapless and quasigapless excitations. This technique will help us to substantiate the last statement in Sect. 3.

The general prescription for deriving an effective low-energy theory is as follows. One expands the Ginzburg-Landau free energy (2.2) in terms of fluctuations of the order parameter, up to the second order in derivatives, near the vortex solution. These fluctuations are assumed to be dependent on the world-sheet coordinates of the vortex,

$$e_{\mu i}(x, y) = e_{\mu i}^{\text{vort}}(x, y) + \delta e_{\mu i}(x, y, z, t) .$$
(4.1)

The variation above is still completely generic. When varying expression (2.2), we observe that the terms proportional to the first order in the variations vanish because of the equations of motion. The remaining terms are as follows:

$$\delta^{2} (F_{GL}) = \hbar i \, \delta e_{\mu i}^{*} \partial_{t} \delta e_{\mu i} + \gamma_{1} \partial_{z} \delta e_{\mu j} \partial_{z} \delta e_{\mu j}^{*} + \gamma_{2} \partial_{z} \delta e_{\mu z} + \gamma_{2} \partial_{z} \delta e_{\mu z} \partial_{z} \partial_{z} \partial_{z} \partial_{z} \partial_{z} \partial_{z} \partial_{z}$$

where the ellipses denote terms of the third and higher order in $\delta e_{\mu i}(x, y, z, t)$ containing at most two derivatives. We will discuss them later.

For the time being we discard the t and z dependences in (4.2) and vary $\delta^2(F_{GL})$ with respect to $\delta e_{\mu i}$. In this way we arrive at a static differential operator acting on $\delta e_{\mu i}$, and define its eigenvalues and eigenmodes,

$$L(x,y)e_{\mu i}^{(n)}(x,y) = E^{(n)}e_{\mu i}^{(n)}(x,y).$$
(4.3)

Then one uses adiabatic approximation to expand $\delta e_{\mu i}(x, y, z, t) = \sum_{n} c_n(t, z) e_{\mu i}^{(n)}(x, y)$. Of all infinite set of modes we are interested only in the gapless and quasi-gapless modes, i.e. those for which $E^{(n)} = 0$ or small. Again, it is convenient to start from the limit $\gamma_{2,3} = 0$.

Let us denote the set of moduli on which $e_{\mu i}^{\text{vort}}(x, y)$ depends as m^a . This set includes two translational moduli ξ_I and three orientational ω^{ℓ} . Then $e_{\mu i}^{\text{vort}}(x, y; m^a)$ determines the gapless modes,

$$e_{\mu i}^{(0,a)}(x,y) \propto \frac{\partial}{\partial m^a} e_{\mu i}^{\text{vort}}(x,y;m), \qquad E^a = 0,$$
(4.4)

where the proportionality coefficient is derived from the normalization of each mode. Substituting these gapless modes in the expansion of $\delta e_{\mu i}(x, y, z, t)$ (and neglecting all nongapless modes),

$$\delta e_{\mu i}(x, y, z, t) = \sum_{a} m^{a}(z, t) e_{\mu i}^{(0, a)} , \qquad (4.5)$$

we arrive at a low-energy effective theory on the vortex in the form

$$F_{\rm LE}^{\rm grad} = \sum_{a,b} G_{a,b}(m)\partial_z m^a(z,t)\partial_z m^b(z,t)$$
(4.6)

plus a similar term with time derivative (quadratic in ∂_t for the type I NG modes and linear for the type II NG modes). In the quadratic approximation, as in (4.2), the target space metric $G_{a,b}(m)$ is replaced by $G_{a,b}(m)|_{m^a \equiv 0}$. The overall m dependence of $G_{a,b}(m)$ is fixed by the pattern of the symmetry breaking, or, alternatively, algebraically

$$G_{a,b}(m) = \int d^2 x_{\perp} \frac{\partial e_{\nu j}^{\text{vort}}(x, y; m)}{\partial m^a} \frac{\partial e_{\mu i}^{\text{vort}}(x, y; m)}{\partial m^b}$$
(4.7)

where convolutions of the spin and angular momentum indices are ignored, so that Eq. (4.7) is somewhat symbolic. Usually, translational moduli are uncorrelated with others. Then the metric (4.7) factors out and acquires a block form

$$F_{\rm LE} = \frac{T}{2} \partial_z \xi_I \partial_z \xi_I + \sum_{\ell,\tilde{\ell}} G_{\ell,\tilde{\ell}}(\omega) \partial_z \omega^\ell(z,t) \partial_z \omega^{\tilde{\ell}}(z,t) + \text{time der.}$$
(4.8)

In the previous sections we explained that our present problem is peculiar since the family of solutions is obtained under simultaneous and concerted variations of rotational and translational moduli.

Now, let us switch on $\gamma_{2,3} \neq 0$ but small. Since SO(3) in this case is explicitly broken by the vortex axis, only the modulus corresponding to the axial SO(2) will remain strictly gapless. Two others – belonging to the coset SO(3)/SO(2) – generally speaking will acquire a mass gap, proportional to $\gamma_{2,3}$, corresponding to a small shift of two eigenvalues in (4.4). In other words, we expect $\Delta F_{\text{LE,m}} \propto \gamma_{2,3} \sum_{\ell_{\perp}} \mu_{\ell_{\perp} \ell_{\perp}} (\omega^{\ell_{\perp}} \omega^{\tilde{\ell}_{\perp}})$ ($\ell, \tilde{\ell} = x, y$). We will show elesewhere that these modes are type-I NG modes with a linear disipersion relation so that all modes are indepentent. Inclusion of quantum effects (not discussed here) may or may not lift the U(1) mode too, depending on dynamical details.

In addition to (4.8) we will have an extra term $\Delta F_{\text{LE}}^{\text{grad}} \propto \gamma_{2,3} (\partial_z \xi_I) (\partial_z \omega^I)$. There is a dissipation of energy from the Kelvin mode to rotational and vice versa with the coefficient $\sim \gamma_{2,3}$.

5 Conclusions

In this paper, we discussed the emergence of non-Abelian (quasi)gapless modes on the vortices in ³He. So far our analyses is purely classical. Quantization needed to determine how many (quasi)gapless excitations appear from given moduli, and whether or not some are lifted by quantum effects is deferred till a more detailed publication.⁵

In conventional superfluids (e.g. ⁴He) the spontaneously broken translational and rotational symmetries on a vortex do not produce independent NG modes; one can say that only translational excitations occur, a single NG excitation, the Kelvon. This is due to the scalar nature of the order parameter.

At the same time, for the mass vortices in the superfluid ³He-B, the rotational symmetry gives rise to independent rotational (quasi)NG excitations. One of them – due to the axial

⁵In the case of non-Abelian vortices in dense QCD, gapless modes localized on the vortex are gapped after taking into account quantum corrections; quantum magnetic monopoles show up [5].

asymmetry of the vortex core – was known for a long time. We argued that extra non-Abelian rotational excitations exist, associated with the full SO(3) symmetry of the ground state. Generically, they are not strictly gapless and can be derivatively coupled with the translational excitations, the Kelvons. Whether or not they will show up as quasi-gapless depends on particular values of $\gamma_{2,3}$ and $\beta_{1,...,5}$. Generically, the parameters γ_2 and γ_3 are not parametrically small compared to γ_1 in ³He-B, and the gap of the new modes could be in principle of the order of the fermionic gap. As already mentioned in the Introduction, this problem can be in principle cured in ultra cold Fermi gases with p-wave pairing.

The tensorial order parameter in ³He-B is crucial for the phenomenon. We outlined the construction of the low-energy effective theory for the (quasi)gapless excitations localized on the vortex, defined as a (1+1)-dimensional theory on the vortex world sheet.

Acknowledgments

We thank Takeshi Mizushima, Yasumasa Tsutsumi for comments and Gregory. E. Volovik for discussions. This work of M.N. is supported in part by Grant-in-Aid for Scientific Research (No. 23740198) and by the "Topological Quantum Phenomena" Grant-in-Aid for Scientific Research on Innovative Areas (No. 23103515) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. The work of M.S. is supported in part by DOE grant DE-FG02- 94ER-40823.

References

- A. Hanany and D. Tong, "Vortices, instantons and branes", JHEP 0307, 037 (2003), hep-th/0306150.
 R. Auzzi, S. Bolognesi, J. Evslin, K. Konishi and A. Yung, "Nonabelian superconductors: Vortices and confinement in N = 2 SQCD", Nucl. Phys. B673, 187 (2003), hep-th/0307287.
 M. Shifman and A. Yung, "Non-Abelian string junctions as confined monopoles", Phys. Rev. D70, 045004 (2004), hep-th/0403149.
 A. Hanany and D. Tong, "Vortex strings and four-dimensional gauge dynamics", JHEP 0404, 066 (2004), hep-th/0403158.
 M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, "Moduli space of non-Abelian vortices", Phys.Rev.Lett. 96, 161601 (2006), hep-th/0511088.
 M. Eto, K. Konishi, G. Marmorini, M. Nitta, K. Ohashi et al., "Non-Abelian Vortices of Higher Winding Numbers", Phys.Rev. D74, 065021 (2006), hep-th/0607070.
- M. Eto, Y. Isozumi, M. Nitta, K. Ohashi and N. Sakai, "Solitons in the Higgs phase: The Moduli matrix approach", J.Phys. A39, R315 (2006), hep-th/0602170.
 M. Shifman and A. Yung, "Supersymmetric Solitons and How They Help Us Understand Non-Abelian Gauge Theories", Rev.Mod.Phys. 79, 1139 (2007), hep-th/0703267.
 M. Shifman and A. Yung, "Supersymmetric solitons, Cambridge, UK: Cambridge Univ. Pr. (2009) 259 p".
- [3] M. G. Alford, A. Schmitt, K. Rajagopal and T. Schafer, "Color superconductivity in dense quark matter", Rev.Mod.Phys. 80, 1455 (2008), 0709.4635.
- [4] A. Balachandran, S. Digal and T. Matsuura, "Semi-superfluid strings in high density QCD", Phys.Rev. D73, 074009 (2006), hep-ph/0509276. • E. Nakano, M. Nitta and T. Matsuura, "Non-Abelian strings in high density QCD: Zero modes and interactions", Phys.Rev. D78, 045002 (2008), 0708.4096. • E. Nakano, M. Nitta and T. Matsuura, "Non-Abelian Strings in Hot or Dense QCD", Prog. Theor. Phys. Suppl. 174, 254 (2008), 0805.4539. • M. Eto and M. Nitta, "Color Magnetic Flux Tubes in Dense QCD", Phys.Rev. D80, 125007 (2009), 0907.1278. • M. Eto, E. Nakano and M. Nitta, "Effective world-sheet theory of color magnetic flux tubes in dense QCD", Phys.Rev. D80, 125011 (2009), 0908.4470. • M. Eto, M. Nitta and N. Yamamoto, "Instabilities of Non-Abelian Vortices in Dense QCD", Phys.Rev.Lett. 104, 161601 (2010), 0912.1352. • Y. Hirono, T. Kanazawa and M. Nitta, "Topological Interactions of Non-Abelian Vortices with Quasi-Particles in High Density QCD", Phys.Rev. D83, 085018 (2011), 1012.6042. • W. Vinci, M. Cipriani and M. Nitta, "Spontaneous Magnetization through Non-Abelian Vortex Formation in Rotating Dense Quark Matter", Phys.Rev. D86, 085018 (2012), 1206.3535. • M. Cipriani, W. Vinci and M. Nitta, "Colorful boojums at the interface of a color superconductor", Phys.Rev. D86, 121704 (2012), 1208.5704.
- [5] A. Gorsky, M. Shifman and A. Yung, "Confined Magnetic Monopoles in Dense QCD", Phys.Rev. D83, 085027 (2011), 1101.1120. • M. Eto, M. Nitta and N. Yamamoto, "Confined Monopoles Induced by Quantum Effects in Dense QCD", Phys.Rev. D83, 085005 (2011), 1101.2574.
- [6] D. D. Osheroff, R. C. Richardson and D. M. Lee, "Evidence for a New Phase of Solid He³", Phys. Rev. Lett. 28, 885 (1972). D. D. Osheroff, W. J. Gully, R. C. Richardson and D. M. Lee, "New Magnetic Phenomena in Liquid He³ below 3 mK", Phys. Rev. Lett. 29, 920 (1972).
- [7] A. J. Leggett, "Interpretation of Recent Results on He³ below 3 mK: A New Liquid Phase?", Phys. Rev. Lett. 29, 1227 (1972).

- [8] A. J. Leggett, "A theoretical description of the new phases of liquid ³He", Rev. Mod. Phys. 47, 331 (1975).
- [9] A. J. Leggett, "Quantum Liquids", Oxford University Press, 408 (2006).
- [10] G. E. Volovik, "The Universe in a helium droplet", Int. Ser. Monogr. Phys., 526 (2006).
- [11] E. V. Thuneberg, "Ginzburg-Landau theory of vortices in superfluid ³He-B", Phys. Rev. B 36, 3583 (1987).
- [12] J. A. Sauls and J. W. Serene, "Potential-scattering models for the quasiparticle interactions in liquid ³He", Phys. Rev. B 24, 183 (1981).
- [13] Y. Nambu, "Quasi-Particles and Gauge Invariance in the Theory of Superconductivity", Phys. Rev. 117, 648 (1960).
 J. Goldstone, "Field Theories with Superconductor Solutions", II Nuovo Cimento 19, 154 (1961).
 J. Goldstone, A. Salam and S. Weinberg, "Broken Symmetries", Phys. Rev. 127, 965 (1962).
 R. P. Feynman, "Excitations in liquid helium", Physica 24, Supplement 1, S18 (1958).
- [14] R. P. Feynman, "Application of Quantum Mechanics to Liquid Helium". A. Abrikosov, "On the Magnetic properties of superconductors of the second group", Sov.Phys.JETP 5, 1174 (1957). • Onsager, Nuovo Cimento 6, Supplement 1, 249 (1949).
- [15] M. M. Salomaa and G. E. Volovik, "Quantized vortices in superfluid ³He", Rev. Mod. Phys. 59, 533 (1987).
- [16] E. B. Sonin, "Vortex oscillations and hydrodynamics of rotating superfluids", Rev. Mod. Phys. 59, 87 (1987).
- [17] A. P. Finne, V. B. Eltsov, R. Hänninen, N. B. Kopnin, J. Kopu, M. Krusius, M. Tsubota and G. E. Volovik, "Dynamics of vortices and interfaces in superfluid 3 He", Reports on Progress in Physics 69, 3157 (2006).
 V. P. Mineev, M. M. Salomaa and O. V. Lounasmaa, "Superfluid 3He in rotation", Nature 324, 333 (1986).
- [18] W. Thomson, Philos. Mag. 10, 155 (1880).
- [19] T. P. Simula, T. Mizushima and K. Machida, "Kelvin Waves of Quantized Vortex Lines in Trapped Bose-Einstein Condensates", Phys. Rev. Lett. 101, 020402 (2008).
- [20] E. Fonda, D. P. Meichle, N. T. Ouellette, S. Hormoz, K. R. Sreenivasan and D. P. Lathrop, "Visualization of Kelvin waves on quantum vortices", arXiv:1210.5194.
- [21] V. Bretin, P. Rosenbusch, F. Chevy, G. V. Shlyapnikov and J. Dalibard, "Quadrupole Oscillation of a Single-Vortex Bose-Einstein Condensate: Evidence for Kelvin Modes", Phys. Rev. Lett. 90, 100403 (2003). H. E. Hall, "An Experimental and Theoretical Study of Torsional Oscillations in Uniformly Rotating Liquid Helium II", Proc. R. Soc. A 69, 043617 (2004). Y. Kondo, J. S. Korhonen, M. Krusius, V. V. Dmitriev, E. V. Thuneberg and G. E. Volovik, "Combined spin-mass vortex with soliton tail in superfluid ³B", Phys. Rev. Lett. 68, 3331 (1992).
- [22] E. V. Thuneberg, "Identification of vortices in superfluid ³He B", Phys. Rev. Lett. 56, 359 (1986). ● M. M. Salomaa and G. E. Volovik, "Vortices with spontaneously broken axisymmetry in ³B", Phys. Rev. Lett. 56, 363 (1986).
- [23] M. Krusius, "Experiments on Rotating Superfluid ³He", Physica 126B, 22 (1984).
 J. P. Pekola and J. T. Simola, "Persistent Currents in Superfluid ³He", J. Low Temp. Phys. 58, 555 (1985).
 P. J. Hakonen, M. Krusius, M. M. Salomaa,

J. T. Simola, Y. M. Bunkov, V. P. Mineev and G. E. Volovik, "Magnetic Vortices in Rotating Superfluid ³He-B", Phys. Rev. Lett. 51, 1362 (1983).

- [24] H. B. Nielsen and S. Chadha, "On How to Count Goldstone Bosons", Nucl.Phys. B105, 445 (1976).
- [25] H. Watanabe and H. Murayama, "Unified Description of Nambu-Goldstone Bosons without Lorentz Invariance", Phys.Rev.Lett. 108, 251602 (2012), 1203.0609. • Y. Hidaka, "Counting rule for Nambu-Goldstone modes in nonrelativistic systems", 1203.1494.
- [26] H. Choi, J. P. Davis, J. Pollanen, T. M. Haard and W. P. Halperin, "Strong coupling corrections to the Ginzburg-Landau theory of superfluid ³He", Phys. Rev. B 75, 174503 (2007).
- [27] I. Low and A. V. Manohar, "Spontaneously broken space-time symmetries and Goldstone's theorem", Phys.Rev.Lett. 88, 101602 (2002), hep-th/0110285.
- [28] E. Ivanov and V. Ogievetsky, "The Inverse Higgs Phenomenon in Nonlinear Realizations", Teor.Mat.Fiz. 25, 164 (1975).
 T. Clark, M. Nitta and T. ter Veldhuis, "Brane dynamics from nonlinear realizations", Phys.Rev. D67, 085026 (2003), hep-th/0208184.
- [29] N. Manton, "A Remark on the Scattering of BPS Monopoles", Phys.Lett. B110, 54 (1982).