



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Flavor-phenomenology of two-Higgs-doublet models with generic Yukawa structure

Andreas Crivellin, Christoph Greub, and Ahmet Kokulu

Phys. Rev. D **87**, 094031 — Published 28 May 2013

DOI: [10.1103/PhysRevD.87.094031](https://doi.org/10.1103/PhysRevD.87.094031)

Flavor-phenomenology of two-Higgs-doublet models with generic Yukawa structure

Andreas Crivellin,¹ Christoph Greub,¹ and Ahmet Kokulu¹

¹*Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics,
University of Bern, CH-3012 Bern, Switzerland*

Abstract

In this article, we perform an extensive study of flavor observables in a two-Higgs-doublet model (2HDM) with generic Yukawa structure (of type III). This model is interesting not only because it is the decoupling limit of the Minimal Supersymmetric Standard Model (MSSM) but also because of its rich flavor phenomenology which also allows for sizable effects not only in FCNC processes but also in tauonic B decays. We examine the possible effects in flavor physics and constrain the model both from tree-level processes and from loop-observables. The free parameters of the model are the heavy Higgs mass, $\tan\beta$ (the ratio of vacuum expectation values) and the "non-holomorphic" Yukawa couplings ϵ_{ij}^f ($f = u, d, \ell$). In our analysis we constrain the elements ϵ_{ij}^f in various ways: In a first step we give order of magnitude constraints on ϵ_{ij}^f from 't Hooft's naturalness criterion, finding that all ϵ_{ij}^f must be rather small unless the third generation is involved. In a second step, we constrain the Yukawa structure of the type-III 2HDM from tree-level FCNC processes ($B_{s,d} \rightarrow \mu^+\mu^-$, $K_L \rightarrow \mu^+\mu^-$, $\bar{D}^0 \rightarrow \mu^+\mu^-$, $\Delta F = 2$ processes, $\tau^- \rightarrow \mu^-\mu^+\mu^-$, $\tau^- \rightarrow e^-\mu^+\mu^-$ and $\mu^- \rightarrow e^-e^+e^-$) and observe that all flavor off-diagonal elements of these couplings, except $\epsilon_{32,31}^u$ and $\epsilon_{23,13}^u$ must be very small in order to satisfy the current experimental bounds. In a third step, we consider Higgs mediated loop contributions to FCNC processes ($b \rightarrow s(d)\gamma$, $B_{s,d}$ mixing, $K - \bar{K}$ mixing and $\mu \rightarrow e\gamma$) finding that also ϵ_{13}^u and ϵ_{23}^u must be very small, while the bounds on ϵ_{31}^u and ϵ_{32}^u are especially weak. Furthermore, considering the constraints from electric dipole moments (EDMs) we obtain constraints on some parameters $\epsilon_{ij}^{u,\ell}$. Taking into account the constraints from FCNC processes we study the size of possible effects in the tauonic B decays ($B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$) as well as in $D_{(s)} \rightarrow \tau\nu$, $D_{(s)} \rightarrow \mu\nu$, $K(\pi) \rightarrow e\nu$, $K(\pi) \rightarrow \mu\nu$ and $\tau \rightarrow K(\pi)\nu$ which are all sensitive to tree-level charged Higgs exchange. Interestingly, the unconstrained $\epsilon_{32,31}^u$ are just the elements which directly enter the branching ratios for $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$. We show that they can explain the deviations from the SM predictions in these processes without fine tuning. Furthermore, $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ can even be explained simultaneously. Finally, we give upper limits on the branching ratios of the lepton flavor-violating neutral B meson decays ($B_{s,d} \rightarrow \mu e$, $B_{s,d} \rightarrow \tau e$ and $B_{s,d} \rightarrow \tau\mu$) and correlate the radiative lepton decays ($\tau \rightarrow \mu\gamma$, $\tau \rightarrow e\gamma$ and $\mu \rightarrow e\gamma$) to the corresponding neutral current lepton decays ($\tau^- \rightarrow \mu^-\mu^+\mu^-$, $\tau^- \rightarrow e^-\mu^+\mu^-$ and $\mu^- \rightarrow e^-e^+e^-$). A detailed appendix contains all relevant information for the considered processes for general scalar-fermion-fermion couplings.

I. INTRODUCTION

Two-Higgs-doublet models (2HDMs) [1] have been under intensive investigation for a long time (see for example Ref. [2] for an introduction or Ref. [3] for a recent review article). There are several reasons for this great interest in 2HDMs: Firstly, 2HDMs are very simple extensions of the Standard Model (SM) obtained by just adding an additional scalar $SU(2)_L$ doublet to the SM particle content. This limits the number of new degrees of freedom and makes the model rather predictive. Secondly, motivation for 2HDMs comes from axion models [4] because a possible CP-violating term in the QCD Lagrangian can be rotated away [5] if the Lagrangian has a global $U(1)$ symmetry which is only possible if there are two Higgs doublets. Also the generation of the baryon asymmetry of the Universe motivates the introduction of a second Higgs doublet because in this way the amount of CP violation can be large enough to accommodate for this asymmetry, while the CP violation in the SM is too small [6]. Finally, probably the best motivation for studying 2HDMs is the Minimal Supersymmetric Standard Model (MSSM) where supersymmetry enforces the introduction of a second Higgs doublet [7] due to the holomorphic superpotential. Furthermore, the 2HDM of type III is also the effective theory obtained by integrating out all super-partners of the SM-like particles (the SM fermion, the gauge boson and the Higgs particles of the 2HDM) from MSSM.

2HDMs are not only interesting for direct searches for additional Higgs bosons at colliders. In addition to these high energy searches at the LHC also low-energy precision flavor observables provide a complementary window to physics beyond the SM, i.e. to the 2HDMs. In this respect, FCNC processes, e.g. neutral meson decays to muon pairs ($B_{s(d)} \rightarrow \mu^+\mu^-$, $D \rightarrow \mu^+\mu^-$ and $K_L \rightarrow \mu^+\mu^-$) are especially interesting because they are very sensitive to flavor changing neutral Higgs couplings. However, also charged current processes like tauonic B -meson decays are affected by the charged Higgs boson and $b \rightarrow s\gamma$ provides currently the best lower limit on the charged Higgs mass in the 2HDM of type II.

Recently, tauonic B decays received special attention because the BABAR collaboration performed an analysis of the semileptonic B decays $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ reporting a discrepancy of 2.0σ and 2.7σ from the SM expectation, respectively. The measurements of both decays exceed the SM predictions, and combining them gives a 3.4σ deviation from the SM [8, 9] expectation, which constitutes first evidence for new physics in semileptonic B decays to tau leptons. This evidence for the violation of lepton flavor universality is further supported by the measurement of $B \rightarrow \tau\nu$ by BABAR [10, 11] and BELLE [12, 13] which exceeds the SM prediction by 1.6σ using V_{ub} from the global fit [14]. Assuming that these deviations from the SM are not statistical fluctuations or underestimated theoretical or systematic uncertainties, it is interesting to ask which model of new physics can explain the measured values. Since, a 2HDM of type II cannot explain $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ simultaneously [8], one must look at 2HDMs with more general Yukawa structures.

Also 2HDMs of type III with Minimal Flavor Violation (MFV) [15] cannot explain these deviations from the SM but a 2HDM of type III (where both Higgs doublets couple to up quarks and down quarks as well) with flavor-violation in the up sector, is capable of explaining $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ without fine tuning [16].

These points motivate us to perform a complete analysis of flavor-violation in 2HDMs of type III in this article. For this purpose we take into account all relevant constraints from FCNC processes (both from tree-level contributions and from loop-induced effects) and consider afterwards the possible effects in charged current processes.

This article is structured as follows: In Sec. II, we review the Yukawa Lagrangian of the 2HDM of type III. In Sec. III we give a general overview on the constraints on 2HDMs and update the bounds on the 2HDM of type II. The following sections discuss in detail the constraints on the 2HDM of type III parameter space from 't Hooft's naturalness argument (Sec. IV), from tree-level FCNC processes (Sec. V) and from loop-induced charged and neutral Higgs mediated contributions to the flavor observables (Sec. VI). Sec. VII studies the possible effects in charged current decays ($B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$, $B \rightarrow D^*\tau\nu$, $D_{(s)} \rightarrow \tau\nu$, $D_{(s)} \rightarrow \mu\nu$, $K(\pi) \rightarrow e\nu$, $K(\pi) \rightarrow \mu\nu$, $\tau \rightarrow K(\pi)\nu$) and Sec. VIII is devoted to the study of the upper limits on the branching ratios $B_{s,d} \rightarrow \tau\mu$, $B_{s,d} \rightarrow \tau e$, $B_{s,d} \rightarrow \mu e$ and the correlations among $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, $\tau^- \rightarrow e^- \mu^+ \mu^-$, $\mu^- \rightarrow e^- e^+ e^-$ and $\tau \rightarrow \mu\gamma$, $\tau \rightarrow e\gamma$, $\mu \rightarrow e\gamma$. Finally, we conclude. A detailed appendix contains some of the input parameters used in our analysis, general expressions for some branching ratios as well as all the relevant Wilson coefficients for $b \rightarrow s(d)\gamma$, $\Delta F = 2$ processes, leptonic neutral meson decays ($\Delta F = 1$), LFV transitions, EDMs, anomalous magnetic moment (AMM) of muon and (semi-) leptonic charged meson decays for general charged and/or neutral scalar-fermion-fermion couplings.

II. SETUP

The SM contains only one scalar weak-isospin doublet, the Higgs doublet. After electroweak symmetry breaking its vacuum expectation value ("vev") gives masses to up quarks, down quarks and charged leptons. The charged (CP-odd neutral) component of this doublet becomes the longitudinal component of the W (Z) boson, and thus we have only one physical CP-even neutral Higgs particle in the SM. In a 2HDM we introduce a second Higgs doublet and obtain four additional physical Higgs particles (in the case of a CP conserving Higgs potential): the neutral heavy CP-even Higgs H^0 , a neutral CP-odd Higgs A^0 and the two charged Higgses H^\pm .

As outlined in the introduction we consider a 2HDM with generic Yukawa structure (2HDM of type III). One motivation is that a 2HDM with natural flavor-conservation (like type I or type II) cannot explain $B \rightarrow D\tau\nu$, $B \rightarrow D^*\tau\nu$ and $B \rightarrow \tau\nu$ simultaneously, while the type III model is capable of doing this [16]. Beside this, our calculations in the 2HDM

III are the most general ones in the sense that they can be applied to models with specific flavor-structures like 2HDMs with MFV[15, 17, 18]. In this sense also our bounds are model independent, because they apply to any 2HDM with specific Yukawa structures as well (in the absence of large cancellations which are unlikely). Finally the type-III 2HDM is the decoupling limit of the MSSM and the calculated bounds can be translated to limits on the MSSM parameter space.

The fact that the 2HDM III is the decoupling limit of the MSSM also motivates us to choose for definiteness a MSSM like Higgs potential¹ which automatically avoids dangerous CP violation. The matching of the MSSM on the 2HDM Yukawa sector has been considered in detail. For the MSSM with MFV it was calculated in Ref. [19–24] and for the MSSM with generic flavor structure in Ref. [25] (neglecting the effects of the A -terms) and in Ref. [26] (including the A -terms). Even the next-to-leading order corrections were calculated for the flavor-conserving case in [27] and for the flavor-changing one in the general MSSM in Ref. [28]. Also the one-loop corrections to the Higgs potential have been considered [29–37], but their effects on flavor-observables were found to be small [38].

Following the notation of Ref. [26, 28, 39] we have the following Yukawa Lagrangian in the 2HDM of type III starting in an electroweak basis:

$$\mathcal{L}_Y = \bar{Q}_{fL}^a [Y_{fi}^{d\text{ew}} \epsilon_{ba} H_d^{b*} - \epsilon_{fi}^{d\text{ew}} H_u^a] d_{iR} + \bar{Q}_{fL}^a [Y_{fi}^{u\text{ew}} \epsilon_{ab} H_u^{b*} - \epsilon_{fi}^{u\text{ew}} H_d^a] u_{iR} + h.c. \quad (1)$$

Here a, b denote $SU(2)_L$ -indices, ϵ_{ab} is the two-dimensional antisymmetric tensor with $\epsilon_{12} = -1$ and the Higgs doublets are defined as :

$$\begin{aligned} H_d &= \begin{pmatrix} H_d^1 \\ H_d^2 \end{pmatrix} = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix} \quad \text{with} \quad \langle H_d \rangle = \begin{pmatrix} v_d \\ 0 \end{pmatrix}, \\ H_u &= \begin{pmatrix} H_u^1 \\ H_u^2 \end{pmatrix} = \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix} \quad \text{with} \quad \langle H_u \rangle = \begin{pmatrix} 0 \\ v_u \end{pmatrix}. \end{aligned} \quad (2)$$

Apart from the holomorphic Yukawa-couplings $Y_{fi}^{u\text{ew}}$ and $Y_{fi}^{d\text{ew}}$, we included the non-holomorphic couplings $\epsilon_{fi}^{q\text{ew}}$ ($q = u, d$) as well.

As a next step we decompose the $SU(2)$ doublets into their components and switch to a

¹ If we would require that the Higgs potential possesses a Z_2 symmetry the results would be very similar (for $v \ll m_H$). The heavy Higgs masses squared would still differ by terms of the order of v^2 and only Higgs self-couplings would be different, but they do not enter the flavor-processes at the loop-level under consideration.

basis in which the holomorphic Yukawa couplings are diagonal:

$$\begin{aligned}
\mathcal{L}_Y = & -\bar{d}_{fL} [Y^{d_i} \delta_{fi} H_d^{0*} + \tilde{\epsilon}_{fi}^d H_u^0] d_{iR} - \bar{u}_{fL} [Y^{u_i} \delta_{fi} H_u^{0*} + \tilde{\epsilon}_{fi}^u H_d^0] u_{iR} \\
& + \bar{u}_{fL} V_{fj} [Y^{d_i} \delta_{ji} - \cot \beta \tilde{\epsilon}_{ji}^d] H_d^{2*} d_{iR} \\
& + \bar{d}_{fL} V_{jf}^* [Y^{u_i} \delta_{ji} - \tan \beta \tilde{\epsilon}_{ji}^u] H_u^{1*} u_{iR} + h.c. .
\end{aligned} \tag{3}$$

where $\tan \beta = v_u/v_d$ is the ratio of the vacuum expectation values v_u and v_d acquired by H_u and H_d , respectively. We perform this intermediate step, because this is the basis which corresponds to the super-CKM basis of the MSSM and the couplings $\tilde{\epsilon}_{ij}^d$ can be directly related to loop-induced non-holomorphic Higgs coupling. The wave-function rotations $U_{fi}^{qL,R}$ necessary to arrive at the physical basis with diagonal quark mass matrices are defined by

$$U_{jf}^{qL*} m_{jk}^q U_{ki}^{qR} = m_{qi} \delta_{fi} . \tag{4}$$

They modify the Yukawa Lagrangian as follows:

$$\begin{aligned}
\mathcal{L}_Y = & -\bar{d}_{fL} \left[\left(\frac{m_{d_i}}{v_d} \delta_{fi} - \epsilon_{fi}^d \tan \beta \right) H_d^{1*} + \epsilon_{fi}^d H_u^2 \right] d_{iR} \\
& - \bar{u}_{fL} \left[\left(\frac{m_{u_i}}{v_u} \delta_{fi} - \epsilon_{fi}^u \cot \beta \right) H_u^{2*} + \epsilon_{fi}^u H_d^1 \right] u_{iR} \\
& + \bar{u}_{fL} V_{fj} \left[\frac{m_{d_i}}{v_d} \delta_{ji} - (\cot \beta + \tan \beta) \epsilon_{ji}^d \right] H_d^{2*} d_{iR} \\
& + \bar{d}_{fL} V_{jf}^* \left[\frac{m_{u_i}}{v_u} \delta_{ji} - (\tan \beta + \cot \beta) \epsilon_{ji}^u \right] H_u^{1*} u_{iR} + h.c. .
\end{aligned} \tag{5}$$

Here, m_{q_i} are the physical running quark masses, H_q^1 and H_q^2 are the components of the Higgs doublets, and

$$V_{fi} = U_{jf}^{uL*} U_{ji}^{dL} , \tag{6}$$

is the CKM matrix. The Higgs doublets H_u and H_d project onto the physical mass eigenstates H^0 (heavy CP-even Higgs), h^0 (light CP-even Higgs), A^0 (CP-odd Higgs) and H^\pm in the following way:

$$\begin{aligned}
H_u^0 &= \frac{1}{\sqrt{2}} (H^0 \sin \alpha + h^0 \cos \alpha + iA^0 \cos \beta) , \\
H_d^0 &= \frac{1}{\sqrt{2}} (H^0 \cos \alpha - h^0 \sin \alpha + iA^0 \sin \beta) , \\
H_u^1 &= \cos \beta H^+ , \\
H_d^2 &= \sin \beta H^- ,
\end{aligned} \tag{7}$$

where α is the mixing angle necessary to diagonalize the neutral CP-even Higgs mass matrix

(see e.g. [40])². Since we assume a MSSM-like Higgs potential³ we have

$$\begin{aligned}\tan \beta &= \frac{v_u}{v_d}, \\ \tan 2\alpha &= \tan 2\beta \frac{m_{A^0}^2 + M_Z^2}{m_{A^0}^2 - M_Z^2}, \\ m_{H^\pm}^2 &= m_{A^0}^2 + M_W^2, \quad m_{H^0}^2 = m_{A^0}^2 + M_Z^2 - m_{h^0}^2,\end{aligned}\tag{8}$$

with $-\frac{\pi}{2} < \alpha < 0$ and $0 < \beta < \frac{\pi}{2}$.

This means that in the phenomenologically interesting and viable limit of large values of $\tan \beta$ and $v \ll m_{A^0}$ we have to a good approximation⁴:

$$\begin{aligned}\tan \beta &\approx -\cot \alpha, \\ m_{H^0} &\approx m_{H^\pm} \approx m_{A^0} \equiv m_H.\end{aligned}\tag{9}$$

Without the non-holomorphic corrections ϵ_{ij}^q , the rotation matrices $U_{fi}^{qL,R}$ would simultaneously diagonalize the mass terms and the neutral Higgs couplings in Eq. (5). However, in the presence of non-holomorphic corrections, this is no longer the case and flavor changing neutral Higgs couplings are present in the basis in which the physical quark mass matrices are diagonal.

The Yukawa Lagrangian in Eq. (5) leads to the following Feynman rules⁵ for Higgs-quark-quark couplings

$$i \left(\Gamma_{q_f q_i}^{LRH} P_R + \Gamma_{q_f q_i}^{RLH} P_L \right)\tag{10}$$

with

$$\begin{aligned}\Gamma_{u_f u_i}^{LRH_k^0} &= x_u^k \left(\frac{m_{u_i}}{v_u} \delta_{fi} - \epsilon_{fi}^u \cot \beta \right) + x_d^{k*} \epsilon_{fi}^u, \\ \Gamma_{d_f d_i}^{LRH_k^0} &= x_d^k \left(\frac{m_{d_i}}{v_d} \delta_{fi} - \epsilon_{fi}^d \tan \beta \right) + x_u^{k*} \epsilon_{fi}^d, \\ \Gamma_{u_f d_i}^{LRH^\pm} &= \sum_{j=1}^3 \sin \beta V_{fj} \left(\frac{m_{d_i}}{v_d} \delta_{ji} - \epsilon_{ji}^d \tan \beta \right), \\ \Gamma_{d_f u_i}^{LRH^\pm} &= \sum_{j=1}^3 \cos \beta V_{jf}^* \left(\frac{m_{u_i}}{v_u} \delta_{ji} - \epsilon_{ji}^u \tan \beta \right).\end{aligned}\tag{11}$$

² Note that we defined α as common in the MSSM. In the 2HDM also a convention with a doubled range for α is used.

³ MSSM-like Higgs potential implies that in the large $\tan \beta$ limit and for $v \ll m_H$ the charged Higgs mass m_{H^\pm} , the heavy CP even Higgs mass m_{H^0} and the CP odd Higgs mass m_{A^0} are equal.

⁴ For the SM-like Higgs boson h^0 we use $m_{h^0} \approx 125$ GeV in our numerical analysis.

⁵ Hermiticity of the Lagrangian implies the relation $\Gamma_{q_f q_i}^{RLH} = \Gamma_{q_i q_f}^{LRH^*}$.

Similarly, for the lepton case, the non-vanishing effective Higgs vertices are

$$\begin{aligned}\Gamma_{\ell_f \ell_i}^{LRH_k^0} &= x_d^k \left(\frac{m_{\ell_i}}{v_d} \delta_{fi} - \epsilon_{fi}^\ell \tan \beta \right) + x_u^{k*} \epsilon_{fi}^\ell, \\ \Gamma_{\nu_f \ell_i}^{LRH^\pm} &= \sum_{j=1}^3 \sin \beta V_{fj}^{\text{PMNS}} \left(\frac{m_{\ell_i}}{v_d} \delta_{ji} - \epsilon_{ji}^\ell \tan \beta \right).\end{aligned}\tag{12}$$

Here, $H_k^0 = (H^0, h^0, A^0)$ and the coefficients x_q^k are given by

$$\begin{aligned}x_u^k &= \left(-\frac{1}{\sqrt{2}} \sin \alpha, -\frac{1}{\sqrt{2}} \cos \alpha, \frac{i}{\sqrt{2}} \cos \beta \right), \\ x_d^k &= \left(-\frac{1}{\sqrt{2}} \cos \alpha, \frac{1}{\sqrt{2}} \sin \alpha, \frac{i}{\sqrt{2}} \sin \beta \right).\end{aligned}\tag{13}$$

This means that flavor-violation (beyond the one already present in the 2HDM of type II) is entirely governed by the couplings $\epsilon_{ij}^{q,\ell}$. If one wants to make the connection to the MSSM, the parameters $\epsilon_{ij}^{q,\ell}$ will depend only on SUSY breaking parameters and $\tan \beta$.

III. CONSTRAINTS ON THE 2HDM PARAMETER SPACE – GENERAL DISCUSSION AND OVERVIEW

In this section we give an overview on flavor observables sensitive to charged Higgs contributions. We review the constraints on the 2HDM of type II and discuss to which extent these bounds will hold in the 2HDM of type III. A detailed analysis of flavor constraints on the type-III 2HDM parameter space will be given in the following sections.

The most common version of 2HDMs, concerning its Yukawa sector, is the 2HDM of type II which respects natural flavor conservation [41] by requiring that one Higgs doublet couples only to up-quarks while the other one gives masses to down-type quarks and charged leptons (like the MSSM at tree-level). Flavor-observables in 2HDMs of type II have been studied in detail [42–44]. In the type II model there are no tree-level flavor-changing neutral currents and all flavor violation is induced by the CKM matrix entering the charged Higgs vertex. In this way the constraints from FCNC processes can be partially avoided. This is true for $\Delta F = 2$ processes where the charged Higgs contribution is small, for $K_L \rightarrow \mu^+ \mu^-$, $D^0 \rightarrow \mu^+ \mu^-$ (due to the tiny Higgs couplings to light quarks) and all flavor observables in the lepton sector. However, the FCNC processes $b \rightarrow s \gamma$ (also to less extent $b \rightarrow d \gamma$) and $B_s \rightarrow \mu^+ \mu^-$ are sensitive to the charged Higgs contributions. In addition, direct searches at the LHC and charged current processes restrict the type-II 2HDM parameter space.

Among the FCNC processes, the constraints from $b \rightarrow s \gamma$ are most stringent due to the necessarily constructive interference with the SM contribution [45–48]. The most recent lower bound on the charged Higgs mass obtained in Ref. [49] is $m_{H^\pm} \geq 360 \text{ GeV}$ which

includes NNLO QCD corrections and is rather independent of $\tan\beta$. In the type-III 2HDM this lower bound on the charged Higgs mass can be weakened due to destructive interference with contributions involving ϵ_{ij}^q . Also in $B_s \rightarrow \mu^+\mu^-$ (and $B_d \rightarrow \mu^+\mu^-$) a sizable loop-induced effect is possible in the 2HDM II, but the constraints are still not very stringent even if the new LHCb measurements are used. The reason for this is that, taking into account the constraints from $b \rightarrow s\gamma$ on the charged Higgs mass, the branching ratio for $B_s \rightarrow \mu^+\mu^-$ in the 2HDM II is even below the SM expectation for larger values of $\tan\beta$ [50–52] due to the destructive interference between the charged Higgs and the SM contribution.

Regarding charged current processes, tauonic B decays are currently most sensitive to charged Higgs effects. Here, the charged-Higgs contribution in the type-II 2HDM to $B \rightarrow \tau\nu$ interferes destructively with the SM contribution [53, 54]. The same is true for $B \rightarrow D^*\tau\nu$ [55] and $B \rightarrow D\tau\nu$ [42, 56, 57]. As outlined in the introduction this leads to the fact that the 2HDM II cannot explain $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ simultaneously [8]. Other charged current observables sensitive to charged Higgses are $D_{(s)} \rightarrow \mu\nu$, $D_{(s)} \rightarrow \tau\nu$ [58–60], $\tau \rightarrow K(\pi)\nu$ and $K \rightarrow \mu\nu/\pi \rightarrow \mu\nu$ [61] (see [44] for a global analysis).

Fig. 1 shows our updated constraints on the 2HDM II parameter space from $b \rightarrow s\gamma$, $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$, $B \rightarrow D^*\tau\nu$, $B_s \rightarrow \mu^+\mu^-$ and $K \rightarrow \mu\nu/\pi \rightarrow \mu\nu$. We see that in order to get agreement within 2σ between the theory prediction and the measurement of $B \rightarrow D^*\tau\nu$, large values of $\tan\beta$ and light Higgs masses would be required which is in conflict with all other processes under consideration.

Concerning direct searches the bounds on the charged Higgs mass are rather weak due to the large background from W events. The search for neutral Higgs bosons is easier and the CMS bounds⁶ on m_{A^0} from $A^0 \rightarrow \tau^+\tau^-$ are shown in Fig. 2. These bounds were obtained in the MSSM, but since the MSSM corrections to $A^0 \rightarrow \tau^+\tau^-$ are rather small and since we consider a MSSM-like Higgs potential, these bounds also hold in the 2HDM III as long as the Peccei-Quinn symmetry breaking in the lepton sector is small⁷.

Going beyond the simple Yukawa structure of the 2HDM of type II, also 2HDMs of type III with MFV [15, 17, 18], alignment [64, 65] or natural flavor conservation [17, 41] have been analyzed in detail. However, flavor-observables in type III models with generic flavor-structure have received much less attention. Ref. [66] considered the possible effects of the flavor-diagonal terms and Ref. [67] considers leptonic observables. As outlined in the introduction, 2HDMs of type II (or type III with MFV) cannot explain $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ simultaneously [8] (and for $B \rightarrow \tau\nu$ fine tuning is needed [18]).

In the following sections we will study in detail the flavor-observables in the 2HDM with generic flavor-structure [68], but for definiteness, with MSSM-like Higgs potential. For this

⁶ Note that we did not use the bounds from unpublished CMS update of the $A^0 \rightarrow \tau^+\tau^-$ analysis.

⁷ For a global analysis of electroweak precision constraints see for example Ref. [62].




FIG. 1: Updated constraints on the 2HDM of type II parameter space. The regions compatible with experiment are shown (the regions are superimposed on each other): $b \rightarrow s\gamma$ (yellow), $B \rightarrow D\tau\nu$ (green), $B \rightarrow \tau\nu$ (red), $B_s \rightarrow \mu^+\mu^-$ (orange), $K \rightarrow \mu\nu/\pi \rightarrow \mu\nu$ (blue) and $B \rightarrow D^*\tau\nu$ (black). Note that no region in parameter space is compatible with all processes. Explaining $B \rightarrow D^*\tau\nu$ would require very small Higgs masses and large values of $\tan\beta$ which is not compatible with the other observables. To obtain this plot, we added the theoretical uncertainty linear on the top of the 2σ experimental error.

purpose, all processes described above are relevant. In addition, $\Delta F = 2$ processes, lepton flavor violating observables (LFV), EDMs, $\tau \rightarrow K(\pi)\nu/K(\pi) \rightarrow \mu\nu$ and $K \rightarrow \mu(e)\nu/\pi \rightarrow \mu(e)\nu$ will turn out to give information on the flavor structure of the 2HDM of type III. Furthermore, we will investigate to which extent contributions to $B_{s,d} \rightarrow \tau\mu$, $B_{s,d} \rightarrow \tau e$, $B_{s,d} \rightarrow \mu e$ and muon anomalous magnetic moment are possible.

IV. CONSTRAINTS FROM 'T HOOFT'S NATURALNESS CRITERION

The naturalness criterion of 't Hooft states that the smallness of a quantity is only natural if a symmetry is gained in the limit in which this quantity is zero. This means on the other hand that large accidental cancellations, which are not enforced by a symmetry, are unnatural and thus not desirable. Let us apply this reasoning to the fermion mass matrices in the 2HDM. We recall from the last section the expressions for the fermion mass matrices




FIG. 2: Plot from the CMS collaboration taken from Ref. [63]: Exclusion limits in the m_{A^0} - $\tan\beta$ plane from $A^0 \rightarrow \tau^+\tau^-$. The analysis was done in the MSSM, but since we consider a 2HDM with MSSM-like Higgs potential and the MSSM corrections to the $A^0\tau\tau$ vertex are small, we can apply this bound to our model. However, a large value of ϵ_{33}^ℓ in the 2HDM of type III could affect the conclusions. Note that in the limit $v \ll m_H$ all heavy Higgs masses (m_{H^0} , m_{A^0} and m_{H^\pm}) are approximately equal.

in the electroweak basis:

$$\begin{aligned}
 m_{ij}^d &= v_d Y_{ij}^{d\text{ew}} + v_u \epsilon_{ij}^{d\text{ew}}, \\
 m_{ij}^u &= v_u Y_{ij}^{u\text{ew}} + v_d \epsilon_{ij}^{u\text{ew}}, \\
 m_{ij}^\ell &= v_d Y_{ij}^{\ell\text{ew}} + v_u \epsilon_{ij}^{\ell\text{ew}}.
 \end{aligned}
 \tag{14}$$

Diagonalizing these fermion mass matrices gives the physical fermion masses and the CKM matrix. Using 't Hooft's naturalness criterion we can demand the absence of fine-tuned cancellations between $v_d Y_{ij}^{d,\ell}$ ($v_u Y_{ij}^u$) and $v_u \epsilon_{ij}^{d,\ell}$ ($v_d \epsilon_{ij}^u$). Thus, we require that the contributions of $v_u \epsilon_{ij}^{d,\ell}$ and $v_d \epsilon_{ij}^u$ to the fermion masses and CKM matrix do not exceed the physical measured quantities.

In first order of a perturbative diagonalization of the fermion mass matrices, the diagonal elements m_{ii}^f give rise to the fermion masses, while (in our conventions) the elements m_{ij}^f with $i < j$ ($i > j$) affect the left-handed (right-handed) rotations necessary to diagonalize the fermion mass matrices. The left-handed rotations of the quark fields are linked to the CKM matrix and can therefore be constrained by demanding that the physical CKM matrix

is generated without a significant degree of fine-tuning. However, the right-handed rotations of the quarks are not known and the mixing angles of the PMNS matrix are big so that for these two cases we can only demand that the fermion masses are generated without too large accidental cancellations. Note, that in Eq. (14) the elements $\epsilon_{ij}^{f\text{ew}}$ enter, while the elements ϵ_{ij}^f which we want to constrain from flavor observables are given in the physical basis with diagonal fermion masses. This means that in order to constrain ϵ_{ij}^f from 't Hooft's naturalness criterion we have to assume in addition that no accidental cancellation occur by switching between the electroweak basis and the physical basis. In conclusion this leads to the following upper bounds

$$\begin{aligned}
|v_{u(d)}\epsilon_{ij}^{d(u)}| &\leq |V_{ij}^{\text{CKM}}| \times \max [m_{d_i(u_i)}, m_{d_j(u_j)}] \text{ for } i < j, \\
|v_{u(d)}\epsilon_{ij}^{d(u)}| &\leq \max [m_{d_i(u_i)}, m_{d_j(u_j)}] \text{ for } i \geq j, \\
|v_u\epsilon_{ij}^\ell| &\leq \max [m_{\ell_i}, m_{\ell_j}].
\end{aligned} \tag{15}$$

In the large $\tan\beta$ limit, inserting the quark masses $m_q(\mu)$ at the Higgs scale (which we choose here to be $\mu_{\text{Higgs}} = 500 \text{ GeV}$), we can immediately read off the upper bounds on $\epsilon_{ij}^{u,d,\ell}$ from Eq. (15):

$$\begin{aligned}
|\epsilon_{ij}^d| &\leq \begin{pmatrix} 1.3 \times 10^{-4} & 5.8 \times 10^{-5} & 5.1 \times 10^{-5} \\ 2.6 \times 10^{-4} & 2.6 \times 10^{-4} & 5.9 \times 10^{-4} \\ 1.4 \times 10^{-2} & 1.4 \times 10^{-2} & 1.4 \times 10^{-2} \end{pmatrix}_{ij}, \\
|\epsilon_{ij}^u| &\leq (\tan\beta/50) \begin{pmatrix} 3.4 \times 10^{-4} & 3.2 \times 10^{-2} & 1.6 \times 10^{-1} \\ 1.4 \times 10^{-1} & 1.4 \times 10^{-1} & 1.9 \\ - & - & - \end{pmatrix}_{ij}, \\
|\epsilon_{ij}^\ell| &\leq \begin{pmatrix} 2.9 \times 10^{-6} & 6.1 \times 10^{-4} & 1.0 \times 10^{-2} \\ 6.1 \times 10^{-4} & 6.1 \times 10^{-4} & 1.0 \times 10^{-2} \\ 1.0 \times 10^{-2} & 1.0 \times 10^{-2} & 1.0 \times 10^{-2} \end{pmatrix}_{ij}.
\end{aligned} \tag{16}$$

Of course, these constraints are not strict bounds in the sense that they must be respected in any viable model. Anyway, big violation of naturalness is not desirable and Eq. (16) gives us a first glance on the possible structure of the elements ϵ_{ij}^f . As we will see later, it is possible to explain $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ using $\epsilon_{31,32}^u$ without violating Eq. (16), while if one wants to explain $B \rightarrow \tau\nu$ with ϵ_{33}^d 't Hooft's naturalness criterion is violated.

V. CONSTRAINTS FROM TREE-LEVEL NEUTRAL-CURRENT PROCESSES

The flavor off-diagonal elements ϵ_{ij}^f (with $i \neq j$) give rise to flavor-changing neutral currents (FCNCs) already at the tree-level. Comparing the Higgs contributions to the loop-suppressed SM contributions, large effects are in principle possible. However, all experimental results are in very good agreement with SM predictions, which put extremely stringent constraints on the non-holomorphic terms ϵ_{ij}^f .

In this section we consider three different kinds of processes:

- Muonic decays of neutral mesons ($B_{s,d} \rightarrow \mu^+\mu^-$, $K_L \rightarrow \mu^+\mu^-$ and $\bar{D}^0 \rightarrow \mu^+\mu^-$).
- $\Delta F = 2$ processes ($D-\bar{D}$, $K-\bar{K}$, $B_s-\bar{B}_s$ and $B_d-\bar{B}_d$ mixing).
- Flavor changing lepton decays ($\tau^- \rightarrow \mu^-\mu^+\mu^-$, $\tau^- \rightarrow e^-\mu^+\mu^-$ and $\mu^- \rightarrow e^-e^+e^-$).

As we will see in detail in Sec. VA, the leptonic neutral meson decays $B_{s,d} \rightarrow \mu^+\mu^-$, $K_L \rightarrow \mu^+\mu^-$ and $\bar{D}^0 \rightarrow \mu^+\mu^-$ put constraints on the elements ϵ_{ij}^d (with $i \neq j$) and $\epsilon_{12,21}^u$ already if one of these elements is non-zero, while $B_d-\bar{B}_d$, $B_s-\bar{B}_s$, $K-\bar{K}$ and $D-\bar{D}$ mixing only provide constraints on the products $\epsilon_{ij}^d\epsilon_{ji}^{d*}$ and $\epsilon_{12}^u\epsilon_{21}^{u*}$ (Sec. VB). This means that the constraints on $\Delta F = 2$ processes can be avoided if one element of the product $\epsilon_{ij}^q\epsilon_{ji}^{q*}$ is zero, while the constraints from the leptonic neutral meson decays can only be avoided if the Peccei Quinn symmetry breaking for the leptons is large such that $\epsilon_{22}^\ell \approx m_\mu/v_u$ is possible.

In Sec. VC we will consider the flavor changing lepton decays $\tau^- \rightarrow \mu^-\mu^+\mu^-$, $\tau^- \rightarrow e^-\mu^+\mu^-$ and $\mu^- \rightarrow e^-e^+e^-$ which constrain the off-diagonal elements $\epsilon_{23,32}^\ell$, $\epsilon_{13,31}^\ell$ and $\epsilon_{12,21}^\ell$, respectively.

A. Leptonic neutral meson decays: $B_{s,d} \rightarrow \mu^+\mu^-$, $K_L \rightarrow \mu^+\mu^-$ and $\bar{D}^0 \rightarrow \mu^+\mu^-$

Muonic decays of neutral mesons ($B_s \rightarrow \mu^+\mu^-$, $B_d \rightarrow \mu^+\mu^-$, $K_L \rightarrow \mu^+\mu^-$ and $\bar{D}^0 \rightarrow \mu^+\mu^-$) are strongly suppressed in the SM for three reasons: they are loop-induced, helicity suppressed and they involve small CKM elements. Therefore, their branching ratios (in the SM) are very small and in fact only $K_L \rightarrow \mu^+\mu^-$ and recently also $B_s \rightarrow \mu^+\mu^-$ [69] have been measured, while for the other decays only upper limits on the branching ratios exist (see Table I). We do not consider decays to electrons (which are even stronger helicity suppressed) nor $B_{d,s} \rightarrow \tau^+\tau^-$ (where the tau leptons are difficult to reconstruct) because the experimental limits are even weaker. The study of meson decays to lepton flavor-violating final states is postponed to Sec. VIII.

We see from Fig. 3 that the off-diagonal elements of $\epsilon_{13,31}^d$, $\epsilon_{23,32}^d$, $\epsilon_{12,21}^d$ and $\epsilon_{12,21}^u$ directly give rise to tree-level neutral Higgs contributions to $B_d \rightarrow \mu^+\mu^-$, $B_s \rightarrow \mu^+\mu^-$, $K_L \rightarrow \mu^+\mu^-$ and $\bar{D}^0 \rightarrow \mu^+\mu^-$, respectively.

Process	Experimental value	SM prediction
$\mathcal{B}[B_s \rightarrow \mu^+\mu^-]$	$3.2^{+1.5}_{-1.2} \times 10^{-9}$ [69]	$(3.23 \pm 0.27) \times 10^{-9}$ [70]
$\mathcal{B}[B_d \rightarrow \mu^+\mu^-]$	$\leq 9.4 \times 10^{-10}$ (95% CL) [69]	$(1.07 \pm 0.10) \times 10^{-10}$ [70]
$\mathcal{B}[K_L \rightarrow \mu^+\mu^-]_{\text{short}}$	$\leq 2.5 \times 10^{-9}$ [71]	$\approx 0.9 \times 10^{-9}$ [71]
$\mathcal{B}[D^0 \rightarrow \mu^+\mu^-]$	$\leq 1.4 \times 10^{-7}$ (90% CL) [72]	–

TABLE I: Experimental values and SM predictions for the branching ratios of neutral meson decays to muon pairs. For $K_L \rightarrow \mu^+\mu^-$ we only give the upper limit on the computable short distance contribution [71] extracted from the experimental value $(6.84 \pm 0.11) \times 10^{-9}$ (90% CL) [72]. The SM prediction for $D^0 \rightarrow \mu^+\mu^-$ cannot be reliably calculated due to hadronic uncertainties.




FIG. 3: Feynman diagram showing the neutral Higgs contribution to $B_{s,d} \rightarrow \mu^+\mu^-$, $K_L \rightarrow \mu^+\mu^-$ and $\bar{D}^0 \rightarrow \mu^+\mu^-$.

In principle, the constraints from these processes could be weakened, or even avoided, if $\epsilon_{22}^\ell \approx m_{\ell_2}/v_u$. Anyway, in this section we will assume that the Peccei Quinn breaking for the leptons is small and neglect the effect of ϵ_{22}^ℓ in our numerical analysis for setting limits on ϵ_{ij}^q .

1. $B_{s,d} \rightarrow \mu^+\mu^-$

For definiteness, consider the decay of a neutral B_s ($\bar{b}s$) meson (the corresponding decay of a B_d meson follow trivially by replacing s with d and 2 with 1) to a muon pair. The

effective Hamiltonian governing this transition is⁸

$$\mathcal{H}_{\text{eff}}^{B_s \rightarrow \mu^+ \mu^-} = -\frac{G_F^2 M_W^2}{\pi^2} [C_A^{bs} O_A^{bs} + C_S^{bs} O_S^{bs} + C_P^{bs} O_P^{bs} + C_A^{\prime bs} O_A^{\prime bs} + C_S^{\prime bs} O_S^{\prime bs} + C_P^{\prime bs} O_P^{\prime bs}] + h.c., \quad (17)$$

where the operators are defined as

$$\begin{aligned} O_A^{bs} &= (\bar{b} \gamma_\mu P_L s) (\bar{\mu} \gamma^\mu \gamma_5 \mu), \\ O_S^{bs} &= (\bar{b} P_L s) (\bar{\mu} \mu), \\ O_P^{bs} &= (\bar{b} P_L s) (\bar{\mu} \gamma_5 \mu), \end{aligned} \quad (18)$$

and the primed operators are obtained replacing P_L with P_R . The corresponding expression for the branching ratio in terms of the Wilson coefficients reads

$$\begin{aligned} \mathcal{B}[B_s \rightarrow \mu^+ \mu^-] &= \frac{G_F^4 M_W^4}{8\pi^5} \sqrt{1 - 4 \frac{m_\mu^2}{M_{B_s}^2}} M_{B_s} f_{B_s}^2 m_\mu^2 \tau_{B_s} \\ &\times \left[\left| \frac{M_{B_s}^2 (C_P^{bs} - C_P^{\prime bs})}{2(m_b + m_s) m_\mu} - (C_A^{bs} - C_A^{\prime bs}) \right|^2 + \left| \frac{M_{B_s}^2 (C_S^{bs} - C_S^{\prime bs})}{2(m_b + m_s) m_\mu} \right|^2 \times \left(1 - 4 \frac{m_\mu^2}{m_{B_s}^2} \right) \right]. \end{aligned} \quad (19)$$

Concerning the running of the Wilson coefficients due to the strong interaction, the operators O_A^{bs} and $O_A^{\prime bs}$ correspond to conserved vector currents with vanishing anomalous dimensions. This means that their Wilson coefficients are scale independent. The scalar and pseudo-scalar Wilson coefficients C_S^{bs} and C_P^{bs} ($C_S^{\prime bs}$ and $C_P^{\prime bs}$) have the same anomalous dimension as quark masses in the SM which means that their scale dependence is given by:

$$C_{S,P}^{(\prime)bs}(\mu_{low}) = \frac{m_q(\mu_{low})}{m_q(\mu_{high})} C_{S,P}^{(\prime)bs}(\mu_{high}), \quad (20)$$

where m_q is the running quark mass with the appropriate number of active flavors. In the SM, C_A is the only non-vanishing Wilson coefficient

$$C_A^{bs} = -V_{tb}^* V_{ts} Y \left(\frac{m_t^2}{M_W^2} \right) - V_{cb}^* V_{cs} Y \left(\frac{m_c^2}{M_W^2} \right), \quad (21)$$

where the function Y is defined as $Y = \eta_Y Y_0$ such that the NLO QCD effects are included in $\eta_Y = 1.0113$ [70] and the one loop Inami-Lim function Y_0 reads [73]

$$Y_0(x) = \frac{x}{8} \left[\frac{4-x}{1-x} + \frac{3x}{(1-x)^2} \ln(x) \right]. \quad (22)$$

The complete Wilson coefficients for general quark-quark-scalar couplings are given in the appendix. In the 2HDM of type III, in the case of large $\tan \beta$ and $v \ll m_H$, the terms

⁸ The complete expression for the Hamiltonian and the branching ratio including lepton flavor-violating final states is given in the appendix.

involving ϵ_{ij}^q simplify to

$$\begin{aligned} C_S^{bs} = C_P^{bs} &= -\frac{\pi^2}{G_F^2 M_W^2} \frac{1}{2m_H^2} \frac{m_{\ell_2} - v_u \epsilon_{22}^\ell}{v} \epsilon_{23}^{d*} \tan^2 \beta, \\ C_S^{bs} = -C_P^{bs} &= -\frac{\pi^2}{G_F^2 M_W^2} \frac{1}{2m_H^2} \frac{m_{\ell_2} - v_u \epsilon_{22}^\ell}{v} \epsilon_{32}^d \tan^2 \beta. \end{aligned} \quad (23)$$

To these Wilson coefficients the well known loop-induced type II 2HDM contributions⁹

$$C_S^{bs} = C_P^{bs} = -\frac{m_b V_{tb}^* V_{ts}}{2} \frac{m_\mu}{2M_W^2} \tan^2 \beta \frac{\log(m_H^2/m_t^2)}{m_H^2/m_t^2 - 1}, \quad (24)$$

have to be added as well [52]. Note that since we give the Wilson coefficients at the matching scale, also m_b and m_t must be evaluated at this scale.

We can now constrain the elements $\epsilon_{23,32}^d$ and $\epsilon_{13,31}^d$ by demanding that the experimental bounds are satisfied within two standard deviations for $B_s \rightarrow \mu^+ \mu^-$ or equivalently at the 95% CL concerning $B_d \rightarrow \mu^+ \mu^-$. The results for the constraints on ϵ_{23}^d and ϵ_{32}^d (ϵ_{13}^d and ϵ_{31}^d) from $B_s \rightarrow \mu^+ \mu^-$ ($B_d \rightarrow \mu^+ \mu^-$) are shown in Fig. 4 (Fig. 5).

All constraints on $\epsilon_{13,31}^d$ and $\epsilon_{23,32}^d$ are very stringent; of the order of 10^{-5} . Both an enhancement or a suppression of $\mathcal{B}[B_{d,s} \rightarrow \mu^+ \mu^-]$ compared to the SM prediction is possible. While in the 2HDM II the minimal value for $\mathcal{B}[B_{d,s} \rightarrow \mu^+ \mu^-]$ is half the SM prediction, in the 2HDM III also a bigger suppression of $B_{d,s} \rightarrow \mu^+ \mu^-$ is possible if $\epsilon_{13,23}^d \neq 0$. In principle, the constraints on ϵ_{23}^d (ϵ_{13}^d) from $B_{s(d)} \rightarrow \mu^+ \mu^-$ are not independent of ϵ_{32}^d (ϵ_{31}^d). Anyway, in the next section it will turn out that the constraints from $\Delta F = 2$ processes are more stringent if both ϵ_{32}^d and ϵ_{23}^d are different from zero (the same conclusions hold for $\epsilon_{31,13}^d$, $\epsilon_{21,12}^d$ and $\epsilon_{21,12}^u$).

$B_s \rightarrow \mu^+ \mu^-$ and $B_d \rightarrow \mu^+ \mu^-$ can also be used to constrain the leptonic parameter ϵ_{22}^ℓ . We will discuss the corresponding subject in Sec.VI.

2. $K_L \rightarrow \mu^+ \mu^-$

Concerning $K_L \rightarrow \mu^+ \mu^-$, the branching ratio and the Wilson coefficients can be obtained by a simple replacement of indices from Eq. (19), Eq. (21) and Eq. (23). Due to the presence of large non-perturbative QCD effects, we require that the 2HDM III contribution together with the short distance piece of the SM contribution does not exceed the upper limit on the short distance contribution to the branching ratio calculated in Ref. [71]. The resulting constraints on $\epsilon_{12,21}^d$ are shown in Fig. 6. They are found to be extremely stringent (of the order of 10^{-6}).

⁹ Since we want to put constraints on the elements $\epsilon_{13,23}^d$ we assume that the loop-induced 2HDM II contribution is not changed by elements ϵ_{i3}^u or ϵ_{33}^d .




FIG. 4: Allowed regions in the complex $\epsilon_{23,32}^d$ -plane from $B_s \rightarrow \mu^+ \mu^-$ for $\tan \beta = 30$, $\tan \beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue). Note that the allowed regions for ϵ_{32}^d -plane are not full circles because in this case a suppression of $\mathcal{B}[B_s \rightarrow \mu^+ \mu^-]$ below the experimental lower bound is possible.

3. $\bar{D}^0 \rightarrow \mu^+ \mu^-$

The analogous expressions for the branching ratio for $\bar{D}^0 \rightarrow \mu^+ \mu^-$ ($\bar{D}^0(\bar{c}u)$) follow by a straightforward replacement of indices in Eq. (19) but the Wilson coefficients in the type-III 2HDM for $\bar{D}^0 \rightarrow \mu^+ \mu^-$ have a different dependence on $\tan \beta$:

$$\begin{aligned}
 C_S^{cu} = -C_P^{cu} &= \frac{\pi^2}{G_F^2 M_W^2} \frac{1}{2m_H^2} \frac{m_{\ell_2} - v_u \epsilon_{22}^\ell}{v} \epsilon_{12}^{u*} \tan \beta, \\
 C_S^{tcu} = C_P^{tcu} &= \frac{\pi^2}{G_F^2 M_W^2} \frac{1}{2m_H^2} \frac{m_{\ell_2} - v_u \epsilon_{22}^\ell}{v} \epsilon_{21}^u \tan \beta.
 \end{aligned}
 \tag{25}$$

Differently than for $B_{d,s} \rightarrow \mu^+ \mu^-$ the SM contribution cannot be calculated due to non-perturbative effects and the 2HDM II contribution is numerically irrelevant. Since we do not know the SM contribution, we require that the 2HDM III contribution alone does not generate more than the experimental upper limit on this branching ratio.

It is then easy to express the constraints on $\epsilon_{12,21}^u$ in terms of the parameters m_H and




FIG. 5: Allowed regions in the complex $\epsilon_{13,31}^d$ -plane from $B_d \rightarrow \mu^+ \mu^-$ for $\tan \beta = 30$, $\tan \beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).

$\tan \beta$:

$$|\epsilon_{12,21}^u| \leq 3.0 \times 10^{-2} \frac{(m_H/500 \text{ GeV})^2}{\tan \beta/50}. \quad (26)$$

The resulting bounds on $\epsilon_{12,21}^u$ (setting one of these elements to zero) are shown in Fig. 7.

B. Tree-level contributions to $\Delta F = 2$ processes

In the presence of non-zero elements ϵ_{ij}^q neutral Higgs mediated contributions to neutral meson mixing ($B_{d,s} - \bar{B}_{d,s}$, $K - \bar{K}$ and $D - \bar{D}$ mixing) arise (see Fig. 8). In these processes, the 2HDM contribution vanishes if the $U(1)_{\text{PQ}}$ symmetry is conserved. This has the consequence that the leading $\tan \beta$ -enhanced tree-level contribution to the $\Delta F = 2$ processes (shown in Fig. 8) is only non-vanishing if ϵ_{ij}^q and ϵ_{ji}^q are simultaneously different from zero (in the approximation $m_{A^0} = m_{H^0}$ and $\cot \beta = 0$). Making use of the effective Hamiltonian defined in Eq. (84) of the appendix we get the following contributions to $B_s - \bar{B}_s$ mixing (the expressions for $B_d - \bar{B}_d$ and $K - \bar{K}$ mixing again follow by a simple replacement of indices):

$$C_4 = -\frac{\epsilon_{23}^d \epsilon_{32}^{d*}}{m_H^2} \tan^2 \beta. \quad (27)$$




FIG. 6: Allowed regions in the complex $\epsilon_{12,21}^d$ -plane from $K_L \rightarrow \mu^+ \mu^-$ for $\tan \beta = 30$, $\tan \beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).

All other Wilson coefficients are sub-leading in $\tan \beta$. For D mixing, again only C_4 is non-zero and given by

$$C_4 = -\frac{\epsilon_{12}^u \epsilon_{21}^{u*}}{m_H^2}. \quad (28)$$

After performing the renormalization group evolution [74–78] (here we used $\mu_H = 500$ GeV at the high scale) it turns out that the dominant contribution to the hadronic matrix elements stems from O_4 . Inserting the bag factors [79, 80] and decay constants from lattice QCD (see Table. XI), we get for the 2HDM of type III contribution

$$\begin{aligned} \langle B_d^0 | C_4 O_4 | \bar{B}_d^0 \rangle &\approx 0.26 C_4 \text{ GeV}^3, \\ \langle B_s^0 | C_4 O_4 | \bar{B}_s^0 \rangle &\approx 0.37 C_4 \text{ GeV}^3, \\ \langle K^0 | C_4 O_4 | \bar{K}^0 \rangle &\approx 0.30 C_4 \text{ GeV}^3, \\ \langle D^0 | C_4 O_4 | \bar{D}^0 \rangle &\approx 0.18 C_4 \text{ GeV}^3, \end{aligned} \quad (29)$$

where we used the normalization of the meson states as defined for example in [77]. In Eq. (29) the Wilson coefficients within the matrix elements are at the corresponding meson




FIG. 7: Allowed regions in the complex $\epsilon_{12,21}^u - \tan \beta$ plane from $\bar{D}^0 \rightarrow \mu^+ \mu^-$ for $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).




FIG. 8: Feynman diagram contributing to $B_{d,s} - \bar{B}_{d,s}$, $K - \bar{K}$ and $D - \bar{D}$ mixing.

scale while C_4 on the right-handed side is given at the matching scale m_H . For computing the constraints on $\epsilon_{13}^d \epsilon_{31}^{d*}$, $\epsilon_{23}^d \epsilon_{32}^{d*}$ and $\epsilon_{12}^d \epsilon_{21}^{d*}$ we use the online update of the analysis of the UTfit collaboration [81]¹⁰. For this purpose we define

$$C_{B_q} e^{2i\varphi_{B_q}} = 1 + \frac{\langle B_q^0 | \mathcal{H}_{eff}^{NP} | \bar{B}_q^0 \rangle}{\langle B_q^0 | \mathcal{H}_{eff}^{SM} | \bar{B}_q^0 \rangle}, \quad (30)$$

for $B_d - \bar{B}_d$ and $B_s - \bar{B}_s$ mixing and

$$C_{\epsilon_K} = 1 + \frac{\text{Im} [\langle K^0 | \mathcal{H}_{eff}^{NP} | \bar{K}^0 \rangle]}{\text{Im} [\langle K^0 | \mathcal{H}_{eff}^{SM} | \bar{K}^0 \rangle]}, \quad (31)$$

$$C_{\Delta M_K} = 1 + \frac{\text{Re} [\langle K^0 | \mathcal{H}_{eff}^{NP} | \bar{K}^0 \rangle]}{\text{Re} [\langle K^0 | \mathcal{H}_{eff}^{SM} | \bar{K}^0 \rangle]},$$

¹⁰ See also the online update of the CKMfitter group for an analogous analysis [14].

for $K - \bar{K}$ mixing. Using for the matrix elements of the SM Hamiltonian¹¹ [82]

$$\begin{aligned}\langle B_d^0 | \mathcal{H}_{SM}^{\Delta F=2} | \bar{B}_d^0 \rangle &\approx (1.08 + 1.25i) \times 10^{-13} \text{ GeV}, \\ \langle B_s^0 | \mathcal{H}_{SM}^{\Delta F=2} | \bar{B}_s^0 \rangle &\approx (59 - 2.2i) \times 10^{-13} \text{ GeV}, \\ \langle K^0 | \mathcal{H}_{SM}^{\Delta F=2} | \bar{K}^0 \rangle &\approx (115 + 1.16i) \times 10^{-17} \text{ GeV},\end{aligned}\tag{32}$$

we can directly read off the bounds on C_4 and thus on $\epsilon_{12}^d \epsilon_{21}^{d*}$, $\epsilon_{13}^d \epsilon_{31}^{d*}$ and $\epsilon_{23}^d \epsilon_{32}^{d*}$:

$$-2.0 \times 10^{-10} \leq \text{Re} \left[\epsilon_{23}^d \epsilon_{32}^{d*} \left(\frac{\tan \beta / 50}{m_H / 500 \text{ GeV}} \right)^2 \right] \leq 6.0 \times 10^{-10},\tag{33}$$

$$-3.0 \times 10^{-10} \leq \text{Im} \left[\epsilon_{23}^d \epsilon_{32}^{d*} \left(\frac{\tan \beta / 50}{m_H / 500 \text{ GeV}} \right)^2 \right] \leq 7.0 \times 10^{-10},\tag{34}$$

$$-3.0 \times 10^{-11} \leq \text{Re} \left[\epsilon_{13}^d \epsilon_{31}^{d*} \left(\frac{\tan \beta / 50}{m_H / 500 \text{ GeV}} \right)^2 \right] \leq 1.5 \times 10^{-11},\tag{35}$$

$$-1.5 \times 10^{-11} \leq \text{Im} \left[\epsilon_{13}^d \epsilon_{31}^{d*} \left(\frac{\tan \beta / 50}{m_H / 500 \text{ GeV}} \right)^2 \right] \leq 2.5 \times 10^{-11},\tag{36}$$

$$-1.0 \times 10^{-12} \leq \text{Re} \left[\epsilon_{12}^d \epsilon_{21}^{d*} \left(\frac{\tan \beta / 50}{m_H / 500 \text{ GeV}} \right)^2 \right] \leq 3.0 \times 10^{-13},\tag{37}$$

$$-4.0 \times 10^{-15} \leq \text{Im} \left[\epsilon_{12}^d \epsilon_{21}^{d*} \left(\frac{\tan \beta / 50}{m_H / 500 \text{ GeV}} \right)^2 \right] \leq 2.5 \times 10^{-15}.\tag{38}$$

We see that if ϵ_{ij}^d is of the same order as ϵ_{ji}^d these bound are even more stringent than the ones from $B_{d,s} \rightarrow \mu^+ \mu^-$ and $K_L \rightarrow \mu^+ \mu^-$ computed in the last subsection.

For $D - \bar{D}$ mixing, the SM predictions is not known due to very large hadronic uncertainties. In order to constrain the NP effects we demand the absence of fine tuning, which means that the NP contribution, which are calculable short distance contributions, should not exceed the measured values. Concerning the 2HDM III contribution, there is no $\tan \beta$ enhancement and taking into account the recent analysis of UTfit collaboration [83] we arrive at the following constraints (for $m_H = 500 \text{ GeV}$):

$$|\epsilon_{12}^u \epsilon_{21}^{u*}| < 2.0 \times 10^{-8}.\tag{39}$$

Note that although these bounds look more stringent than the corresponding $\Delta F = 1$ constraints, they scale differently with $\tan \beta$ and also involve products of pairs of ϵ_{ij}^u . Therefore, contrary to the $\Delta F = 1$ case, in principle all of these limits can be evaded for one of the couplings by suppressing the other one. Fig. 9 and Fig. 10 show the allowed regions for these parameters obtained from neutral Higgs contribution to $B_{d,s} - \bar{B}_{d,s}$, $K - \bar{K}$ and $D - \bar{D}$ mixing (see the Feynman diagram in Fig. 8).

¹¹ To obtain a value consistent with the NP analysis of the UTfit collaboration, we also used their input for computing the matrix elements of the SM $\Delta F = 2$ Hamiltonian in Eq. (32).




FIG. 9: Allowed regions in the complex ϵ_{ij}^d -plane from $B_{d,s}$ - $\bar{B}_{d,s}$ mixing for $\tan\beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).




FIG. 10: Allowed regions in the complex ϵ_{12}^q - ϵ_{21}^{q*} -plane from $K-\bar{K}$ and $D-\bar{D}$ mixing for $\tan\beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).

C. Lepton-flavor-violating decays: $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, $\tau^- \rightarrow e^- \mu^+ \mu^-$ and $\mu \rightarrow e^- e^+ e^-$

In this section, we investigate the constraints that $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, $\tau^- \rightarrow e^- \mu^+ \mu^-$ and $\mu \rightarrow e^- e^+ e^-$ place on the flavor changing couplings $\epsilon_{32,23}^\ell$, $\epsilon_{31,13}^\ell$ and $\epsilon_{21,12}^\ell$, respectively.

For these decays, the experimental upper limits [84, 85] are

$$\begin{aligned}
 \mathcal{B}[\tau^- \rightarrow \mu^- \mu^+ \mu^-] &\leq 2.1 \times 10^{-8}, \\
 \mathcal{B}[\tau^- \rightarrow e^- \mu^+ \mu^-] &\leq 2.7 \times 10^{-8}, \\
 \mathcal{B}[\mu^- \rightarrow e^- e^+ e^-] &\leq 1.0 \times 10^{-12},
 \end{aligned}
 \tag{40}$$

at 90% CL. Let us consider the processes $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ and $\tau^- \rightarrow e^- \mu^+ \mu^-$ which are shown in Fig. 11. The expressions for the branching-ratio for $\tau^- \rightarrow e^- \mu^+ \mu^-$ can be written as




FIG. 11: Feynman diagrams contributing to $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ and $\tau^- \rightarrow e^- \mu^+ \mu^-$ via neutral Higgs exchange. Note that for $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ (or $\mu^- \rightarrow e^- e^+ e^-$) two distinct diagrams exist which come with a relative minus sign due to the exchange of the two fermion lines.




FIG. 12: Allowed regions for the absolute value of $\epsilon_{13,31}^l$, $\epsilon_{23,32}^l$ and $\epsilon_{12,21}^l$ for $\tan \beta = 30$ (yellow), $\tan \beta = 40$ (red) and $\tan \beta = 50$ (blue) from $\tau^- \rightarrow e^- \mu^+ \mu^-$, $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ and $\mu^- \rightarrow e^- e^+ e^-$, respectively. In each plot only one of the elements ϵ_{if}^l or ϵ_{fi}^l is assumed to be different from zero.

$$\mathcal{B} [\tau^- \rightarrow e^- \mu^+ \mu^-] = \frac{m_\tau^5}{12(8\pi)^3 \Gamma_\tau} \frac{\tan^4 \beta}{m_H^4} \left| \left(\frac{m_\mu}{v} - \varepsilon_{22}^l \right) \right|^2 \left(|\varepsilon_{31}^l|^2 + |\varepsilon_{13}^l|^2 \right) \quad (41)$$

where Γ_τ is the total decay width of the τ -lepton. The branching ratios for $\tau^- \rightarrow e^- e^+ e^-$ and $\mu^- \rightarrow e^- e^+ e^-$ can be obtained by an obvious replacement of masses, indices and total decays widths. Note that the full expression for general scalar couplings given in Eq. (116) of the appendix is different for $\tau^- \rightarrow e^- \mu^+ \mu^-$ than for $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ and only approaches a common expression in the limit of large $\tan \beta$ and large Higgs masses.

Comparing the type-III 2HDM expression with experiment we obtain the following con-

straints on ϵ_{fi}^ℓ (assuming $\epsilon_{jj}^\ell = 0$)

$$\begin{aligned}
|\epsilon_{12}^\ell|^2 + |\epsilon_{21}^\ell|^2 &\leq (2.3 \times 10^{-3})^2 \left(\frac{m_H/500 \text{ GeV}}{\tan \beta/50} \right)^4 \frac{\mathcal{B}[\mu^- \rightarrow e^- e^+ e^-]}{1.0 \times 10^{-12}}, \\
|\epsilon_{13}^\ell|^2 + |\epsilon_{31}^\ell|^2 &\leq (4.2 \times 10^{-3})^2 \left(\frac{m_H/500 \text{ GeV}}{\tan \beta/50} \right)^4 \frac{\mathcal{B}[\tau^- \rightarrow e^- \mu^+ \mu^-]}{2.7 \times 10^{-8}}, \\
|\epsilon_{23}^\ell|^2 + |\epsilon_{32}^\ell|^2 &\leq (3.7 \times 10^{-3})^2 \left(\frac{m_H/500 \text{ GeV}}{\tan \beta/50} \right)^4 \frac{\mathcal{B}[\tau^- \rightarrow \mu^- \mu^+ \mu^-]}{2.1 \times 10^{-8}}.
\end{aligned} \tag{42}$$

These constraints are also illustrated in Fig. 12 for the experimental limits given in Eq. (40).

VI. LOOP-CONTRIBUTIONS TO FCNC PROCESSES

We observed in the previous section that all elements ϵ_{ij}^d , ϵ_{ij}^ℓ (with $i \neq j$) and $\epsilon_{12,21}^u$ must be extremely small due to the constraints from tree-level neutral Higgs contributions to FCNC processes. Furthermore, the constraints on ϵ_{ij}^q and ϵ_{ji}^q get even more stringent if both of them are non-zero at the same time due to the bounds from $\Delta F = 2$ processes. Nevertheless, the elements $\epsilon_{13,23}^u$ and $\epsilon_{31,32}^u$ are still unconstrained because we have no data from neutral current top decays. In addition, also the flavor-conserving elements ϵ_{ii}^f are not constrained from neutral Higgs contributions to FCNC processes.

In this section, we study the constraints from Higgs mediated loop contributions to FCNC observables. First, in Sec. VIA we consider the $\Delta F = 2$ processes, $B_s - \bar{B}_s$, $B_d - \bar{B}_d$ and $K - \bar{K}$ mixing and then examine the constraints on $\epsilon_{13,23}^u$ and $\epsilon_{31,32}^u$ from $b \rightarrow s(d)\gamma$. Also ϵ_{22}^u (ϵ_{33}^u) can be constrained from these processes due to the relative $\tan \beta$ enhancement compared to m_c (m_t) in the quark-quark-Higgs vertices. In this analysis, we neglect the effects of the elements ϵ_{ij}^d , which means that we assume the absence of large accidental cancellations between different contributions.

Also $\Delta F = 0$ processes (electric dipole moments) place relevant constraints on the type-III 2HDM parameter space, as we will see in Sec VIF.

A. $B_s - \bar{B}_s$, $B_d - \bar{B}_d$ and $K - \bar{K}$ mixing

For the charged Higgs contributions to $\Delta F = 2$ processes we calculated the complete set of Wilson Coefficients in a general R_ξ -gauge. The result is given, together with our conventions for the Hamiltonian, in the appendix. For the QCD evolution we used the NLO running of the Wilson coefficients of Ref. [74, 75].

For computing the allowed regions in parameter space we used the same procedure as explained in the last section. The results are shown in Fig. 13, 14 and 15 and can be summarized as follows: $B_s - \bar{B}_s$ ($B_d - \bar{B}_d$) mixing gives constraints on ϵ_{23}^u (ϵ_{13}^u) which are of




FIG. 13: Allowed regions in the complex ϵ_{ij}^u -plane from B_s mixing for $\tan\beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).




FIG. 14: Allowed regions in the complex ϵ_{ij}^u -plane from B_d mixing for $\tan\beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).

the order of 10^{-1} (10^{-2}) for our typical values of $\tan\beta$ and m_H . In addition, $B_d-\bar{B}_d$ mixing also constrains ϵ_{23}^u to a similar extent as $B_s-\bar{B}_s$ mixing. The constraints on ϵ_{33}^u , ϵ_{32}^u and ϵ_{31}^u are all very weak (of order one). Also Kaon mixing gives comparable bounds on $\text{Abs}[\epsilon_{23}^u]$ and the bounds on $\text{Abs}[\epsilon_{22}^u]$ are of the order 10^{-1} .

B. Radiative B meson decays: $b \rightarrow s\gamma$ and $b \rightarrow d\gamma$

The radiative B decay $b \rightarrow s\gamma$ ($b \rightarrow d\gamma$) imposes stringent constraints on the element ϵ_{23}^u (ϵ_{13}^u) while also in this case the constraints on ϵ_{32}^u (ϵ_{31}^u) are very weak due to the light charm (up) quark involved (see left diagram in Fig. 16). For these processes both a neutral and a charged Higgs contribution occur. Since the flavor off-diagonal elements $\epsilon_{13,23}^d$ and $\epsilon_{31,32}^d$ are already stringently constrained from tree-level decays we neglect the neutral Higgs contribution here. We give the explicit results for the Higgs contributions to the Wilson




FIG. 15: Allowed regions in the complex ϵ_{ij}^u -plane from $K-\bar{K}$ mixing for $\tan\beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue). The constraints are practically independent of $\tan\beta$.




FIG. 16: Left: Feynman diagram contributing to $b \rightarrow s\gamma$ via a charm-loop containing ϵ_{32}^{u*} . The contribution is suppressed, since the small charm mass enters either from the propagator or from the charged Higgs coupling to the charm and strange quark.

Right: Feynman diagram showing a neutral Higgs box contribution to $D-\bar{D}$ mixing arising if ϵ_{31}^u and ϵ_{32}^u are simultaneously different from zero.

coefficients governing $b \rightarrow s(d)\gamma$ in the appendix.

For $B \rightarrow X_s\gamma$, we obtain the constraints on the 2HDM of type III parameters ϵ_{ij}^u by using $\mathcal{B}[B \rightarrow X_s\gamma]$ from Ref. [86] (BABAR) and Ref. [87, 88] (BELLE). Combined and extrapolated to a photon energy cut of 1.6 GeV, the HFAG value is [89]

$$\mathcal{B}[B \rightarrow X_s\gamma]_{E_\gamma > 1.6 \text{ GeV}}^{\text{exp}} = (3.43 \pm 0.21 \pm 0.07) \times 10^{-4}. \quad (43)$$

In order to estimate the possible size of NP we use the NNLO SM calculation of Ref. [48] (again for a photon energy cut of 1.6 GeV)

$$\mathcal{B}[B \rightarrow X_s\gamma]^{\text{SM}} = (3.15 \pm 0.23) \times 10^{-4}, \quad (44)$$




FIG. 17: Allowed regions for ϵ_{ij}^u from $B \rightarrow X_{s(d)}\gamma$, obtained by adding the 2σ experimental error and theoretical uncertainty linear for $\tan\beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).




FIG. 18: Allowed regions for ϵ_{ij}^u from $B \rightarrow X_{s(d)}\gamma$, obtained by adding the 2σ experimental error and theoretical uncertainty linear for $\tan\beta = 30$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).

and calculate the ratio

$$R_{\text{exp}}^{b \rightarrow s\gamma} = \frac{\mathcal{B}[B \rightarrow X_s\gamma]^{exp}}{\mathcal{B}[B \rightarrow X_s\gamma]^{SM}}. \quad (45)$$

This leads to a certain range for $R_{\text{exp}}^{b \rightarrow s\gamma}$. Now, we require that in our leading-order calculation the ratio

$$R_{\text{theory}}^{b \rightarrow s\gamma} = \frac{\mathcal{B}[B \rightarrow X_s\gamma]^{2\text{HDM}}}{\mathcal{B}[B \rightarrow X_s\gamma]^{SM}} \quad (46)$$

lies within this range. In this way, we obtain the constraints on our model parameters ϵ_{ij}^u as illustrated in Fig. 17 and Fig. 18.

The analysis for $b \rightarrow d\gamma$ is performed in an analogous way. In addition we use here the fact that most of the hadronic uncertainties cancel in the CP-averaged branching ratio for

$B \rightarrow X_d \gamma$ [90, 91]. The current experimental value of the BABAR collaboration [92, 93] for the CP averaged branching ratio reads

$$\mathcal{B} [B \rightarrow X_d \gamma] |_{E_\gamma > 1.6 \text{ GeV}}^{\text{exp}} = (1.41 \pm 0.57) \times 10^{-5}. \quad (47)$$

Here we take into account a conservative estimate of the uncertainty coming from the extrapolation in the photon energy cut [94]. For the theory prediction we use the NLL SM predictions of the CP-averaged branching ratio $\mathcal{B}(B \rightarrow X_d \gamma) |_{E_\gamma > 1.6 \text{ GeV}}$ of Ref. [95, 96], which was recently updated in Ref. [94] and reads

$$\mathcal{B} [B \rightarrow X_d \gamma] |_{E_\gamma > 1.6 \text{ GeV}}^{\text{SM}} = (1.54_{-0.31}^{+0.26}) \times 10^{-5}. \quad (48)$$

After defining the ratios $R_{\text{exp}}^{b \rightarrow d \gamma}$ and $R_{\text{theory}}^{b \rightarrow d \gamma}$ we continue as in the case of $\mathcal{B}[B \rightarrow X_s \gamma]$ in order to constrain ϵ_{13}^u .

As can be seen from Fig.17 and Fig. 18, the constraints that $B \rightarrow X_{s(d)} \gamma$ enforces on $\epsilon_{23(13)}^u$ are stronger than the ones from $B_{s(d)}$ mixing. Even ϵ_{33}^u can be restricted to a rather small range.

While in the 2HDM of type II $b \rightarrow s \gamma$ enforces a lower limit on the charged Higgs mass of 360 GeV [49] this constraint can get weakened in the 2HDM of type III: The off-diagonal element ϵ_{23}^u can lead to a destructive interference with the SM (depending on its phase) and thus reduce the 2HDM contribution. Lighter charged Higgs masses are also constrained from $b \rightarrow d \gamma$ but also this constraint can be avoided by ϵ_{13}^u .

C. Neutral Higgs box contributions to $D-\bar{D}$ mixing

Nearly all the loop-induced neutral Higgs contributions to FCNC processes can be neglected because the elements involved are already stringently constrained from tree-level processes. However, there is one exception: since the constraints on $\epsilon_{31,32}^u$ are particularly weak (because of the light charm or up quark entering the loop) this can give a sizable effect in $D-\bar{D}$ mixing via a neutral Higgs box¹² (see Fig. 16). As we will use ϵ_{31}^u and ϵ_{32}^u in Sec. VII for explaining the mentioned deviations from the SM prediction in $B \rightarrow \tau \nu$, $B \rightarrow D \tau \nu$ and $B \rightarrow D^* \tau \nu$ it is interesting to ask if all processes can be explained simultaneously without violating $D-\bar{D}$ mixing. In principle also charged Higgs contributions to $D-\bar{D}$ mixing arise but we find that they are very small compared to the H_k^0 contributions. The explicit expression for the Wilson coefficients can be found in the appendix.

Fig. 19 shows the allowed regions in the complex $\epsilon_{32}^u \epsilon_{31}^{u*}$ -plane. The constraints are again obtained by using the recent UTFit [83] analysis for the $D-\bar{D}$ system.

¹² In principle, one can also get contribution to $\bar{D}^0 \rightarrow \mu^+ \mu^-$ through H_k^0 box and penguin contributions if the elements ϵ_{32}^u and ϵ_{31}^u are simultaneously non-zero. However, we observe that they are negligible.




FIG. 19: Allowed region in the complex $\epsilon_{32}^u \epsilon_{31}^{u*}$ -plane obtained from neutral Higgs box contributions to $D-\bar{D}$ mixing for $\tan\beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).

D. Radiative lepton decays : $\mu \rightarrow e\gamma$, $\tau \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma$

The bounds on $\epsilon_{13,31}^\ell$ and $\epsilon_{23,32}^\ell$ from the radiative lepton decays $\tau \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma$ (using the experimental values given in Table II) turn out to be significantly weaker than the ones from $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ and $\tau^- \rightarrow e^- \mu^+ \mu^-$. Concerning $\mu \rightarrow e\gamma$ we expect constraints which are at least comparable to the ones from $\mu^- \rightarrow e^- e^+ e^-$ since $\mu \rightarrow e\gamma$ does not involve the small electron Yukawa coupling entering $\mu^- \rightarrow e^- e^+ e^-$. In fact, using the new MEG results [97] the constraints from $\mu \rightarrow e\gamma$ turn out to be stronger than the ones from $\mu^- \rightarrow e^- e^+ e^-$ (see Fig. 20). Note that the constraints from $\mu^- \rightarrow e^- e^+ e^-$ can be avoided if $v_u \epsilon_{11}^\ell \approx m_e$ while the leading contribution to $\mu \rightarrow e\gamma$ vanishes for $v_u \epsilon_{22}^\ell \approx m_\mu$.

Process	Experimental bounds
$\mathcal{B}[\tau \rightarrow \mu\gamma]$	$\leq 4.5 \times 10^{-8}$ [98, 99]
$\mathcal{B}[\tau \rightarrow e\gamma]$	$\leq 1.1 \times 10^{-7}$ [98]
$\mathcal{B}[\mu \rightarrow e\gamma]$	$\leq 5.7 \times 10^{-13}$ [97]

TABLE II: Experimental upper limits on the branching ratios of lepton-flavor violating decays.

In principle, for $\mu \rightarrow e\gamma$ a simplified expression for the branching ratio in the large $\tan\beta$ limit and $v \ll m_H$ could also be given. However, due to the large logarithm with a relative big prefactor (last term of Eq. (96)) this is only a good approximation for very heavy Higgses




FIG. 20: Allowed region for ϵ_{12}^ℓ (left plot) and ϵ_{21}^ℓ (right plot) from $\mu \rightarrow e\gamma$ for $\tan\beta = 30$ (yellow), $\tan\beta = 40$ (red) and $\tan\beta = 50$ (blue).

and we therefore use the full expression in our numerical analysis.

We will return to the radiative lepton decays in Sec. VIII and correlate them to the decays $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, $\tau^- \rightarrow e^- \mu^+ \mu^-$ and $\mu^- \rightarrow e^- e^+ e^-$.

E. $B_s \rightarrow \mu^+ \mu^-$

Setting $\epsilon_{ij}^q = 0$ only the loop induced charged Higgs contribution to $B_s \rightarrow \mu^+ \mu^-$ (and $B_d \rightarrow \mu^+ \mu^-$) exist. This contribution (see Eq. (24)) gets altered in the presence of non-zero elements ϵ_{ij}^ℓ , e.g. ϵ_{22}^ℓ . In the large $\tan\beta$ limit, the loop induced result in Eq. (24) is modified to

$$C_S^{bs} = C_P^{bs} = -\frac{m_b V_{tb}^* V_{ts}}{2} \frac{m_\mu - v_u \epsilon_{22}^\ell}{2M_W^2} \tan^2 \beta \frac{\log(m_H^2/m_t^2)}{m_H^2/m_t^2 - 1}. \quad (49)$$

The resulting constraints on ϵ_{22}^ℓ from $B_s \rightarrow \mu^+ \mu^-$ are shown in Fig. 21 and the ones from $B_d \rightarrow \mu^+ \mu^-$ are found to be weaker.

F. Electric dipole moments and anomalous magnetic moments

1. Charged leptons

The same diagrams which contribute to the radiative lepton decays for $\ell_i \neq \ell_f$ also affect the electric dipole moments and the anomalous magnetic moments of leptons for $\ell_i = \ell_f$. For this reason we use the same conventions as in Eq. (93) and express the EDMs of leptons in terms of the coefficients $c_{L,R}^{\ell_f \ell_i}$ of the magnetic dipole operators $O_{L,R}^{\ell_f \ell_i}$ in the following way (using that for flavor conserving transitions $c_L^{\ell_i \ell_i} = c_R^{\ell_i \ell_i^*}$)

$$d_{\ell_i} = 2m_{\ell_i} \text{Im} [c_R^{\ell_i \ell_i}]. \quad (50)$$




FIG. 21: Left: Allowed regions in the complex ϵ_{22}^ℓ -plane from $B_s \rightarrow \mu^+ \mu^-$ for $\tan \beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue). Right: Allowed regions in the ϵ_{22}^ℓ - m_H plane from $B_s \rightarrow \mu^+ \mu^-$ for $\tan \beta = 30$ (yellow), $\tan \beta = 40$ (red), $\tan \beta = 50$ (blue) and real values of ϵ_{22}^ℓ .




FIG. 22: Left: Feynman diagram contributing to EDMs (for $i = f$) or LFV decays (for $i \neq f$) involving a neutral-Higgs boson. Right: Feynman diagram contributing to EDMs (for $i = f$) or LFV decays (for $i \neq f$) involving a charged-Higgs boson.

In SM there is no contribution to the EDMs of leptons at the one-loop level. This is also true in the 2HDM of type II, because the Wilson coefficients are purely real since the phases of the PMNS matrix drop out in the charged Higgs contributions after summing over the massless neutrinos. However, in a 2HDM of type III, one can have neutral Higgs mediated contributions to EDMs. Note that there is no charged Higgs contribution to the charged lepton EDMs also in the 2HDM of type III because the Wilson coefficients are purely real in this case. Comparing the expression for the EDMs in the 2HDM of type III with the experimental upper bounds on d_e , d_μ and d_τ (see Table III), one can constrain the parameters ϵ_{ij}^ℓ (or combination of them) if they are complex.

We observe that while d_e enforces strong constraints on the products $\text{Im} [\epsilon_{13}^\ell \epsilon_{31}^\ell]$ and $\text{Im} [\epsilon_{12}^\ell \epsilon_{21}^\ell]$ (see Fig. 23), d_μ and d_τ are not capable of placing good constraints on our model

EDMs	$ d_e $	$ d_\mu $	d_τ	$ d_n $
Bounds (e cm)	10.5×10^{-28} [100]	1.9×10^{-19} [101]	$\in [-2.5, 0.8] \times 10^{-17}$ [102]	2.9×10^{-26} [103]

TABLE III: Experimental (upper) bounds on electric dipole moments.




FIG. 23: Allowed regions in the $\text{Im}[\epsilon'_{13}\epsilon'_{31}]-m_H$ and $\text{Im}[\epsilon'_{12}\epsilon'_{21}]-m_H$ planes from neutral Higgs contribution to d_e for $\tan\beta = 50$ (blue), $\tan\beta = 40$ (red) and $\tan\beta = 30$ (yellow). The constraints on $\text{Im}[\epsilon'_{11}]$ are not sizable.

parameters.

Similarly, following the conventions in Eq. (93), the anomalous magnetic moments (AMMs) can be written in terms of $c_R^{\ell_i \ell_i}$ as ($e > 0$)

$$a_{\ell_i} = -\frac{4m_{\ell_i}^2}{e} \text{Re}[c_R^{\ell_i \ell_i}]. \quad (51)$$

The discrepancy between experiment and the SM prediction for the muon magnetic moment $a_\mu = (g - 2)/2$ is [104–108]

$$\Delta a_\mu = a_\mu^{exp} - a_\mu^{SM} \approx (3 \pm 1) \times 10^{-9}. \quad (52)$$

In the 2HDM of type II, the sum of the neutral and charged Higgs mediated diagrams gives the following contribution to a_μ (for $\tan\beta = 50$ and $m_H = 500$ GeV):

$$a_\mu^{2\text{HDM II}} \approx 2.7 \times 10^{-13}, \quad (53)$$

which is interfering constructively with the SM. Anyway, it can be seen that the effect is orders of magnitude smaller than the actual sensitivity and it even gets smaller for higher Higgs masses.




FIG. 24: Left: Allowed region in the $\epsilon_{23}^{\ell}-\epsilon_{32}^{\ell}$ plane from Δa_{μ} for real values of ϵ_{23}^{ℓ} , ϵ_{32}^{ℓ} and $\tan \beta = 50$, $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue). Right: Allowed region in $\epsilon_{22}^{\ell}-m_H$ plane from Δa_{μ} for real values of ϵ_{22}^{ℓ} and $\tan \beta = 50$ (blue), $\tan \beta = 40$ (red) and $\tan \beta = 30$ (yellow).

Concerning the 2HDM of type III the discrepancy between experiment and the SM prediction given in Eq. (52) could be explained but only with severe fine-tuning. One would need to allow for very large values of ϵ_{22}^{ℓ} which would not only violate 't Hooft's naturalness criterion but also enhance $B_s \rightarrow \mu^+ \mu^-$ by orders of magnitude above the experimental limit. If one would try to explain the anomaly using ϵ_{23}^{ℓ} and ϵ_{32}^{ℓ} (ϵ_{12}^{ℓ} and ϵ_{21}^{ℓ}) one would violate the bounds from $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ ($\mu^- \rightarrow e^- e^+ e^-$ or $\mu \rightarrow e \gamma$) as illustrated in Fig. 24.

In conclusion, neither a type-II nor a type-III 2HDM can give a sizable effect in a_{μ} and both models are not capable of explaining the deviation from the SM.

2. Electric dipole moment of the neutron

The neutron electric dipole moment d_n can also provide constraints on the parameters ϵ_{ij}^a . In the SM, there is no contribution to d_n at the 1-loop level since the coefficients are real. This is also true in the type-II 2HDM.

Using the theory estimate of Ref. [109], which is based on the QCD sum-rules calculations of Refs. [110–113], the neutron EDM can be written as

$$d_n = (1 \pm 0.5) [1.4(d_d - 0.25d_u) + 1.1e(d_d^g + 0.5d_u^g)] , \quad (54)$$

where d_u (d_d) is the EDM of the up (down) quark and $d_{u(d)}^g$ define the corresponding chromoelectric dipole moments which stem from the chromomagnetic dipole operator

$$O_{R(L)}^{q_f q_i} = m_{q_i} \bar{q}_f \sigma^{\mu\nu} T^a P_{R(L)} q_i G_{\mu\nu}^a . \quad (55)$$




FIG. 25: Allowed regions in the $\text{Im}[\epsilon_{11,22}^u]-m_H$ planes from the electric dipole moment of the neutron for $\tan\beta = 50$ (blue), $\tan\beta = 40$ (red) and $\tan\beta = 30$ (yellow). We observe that d_n can not provide good constraints on the real parts of $\epsilon_{11,22}^u$.




FIG. 26: Allowed regions in the complex $\epsilon_{21,31}^u$ -planes from d_n for $\tan\beta = 50$ and $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue). We see that the absolute value of ϵ_{31}^u can only be large if it is aligned to V_{ub} , i.e. $\text{Arg}[V_{ub}] = \text{Arg}[\epsilon_{31}^u] \pm \pi$ which is very important when we consider later $B \rightarrow \tau\nu$.

Similar to EDMs, the (chromo) electric dipole moments of quarks are given as

$$d_{q_i}^{(g)} = 2m_{q_i} \text{Im} \left[c_{R,(g)}^{q_i q_i} \right]. \quad (56)$$

Using the upper limit on d_n (see Table III) we can constrain some of ϵ_{ij}^u (for $\epsilon_{ij}^d = 0$) as shown in Fig. 25 and 26. These constraints are obtained for the conservative case of assuming a prefactor of 0.5 in Eq. (54). The explicit expressions for $c_{R,(g)}^{q_i q_i}$ stemming from neutral and charged Higgs contributions to $d_{q_i}^{(g)}$ are relegated to the appendix. Note that for the neutron EDM we did not include QCD corrections.

VII. TREE-LEVEL CHARGED CURRENT PROCESSES

In this section we study the constraints from processes which are mediated in the SM by a tree-level W exchange and which receive additional contributions from charged Higgs exchange in 2HDMs. We study purely leptonic meson decays, semileptonic meson decays and tau lepton decays. Concerning B meson decays we consider $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ which are, as outlined in the introduction, very interesting in the light of the observed deviation from the SM. We consider in addition $D_{(s)} \rightarrow \tau\nu$, $D_{(s)} \rightarrow \mu\nu$, $K(\pi) \rightarrow e\nu$, $K(\pi) \rightarrow \mu\nu$ and $\tau \rightarrow K(\pi)\nu$ and look for violation of lepton flavor universality via $K(\pi) \rightarrow e\nu/K(\pi) \rightarrow \mu\nu$ and $\tau \rightarrow K(\pi)\nu/K(\pi) \rightarrow \mu\nu$. Even though no deviations from the SM have been observed in these channels, they put relevant constraints on the parameter space of the type-III 2HDM.

For purely leptonic decays of a pseudoscalar meson M (and also tau decays to mesons) to a lepton ℓ_j and a neutrino ν (which is not detected) the SM prediction is given by

$$\mathcal{B}_{SM} [M \rightarrow \ell_j\nu] = \frac{m_M}{8\pi} G_F^2 m_{\ell_j}^2 \tau_M f_M^2 |V_{u_f d_i}|^2 \left(1 - \frac{m_{\ell_j}^2}{m_M^2}\right)^2 \left(1 + \delta_{EM}^{M\ell_j}\right), \quad (57)$$

where $\delta_{EM}^{M\ell_j}$ stands for channel dependent electromagnetic corrections (see Table IV), m_M is the mass of the meson involved and m_{u_f} (m_{d_i}) refers to the mass of its constituent up (down) type quark. The expression for $\tau \rightarrow M\nu$ differs by the exchange of the meson masses (life time) with the tau masses (life time) and by a factor of 1/2 stemming from spin averaging.

NP via scalar operators can be included very easily:

$$\mathcal{B}_{NP} = \mathcal{B}_{SM} \left| 1 + \frac{m_M^2}{(m_{u_f} + m_{d_i}) m_{\ell_j}} \frac{C_R^{u_f d_i, \ell_j} - C_L^{u_f d_i, \ell_j}}{C_{SM}^{u_f d_i, \ell_j}} \right|^2 \quad (58)$$

with

$$C_{SM}^{u_f d_i, \ell_j} = 4G_F V_{u_f d_i} / \sqrt{2}. \quad (59)$$

All quantities in Eq. (58) are understood to be at the meson scale m_M . Like for $B_s \rightarrow \mu^+\mu^-$, the SM Wilson coefficient is renormalization scale independent and the scalar Wilson coefficients evolve in the same way as the quark masses.

In the 2HDM III the Wilson coefficients $C_L^{u_f d_i, \ell_j}$ and $C_R^{u_f d_i, \ell_j}$ are given by (neglecting terms which are not $\tan\beta$ enhanced)

$$\begin{aligned} C_R^{u_f d_i, \ell_j} &= -\frac{\tan^2\beta}{m_{H^\pm}^2} \left(V_{fi} \frac{m_{d_i}}{v} - \sum_{j=1}^3 V_{fj} \epsilon_{ji}^d \right) \left(\frac{m_{\ell_j}}{v} - \sum_{k=1}^3 \epsilon_{kj}^{\ell^*} \right), \\ C_L^{u_f d_i, \ell_j} &= \frac{\tan\beta}{m_{H^\pm}^2} \sum_{j=1}^3 V_{ji} \epsilon_{jf}^{\star u} \left(\frac{m_{\ell_j}}{v} - \sum_{k=1}^3 \epsilon_{kj}^{\ell^*} \right). \end{aligned} \quad (60)$$

Ratio	Experimental value	SM prediction	$\delta_{EM}^{M\ell_j}$
$\mathcal{B}[K \rightarrow e\nu]/\mathcal{B}[K \rightarrow \mu\nu]$	$(2.488 \pm 0.013) \times 10^{-5}$	$(2.472 \pm 0.001) \times 10^{-5}$	-0.0378 ± 0.0004 [61]
$\mathcal{B}[K \rightarrow \mu\nu]/\mathcal{B}[\pi \rightarrow \mu\nu]$	$(63.55 \pm 0.11) \times 10^{-2}$	$(63.48 \pm 1.37) \times 10^{-2}$	-0.0070 ± 0.0018 [114]
$\mathcal{B}[K \rightarrow e\nu]/\mathcal{B}[\pi \rightarrow e\nu]$	$(1.285 \pm 0.008) \times 10^{-1}$	$(1.270 \pm 0.027) \times 10^{-1}$	-0.0070 ± 0.0018 [114]
$\mathcal{B}[\pi \rightarrow e\nu]/\mathcal{B}[\pi \rightarrow \mu\nu]$	$(1.230 \pm 0.004) \times 10^{-4}$	1.234×10^{-4}	-3.85% [115]
$\mathcal{B}[\tau \rightarrow K\nu]/\mathcal{B}[\tau \rightarrow \pi\nu]$	$(6.46 \pm 0.10) \times 10^{-2}$	$(6.56 \pm 0.16) \times 10^{-2}$	0.0003 ± 0.0044 [116]
$\mathcal{B}[\tau \rightarrow \pi\nu]/\mathcal{B}[\pi \rightarrow \mu\nu]$	$(10.83 \pm 0.06) \times 10^{-2}$	10.87×10^{-2}	$+1.2\%$ [115]
$\mathcal{B}[\tau \rightarrow K\nu]/\mathcal{B}[K \rightarrow \mu\nu]$	$(1.102 \pm 0.016) \times 10^{-2}$	1.11×10^{-2}	$+2.0\%$ [115]

TABLE IV: Experimental values, SM predictions and electromagnetic corrections (in the SM) for the ratios of charged current processes. The experimental values are obtained by adding the errors of the individual branching ratios given in Ref. [72] in quadrature. The SM predictions include the uncertainties from $\delta_{EM}^{M\ell_j}$ and (if involved) as well as the uncertainties due to CKM factors and decay constants. As always, we add the theory error linear to the experimental ones.

Note that $C_L^{u_f d_i, \ell_j}$ is only proportional to one power of $\tan\beta$ while $C_R^{u_f d_i, \ell_j}$ is proportional to $\tan^2\beta$. The Hamiltonian governing $M \rightarrow \ell_j\nu$ ($\tau \rightarrow M\nu$) and the Wilson coefficients for general scalar interactions are given in the appendix. It is important to keep in mind that, since we are dealing with lepton flavour-violating terms, we must sum over the neutrinos in the final state because the neutrino is not detected. Note that we did not include the PMNS matrix in both $C_{SM}^{u_f d_i, \ell_j}$ and $C_{L,R}^{u_f d_i, \ell_j}$ for simplifying the expressions, since it cancels in the final expression after summing over the neutrinos.

For semileptonic meson decays $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$, which have a three-body final state, both the SM prediction and the inclusion of NP is more complicated, as will be discussed in subsection VII A 1.

A. Tauonic charged B meson decays: $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$

As discussed in the introduction the BABAR collaboration performed an analysis of the semileptonic B decays $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ using the full available data set [8, 9].

They find for the ratios

$$\mathcal{R}(D^{(*)}) = \mathcal{B}(B \rightarrow D^{(*)}\tau\nu)/\mathcal{B}(B \rightarrow D^{(*)}\ell\nu), \quad (61)$$

(with $\ell = e, \mu$) the following results:

$$\mathcal{R}(D) = 0.440 \pm 0.058 \pm 0.042, \quad (62)$$

$$\mathcal{R}(D^*) = 0.332 \pm 0.024 \pm 0.018. \quad (63)$$

Here the first error is statistical and the second one is systematic. Comparing these measurements to the SM predictions

$$\mathcal{R}_{\text{SM}}(D) = 0.297 \pm 0.017, \quad (64)$$

$$\mathcal{R}_{\text{SM}}(D^*) = 0.252 \pm 0.003, \quad (65)$$

we see that there is a discrepancy of 2.0σ for $\mathcal{R}(D)$ and 2.7σ for $\mathcal{R}(D^*)$. For the theory predictions we used the updated results of [8], which rely on the calculations of Refs. [55, 117] based on the results of Refs. [118–122]. The measurements of both ratios $\mathcal{R}(D)$ and $\mathcal{R}(D^*)$ exceed the SM prediction, and combining them gives a 3.4σ deviation from the SM [8, 9] expectation.

This evidence for the violation of lepton flavour universality in $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ is further supported by the measurement of $B \rightarrow \tau\nu$ by BABAR [10, 11] and BELLE [12]. Until recently, all measurements of $B \rightarrow \tau\nu$ (the hadronic tag and the leptonic tag both from BABAR and BELLE) were significantly above the SM prediction. However, the latest BELLE result for the hadronic tag [13] of $\mathcal{B}[B \rightarrow \tau\nu] = (0.72_{-0.25}^{+0.27} \pm 0.11) \times 10^{-4}$ is in agreement with the SM prediction [14]:

$$\mathcal{B}_{\text{SM}}[B \rightarrow \tau\nu] = (0.796_{-0.087}^{+0.088}) \times 10^{-4}. \quad (66)$$

Averaging all measurements, one obtains the branching ratio

$$\mathcal{B}_{\text{exp}}[B \rightarrow \tau\nu] = (1.15 \pm 0.23) \times 10^{-4}. \quad (67)$$

which now disagrees with the SM prediction by 1.6σ using V_{ub} from the global fit [14].

Combining $\mathcal{R}(D)$, $\mathcal{R}(D^*)$ and $B \rightarrow \tau\nu$, we have evidence for violation of lepton flavor universality. Assuming that these deviations from the SM are not statistical fluctuations or underestimated theoretical or systematic uncertainties, it is interesting to ask which model of new physics can explain the measured values [16, 123–132].

1. $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$

Let us first consider the semileptonic decays $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$. Here the Wilson coefficients $C_R^{qb,\tau}$ and $C_L^{qb,\tau}$ affect $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ in the following way [54, 55, 133]:




FIG. 27: Feynman diagram showing a charged Higgs contributing to $B \rightarrow \tau\nu$ and $B \rightarrow D^{(*)}\tau\nu$ involving the flavour changing parameters ϵ_{31}^u and ϵ_{32}^u which affect $B \rightarrow \tau\nu$ and $B \rightarrow D^{(*)}\tau\nu$, respectively.

$$\mathcal{R}(D) = \mathcal{R}_{SM}(D) \left(1 + 1.5\Re \left[\frac{C_R^{cb,\tau} + C_L^{cb,\tau}}{C_{SM}^{cb,\tau}} \right] + 1.0 \left| \frac{C_R^{cb,\tau} + C_L^{cb,\tau}}{C_{SM}^{cb,\tau}} \right|^2 \right), \quad (68)$$

$$\mathcal{R}(D^*) = \mathcal{R}_{SM}(D^*) \left(1 + 0.12\Re \left[\frac{C_R^{cb,\tau} - C_L^{cb,\tau}}{C_{SM}^{cb,\tau}} \right] + 0.05 \left| \frac{C_R^{cb,\tau} - C_L^{cb,\tau}}{C_{SM}^{cb,\tau}} \right|^2 \right). \quad (69)$$

For our analysis we add the experimental errors in quadrature and the theoretical uncertainty linear on top of this. There are also efficiency corrections to $\mathcal{R}(D)$ due to the BABAR detector [8] which are important in the case of large contributions from the scalar Wilson coefficients $C_{R,L}^{cb,\tau}$ (i.e. if one wants to explain $\mathcal{R}(D)$ with destructive interference with the SM contribution). As shown in Ref. [123], these corrections can be effectively taken into account by multiplying the quadratic term in $C_{R,L}^{cb,\tau}$ of Eq. (68) by an approximate factor of 1.5 (not included in Eq. (68)).

Since ϵ_{33}^d contributes to $C_R^{cb,\tau}$ (the same Wilson coefficient generated in the type-II 2HDM) it cannot simultaneously explain $\mathcal{R}(D)$ and $\mathcal{R}(D^*)$. Therefore, we are left with ϵ_{32}^u , which contributes to $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$. In the left frame of Fig. 28 we see the allowed region in the complex ϵ_{32}^u -plane, which gives the correct values for $\mathcal{R}(D)$ and $\mathcal{R}(D^*)$ within the 1σ uncertainties for $\tan\beta = 50$ and $m_H = 500$ GeV, and the middle and the right frames correspond to the allowed regions on ϵ_{31}^u from $B \rightarrow \tau\nu$.

2. $B \rightarrow \tau\nu$

In principle, $B \rightarrow \tau\nu$ can be explained either by using ϵ_{33}^d (as in 2HDMs with MFV) or by ϵ_{31}^u (or by a combination of both of them). However, ϵ_{33}^d alone cannot explain the deviation from the SM without fine tuning, while ϵ_{31}^u is capable of doing this [16].

$B \rightarrow \tau\nu$ can also be used to constrain ϵ_{13}^ℓ , ϵ_{23}^ℓ and ϵ_{33}^ℓ as illustrated in Fig. 29. In order to obtain these constraints, we assumed that all other relevant elements (ϵ_{33}^d and ϵ_{31}^u) are zero.




FIG. 28: Left: Allowed regions in the complex ϵ_{32}^u -plane from $\mathcal{R}(D)$ (blue) and $\mathcal{R}(D^*)$ (yellow) for $\tan \beta = 50$ and $m_H = 500$ GeV. Middle: Allowed regions in the complex ϵ_{31}^u -plane combining the constraints from $B \rightarrow \tau\nu$ (1σ (yellow) and 2σ (blue)) and neutron EDM (green) for $\tan \beta = 50$ and $m_H = 500$ GeV. Right: Allowed regions in the m_H - ϵ_{31}^u plane from $B \rightarrow \tau\nu$ for real values of ϵ_{31}^u and $\tan \beta = 50$ (green), $\tan \beta = 30$ (orange).




FIG. 29: Left: Allowed regions in the m_H - ϵ_{33}^l plane from $B \rightarrow \tau\nu$ for real values of ϵ_{33}^l and $\tan \beta = 30$ (yellow), $\tan \beta = 40$ (red), $\tan \beta = 50$ (blue). Right: Allowed regions in the complex ϵ_{13}^l , ϵ_{23}^l and ϵ_{33}^l -planes from $B \rightarrow \tau\nu$ for $m_H = 700$ GeV (yellow), $m_H = 500$ GeV (red) and $m_H = 300$ GeV (blue).

B. $D_{(s)} \rightarrow \tau\nu$ and $D_{(s)} \rightarrow \mu\nu$

Previously, there were some indications for NP in $D_s \rightarrow \tau\nu$ [134–136]. However, using the new experimental values for $\mathcal{B}[D_s \rightarrow \tau\nu]$ (see Table V) and the improved lattice determination for the decay constant f_{D_s} [137, 138] we find agreement between the SM predictions and experiment. Nevertheless, it is interesting to consider the constraints on the 2HDM of type III parameter space. Charged Higgs contributions to $D_{(s)} \rightarrow \tau\nu$ and $D_{(s)} \rightarrow \mu\nu$ have




FIG. 30: Left: Allowed region in the m_H - ϵ_{22}^u -plane (for real values of ϵ_{22}^u) obtained by combining the constraints from $D \rightarrow \mu\nu$ and $D_s \rightarrow \mu\nu$ for $\tan \beta = 30$ (yellow), $\tan \beta = 40$ (red) and $\tan \beta = 50$ (blue). While the upper bound on ϵ_{22}^u comes from $D_s \rightarrow \mu\nu$, $D \rightarrow \mu\nu$ is more constraining for negative values of ϵ_{22}^u . The bound on the imaginary part of ϵ_{22}^u are very weak. The constraints from $D_s \rightarrow \tau\nu$ turn out to be comparable (but a bit weaker) while the ones from $D \rightarrow \tau\nu$ are weak.

Process	Experimental value (bound)	SM prediction
$\mathcal{B}[D_s \rightarrow \tau\nu]$	$(5.43 \pm 0.31) \times 10^{-2}$	$(5.36_{-0.50}^{+0.54}) \times 10^{-2}$
$\mathcal{B}[D_s \rightarrow \mu\nu]$	$(5.90 \pm 0.33) \times 10^{-3}$	$(5.50_{-0.52}^{+0.55}) \times 10^{-3}$
$\mathcal{B}[D \rightarrow \tau\nu]$	$\leq 1.2 \times 10^{-3}$	$(1.10 \pm 0.06) \times 10^{-3}$
$\mathcal{B}[D \rightarrow \mu\nu]$	$(3.82 \pm 0.33) \times 10^{-4}$	$(4.15_{-0.21}^{+0.22}) \times 10^{-4}$

TABLE V: Experimental values (upper bounds) and SM predictions for $D_{(s)} \rightarrow \tau\nu$ and $D_{(s)} \rightarrow \mu\nu$ processes. The SM prediction for $D_s \rightarrow \mu\nu$ mode takes into account the EM correction effects of +1.0% [134, 135, 140].

been investigated in Ref. [58–60, 139].

The most important constraints on the 2HDM of type III parameter space are the ones on ϵ_{22}^u (shown in Fig. 30). $D_{(s)} \rightarrow \tau\nu$ and $D_{(s)} \rightarrow \mu\nu$ constrains $\text{Re}[\epsilon_{22}^u]$ while the constraints on $\text{Im}[\epsilon_{22}^u]$ are very weak. In principle, also the ratio $D_{(s)} \rightarrow \tau\nu / D_{(s)} \rightarrow \mu\nu$ could be used for constraining deviations from lepton flavor universality, but the constraints from $K(\pi) \rightarrow e\nu / K(\pi) \rightarrow \mu\nu$ and $\tau \rightarrow K(\pi)\nu / K(\pi) \rightarrow \mu\nu$ turn out to be stronger.




FIG. 31: Allowed regions in the m_H - $\epsilon_{i1, i2}^\ell$ -plane from $K \rightarrow \mu\nu/\pi \rightarrow \mu\nu$ and $K \rightarrow e\nu/\pi \rightarrow e\nu$ for real values of $\epsilon_{i1, i2}^\ell$ and $\tan\beta = 30$ (yellow), $\tan\beta = 40$ (red) and $\tan\beta = 50$ (blue). The constraints are weaker than the ones from $K(\pi) \rightarrow e\nu/K(\pi) \rightarrow \mu\nu$ and $\tau \rightarrow K(\pi)\nu/K(\pi) \rightarrow \mu\nu$ but cannot be avoided assuming the MFV limit ($\frac{m_{\ell_i}}{m_{\ell_j}} = \frac{\epsilon_{ij}^\ell}{\epsilon_{jj}^\ell}$).

C. $K \rightarrow \mu\nu/\pi \rightarrow \mu\nu$ and $K \rightarrow e\nu/\pi \rightarrow e\nu$

The ratio $R_{K_{e2}, \pi_{e2}} = \mathcal{B}[K \rightarrow \ell\nu]/\mathcal{B}[\pi \rightarrow \ell\nu]$ ($\ell = e, \mu$) is useful for constraining ϵ_{22}^d , ϵ_{i1}^ℓ and ϵ_{i2}^ℓ because the ratio of the decay constants f_K/f_π is known more precisely than the single decay constants [61].

For obtaining the experimental values we add the errors of the individual branching ratios in quadrature and the SM values take into account the electromagnetic correction. The corresponding values are given in Table. IV. The errors are due to the combined uncertainties in f_K/f_π , the CKM elements and the EM corrections. We obtained the value for V_{us} from $K \rightarrow \pi\ell\nu$ (which is much less sensitive to charged Higgs contributions than $K \rightarrow \mu\nu/\pi \rightarrow \mu\nu$) and V_{ud} by exploiting CKM unitarity.

Fig. 32 illustrates the allowed regions for ϵ_{22}^d by combining the constraints from $K \rightarrow \mu\nu/\pi \rightarrow \mu\nu$ and $K \rightarrow e\nu/\pi \rightarrow e\nu$. Like in $D_{(s)} \rightarrow \tau\nu$ and $D_{(s)} \rightarrow \mu\nu$ the constraints are on the real part of ϵ_{22}^d while the constraints on the imaginary part are very weak. Concerning ϵ_{i1}^ℓ and ϵ_{i2}^ℓ the constraints from $K(\pi) \rightarrow e\nu/K(\pi) \rightarrow \mu\nu$ will turn out to be more stringent but the latter ones can be avoided in the limit $\frac{m_{\ell_i}}{m_{\ell_j}} = \frac{\epsilon_{ij}^\ell}{\epsilon_{jj}^\ell}$ (see Fig. 31 and Fig. 33).

D. $\tau \rightarrow K\nu/\tau \rightarrow \pi\nu$

The τ is the only lepton which is heavy enough to decay into hadrons. The ratio $\mathcal{B}[\tau \rightarrow K\nu]/\mathcal{B}[\tau \rightarrow \pi\nu]$ can be considered for putting constraints on ϵ_{21}^u , ϵ_{12}^d and ϵ_{i3}^ℓ .




FIG. 32: Left: Allowed regions in the m_H - ϵ_{i3}^ℓ -plane from $\tau \rightarrow K\nu/\tau \rightarrow \pi\nu$. Right: Allowed regions in the m_H - ϵ_{22}^d -plane obtained by combining the constraints from $K \rightarrow \mu\nu/\pi \rightarrow \mu\nu$ and $K \rightarrow e\nu/\pi \rightarrow e\nu$ for real values of ϵ_{22}^d . In both plots $\tan\beta = 30$ (yellow), $\tan\beta = 40$ (red) and $\tan\beta = 50$ (blue).

The experimental and theoretical values for this ratio are given in Table. IV. We observe that the constraints from $\bar{D}^0 \rightarrow \mu^+\mu^-$ and $D - \bar{D}$ mixing on ϵ_{21}^u and $K_L \rightarrow \mu^+\mu^-$ on ϵ_{12}^d are too stringent so that no sizable effects stemming from these elements are possible. Also concerning ϵ_{i3}^ℓ , as we will see in the following sections, the constraints from $\tau \rightarrow \pi\nu/\pi \rightarrow \mu\nu$ will be stronger but again the latter ones can be avoided in the MFV limit $\frac{m_{\ell_i}}{m_{\ell_j}} = \frac{\epsilon_{ij}^\ell}{\epsilon_{jj}^\ell}$ (see Fig. 32).

E. Tests for lepton flavour universality: $K(\pi) \rightarrow e\nu/K(\pi) \rightarrow \mu\nu$ and

$\tau \rightarrow K(\pi)\nu/K(\pi) \rightarrow \mu\nu$

$K_{\ell 2}(K \rightarrow \ell\nu)$ decays ($\ell = e, \mu$) are helicity suppressed in the SM and suffers from large theoretical uncertainties due to the decay constants. However, considering the ratio $R_{K_{\ell 2}} = \mathcal{B}[K \rightarrow e\nu]/\mathcal{B}[K \rightarrow \mu\nu]$ the dependence on decay constants drops out.

In the 2HDM of type II the charged Higgs contributions to $K(\pi) \rightarrow e\nu/K(\pi) \rightarrow \mu\nu$ and $\tau \rightarrow K(\pi)\nu/K(\pi) \rightarrow \mu\nu$ drop out. This is also true in the 2HDM of type III (for $\epsilon_{ij}^\ell = 0$ with $i \neq j$) as long as the MFV-like relation $\epsilon_{22}^\ell/m_\mu = \epsilon_{11}^\ell/m_e$ is not violated.

1. $K \rightarrow e\nu/K \rightarrow \mu\nu$ and $\pi \rightarrow e\nu/\pi \rightarrow \mu\nu$

$K \rightarrow e\nu/K \rightarrow \mu\nu$ is a very precise test of lepton flavor universality [141] (see table. IV). Including NP entering via scalar operators modifies this ratio according to Eq. (58).




FIG. 33: Allowed regions in the m_H - ϵ_{ij}^l -plane obtained by combining the constraints from $K \rightarrow e\nu/K \rightarrow \mu\nu$ and $\pi \rightarrow e\nu/\pi \rightarrow \mu\nu$ for real values of ϵ_{ij}^l and $\tan\beta = 30$ (yellow), $\tan\beta = 40$ (red) and $\tan\beta = 50$ (blue). The constraints on ϵ_{i1}^l (affecting the electron coupling) are more stringent than the constraints on ϵ_{i2}^l (which affect the muon coupling).

We find strong constraints on ϵ_{i2}^l (which affect the coupling to the muon) and the constraints on ϵ_{i1}^l (where the coupling of the electron is involved) are even more stringent. Like for $D_{(s)} \rightarrow \tau\nu$ and $D_{(s)} \rightarrow \mu\nu$ the constraints are much better for the real part of ϵ_{ij}^l than the imaginary part. Note that these constraints are obtained assuming that only one element ϵ_{ij}^l is non-zero. In the case $\epsilon_{22}^l/m_\mu = \epsilon_{11}^l/m_e$ where lepton flavor universality is restored no constraints can be obtained.

Alternatively, the ratio $\pi \rightarrow e\nu/\pi \rightarrow \mu\nu$ can test lepton flavor universality. We find that the constraints from $\pi \rightarrow e\nu/\pi \rightarrow \mu\nu$ are comparable with the ones from $K \rightarrow e\nu/K \rightarrow \mu\nu$. Our results are illustrated in Fig. 33.

2. $\tau \rightarrow K\nu/K \rightarrow \mu\nu$ and $\tau \rightarrow \pi\nu/\pi \rightarrow \mu\nu$

The ratios $\tau \rightarrow K\nu/K \rightarrow \mu\nu$ and $\tau \rightarrow \pi\nu/\pi \rightarrow \mu\nu$ are very similar to $K(\pi) \rightarrow e\nu/K(\pi) \rightarrow \mu\nu$: all dependences on decay constants and CKM elements drop out and they are only sensitive to NP which violates lepton-flavour universality. The corresponding experimental and the theoretical values for these ratios are given in Table. IV.

We find that the constraints on ϵ_{i3}^l from $\tau \rightarrow \pi\nu/\pi \rightarrow \mu\nu$ are stronger than the ones from $\tau \rightarrow K\nu/K \rightarrow \mu\nu$ and they are shown in Fig. 34.




FIG. 34: Allowed regions in the $m_{H^-}-\epsilon_{i3}^{\ell}$ -plane from $\tau \rightarrow \pi\nu/\pi \rightarrow \mu\nu$ for real values of ϵ_{i3}^{ℓ} and $\tan\beta = 30$ (yellow), $\tan\beta = 40$ (red), $\tan\beta = 50$ (blue). The bounds on the imaginary parts are very weak.

VIII. UPPER LIMITS AND CORRELATION FOR LFV PROCESSES

In Sec. V we found that the neutral current lepton decays $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ and $\tau^- \rightarrow e^- \mu^+ \mu^-$ give more stringent bounds on the elements $\epsilon_{32,23}^{\ell}$ and $\epsilon_{31,13}^{\ell}$ than the radiative decays $\tau \rightarrow \mu\gamma$ and $\tau \rightarrow e\gamma$. Also the LFV neutral meson decays $B_{s,d} \rightarrow \tau\mu$, $B_{s,d} \rightarrow \tau e$, $B_{s,d} \rightarrow \mu e$ cannot be arbitrarily large in the type-III 2HDM due to the constraints from $B_{s,d} \rightarrow \mu^+ \mu^-$ and $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, $\tau^- \rightarrow e^- \mu^+ \mu^-$, $\mu^- \rightarrow e^- e^+ e^-$ (assuming again the absence of large cancellations)¹³.

Therefore, in this section we study the upper limits on $B_{s,d} \rightarrow \tau\mu$, $B_{s,d} \rightarrow \tau e$, $B_{s,d} \rightarrow \mu e$ and the correlation among $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, $\tau^- \rightarrow e^- \mu^+ \mu^-$, $\mu^- \rightarrow e^- e^+ e^-$ and $\tau \rightarrow \mu\gamma$, $\tau \rightarrow e\gamma$, $\mu \rightarrow e\gamma$ in the type-III 2HDM.

A. Neutral meson decays: $B_{s,d} \rightarrow \tau\mu$, $B_{s,d} \rightarrow \tau e$ and $B_{s,d} \rightarrow \mu e$

In the SM (with massless neutrinos) the branching ratios for these decays vanish. Also in the 2HDM of type II these decays are not possible (even beyond tree-level). In the type-III 2HDM, these decay modes are generated in the presence of flavor-violating terms ϵ_{ij}^{ℓ} and there exists even a tree-level neutral Higgs contribution to $B_s \rightarrow \ell_i^+ \ell_j^-$ ($B_d \rightarrow \ell_i^+ \ell_j^-$) if also $\epsilon_{23,32}^d \neq 0$ ($\epsilon_{13,31}^d \neq 0$).

¹³ see e.g. Ref. [142–144] for an analysis of NP in $B_{s,d} \rightarrow \tau\mu$.

Observables	$\mathcal{B}(B_s \rightarrow \mu e)$	$\mathcal{B}(B_d \rightarrow \mu e)$	$\mathcal{B}(B_d \rightarrow \tau \mu)$	$\mathcal{B}(B_d \rightarrow \tau e)$
Upper bounds	2.0×10^{-7} [145]	6.4×10^{-8} [145]	2.2×10^{-5} [146]	2.8×10^{-5} [146]

TABLE VI: Upper limits (90 % CL) on the branching ratios of the lepton flavor-violating B meson decays.

In the large $\tan \beta$ limit, $v \ll m_H$ and neglecting the smaller lepton mass, the corresponding expressions for these branching ratios take the simple form

$$\mathcal{B}[B_q \rightarrow \ell_i^+ \ell_j^-] \approx N_{ij}^q \left(\frac{\tan \beta / 50}{m_H / 500 \text{ GeV}} \right)^4 2 \left[|\epsilon_{ji}^\ell|^2 |\epsilon_{q3}^d|^2 + |\epsilon_{ij}^\ell|^2 |\epsilon_{3q}^d|^2 \right], \quad (70)$$

with $q = d, s$, $N_{ji}^q = N_{ij}^q$ and

$$\begin{aligned} N_{21}^s &\approx 2.1 \times 10^7 \frac{f_{B_s}}{0.229 \text{ GeV}}, \\ N_{21}^d &\approx 1.6 \times 10^7 \frac{f_{B_d}}{0.196 \text{ GeV}}, \\ N_{31,32}^s &\approx 1.7 \times 10^7 \frac{f_{B_s}}{0.229 \text{ GeV}}, \\ N_{31,32}^d &\approx 1.2 \times 10^7 \frac{f_{B_d}}{0.196 \text{ GeV}}. \end{aligned} \quad (71)$$

Note that the expressions for the branching ratios are not symmetric in ϵ_{ij}^ℓ and ϵ_{ji}^ℓ . Since experimentally both $B_q \rightarrow \ell_i^+ \ell_j^-$ and $B_q \rightarrow \ell_i^- \ell_j^+$ are combined we compute the average

$$\mathcal{B}[B_q \rightarrow \ell_i \ell_j] = (\mathcal{B}[B_q \rightarrow \ell_i^+ \ell_j^-] + \mathcal{B}[B_q \rightarrow \ell_j^+ \ell_i^-]) / 2.$$

In order to obtain the upper limits we insert the biggest allowed values for $\text{Abs}[\epsilon_{ij}^{d,\ell}]$. For $\epsilon_{23,32}^d$ ($\epsilon_{13,31}^d$) we use the biggest allowed absolute value compatible with the bounds from $B_s \rightarrow \mu^+ \mu^-$ ($B_d \rightarrow \mu^+ \mu^-$). As we can see from Fig. 4 (Fig. 5) the absolute value for ϵ_{32}^d (ϵ_{31}^d) can be bigger than ϵ_{23}^d (ϵ_{13}^d). For the leptonic parameters $\epsilon_{13,31}^\ell$ and $\epsilon_{23,32}^\ell$ we use the constraints obtained from $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, $\tau^- \rightarrow e^- \mu^+ \mu^-$ (see Sec. VC)

$$\begin{aligned} |\epsilon_{31,13}^\ell| &\leq 4.2 \times 10^{-3} \left(\frac{m_H / 500 \text{ GeV}}{\tan \beta / 50} \right)^2, \\ |\epsilon_{32,23}^\ell| &\leq 3.7 \times 10^{-3} \left(\frac{m_H / 500 \text{ GeV}}{\tan \beta / 50} \right)^2, \end{aligned} \quad (72)$$

while for $\epsilon_{12,21}^\ell$ we use the combined constraints from $\mu^- \rightarrow e^- e^+ e^-$ and from $\mu \rightarrow e \gamma$ (see Sec. VID).

Our results are shown in Fig. 35. We see that for bigger Higgs masses larger values for the branching ratios are possible.




FIG. 35: Upper limits on the branching ratios of the lepton flavor violating B meson decays as a function of m_H for $\tan \beta = 30$ (yellow), $\tan \beta = 40$ (red) and $\tan \beta = 50$ (blue).

B. Radiative lepton decays: $\tau \rightarrow \mu\gamma$, $\tau \rightarrow e\gamma$ and $\mu \rightarrow e\gamma$.

In Sec. VC and Sec. VID we found that the radiative lepton decays $\tau \rightarrow \mu\gamma$ and $\tau \rightarrow e\gamma$ give less stringent bounds on the parameters $\epsilon_{23,32}^\ell$ and $\epsilon_{13,31}^\ell$ than the processes $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ and $\tau^- \rightarrow e^- \mu^+ \mu^-$ while the constraints on $\epsilon_{12,21}^\ell$ from $\mu \rightarrow e\gamma$ are stronger than the ones from $\mu^- \rightarrow e^- e^+ e^-$.

There are however interesting correlations between these decays in the type-III 2HDM. In the large $\tan \beta$ limit and for $v \ll m_H$ we obtain the following relation

$$\frac{\mathcal{B}[\ell_i \rightarrow \ell_f \gamma]}{\mathcal{B}[\ell_i^- \rightarrow \ell_f^- \ell_j^+ \ell_j^-]} = \frac{\alpha_{em}}{24\pi} \frac{|m_{\ell_i}/v - \epsilon_{ii}^\ell|^2 \left(|\epsilon_{if}^\ell|^2 + 4 |\epsilon_{fi}^\ell|^2 \right)}{|m_{\ell_j}/v - \epsilon_{jj}^\ell|^2 \left(|\epsilon_{if}^\ell|^2 + |\epsilon_{fi}^\ell|^2 \right)}. \quad (73)$$

As already noted in Sec. VID, we stress that this formula is only a good approximation for very heavy Higgs due to the large logarithmic term in the expression for $\ell_i \rightarrow \ell_f \gamma$ (see Eq. (96)). Therefore, the relation in Eq. (73) gets modified for lighter Higgs masses as shown in Fig. 36. We see that, as expected, for very large Higgs masses the ratios approach




FIG. 36: Left: $\frac{\mathcal{B}[\mu \rightarrow e\gamma]}{\mathcal{B}[\mu^- \rightarrow e^- e^+ e^-]}$ as a function of m_H assuming that only ϵ_{12}^l (red) or ϵ_{21}^l (blue) is different from zero for $\tan\beta = 50$. Right: $\frac{\mathcal{B}[\tau \rightarrow \mu\gamma]}{\mathcal{B}[\tau^- \rightarrow \mu^- \mu^+ \mu^-]}$ as a function of m_H assuming that only ϵ_{23}^l (red) or ϵ_{32}^l (blue) is different from zero for $\tan\beta = 50$.

For scenarios in which both ϵ_{23}^l and ϵ_{32}^l (ϵ_{12}^l and ϵ_{21}^l) are different from zero the 2HDM of type III predicts the ratio $\frac{\mathcal{B}[\tau \rightarrow \mu\gamma]}{\mathcal{B}[\tau^- \rightarrow \mu^- \mu^+ \mu^-]}$ $\left(\frac{\mathcal{B}[\mu \rightarrow e\gamma]}{\mathcal{B}[\mu^- \rightarrow e^- e^+ e^-]} \right)$ to be within the yellow region. These ratios are to a good approximation independent of $\tan\beta$ for $\tan\beta \gtrsim 20$. The behavior of $\frac{\mathcal{B}[\tau \rightarrow e\gamma]}{\mathcal{B}[\tau^- \rightarrow e^- \mu^+ \mu^-]}$ (not shown here) is very similar to the case of $3 \rightarrow 2$ transitions.

$$\begin{aligned} \frac{\mathcal{B}[\ell_i \rightarrow \ell_f \gamma]}{\mathcal{B}[\ell_i^- \rightarrow \ell_f^- \ell_j^+ \ell_j^-]} &= \frac{\alpha_{em}}{24\pi} \frac{m_{\ell_i}^2}{m_{\ell_j}^2} \text{ for } \epsilon_{if}^\ell \neq 0, \\ \frac{\mathcal{B}[\ell_i \rightarrow \ell_f \gamma]}{\mathcal{B}[\ell_i^- \rightarrow \ell_f^- \ell_j^+ \ell_j^-]} &= \frac{\alpha_{em}}{6\pi} \frac{m_{\ell_i}^2}{m_{\ell_j}^2} \text{ for } \epsilon_{fi}^\ell \neq 0, \end{aligned} \quad (74)$$

where, we assumed that $\epsilon_{jj}^\ell/\epsilon_{ii}^\ell = m_{\ell_j}/m_{\ell_i}$ and that only one flavor changing element $\epsilon_{fi}^\ell, \epsilon_{if}^\ell$ is different from zero.

IX. CONCLUSIONS

In this article we studied in detail the flavor phenomenology of a 2HDM with general Yukawa couplings. Motivated by the fact that the 2HDM of type III is the decoupling limit of the MSSM we assumed a MSSM-like Higgs potential. In our analysis we proceeded in several steps:

1. We gave order of magnitude constraints on the parameters $\epsilon_{ij}^{q,\ell}$ from 't Hooft's naturalness criterion and found that all couplings except $\epsilon_{i3,3i}^u$ and $\epsilon_{21,22}^u$ should be much smaller than one.
2. Considering tree-level FCNC processes we constrained the elements ϵ_{ij}^d ($i \neq j$) and $\epsilon_{12,21}^u$ from neutral meson decays to muons and from $\Delta F = 2$ processes, finding that they are tiny for the values of m_H and $\tan\beta$ under investigation (assuming $\epsilon_{ij}^\ell = 0$).

In the lepton sector the absolute values of all flavor off-diagonal elements ϵ_{ij}^ℓ were constrained from $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, $\tau^- \rightarrow e^- \mu^+ \mu^-$ and $\mu^- \rightarrow e^- e^+ e^-$ to be very small.

3. After having found that the off-diagonal elements ϵ_{ij}^d must be very small due to constraints from tree-level contributions to FCNC processes we considered charged Higgs contributions to $K-\bar{K}$, $B_s-\bar{B}_s$, $B_d-\bar{B}_d$ mixing and $b \rightarrow s(d)\gamma$ arising at the one-loop level. In these contributions the so far unconstrained elements $\epsilon_{i3,3i}^u$ (and also ϵ_{22}^u) enter for the first time and we found that, setting $\epsilon_{ij}^d = 0$ (with $i \neq j$), $\epsilon_{13,23}^u$ should be rather small. Furthermore, the electric dipole moment of the neutron and of the charged leptons constrain ϵ_{11}^u , ϵ_{22}^u , ϵ_{21}^u , ϵ_{31}^u and ϵ_{ij}^ℓ , respectively. Respecting all other constraints, no sizable effect in a_μ is possible.
4. Keeping in mind the constraints from the previous steps, we considered the possible effects in charged current processes. Here we found that tests for lepton flavor universality constrain the differences $\epsilon_{ii}^\ell/m_{\ell_i} - \epsilon_{jj}^\ell/m_{\ell_j}$. Most importantly, the unconstrained elements ϵ_{31}^u and ϵ_{32}^u enter the processes $B \rightarrow \tau\nu$ and $B \rightarrow D^{(*)}\tau\nu$ directly (without CKM suppression) and can remove the tension between experiment and theory prediction observed in the SM simultaneously.
5. Finally we gave upper limits on the lepton flavor violating neutral B meson decays in the 2HDM of type III and correlated the radiative lepton decays to $\tau^- \rightarrow \mu^- \mu^+ \mu^-$, $\tau^- \rightarrow e^- \mu^+ \mu^-$ and $\mu^- \rightarrow e^- e^+ e^-$.

In Table VII and VIII we list all processes which have been under consideration and quote the constraints placed on the parameters $\epsilon_{ij}^{q,\ell}$ for our benchmark point $m_H = 500$ GeV and $\tan\beta = 50$.

In summary, combining the constraints from Table VII and VIII the following bounds on the absolute values of the parameters ϵ_{ij}^q and ϵ_{ij}^ℓ (for our benchmark point with $m_H =$

Observable	Results
Neutral meson decays to muons	
$B_s \rightarrow \mu^+ \mu^-$	$ \epsilon_{32}^d \leq 3.0 \times 10^{-5}$, $ \epsilon_{23}^d \leq 1.9 \times 10^{-5}$, $ \epsilon_{22}^\ell \leq 2.0 \times 10^{-3}$
$B_d \rightarrow \mu^+ \mu^-$	$ \epsilon_{31}^d \leq 1.1 \times 10^{-5}$, $ \epsilon_{13}^d \leq 9.4 \times 10^{-6}$
$K_L \rightarrow \mu^+ \mu^-$	$ \epsilon_{21}^d \leq 1.6 \times 10^{-6}$, $ \epsilon_{12}^d \leq 1.6 \times 10^{-6}$
$\bar{D}^0 \rightarrow \mu^+ \mu^-$	$ \epsilon_{21}^u \leq 3.0 \times 10^{-2}$, $ \epsilon_{12}^u \leq 3.0 \times 10^{-2}$
$\Delta F = 2$ processes	
$B_s - \bar{B}_s$ mixing	$ \epsilon_{23}^d \epsilon_{32}^{d*} \leq 9.2 \times 10^{-10}$, $ \epsilon_{23}^u \leq 0.18$, $ \epsilon_{32}^u \leq 1.7$, $ \epsilon_{33}^u \leq 0.7$
$B_d - \bar{B}_d$ mixing	$ \epsilon_{13}^d \epsilon_{31}^{d*} \leq 3.9 \times 10^{-11}$, $ \epsilon_{23}^u \leq 0.2$, $ \epsilon_{13}^u \leq 0.04$, $ \epsilon_{31}^u \leq 1.9$
$K - \bar{K}$ mixing	$ \epsilon_{12}^d \epsilon_{21}^{d*} \leq 1.0 \times 10^{-12}$, $ \epsilon_{22}^u \leq 0.25$, $ \epsilon_{23}^u \leq 0.14$
$D - \bar{D}$ mixing	$ \epsilon_{12}^u \epsilon_{21}^{u*} \leq 2.0 \times 10^{-8}$, $ \epsilon_{32}^u \epsilon_{31}^{u*} \leq 0.02$
Radiative B decays	
$b \rightarrow s\gamma$	$ \epsilon_{23}^u \leq 0.024$, $ \epsilon_{33}^u \leq 0.55$
$b \rightarrow d\gamma$	$ \epsilon_{13}^u \leq 7.0 \times 10^{-3}$
Radiative lepton decays	
$\mu \rightarrow e\gamma$	$ \epsilon_{12}^\ell \leq 1.7 \times 10^{-4}$, $ \epsilon_{21}^\ell \leq 2.2 \times 10^{-4}$, $55 \leq \frac{\mathcal{B}[\mu \rightarrow e\gamma]}{\mathcal{B}[\mu^- \rightarrow e^- e^+ e^-]} \leq 86$
$\tau \rightarrow e\gamma$	$0.19 \leq \frac{\mathcal{B}[\tau \rightarrow e\gamma]}{\mathcal{B}[\tau^- \rightarrow e^- \mu^+ \mu^-]} \leq 0.35$
$\tau \rightarrow \mu\gamma$	$0.19 \leq \frac{\mathcal{B}[\tau \rightarrow \mu\gamma]}{\mathcal{B}[\tau^- \rightarrow \mu^- \mu^+ \mu^-]} \leq 0.35$
Neutral current lepton decays	
$\mu^- \rightarrow e^- e^+ e^-$	$ \epsilon_{12,21}^\ell \leq 2.3 \times 10^{-3}$
$\tau^- \rightarrow e^- \mu^+ \mu^-$	$ \epsilon_{13,31}^\ell \leq 4.2 \times 10^{-3}$
$\tau^- \rightarrow \mu^- \mu^+ \mu^-$	$ \epsilon_{23,32}^\ell \leq 3.7 \times 10^{-3}$

TABLE VII: Results obtained in the type-III 2HDM from various processes for $\tan\beta = 50$ and $m_H = 500$ GeV.

Observable	Results
Charged current processes	
$B \rightarrow \tau\nu$	$2.7 \times 10^{-3} \leq \epsilon_{31}^u \leq 2.0 \times 10^{-2}, \epsilon_{i3}^\ell \leq 6.0 \times 10^{-2}$
$B \rightarrow D\tau\nu$ & $B \rightarrow D^*\tau\nu$	$0.43 \leq \epsilon_{32}^u \leq 0.74$
$D_s \rightarrow \tau\nu$ & $D_{(s)} \rightarrow \mu\nu$	$ \text{Re}[\epsilon_{22}^u] \leq 0.2$
$D \rightarrow \tau\nu$	–
$K \rightarrow \mu(e)\nu/\pi \rightarrow \mu(e)\nu$	$ \text{Re}[\epsilon_{22}^d] \leq 1.0 \times 10^{-3}$
$K(\pi) \rightarrow e\nu/K(\pi) \rightarrow \mu\nu$	$ \text{Re}[\epsilon_{i1}^\ell] \leq 2.0 \times 10^{-6}, \text{Re}[\epsilon_{i2}^\ell] \leq 5.0 \times 10^{-4}$
$\tau \rightarrow K(\pi)\nu/K(\pi) \rightarrow \mu\nu$	$-4.0 \times 10^{-2} \leq \text{Re}[\epsilon_{i3}^\ell] \leq 2.0 \times 10^{-2}$
$\tau \rightarrow K\nu/\tau \rightarrow \pi\nu$	$ \epsilon_{i3}^\ell \leq 0.14$
EDMs and anomalous magnetic moments	
d_e	$ \text{Im}[\epsilon_{12}^\ell\epsilon_{21}^\ell] \leq 2.5 \times 10^{-8}, \text{Im}[\epsilon_{13}^\ell\epsilon_{31}^\ell] \leq 2.5 \times 10^{-9}$
d_μ	–
d_τ	–
d_n	$ \text{Im}[\epsilon_{11}^u] \leq 2.2 \times 10^{-2}, \text{Im}[\epsilon_{22}^u] \leq 1.1 \times 10^{-1}, \text{Arg}[\epsilon_{31}^u] = \text{Arg}[V_{ub}] \pm \pi$
a_μ	Deviation from the SM cannot be explained
LVF B meson decays	
$B_s \rightarrow \tau\mu$	$\mathcal{B}[B_s \rightarrow \tau\mu] \leq 2.0 \times 10^{-7}$
$B_s \rightarrow \mu e$	$\mathcal{B}[B_s \rightarrow \mu e] \leq 9.2 \times 10^{-10}$
$B_s \rightarrow \tau e$	$\mathcal{B}[B_s \rightarrow \tau e] \leq 2.8 \times 10^{-7}$
$B_d \rightarrow \tau\mu$	$\mathcal{B}[B_d \rightarrow \tau\mu] \leq 2.1 \times 10^{-8}$
$B_d \rightarrow \mu e$	$\mathcal{B}[B_d \rightarrow \mu e] \leq 9.2 \times 10^{-11}$
$B_d \rightarrow \tau e$	$\mathcal{B}[B_d \rightarrow \tau e] \leq 2.8 \times 10^{-8}$

TABLE VIII: Results obtained in the type-III 2HDM from various processes for $\tan\beta = 50$ and $m_H = 500$ GeV.

500 GeV and $\tan \beta = 50$) are obtained:

$$\begin{aligned}
|\epsilon_{ij}^u| &\leq \begin{pmatrix} 3.4 \times 10^{-4} & 3.0 \times 10^{-2} & 7.0 \times 10^{-3} \\ 3.0 \times 10^{-2} & 1.4 \times 10^{-1} & 2.4 \times 10^{-2} \\ 2.0 \times 10^{-2} & 7.4 \times 10^{-1} & 5.5 \times 10^{-1} \end{pmatrix}_{ij} \\
|\epsilon_{ij}^d| &\leq \begin{pmatrix} 1.3 \times 10^{-4} & 1.6 \times 10^{-6} & 9.4 \times 10^{-6} \\ 1.6 \times 10^{-6} & 2.6 \times 10^{-4} & 2.0 \times 10^{-5} \\ 1.1 \times 10^{-5} & 3.0 \times 10^{-5} & 1.4 \times 10^{-2} \end{pmatrix}_{ij} \\
|\epsilon_{ij}^\ell| &\leq \begin{pmatrix} 2.9 \times 10^{-6} & 1.7 \times 10^{-4} & 4.2 \times 10^{-3} \\ 2.2 \times 10^{-4} & 6.1 \times 10^{-4} & 3.7 \times 10^{-3} \\ 4.2 \times 10^{-3} & 3.7 \times 10^{-3} & 1.0 \times 10^{-2} \end{pmatrix}_{ij}
\end{aligned} \tag{75}$$

These bounds hold in the absence of large cancellations between different contributions. Note that in Eq. (75) we applied the naturalness bounds in case they were stronger than the experimental limits.

It is interesting that $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$ can be explained simultaneously in the 2HDM of type III without violating bounds from other observables and without significant fine-tuning. It remains to be seen if these tensions with the SM remain when updated experimental results and improved theory predictions will be available in the future. In order to further test the model and constrain the parameters ϵ_{32}^u (ϵ_{31}^u) we propose to study $H^0, A^0 \rightarrow \bar{t}c$ ($H^0, A^0 \rightarrow \bar{t}u$) at the LHC.

Acknowledgments : This work is supported by the Swiss National Science Foundation. We thank Jernej Kamenik and Ulrich Nierste for collaboration in the early stages of this work. A.C. also thanks Jernej Kamenik for useful comments on the manuscript.

X. APPENDIX

In this appendix, we collect the Wilson coefficients (to the relevant precision at the matching scale) which are needed for the calculation of $b \rightarrow s(d)\gamma$, $\Delta F = 2$ processes (i.e. neutral meson mixing), leptonic neutral meson decays ($\Delta F = 1$ processes), $B \rightarrow \tau\nu$, $B \rightarrow D\tau\nu$, $B \rightarrow D^*\tau\nu$, $D_{(s)} \rightarrow \ell\nu_\ell$, $K(\pi) \rightarrow \ell\nu_\ell$, $\tau \rightarrow K(\pi)\nu$, LFV radiative lepton transitions, EDMs of charged leptons and neutron, as well as the AMM of the muon. In addition, we give general expressions for some branching ratios, the explicit form of the loop functions entering our results and summarize the input parameters used in our analysis in tabular form.

A. Loop functions

We give the explicit form of the loop functions entering our results. In the limit of vanishing external momentum the one and two-point Passarino Veltman functions [147] are defined as

$$\begin{aligned} A_0(m^2) &= \frac{16\pi^2}{i}\mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2 - m^2)}, \\ B_0(m_1^2, m_2^2) &= \frac{16\pi^2}{i}\mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{1}{(k^2 - m_1^2)(k^2 - m_2^2)}, \end{aligned} \quad (76)$$

where μ is the renormalization scale.

The loop functions C_0 (three-point) and D_0 (four-point) are defined in analogy to B_0 , but with three and four propagators, respectively. Evaluating these loop functions yields (with $d = 4 - 2\varepsilon$)

$$\begin{aligned} A_0(m^2) &= m^2 \left[1 + \frac{1}{\varepsilon} - \gamma_E + \ln(4\pi) + \ln\left(\frac{\mu^2}{m^2}\right) \right] + O(\varepsilon), \\ B_0(m_1^2, m_2^2) &= 1 + \frac{1}{\varepsilon} - \gamma_E + \ln(4\pi) + \frac{m_1^2 \ln\left(\frac{\mu^2}{m_1^2}\right) - m_2^2 \ln\left(\frac{\mu^2}{m_2^2}\right)}{m_1^2 - m_2^2} + O(\varepsilon), \end{aligned} \quad (77)$$

$$\begin{aligned} C_0(m_1^2, m_2^2, m_3^2) &= \frac{B_0(m_1^2, m_2^2) - B_0(m_1^2, m_3^2)}{m_2^2 - m_3^2} \\ &= \frac{m_1^2 m_2^2 \ln\left(\frac{m_1^2}{m_2^2}\right) + m_3^2 m_2^2 \ln\left(\frac{m_2^2}{m_3^2}\right) + m_3^2 m_1^2 \ln\left(\frac{m_3^2}{m_1^2}\right)}{(m_1^2 - m_2^2)(m_3^2 - m_1^2)(m_2^2 - m_3^2)}, \\ D_0(m_1^2, m_2^2, m_3^2, m_4^2) &= \frac{C_0(m_1^2, m_2^2, m_3^2) - C_0(m_1^2, m_2^2, m_4^2)}{m_3^2 - m_4^2}. \end{aligned} \quad (78)$$

Here, the one and the two-point loop functions A_0, B_0 are UV-divergent and ε is the UV-regulator.

At various places also the functions C_2 and D_2 appear, which have, compared to C_0 and D_0 , an additional factor k^2 in the numerator of the integrand. These functions read

$$\begin{aligned} C_2(m_1^2, m_2^2, m_3^2) &= B_0(m_1^2, m_2^2) + m_3^2 C_0(m_1^2, m_2^2, m_3^2), \\ D_2(m_1^2, m_2^2, m_3^2, m_4^2) &= C_0(m_1^2, m_2^2, m_3^2) + m_4^2 D_0(m_1^2, m_2^2, m_3^2, m_4^2). \end{aligned} \quad (79)$$

B. Radiative $b \rightarrow s(d)\gamma$ decays

Concerning new physics contributions to $b \rightarrow s(d)\gamma$, we work in leading logarithmic (LL) precision in this paper. As mentioned before, we use these processes to constrain certain elements ϵ_{ij}^u . For this purpose, we put the ϵ_{ij}^d -couplings (which are already constrained to be very small) to zero. When also neglecting the mass of the strange quark and further neglecting operators with mass dimension higher than six, we obtain the same effective Hamiltonian as in the SM, reading for $b \rightarrow s\gamma$ (see e.g. Ref. [47]).

$$\mathcal{H}_{eff}^{b \rightarrow s\gamma} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum_i C_i O_i. \quad (80)$$

For $b \rightarrow d\gamma$ the CKM structure is slightly more complicated (see e.g. Ref. [95]). In our approximation only the Wilson coefficients C_7 and C_8 of the operators

$$O_7 = \frac{e}{16\pi^2} m_b \bar{s} \sigma^{\mu\nu} P_R b F_{\mu\nu} \quad ; \quad O_8 = \frac{g_s}{16\pi^2} m_b \bar{s} \sigma^{\mu\nu} T^a P_R b G_{\mu\nu}^a \quad (81)$$

get new physics contributions. They are induced through charged Higgs bosons propagating in the loop (neutral Higgs boson exchange leads to power suppressed contributions which we neglect). For $b \rightarrow s\gamma$ the new physics contributions read (with $y_j = m_{u_j}^2/m_{H^\pm}^2$ and $\lambda_t = V_{tb} V_{ts}^*$)

$$\begin{aligned} C_7^{NP} &= \frac{v^2}{\lambda_t m_b} \sum_{j=1}^3 \Gamma_{u_j d_2}^{RLH^{\pm*}} \Gamma_{u_j d_3}^{LRH^\pm} \frac{C_{7,XY}^0(y_j)}{m_{u_j}} \\ &+ \frac{v^2}{\lambda_t} \sum_{j=1}^3 \Gamma_{u_j d_2}^{RLH^{\pm*}} \Gamma_{u_j d_3}^{RLH^\pm} \frac{C_{7,YY}^0(y_j)}{m_{u_j}^2}, \\ C_8^{NP} &= \frac{v^2}{\lambda_t m_b} \sum_{j=1}^3 \Gamma_{u_j d_2}^{RLH^{\pm*}} \Gamma_{u_j d_3}^{LRH^\pm} \frac{C_{8,XY}^0(y_j)}{m_{u_j}} \\ &+ \frac{v^2}{\lambda_t} \sum_{j=1}^3 \Gamma_{u_j d_2}^{RLH^{\pm*}} \Gamma_{u_j d_3}^{RLH^\pm} \frac{C_{8,YY}^0(y_j)}{m_{u_j}^2}, \end{aligned} \quad (82)$$

while for $b \rightarrow d\gamma$ the label d_2 and $\lambda_t = V_{tb} V_{ts}^*$ have to be replaced by d_1 and $\lambda_t = V_{tb} V_{td}^*$, respectively. The functions $C_{7,XY}^0$, $C_{7,YY}^0$, $C_{8,XY}^0$ and $C_{8,YY}^0$ were introduced in Ref. [47];

their explicit form reads

$$\begin{aligned}
C_{7,XY}^0(y_j) &= \frac{y_j}{12} \left[\frac{-5y_j^2 + 8y_j - 3 + (6y_j - 4) \ln y_j}{(y_j - 1)^3} \right] , \\
C_{8,XY}^0(y_j) &= \frac{y_j}{4} \left[\frac{-y_j^2 + 4y_j - 3 - 2 \ln y_j}{(y_j - 1)^3} \right] , \\
C_{7,YY}^0(y_j) &= \frac{y_j}{72} \left[\frac{-8y_j^3 + 3y_j^2 + 12y_j - 7 + (18y_j^2 - 12y_j) \ln y_j}{(y_j - 1)^4} \right] , \\
C_{8,YY}^0(y_j) &= \frac{y_j}{24} \left[\frac{-y_j^3 + 6y_j^2 - 3y_j - 2 - 6y_j \ln y_j}{(y_j - 1)^4} \right] .
\end{aligned} \tag{83}$$

In Eq. (82) we retained the contributions from internal up- and charm-quarks, although these contributions are subleading.

C. Wilson coefficients for $\Delta F = 2$ processes

The extended Higgs sector of our 2HDM of type-III also leads to extra contributions to $\Delta F = 2$ processes (B_s , B_d , Kaon and D mixing) which can be matched onto the effective Hamiltonian

$$\mathcal{H}_{eff}^{\Delta F=2} = \sum_{j=1}^5 C_j O_j + \sum_{j=1}^3 C'_j O'_j + h.c. , \tag{84}$$

where the operators read in the case of B_s mixing

$$\begin{aligned}
O_1 &= (\bar{s}_\alpha \gamma^\mu P_L b_\alpha) (\bar{s}_\beta \gamma^\mu P_L b_\beta) , & O_2 &= (\bar{s}_\alpha P_L b_\alpha) (\bar{s}_\beta P_L b_\beta) , \\
O_3 &= (\bar{s}_\alpha P_L b_\beta) (\bar{s}_\beta P_L b_\alpha) , & O_4 &= (\bar{s}_\alpha P_L b_\alpha) (\bar{s}_\beta P_R b_\beta) , \\
O_5 &= (\bar{s}_\alpha P_L b_\beta) (\bar{s}_\beta P_R b_\alpha) .
\end{aligned} \tag{85}$$

α and β are color indices and the primed operators can be obtained from $O_{1,2,3}$ by interchanging L and R . Similarly, the corresponding operator bases for B_d , Kaon and D mixing follow from Eq. (85) through simple adjustment of the indices.

In the following subsections we present the contributions to these Wilson coefficients arising from: 1.) one-loop box diagrams with charged Higgs boson exchange; 2.) tree-level contributions induced by neutral Higgs boson exchange; 3.) box diagrams involving neutral Higgs bosons, relevant in the case of D mixing.

1. Charged Higgs box contributions

For definiteness, let us consider B_s mixing. The corresponding Wilson coefficients for B_d and Kaon mixing follow by a simple adjustment of the indices. We have performed our

calculation in a general R_ξ gauge. The non-vanishing Wilson coefficients from pure charged Higgs boxes are given by

$$\begin{aligned}
C_1 &= \frac{-1}{128\pi^2} \sum_{j,k=1}^3 \Gamma_{u_j d_2}^{RL H^\pm \star} \Gamma_{u_j d_3}^{RL H^\pm} \Gamma_{u_k d_2}^{RL H^\pm \star} \Gamma_{u_k d_3}^{RL H^\pm} D_2 \left(m_{u_j}^2, m_{u_k}^2, m_{H^\pm}^2, m_{H^\pm}^2 \right), \\
C_2 &= \frac{-1}{32\pi^2} \sum_{j,k=1}^3 m_{u_j} m_{u_k} \Gamma_{u_j d_2}^{LR H^\pm \star} \Gamma_{u_j d_3}^{RL H^\pm} \Gamma_{u_k d_2}^{LR H^\pm \star} \Gamma_{u_k d_3}^{RL H^\pm} D_0 \left(m_{u_j}^2, m_{u_k}^2, m_{H^\pm}^2, m_{H^\pm}^2 \right), \\
C_4 &= \frac{-1}{16\pi^2} \sum_{j,k=1}^3 m_{u_j} m_{u_k} \Gamma_{u_j d_2}^{LR H^\pm \star} \Gamma_{u_j d_3}^{RL H^\pm} \Gamma_{u_k d_2}^{RL H^\pm \star} \Gamma_{u_k d_3}^{LR H^\pm} D_0 \left(m_{u_j}^2, m_{u_k}^2, m_{H^\pm}^2, m_{H^\pm}^2 \right), \\
C_5 &= \frac{1}{32\pi^2} \sum_{j,k=1}^3 \Gamma_{u_j d_2}^{LR H^\pm \star} \Gamma_{u_j d_3}^{LR H^\pm} \Gamma_{u_k d_2}^{RL H^\pm \star} \Gamma_{u_k d_3}^{RL H^\pm} D_2 \left(m_{u_j}^2, m_{u_k}^2, m_{H^\pm}^2, m_{H^\pm}^2 \right).
\end{aligned} \tag{86}$$

The sum of the charged Higgs– W^\pm and charged Higgs–Goldstone-boson boxes is given by

$$\begin{aligned}
C_1 &= \frac{g_2^2}{32\pi^2} \sum_{j,k=1}^3 \left(m_{u_j} m_{u_k} V_{j2}^\star V_{k3} \Gamma_{u_j d_3}^{RL H^\pm} \Gamma_{u_k d_2}^{RL H^\pm \star} \right. \\
&\quad \left. \times \frac{4M_W^2 D_0 \left(M_W^2, m_{H^\pm}^2, m_{u_j}^2, m_{u_k}^2 \right) - D_2 \left(M_W^2, m_{H^\pm}^2, m_{u_j}^2, m_{u_k}^2 \right)}{4M_W^2} \right) \\
C_4 &= \frac{1}{16\pi^2} \frac{g_2^2}{2} \sum_{j,k=1}^3 \left(V_{j3} V_{k2}^\star \Gamma_{u_j d_2}^{LR H^\pm \star} \Gamma_{u_k d_3}^{LR H^\pm} \right. \\
&\quad \left. \times \frac{C_2 \left(\xi M_W^2, m_{H^+}^2, m_{u_j}^2 \right) - C_2 \left(m_{H^+}^2, m_{u_j}^2, m_{u_k}^2 \right) + m_{u_k}^2 C_0 \left(\xi M_W^2, m_{H^\pm}^2, m_{u_k}^2 \right)}{M_W^2} \right)
\end{aligned} \tag{87}$$

We stress here, that we want to use B_s mixing only to constrain certain ϵ_{ij}^u –couplings, because the ϵ_{ij}^d –quantities are already constrained to be very small. We therefore only took systematically into account those contributions to the Wilson coefficients which stay different from zero in the limit $\epsilon_{ij}^d \rightarrow 0$. At first sight, the Wilson coefficient C_4 seems to be gauge dependent. However, when using the unitarity of the CKM matrix (entering the expression for C_4 both, explicitly and implicitly through the Γ –quantities), we find that the ξ –dependent terms are always proportional to an element ϵ_{ij}^d , which we put to zero in our analysis. Also note that our result agrees with the one of Ref. [148]. The only difference is that we neglected gauge dependent terms corresponding to higher dimensional operators. The Wilson coefficients of the primed operators can be obtained by interchanging L and R in the corresponding unprimed ones.

2. Tree-level H_k^0 contribution

The Wilson coefficients from neutral Higgs mediated tree-level contributions to B_s mixing read:

$$\begin{aligned}
C_2^{H_k^0} &= \sum_{k=1}^3 \frac{-1}{2m_{H_k^0}^2} (\Gamma_{d_3 d_2}^{LR H_k^0 \star})^2 \\
C_2^{\prime H_k^0} &= \sum_{k=1}^3 \frac{-1}{2m_{H_k^0}^2} (\Gamma_{d_2 d_3}^{LR H_k^0})^2 \\
C_4^{H_k^0} &= \sum_{k=1}^3 \frac{-1}{m_{H_k^0}^2} \Gamma_{d_2 d_3}^{LR H_k^0} \Gamma_{d_3 d_2}^{LR H_k^0 \star}
\end{aligned} \tag{88}$$

The corresponding coefficients for B_d , Kaon and D mixing follow by a careful adjustment of the indices.

Note that in the limit of large $\tan \beta$ and $m_A \gg v$, $C_2^{H_k^0}$ and $C_2^{\prime H_k^0}$ vanish and we only get a contribution to $C_4^{H_k^0}$.

3. Neutral Higgs box contribution to D mixing

The Wilson coefficients resulting from the neutral Higgs box contribution to D mixing are given as

$$\begin{aligned}
C_1 &= \frac{-1}{128\pi^2} \sum_{j_1, j_2=1}^3 \sum_{k_1, k_2=1}^3 \Gamma_{u_2 u_{j_1}}^{LR H_{k_1}^0 \star} \Gamma_{u_1 u_{j_1}}^{LR H_{k_2}^0} \Gamma_{u_2 u_{j_2}}^{LR H_{k_2}^0 \star} \Gamma_{u_1 u_{j_2}}^{LR H_{k_1}^0} D_2 \left(m_{u_{j_1}}^2, m_{u_{j_2}}^2, m_{H_{k_1}^0}^2, m_{H_{k_2}^0}^2 \right), \\
C_2 &= \frac{-1}{32\pi^2} \sum_{j_1, j_2=1}^3 \sum_{k_1, k_2=1}^3 m_{u_{j_1}} m_{u_{j_2}} \Gamma_{u_{j_1} u_1}^{LR H_{k_1}^0 \star} \Gamma_{u_2 u_{j_1}}^{LR H_{k_2}^0 \star} \Gamma_{u_{j_2} u_1}^{LR H_{k_1}^0 \star} \Gamma_{u_2 u_{j_2}}^{LR H_{k_2}^0 \star} \\
&\quad \times D_0 \left(m_{u_{j_1}}^2, m_{u_{j_2}}^2, m_{H_{k_1}^0}^2, m_{H_{k_2}^0}^2 \right), \\
C_3 &= 0, \\
C_4 &= \frac{-1}{16\pi^2} \sum_{j_1, j_2=1}^3 \sum_{k_1, k_2=1}^3 m_{u_{j_1}} m_{u_{j_2}} \Gamma_{u_{j_1} u_1}^{LR H_{k_1}^0 \star} \Gamma_{u_2 u_{j_1}}^{LR H_{k_2}^0 \star} \Gamma_{u_{j_2} u_2}^{LR H_{k_2}^0} \Gamma_{u_1 u_{j_2}}^{LR H_{k_1}^0} \\
&\quad \times D_0 \left(m_{u_{j_1}}^2, m_{u_{j_2}}^2, m_{H_{k_1}^0}^2, m_{H_{k_2}^0}^2 \right), \\
C_5 &= \frac{-1}{128\pi^2} \sum_{j_1, j_2=1}^3 \sum_{k_1, k_2=1}^3 \Gamma_{u_{j_1} u_1}^{LR H_{k_1}^0 \star} \Gamma_{u_{j_1} u_2}^{LR H_{k_2}^0} \Gamma_{u_1 u_{j_2}}^{LR H_{k_1}^0} \Gamma_{u_2 u_{j_2}}^{LR H_{k_2}^0 \star} D_2 \left(m_{u_{j_1}}^2, m_{u_{j_2}}^2, m_{H_{k_1}^0}^2, m_{H_{k_2}^0}^2 \right).
\end{aligned} \tag{89}$$

The indices j_1, j_2 describe the internal up-type quarks while k_1, k_2 stand for neutral Higgs indices (H^0, h^0, A^0). Moreover, the primed Wilson coefficients can be obtained from above by the replacement $L \leftrightarrow R$ in the couplings.

D. Semileptonic and leptonic meson decays and tau decays: $B \rightarrow (D^{(*)})\tau\nu$, $D_{(s)} \rightarrow \ell\nu_\ell$, $K(\pi) \rightarrow \ell\nu_\ell$ and $\tau \rightarrow K(\pi)\nu$ processes

These processes are governed by the effective Hamiltonian

$$\mathcal{H}_{\text{eff}} = C_{\text{SM}}^{u_f d_i, \ell_j} O_{\text{SM}}^{u_f d_i, \ell_j} + C_L^{u_f d_i, \ell_j} O_L^{u_f d_i, \ell_j} + C_R^{u_f d_i, \ell_j} O_R^{u_f d_i, \ell_j} + \text{h.c.}, \quad (90)$$

with the operators defined as

$$\begin{aligned} O_{\text{SM}}^{u_f d_i, \ell_j} &= \bar{u}_f \gamma_\mu P_L d_i \bar{\ell}_j \gamma_\mu P_L \nu, \\ O_R^{u_f d_i, \ell_j} &= \bar{u}_f P_R d_i \bar{\ell}_j P_L \nu, \\ O_L^{u_f d_i, \ell_j} &= \bar{u}_f P_L d_i \bar{\ell}_j P_L \nu. \end{aligned} \quad (91)$$

Here, for tauonic B meson decays $\ell_j = \tau$, $d_i = b$ and $u_f = u$ ($u_f = c$) for $B \rightarrow \tau\nu$ ($B \rightarrow D\tau\nu$ and $B \rightarrow D^*\tau\nu$). For $D_s \rightarrow \ell_j\nu$ ($D \rightarrow \ell_j\nu$), $u_f = c$ and $d_i = s$ (d), for $\tau \rightarrow K(\pi)\nu$, $\ell_j = \tau$, $u_f = u$ and $d_i = s$ (d) and for $K(\pi) \rightarrow \ell_j\nu$ we have $\ell_j = \mu, e$, $u_f = u$ and $d_i = s$ (d). The Wilson coefficients in 2HDM of type III at the matching scale read

$$\begin{aligned} C_{\text{SM}}^{u_f d_i, \ell_j} &= \frac{4G_F}{\sqrt{2}} V_{u_f d_i}, \\ C_R^{u_f d_i, \ell_j} &= \frac{-1}{m_{H^\pm}^2} \Gamma_{u_f d_i}^{LRH^\pm} \Gamma_{\nu \ell_j}^{LRH^{\pm*}}, \\ C_L^{u_f d_i, \ell_j} &= \frac{-1}{m_{H^\pm}^2} \Gamma_{u_f d_i}^{RLH^\pm} \Gamma_{\nu \ell_j}^{LRH^{\pm*}}. \end{aligned} \quad (92)$$

E. Lepton flavour violation (LFV): $\ell_i \rightarrow \ell_f \gamma$ processes

The radiative lepton decays $\ell_i \rightarrow \ell_f \gamma$ ($\ell = e, \mu$ or τ) are induced by one-loop penguin diagrams with internal neutral or charged Higgs bosons. The result for the one-loop decay amplitude can be written as a tree-level matrix element of the effective Hamiltonian

$$\mathcal{H}_{\text{eff}} = c_R^{\ell_f \ell_i} O_R^{\ell_f \ell_i} + c_L^{\ell_f \ell_i} O_L^{\ell_f \ell_i}, \quad (93)$$

where $c_R^{\ell_f \ell_i}$ and $c_L^{\ell_f \ell_i}$ are the effective Wilson coefficients of the magnetic dipole operators

$$O_{R(L)}^{\ell_f \ell_i} = m_{\ell_i} \bar{\ell}_f \sigma_{\mu\nu} P_{R(L)} \ell_i F^{\mu\nu}. \quad (94)$$

With these conventions, the branching ratio for the radiative lepton decays $\ell_i \rightarrow \ell_f \gamma$ reads

$$\mathcal{B}[\ell_i \rightarrow \ell_f \gamma] = \frac{m_{\ell_i}^5}{4\pi \Gamma_{\ell_i}} \left(|c_R^{\ell_f \ell_i}|^2 + |c_L^{\ell_f \ell_i}|^2 \right). \quad (95)$$

The neutral Higgs ($H_k^0 = H^0, h^0, A^0$) penguin contribution to $c_R^{\ell_f \ell_i}$ is given by

$$c_{RH_k^0}^{\ell_f \ell_i} = \sum_{k,j=1}^3 \frac{-e}{192\pi^2 m_{H_k^0}^2} \left[\Gamma_{\ell_f \ell_j}^{LRH_k^0} \Gamma_{\ell_i \ell_j}^{LRH_k^{0*}} + \frac{m_{\ell_f}}{m_{\ell_i}} \Gamma_{\ell_j \ell_f}^{LRH_k^0} \Gamma_{\ell_j \ell_i}^{LRH_k^0} - \frac{m_{\ell_j}}{m_{\ell_i}} \Gamma_{\ell_f \ell_j}^{LRH_k^0} \Gamma_{\ell_j \ell_i}^{LRH_k^0} \left(9 + 6 \ln \left(\frac{m_{\ell_j}^2}{m_{H_k^0}^2} \right) \right) \right], \quad (96)$$

and $c_L^{\ell_f \ell_i}$ can be obtained from $c_R^{\ell_f \ell_i}$ by interchanging L and R . Similarly, for the charged Higgs penguin contributions we find

$$c_{LH^\pm}^{\ell_f \ell_i} = \frac{e}{384\pi^2 m_{H^\pm}^2} \sum_{j=1}^3 \Gamma_{\nu_j \ell_i}^{LRH^\pm} \Gamma_{\nu_j \ell_f}^{LRH^{\pm*}}, \quad (97)$$

$$c_{RH^\pm}^{\ell_f \ell_i} = \frac{m_{\ell_f}}{m_{\ell_i}} \frac{e}{384\pi^2 m_{H^\pm}^2} \sum_{j=1}^3 \Gamma_{\nu_j \ell_i}^{LRH^\pm} \Gamma_{\nu_j \ell_f}^{LRH^{\pm*}}.$$

F. Wilson coefficients for EDMs and the anomalous magnetic moment of the muon

1. Wilson coefficients for EDMs of charged leptons and the anomalous magnetic moment of the muon

As in the case of the LFV processes discussed in the previous section, we again have both neutral and charged Higgs penguin contributions to the flavor conserving radiative transitions $\ell_i \rightarrow \ell_i \gamma$. The corresponding effective Hamiltonian is obtained from Eq. (93) and Eq. (94) by identifying ℓ_f with ℓ_i . The contribution to the effective Wilson coefficients related to neutral Higgs bosons (propagating in the loop) reads

$$c_{RH_k^0}^{\ell_i \ell_i} = \sum_{k,j=1}^3 \frac{-e}{192\pi^2 m_{H_k^0}^2} \left[\Gamma_{\ell_i \ell_j}^{LRH_k^0} \Gamma_{\ell_i \ell_j}^{LRH_k^{0*}} + \Gamma_{\ell_j \ell_i}^{LRH_k^0} \Gamma_{\ell_j \ell_i}^{LRH_k^0} - \frac{m_{\ell_j}}{m_{\ell_i}} \Gamma_{\ell_i \ell_j}^{LRH_k^0} \Gamma_{\ell_j \ell_i}^{LRH_k^0} \left(9 + 6 \ln \left(\frac{m_{\ell_j}^2}{m_{H_k^0}^2} \right) \right) \right], \quad (98)$$

$$c_{LH_k^0}^{\ell_i \ell_i} = c_{RH_k^0}^{\ell_i \ell_i}, \quad (99)$$

while the charged Higgs penguin contribution leads to the (real) coefficients

$$c_{LH^\pm}^{\ell_i \ell_i} = c_{RH^\pm}^{\ell_i \ell_i} = \frac{e}{384\pi^2 m_{H^\pm}^2} \sum_{j=1}^3 \Gamma_{\nu_j \ell_i}^{LRH^\pm} \Gamma_{\nu_j \ell_i}^{LRH^{\pm*}}. \quad (100)$$

2. Wilson coefficients for neutron EDM

In this section we consider the transitions $d \rightarrow d\gamma(g)$ and $u \rightarrow u\gamma(g)$ (denoted by $d_d^{(g)}$ and $d_u^{(g)}$) which are the building blocks for the electric dipole moment d_n of the neutron.

As we are only interested in a rough estimate of d_n , we do not include QCD corrections to these building blocks. In this approximation the latter can be described by the effective Hamiltonian

$$\begin{aligned} \mathcal{H}_{eff}^{dd,uu} = & c_R^{dd} m_d \bar{d} \sigma_{\mu\nu} P_R d F^{\mu\nu} + c_L^{dd} m_d \bar{d} \sigma_{\mu\nu} P_L d F^{\mu\nu} + \\ & c_{R,g}^{dd} m_d \bar{d} \sigma_{\mu\nu} P_R T^a d G^{a,\mu\nu} + c_{L,g}^{dd} m_d \bar{d} \sigma_{\mu\nu} P_L T^a d G^{a,\mu\nu} + (d \rightarrow u). \end{aligned} \quad (101)$$

The effective Wilson coefficients $c_{R,L}^{dd,uu}$ and $c_{R,L,g}^{dd,uu}$ again receive neutral and charged Higgs contributions. The neutral contributions of the Wilson coefficients (involved in $d_d^{(g)}$) read

$$\begin{aligned} c_R^{dd,H_k^0} = & \sum_{k,j=1}^3 \frac{e Q_d}{192 \pi^2 m_{H_k^0}^2} \left[\Gamma_{ddj}^{LRH_k^0*} \Gamma_{ddj}^{LRH_k^0} + \Gamma_{d_j d}^{LRH_k^0*} \Gamma_{d_j d}^{LRH_k^0} \right. \\ & \left. - \frac{m_{d_j}}{m_d} \Gamma_{ddj}^{LRH_k^0} \Gamma_{d_j d}^{LRH_k^0} \left(9 + 6 \ln \left(\frac{m_{d_j}^2}{m_{H_k^0}^2} \right) \right) \right], \end{aligned} \quad (102)$$

$$\begin{aligned} c_{R,g}^{dd,H_k^0} = & \sum_{k,j=1}^3 \frac{g_s}{192 \pi^2 m_{H_k^0}^2} \left[\Gamma_{ddj}^{LRH_k^0*} \Gamma_{ddj}^{LRH_k^0} + \Gamma_{d_j d}^{LRH_k^0*} \Gamma_{d_j d}^{LRH_k^0} \right. \\ & \left. - \frac{m_{d_j}}{m_d} \Gamma_{ddj}^{LRH_k^0} \Gamma_{d_j d}^{LRH_k^0} \left(9 + 6 \ln \left(\frac{m_{d_j}^2}{m_{H_k^0}^2} \right) \right) \right], \end{aligned} \quad (103)$$

and $c_{L,(g)}^{dd,H_k^0} = c_{R,(g)}^{dd,H_k^0*}$. The charged Higgs penguin contributions to the Wilson coefficients (involved in $d_d^{(g)}$) read

$$\begin{aligned} c_R^{dd,H^\pm} = & \sum_{j=1}^3 \frac{-e}{16 \pi^2 m_{u_j}^2} \left[\Gamma_{du_j}^{LRH^\pm*} \Gamma_{du_j}^{LRH^\pm} C_{7,YY}^0 \left(\frac{m_{u_j}^2}{m_{H^\pm}^2} \right) + \Gamma_{u_j d}^{LRH^\pm*} \Gamma_{u_j d}^{LRH^\pm} C_{7,YY}^0 \left(\frac{m_{u_j}^2}{m_{H^\pm}^2} \right) \right. \\ & \left. + \Gamma_{du_j}^{LRH^\pm} \Gamma_{u_j d}^{LRH^\pm} \frac{m_{u_j}}{m_d} C_{7,XY}^0 \left(\frac{m_{u_j}^2}{m_{H^\pm}^2} \right) \right], \end{aligned} \quad (104)$$

$$\begin{aligned} c_{R,g}^{dd,H^\pm} = & \sum_{j=1}^3 \frac{-g_s}{16 \pi^2 m_{u_j}^2} \left[\Gamma_{du_j}^{LRH^\pm*} \Gamma_{du_j}^{LRH^\pm} C_{8,YY}^0 \left(\frac{m_{u_j}^2}{m_{H^\pm}^2} \right) + \Gamma_{u_j d}^{LRH^\pm*} \Gamma_{u_j d}^{LRH^\pm} C_{8,YY}^0 \left(\frac{m_{u_j}^2}{m_{H^\pm}^2} \right) \right. \\ & \left. + \Gamma_{du_j}^{LRH^\pm} \Gamma_{u_j d}^{LRH^\pm} \frac{m_{u_j}}{m_d} C_{8,XY}^0 \left(\frac{m_{u_j}^2}{m_{H^\pm}^2} \right) \right], \end{aligned} \quad (105)$$

and $c_{L,(g)}^{dd,H^\pm} = c_{R,(g)}^{dd,H^\pm*}$.

The analogous expressions for $c_{R,(g)}^{uu,H^\pm,H_k^0}$, which are involved in the expressions of $d_u^{(g)}$ are given as

$$\begin{aligned} c_R^{uu,H_k^0} = & \sum_{j,k=1}^3 \frac{-e Q_u}{16 \pi^2 m_{u_j}^2} \left[\Gamma_{uu_j}^{LRH_k^0*} \Gamma_{uu_j}^{LRH_k^0} C_{8,YY}^0 \left(\frac{m_{u_j}^2}{m_{H_k^0}^2} \right) + \Gamma_{u_j u}^{LRH_k^0*} \Gamma_{u_j u}^{LRH_k^0} C_{8,YY}^0 \left(\frac{m_{u_j}^2}{m_{H_k^0}^2} \right) \right. \\ & \left. + \Gamma_{uu_j}^{LRH_k^0} \Gamma_{u_j u}^{LRH_k^0} \frac{m_{u_j}}{m_u} C_{8,XY}^0 \left(\frac{m_{u_j}^2}{m_{H_k^0}^2} \right) \right], \end{aligned} \quad (106)$$

$$c_{R,g}^{uu,H_k^0} = \sum_{j,k=1}^3 \frac{-g_s}{16\pi^2 m_{u_j}^2} \left[\Gamma_{uu_j}^{LRH_k^0*} \Gamma_{uu_j}^{LRH_k^0} C_{8,YY}^0 \left(\frac{m_{u_j}^2}{m_{H_k^0}^2} \right) + \Gamma_{u_j u}^{LRH_k^0*} \Gamma_{u_j u}^{LRH_k^0} C_{8,YY}^0 \left(\frac{m_{u_j}^2}{m_{H_k^0}^2} \right) + \Gamma_{uu_j}^{LRH_k^0} \Gamma_{u_j u}^{LRH_k^0} \frac{m_{u_j}}{m_u} C_{8,XY}^0 \left(\frac{m_{u_j}^2}{m_{H_k^0}^2} \right) \right], \quad (107)$$

$$c_R^{uu,H^\pm} = \sum_{j=1}^3 \frac{-e}{1152\pi^2 m_{H^+}^2} \left[5 \Gamma_{ud_j}^{LRH^\pm*} \Gamma_{ud_j}^{LRH^\pm} + 5 \Gamma_{d_j u}^{LRH^\pm*} \Gamma_{d_j u}^{LRH^\pm} - \Gamma_{ud_j}^{LRH^\pm} \Gamma_{d_j u}^{LRH^\pm} \frac{m_{d_j}}{m_u} 12 \ln \left(\frac{m_{d_j}^2}{m_{H^+}^2} \right) \right], \quad (108)$$

$$c_{R,g}^{uu,H^\pm} = \sum_{j=1}^3 \frac{g_s}{192\pi^2 m_{H^+}^2} \left[\Gamma_{ud_j}^{LRH^\pm*} \Gamma_{ud_j}^{LRH^\pm} + \Gamma_{d_j u}^{LRH^\pm*} \Gamma_{d_j u}^{LRH^\pm} - \Gamma_{ud_j}^{LRH^\pm} \Gamma_{d_j u}^{LRH^\pm} \frac{m_{d_j}}{m_u} \left(9 + 6 \ln \left(\frac{m_{d_j}^2}{m_{H^+}^2} \right) \right) \right]. \quad (109)$$

Again, we have $c_{L,(g)}^{uu,H_k^0} = c_{R,(g)}^{uu,H_k^0*}$ and $c_{L,(g)}^{uu,H^\pm} = c_{R,(g)}^{uu,H^\pm*}$. The loop functions $C_{7,8,XY,YY}^0(y_j)$ are given in Eq. (83).

G. Leptonic decays of neutral mesons

The effective Hamiltonian \mathcal{H}_{eff} which includes the full set of operators for the general decays $PS(\bar{q}_f q_i) \rightarrow \ell_A^+ \ell_B^-$ (PS refers to the pseudo-scalar meson) reads

$$\mathcal{H}_{eff}^{\Delta F=1} = -\frac{G_F^2 M_W^2}{\pi^2} [C_V^{q_f q_i} O_V^{q_f q_i} + C_A^{q_f q_i} O_A^{q_f q_i} + C_S^{q_f q_i} O_S^{q_f q_i} + C_P^{q_f q_i} O_P^{q_f q_i} + \text{primed}] + \text{h.c.} \quad (110)$$

where the operators (together with their primed counterparts) are defined as

$$\begin{aligned} \mathcal{O}_V^{q_f q_i} &= (\bar{q}_f \gamma_\mu P_L q_i) (\bar{\ell}_B \gamma^\mu \ell_A), & \mathcal{O}_A^{q_f q_i} &= (\bar{q}_f \gamma_\mu P_L q_i) (\bar{\ell}_B \gamma^\mu \gamma_5 \ell_A), \\ \mathcal{O}_V^{\prime q_f q_i} &= (\bar{q}_f \gamma_\mu P_R q_i) (\bar{\ell}_B \gamma^\mu \ell_A), & \mathcal{O}_A^{\prime q_f q_i} &= (\bar{q}_f \gamma_\mu P_R q_i) (\bar{\ell}_B \gamma^\mu \gamma_5 \ell_A), \\ \mathcal{O}_S^{q_f q_i} &= (\bar{q}_f P_L q_i) (\bar{\ell}_B \ell_A), & \mathcal{O}_P^{q_f q_i} &= (\bar{q}_f P_L q_i) (\bar{\ell}_B \gamma_5 \ell_A), \\ \mathcal{O}_S^{\prime q_f q_i} &= (\bar{q}_f P_R q_i) (\bar{\ell}_B \ell_A), & \mathcal{O}_P^{\prime q_f q_i} &= (\bar{q}_f P_R q_i) (\bar{\ell}_B \gamma_5 \ell_A). \end{aligned} \quad (111)$$

Making use of the hadronic matrix elements

$$\begin{aligned} \langle 0 | \bar{q}_f \gamma_\mu \gamma_5 q_i | PS \rangle &= i f_{PS} p_{PS}^\mu, \\ \langle 0 | \bar{q}_f \gamma_5 q_i | PS \rangle &= -i f_{PS} \frac{M_{PS}^2}{(m_{q_f} + m_{q_i})}, \end{aligned} \quad (112)$$

one obtains the branching ratio

$$\begin{aligned}
\mathcal{B} [PS(\bar{q}_f q_i) \rightarrow \ell_A^+ \ell_B^-] &= \frac{G_F^4 M_W^4}{32\pi^5} f(x_A^2, x_B^2) M_{PS} f_{PS}^2 (m_{l_A} + m_{l_B})^2 \tau_{PS} \\
&\times \left\{ \left| \frac{M_{PS}^2 (C_P^{q_f q_i} - C_P'^{q_f q_i})}{(m_{q_f} + m_{q_i})(m_{l_A} + m_{l_B})} - (C_A^{q_f q_i} - C_A'^{q_f q_i}) \right|^2 \times [1 - (x_A - x_B)^2] \right. \\
&+ \left. \left| \frac{M_{PS}^2 (C_S^{q_f q_i} - C_S'^{q_f q_i})}{(m_{q_f} + m_{q_i})(m_{l_A} + m_{l_B})} + \frac{(m_{l_A} - m_{l_B})}{(m_{l_A} + m_{l_B})} (C_V^{q_f q_i} - C_V'^{q_f q_i}) \right|^2 \times [1 - (x_A + x_B)^2] \right\}, \tag{113}
\end{aligned}$$

where the function $f(x_i, x_j)$ and the ratio x_i are defined as [149]

$$f(x_i, x_j) = \sqrt{1 - 2(x_i + x_j) + (x_i - x_j)^2}, \quad x_i = \frac{m_{l_i}}{M_{PS}}.$$

1. Wilson coefficients

- *Tree level neutral Higgs contributions to $PS(\bar{q}_f q_i) \rightarrow \ell_A^+ \ell_B^-$ in the 2HDM of type III*

The non-vanishing Wilson coefficients of the operators in Eq. (110) induced through tree-level neutral Higgs ($H_k^0 = H^0, h^0, A^0$) exchange read

$$\begin{aligned}
C_S^{q_f q_i} &= \frac{\pi^2}{2G_F^2 M_W^2} \sum_{k=1}^3 \frac{1}{m_{H_k^0}^2} \left(\Gamma_{\ell_B \ell_A}^{LR H_k^0} + \Gamma_{\ell_B \ell_A}^{RL H_k^0} \right) \Gamma_{q_f q_i}^{RL H_k^0} \\
C_P^{q_f q_i} &= \frac{\pi^2}{2G_F^2 M_W^2} \sum_{k=1}^3 \frac{1}{m_{H_k^0}^2} \left(\Gamma_{\ell_B \ell_A}^{LR H_k^0} - \Gamma_{\ell_B \ell_A}^{RL H_k^0} \right) \Gamma_{q_f q_i}^{RL H_k^0} \\
C_S'^{q_f q_i} &= \frac{\pi^2}{2G_F^2 M_W^2} \sum_{k=1}^3 \frac{1}{m_{H_k^0}^2} \left(\Gamma_{\ell_B \ell_A}^{LR H_k^0} + \Gamma_{\ell_B \ell_A}^{RL H_k^0} \right) \Gamma_{q_f q_i}^{LR H_k^0} \\
C_P'^{q_f q_i} &= \frac{\pi^2}{2G_F^2 M_W^2} \sum_{k=1}^3 \frac{1}{m_{H_k^0}^2} \left(\Gamma_{\ell_B \ell_A}^{LR H_k^0} - \Gamma_{\ell_B \ell_A}^{RL H_k^0} \right) \Gamma_{q_f q_i}^{LR H_k^0}. \tag{114}
\end{aligned}$$

- *Loop-induced charged Higgs contributions to $B_s \rightarrow \mu^+ \mu^-$ in the 2HDM of type II*

As mentioned earlier, we also include in our analysis the 2HDM of type II loop-induced charged Higgs contributions to $B_s \rightarrow \mu^+ \mu^-$ from Ref. [52]:

$$C_S^{bs} = C_P^{bs} = -\frac{m_b V_{tb}^* V_{ts}}{2} \frac{m_\mu}{2M_W^2} \tan^2 \beta \frac{\log(m_H^2/m_t^2)}{m_H^2/m_t^2 - 1}, \tag{115}$$

where m_b and m_t are understood to be running masses evaluated at the matching scale.

H. Flavour-changing lepton decays

The general expressions for the branching ratios of $\tau^- \rightarrow e^- \mu^+ \mu^-$ and $\tau^- \rightarrow \mu^- \mu^+ \mu^-$ have the form

$$\begin{aligned}
\mathcal{B}[\tau^- \rightarrow e^- \mu^+ \mu^-] &= \frac{m_\tau^5}{12(8\pi)^3 \Gamma_\tau} \left(\left| \frac{\Gamma_{\tau e}^{LR H_k^0} \Gamma_{\mu\mu}^{LR H_k^0}}{m_{H_k^0}^2} \right|^2 + \left| \frac{\Gamma_{e\tau}^{LR H_k^0} \Gamma_{\mu\mu}^{LR H_k^0}}{m_{H_k^0}^2} \right|^2 \right. \\
&\quad \left. + \left| \frac{\Gamma_{\tau e}^{LR H_k^0} \Gamma_{\mu\mu}^{LR H_k^0}}{m_{H_k^0}^2} \right|^2 + \left| \frac{\Gamma_{e\tau}^{LR H_k^0} \Gamma_{\mu\mu}^{LR H_k^0}}{m_{H_k^0}^2} \right|^2 \right) \\
\mathcal{B}[\tau^- \rightarrow \mu^- \mu^+ \mu^-] &= \frac{m_\tau^5}{12(8\pi)^3 \Gamma_\tau} \frac{1}{2} \left(2 \left| \frac{\Gamma_{\tau\mu}^{LR H_k^0} \Gamma_{\mu\mu}^{LR H_k^0}}{m_{H_k^0}^2} \right|^2 + 2 \left| \frac{\Gamma_{\mu\tau}^{LR H_k^0} \Gamma_{\mu\mu}^{LR H_k^0}}{m_{H_k^0}^2} \right|^2 \right. \\
&\quad \left. + \left| \frac{\Gamma_{\tau\mu}^{LR H_k^0} \Gamma_{\mu\mu}^{LR H_k^0}}{m_{H_k^0}^2} \right|^2 + \left| \frac{\Gamma_{\mu\tau}^{LR H_k^0} \Gamma_{\mu\mu}^{LR H_k^0}}{m_{H_k^0}^2} \right|^2 \right).
\end{aligned} \tag{116}$$

Note that the (not explicitly denoted) sum over the Higgses must be performed before taking the various absolute values in Eq. (116).

I. Input parameters

In this section we list our input parameters in tabular form.

Parameter	Value (GeV)	Parameter	Value
$\overline{m}_u(2 \text{ GeV})$	0.00219 ± 0.00015 [150]	M_W	80.40 GeV
$\overline{m}_d(2 \text{ GeV})$	0.00467 ± 0.00020 [150]	M_Z	91.19 GeV
$\overline{m}_s(2 \text{ GeV})$	0.095 ± 0.006 [150]	$\alpha_s(M_Z)$	0.119
$\overline{m}_c(m_c)$	1.28 ± 0.04 [151]	G_F	$1.16637 \times 10^{-5} \text{ GeV}^{-2}$
$\overline{m}_b(m_b)$	4.243 ± 0.043 [89]	α_{em}^{-1}	137
$\overline{m}_t(m_t)$	$165.80 \pm 0.54 \pm 0.72$ [14]	v	174.10 GeV

TABLE IX: Left: Input values for the quark masses used in our article. In the numerical analysis, we used the NNLO expressions in α_s for the running (see for example Ref. [152]) in order to obtain the quark-mass values at higher scales. Right: Electroweak parameters and the strong coupling constant used in our analysis. Concerning the running of α_s we used NNLO expressions (given for example in Ref. [72]).

Parameter	Value
f_{B_s}/f_B	$1.221 \pm 0.010 \pm 0.033$ [14]
f_D	218.9 ± 11.3 MeV [138]
f_{D_s}	$249 \pm 2 \pm 5$ MeV [14]
f_{D_s}/f_D	1.188 ± 0.025 [138]
f_K	$156.3 \pm 0.3 \pm 1.9$ MeV [14]
f_K/f_π	1.193 ± 0.005 [150]

Meson masses	Values (GeV)
$m_{B^\pm(B^0)}$	5.279
m_{B_s}	5.367
$m_{D^\pm(D^0)}$	1.870 (1.865)
m_{D_s}	1.969
$m_{K^\pm(K^0)}$	0.494 (0.498)
$m_{\pi^\pm(\pi^0)}$	0.140 (0.135)

TABLE X: Left: Values for decay constants of Ref. [14] obtained by averaging the lattice results of Ref. [138, 153–165]. Right: Meson masses according to the particle data group (see online update of Ref. [72]).

-
- [1] T. Lee, “A Theory of Spontaneous T Violation,” *Phys.Rev.*, vol. D8, pp. 1226–1239, 1973.
- [2] J. F. Gunion, H. E. Haber, G. L. Kane, and S. Dawson, “THE HIGGS HUNTER’S GUIDE,” *Front.Phys.*, vol. 80, pp. 1–448, 2000.
- [3] G. Branco, P. Ferreira, L. Lavoura, M. Rebelo, M. Sher, *et al.*, “Theory and phenomenology of two-Higgs-doublet models,” *Phys.Rept.*, vol. 516, pp. 1–102, 2012, 1106.0034.
- [4] J. E. Kim, “Light Pseudoscalars, Particle Physics and Cosmology,” *Phys.Rept.*, vol. 150, pp. 1–177, 1987.
- [5] R. Peccei and H. R. Quinn, “CP Conservation in the Presence of Instantons,” *Phys.Rev.Lett.*, vol. 38, pp. 1440–1443, 1977.
- [6] M. Trodden, “Electroweak baryogenesis: A Brief review,” 1998, hep-ph/9805252.
- [7] H. E. Haber and G. L. Kane, “The Search for Supersymmetry: Probing Physics Beyond the Standard Model,” *Phys.Rept.*, vol. 117, pp. 75–263, 1985.
- [8] B. Aubert *et al.*, “Evidence for an excess of $B \rightarrow D^{(*)}\tau\nu$ decays,” 2012, 1205.5442.
- [9] J. Lees *et al.*, “Measurement of an Excess of $B \rightarrow D^{(*)}\tau\nu$ Decays and Implications for Charged Higgs Bosons,” 2013, 1303.0571.
- [10] J. Lees *et al.*, “Evidence of $B \rightarrow \tau\nu$ decays with hadronic B tags,” 2012, 1207.0698.
- [11] B. Aubert *et al.*, “A Search for $B^+ \rightarrow \ell^+\nu_\ell$ Recoiling Against $B^- \rightarrow D^0\ell^-\bar{\nu}X$,” *Phys.Rev.*, vol. D81, p. 051101, 2010.
- [12] K. Hara *et al.*, “Evidence for $B^- \rightarrow \tau^-\bar{\nu}$ with a Semileptonic Tagging Method,” *Phys.Rev.*, vol. D82, p. 071101, 2010, 1006.4201.
- [13] I. Adachi *et al.*, “Measurement of $B^- \rightarrow \tau^-\bar{\nu}_\tau$ with a Hadronic Tagging Method Using the Full Data Sample of Belle,” 2012, 1208.4678.
- [14] J. Charles *et al.*, “CP violation and the CKM matrix: Assessing the impact of the asymmetric B factories,” *Eur.Phys.J.*, vol. C41, pp. 1–131, 2005, hep-ph/0406184.
- [15] G. D’Ambrosio, G. F. Giudice, G. Isidori, and A. Strumia, “Minimal flavour violation: An effective field theory approach,” *Nucl. Phys.*, vol. B645, pp. 155–187, 2002, hep-ph/0207036.
- [16] A. Crivellin, C. Greub, and A. Kokulu, “Explaining $B \rightarrow D\tau\nu$, $B \rightarrow D^*\tau\nu$ and $B \rightarrow \tau\nu$ in a 2HDM of type III,” *Phys.Rev.*, vol. D86, p. 054014, 2012, 1206.2634.
- [17] A. J. Buras, M. V. Carlucci, S. Gori, and G. Isidori, “Higgs-mediated FCNCs: Natural Flavour Conservation vs. Minimal Flavour Violation,” *JHEP*, vol. 1010, p. 009, 2010, 1005.5310.
- [18] G. Blankenburg and G. Isidori, “ $B \rightarrow \tau\nu$ in multi-Higgs models with MFV,” 2011, 1107.1216.
- [19] C. Hamzaoui, M. Pospelov, and M. Toharia, “Higgs mediated FCNC in supersymmetric models with large tan Beta,” *Phys.Rev.*, vol. D59, p. 095005, 1999, hep-ph/9807350.
- [20] K. S. Babu and C. F. Kolda, “Higgs mediated $B^0 \rightarrow \mu^+\mu^-$ in minimal supersymmetry,” *Phys. Rev. Lett.*, vol. 84, pp. 228–231, 2000, hep-ph/9909476.

- [21] M. S. Carena, D. Garcia, U. Nierste, and C. E. Wagner, “Effective Lagrangian for the $\bar{t}bH^+$ interaction in the MSSM and charged Higgs phenomenology,” *Nucl.Phys.*, vol. B577, pp. 88–120, 2000, hep-ph/9912516.
- [22] G. Isidori and A. Retico, “Scalar flavor changing neutral currents in the large tan beta limit,” *JHEP*, vol. 0111, p. 001, 2001, hep-ph/0110121.
- [23] A. J. Buras, P. H. Chankowski, J. Rosiek, and L. Slawianowska, “ $\Delta M_{d,s}, B_{d,s}^0 \rightarrow \mu^+\mu^-$ and $B \rightarrow X_s\gamma$ in supersymmetry at large tan β ,” *Nucl. Phys.*, vol. B659, p. 3, 2003, hep-ph/0210145.
- [24] L. Hofer, U. Nierste, and D. Scherer, “Resummation of tan(β)-enhanced supersymmetric loop corrections beyond the decoupling limit,” *JHEP*, vol. 0910, p. 081, 2009, 0907.5408.
- [25] G. Isidori and A. Retico, “ $B_{s,d} \rightarrow \ell^+\ell^-$ and $K_L \rightarrow \ell^+\ell^-$ in SUSY models with nonminimal sources of flavor mixing,” *JHEP*, vol. 0209, p. 063, 2002, hep-ph/0208159.
- [26] A. Crivellin, “Effective Higgs Vertices in the generic MSSM,” *Phys. Rev.*, vol. D83, p. 056001, 2011, 1012.4840.
- [27] D. Noth and M. Spira, “Supersymmetric Higgs Yukawa Couplings to Bottom Quarks at next-to-next-to-leading Order,” *JHEP*, vol. 06, p. 084, 2011, 1001.1935.
- [28] A. Crivellin and C. Greub, “Two-loop SQCD corrections to Higgs-quark-quark couplings in the generic MSSM,” 2012, 1210.7453.
- [29] Y. Okada, M. Yamaguchi, and T. Yanagida, “Upper bound of the lightest Higgs boson mass in the minimal supersymmetric standard model,” *Prog.Theor.Phys.*, vol. 85, pp. 1–6, 1991.
- [30] H. E. Haber and R. Hempfling, “Can the mass of the lightest Higgs boson of the minimal supersymmetric model be larger than m_Z ?,” *Phys.Rev.Lett.*, vol. 66, pp. 1815–1818, 1991.
- [31] J. R. Ellis, G. Ridolfi, and F. Zwirner, “Radiative corrections to the masses of supersymmetric Higgs bosons,” *Phys.Lett.*, vol. B257, pp. 83–91, 1991.
- [32] A. Brignole, “Radiative corrections to the supersymmetric charged Higgs boson mass,” *Phys.Lett.*, vol. B277, pp. 313–323, 1992.
- [33] P. H. Chankowski, S. Pokorski, and J. Rosiek, “Complete on-shell renormalization scheme for the minimal supersymmetric Higgs sector,” *Nucl.Phys.*, vol. B423, pp. 437–496, 1994, hep-ph/9303309.
- [34] A. Dabelstein, “The One loop renormalization of the MSSM Higgs sector and its application to the neutral scalar Higgs masses,” *Z.Phys.*, vol. C67, pp. 495–512, 1995, hep-ph/9409375.
- [35] A. Pilaftsis and C. E. Wagner, “Higgs bosons in the minimal supersymmetric standard model with explicit CP violation,” *Nucl.Phys.*, vol. B553, pp. 3–42, 1999, hep-ph/9902371.
- [36] M. S. Carena, J. R. Ellis, A. Pilaftsis, and C. Wagner, “Renormalization group improved effective potential for the MSSM Higgs sector with explicit CP violation,” *Nucl.Phys.*, vol. B586, pp. 92–140, 2000, hep-ph/0003180.
- [37] A. Freitas, E. Gasser, and U. Haisch, “Supersymmetric large tan(β) corrections to $\Delta M_{d,s}$ and $B_{d,s} \rightarrow \mu^+\mu^-$ revisited,” *Phys.Rev.*, vol. D76, p. 014016, 2007, hep-ph/0702267.

- [38] M. Gorbahn, S. Jager, U. Nierste, and S. Trine, “The supersymmetric Higgs sector and $B-\bar{B}$ mixing for large $\tan\beta$,” *Phys.Rev.*, vol. D84, p. 034030, 2011, 0901.2065.
- [39] A. Crivellin, L. Hofer, and J. Rosiek, “Complete resummation of chirally-enhanced loop-effects in the MSSM with non-minimal sources of flavor-violation,” 2011, 1103.4272.
- [40] J. Rosiek, “Complete set of Feynman rules for the MSSM: Erratum,” 1995, hep-ph/9511250.
- [41] S. L. Glashow and S. Weinberg, “Natural Conservation Laws for Neutral Currents,” *Phys.Rev.*, vol. D15, p. 1958, 1977.
- [42] T. Miki *et al.*, “Effects of charged Higgs boson and QCD corrections in $\bar{B} \rightarrow D\tau\bar{\nu}_\tau$,” pp. 116–124, 2002, hep-ph/0210051.
- [43] A. Wahab El Kaffas, P. Osland, and O. M. Ogreid, “Constraining the Two-Higgs-Doublet-Model parameter space,” *Phys.Rev.*, vol. D76, p. 095001, 2007, 0706.2997.
- [44] O. Deschamps, S. Descotes-Genon, S. Monteil, V. Niess, S. T’Jampens, *et al.*, “The Two Higgs Doublet of Type II facing flavour physics data,” *Phys.Rev.*, vol. D82, p. 073012, 2010, 0907.5135.
- [45] S. Bertolini, F. Borzumati, A. Masiero, and G. Ridolfi, “Effects of supergravity induced electroweak breaking on rare B decays and mixings,” *Nucl.Phys.*, vol. B353, pp. 591–649, 1991.
- [46] M. Ciuchini, G. Degrossi, P. Gambino, and G. Giudice, “Next-to-leading QCD corrections to $B \rightarrow X_s\gamma$: Standard model and two Higgs doublet model,” *Nucl.Phys.*, vol. B527, pp. 21–43, 1998, hep-ph/9710335.
- [47] F. Borzumati and C. Greub, “2HDMs predictions for $\bar{B} \rightarrow X_s\gamma$ in NLO QCD,” *Phys.Rev.*, vol. D58, p. 074004, 1998, hep-ph/9802391.
- [48] M. Misiak, H. Asatrian, K. Bieri, M. Czakon, A. Czarnecki, *et al.*, “Estimate of $\mathcal{B}(\bar{B} \rightarrow X_s\gamma)$ at $O(\alpha_s^2)$,” *Phys.Rev.Lett.*, vol. 98, p. 022002, 2007, hep-ph/0609232.
- [49] T. Hermann, M. Misiak, and M. Steinhauser, “ $\bar{B} \rightarrow X_s\gamma$ in the Two Higgs Doublet Model up to Next-to-Next-to-Leading Order in QCD,” 2012, 1208.2788.
- [50] X. He, T. Nguyen, and R. Volkas, “B MESON RARE DECAYS IN TWO HIGGS DOUBLETS MODELS,” *Phys.Rev.*, vol. D38, p. 814, 1988.
- [51] W. Skiba and J. Kalinowski, “ $B_s \rightarrow \tau^+\tau^-$ decay in a two Higgs doublet model,” *Nucl.Phys.*, vol. B404, pp. 3–19, 1993.
- [52] H. E. Logan and U. Nierste, “ $B_{s,d} \rightarrow \ell^+\ell^-$ in a two Higgs doublet model,” *Nucl.Phys.*, vol. B586, pp. 39–55, 2000, hep-ph/0004139.
- [53] W.-S. Hou, “Enhanced charged Higgs boson effects in $B^- \rightarrow \tau\bar{\nu}, \mu\bar{\nu}$, and $b \rightarrow \tau\bar{\nu}X$,” *Phys.Rev.*, vol. D48, pp. 2342–2344, 1993.
- [54] A. Akeroyd and S. Recksiegel, “The Effect of H^\pm on $B^\pm \rightarrow \tau^\pm\nu_\tau$ and $B^\pm \rightarrow \mu^\pm\nu_\mu$,” *J.Phys.G*, vol. G29, pp. 2311–2317, 2003, hep-ph/0306037.
- [55] S. Fajfer, J. F. Kamenik, and I. Nisandzic, “On the $B \rightarrow D^*\tau\nu_\tau$ Sensitivity to New Physics,” 2012, 1203.2654.

- [56] M. Tanaka, “Charged Higgs effects on exclusive semitauonic B decays,” *Z.Phys.*, vol. C67, pp. 321–326, 1995, hep-ph/9411405.
- [57] U. Nierste, S. Trine, and S. Westhoff, “Charged-Higgs effects in a new $B \rightarrow D\tau\nu$ differential decay distribution,” *Phys.Rev.*, vol. D78, p. 015006, 2008, 0801.4938.
- [58] A. Akeroyd, “Effect of H^\pm on $D_s^+ \rightarrow \mu\nu$ and $D_s^+ \rightarrow \tau\nu$,” *Prog.Theor.Phys.*, vol. 111, pp. 295–299, 2004, hep-ph/0308260.
- [59] A. Akeroyd and C. H. Chen, “Effect of H^\pm on $B \rightarrow \tau\nu_\tau$ and $D_s^+ \rightarrow \tau, \mu\nu$,” *Phys.Rev.*, vol. D75, p. 075004, 2007, hep-ph/0701078.
- [60] A. Akeroyd and F. Mahmoudi, “Constraints on charged Higgs bosons from $D_s^+ \rightarrow \mu\nu$ and $D_s^+ \rightarrow \tau\nu$,” *JHEP*, vol. 0904, p. 121, 2009, 0902.2393.
- [61] M. Antonelli *et al.*, “Precision tests of the Standard Model with leptonic and semileptonic kaon decays,” 2008, 0801.1817.
- [62] M. Baak, M. Goebel, J. Haller, A. Hoecker, D. Ludwig, *et al.*, “Updated Status of the Global Electroweak Fit and Constraints on New Physics,” *Eur.Phys.J.*, vol. C72, p. 2003, 2012, 1107.0975.
- [63] S. Chatrchyan *et al.*, “Search for neutral Higgs bosons decaying to tau pairs in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys.Lett.*, vol. B713, pp. 68–90, 2012, 1202.4083.
- [64] A. Pich and P. Tuzon, “Yukawa Alignment in the Two-Higgs-Doublet Model,” *Phys.Rev.*, vol. D80, p. 091702, 2009, 0908.1554.
- [65] M. Jung, A. Pich, and P. Tuzon, “Charged-Higgs phenomenology in the Aligned two-Higgs-doublet model,” *JHEP*, vol. 1011, p. 003, 2010, 1006.0470.
- [66] F. Mahmoudi and O. Stal, “Flavor constraints on the two-Higgs-doublet model with general Yukawa couplings,” *Phys.Rev.*, vol. D81, p. 035016, 2010, 0907.1791.
- [67] E. Iltan, “Electric dipole moments of charged leptons and lepton flavor violating interactions in the general two Higgs doublet model,” *Phys.Rev.*, vol. D64, p. 013013, 2001, hep-ph/0101017.
- [68] T. Cheng and M. Sher, “Mass Matrix Ansatz and Flavor Nonconservation in Models with Multiple Higgs Doublets,” *Phys.Rev.*, vol. D35, p. 3484, 1987.
- [69] R. Aaij *et al.*, “First evidence for the decay $B_s \rightarrow \mu^+\mu^-$,” 2012, 1211.2674.
- [70] A. J. Buras, J. Girrbach, D. Guadagnoli, and G. Isidori, “On the Standard Model prediction for $BR(B_{s,d} \rightarrow \mu^+\mu^-)$,” 2012, 1208.0934.
- [71] G. Isidori and R. Unterdorfer, “On the short distance constraints from $K_{L,S} \rightarrow \mu^+\mu^-$,” *JHEP*, vol. 0401, p. 009, 2004, hep-ph/0311084.
- [72] J. Beringer *et al.*, “Review of Particle Physics (RPP),” *Phys.Rev.*, vol. D86, p. 010001, 2012.
- [73] T. Inami and C. Lim *Prog. Theor. Phys*, vol. 65, p. 297, 1981.
- [74] M. Ciuchini, E. Franco, V. Lubicz, G. Martinelli, I. Scimemi, *et al.*, “Next-to-leading order QCD corrections to $\Delta F = 2$ effective Hamiltonians,” *Nucl.Phys.*, vol. B523, pp. 501–525, 1998, hep-ph/9711402.

- [75] A. J. Buras, M. Misiak, and J. Urban, “Two loop QCD anomalous dimensions of flavor changing four quark operators within and beyond the standard model,” *Nucl.Phys.*, vol. B586, pp. 397–426, 2000, hep-ph/0005183.
- [76] M. Bona *et al.*, “Model-independent constraints on $\Delta F=2$ operators and the scale of new physics,” *JHEP*, vol. 0803, p. 049, 2008, 0707.0636.
- [77] D. Becirevic, M. Ciuchini, E. Franco, V. Gimenez, G. Martinelli, *et al.*, “ $B_d - \bar{B}_d$ mixing and the $B_d \rightarrow J/\psi K_s$ asymmetry in general SUSY models,” *Nucl.Phys.*, vol. B634, pp. 105–119, 2002, hep-ph/0112303.
- [78] M. Ciuchini, V. Lubicz, L. Conti, A. Vladikas, A. Donini, *et al.*, “Delta M(K) and epsilon(K) in SUSY at the next-to-leading order,” *JHEP*, vol. 9810, p. 008, 1998, hep-ph/9808328.
- [79] V. Lubicz and C. Tarantino, “Flavour physics and Lattice QCD: Averages of lattice inputs for the Unitarity Triangle Analysis,” *Nuovo Cim.*, vol. B123, pp. 674–688, 2008, 0807.4605.
- [80] D. Becirevic, V. Gimenez, G. Martinelli, M. Papinutto, and J. Reyes, “Combined relativistic and static analysis for all Delta B=2 operators,” *Nucl.Phys.Proc.Suppl.*, vol. 106, pp. 385–387, 2002, hep-lat/0110117.
- [81] M. Ciuchini, G. D’Agostini, E. Franco, V. Lubicz, G. Martinelli, *et al.*, “2000 CKM triangle analysis: A Critical review with updated experimental inputs and theoretical parameters,” *JHEP*, vol. 0107, p. 013, 2001, hep-ph/0012308.
- [82] A. Lenz, U. Nierste, J. Charles, S. Descotes-Genon, A. Jantsch, *et al.*, “Anatomy of New Physics in $B - \bar{B}$ mixing,” *Phys.Rev.*, vol. D83, p. 036004, 2011, 1008.1593.
- [83] M. Bona *et al.*, “The UTfit Collaboration Average of D meson mixing data: Spring 2012,” 2012, 1206.6245.
- [84] K. Hayasaka, K. Inami, Y. Miyazaki, K. Arinstein, V. Aulchenko, *et al.*, “Search for Lepton Flavor Violating Tau Decays into Three Leptons with 719 Million Produced $\tau^+\tau^-$ Pairs,” *Phys.Lett.*, vol. B687, pp. 139–143, 2010, 1001.3221.
- [85] U. Bellgardt *et al.*, “Search for the Decay $\mu^+ \rightarrow e^+e^+e^-$,” *Nucl.Phys.*, vol. B299, p. 1, 1988.
- [86] S. Stone, “Planery talk at the international conference on high energy physics (ichep 2012), melbourne, australia,” *Planery talk at the International Conference on High Energy Physics (ICHEP 2012), Melbourne, Australia., July 4-11th, 2012.*
- [87] B. Collaboration, “Precision measurement of the $b \rightarrow x_s \gamma$ photon energy spectrum, branching fraction, and direct cp asymmetry $a_{CP}(b \rightarrow x_{s+d} \gamma)$,” 07 2012, 1207.2690v1,1207.5772.
- [88] J. Lees *et al.*, “Exclusive Measurements of $b \rightarrow s \gamma$ Transition Rate and Photon Energy Spectrum,” 2012, 1207.2520.
- [89] Y. Amhis *et al.*, “Averages of b -hadron, c -hadron, and τ -lepton properties as of early 2012,” 2012, 1207.1158.
- [90] M. Benzke, S. J. Lee, M. Neubert, and G. Paz, “Factorization at Subleading Power and Irreducible Uncertainties in $\bar{B} \rightarrow X_s \gamma$ Decay,” *JHEP*, vol. 1008, p. 099, 2010, 1003.5012.
- [91] T. Hurth and M. Nakao, “Radiative and Electroweak Penguin Decays of B Mesons,”

- Ann.Rev.Nucl.Part.Sci.*, vol. 60, pp. 645–677, 2010, 1005.1224.
- [92] P. del Amo Sanchez *et al.*, “Study of $B \rightarrow X\gamma$ decays and determination of $|V_{td}/V_{ts}|$,” *Phys.Rev.*, vol. D82, p. 051101, 2010, 1005.4087.
- [93] W. Wang, “ $b \rightarrow s\gamma$ and $b \rightarrow d\gamma$ (B factories),” 2011, 1102.1925.
- [94] A. Crivellin and L. Mercolli, “ $B \rightarrow X_d\gamma$ and constraints on new physics,” *Phys.Rev.*, vol. D84, p. 114005, 2011, 1106.5499.
- [95] A. Ali, H. Asatrian, and C. Greub, “Inclusive decay rate for $B \rightarrow X_d\gamma$ in next-to-leading logarithmic order and CP asymmetry in the standard model,” *Phys.Lett.*, vol. B429, pp. 87–98, 1998, hep-ph/9803314.
- [96] T. Hurth, E. Lunghi, and W. Porod, “Untagged $\bar{B} \rightarrow X_{s+d}\gamma$ CP asymmetry as a probe for new physics,” *Nucl.Phys.*, vol. B704, pp. 56–74, 2005, hep-ph/0312260.
- [97] J. Adam *et al.*, “New constraint on the existence of the $\mu^+ \rightarrow e^+\gamma$ decay,” 2013, 1303.0754.
- [98] B. Aubert *et al.*, “Searches for Lepton Flavor Violation in the Decays $\tau^\pm \rightarrow e^\pm\gamma$ and $\tau^\pm \rightarrow \mu^\pm\gamma$,” *Phys.Rev.Lett.*, vol. 104, p. 021802, 2010, 0908.2381.
- [99] K. Hayasaka *et al.*, “New search for $\tau \rightarrow \mu\gamma$ and $\tau \rightarrow e\gamma$ decays at Belle,” *Phys.Lett.*, vol. B666, pp. 16–22, 2008, 0705.0650.
- [100] J. Hudson, D. Kara, I. Smallman, B. Sauer, M. Tarbutt, *et al.*, “Improved measurement of the shape of the electron,” *Nature*, vol. 473, pp. 493–496, 2011.
- [101] G. Bennett *et al.*, “An Improved Limit on the Muon Electric Dipole Moment,” *Phys.Rev.*, vol. D80, p. 052008, 2009, 0811.1207.
- [102] K. Inami *et al.*, “Search for the electric dipole moment of the tau lepton,” *Phys.Lett.*, vol. B551, pp. 16–26, 2003, hep-ex/0210066.
- [103] C. Baker, D. Doyle, P. Geltenbort, K. Green, M. van der Grinten, *et al.*, “An Improved experimental limit on the electric dipole moment of the neutron,” *Phys.Rev.Lett.*, vol. 97, p. 131801, 2006, hep-ex/0602020.
- [104] J. Hisano, M. Nagai, P. Paradisi, and Y. Shimizu, “Waiting for $\mu \rightarrow e\gamma$ from the MEG experiment,” *JHEP*, vol. 0912, p. 030, 2009, 0904.2080.
- [105] M. Passera, “The Standard model prediction of the muon anomalous magnetic moment,” *J.Phys.*, vol. G31, pp. R75–R94, 2005, hep-ph/0411168.
- [106] M. Passera, “Status of the standard model prediction of the muon $g - 2$,” *Nucl.Phys.Proc.Suppl.*, vol. 155, pp. 365–368, 2006, hep-ph/0509372.
- [107] M. Davier, “The Hadronic contribution to $(g-2)_\mu$,” *Nucl.Phys.Proc.Suppl.*, vol. 169, pp. 288–296, 2007, hep-ph/0701163.
- [108] K. Hagiwara, A. Martin, D. Nomura, and T. Teubner, “Improved predictions for $g - 2$ of the muon and $\alpha_{QED}(M_Z^2)$,” *Phys.Lett.*, vol. B649, pp. 173–179, 2007, hep-ph/0611102.
- [109] A. J. Buras, G. Isidori, and P. Paradisi, “EDMs versus CPV in $B_{s,d}$ mixing in two Higgs doublet models with MFV,” *Phys.Lett.*, vol. B694, pp. 402–409, 2011, 1007.5291.
- [110] D. A. Demir, O. Lebedev, K. A. Olive, M. Pospelov, and A. Ritz, “Electric dipole moments

- in the MSSM at large $\tan\beta$,” *Nucl.Phys.*, vol. B680, pp. 339–374, 2004, hep-ph/0311314.
- [111] M. Pospelov and A. Ritz, “Neutron EDM from electric and chromoelectric dipole moments of quarks,” *Phys.Rev.*, vol. D63, p. 073015, 2001, hep-ph/0010037.
- [112] D. A. Demir, M. Pospelov, and A. Ritz, “Hadronic EDMs, the Weinberg operator, and light gluinos,” *Phys.Rev.*, vol. D67, p. 015007, 2003, hep-ph/0208257.
- [113] K. A. Olive, M. Pospelov, A. Ritz, and Y. Santoso, “CP-odd phase correlations and electric dipole moments,” *Phys.Rev.*, vol. D72, p. 075001, 2005, hep-ph/0506106.
- [114] M. Antonelli, V. Cirigliano, G. Isidori, F. Mescia, M. Moulson, *et al.*, “An Evaluation of $|V_{us}|$ and precise tests of the Standard Model from world data on leptonic and semileptonic kaon decays,” *Eur.Phys.J.*, vol. C69, pp. 399–424, 2010, 1005.2323.
- [115] R. Decker and M. Finkemeier, “Radiative corrections to the decay $\tau \rightarrow \pi(K)\nu_\tau$,” *Phys.Lett.*, vol. B316, pp. 403–406, 1993, hep-ph/9307372.
- [116] S. Banerjee, “Lepton Universality, $|V_{us}|$ and search for second class current in tau decays,” 2008, 0811.1429.
- [117] J. F. Kamenik and F. Mescia, “ $B \rightarrow D\tau\nu$ Branching Ratios: Opportunity for Lattice QCD and Hadron Colliders,” *Phys.Rev.*, vol. D78, p. 014003, 2008, 0802.3790.
- [118] J. Korner and G. Schuler, “Exclusive Semileptonic Decays of Bottom Mesons in the Spectator Quark Model,” *Z.Phys.*, vol. C38, p. 511, 1988.
- [119] J. Korner and G. Schuler, “LEPTON MASS EFFECTS IN SEMILEPTONIC B MESON DECAYS,” *Phys.Lett.*, vol. B231, p. 306, 1989.
- [120] J. Korner and G. Schuler, “Exclusive Semileptonic Heavy Meson Decays Including Lepton Mass Effects,” *Z.Phys.*, vol. C46, p. 93, 1990.
- [121] P. Heiliger and L. Sehgal, “SEMILEPTONIC DECAYS OF B MESONS INTO $\tau\nu_\tau$,” *Phys.Lett.*, vol. B229, p. 409, 1989.
- [122] X.-Y. Pham, “New method for determination of the $D \rightarrow \bar{K}^*e\nu$ axial form-factors without resort to angular distributions,” *Phys.Rev.*, vol. D46, p. R1909, 1992.
- [123] S. Fajfer, J. F. Kamenik, I. Nisandzic, and J. Zupan, “Implications of lepton flavor universality violations in B decays,” 2012, 1206.1872.
- [124] N. Deshpande and A. Menon, “Hints of R-parity violation in B decays into $\tau\nu$,” 2012, 1208.4134.
- [125] X.-G. He and G. Valencia, “B decays with τ -leptons in non-universal left-right models,” 2012, 1211.0348.
- [126] A. Celis, M. Jung, X.-Q. Li, and A. Pich, “Sensitivity to charged scalars in $B \rightarrow D^{(*)}\tau\nu_\tau$ and $B \rightarrow \tau\nu_\tau$ decays,” 2012, 1210.8443.
- [127] M. Tanaka and R. Watanabe, “New physics in the weak interaction of $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$,” 2012, 1212.1878.
- [128] J. A. Bailey, A. Bazavov, C. Bernard, C. Bouchard, C. DeTar, *et al.*, “Refining new-physics searches in $B \rightarrow D\tau\nu$ decay with lattice QCD,” *Phys.Rev.Lett.*, vol. 109, p. 071802, 2012,

1206.4992.

- [129] A. Datta, M. Duraisamy, and D. Ghosh, “Diagnosing New Physics in $b \rightarrow c \tau \nu_\tau$ decays in the light of the recent BaBar result,” *Phys.Rev.*, vol. D86, p. 034027, 2012, 1206.3760.
- [130] D. Becirevic, N. Kosnik, and A. Tayduganov, “ $\bar{B} \rightarrow D \tau \bar{\nu}_\tau$ vs. $\bar{B} \rightarrow D \mu \bar{\nu}_\mu$,” *Phys.Lett.*, vol. B716, pp. 208–213, 2012, 1206.4977.
- [131] P. Ko, Y. Omura, and C. Yu, “ $B \rightarrow D^{(*)} \tau \nu$ and $B \rightarrow \tau \nu$ in chiral $U(1)'$ models with flavored multi Higgs doublets,” 2012, 1212.4607.
- [132] P. Biancofiore, P. Colangelo, and F. De Fazio, “On the anomalous enhancement observed in $B \rightarrow D^{(*)} \tau \bar{\nu}_\tau$ decays,” 2013, 1302.1042.
- [133] Y. Sakaki and H. Tanaka, “Constraints of the Charged Scalar Effects Using the Forward-Backward Asymmetry on $B \rightarrow D^{(*)} \tau \bar{\nu}_\tau$,” 2012, 1205.4908.
- [134] B. A. Dobrescu and A. S. Kronfeld, “Accumulating evidence for nonstandard leptonic decays of D_s mesons,” *Phys.Rev.Lett.*, vol. 100, p. 241802, 2008, 0803.0512.
- [135] J. Alexander *et al.*, “Measurement of $B(D_s^+ \rightarrow \ell^+ \nu)$ and the Decay Constant $f_{D_s^+}$ From 600 $/pb^{-1}$ of e^\pm Annihilation Data Near 4170 MeV,” *Phys.Rev.*, vol. D79, p. 052001, 2009, 0901.1216.
- [136] P. Onyisi *et al.*, “Improved Measurement of Absolute Branching Fraction of $D_s^+ \rightarrow \tau \nu_\tau$,” *Phys.Rev.*, vol. D79, p. 052002, 2009, 0901.1147.
- [137] J. N. Simone, “The decay constants f_{D_s} and f_{D^+} from lattice QCD,” *Int.J.Mod.Phys.Conf.Ser.*, vol. 02, pp. 92–96, 2011.
- [138] A. Bazavov *et al.*, “ B - and D -meson decay constants from three-flavor lattice QCD,” *Phys.Rev.*, vol. D85, p. 114506, 2012, 1112.3051.
- [139] V. G. M. J. Barranco, D. Delepine and L. Lopez-Lozano, “Constraining New Physics with D meson decays,” 2013, 1303.3896.
- [140] G. Burdman, J. T. Goldman, and D. Wyler, “Radiative leptonic decays of heavy mesons,” *Phys.Rev.*, vol. D51, pp. 111–117, 1995, hep-ph/9405425.
- [141] A. Masiero and P. Paradisi, “The B - TAU FCNC connection in SUSY Unified Theories,” *J.Phys.Conf.Ser.*, vol. 53, pp. 248–254, 2006, hep-ph/0609262.
- [142] A. Dedes, J. R. Ellis, and M. Raidal, “Higgs mediated $B_{s,d}^0 \rightarrow \mu \tau$, $e \tau$ and $\tau \rightarrow 3 \mu$, $e \mu \mu$ decays in supersymmetric seesaw models,” *Phys.Lett.*, vol. B549, pp. 159–169, 2002, hep-ph/0209207.
- [143] A. Dedes, J. Rosiek, and P. Tanedo, “Complete One-Loop MSSM Predictions for $B^0 \rightarrow \ell^+ \ell'^-$ at the Tevatron and LHC,” *Phys.Rev.*, vol. D79, p. 055006, 2009, 0812.4320.
- [144] D. Boubaa, A. Datta, M. Duraisamy, and S. Khalil, “Predictions for $B \rightarrow \tau \bar{\mu} + \mu \bar{\tau}$,” 2012, 1211.5168.
- [145] T. Aaltonen *et al.*, “Search for the Decays $B_{(s)}^0 \rightarrow e^+ \mu^-$ and $B_{(s)}^0 \rightarrow e^+ e^-$ in CDF Run II,” *Phys.Rev.Lett.*, vol. 102, p. 201801, 2009, 0901.3803.
- [146] B. Aubert *et al.*, “Searches for the decays $B^0 \rightarrow \ell^\pm \tau^\mp$ and $B^+ \rightarrow \ell^+ \nu$ ($\ell = e, \mu$) using

- hadronic tag reconstruction,” *Phys.Rev.*, vol. D77, p. 091104, 2008, 0801.0697.
- [147] G. Passarino and M. Veltman, “One Loop Corrections for $e^+ e^-$ Annihilation Into $\mu^+ \mu^-$ in the Weinberg Model,” *Nucl.Phys.*, vol. B160, p. 151, 1979.
- [148] W. Altmannshofer, A. J. Buras, and D. Guadagnoli, “The MFV limit of the MSSM for low $\tan(\beta)$: Meson mixings revisited,” *JHEP*, vol. 0711, p. 065, 2007, hep-ph/0703200.
- [149] P. H. Chankowski and L. Slawianowska, “ $B_{s,d}^0 \rightarrow \mu^+ \mu^-$ decay in the MSSM,” *Phys.Rev.*, vol. D63, p. 054012, 2001, hep-ph/0008046.
- [150] G. Colangelo, S. Durr, A. Juttner, L. Lellouch, H. Leutwyler, *et al.*, “Review of lattice results concerning low energy particle physics,” *Eur.Phys.J.*, vol. C71, p. 1695, 2011, 1011.4408.
- [151] B. Blossier *et al.*, “Average up/down, strange and charm quark masses with $N_f=2$ twisted mass lattice QCD,” *Phys.Rev.*, vol. D82, p. 114513, 2010, 1010.3659.
- [152] A. J. Buras, “Weak Hamiltonian, CP violation and rare decays,” pp. 281–539, 1998, hep-ph/9806471.
- [153] B. Blossier *et al.*, “Pseudoscalar decay constants of kaon and D-mesons from $N_f = 2$ twisted mass Lattice QCD,” *JHEP*, vol. 0907, p. 043, 2009, 0904.0954.
- [154] C. Bernard, C. E. DeTar, L. Levkova, S. Gottlieb, U. Heller, *et al.*, “Status of the MILC light pseudoscalar meson project,” *PoS*, vol. LAT2007, p. 090, 2007, 0710.1118.
- [155] E. Follana, C. Davies, G. Lepage, and J. Shigemitsu, “High Precision determination of the π , K, D and D(s) decay constants from lattice QCD,” *Phys.Rev.Lett.*, vol. 100, p. 062002, 2008, 0706.1726.
- [156] C. Aubin, J. Laiho, and R. S. Van de Water, “Light pseudoscalar meson masses and decay constants from mixed action lattice QCD,” *PoS*, vol. LATTICE2008, p. 105, 2008, 0810.4328.
- [157] S. Beane, P. Bedaque, K. Orginos, and M. Savage, “ $f(K)/f(\pi)$ in Full QCD with Domain Wall Valence Quarks,” *Phys.Rev.*, vol. D75, p. 094501, 2007, hep-lat/0606023.
- [158] S. Durr, Z. Fodor, C. Hoelbling, S. Katz, S. Krieg, *et al.*, “The ratio $FK/F\pi$ in QCD,” *Phys.Rev.*, vol. D81, p. 054507, 2010, 1001.4692.
- [159] C. Bernard *et al.*, “Lattice calculation of heavy light decay constants with two flavors of dynamical quarks,” *Phys.Rev.*, vol. D66, p. 094501, 2002, hep-lat/0206016.
- [160] A. Bazavov *et al.*, “The D_s and D^+ Leptonic Decay Constants from Lattice QCD,” *PoS*, vol. LAT2009, p. 249, 2009, 0912.5221.
- [161] C. Davies, C. McNeile, E. Follana, G. Lepage, H. Na, *et al.*, “Update: Precision D_s decay constant from full lattice QCD using very fine lattices,” *Phys.Rev.*, vol. D82, p. 114504, 2010, 1008.4018.
- [162] P. Dimopoulos *et al.*, “Lattice QCD determination of m_b , f_B and f_{B_s} with twisted mass Wilson fermions,” *JHEP*, vol. 1201, p. 046, 2012, 1107.1441.
- [163] E. Gamiz, C. T. Davies, G. P. Lepage, J. Shigemitsu, and M. Wingate, “Neutral B Meson Mixing in Unquenched Lattice QCD,” *Phys.Rev.*, vol. D80, p. 014503, 2009, 0902.1815.
- [164] C. McNeile, C. Davies, E. Follana, K. Hornbostel, and G. Lepage, “High-Precision f_{B_s} and

- HQET from Relativistic Lattice QCD,” *Phys.Rev.*, vol. D85, p. 031503, 2012, 1110.4510.
- [165] C. Albertus, Y. Aoki, P. Boyle, N. Christ, T. Dumitrescu, *et al.*, “Neutral B-meson mixing from unquenched lattice QCD with domain-wall light quarks and static b-quarks,” *Phys.Rev.*, vol. D82, p. 014505, 2010, 1001.2023.