



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Top squark searches using dilepton invariant mass distributions and bino-Higgsino dark matter at the LHC

Bhaskar Dutta, Teruki Kamon, Nikolay Kolev, Kuver Sinha, Kechen Wang, and Sean Wu

Phys. Rev. D **87**, 095007 — Published 10 May 2013

DOI: [10.1103/PhysRevD.87.095007](https://doi.org/10.1103/PhysRevD.87.095007)

Top Squark Searches Using Dilepton Invariant Mass Distributions and Bino-Higgsino Dark Matter at the LHC

Bhaskar Dutta¹, Teruki Kamon^{1,2}, Nikolay Kolev³, Kuver Sinha¹, Kechen Wang¹, and Sean Wu¹

¹ *Mitchell Institute for Fundamental Physics and Astronomy, Department of Physics and Astronomy, Texas A&M University, College Station, TX 77843-4242, USA*

² *Department of Physics, Kyungpook National University, Daegu 702-701, South Korea*

³ *Department of Physics, University of Regina, SK, S4S 0A2, Canada*

Pair production of light top squarks at the 8-TeV LHC can be used to probe the gaugino-Higgsino sector of the Minimal Supersymmetric Standard Model. The case where the lightest neutralino is a mixture of Bino and Higgsino, satisfying the thermal dark matter relic density, is investigated. In such a scenario, the lightest top squark decays mostly into (i) a top quark plus the second or third lightest neutralino, and (ii) a bottom quark plus the lightest chargino, instead of a decay scenario of the lightest top squark into a top quark and the lightest neutralino. Final states with ≥ 2 jets, dileptons, and missing energy are expected in a subsequent decay of the second or third lightest neutralinos into the lightest neutralino via an intermediate slepton (“light sleptons” case) or Z boson (“heavy sleptons” case). The opposite-sign same flavor dilepton mass distribution after subtracting the opposite-sign different flavor distribution shows a clear edge in the case of light sleptons. The significance for discovering such a scenario is calculated with optimized cuts in both light and heavy sleptons cases.

I. INTRODUCTION

Results from the 8-TeV LHC (LHC8) have put bounds on the masses of the gluino (\tilde{g}) and first two generation squarks (\tilde{q}) in models of supersymmetry (SUSY). If they have comparable masses, the exclusion limits reach to approximately 1.5 TeV at 95% CL with 13 fb^{-1} of integrated luminosity [1–3]. The third-generation squarks have also actively been searched, putting experimental bounds on parameter space in certain decay modes [4, 5].

Light top squark searches so far have been carried out on the case where the lightest neutralino ($\tilde{\chi}_1^0$) is mainly a Bino and the second lightest neutralino ($\tilde{\chi}_2^0$) mainly a Wino. In such a scenario, the lightest top squark (\tilde{t}_1) decays to $\tilde{\chi}_1^0$ and a top (t) quark at a branching fraction (\mathcal{B}) of nearly 100%. In a recent paper by some of the current authors [7], the trijet invariant mass $M3$ was used to reconstruct top quarks in fully hadronic final state of events with at least four non- b jets, at least two b jets and large missing energy. There have been other approaches to probe this decay by different groups [8]. Reference [9] has studied the \tilde{t}_1 decay in the scenario where $\tilde{\chi}_1^0$ and the lightest chargino ($\tilde{\chi}_1^\pm$) are purely Higgsino.

Due to small electroweak (EW) production the bounds on the neutralinos and charginos are much weaker. This sector, along with the sleptons, plays a crucial role in the dark matter physics of supersymmetric models. In the R -parity conserving Minimal Supersymmetric Standard Model (MSSM), $\tilde{\chi}_1^0$ is typically the dark matter (DM) candidate. If $\tilde{\chi}_1^0$ is purely a Bino, its relic density tends to be large since the annihilation cross-section is smaller than the required thermal annihilation rate 3×10^{-26}

cm^3/sec .¹ One way to obtain the correct relic density is to consider a thermal, well-tempered $\tilde{\chi}_1^0$ which is a mixture of Bino and Higgsino [12, 13], while having $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ as primarily Higgsinos.²

The purpose of this paper is to utilize \tilde{t}_1 decay to probe the dark matter sector in a scenario with $\tilde{\chi}_1^0$ as a Bino-Higgsino mixture and $\tilde{\chi}_{2,3}^0$ as mainly Higgsinos. All three are lighter than the lightest top squark, which is in the sub-TeV range. The main theoretical motivation for considering a light top squark as well as light Higgsinos is Naturalness, while the motivation for the presence of a light Bino is to obtain the correct relic density for $\tilde{\chi}_1^0$, since if a sub-TeV $\tilde{\chi}_1^0$ is purely Higgsino, the relic density is too small [17].

In such a scenario, the lightest top squark mainly decays into $t\tilde{\chi}_{2,3}^0$ and $b\tilde{\chi}_1^\pm$, followed by $\tilde{\chi}_{2,3}^0 \rightarrow Z\tilde{\chi}_1^0$ or $l\tilde{\chi}_1^0$ (via an intermediate slepton state) and $\tilde{\chi}_1^\pm \rightarrow l\nu\tilde{\chi}_1^0$. The final state in $\tilde{t}_1\tilde{t}_1^*$ events has dileptons with jets and missing energy (\cancel{E}_T). The cases of light slepton (whose mass is between $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$) and heavy sleptons are considered in events with at least two leptons, jets and \cancel{E}_T .

The dilepton final states investigated in this paper can lead to a quite robust \tilde{t}_1 search. The cross section for $\tilde{t}_1\tilde{t}_1^*$ production is appreciable at the LHC8 for the mass

¹ The correct relic density for pure Bino can still be obtained by different methods, such as coannihilation [10], slepton exchange for lighter Bino masses, resonance, or non-thermal production without further annihilation [11].

² Other options include non-thermal Winos or Higgsinos [14], multi-component dark matter [15], or well-tempering with Bino and Wino. We note that if $\tilde{\chi}_1^0$ is purely a Wino, the annihilation into W^+W^- final states is in tension with Fermi data for Wino mass below $\sim 250 \text{ GeV}$ [16].

range between 300 and 700 GeV. It is shown that the SUSY combinatoric and SM backgrounds are reduced by performing a opposite-sign same flavor (OSSF) minus opposite-sign different flavor (OSDF) subtraction. If slepton masses are between $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$, an edge in the dilepton mass distribution could be visible due to higher branching fractions of $\tilde{\chi}_{2,3}^0 \rightarrow l\tilde{\chi}_1^0$ decays. The presence of a b -tagged jet in the final state is key to inferring the production of a third-generation squark.

The most dominant background for $\tilde{t}_1\tilde{t}_1^*$ events is $t\bar{t}$ plus n -jets processes. ATLAS has reported bounds on the $m_{\tilde{t}}-m_{\tilde{\chi}_1^0}$ plane for single lepton, dilepton, and fully hadronic final states using 13 fb^{-1} of data at $\sqrt{s} = 8\text{ TeV}$ [4, 5]. Bounds have also been reported for the $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$ decay mode.

It is worthwhile to point out that for probing the dark matter sector with mostly-Bino $\tilde{\chi}_1^0$ and mostly-Wino $\tilde{\chi}_2^0$, vector boson fusion processes are a powerful tool [18]. This is somewhat complementary to the strategy in this paper.

The rest of the paper is structured as follows. In Section II, the results for the “light slepton case” $m_{\tilde{t}_1} > m_{\tilde{\chi}_2^0, \tilde{\chi}_1^\pm} > m_{\tilde{l}} > m_{\tilde{\chi}_1^0}$ are described, while results for the “heavy slepton case” $m_{\tilde{l}} > m_{\tilde{t}_1} > m_{\tilde{\chi}_2^0, \tilde{\chi}_1^\pm} > m_{\tilde{\chi}_1^0}$ are in Section III. The dark matter relic density for the different scenarios are discussed in Section IV, followed by our conclusions in Section V.

II. SCENARIOS WITH LIGHT SLEPTONS

In this section, scenarios which satisfy the following mass relation are studied:

$$m_{\tilde{t}_1} > m_{\tilde{\chi}_3^0}, m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^\pm} > m_{\tilde{l}} > m_{\tilde{\chi}_1^0}. \quad (1)$$

The possible \tilde{t}_1 decay modes are:

$$\tilde{t}_1 \rightarrow t \tilde{\chi}_1^0 \quad (2)$$

$$\tilde{t}_1 \rightarrow t \tilde{\chi}_2^0 \rightarrow t l \bar{\nu} \tilde{l}^{(*)\pm} \rightarrow t l \bar{\nu} l^\pm \tilde{\chi}_1^0, \quad (3)$$

$$\tilde{t}_1 \rightarrow b \tilde{\chi}_1^+ \rightarrow b l \bar{\nu} \tilde{\chi}_1^0 \text{ (or } b q \bar{q}' \tilde{\chi}_1^0) \quad (4)$$

$$\tilde{t}_1 \rightarrow b \tilde{\chi}_2^+ \rightarrow b Z \tilde{\chi}_1^+ \quad (5)$$

Throughout this paper inclusion of charge conjugate modes is implied. The last mode is allowed when the $\tilde{\chi}_2^\pm$ is lighter than \tilde{t}_1 .

It is clear that one obtains an edge in the dilepton invariant mass distribution as well as Z -peak depending on the size of $\mathcal{B}(\tilde{t}_1 \rightarrow b \tilde{\chi}_2^\pm)$ value. A mass spectrum at our benchmark point is displayed in Table I.

The mass spectrum of the model is determined using ISASUGRA [20]. The spectrum is then fed to PYTHIA [21], which generates the Monte Carlo hard scattering events and hadron cascade. These events are passed to the detector simulator PGS4 [22]. We use darkSUSY [23] for relic density computations.

The latest ATLAS exclusion limits on the chargino-neutralino plane are given in [24], in events with $3l + \cancel{E}_T$

in 13.0 fb^{-1} of data at $\sqrt{s} = 8\text{ TeV}$. Exclusion plots for the direct production of charginos, neutralinos, and sleptons are also given in [25], in final states with at least two hadronically decaying τ s and \cancel{E}_T . Our benchmark values of $\tilde{\chi}_1^0, \tilde{\chi}_1^\pm$ are outside the exclusion limits given by ATLAS.

TABLE I: SUSY masses (in GeV) at “light slepton” benchmark point.

Particle Mass (GeV)		\mathcal{B}
\tilde{t}_1	500	17% ($t\tilde{\chi}_2^0$), 22% ($t\tilde{\chi}_3^0$), 8% ($t\tilde{\chi}_1^0$) 53% ($b\tilde{\chi}_1^\pm$)
$\tilde{\chi}_2^0$	175	100% ($l\bar{l}$)
$\tilde{\chi}_3^0$	176	88% ($l\bar{l}$)
$\tilde{\chi}_1^\pm$	164	22% ($l\nu\tilde{\chi}_1^0$)
\tilde{l}	144	100% ($l\tilde{\chi}_1^0$)
$\tilde{\chi}_1^0$	113	

In the benchmark scenario, the mass difference between $\tilde{\chi}_{2,3}^0$ and $\tilde{\chi}_1^0$ is around 63 GeV, and thus an edge in the dilepton invariant mass distribution may be expected around this value. The final state of 2 jets + 2 leptons + \cancel{E}_T events arises mostly from a combination of the $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$ and $\tilde{t}_1 \rightarrow t\tilde{\chi}_{2,3}^0$ decays. If both top squarks decay into a b and a $\tilde{\chi}_1^\pm$, then $2b + 2l + \cancel{E}_T$ events are expected.

For the light slepton case, results will be shown for $m_{\tilde{t}_1} = 390, 440, 500, 550$ and 600 GeV (with masses of $\tilde{\chi}_1^0, \tilde{\chi}_2^0, \tilde{\chi}_3^0$ at the benchmark values in Table I). The heavy slepton case considered in the next section has $m_{\tilde{t}_1} = 390$ GeV, with chargino and neutralinos similar to Table I).

We note that there is no stringent constraint on the \tilde{t}_1 mass from $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$ mode for the benchmark point in Table I) with $m_{\tilde{t}_1} = 500$ GeV. The ATLAS search [5] in final states with two leptons places limits upto a stop mass of ~ 450 GeV in the case where the mass difference between \tilde{t}_1 and $\tilde{\chi}_1^\pm$ is 10 GeV. The ATLAS search [4] in 1 lepton + 4 jets + \cancel{E}_T final state places bounds for \tilde{t}_1 mass less than ~ 390 GeV.

We note, moreover, that these bounds assume $\mathcal{B}(\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm) = 100\%$ in $\tilde{\chi}_1^\pm \rightarrow W^{(*)}\tilde{\chi}_1^0$, whereas for the benchmark scenario considered in Table III in the next Section (“heavy slepton” scenario) with $m_{\tilde{t}_1} = 390$ GeV, the branching fraction to $b\tilde{\chi}_1^\pm$ is $\sim 60\%$. Thus, the benchmark point for the “heavy slepton” scenario too is beyond the ATLAS bounds.

The ATLAS top squark searches [4] in single lepton + ≥ 4 jet final state assume $\mathcal{B}(\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0) = 100\%$. For the top squark masses shown in the paper, the \tilde{t}_1 branching fraction into $t\tilde{\chi}_{2,3}^0$ and $t\tilde{\chi}_1^0$ are $\sim 40\%$ and $\sim 8\%$ respectively. Thus, these benchmark points are not ruled out by the relevant ATLAS searches.

ATLAS top squark searches in the fully hadronic final state [6] also assume $\mathcal{B}(\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0) = 100\%$ and for the

benchmark points in this paper, the exclusion limits are weaker than the limits in the semileptonic final state. Thus, the benchmark points are not ruled out by the fully hadronic searches either.

The analysis begins with selecting events with the following selection cuts:

- (i) At least 2 isolated leptons (e or μ) with $p_T > 20$ and 10 GeV in $|\eta| < 2.5$, where the isolation is defined as $\sum p_T^{\text{track}} < 5$ GeV with $\Delta R = 0.4$;
- (ii) At least 2 jets with $p_T > 30$ GeV in $|\eta| < 2.5$;
- (iii) At least 1 b -tagged jet with $p_T > 30$ GeV in $|\eta| < 2.5$;
- (iv) $\cancel{E}_T > 150$ GeV;
- (v) $H_T > 100$ GeV.

At this stage, the dominant SM background is $t\bar{t}$ events. OSSF dileptons arising from the $\tilde{\chi}_2^0$ decay are kinematically correlated and its dilepton invariant mass distribution is expected to have an edge given by

$$M_{ll}^{\text{edge}} \sim m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}. \quad (6)$$

The OSSF dilepton mass distribution from $t\bar{t}$ events can be modelled by the dilepton distribution of OSDF dilepton events [19]. The OSSF dilepton mass distribution from supersymmetric combinatoric background (*i.e.*, uncorrelated leptonic pairs) can also be modelled by OSDF dilepton mass distribution. This leads to adopting subtracting OSDF distribution from OSSF distribution. The ‘‘light slepton’’ benchmark events would arise in an excess in OSSF–OSDF dilepton mass distribution.

The OSDF dilepton mass distributions for the SUSY benchmark point in Table I along with SM $t\bar{t} + (0-4)$ jets background is shown in shaded histogram in Fig. 1, while its OSSF distribution (blank histogram) is overlaid. A clear edge is seen at around 63 GeV for 30 fb^{-1} luminosity.

Figure 2 shows the flavor subtracted distributions at $\Delta M = m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 53, 63, 70, 77$ and 100 GeV for $\tilde{t}_1 = 500$ GeV, $\tilde{\chi}_1^0 = 113$ GeV and $m_{\tilde{\chi}_3^0} \sim m_{\tilde{\chi}_2^0}$. The dilepton mass distribution edge for all these mass differences except $\Delta M = 100$ GeV can be seen clearly. For $\Delta M = 100$ GeV, the signal acceptance is lower and the dilepton study does not have sensitivity at 30 fb^{-1} luminosity.

Figure 3 shows the flavor subtracted distribution for $t\bar{t} + (0-4)$ jets background plus signal events for $m_{\tilde{t}} = 390, 440, 500, 550$ and 600 GeV, with $m_{\tilde{\chi}_1^0} = 113$ GeV and $m_{\tilde{\chi}_3^0} \sim m_{\tilde{\chi}_2^0} = 175$ GeV. The edge of the dilepton distribution for $m_{\tilde{t}}$ mass upto 550 GeV can be distinguished from the background for 30 fb^{-1} luminosity.

An excess for each top squark mass case in Fig. 3 is evaluated in terms of significances (\mathcal{S}) in Table II. We show significances in cases where at least one of the jets is required to be a b -jet. Here $\mathcal{S} = N_S / \sqrt{N_S + N_B}$, where




FIG. 1: The dilepton invariant mass distributions for $t\bar{t} + (0-4)$ jets background and the benchmark point in Table I are displayed for 30 fb^{-1} luminosity. The unshaded histogram shows the M_{ll}^{OSSF} distribution, while the shaded histogram shows the M_{ll}^{OSDF} distribution, which is fitted with the dot-dashed curve. The solid curve shows the subtracted M_{ll}^{diff} distribution.




FIG. 2: The subtracted dilepton invariant mass distribution M_{ll}^{diff} as $\Delta M = M_{\tilde{\chi}_2^0} - M_{\tilde{\chi}_1^0}$ is varied, for $\tilde{t}_1 = 500$ GeV and $\tilde{\chi}_1^0 = 113$ GeV for 30 fb^{-1} luminosity.




FIG. 3: The subtracted dilepton invariant mass distribution M_{ll}^{diff} as the \tilde{t}_1 mass is varied, all other masses remaining at the benchmark value in Table I for 30 fb^{-1} luminosity.

N_S and N_B are the number of OSSF dilepton events in range of $20 \text{ GeV} < M_{ll} < 70 \text{ GeV}$ for signal (S) and background (B), respectively.

N_B is determined by fitting the entire (SUSY plus $t\bar{t}$) OSDF dilepton distribution to a polynomial function and calculating the number of events in $20 \text{ GeV} < M_{ll} < 70 \text{ GeV}$. N_S is the the number of events in excess above N_B . We find that the significance of the benchmark scenario for $m_{\tilde{t}_1} = 500 \text{ GeV}$ is above 3σ for 30 fb^{-1}

TABLE II: [Light slepton case] Signal and background cross sections in fb for various \tilde{t}_1 masses with $\tilde{\chi}_1^0 = 113 \text{ GeV}$ and $\tilde{\chi}_2^0, \tilde{\chi}_3^0 = 175 \text{ GeV}$. Significances (\mathcal{S}) are given at 30 fb^{-1}

\tilde{t}_1 Mass (GeV)	Signal	Background	\mathcal{S} ($\geq 1 b$)
390	7.08	46.4	5.3
440	6.0	45.6	4.6
500 (benchmark)	3.90	45.1	3.1
550	2.60	44.9	2.1
600	1.70	44.8	1.4

III. SCENARIOS WITH HEAVY SLEPTONS

In this section the case where the sleptons are heavier than the lightest top squark in the following relations:

$$m_{\tilde{t}_1} > m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^\pm} > m_{\tilde{\chi}_1^0} \quad (7)$$

The possible \tilde{t}_1 decay modes are:

$$\tilde{t}_1 \rightarrow t \tilde{\chi}_2^0 \rightarrow t l \tilde{\chi}_1^0 \text{ (or } t q \bar{q} \tilde{\chi}_1^0 \text{)} \quad (8)$$

$$\tilde{t}_1 \rightarrow b \tilde{\chi}_1^\pm \rightarrow b l \bar{\nu} \tilde{\chi}_1^0 \text{ (or } b q \bar{q}' \tilde{\chi}_1^0 \text{)} \quad (9)$$

$$\tilde{t}_1 \rightarrow t \tilde{\chi}_1^0 \quad (10)$$

The leptons and quarks from the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ decays are through off-shell Z and W bosons. SUSY masses at a benchmark point are shown in Table III.

TABLE III: SUSY masses (in GeV) at “heavy slepton” benchmark point.

Particle	Mass	\mathcal{B}
\tilde{t}_1	390	17% ($t\tilde{\chi}_2^0$), 14% ($t\tilde{\chi}_3^0$) 7% ($t\tilde{\chi}_1^0$), 62% ($b\tilde{\chi}_1^\pm$)
$\tilde{\chi}_2^0$	174	7% ($l\tilde{\chi}_1^0$)
$\tilde{\chi}_3^0$	175	7% ($l\tilde{\chi}_1^0$)
$\tilde{\chi}_1^\pm$	164	22% ($l\nu\tilde{\chi}_1^0$)
$\tilde{\chi}_1^0$	112	

The same final state like in the “light slepton” scenario, i.e., 2 jets + 2 leptons + \cancel{E}_T is still the key in this scenario. Therefore the same event selection with re-optimized cuts on the \cancel{E}_T and H_T variables.

- (i) At least 2 isolated leptons with $p_T > 20$ and 10 GeV in $|\eta| < 2.5$;
- (ii) At least 2 jets with $p_T > 30 \text{ GeV}$ in $|\eta| < 2.5$;
- (iii) At least one b -tagged jet with $p_T > 30 \text{ GeV}$ in $|\eta| < 2.5$;
- (iv) $\cancel{E}_T > 190 \text{ GeV}$;
- (v) $H_T > 180 \text{ GeV}$;
- (vi) $20 < M_{ll} < 70 \text{ GeV}$.

In Table IV, we list our signal and background at different stages of cuts and flavor subtraction. The final significance at 30 fb^{-1} is 0.97 if a b jet is required in the event sample. Small value of $\mathcal{B}(Z \rightarrow ll)$ causes smaller significance in the “heavy slepton” case compared to the “light slepton” case.

TABLE IV: [Heavy slepton case] Cross section (fb) for signal and background at different stages of event selection and flavor subtractions are shown for the benchmark point in Table III.

Event Selection	$\tilde{t}_1 \tilde{t}_1^*$	$t\bar{t}$ + jets
$N_l \geq 2, N_j \geq 2, N_b \geq 1,$ $\cancel{E}_T > 190 \text{ GeV}, H_T > 180 \text{ GeV}$	2.1	84.7
OSSF dileptons with $20 < M_{ll}^{\text{OSSF}} < 70 \text{ GeV}$	0.70	13.2
OSDF dileptons with $20 < M_{ll}^{\text{OSDF}} < 70 \text{ GeV}$	0.44	12.8
OSSF – OSDF dilepton with $20 < M_{ll}^{\text{OSSF-OSDF}} < 70 \text{ GeV}$	0.26	0.40

IV. DARK MATTER RELIC DENSITY

For the spectrum (light slepton case) in Table I, $\tilde{\chi}_1^0$ is a mixture of Bino and Higgsino. In this case, one has $m_{\tilde{\chi}_2^0} \sim m_{\tilde{\chi}_3^0}$ and this proximity of masses increases the branching ratio to dileptonic final states. The correct relic density may be obtained by choosing the relative amounts of Bino and Higgsino in $\tilde{\chi}_1^0$ correctly. This choice depends on the mass splitting between $\tilde{\chi}_{2,3}^0$ and $\tilde{\chi}_1^0$ and hence the position of the edge in the dilepton invariant mass distribution. For the benchmark point of Table I, $\tilde{\chi}_1^0$ is 72% Bino and 28% Higgsino, and the relic density is obtained as $\Omega h^2 = 0.11$, as displayed in the first row of Table V. The significance of the analysis is also given.

We note that constraints on DM direct detection for Bino-Higgsino dark matter depends on the sign of μ , as shown for example in [13]. The spectrum considered in this paper may be obtained for either sign of μ , and in

fact, for $\mu < 0$, the DM masses considered in the paper are not constrained by current experiments. We note that for $\mu > 0$, the mixed Bino-Higgsino DM in the mass range considered here is ruled out by XENON100.

If the mass splitting between $\tilde{\chi}_{2,3}^0$ and $\tilde{\chi}_1^0$ is increased, the relic density drops due to smaller Higgsino component in $\tilde{\chi}_1^0$. The correct relic density may be obtained in the light slepton case by coannihilation effects, as displayed in the second row of Table V. Due to a low p_T lepton, the significance for this case is smaller.

For the spectrum (heavy slepton case) in Table III, $\tilde{\chi}_1^0$ is again a mixture of Bino and Higgsino, but due to heavier slepton, the annihilation of $\tilde{\chi}_1^0$ through t -channel slepton exchange is not open. Consequently, a larger Higgsino component in $\tilde{\chi}_1^0$ is required to fulfil the relic density, and hence the mass splitting between $\tilde{\chi}_{2,3}^0$ and $\tilde{\chi}_1^0$ is smaller than the light slepton case. The significance is presented in the third row of Table V.

TABLE V: Significances (\mathcal{S}), relic density (Ωh^2), and Bino-Higgsino composition in three representing SUSY benchmark points for 30 fb^{-1} luminosity. ΔM denotes the mass splitting between $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$. For all points, $m_{\tilde{\chi}_1^0} = 113 \text{ GeV}$.

Masses (GeV)	$\{B, H\}$ (%)	Ωh^2	\mathcal{S}	Comments
$\Delta M = 64$ $m_{\tilde{t}} = 157$ $m_{\tilde{\tau}_1} = 500$	{72, 28}	0.11	3.1	Bino-Higgsino DM (Light slepton, Table I)
$\Delta M = 160$ $m_{\tilde{t}} = 123$ $m_{\tilde{\tau}_1} = 500$	{96, 4}	0.11	0.44	Mainly Bino DM (Coannihilation)
$\Delta M = 62$ $m_{\tilde{t}} = 4000$ $m_{\tilde{\tau}_1} = 390$	{67, 33}	0.11	0.97	Bino-Higgsino DM (Heavy slepton, Table III)

If $\tilde{\chi}_1^0$ is totally a Bino, $\tilde{\chi}_2^0$ a Wino, and the Higgsinos are heavy, the relic density tends to be high unless the sleptons are light, in which case $\tilde{\chi}_1^0$ annihilates through t -channel slepton exchange ($\tilde{\chi}_1^0$ also need to be less than 100 GeV). However, in this case the branching fraction of \tilde{t} to $\tilde{\chi}_2^0$ is quite low, and a very large luminosity is required for this analysis.

Table V is a summary of the significances for three DM scenarios in this study. We see that the signal ≥ 2

jets + 2 leptons + \cancel{E}_T with OSSF–OSDF shows larger significance when there is a slepton between $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$ which is not unnatural in SUSY models..

V. CONCLUSION

In this paper, we show that top squark searches at the LHC8 in dilepton final state could be a powerful probe of the composition of $\tilde{\chi}_1^0$ if $m_{\tilde{t}_1} \lesssim 600 \text{ GeV}$. Motivated by Naturalness and the relic density constraint, the specific case of a well-tempered neutralino that is a mixture of Bino and Higgsino is probed using the decay of top squarks. A typical benchmark point has $m_{\tilde{\chi}_1^0} \sim 113 \text{ GeV}$, $m_{\tilde{\chi}_2^0} \sim m_{\tilde{\chi}_3^0} \sim 175 \text{ GeV}$, and $m_{\tilde{t}} \sim 500 \text{ GeV}$.

In such scenarios, the lightest top squark decays predominantly into $t\tilde{\chi}_{2,3}^0$ and $b\tilde{\chi}_1^\pm$. Both $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ decay to $l\tilde{\chi}_1^0$ (via an intermediate slepton state) or $Z\tilde{\chi}_1^0$, while $\tilde{\chi}_1^\pm$ decays into $l\nu\tilde{\chi}_1^0$ or $W\tilde{\chi}_1^0$. Therefore, the final state has dileptons, jets and missing energy with a clear edge in the OSSF–OSDF dilepton mass distribution. This is different from the most studied scenarios where $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$ are mostly Bino and Wino, respectively, and \tilde{t}_1 decays into $t\tilde{\chi}_1^0$ with branching fraction of 100%.

In “light slepton” scenario where high yield of dilepton events is expected, a discovery sensitivity up to 600 GeV of $m_{\tilde{t}_1}$ with 30 fb^{-1} of integrated luminosity at the LHC8 is expected. If the Higgsino component in $\tilde{\chi}_1^0$ is reduced, then one needs coannihilation processes to satisfy the relic density. In such a case, the p_T of leptons becomes lower and the significance of the study is decreased. In “heavy slepton” case, dileptons are produced from the Z boson decays. Small branching fraction of $\mathcal{B}(Z \rightarrow ll)$ results in decreasing the discovery sensitivity, compared to the “light slepton” case.

VI. ACKNOWLEDGMENTS

This work is supported in part by DOE Grant No. DE-FG02-95ER40917 and by the World Class University (WCU) project through the National Research Foundation (NRF) of Korea funded by the Ministry of Education, Science, and Technology (Grant No. R32-2008-000-20001-0).

-
- [1] ATLAS Collaboration, “Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7 fb^{-1} of $\sqrt{s} = 7 \text{ TeV}$ proton-proton collision data,” arXiv:1208.0949 [hep-ex].
- [2] ATLAS Collaboration, “Hunt for new phenomena using large jet multiplicities and missing transverse momentum

with ATLAS in 4.7 fb^{-1} of $\sqrt{s} = 7 \text{ TeV}$ proton-proton collisions,” JHEP **07**, 167 (2012) [arXiv:1206.1760 [hep-ex]].

- [3] CMS Collaboration, “Search for new physics in the multijet and missing transverse momentum final state in proton-proton collisions at $\sqrt{s} = 7 \text{ TeV}$,” arXiv:1207.1898 [hep-ex].

- [4] ATLAS Collaboration, “Search for direct top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions using 13 fb^{-1} of ATLAS data,” ATLAS-CONF-2012-166.
- [5] ATLAS Collaboration, “Search for the charged current decay of a supersymmetric top-quark partner in final states with two leptons in $\sqrt{s} = 8$ TeV pp collisions using 13 fb^{-1} of ATLAS data,” ATLAS-CONF-2012-167.
- [6] ATLAS Collaboration, “Search for direct production of the top squark in the all-hadronic $t\bar{t} + \cancel{E}_T$ final state in 21 fb^{-1} of pp collisions $\sqrt{s} = 8$ TeV with the ATLAS detector,” ATLAS-CONF-2013-024.
- [7] B. Dutta, T. Kamon, N. Kolev, K. Sinha and K. Wang, “Searching for Top Squarks at the LHC in Fully Hadronic Final State,” arXiv:1207.1873 [hep-ph].
- [8] T. Plehn and M. Spannowsky, “Top Tagging,” arXiv:1112.4441 [hep-ph]. T. Plehn, M. Spannowsky and M. Takeuchi, “Stop searches in 2012,” arXiv:1205.2696 [hep-ph]. D. E. Kaplan, K. Rehermann and D. Stolarski, “Searching for Direct Stop Production in Hadronic Top Data at the LHC,” arXiv:1205.5816 [hep-ph]. D. S. M. Alves, M. R. Buckley, P. J. Fox, J. D. Lykken and C. -T. Yu, “Stops and MET: the shape of things to come,” arXiv:1205.5805 [hep-ph]. T. Han, R. Mahbubani, D. G. E. Walker and L. -T. Wang, “Top Quark Pair plus Large Missing Energy at the LHC,” JHEP **05**, 117 (2009) [arXiv:0803.3820 [hep-ph]]. Y. Bai, H. -C. Cheng, J. Gallicchio and J. Gu, “Stop the Top Background of the Stop Search,” JHEP **07**, 110 (2012) [arXiv:1203.4813 [hep-ph]]. D. S. M. Alves, M. R. Buckley, P. J. Fox, J. D. Lykken and C. -T. Yu, “Stops and MET: The Shape of Things to Come,” arXiv:1205.5805 [hep-ph]. C. Kilic and B. Tweedie, “Cornering Light Stops with Dileptonic mT_2 ,” arXiv:1211.6106 [hep-ph].
- [9] M. L. Graesser and J. Shelton, “Hunting Asymmetric Stops,” arXiv:1212.4495 [hep-ph].
- [10] K. Griest and D. Seckel, “Three exceptions in the calculation of relic abundances,” Phys. Rev. D **43**, 3191 (1991). R. L. Arnowitt, B. Dutta and Y. Santoso, “Coannihilation effects in supergravity and D-brane models,” Nucl. Phys. B **606**, 59 (2001) [hep-ph/0102181].
- [11] R. Allahverdi, B. Dutta and K. Sinha, “Successful Supersymmetric Dark Matter with Thermal Over/Under-Abundance from Late Decay of a Visible Sector Scalar,” arXiv:1212.6948 [hep-ph].
- [12] N. Arkani-Hamed, A. Delgado and G. F. Giudice, “The Well-tempered neutralino,” Nucl. Phys. B **741**, 108 (2006) [hep-ph/0601041].
- [13] C. Cheung, L. J. Hall, D. Pinner and J. T. Ruderman, “Prospects and Blind Spots for Neutralino Dark Matter,” arXiv:1211.4873 [hep-ph]. P. Grothaus, M. Lindner and Y. Takanishi, “Naturalness of Neutralino Dark Matter,” arXiv:1207.4434 [hep-ph]. M. Perelstein and B. Shakya, “XENON100 Implications for Naturalness in the MSSM, NMSSM and lambda-da-SUSY,” arXiv:1208.0833 [hep-ph]. M. Farina, M. Kadastik, D. Pappadopulo, J. Pata, M. Raidal and A. Strumia, “Implications of XENON100 and LHC results for Dark Matter models,” Nucl. Phys. B **853**, 607 (2011) [arXiv:1104.3572 [hep-ph]].
- [14] B. Dutta, L. Leblond and K. Sinha, “Mirage in the Sky: Non-thermal Dark Matter, Gravitino Problem, and Cosmic Ray Anomalies,” Phys. Rev. D **80**, 035014 (2009) [arXiv:0904.3773 [hep-ph]]. B. S. Acharya, G. Kane, S. Watson and P. Kumar, “A Non-thermal WIMP Miracle,” Phys. Rev. D **80**, 083529 (2009) [arXiv:0908.2430 [astro-ph.CO]].
- [15] H. Baer, V. Barger, P. Huang, D. Mickelson, A. Mustafayev and X. Tata, “Radiative natural supersymmetry: Reconciling electroweak fine-tuning and the Higgs boson mass,” arXiv:1212.2655 [hep-ph].
- [16] A. Geringer-Sameth and S. M. Koushiappas, “Exclusion of canonical WIMPs by the joint analysis of Milky Way dwarfs with Fermi,” Phys. Rev. Lett. **107**, 241303 (2011) [arXiv:1108.2914 [astro-ph.CO]].
- [17] R. Allahverdi, B. Dutta and K. Sinha, “Non-thermal Higgsino Dark Matter: Cosmological Motivations and Implications for a 125 GeV Higgs,” Phys. Rev. D **86**, 095016 (2012) [arXiv:1208.0115 [hep-ph]]. I. Gogoladze, F. Nasir and Q. Shafi, “Non-Universal Gaugino Masses and Natural Supersymmetry,” arXiv:1212.2593 [hep-ph].
- [18] B. Dutta, A. Gurrola, W. Johns, T. Kamon, P. Sheldon and K. Sinha, “Vector Boson Fusion Processes as a Probe of Supersymmetric Electroweak Sectors at the LHC,” arXiv:1210.0964 [hep-ph].
- [19] CMS Collaboration, “Search for new physics in events with opposite-sign leptons, jets, and missing transverse energy in pp collisions at $\sqrt{s} = 7$ TeV,” Phys. Lett. B **718**, 815 (2013) [arXiv:1206.3949 [hep-ex]].
- [20] F. E. Paige, S. D. Protopopescu, H. Baer and X. Tata, “ISAJET 7.69: A Monte Carlo event generator for p p, anti-p p, and e+ e- reactions,” [hep-ph/0312045]. We use ISAJET version 7.74.
- [21] T. Sjostrand, S. Mrenna, and P. Skands, “PYTHIA 6.4 Physics and Manual.” J. High Energy Phys. **05** (2006) 026. We use PYTHIA version 6.411 with TAUOLA.
- [22] PGS4 is a parameterized detector simulator. We use version 4 (<http://www.physics.ucdavis.edu/~conway/research/software/pgs/pgs4-general.htm>) in the CMS detector configuration.
- [23] P. Gondolo, J. Edsjo, P. Ullio, L. Bergstrom, M. Schelke, E. A. Baltz, “Darksusy - a numerical package for supersymmetric dark matter calculations,” [astro-ph/0211238].
- [24] ATLAS Collaboration, “Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in 13 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector,” ATLAS-CONF-2012-154.
- [25] ATLAS Collaboration, “Search for electroweak production of supersymmetric particles in final states with at least two hadronically decaying taus and missing transverse momentum with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV,” ATLAS-CONF-2013-028.