



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Automatic computations at next-to-leading order in QCD for top-quark flavor-changing neutral processes

Celine Degrande, Fabio Maltoni, Jian Wang, and Cen Zhang

Phys. Rev. D **91**, 034024 — Published 23 February 2015

DOI: [10.1103/PhysRevD.91.034024](https://doi.org/10.1103/PhysRevD.91.034024)

Automatic Computations at NLO in QCD for Top-Quark Flavor-Changing Neutral Processes

Celine Degrande,¹ Fabio Maltoni,² Jian Wang,³ and Cen Zhang⁴

¹*Institute for Particle Physics Phenomenology, Department of Physics
Durham University, Durham DH1 3LE, United Kingdom*

²*Centre for Cosmology, Particle Physics and Phenomenology,
Université Catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium*

³*PRISMA Cluster of Excellence & Mainz Institute for Theoretical Physics,
Johannes Gutenberg University, D-55099 Mainz, Germany*

⁴*Department of Physics, Brookhaven National Laboratory, Upton, N.Y., 11973, U.S.A.*

Computations at next-to-leading order in the Standard Model offer new technical challenges in presence of higher dimensional operators. We introduce a framework that, starting from the top-quark effective field theory at dimension six, allows to make predictions for cross sections as well as distributions in a fully automatic way. As an application, we present the first complete results at NLO in QCD for flavor changing neutral interactions including parton shower effects, for tZ , th , $t\gamma$ associated production at the LHC.

PACS numbers: 14.65.Ha, 12.38.Bx

Introduction. The millions of top quarks already produced at the LHC together with the tens of millions expected in the coming years will provide a unique opportunity to search for interactions beyond the Standard Model (SM). Among them Flavor-Changing Neutral (FCN) interactions are of special interest. In the SM, FCN interactions can be generated at one loop, yet they turn out to be suppressed by the Glashow-Iliopoulos-Maiani mechanism [1]. The resulting FCN decay modes of the top quark have branching ratios of order $10^{-12} - 10^{-15}$ [2–4]. Thus, any signal for top-quark FCN processes at a measurable rate would immediately indicate new physics in the top-quark sector. These processes have been searched for already at different colliders, including LEP2, HERA, Tevatron and more recently at the LHC [5]. So far no signal has been observed and limits have been set on the coupling strengths.

The most important top-quark FCN processes at the LHC include decay processes such as $t \rightarrow qB$ and production processes such as $qg \rightarrow t$ and $qg \rightarrow tB$, where q is a u or c quark and B is a neutral boson, *i.e.*, $B = \gamma, Z, h$. In general, the decay processes are equally sensitive to utB and ctB couplings, while the production modes are less sensitive to ctB , but may provide a better handle on a certain class of utB couplings [6]. In addition, compared to decay modes, single-top production can provide more information. First, it makes a wider range of kinematic variables accessible, helping in the discrimination of the light quark flavors involved and the structure of the qtB couplings. Second, it probes interactions at higher scales where new physics effects could be enhanced. In general, being somewhat complementary, both decay and production processes are used for setting the most stringent constraints.

Leading order (LO) predictions for the production processes suffer from large uncertainties due to missing higher order corrections. To curb such uncertainties,

next-to-leading order (NLO) predictions in QCD for this class of processes have started to be calculated in recent years [7–11], providing a much better, yet incomplete, picture of their relevance. In general, corrections are found to be large, of order 30% to 80% and to lead to considerable reductions of the residual theoretical uncertainties. Both aspects are important in bounding and possibly extracting top-quark FCN couplings at the LHC.

In this paper we present the first automatic computations for top-quark FCN production processes, $qg \rightarrow tB$ with $B = \gamma, Z, h$, at NLO in QCD, by implementing all flavor-changing dimension-six fully gauge-invariant operators in FEYNRULES [12] and then passing this information into the MADGRAPH5_AMC@NLO framework [13]. Compared to previous works [9–11, 14–17], the salient features of our results can be summarized as follows. Our study is the first where all relevant dimension-six operators for this class of processes (associated production with a boson) are consistently taken into account. In particular the vector-current like tqZ coupling in $ug \rightarrow tZ$, and the tug and $tugh$ couplings in $ug \rightarrow th$ are included here for the first time. Second, our results are obtained via a fully automatic chain of tools that allows to go directly from the Lagrangian to the hard events by performing its renormalization at one loop, and then passing the corresponding Feynman rules to the MADGRAPH5_AMC@NLO to generate all the elements necessary for a computation at NLO in QCD. In particular, other processes triggered by the same set of operators can be seamlessly computed within the same framework. Third, event generation is also automatically available at NLO accuracy, by matching it to parton shower (PS) via the MC@NLO formalism [18] so that results can be employed in realistic experimental simulations. Finally, another important aspect of this work is that it provides a proof-of-principle that fully automatic computation of cross sections at NLO in QCD is possible in the context

of the full dimension-six Lagrangian of the SM. Higher order computations in effective field theories, which are renormalizable only order by order in $1/\Lambda$, Λ being the scale of new physics, present novel technical challenges. In general, UV divergences generated by one operator at a certain order of $1/\Lambda$ have to be absorbed also by other effective operators. As a result, the full set of relevant operators together with their operator mixing effects need to be considered simultaneously, and appropriate UV counterterms have to be implemented in the calculation. Our method and its implementation are fully general and can cover arbitrary NLO calculations in the complete dimension-six Lagrangian of the SM.

Framework. The FCN couplings of the top quark can be parameterized using either fully gauge-symmetric dimension-six operators [19, 20] or dimension-four and five operators in the electroweak broken phase [6, 21]. The latter approach has some intrinsic limitations [22], and we will use the dimension-six operators throughout the paper. The effective Lagrangian can be written as

$$\mathcal{L}_{\text{EFT}} = \mathcal{L}_{\text{SM}} + \sum_i \frac{C_i}{\Lambda^2} O_i + h.c. \quad (1)$$

In this work we consider qtB couplings at the dimension-six level. The relevant operators must involve one top quark and one light quark. They are

$$\begin{aligned} O_{\varphi q}^{(3,i+3)} &= i \left(\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{q}_i \gamma^\mu \tau^I Q) \\ O_{\varphi q}^{(1,i+3)} &= i \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{q}_i \gamma^\mu Q) \\ O_{\varphi u}^{(i+3)} &= i \left(\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \right) (\bar{u}_i \gamma^\mu t) \\ O_{uB}^{(i3)} &= g_Y (\bar{q}_i \sigma^{\mu\nu} t) \tilde{\varphi} B_{\mu\nu}, \quad O_{uW}^{(i3)} = g_W (\bar{q}_i \sigma^{\mu\nu} \tau^I t) \tilde{\varphi} W_{\mu\nu}^I \\ O_{uG}^{(i3)} &= g_s (\bar{q}_i \sigma^{\mu\nu} T^A t) \tilde{\varphi} G_{\mu\nu}^A, \quad O_{u\varphi}^{(i3)} = (\varphi^\dagger \varphi) (\bar{q}_i t) \tilde{\varphi}, \end{aligned}$$

where the operator notation is consistent with Ref. [23], with additional flavor indices. On the right hand side, the subscript $i = 1, 2$ represents the generation of the light quark fields. u_i and q_i are single and doublet quark fields of the first two generation, respectively, while t and Q are of the third-generation. φ is the Higgs doublet. A diagonal CKM matrix is assumed. The group generators are normalized such that $\text{Tr}(T^A T^B) = \delta^{AB}/2$ and $\text{Tr}(\tau^I \tau^J) = 2\delta^{IJ}$, and $\varphi^\dagger \overleftrightarrow{D}_\mu \varphi \equiv \varphi^\dagger D_\mu \varphi - D_\mu \varphi^\dagger \varphi$, $\varphi^\dagger \overleftrightarrow{D}_\mu^I \varphi \equiv \varphi^\dagger \tau^I D_\mu \varphi - D_\mu \varphi^\dagger \tau^I \varphi$. For operators with $(i3)$ superscript, a similar set of operators with $(3i)$ flavor structure can be obtained by interchanging $(i3) \leftrightarrow (3i)$, $t \leftrightarrow u_i$ and $Q \leftrightarrow q_i$. The first three operators give rise to V/A couplings of Z to a flavor-changing current, which were not considered in previous calculations of Ref. [10]. The $O_{uB}^{(i3,3i)}$, $O_{uW}^{(i3,3i)}$ and $O_{uG}^{(i3,3i)}$ operators correspond to weak- and color-dipole couplings. In particular, $O_{uG}^{(i3,3i)}$ could induce the production $pp \rightarrow th$, and it was not included in [11]. The last operator gives rise to flavor-changing Yukawa couplings. This operator is actually implemented as $O_{u\varphi}^{(i3)} = (\varphi^\dagger \varphi - v^2/2) (\bar{q}_i t) \tilde{\varphi}$ to

avoid any need for a field redefinition in order to remove the tree-level $q-t$ mixing. It is interesting to note that all qtB interactions receive contributions from operators that involve the Higgs field, therefore they are also relevant for constraining new physics in the Higgs sector. Finally, we stress that four-fermion operators should also be taken into account for a complete phenomenological study of FCN interactions, see Ref. [22]. Their implementation in the current framework is possible and left for future work.

In calculations at NLO in QCD, a renormalization scheme needs to be specified, in particular for the dimension-six operators. We adopt the $\overline{\text{MS}}$ scheme in general, except for masses and wave functions that are renormalized on shell. Specifically, this requires the introduction of off-diagonal wave function counterterms to cancel the $u-t$ or $c-t$ two-point functions generated by $O_{uG}^{(i3,3i)}$. We work in the five-flavour scheme where the b -quark mass is neglected, and we subtract the massless modes according to the $\overline{\text{MS}}$ scheme and the top at zero momentum for the strong coupling constant renormalization [24]. At order α_S these operators will not mix with the SM terms, but mix among themselves. The running of these coefficients is given by the renormalization group equations

$$\frac{dC_i(\mu)}{d\ln\mu} = \gamma_{ij} C_j(\mu), \quad (2)$$

where γ_{ij} for $C_{uG}^{(13)}$, $C_{uW}^{(13)}$, $C_{uB}^{(13)}$ and $C_{u\varphi}^{(13)}$ can be written as a matrix [17, 25]:

$$\gamma = \frac{\alpha_S}{\pi} \begin{pmatrix} \frac{1}{3} & 0 & 0 & 0 \\ \frac{2}{3} & \frac{2}{3} & 0 & 0 \\ \frac{10}{9} & 0 & \frac{2}{3} & 0 \\ 4y_t^2 & 0 & 0 & -2 \end{pmatrix}, \quad (3)$$

where y_t is the top-quark Yukawa coupling. The same γ_{ij} matrix applies for the operators with either $(i3)$ or $(3i)$ superscript. The operators $O_{\varphi q}^{(3,i+3)}$, $O_{\varphi q}^{(1,i+3)}$ and $O_{\varphi u}^{(i+3)}$ do not have any anomalous dimension due to current conservation and do not mix with other operators.

Implementation and checks. The operators are implemented in the UFO format [26], using the FEYNRULES package [12]. The evaluation of the loop corrections in MADGRAPH5_AMC@NLO requires two additional elements, the UV counterterms and the rational R2 terms which are required by the OPP technique [27]. These are computed fully automatically by the NLOCT [28] package, which has been extended to handle EFT's *i.e.*, to compute the R2 and UV divergent parts of amplitudes with integrals of arbitrary high ranks. Currently, such calculations are limited to operators with up to two fermion fields. The determination of the UV divergent part of the counterterms is obtained by simply changing the sign of the UV divergent part of the corresponding amplitude. This avoids the translations of the counterterms vertices in the operator renormalization constants




FIG. 1. Tree-level diagrams for $pp \rightarrow tV$ and $pp \rightarrow th$. The black dots represent contributions from color dipole operators $O_{uG}^{(i3,3i)}$, while the shaded squares represent other operators.

and the associated basis reduction. However, it is only valid when the dimension-six operators are renormalized in the $\overline{\text{MS}}$ scheme.

We have extensively checked our implementation by evaluating the virtual contributions of $ug \rightarrow t$, $u\gamma \rightarrow t$, $uZ \rightarrow t$, $uh \rightarrow t$ and $ug \rightarrow th$ (with uth coupling only) and comparing them with corresponding known analytical expressions numerically. In each case the results agree. In addition we have checked the gauge-invariance of all virtual contributions, as well as the pole cancellation when combining virtual and real contributions. When possible, we have also made comparisons with the results for total cross sections for $pp \rightarrow t\gamma$, tZ , th at fixed order of Refs. [9–11], finding consistent results.

Calculation. As an application of our general framework to the phenomenology of the top quark FCN at the LHC, we consider three processes, $pp \rightarrow t\gamma$, $pp \rightarrow tZ$ and $pp \rightarrow th$. The LO diagrams are shown in Fig. 1. Each process receives contributions from two different interactions, one from utg coupling and the other from utB coupling. At NLO in QCD the utg operator will mix with other operators, and as a result a NLO calculation needs to be carried out with the full set of operators.

Our numerical results are obtained by employing the following input parameters

$$\begin{aligned} m_Z &= 91.1876 \text{ GeV}, & \alpha &= 1/127.9, \\ G_F &= 1.166370 \times 10^{-5} \text{ GeV}^{-2}, \\ m_t &= 172.5 \text{ GeV}, & m_h &= 125 \text{ GeV}, & \Lambda &= 1 \text{ TeV}. \end{aligned} \quad (4)$$

We use CTEQ6M for NLO and CTEQ6L for LO calculations respectively, with their respective values of α_S [29]. The renormalization scale μ_r and factorization scale μ_f are chosen to be $m_t + m_B$ for $pp \rightarrow tB$ process, and are allowed to vary independently by a factor of 0.5 to 2. In $pp \rightarrow t\gamma$, we require the photon $p_T > 50$ GeV and its pseudorapidity $|\eta| < 2.5$. For the photon, we employ the isolation criterium of Ref. [30] with a radius of 0.4. The events are then showered with PYTHIA6 [31] or HERWIG6 [32]. Finally, we have checked that the doubly resonant diagrams with the anti-top decaying through FCN interactions have a small impact, yet they have been removed from the real contributions, see Ref. [33].

Currently the best limits on top FCN couplings are from the decay searches of $t \rightarrow qZ$ [34], $t \rightarrow qh$ [35, 36],

	LO		NLO	
coefficient	σ [fb]	scale uncertainty	σ [fb]	scale uncertainty
$C_{uB}^{(13)}$ = 1.0	546	+14.4% -11.8%	764	+6.9% -6.4%
$C_{uG}^{(13)}$ = 0.04	1.00	+12.0% -10.2%	2.34	+15.2% -11.5%
$C_{uG}^{(13)}$, veto	0.739	+11.50% -9.8%	1.19	+7.7% -6.5%
$C_{uB}^{(23)}$ = 1.9	152	+10.6% -9.6%	258	+6.8% -6.0%
$C_{uG}^{(23)}$ = 0.09	0.590	+12.1% -11.1%	1.95	+16.4% -12.3%
$C_{uG}^{(23)}$, veto	0.457	+12.2% -11.2%	1.04	+10.3% -8.9%

TABLE I. Total cross sections for $pp \rightarrow t\gamma$. Contributions from operators with (31), (32) superscripts are not displayed, but they are the same as their (13), (23) counterparts. Contributions from $O_{uW}^{(i3),(3i)}$ are equal to those from $O_{uB}^{(i3),(3i)}$.

and the production searches of $qg \rightarrow t$ [37] and $qg \rightarrow t\gamma$ [38]. To make a viable choice for the operator coefficients in our calculation, we exploit the results of Ref. [22] that are based on a global fit on the full set of current limits:

coefficient	limit	coefficient	limit	relevant production
$C_{\varphi q}^{(j,i+3)}$	1.05	$C_{\varphi u}^{(i+3)}$	1.05	tZ
$C_{uG}^{(13,31)}$	0.041	$C_{uG}^{(23,32)}$	0.093	$t\gamma, tZ, th$
$C_{uW}^{(13,31)}$	0.92	$C_{uW}^{(23,32)}$	1.1	$t\gamma, tZ$
$C_{uB}^{(13,31)}$	1.0	$C_{uB}^{(23,32)}$	1.9	$t\gamma, tZ$
$C_{u\varphi}^{(13,31)}$	3.5	$C_{u\varphi}^{(23,32)}$	3.5	th

where $i = 1, 2$, $j = 1, 3$, and the limits apply to the moduli of the coefficients, assuming $\Lambda = 1$ TeV. Each limit is obtained by marginalizing over all the other operator coefficients. In this work, we choose real and positive values for the coefficients that do not exceed these bounds. The total cross sections at the LHC at $\sqrt{s} = 8$ TeV corresponding to each operator are displayed in Tables I, II and III. The scale uncertainties are also displayed. As expected the K -factors are generally sizeable and the scale uncertainties are significantly reduced at NLO. This is the case for all operators except for $O_{uG}^{(i3,3i)}$ in $t\gamma$ production. This process has an unusually large K -factor when the flavor-changing coupling is coming from $O_{uG}^{(i3,3i)}$. As shown in Table I, vetoing any extra jet with $p_T > 50$ GeV reduces the K -factor from 2.3 (3.3) to 1.6 (2.3) for utg (ctg) coupling as well as the uncertainties for this production mechanism. Note also that a jet veto can help to improve the signal over the SM background ratio, for all three processes.

Differential cross sections. The p_T distributions of the top quark in $pp \rightarrow t\gamma$ and $pp \rightarrow th$ are shown in Fig. 2. Both LO and NLO signals are displayed, together with the SM backgrounds from $pp \rightarrow t\gamma j$ and thj , which are generated at NLO with same parameters. In all cases the $O_{uG}^{(13)}$ contributions are very small due to the stringent limit from $ug \rightarrow t$ production. Therefore, the $pp \rightarrow tX$ processes appear more as confirmation than as discovery channel for the chromomagnetic operator.

To illustrate the importance of keeping all operators

coefficient	LO		NLO	
	σ [fb]	scale uncertainty	σ [fb]	scale uncertainty
$C_{\varphi u}^{(1+3)} = 1.0$	905	+12.9% -10.9%	1163	+6.2% -5.6%
$C_{uW}^{(13)} = 0.9$	1737	+11.5% -9.8%	2270	+6.6% -6.2%
$C_{uG}^{(13)} = 0.04$	30.1	+17.5% -13.8%	36.0	+3.8% -5.2%
$C_{uG}^{(31)} = 0.04$	29.4	+17.7% -13.9%	34.9	+3.4% -5.1%
$C_{\varphi u}^{(2+3)} = 1.0$	73.2	+10.4% -9.3%	107	+6.5% -5.9%
$C_{uW}^{(23)} = 1.1$	172	+7.5% -7.2%	255	+6.1% -5.2%
$C_{uG}^{(23)} = 0.09$	6.92	+11.3% -9.9%	10.6	+5.8% -5.4%
$C_{uG}^{(32)} = 0.09$	6.58	+11.5% -10.1%	10.0	+5.7% -5.3%

TABLE II. Total cross sections for $pp \rightarrow tZ$. Contributions from operators $O_{uW}^{(31),(32)}$ are the same as those from $O_{uW}^{(13),(23)}$. Contributions from $O_{uB}^{(i3),(3i)}$ are equal to those from $O_{uW}^{(i3),(3i)}$ times $\tan^4 \theta_W$. Contributions from $O_{\varphi q}^{(j,i+3)}$ are the same as those from $O_{\varphi u}^{(i+3)}$.

coefficient	LO		NLO	
	σ [fb]	scale uncertainty	σ [fb]	scale uncertainty
$C_{u\varphi}^{(13)} = 3.5$	2603	+13.0% -11.0%	3858	+7.4% -6.7%
$C_{uG}^{(13)} = 0.04$	40.1	+16.5% -13.2%	50.7	+4.0% -5.2%
$C_{uG}^{(23)} = 3.5$	171	+9.7% -8.7%	310	+7.3% -6.3%
$C_{uG}^{(32)} = 0.09$	9.53	+11.0% -9.7%	16.6	+5.5% -5.1%

TABLE III. Total cross sections for $pp \rightarrow th$. Contributions from operators $O_{u\varphi}^{(3i)}$ and $O_{uG}^{(3i)}$ are equal to those from $O_{u\varphi}^{(i3)}$ and $O_{uG}^{(i3)}$, respectively.

possibly contributing to a given final state, we illustrate in Fig. 3 the interference effect between $O_{uW}^{(23)}$ and $O_{uG}^{(23)}$, in $pp \rightarrow tZ$ production. As a matter of fact, the interference between these two operators is large and gives rise to a significant change in the rate as well as in the distributions.

Finally, Fig. 4 shows an example where kinematic variables can be used to distinguish the contributions between different operators. The Higgs boson rapidity distribution in $pp \rightarrow th$ for tuh coupling induced production is more forward than that induced by the tug coupling. The reason is that an incoming up quark, which is in general more energetic than a gluon, can emit a forward Higgs boson and turn into an off-shell top quark via a uth vertex, while the same mechanism is not possible for the utg mediated production. The same observable may also be used to discriminate between uth and cth couplings, as proposed in [39], because c and g have similar PDFs.

Summary. Precision top-quark physics will be one of the priorities at the next run of the LHC. The detection of new interactions and in particular of FCN ones, will be among the most promising searches for new physics. A consistent framework to perform such searches is provided by the dimension-six SM, *i.e.*, the SM Lagrangian augmented by all operators of dimension-six compatible with the gauge symmetries of the SM. Bounding the coef-




FIG. 2. The p_T distribution of top quark in $pp \rightarrow t\gamma$ (top) and in $pp \rightarrow th$ (bottom).

ficients of such operators first (and possibly determining them in case of deviations) requires accurate predictions for both SM and beyond processes. NLO accurate predictions in QCD are required in order to correctly extract FCN couplings from measurements or to set reasonable limits on their sizes. In this paper we have presented the first complete NLO computation for the single top production processes $u(c) + g \rightarrow t + B$, $B = \gamma, Z, h$, *i.e.*, including all dimension-six flavor-changing (two fermion) operators. In particular, the chromomagnetic operators with their extra non-trivial effects have been added. Our computation is based on the MADGRAPH5_AMC@NLO framework, thus the computation is fully automatic and can be applied to other FCN processes. The matching of the NLO results to parton shower is included as well. The K -factors in all the FCN processes are found to be large, and are in general not constant over the phase-space. Our work is a first step towards the automation of NLO computations relevant for searches of new interactions through the effective field theory framework.




FIG. 3. The interference effect in $pp \rightarrow tZ$, as a function of top p_T .




FIG. 4. Normalized rapidity distribution of the Higgs boson in $pp \rightarrow th$ induced by operators $O_{u\varphi}^{(13)}$ and $O_{uG}^{(13)}$.

I. ACKNOWLEDGEMENTS

We would like to thank G. Durieux, R. Frederix, V. Hirschi and O. Mattelaer and Y. Wang for many discussions. C. D. is a Durham International Junior Research Fellow. This work has been performed in the framework of the ERC grant 291377 “LHCTheory” and of the FP7 Marie Curie Initial Training Network MCnetITN (PITN-GA-2012-315877). The research of J. W. has been supported by the Cluster of Excellence *Precision Physics, Fundamental Interactions and Structure of Matter* (PRISMA-EXC 1098). C. Z. has been supported by the IISN “Fundamental interactions” convention 4.4517.08, and by US Department of Energy under Grant DE-AC02-98CH10886.

-
- [1] S. Glashow, J. Iliopoulos, and L. Maiani, *Weak Interactions with Lepton-Hadron Symmetry*, Phys.Rev. **D2**, 1285 (1970).
- [2] B. Mele, S. Petrarca, and A. Soddu, *A New evaluation of the $t \rightarrow cH$ decay width in the standard model*, Phys.Lett. **B435**, 401 (1998), arXiv:hep-ph/9805498.
- [3] G. Eilam, J. Hewett, and A. Soni, *Rare decays of the top quark in the standard and two Higgs doublet models*, Phys.Rev. **D44**, 1473 (1991).
- [4] J. Aguilar-Saavedra and B. Nobre, *Rare top decays $t \rightarrow c\gamma$, $t \rightarrow cg$ and CKM unitarity*, Phys.Lett. **B553**, 251 (2003), arXiv:hep-ph/0210360.
- [5] K. Olive *et al.* (Particle Data Group), *Review of Particle Physics*, Chin.Phys. **C38**, 090001 (2014).
- [6] J. Aguilar-Saavedra, *Top flavor-changing neutral interactions: Theoretical expectations and experimental detection*, Acta Phys.Polon. **B35**, 2695 (2004), arXiv:hep-ph/0409342.
- [7] J. J. Liu, C. S. Li, L. L. Yang, and L. G. Jin, *Next-to-leading order QCD corrections to the direct top quark production via model-independent FCNC couplings at hadron colliders*, Phys.Rev. **D72**, 074018 (2005), arXiv:hep-ph/0508016.
- [8] J. Gao, C. S. Li, J. J. Zhang, and H. X. Zhu, *Next-to-leading order QCD corrections to the single top quark production via model-independent t - q - g flavor-changing neutral-current couplings at hadron colliders*, Phys.Rev. **D80**, 114017 (2009), arXiv:0910.4349 [hep-ph].
- [9] Y. Zhang, B. H. Li, C. S. Li, J. Gao, and H. X. Zhu, *Next-to-leading order QCD corrections to the top quark associated with γ production via model-independent flavor-changing neutral-current couplings at hadron colliders*, Phys.Rev. **D83**, 094003 (2011), arXiv:1101.5346 [hep-ph].
- [10] B. H. Li, Y. Zhang, C. S. Li, J. Gao, and H. X. Zhu, *Next-to-leading order QCD corrections to tZ associated production via the flavor-changing neutral-current couplings at hadron colliders*, Phys.Rev. **D83**, 114049 (2011), arXiv:1103.5122 [hep-ph].
- [11] Y. Wang, F. P. Huang, C. S. Li, B. H. Li, D. Y. Shao, *et al.*, *Constraints on flavor-changing neutral-current Htq couplings from the signal of tH associated production*

- with QCD next-to-leading order accuracy at the LHC, *Phys.Rev.* **D86**, 094014 (2012), arXiv:1208.2902 [hep-ph].
- [12] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr, and B. Fuks, *FeynRules 2.0 - A complete toolbox for tree-level phenomenology*, *Comput.Phys.Commun.* **185**, 2250 (2014), arXiv:1310.1921 [hep-ph].
- [13] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **1407**, 079 (2014), arXiv:1405.0301 [hep-ph].
- [14] J. Drobnak, S. Fajfer, and J. F. Kamenik, *Flavor Changing Neutral Coupling Mediated Radiative Top Quark Decays at Next-to-Leading Order in QCD*, *Phys.Rev.Lett.* **104**, 252001 (2010), arXiv:1004.0620 [hep-ph].
- [15] J. Drobnak, S. Fajfer, and J. F. Kamenik, *QCD Corrections to Flavor Changing Neutral Coupling Mediated Rare Top Quark Decays*, *Phys.Rev.* **D82**, 073016 (2010), arXiv:1007.2551 [hep-ph].
- [16] C. Zhang and F. Maltoni, *Top-quark decay into Higgs boson and a light quark at next-to-leading order in QCD*, *Phys.Rev.* **D88**, 054005 (2013), arXiv:1305.7386 [hep-ph].
- [17] C. Zhang, *Effective field theory approach to top-quark decay at next-to-leading order in QCD*, *Phys.Rev.* **D90**, 014008 (2014), arXiv:1404.1264 [hep-ph].
- [18] S. Frixione and B. R. Webber, *Matching NLO QCD computations and parton shower simulations*, *JHEP* **0206**, 029 (2002), arXiv:hep-ph/0204244.
- [19] J. Aguilar-Saavedra, *A Minimal set of top anomalous couplings*, *Nucl.Phys.* **B812**, 181 (2009), arXiv:0811.3842 [hep-ph].
- [20] J. Aguilar-Saavedra, *A Minimal set of top-Higgs anomalous couplings*, *Nucl.Phys.* **B821**, 215 (2009), arXiv:0904.2387 [hep-ph].
- [21] M. Beneke, I. Efthymiopoulos, M. L. Mangano, J. Womersley, A. Ahmadov, et al., *Top quark physics*, (2000), arXiv:hep-ph/0003033.
- [22] G. Durieux, F. Maltoni, and C. Zhang, *A global approach to top-quark flavor-changing interactions*, in progress.
- [23] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **1010**, 085 (2010), arXiv:1008.4884 [hep-ph].
- [24] J. C. Collins, F. Wilczek, and A. Zee, *Low-Energy Manifestations of Heavy Particles: Application to the Neutral Current*, *Phys.Rev.* **D18**, 242 (1978).
- [25] R. Alonso, E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators III: Gauge Coupling Dependence and Phenomenology*, *JHEP* **1404**, 159 (2014), arXiv:1312.2014 [hep-ph].
- [26] C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer, et al., *UFO - The Universal FeynRules Output*, *Comput.Phys.Commun.* **183**, 1201 (2012), arXiv:1108.2040 [hep-ph].
- [27] G. Ossola, C. G. Papadopoulos, and R. Pittau, *Reducing full one-loop amplitudes to scalar integrals at the integrand level*, *Nucl.Phys.* **B763**, 147 (2007), arXiv:hep-ph/0609007.
- [28] C. Degrande, *Automatic evaluation of UV and R2 terms for beyond the Standard Model Lagrangians: a proof-of-principle*, (2014), arXiv:1406.3030 [hep-ph].
- [29] J. Pumplin, D. Stump, J. Huston, H. Lai, P. M. Nadolsky, et al., *New generation of parton distributions with uncertainties from global QCD analysis*, *JHEP* **0207**, 012 (2002), arXiv:hep-ph/0201195.
- [30] S. Frixione, *Isolated photons in perturbative QCD*, *Phys.Lett.* **B429**, 369 (1998), arXiv:hep-ph/9801442.
- [31] T. Sjostrand, S. Mrenna, and P. Z. Skands, *PYTHIA 6.4 Physics and Manual*, *JHEP* **0605**, 026 (2006), arXiv:hep-ph/0603175.
- [32] G. Corcella, I. Knowles, G. Marchesini, S. Moretti, K. Odagiri, et al., *HERWIG 6: An Event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)*, *JHEP* **0101**, 010 (2001), arXiv:hep-ph/0011363.
- [33] S. Frixione, E. Laenen, P. Motylinski, B. R. Webber, and C. D. White, *Single-top hadroproduction in association with a W boson*, *JHEP* **0807**, 029 (2008), arXiv:0805.3067 [hep-ph].
- [34] S. Chatrchyan et al. (CMS Collaboration), *Search for Flavor-Changing Neutral Currents in Top-Quark Decays $t \rightarrow Zq$ in pp Collisions at $\sqrt{s} = 8$ TeV*, *Phys.Rev.Lett.* **112**, 171802 (2014), arXiv:1312.4194 [hep-ex].
- [35] CMS Collaboration, *Combined multilepton and diphoton limit on $t \rightarrow cH$* , CMS-PAS-HIG-13-034 (2014).
- [36] G. Aad et al. (ATLAS Collaboration), *Search for top quark decays $t \rightarrow qH$ with $H \rightarrow \gamma\gamma$ using the ATLAS detector*, *JHEP* **1406**, 008 (2014), arXiv:1403.6293 [hep-ex].
- [37] The ATLAS Collaboration, *Search for single top-quark production via FCNC in strong interaction in $\sqrt{s} = 8$ TeV ATLAS data*, (2013).
- [38] CMS Collaboration, *Search for anomalous single top quark production in association with a photon*, CMS-PAS-TOP-14-003 (2014).
- [39] A. Greljo, J. F. Kamenik, and J. Kopp, *Disentangling Flavor Violation in the Top-Higgs Sector at the LHC*, *JHEP* **1407**, 046 (2014), arXiv:1404.1278 [hep-ph].