

This is the accepted manuscript made available via CHORUS. The article has been published as:

Nonzero θ_{13} in $SO(3) \rightarrow A_4$ lepton models

Yuval Grossman and Wee Hao Ng

Phys. Rev. D **91**, 073005 — Published 8 April 2015

DOI: [10.1103/PhysRevD.91.073005](https://doi.org/10.1103/PhysRevD.91.073005)

Nonzero θ_{13} in $SO(3) \rightarrow A_4$ lepton models

Yuval Grossman^{*} and Wee Hao Ng[†]

Laboratory for Elementary-Particle Physics, Cornell University, Ithaca, N.Y.

Abstract

The simplest neutrino mass models based on A_4 symmetry predict $\theta_{13} = 0$ at tree level, a value that contradicts recent data. We study models that arise from the spontaneous breaking of an $SO(3)$ symmetry to its A_4 subgroup, and find that such models can naturally accommodate a nonzero θ_{13} at tree level. Standard Model charged leptons mix with additional heavy ones to generate a θ_{13} that scales with the ratio of the A_4 -breaking to $SO(3)$ -breaking scales. A suitable choice of energy scales hence allows one to reproduce the correct lepton mixing angles. We also consider an alternative approach where we modify the alignment of flavons associated with the charged lepton masses, and find that the effects on θ_{13} are enhanced by a factor that scales as m_τ/m_μ .

^{*} yg73@cornell.edu

[†] wn68@cornell.edu

I. INTRODUCTION

For a long time, the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix U_{PMNS} [1, 2] was believed to be consistent with the tri-bimaximal mixing matrix [3]:

$$U_{\text{TBM}} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \end{pmatrix}. \quad (1)$$

The pattern exhibited by the tri-bimaximal mixing matrix seemed to suggest some underlying symmetry in the lepton sector, thus motivating the development of lepton models based on discrete flavor symmetries. One class of models was based on the discrete group A_4 [4–47] (see [48–51] for reviews). The most basic implementation of these A_4 models comprise the SM leptons and Higgs, right-handed singlet neutrinos, as well as two scalar flavons ϕ and ϕ' . These fields are assigned into representations of A_4 , where the flavons, in particular, are three-dimensional representations. The Lagrangian is invariant under A_4 , but this symmetry is spontaneously broken when the flavons acquire VEVs, thus generating mass terms for the leptons. To reproduce the tri-bimaximal mixing at tree level, the lepton mass matrices have to take specific forms, which imply specific alignments $\langle\phi\rangle = (v, v, v)$ and $\langle\phi'\rangle = (v', 0, 0)$ for the flavon VEVs. Such alignments may be explained by various UV completions based on supersymmetry or extra dimensions [52–59].

The recent discovery of nonzero θ_{13} by the Daya Bay [60] and RENO [61] experiments have thrown the tri-bimaximal mixing pattern into question. The current experimental status of the elements of U_{PMNS} is shown in Table I. The best fit value of $|\sin(\theta_{13})|$ for both hierarchies is approximately 0.16, significantly different from zero. This can be interpreted in two different ways, the first being that there is really no symmetry behind the lepton mixing (anarchy), and the second that there is a large modification to tri-bimaximal mixing in the underlying symmetry models.

In accordance with the second viewpoint, various ideas have been proposed to modify the A_4 models to reproduce a nonzero θ_{13} . One way is to consider higher dimension operators, which introduce correction terms to the mass matrices of relative size given by v/Λ and v'/Λ (where Λ is the cutoff scale) [50, 59, 63]. Another approach is to extend the A_4 model to include more flavons that contribute to the lepton mass matrices [64–68]. Yet another avenue is to introduce perturbations in the flavon sector that modify their vacuum alignments and hence the form of the lepton mass matrices [69–72]. Radiative corrections as a way to generate nonzero θ_{13} have also been considered in [73–77].

In this paper, we focus on a specific class of models [78–80] that can be regarded as UV comple-

Parameter	Best fit	1σ range	3σ range
$\sin^2(\theta_{12})$ (NH, IH)	0.307	$0.291 - 0.325$	$0.259 - 0.359$
$\sin^2(\theta_{13})$ (NH)	0.0243	$0.0216 - 0.0266$	$0.0169 - 0.0313$
$\sin^2(\theta_{13})$ (IH)	0.0242	$0.0219 - 0.0267$	$0.0171 - 0.0315$
$\sin^2(\theta_{23})$ (NH)	0.386	$0.365 - 0.410$	$0.331 - 0.637$
$\sin^2(\theta_{23})$ (IH)	0.392	$0.370 - 0.431$	$0.335 - 0.663$
δ (NH)	1.08π	$0.77\pi - 1.36\pi$	—
δ (IH)	1.09π	$0.83\pi - 1.47\pi$	—

TABLE I. Current experimental status of the mixing angles in U_{PMNS} [62]. NH and IH stands for normal and inverted hierarchy respectively.

tions of certain A_4 models. These UV models are invariant under a continuous symmetry group, for example $SO(3)$, of which A_4 is a subgroup. This symmetry is spontaneously broken to A_4 by certain flavons that acquire a specific pattern of VEVs, generating A_4 models as effective low energy theories. In light of the recent measurements, it is worth investigating how a nonzero θ_{13} can be accommodated in such models.

An interesting observation is that that such models actually already predict θ_{13} to be nonzero even with the usual vacuum alignment. $SO(3)$ -based models in general require additional heavy charged leptons to complete the $SO(3)$ representations the Standard Model (SM) charged leptons belong to. While the mixing between SM and heavy charged leptons is very small, it is enough to modify the pattern of the light charged lepton mass matrix, which breaks the tri-bimaximal mixing pattern. The idea of modifying the charged-lepton mass matrix to obtain a nonzero θ_{13} is certainly not new; however, seldom has the context been that of mixing between SM and heavy charged leptons. This will be the main focus of our work, using a model motivated by [79] as an illustration. An interesting result is that the size of θ_{13} scales with the ratio of A_4 -breaking to $SO(3)$ -breaking scales. In other words, θ_{13} may reflect certain features of the UV physics, rather than simply arising from some arbitrary coefficients.

A second way to obtain a nonzero θ_{13} , with a clear parallel in typical A_4 models, is to allow the VEVs to deviate from the usual alignment that reproduces U_{TBM} . An interesting feature of this approach is the presence of an enhancement factor that scales as m_τ/m_μ should the flavons involved be those associated with the charged lepton masses. In other words, a small angular deviation of these flavons from the usual alignment can give rise to a much larger θ_{13} .

This paper is organized as follows. In Section II we provide an overview of the $SO(3) \rightarrow A_4$

model of [79]. In Section III we present our main results, where we demonstrate that mixing of the SM charged leptons with heavy ones give rise to a nonzero tree-level θ_{13} , the size of which is related to the ratio of scales. In Section IV, we discuss the second approach of modifying the flavon alignment associated with charged leptons and demonstrate the existence of the enhancement factor. We summarize our work in Section V. Details of the model and the numerical simulations are given in the appendix.

II. REVIEW OF THE $SO(3) \rightarrow A_4$ MODEL

A. Field content

We review a lepton model motivated by [79] where a larger continuous flavour symmetry is spontaneously broken to the A_4 subgroup. The symmetries of this model are the electroweak gauge symmetry $SU(2)_L \times U(1)_Y$ as well as a global $SO(3)_F$. The fields and their representations are summarized in Table II.

For the lepton sector, the three SM left-handed $SU(2)_L$ doublets ψ_l form a **3** under $SO(3)_F$. Among the three SM charged right-handed $SU(2)_L$ singlets, ψ_e is a **1**, while the other two have been subsumed into a **5** denoted by ψ_m . In doing so, we now have three extra charged right-handed $SU(2)_L$ singlets from ψ_m , which we give large masses to by introducing a charged left-handed $SU(2)_L$ singlet ψ_f that form a **3**. Finally we introduce three right-handed neutrinos ψ_n which form a **3**.

In the scalar sector, we have the SM Higgs, H , which is a singlet of the flavor group, and four flavons ϕ , ϕ' , ϕ_5 , and T which are **3**, **3**, **5** and **7** respectively. The flavon T is responsible for the $SO(3)_F \rightarrow A_4$ breaking, and is required to be at least a **7** since that is the smallest representation of $SO(3)$ that can have an A_4 -invariant VEV. While ϕ and ϕ' can be identified with the usual flavons in the minimal A_4 model, the extra flavon ϕ_5 is required here to prevent the muon and tau from becoming degenerate. This degeneracy is due to the right-handed muon and tau being part of the same $SO(3)_F$ multiplet and hence sharing the same Yukawa coupling with ϕ . An extra flavon ϕ_5 in a different $SO(3)_F$ representation from ϕ is needed to lift this degeneracy. (We note that this degeneracy is actually also lifted by the block-diagonalization process to be discussed later, but the resulting mass differences are in general too small.)

Field	$SU(2)_L$	$U(1)_Y$	$SO(3)_F$
ψ_l	2	$-\frac{1}{2}$	3
ψ_f	1	-1	3
ψ_e	1	-1	1
ψ_m	1	-1	5
ψ_n	1	0	3
H	2	$\frac{1}{2}$	1
ϕ	1	0	3
ϕ'	1	0	3
ϕ_5	1	0	5
T	1	0	7

TABLE II. Matter fields and the representations they transform as under the gauge symmetry $SU(2)_L \times U(1)_Y$ and global symmetry $SO(3)_F$. We have partitioned the fields into left-handed leptons, right-handed leptons and scalars.

B. Lagrangian

We now assume the following terms in the Lagrangian for the charged leptons and neutrinos:

$$\mathcal{L}_e = -y_e \overline{\psi_l}^a \frac{H}{\Lambda} \phi^a \psi_e - y_m \overline{\psi_l}^a \frac{H}{\Lambda} \phi^b \psi_m^{ab} - y_m^T \overline{\psi_l}^a \frac{H}{\Lambda} T^{abc} \psi_m^{bc} - y_m^5 \epsilon^{abc} \overline{\psi_l}^a \frac{H}{\Lambda} \phi_5^{bd} \psi_m^{cd} \quad (2)$$

$$-y_e' \overline{\psi_f}^a \phi^a \psi_e - y_m' \overline{\psi_f}^a \phi^b \psi_m^{ab} - y_m^{T'} \overline{\psi_f}^a T^{abc} \psi_m^{bc} - y_m^{5'} \epsilon^{abc} \overline{\psi_f}^a \phi_5^{bd} \psi_m^{cd} + \text{h.c.},$$

$$\mathcal{L}_\nu = -M \overline{\psi_n^c}^a \psi_n^a - \frac{x_\nu}{\Lambda} \overline{\psi_n^c}^a \psi_n^b \phi'^c T^{abc} - y_\nu \overline{\psi_l}^a H^c \psi_n^a + \text{h.c.}, \quad (3)$$

where a , b , and c are $SO(3)_F$ indices running from 1 to 3, and Λ is the cutoff scale of the model. This Lagrangian is not renormalizable and includes certain dimension-five operators. These are required to give masses to the light charged leptons, and to lift the degeneracy of the light neutrinos.

There are other terms in the Lagrangian involving only the scalars, of which we will just focus on the renormalizable self-interactions of the flavon T :

$$V(T) = -\frac{\mu^2}{2} T^{abc} T^{abc} + \frac{\lambda}{4} (T^{abc} T^{abc})^2 + c T^{abc} T^{bcd} T^{def} T^{efa}. \quad (4)$$

It is shown in [79] that conditions on λ and c exist such that $V(T)$ has an A_4 invariant minimum, which breaks $SO(3)_F$ into its A_4 subgroup. We then end up with an effective non-minimal A_4 model, with three more pairs of left- and right-handed charged leptons, and one more flavon ϕ_5 .

Before proceeding further, we acknowledge two issues with the lepton Lagrangian. First, this is not the most general Lagrangian consistent with the gauge and global symmetries. [79] proposed

an auxiliary Z_2 symmetry to forbid the excluded terms, but a careful check shows that it does not work. Nonetheless, we have been able to identify modified models that can reproduce the same lepton mass matrices as this Lagrangian, the details of which are provided in Appendix A. The second issue is that since the flavons can acquire VEVs of different scales, mass terms arising from higher-dimension operators need not be smaller than those from lower-dimension ones, especially if they contain multiple factors of the larger VEVs. Therefore, the errors associated with truncation of the Lagrangian can be significant. This issue will also be addressed in Appendix A. For the rest of our work, we will neglect both issues and continue to work with the given Lagrangian to demonstrate the key ideas behind our approach.

C. Lepton mass matrices and U_{PMNS}

We assume that the flavons ϕ , ϕ_5 and ϕ' acquire VEVs with the following alignments:

$$\langle \phi \rangle = \begin{pmatrix} v \\ v \\ v \end{pmatrix}, \quad \langle \phi_5 \rangle = \begin{pmatrix} 0 & v_5 & v_5 \\ v_5 & 0 & v_5 \\ v_5 & v_5 & 0 \end{pmatrix}, \quad \langle \phi' \rangle = \begin{pmatrix} v' \\ 0 \\ 0 \end{pmatrix}. \quad (5)$$

We also assume that the VEV of T satisfies $v_T \gg v, v', v_5$, in accordance to the picture of $SO(3)_F$ broken to A_4 . After electroweak symmetry breaking, the Higgs boson H acquires a VEV $v_H = (246/\sqrt{2})$ GeV, and we obtain two 6×6 matrices: $M_l^{6 \times 6}$ for the charged-lepton Dirac masses and $M_\nu^{6 \times 6}$ for the neutrino Majorana masses.

In [79], the mixing between the SM and the new charged leptons were considered to be small and hence neglected. In that case, the leading mass matrix M_l for the three light charged leptons is simply given by the upper-left 3×3 block of $M_l^{6 \times 6}$:

$$M_l = \frac{v_H}{\Lambda} \begin{pmatrix} y_e v & y_m v + y_m^5 v_5 (\omega^2 - \omega) & y_m v + y_m^5 v_5 (\omega - \omega^2) \\ y_e v & [y_m v + y_m^5 v_5 (\omega^2 - \omega)] \omega & [y_m v + y_m^5 v_5 (\omega - \omega^2)] \omega^2 \\ y_e v & [y_m v + y_m^5 v_5 (\omega^2 - \omega)] \omega^2 & [y_m v + y_m^5 v_5 (\omega - \omega^2)] \omega \end{pmatrix}, \quad (6)$$

where $\omega = e^{2\pi i/3}$. The charged lepton masses can be obtained by diagonalizing $M_l(M_l)^\dagger$ and taking the square root, and we obtain

$$m_e = \left| \sqrt{3} \frac{y_e v_H v}{\Lambda} \right|, \quad m_\mu, m_\tau = \left| \sqrt{3} \frac{y_m v_H v}{\Lambda} \pm 3i \frac{y_m^5 v_H v_5}{\Lambda} \right|. \quad (7)$$

We note that the Yukawas have to be fine-tuned to generate the correct charged lepton masses. Therefore, this model does not ameliorate the fine-tuning issue also present in minimal A_4 models¹.

¹ Certain A_4 models [51, 52] resolve the fine-tuning issue by relegating the electron mass to higher-dimensional

The unitary transformation required to diagonalize $M_l(M_l)^\dagger$ is

$$U_l = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}. \quad (8)$$

It is important to note that (8) is a result of M_l taking the form

$$M_{\text{aligned}} = \begin{pmatrix} a & b & c \\ a & b\omega & c\omega^2 \\ a & b\omega^2 & c\omega \end{pmatrix}, \quad (9)$$

where a , b and c are constants.

For the neutrino sector, the 6×6 Majorana mass matrix $M_\nu^{6 \times 6}$ can be block-diagonalized, and the resulting upper-left 3×3 block is identified with the mass matrix M_ν of the three light neutrinos

$$M_\nu = -y_\nu^2 v_H^2 \begin{pmatrix} -\frac{1}{M} & 0 & 0 \\ 0 & -\frac{M}{M^2 - x_\nu^2 (v' v_T / \Lambda)^2} & \frac{x_\nu (v' v_T / \Lambda)}{M^2 - x_\nu^2 (v' v_T / \Lambda)^2} \\ 0 & \frac{x_\nu (v' v_T / \Lambda)}{M^2 - x_\nu^2 (v' v_T / \Lambda)^2} & -\frac{M}{M^2 - x_\nu^2 (v' v_T / \Lambda)^2} \end{pmatrix}. \quad (10)$$

Note that this is exactly the see-saw mechanism, as M_ν becomes very small if the Majorana mass parameters M and $v' v_T / \Lambda$ for ψ_n are much larger than v_H . The light neutrino masses can be obtained by diagonalizing M_ν . We choose the following assignment for the mass eigenvalues:

$$m_1 = \left| \frac{y_\nu^2 v_H^2}{M + x_\nu (v' v_T / \Lambda)} \right|, \quad m_2 = \left| \frac{y_\nu^2 v_H^2}{M} \right|, \quad m_3 = \left| \frac{y_\nu^2 v_H^2}{M - x_\nu (v' v_T / \Lambda)} \right|. \quad (11)$$

Such an assignment can accommodate both normal and inverted hierarchies, but the latter requires fine-tuning between the magnitude and phase of the combination $x_\nu v' v_T / (M \Lambda)$ [59]. Therefore, for the rest of our work, we will only focus on normal hierarchy.

The unitary transformation required to diagonalize M_ν is:

$$U_\nu = i \begin{pmatrix} 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{pmatrix}. \quad (12)$$

The PMNS matrix is then given by:

$$U_{\text{PMNS}} = U_l (U_\nu)^\dagger = \begin{pmatrix} -i\sqrt{\frac{2}{3}} & -i\frac{1}{\sqrt{3}} & 0 \\ i\frac{1}{\sqrt{6}} & -i\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ i\frac{1}{\sqrt{6}} & -i\frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix}, \quad (13)$$

operators, through the use of additional symmetries and flavons. While we do not show it here, an analogous approach can be adopted in our model to naturally suppress y_e . However, as we see later, this does not fully resolve the issue since subleading contributions from block diagonalization do not scale with y_e in general.

which can be brought into the form U_{TBM} by redefining the phases of ν_1 , ν_2 and τ . Note that the tri-bimaximal mixing pattern obtained above depends on U_l taking the form Eq. (8). Any deviation of M_l from M_{aligned} would result in deviation of U_{PMNS} from U_{TBM} . We will exploit this fact in the next section in order to generate a nonzero θ_{13} . We also note that Eq. (13) omits certain nonunitary matrix factors which we show in Appendix B to be negligible.

We now briefly mention the masses of the heavy leptons, which can be obtained from the corresponding full 6×6 mass matrices. The heavy charged lepton masses are typically of order $y_m^{T'} v_T$, and the heavy neutrino masses of order $M \sim x_\nu v' v_T / \Lambda$. (We will demonstrate shortly that M and $x_\nu v' v_T / \Lambda$ are typically of the same scale.)

D. Energy scales

We can use our results for the light lepton masses to obtain a rough picture of the energy scales in this model. We first consider the neutrino masses given in Eq. (11). The current experimental results are as follows [62]:

$$\delta m^2 \equiv m_2^2 - m_1^2 = 7.54_{-0.22}^{+0.26} \times 10^{-5} \text{ eV}^2, \quad (14)$$

$$|\Delta m^2| \equiv \left| m_3^2 - \frac{m_1^2 + m_2^2}{2} \right| = \begin{matrix} 2.43_{-0.10}^{+0.06} \text{ (NH)} \\ 2.42_{-0.11}^{+0.07} \text{ (IH)} \end{matrix} \times 10^{-3} \text{ eV}^2. \quad (15)$$

Assuming normal hierarchy, we find that $M \sim x_\nu v' v_T / \Lambda \sim 10^{15} |y_\nu| \text{ GeV}$, with neutrino masses $m_1 \sim 6 \text{ meV}$, $m_2 \sim 10 \text{ meV}$ and $m_3 \sim 50 \text{ meV}$ independent of y_ν . The fact that $M \sim x_\nu v' v_T / \Lambda$ is not surprising since we require large cancellations in $M - x_\nu v' v_T / \Lambda$ to ensure that $m_3 \gg m_1, m_2$, as implied by the experimental results. This gives rise to the following hierarchy of energy scales:

$$v_H \sim 100 \text{ GeV} \ll \frac{M}{x_\nu} \sim \frac{v' v_T}{\Lambda} \ll \{v, v_5, v'\} \ll v_T \ll \Lambda. \quad (16)$$

The charged-lepton masses provide further constraints on the hierarchy. Since $m_\tau \sim 1 \text{ GeV}$, Eq. (7) implies that $\{v, v_5, v'\} / \Lambda \gtrsim O(10^{-3})$ so that the associated Yukawas remain perturbatively small. This somewhat restricts the ratio of symmetry breaking scales $\epsilon \equiv \{v, v_5, v'\} / v_T$, if we do not want the scales $\{v, v_5, v'\}$, v_T and Λ to be too close to one another.

We now consider an example to illustrate the typical energy scales involved. Assuming all Yukawas to be $O(1)$, we find that $M \sim 10^{15} \text{ GeV}$, $\{v, v_5, v'\} \sim 10^{16} \text{ GeV}$, $v_T \sim 10^{18} \text{ GeV}$ and $\Lambda \sim 10^{19} \text{ GeV}$. Other values can be obtained by varying the Yukawas, although this is limited by the requirement that the Yukawas remain perturbative.

III. EFFECTS OF MIXING IN THE CHARGED LEPTON SECTOR

A. Obtaining the light charged lepton mass-squared matrix

The tri-bimaximal mixing pattern of this model is the result of the unitary matrix U_l that diagonalizes $M_l(M_l)^\dagger$ taking the form Eq. (8). This in turn requires the light charged lepton mass matrix M_l to take the form M_{aligned} given in Eq. (9). As we shall see below, subleading corrections from block diagonalisation modifies the form of M_l and hence lead to a nonzero θ_{13} .

To obtain the exact form of $M_l(M_l)^\dagger$ in the general case, we start with the full 6×6 mass matrix $M_l^{6 \times 6}$ obtained from the Lagrangian Eq. (2). We express $M_l^{6 \times 6}$ in terms of 3×3 matrices A , B , C and D :

$$M_l^{6 \times 6} \equiv \begin{pmatrix} \frac{v_H}{\Lambda} A & \frac{v_H}{\Lambda} B \\ C & D \end{pmatrix}. \quad (17)$$

where

$$A = \begin{pmatrix} y_e v & [y_m v + y_m^5 v_5 (\omega^2 - \omega)] & [y_m v + y_m^5 v_5 (\omega - \omega^2)] \\ y_e v & [y_m v + y_m^5 v_5 (\omega^2 - \omega)] \omega & [y_m v + y_m^5 v_5 (\omega - \omega^2)] \omega^2 \\ y_e v & [y_m v + y_m^5 v_5 (\omega^2 - \omega)] \omega^2 & [y_m v + y_m^5 v_5 (\omega - \omega^2)] \omega \end{pmatrix}, \quad (18)$$

$$B = \begin{pmatrix} y_m v + 2y_m^T v_T & y_m v + y_m^5 v_5 & -y_m^5 v_5 \\ y_m v & 2y_m^T v_T & y_m v \\ y_m v + y_m^5 v_5 + y_m^T v_T & y_m v - y_m^5 v_5 & y_m^T v_T \end{pmatrix}, \quad (19)$$

$$C = \begin{pmatrix} y'_e v & [y'_m v + y_m^{5'} v_5 (\omega^2 - \omega)] & [y'_m v + y_m^{5'} v_5 (\omega - \omega^2)] \\ y'_e v & [y'_m v + y_m^{5'} v_5 (\omega^2 - \omega)] \omega & [y'_m v + y_m^{5'} v_5 (\omega - \omega^2)] \omega^2 \\ y'_e v & [y'_m v + y_m^{5'} v_5 (\omega^2 - \omega)] \omega^2 & [y'_m v + y_m^{5'} v_5 (\omega - \omega^2)] \omega \end{pmatrix}, \quad (20)$$

$$D = \begin{pmatrix} y'_m v + 2y_m^{T'} v_T & y'_m v + y_m^{5'} v_5 & -y_m^{5'} v_5 \\ y'_m v & 2y_m^{T'} v_T & y'_m v \\ y'_m v + y_m^{5'} v_5 + y_m^{T'} v_T & y'_m v - y_m^{5'} v_5 & y_m^{T'} v_T \end{pmatrix}. \quad (21)$$

We then block-diagonalize $M_l^{6 \times 6} (M_l^{6 \times 6})^\dagger$ and obtain $M_l(M_l)^\dagger$ from the upper-left 3×3 block:

$$M_l(M_l)^\dagger = \frac{v_H^2}{\Lambda^2} \left[AA^\dagger + BB^\dagger - (AC^\dagger + BD^\dagger)(CC^\dagger + DD^\dagger)^{-1}(CA^\dagger + DB^\dagger) \right]. \quad (22)$$

We see that when we previously assumed M_l to be given by Eq. (6), we have kept only the leading term $v_H^2 AA^\dagger/\Lambda^2$.

We now examine the effects of the other terms. We first note that A and C are both of the form M_{aligned} . We further define

$$E \equiv B - \frac{y_m^T}{y_m^{T'}} D, \quad (23)$$

and the small parameter

$$\epsilon \equiv O(v/v_T) \sim O(v_5/v_T). \quad (24)$$

Assuming all Yukawa couplings to be of the same order y , we find that the scales of B and D are of order yv_T , while those of A , C and E are of order yv and hence ϵ smaller. (We quantify the scale of a matrix by the characteristic size of the eigenvalues). We can thus expand Eq. (22) in ϵ . To the lowest nontrivial order we find that $M_l(M_l)^\dagger$ is factorizable with

$$M_l = \frac{v_H}{\Lambda} (A - BD^{-1}C) = \frac{v_H}{\Lambda} \left(A - \frac{y_m^T}{y_m^{T'}} C - ED^{-1}C \right). \quad (25)$$

Since both A and C are of the form M_{aligned} , so is any linear superposition of them, and thus the first correction term $-(y_m^T/y_m^{T'})C$ does not give a nonzero θ_{13} . The second correction term $-ED^{-1}C$, of order ϵ , is what generate deviations of M_l from M_{aligned} . This in turn suggests that the size of θ_{13} is also of order ϵ . In other words, θ_{13} reflects the ratio of the A_4 -breaking to $SO(3)_F$ -breaking scales.

B. Amplification from nearly-degenerate mass eigenvalues

While the above analysis seems to suggest that $\theta_{13} \sim \epsilon$, there is actually a numerical factor that enhances the size of θ_{13} . From perturbation theory, the mixing angle between two eigenvectors is given approximately by the ratio of the small perturbation mixing them to the difference between their eigenvalues. Our previous result of $\theta_{13} \sim \epsilon$ implicitly assumes that the difference between mass eigenvalues are of order m_τ . However, in the actual charged-lepton mass spectrum, m_e and m_μ are nearly degenerate relative to m_τ , suggesting an enhancement in the mixing angle.

To illustrate this enhancement, it is useful to write $M_l(M_l)^\dagger$ in a different basis:

$$U_l M_l (M_l)^\dagger (U_l)^\dagger = \frac{v_H^2}{\Lambda^2} U_l \left(A - \frac{y_m^T}{y_m^{T'}} C \right) \left(A - \frac{y_m^T}{y_m^{T'}} C \right)^\dagger (U_l)^\dagger + \Delta \quad (26)$$

$$= \begin{pmatrix} m_a^2 & 0 & 0 \\ 0 & m_b^2 & 0 \\ 0 & 0 & m_c^2 \end{pmatrix} + \begin{pmatrix} \Delta_{11} & \Delta_{21}^* & \Delta_{31}^* \\ \Delta_{21} & \Delta_{22} & \Delta_{32}^* \\ \Delta_{31} & \Delta_{32} & \Delta_{33} \end{pmatrix}, \quad (27)$$

where U_l takes the specific form given in Eq. (8) and

$$\Delta \equiv -\frac{v_H^2}{\Lambda^2} U_l \left[\left(A - \frac{y_m^T}{y_m^{T'}} C \right) C^\dagger (D^\dagger)^{-1} E^\dagger + E D^{-1} C \left(A - \frac{y_m^T}{y_m^{T'}} C \right)^\dagger \right] U_l^\dagger, \quad (28)$$

comes from the order ϵ corrections in Eq. (25). In this basis, θ_{13} is determined by how much the perturbation Δ changes the zeroth-order eigenvector $(1, 0, 0)$. From perturbation theory, θ_{13} is roughly $\Delta_{21}/(m_b^2 - m_a^2)$ or $\Delta_{31}/(m_c^2 - m_a^2)$, whichever is larger.

Since the actual eigenvalues are given by the charged lepton masses, we assume the following sizes for the zeroth-order eigenvalues:

$$m_a^2 \lesssim m_\mu^2, \quad m_b^2 \sim m_\mu^2, \quad m_c^2 \sim m_\tau^2. \quad (29)$$

The perturbation matrix Δ is of the scale $O(m_c^2 \epsilon)$, and so naively we might expect $\Delta_{21}/(m_b^2 - m_a^2) \sim O(m_\tau^2 \epsilon / m_\mu^2)$ and $\Delta_{31}/(m_c^2 - m_a^2) \sim O(\epsilon)$. This will imply that θ_{13} is enhanced compared to the naive expectation ϵ by m_τ^2 / m_μ^2 . However, as we show below, while indeed $\Delta_{31} \sim O(m_c^2 \epsilon)$, we instead find that $\Delta_{21} \sim O(m_b m_c \epsilon)$. This is due to $M_l(M_l)^\dagger$ being factorizable at a well-defined order in ϵ , as we have demonstrated in Eq. (25). As a result, $\Delta_{21}/(m_b^2 - m_a^2) \sim O(m_\tau \epsilon / m_\mu)$, from which we conclude that

$$\theta_{13} \sim O\left(\frac{m_\tau}{m_\mu} \epsilon\right). \quad (30)$$

Thus the size of θ_{13} is enhanced by a smaller factor m_τ / m_μ .

We now explain why the factorizability of $M_l(M_l)^\dagger$ at a well-defined order in ϵ sets the sizes of Δ_{21} and Δ_{31} . In such a scenario, we expect $U_l M_l (U_l)^\dagger$ to be given by

$$U_l M_l (U_l)^\dagger = \begin{pmatrix} m_a & 0 & 0 \\ 0 & m_b & 0 \\ 0 & 0 & m_c \end{pmatrix} + \epsilon m_c R \quad (31)$$

where R is an $O(1)$ matrix. Substituting this into $U_l M_l (M_l)^\dagger (U_l)^\dagger$, we find that

$$\Delta = \epsilon m_c \begin{pmatrix} m_a R_{11}^* + m_a R_{11} & m_a R_{21}^* + m_b R_{12} & m_a R_{31}^* + m_c R_{13} \\ m_b R_{12}^* + m_a R_{21} & m_b R_{22}^* + m_b R_{22} & m_b R_{32}^* + m_c R_{23} \\ m_c R_{13}^* + m_a R_{31} & m_c R_{23}^* + m_b R_{32} & m_c R_{33}^* + m_c R_{33} \end{pmatrix} \quad (32)$$

Therefore, we conclude that

$$\Delta_{21} = \epsilon m_c (m_b R_{12}^* + m_a R_{21}) \sim O(m_b m_c \epsilon), \quad (33)$$

$$\Delta_{31} = \epsilon m_c (m_c R_{13}^* + m_a R_{31}) \sim O(m_c^2 \epsilon). \quad (34)$$

in agreement with our assertions above.

A few points to note: First, the analysis above breaks down when $\epsilon \gg m_\mu/m_\tau$, since this implies that the perturbation $\Delta_{21} \gg m_\mu^2$, in which case we also require the zeroth-order eigenvalues m_a^2 , $m_b^2 \gg m_\mu^2$ so that large cancellations can occur to give two small eigenvalues m_e^2 and m_μ^2 . Second, as we shall see from the simulation results next section, there are various “large” $O(1)$ factors that we have not taken into account in our analysis, so the exact θ_{13} may be a few times smaller than the prediction above.

C. Verifying the results via simulation

To verify the above results, we compute the exact tree-level U_{PMNS} for two large collections of random parameter sets \mathcal{C}_1 and \mathcal{C}_2 . Details of how they are generated are provided in Appendix C 1. The mass eigenvalues for parameter sets in \mathcal{C}_1 are unconstrained, whereas those in \mathcal{C}_2 are required to have the correct charged-lepton mass ratios. We note that only a very small fraction of random parameter sets satisfy the conditions for \mathcal{C}_2 , a consequence of the charged lepton mass fine-tuning.

Figure 1(a) shows the value of $\sin(\theta_{13})$ against $\sup\{v/v_T, v_5/v_T\}$ for \mathcal{C}_1 , which in general agrees with the expectation that $\theta_{13} \sim O(\epsilon)$. Since no conditions have been imposed on the mass ratios, all three mass eigenvalues are typically of the same order of magnitude and hence there is no significant amplification effect.

Figure 1(b) shows the value of $\sin(\theta_{13})$ against $\sup\{v/v_T, v_5/v_T\}$ for \mathcal{C}_2 . While we still have $\theta_{13} \propto O(\epsilon)$, the proportionality constant is now about five times that of \mathcal{C}_1 . Since the eigenvalues of \mathcal{C}_2 are now of the correct ratios, we attribute the larger proportionality constant to the amplification effect, although the amplification is smaller than our original prediction due to “large” $O(1)$ factors that we have neglected in our analysis. We hence conclude that the experimental best fit value of $|\sin(\theta_{13})| \simeq 0.16$ corresponds to the ratio of symmetry-breaking scales $\epsilon \sim 0.05$.

D. Compatibility with experimental constraints

We now discuss whether this model can satisfy the experimental constraints on lepton masses and mixing angles. We first consider lepton masses. As we have shown in Section II D, the measured




FIG. 1. Graphs of $\sin(\theta_{13})$ against $\sup\{v/v_T, v_5/v_T\}$ for two collections (a) \mathcal{C}_1 and (b) \mathcal{C}_2 of random parameter sets. Various lines have been included for reference. Both graphs demonstrate the linear dependence predicted in our analysis. Collection \mathcal{C}_2 has a charged lepton mass spectrum much closer to the actual hierarchy than \mathcal{C}_1 , as a result of which the constant of proportionality is significantly enhanced by the amplification effect.

neutrino mass differences δm^2 and $|\Delta m^2|$ are certainly compatible with the model provided that $M \sim x_\nu v' v_T / \Lambda$. We have also demonstrated that the correct charged-lepton mass spectrum can be reproduced in Section III C, although significant fine-tuning is required.

We now focus on the mixing angles. A major concern is that corrections that reproduce the measured θ_{13} might also affect the other mixing angles θ_{12} and θ_{23} to the extent that they no longer remain compatible with experimental observations. In particular, many models predict the same size of corrections to θ_{12} and θ_{13} , in which case a large θ_{13} will imply a large correction to θ_{12} .

Figure 2 shows plots of $\sin(\theta_{12})$ and $\sin(\theta_{23})$ against $\sin(\theta_{13})$, using parameter sets from \mathcal{C}_2 and zoomed into the regions around $\sin(\theta_{13}) \sim 0.15$. We see that for a large $\sin(\theta_{13}) \sim 0.15$, $\sin(\theta_{12})$ is fairly evenly distributed between 0.45 and 0.7, with about 25% of the points within the 3σ range, as opposed to a bimodal distribution peaked at the two extremes. We hence conclude that corrections to θ_{12} need not be of the same size as θ_{13} , and so this model can be made compatible with all three experimental mixing angles.




FIG. 2. Graphs of $\sin(\theta_{12})$ and $\sin(\theta_{23})$ against $\sin(\theta_{13})$. The error ellipses for the mixing angles are roughly derived from the 1σ to 3σ ranges in [62].

IV. MODIFYING THE FLAVON VACUUM ALIGNMENT

In this section, we consider the idea of changing the alignments of the flavon VEVs to obtain a nonzero θ_{13} . In particular, we focus on flavons associated with the charged lepton masses, and show that the effects on θ_{13} are enhanced by a factor that scales with m_τ/m_μ . We assume that the corrections discussed in the previous section are not important, e.g. when ϵ is extremely small², so that $M_l \propto A$. With a modified alignment, A is no longer given by Eq. (18). In particular, it is not of the form M_{aligned} and hence a nonzero θ_{13} can be generated. We do not attempt to explain the origin of the modified alignment, and will just focus on the consequence of such a modification.

In general, the relative alignments between all the flavons can be varied. However, we can illustrate most of the important features by just varying the alignment of $\langle\phi\rangle$:

$$\langle\phi\rangle = \sqrt{3}v \begin{pmatrix} \sin(a) \cos(b) \\ \sin(a) \sin(b) \\ \cos(a) \end{pmatrix}. \quad (35)$$

We recover the original alignment when $a = \arcsin(\sqrt{2/3})$ and $b = \frac{\pi}{4}$. With this alignment we

² Actually even for very small ϵ , the zeroth-order approximation is really $M_l = \frac{v_H}{\Lambda} \left(A - \frac{y_m^T}{y_r^T} C \right)$. However, since A and C are of the same form, this is equivalent to a different choice of Yukawas for A in $M_l = \frac{v_H}{\Lambda} A$. Henceforth, for notational simplicity, we ignore the C correction.

now have

$$A = \sqrt{3} \begin{pmatrix} y_e v s_a c_b & y_m v s_a c_b + \frac{y_m^5 v_5}{\sqrt{3}}(\omega^2 - \omega) & y_m v s_a c_b + \frac{y_m^5 v_5}{\sqrt{3}}(\omega - \omega^2) \\ y_e v s_a s_b & y_m v s_a s_b \omega + \frac{y_m^5 v_5}{\sqrt{3}}(1 - \omega^2) & y_m v s_a s_b \omega^2 + \frac{y_m^5 v_5}{\sqrt{3}}(1 - \omega) \\ y_e v c_a & y_m v c_a \omega^2 + \frac{y_m^5 v_5}{\sqrt{3}}(\omega - 1) & y_m v c_a \omega + \frac{y_m^5 v_5}{\sqrt{3}}(\omega^2 - 1), \end{pmatrix} \quad (36)$$

where $s_x \equiv \sin x$ and $c_x \equiv \cos x$. The angle between the original and modified alignment, which we denote as χ , can be thought of as the small parameter in this approach. At first glance, we might expect the size of θ_{13} to be given by χ . However, the near-degeneracy of m_e and m_μ relative to m_τ again comes into play, and so θ_{13} is amplified by a factor of $O(m_\tau/m_\mu)$. As discussed in Sec. III B, the amplification is not $O(m_\tau^2/m_\mu^2)$ since $M_l(M_l)^\dagger$ is obviously factorizable at a well-defined order in χ .

As an aside, it is not particularly difficult to perform a parameter scan to find values of VEVs, Yukawas and alignment angles that generate the correct mass spectrum and mixing angles. A useful observation is that the relation

$$m_e^2 + m_\nu^2 + m_\tau^2 = \text{Tr} [M_l(M_l)^\dagger] = 3 \frac{v_H^2 v^2}{\Lambda^2} (|y_e|^2 + 2|y_m|^2) + 18 \frac{v_H^2 v_5^2}{\Lambda^2} |y_m^5|^2, \quad (37)$$

is independent of the alignment, hence allowing us to reduce the number of parameters by one. However, such a scan is not very useful, since the effects of block diagonalization discussed in the previous section are expected to be significant given the constraints on the hierarchy of energy scales. Therefore, we will only focus on demonstrating the enhancement effects.

We compute the tree-level U_{PMNS} for four large collections of random parameter sets \mathcal{C}_3 , \mathcal{C}_4 , \mathcal{C}_5 and \mathcal{C}_6 . Details of their generation are provided in Appendix C 2. The collections differ in the conditions imposed on the ratio of mass eigenvalues: No conditions have been imposed on \mathcal{C}_3 , so the mass eigenvalues are typically of the same order, while the correct mass ratios have been imposed on \mathcal{C}_4 . The conditions on mass ratios m_μ^2/m_τ^2 and m_e^2/m_τ^2 have been (unphysically) modified to be 10 times smaller in \mathcal{C}_5 , and 100 times smaller in \mathcal{C}_6 . In other words, the mass ratio m_τ/m_μ relevant to the enhancement effect is larger in \mathcal{C}_5 and even more so in \mathcal{C}_6 .

Figure 3 shows the graphs of $\sin(\theta_{13})$ against χ for all the four collections. Just as in Section III C, we observe an enhancement effect in \mathcal{C}_4 relative in \mathcal{C}_3 , although it is smaller than the predicted enhancement of m_τ/m_μ due to large $O(1)$ factors that we have not taken into account. Nonetheless, the graphs for \mathcal{C}_5 and \mathcal{C}_6 clearly demonstrate that the enhancement factor scales as m_τ/m_μ , in agreement with our predictions.

To conclude, we have demonstrated that modifying the alignment of flavons associated with charged lepton masses give rise to a nonzero θ_{13} , with an enhancement factor that scales as m_τ/m_μ .




FIG. 3. Graphs of $\sin(\theta_{13})$ against the change in flavon alignment χ for collections (a) \mathcal{C}_3 , (b) \mathcal{C}_4 , (c) \mathcal{C}_5 and (d) \mathcal{C}_6 . The collections differ in the conditions imposed on the ratio of mass eigenvalues.

This enhancement may be applicable to a large class of A_4 models since the only feature we have alluded to beyond the minimal A_4 model are the additional mass terms involving v_5 , which can be easily reproduced with an additional A_4 flavon.

V. DISCUSSION AND CONCLUSION.

Having discussed the two approaches of obtaining a nonzero θ_{13} , there remains various issues that we did not touch on and are worth further investigation. First, we have not addressed one shortcoming of the model originally mentioned in [79]. This is the issue of Goldstone bosons when the global $SO(3)_F$ symmetry is broken, and the issue of mixed anomalies involving $U(1)_Y$ should we gauge the $SO(3)_F$ symmetry to eat these Goldstone bosons. However, the variety of modified models in Appendix A suggests that it should be possible to introduce additional heavy leptons to

address the issue of anomalies, and yet suppress their mass couplings to the existing leptons using auxiliary symmetries.

Second, our analysis so far is only at the classical level. We have yet to consider the running of parameters down to the electroweak scale [73–77, 81–88]. Nonetheless, since our neutrino mass spectrum is not quasi-degenerate, the classical results should hold as a first approximation.

Last is the issue of fine-tuning of the charged-lepton masses. In the minimal A_4 model, this can be resolved by modifying the model to give naturally small electron Yukawas. In the $SO(3)_F \rightarrow A_4$ model however, suppressing particular Yukawas do not guarantee the correct mass hierarchy, since the subleading corrections from block diagonalization can significantly affect the small eigenvalues. Still, we have demonstrated with our simulation in Section III C that small electron masses can be achieved, although the small fraction of successful parameter sets imply that specific relations between the Yukawas are required. Unfortunately, the exact forms of these relations are far from obvious, hence obscuring any UV explanation of the fine-tuning.

To conclude, the $SO(3)_F \rightarrow A_4$ model of [79] is the UV completion of an effective A_4 model with the purpose of reproducing the tri-bimaximal mixing pattern in U_{PMNS} . However, due to mixing between heavy and SM charged leptons, we find that the model actually predicts a nonzero θ_{13} , with the size of θ_{13} a measure of the ratio of the A_4 -breaking to $SO(3)_F$ -breaking scales. We have also shown that this model can reproduce both the measured light lepton spectrum and the mixing angles, and is hence compatible with experimental observations. Nonetheless, there exists various unattractive aspects of the model, in particular the fine-tuning of the charged-lepton eigenvalues and the need of an auxiliary symmetry on top of the original $SO(3)_F$ symmetry. We hope to resolve these issues in a future work.

ACKNOWLEDGMENTS

We thank Josh Berger and Jeff Dror for helpful discussions. YG is a Weston Visiting Professor at the Weizmann Institute. This work was partially supported by a grant from the Simons Foundation (#267432 to Yuval Grossman). The work of YG is supported in part by the U.S. National Science Foundation through grant PHY-0757868 and by the United States-Israel Binational Science Foundation (BSF) under grant No. 2010221.

Appendix A: Modified models with similar SM lepton phenomenology

1. Overview

As pointed out in Section IIB, there are two issues with the Lagrangian given by Eq. (2) and (3). First, it is not the most general one consistent with $SU(2)_L \times U(1)_Y$ gauge and global $SO(3)_F$ symmetries. Second, it is not clear whether the truncation of the Lagrangian is consistent with our hierarchy of scales. As an example, we have omitted dimension-six terms like $-\overline{\psi}_l^a \psi_m^{bc} \epsilon^{bde} H T^{adf} T^{cef} / \Lambda^2$ while keeping dimension-five terms like $-\overline{\psi}_l^a \psi_m^{ab} H \phi^b / \Lambda$, both of which contribute to the Dirac mass of ψ_l by an amount $\sim v_H v_T^2 / \Lambda^2$ and $\sim v_H v / \Lambda$ respectively. However, since $\{v, v_5, v'\} \ll v_T \ll \Lambda$, it is not immediately clear that the former is necessarily smaller than the latter, unless we impose the additional restriction that $v_T / \Lambda \ll \{v, v_5, v'\} / v_T$.

It turns out that both issues can be addressed if we modify the model to include an auxiliary Z_n symmetry and a Z_n flavon S . The modified model is designed to reproduce the same lepton mass matrices as the original Lagrangian. The auxiliary Z_n symmetry forbids lower-dimension terms otherwise allowed by the gauge and $SO(3)_F$ symmetry that would have changed the mass matrices, as well as certain higher-dimension terms (such as those quadratic in T) that if neglected, would have led to large truncation errors. The flavon S is a gauge and $SO(3)_F$ singlet, the VEV of which is related to the neutrino Majorana mass parameter M .

Two versions of modified models are discussed below, the main difference being the effective size of y_ν in the original Lagrangian, and hence the neutrino seesaw scale.

2. Model 1: Z_8 , with typical seesaw scale

We assign the following Z_8 representations to the matter fields:

Field	ψ_l	ψ_f	ψ_e	ψ_m	ψ_n	H	ϕ	ϕ'	ϕ_5	S	T
Z_8 rep.	$e^{i\frac{\pi}{4}}$	$e^{i\frac{\pi}{4}}$	$e^{-i\frac{3\pi}{4}}$	$e^{-i\frac{3\pi}{4}}$	$e^{i\frac{\pi}{4}}$	+1	-1	+i	-1	-i	-1

Note that ϕ' and S have to be complex fields since they are in complex Z_8 representations.

For the charged-lepton sector, since Dirac masses for ψ_f^- and ψ_l^- are generated by operators that are at least dimension-four and five respectively, with the latter always requiring a Higgs field, it is natural to use these minimum criteria as the truncation scheme. The most general Lagrangian turns out to be same as the original \mathcal{L}_e given in Eq. (2). The higher-dimension terms we have

neglected are given heuristically (coefficients and $SO(3)$ indices suppressed) by

$$\begin{aligned}\mathcal{L}_e^{\text{h.o.}} \sim & -\frac{1}{\Lambda} (\overline{\psi}_f \psi_e \phi' S^* + \overline{\psi}_f \psi_m \phi' \phi' + \dots) \\ & -\frac{1}{\Lambda^2} (\overline{\psi}_l \psi_e H \phi' S^* + \overline{\psi}_l \psi_m H \phi' \phi' + \dots + \overline{\psi}_f \psi_e TTT + \overline{\psi}_f \psi_m TTT + \dots) \\ & -\frac{1}{\Lambda^3} (\overline{\psi}_l \psi_e HTTT + \overline{\psi}_l \psi_m HTTT + \dots) - \dots + \text{h.c.}\end{aligned}\quad (\text{A1})$$

We note that terms like $-\overline{\psi}_f \psi_m TT/\Lambda$ that may lead to large truncation errors (if neglected) are explicitly forbidden by the Z_8 symmetry.

We now discuss the neutrino sector. Neutrino Dirac masses are generated by operators that are at least dimension four and always require a Higgs field. While neutrino Majorana masses can be generated by dimension-four operators, we also allow dimension-five operators that can potentially give comparable contributions as a result of the hierarchy of scales. With the above as the truncation scheme, the neutrino Lagrangian is then given by $\mathcal{L}_\nu^{\text{l.o.}} = \mathcal{L}_\nu^{\text{l.o.}(\text{old})} + \mathcal{L}_\nu^{\text{l.o.}(\text{new})}$, where

$$\begin{aligned}\mathcal{L}_\nu^{\text{l.o.}(\text{old})} &= -x_\nu^S S \overline{\psi}_n^c \psi_n^a - x_\nu \frac{1}{\Lambda} \overline{\psi}_n^c \psi_n^b \phi'^c T^{abc} - y_\nu \overline{\psi}_l^a \tilde{H} \psi_n^a + \text{h.c.} \\ \mathcal{L}_\nu^{\text{l.o.}(\text{new})} &\sim -\frac{1}{\Lambda} (\overline{\psi}_n^c \psi_n \phi' \phi + \overline{\psi}_n^c \psi_n \phi' \phi_5) + \text{h.c.}\end{aligned}\quad (\text{A2})$$

The higher dimension terms that we have neglected are

$$\mathcal{L}_\nu^{\text{h.o.}} \sim -\frac{1}{\Lambda^2} (\overline{\psi}_l \psi_n HTT + \dots + \overline{\psi}_n \psi_n STT + \dots) - \dots + \text{h.c.}\quad (\text{A3})$$

When the flavon S gains a VEV v_S , if we identify $x_\nu^S v_S$ with M , $\mathcal{L}_\nu^{\text{l.o.}(\text{old})}$ then generates the same neutrino mass matrix as the original Lagrangian. Therefore, the largest contributions that we have omitted from our original mass matrix come from $\mathcal{L}_\nu^{\text{l.o.}(\text{new})}$ and $\mathcal{L}_\nu^{\text{h.o.}}$. Note that $\mathcal{L}_\nu^{\text{l.o.}(\text{old})}$ cannot be eliminated through a different implementation of auxiliary symmetries without significantly modifying the charged-lepton mass matrix.

We now analyze the fractional errors in both the charged lepton and neutrino mass matrices as a result of the various omitted contributions discussed above. For simplicity, we assume that all the Yukawas of terms that contribute to the same mass type, omitted or otherwise, are of the same order. As a result, the Yukawas (heuristically denoted as y) cancel out in the fractional errors, which we summarize in the table below. Note that we have defined $\epsilon_T \equiv v_T/\Lambda$.

Mass types	Smallest contributions included	Largest contributions omitted	Fractional error
ψ_l^- , Dirac	$y \frac{v_H v}{\Lambda}$	$\sup\{y \frac{v_H v^2}{\Lambda^2}, y \frac{v_H v_T^3}{\Lambda^3}\}$	$\sup\{\epsilon_T, \frac{\epsilon_T^2}{\epsilon}\}$
ψ_f^- , Dirac	$y v$	$\sup\{y \frac{v^2}{\Lambda}, y \frac{v_T^3}{\Lambda^2}\}$	$\sup\{\epsilon_T, \frac{\epsilon_T^2}{\epsilon}\}$
Neutrino, Dirac	$y v_H$	$y \frac{v_H v_T^2}{\Lambda^2}$	ϵ_T^2
Neutrino, Majorana	$y \frac{v v_T}{\Lambda}$	$y \frac{v^2}{\Lambda}$	ϵ

We want all fractional errors to be smaller than ϵ so that the omitted contributions generate smaller corrections to θ_{13} than what we have discussed in Section III. Except for neutrino Majorana masses, this can be achieved by choosing a hierarchy where $\epsilon_T < \epsilon$. For neutrino Majorana masses, fine-tuning may be required to suppress the Yukawas associated with the omitted contributions to reduce the fractional errors. We have not taken into account any enhancement effects which may ameliorate or exacerbate the fine-tuning.

3. Model 2: Z_8 , with lower seesaw scale

In this model, we assign different Z_8 representations to the matter fields.

Field	ψ_l	ψ_f	ψ_e	ψ_m	ψ_n	H	ϕ	ϕ'	ϕ_5	S	T
Z_8 rep.	-1	-1	+1	+1	$e^{i\frac{\pi}{4}}$	+1	-1	+i	-1	$e^{i\frac{3\pi}{4}}$	-1

Again, we note that ϕ' and S have to be complex fields. The charged-lepton Lagrangian is the same as the one in the previous model. For the neutrino sector, since neutrino Dirac masses now only arise at dimension five, the truncation scheme is modified accordingly. The neutrino Lagrangian is given by $\mathcal{L}_\nu^{\text{l.o.}} = \mathcal{L}_\nu^{\text{l.o.}(\text{old})} + \mathcal{L}_\nu^{\text{l.o.}(\text{new})}$, where

$$\begin{aligned}\mathcal{L}_\nu^{\text{l.o.}(\text{old})} &= -x_\nu^S \frac{1}{\Lambda} S^2 \bar{\psi}_n^c{}^a \psi_n^a - x_\nu \frac{1}{\Lambda} \bar{\psi}_n^c{}^a \psi_n^b \phi'^c T^{abc} - y'_\nu \frac{1}{\Lambda} S \bar{\psi}_l^a \tilde{H} \psi_n^a \\ \mathcal{L}_\nu^{\text{l.o.}(\text{new})} &\sim -\frac{1}{\Lambda} (\bar{\psi}_n^c \psi_n \phi' \phi + \bar{\psi}_n^c \psi_n \phi' \phi_5) + \text{h.c.}\end{aligned}\tag{A4}$$

The higher dimension terms that we have neglected are

$$\mathcal{L}_\nu^{\text{h.o.}} \sim -\frac{1}{\Lambda^2} (\bar{\psi}_n^c \psi_n S^* S^* T + \bar{\psi}_n^c \psi_n \phi'^* T T + \dots) - \frac{1}{\Lambda^3} (\bar{\psi}_l \psi_n H S T T + \dots) - \dots + \text{h.c.}\tag{A5}$$

When the flavon S gains a VEV v_S , if we identify $x_\nu^S v_S^2/\Lambda$ with M and $y'_\nu v_S/\Lambda$ with y_ν , $\mathcal{L}_\nu^{\text{l.o.}(\text{old})}$ then generates the same neutrino mass matrix as the original Lagrangian, but with y_ν naturally suppressed by v_S/Λ . This allows for a seesaw scale M that is roughly two orders of magnitude lower than usual.

The fractional errors for the different mass types are given in the table below. Again the preferred hierarchy is one where $\epsilon_T < \epsilon$, and fine-tuning is still required to suppress the Yukawas associated with neutrino Majorana mass contributions that have been omitted.

Mass types	Smallest contributions included	Largest contributions omitted	Fractional error
ψ_l^- , Dirac	$\frac{v_H v}{\Lambda}$	$\sup\{\frac{v_H v^2}{\Lambda^2}, \frac{v_H v_T^3}{\Lambda^3}\}$	$\sup\{\epsilon\epsilon_T, \frac{\epsilon_T^2}{\epsilon}\}$
ψ_f^- , Dirac	v	$\sup\{\frac{v^2}{\Lambda}, \frac{v_T^3}{\Lambda^2}\}$	$\sup\{\epsilon\epsilon_T, \frac{\epsilon_T^2}{\epsilon}\}$
Neutrino, Dirac	$\frac{v_H \sqrt{v v_T}}{\Lambda}$	$\frac{v_H \sqrt{v v_T} v_T^2}{\Lambda^3}$	ϵ_T^2
Neutrino, Majorana	$\frac{v v_T}{\Lambda}$	$\sup\{\frac{v^2}{\Lambda}, \frac{v v_T^2}{\Lambda^2}\}$	$\sup\{\epsilon, \epsilon_T\}$

Appendix B: Nonunitary factors in U_{PMNS}

In this Appendix, we discuss the origin of nonunitary factors mentioned in Section II and why they turn out to be negligible. The charged current weak interaction acts between the left-handed SM charged leptons and neutrinos, both of which are linear combinations of light and heavy mass eigenstates. U_{PMNS} characterizes this interaction between only the light mass eigenstates.

Define 6×6 unitary matrices $U_{l,\text{full}}^{6 \times 6}$ and $U_{\nu,\text{full}}^{6 \times 6}$ that are required to fully diagonalize $M_l^{6 \times 6} (M_l^{6 \times 6})^\dagger$ and $M_\nu^{6 \times 6}$:

$$U_{\nu,\text{full}}^{6 \times 6} M_\nu^{6 \times 6} (U_{\nu,\text{full}}^{6 \times 6})^T = \begin{pmatrix} m_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & m_2 & 0 & 0 & 0 & 0 \\ 0 & 0 & m_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & m_{1'} & 0 & 0 \\ 0 & 0 & 0 & 0 & m_{2'} & 0 \\ 0 & 0 & 0 & 0 & 0 & m_{3'} \end{pmatrix}, \quad (\text{B1})$$

$$U_{l,\text{full}}^{6 \times 6} M_l^{6 \times 6} (M_l^{6 \times 6})^\dagger (U_{l,\text{full}}^{6 \times 6})^\dagger = \begin{pmatrix} m_e^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & m_\mu^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & m_\tau^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & m_{e'}^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & m_{\mu'}^2 & 0 \\ 0 & 0 & 0 & 0 & 0 & m_{\tau'}^2 \end{pmatrix}, \quad (\text{B2})$$

where ' indicates a heavy lepton. We can write $U_{l,\text{full}}^{6 \times 6}$ and $U_{\nu,\text{full}}^{6 \times 6}$ in terms of 3×3 blocks as shown here:

$$U_{l,\text{full}}^{6 \times 6} = \begin{pmatrix} U_{l,\text{full}} & U'_{l,\text{full}} \\ U''_{l,\text{full}} & U'''_{l,\text{full}} \end{pmatrix}, \quad U_{\nu,\text{full}}^{6 \times 6} = \begin{pmatrix} U_{\nu,\text{full}} & U'_{\nu,\text{full}} \\ U''_{\nu,\text{full}} & U'''_{\nu,\text{full}} \end{pmatrix}. \quad (\text{B3})$$

U_{PMNS} is then given by

$$U_{\text{PMNS}} = U_{l,\text{full}} (U_{\nu,\text{full}})^\dagger, \quad (\text{B4})$$

Since the 3×3 blocks are nonunitary in general, we expect the same for U_{PMNS} .

It is perhaps more illustrative to regard the diagonalization as a two-step process, which we demonstrate here with the neutrino sector. Define a 6×6 unitary matrix $U_{\nu,\text{bd}}^{6 \times 6}$ that is required to block-diagonalize $M_\nu^{6 \times 6}$:

$$U_{\nu,\text{bd}}^{6 \times 6} M_\nu^{6 \times 6} (U_{\nu,\text{bd}}^{6 \times 6})^T = \begin{pmatrix} M_\nu & 0 \\ 0 & M_{\nu'} \end{pmatrix}, \quad (\text{B5})$$

where M_ν and $M_{\nu'}$ are the 3×3 Majorana mass matrices for the light and heavy neutrinos. Again we can write $U_{\nu,\text{bd}}^{6 \times 6}$ in terms of 3×3 blocks:

$$U_{\nu,\text{bd}}^{6 \times 6} = \begin{pmatrix} U_{\nu,\text{bd}} & U'_{\nu,\text{bd}} \\ U''_{\nu,\text{bd}} & U'''_{\nu,\text{bd}} \end{pmatrix}. \quad (\text{B6})$$

Let U_ν be the 3×3 unitary matrix required to diagonalize M_ν :

$$U_\nu M_\nu (U_\nu)^T = \begin{pmatrix} m_1 & 0 & 0 \\ 0 & m_2 & 0 \\ 0 & 0 & m_3 \end{pmatrix}, \quad (\text{B7})$$

We can then show that

$$U_{\nu,\text{full}} = U_\nu U_{\nu,\text{bd}}. \quad (\text{B8})$$

In other words, $U_{\nu,\text{full}}$ can be decomposed into a unitary factor associated with the diagonalization of M_ν , and a non-unitary factor associated with the block-diagonalization of $M_\nu^{6 \times 6}$. Applying a similar two-step process to the charged lepton sector gives us the factorization

$$U_{l,\text{full}} = U_l U_{l,\text{bd}}. \quad (\text{B9})$$

U_{PMNS} is then given by

$$U_{\text{PMNS}} = U_l U_{l,\text{bd}} (U_{\nu,\text{bd}})^\dagger (U_\nu)^\dagger. \quad (\text{B10})$$

This expression differs from Eq. (13) by the nonunitary factor $U_{l,\text{bd}} (U_{\nu,\text{bd}})^\dagger$ associated with the block-diagonalization process. However, we can show that $U_{l,\text{bd}}$ and $(U_{\nu,\text{bd}})^\dagger$ deviate from the identity matrix by terms of order $O(\frac{v_H^2}{\Lambda^2})$ and $O(\frac{v_H^2}{M^2})$ respectively. Based on the energy scales in Eq. (16), these are exceedingly small deviations. Hence, their effects on U_{PMNS} are negligible and U_{PMNS} can be considered to be unitary.

Appendix C: Generation of random parameter sets

In this appendix, we discuss how we generate the various collections of random parameter sets used in the simulations.

1. \mathcal{C}_1 and \mathcal{C}_2

The collections \mathcal{C}_1 and \mathcal{C}_2 are used in Figure 1. In each collection, the VEV v_T is a log flat random variable between 10^{16} and 10^{19} GeV, while v and v_5 are uniform random variables between 10^{15} and 10^{16} GeV.

In \mathcal{C}_1 , which consists of 20,000 sets, all eight charged-lepton Yukawas are simply $O(1)$ uniform random complex variables, with real and imaginary parts between -3 and 3 . In \mathcal{C}_2 , we want to restrict the parameter sets to only those that produce the correct charged-lepton mass ratios. Ideally, we would like to use the same definitions of random variables as \mathcal{C}_1 , and simply reject the parameter sets that fail the cut. However, the very small measure of the allowed parameter space makes this computationally prohibitive, so we instead adopt an alternative procedure for \mathcal{C}_2 which we outline below.

First, we define two new uniform random complex variables α_1 and α_2 of magnitudes $O(\frac{1}{1000})$ and $O(\frac{1}{10})$ respectively, that satisfy the relations

$$y'_e = \frac{y_m^{T'}}{y_m^T} (y_e - \alpha_1), \quad (\text{C1})$$

$$y_m^{5'} = \frac{y_m^{T'}}{y_m^T} y_m^5 + \frac{y_m^{T'}}{y_m^T} \frac{i}{\sqrt{3}} \frac{v}{v_5} \left(y_m - \frac{y_m^T}{y_m^{T'}} y'_m \right) (1 - \alpha_2) \quad (\text{C2})$$

Instead of generating all eight charged-lepton Yukawas randomly, we now generate only six of them (excluding y'_e and $y_m^{5'}$), together with α_1 and α_2 . y'_e and $y_m^{5'}$ are then obtained from the relations above. Since we still want all Yukawas to be $O(1)$, we discard the parameter set should the resulting y'_e and $y_m^{5'}$ not be $O(1)$. We also discard parameter sets where the mass spectra do not satisfy $10^{-3} \leq m_\mu^2/m_\tau^2 \leq 10^{-2}$ and $10^{-8} \leq m_e^2/m_\tau^2 \leq 10^{-6}$. Only parameter sets that satisfy both conditions are included in \mathcal{C}_2 .

Second, we notice that parameter sets that satisfy the conditions tend to be concentrated around very small $\sup\{v/v_T, v_5/v_T\}$. Since we want to study θ_{13} over a large range of ϵ , we generate 10,000 parameter sets (satisfying the conditions) for v_T a log flat random variable between 10^{16} and 10^{19} GeV, 6,000 sets for v_T between 10^{16} and 10^{18} GeV and 4,000 sets for v_T between 10^{16} and $10^{17.5}$ GeV. This ensures that the combined 20,000 sets in \mathcal{C}_2 span a useful range of

$\sup\{v/v_T, v_5/v_T\}$ that we can work with.

Finally, we explain the motivation behind Eq. (C1) and (C2). From Eq. (25), the zeroth-order term of M_l is $\frac{v_H}{\Lambda} \left(A - \frac{y_m^T}{y_m^{T'}} C \right)$. This has eigenvalues

$$m_a = \sqrt{3} \frac{v_H}{\Lambda} \left| y_e v - \frac{y_m^T}{y_m^{T'}} y_e' v \right|, \quad m_b, m_c = \frac{v_H}{\Lambda} \left| \sqrt{3} \left(y_m - \frac{y_m^T}{y_m^{T'}} y_m' \right) v \pm 3i \left(y_m^5 - \frac{y_m^T}{y_m^{T'}} y_m^{5'} \right) v_5 \right|. \quad (\text{C3})$$

Eq. (C1) and (C2) hence ensures that the zeroth-order eigenvalues satisfy the mass ratio conditions. Nonetheless, we note that only a very small fraction of random parameter sets generated this way end up being included in \mathcal{C}_2 . The reason is that subleading corrections to the small eigenvalues from block diagonalisation can be much larger than the zeroth-order small eigenvalues themselves, especially for larger values of $\sup\{v/v_T, v_5/v_T\}$. As a result, the mass ratio conditions may be violated.

2. $\mathcal{C}_3, \mathcal{C}_4, \mathcal{C}_5$ and \mathcal{C}_6

The collections $\mathcal{C}_3, \mathcal{C}_4, \mathcal{C}_5$ and \mathcal{C}_6 are used in Figure 3. The random parameters of interest are the VEVs v and v_5 , the Yukawas y_e, y_m and y_m^5 , and the changes δa and δb from the original values of a and b . In all four collections, the VEVs are uniform random variables between 10^{15} and 10^{16} GeV. For δa and δb , we first simulate the deviation angle χ as log flat random variable between 10^{-4} and 10^{-1} . Since $\chi^2 \simeq (\delta a)^2 + 2(\delta b)^2/3$, we simulate δa as a uniform random variable between $-\chi$ and χ , and then derive δb using $\delta b = \pm \sqrt{3(\chi^2 - (\delta a)^2)/2}$, with the signs randomly generated.

The difference between the four collections lie in the Yukawas, since the size of Yukawas is directly related to the size of the mass eigenvalues. For \mathcal{C}_3 , all Yukawas are simply $O(1)$ uniform random complex variables, with real and imaginary parts between -3 and 3 . For \mathcal{C}_4 , we generate y_e and y_m as $O(\frac{1}{1000})$ and $O(1)$ random complex variables. We also first generate a $O(0.1)$ random complex variable α , from which y_m^5 is derived using the relation

$$y_m^5 = -\frac{i}{\sqrt{3}} \frac{v}{v_5} y_m (1 - \alpha) \quad (\text{C4})$$

These choices are made to increase the likelihood of the eigenvalues satisfying the correct mass ratio. For \mathcal{C}_5 and \mathcal{C}_6 , the procedures are similar to that of \mathcal{C}_4 , except that y_e and α are further reduced to increase the likelihood of satisfying the (unphysical) smaller mass ratios.

[1] B. Pontecorvo, Sov. Phys. JETP **7**, 172 (1958).

- [2] Z. Maki, M. Nakagawa, and S. Sakata, Prog. Theo. Phys. **28**, 870 (1962).
- [3] P. Harrison, D. Perkins, and W. Scott, Phys. Lett. **B530**, 167 (2002), arXiv:hep-ph/0202074.
- [4] E. Ma and G. Rajasekaran, Phys. Rev. D **64**, 113012 (2001), arXiv:hep-ph/0106291.
- [5] E. Ma, Mod. Phys. Lett. A **17**, 627 (2002), arXiv:hep-ph/0203238.
- [6] K. S. Babu, E. Ma, and J. W. F. Valle, Phys. Lett. B **552**, 207 (2003), arXiv:hep-ph/0206292.
- [7] M. Hirsch *et al.*, (2003), arXiv:hep-ph/0312244.
- [8] M. Hirsch *et al.*, Phys. Rev. D **69**, 093006 (2004), arXiv:hep-ph/0312265.
- [9] E. Ma, Phys. Rev. D **70**, 031901 (2004), arXiv:hep-ph/0404199.
- [10] E. Ma, New J. Phys. **6**, 104 (2004), arXiv:hep-ph/0405152.
- [11] E. Ma, (2004), arXiv:hep-ph/0409075.
- [12] S. L. Chen, M. Frigerio, and E. Ma, Nucl. Phys. B **724**, 423 (2005), arXiv:hep-ph/0504181.
- [13] E. Ma, Phys. Rev. D **72**, 037301 (2005), arXiv:hep-ph/0505209.
- [14] M. Hirsch *et al.*, Phys. Rev. D **72**, 091301 (2005), arXiv:hep-ph/0507148.
- [15] K. S. Babu and X. G. He, (2005), arXiv:hep-ph/0507217.
- [16] E. Ma, Mod. Phys. Lett. A **20**, 2601 (2005), arXiv:hep-ph/0508099.
- [17] A. Zee, Phys. Lett. B **630**, 58 (2005), arXiv:hep-ph/0508278.
- [18] X. G. He, Y. Y. Keum, and R. R. Volkas, JHEP **0604**, 039 (2006), arXiv:hep-ph/0601001.
- [19] B. Adhikary *et al.*, Phys. Lett. B **638**, 345 (2006), arXiv:hep-ph/0603059.
- [20] L. Lavoura and H. Kuhbock, Mod. Phys. Lett. A **22**, 181 (2007), arXiv:hep-ph/0610050.
- [21] S. F. King and M. Malinsky, Phys. Lett. B **645**, 351 (2007), arXiv:hep-ph/0610250.
- [22] S. Morisi, M. Picariello, and E. Torrente-Lujan, Phys. Rev. D **75**, 075015 (2007), arXiv:hep-ph/0702034.
- [23] M. Hirsch *et al.*, Phys. Rev. Lett. **99**, 151802 (2007), arXiv:hep-ph/0703046.
- [24] F. Yin, Phys. Rev. D **75**, 073010 (2007), arXiv:0704.3827 [hep-ph].
- [25] M. Honda and M. Tanimoto, Prog. Theo. Phys. **119**, 583 (2008), arXiv:0801.0181 [hep-ph].
- [26] B. Brahmachari, S. Choubey, and M. Mitra, Phys. Rev. D **77**, 073008 (2008), arXiv:0801.3554 [hep-ph].
- [27] G. Altarelli, F. Feruglio, and C. Hagedorn, JHEP **0803**, 052 (2008), arXiv:0802.0090 [hep-ph].
- [28] B. Adhikary and A. Ghosal, Phys. Rev. D **78**, 073007 (2008), arXiv:0803.3582 [hep-ph].
- [29] M. Hirsch, S. Morisi, and J. W. F. Valle, Phys. Rev. D **78**, 093007 (2008), arXiv:0804.1521 [hep-ph].
- [30] Y. Lin, Nucl. Phys. B **813**, 025 (2008), arXiv:0804.2867 [hep-ph].
- [31] F. Feruglio *et al.*, Nucl. Phys. B **809**, 218 (2009), arXiv:0807.3160 [hep-ph].
- [32] F. Bazzocchi, M. Frigerio, and S. Morisi, Phys. Rev. D **78**, 116018 (2008), arXiv:0809.3573 [hep-ph].
- [33] S. Morisi, Phys. Rev. D **79**, 033008 (2009), arXiv:0901.1080 [hep-ph].
- [34] P. Ciafaloni *et al.*, Phys. Rev. D **79**, 116010 (2009), arXiv:0901.2236 [hep-ph].
- [35] M. C. Chen and S. King, JHEP **0906**, 072 (2009), arXiv:0903.0125 [hep-ph].
- [36] Y. Lin, Phys. Rev. D **80**, 076011 (2009), arXiv:0903.0831 [hep-ph].
- [37] G. C. Branco *et al.*, Phys. Rev. D **79**, 093008 (2009), arXiv:0904.3076 [hep-ph].

- [38] T. Araki, J. Mei, and Z. Z. Xing, Phys. Lett. B **695**, 165 (2011), arXiv:1010.3065 [hep-ph].
- [39] D. Meloni, S. Morisi, and E. Peinado, Phys. Lett. B **697**, 339 (2011), arXiv:1011.1371 [hep-ph].
- [40] A. Adulpravitchai and R. Takahashi, JHEP **1109**, 127 (2011), arXiv:1107.3829 [hep-ph].
- [41] G. J. Ding and D. Meloni, Nucl. Phys. B **855**, 21 (2012), arXiv:1108.2733 [hep-ph].
- [42] Y. BenTov, X. G. He, and A. Zee, JHEP **1212**, 093 (2012), arXiv:1208.1062 [hep-ph].
- [43] M. Holthausen, M. Lindner, and M. A. Schmidt, Phys. Rev. D **87**, 033006 (2013), arXiv:1211.5143 [hep-ph].
- [44] R. G. Felipe, H. Serodio, and J. P. Silva, Phys. Rev. D **87**, 055010 (2013), arXiv:1302.0861 [hep-ph].
- [45] R. G. Felipe, H. Serodio, and J. P. Silva, Phys. Rev. D **88**, 015015 (2013), arXiv:1304.3468 [hep-ph].
- [46] D. V. Forero *et al.*, Phys. Rev. D **88**, 016003 (2013), arXiv:1305.6774 [hep-ph].
- [47] P. M. Ferreira, L. Lavoura, and P. O. Ludl, (2013), arXiv:1306.1500 [hep-ph].
- [48] G. Altarelli, (2009), arXiv:0905.2350 [hep-ph].
- [49] G. Altarelli and F. Feruglio, Rev. Mod. Phys. **82**, 2701 (2010), arXiv:1002.0211 [hep-ph].
- [50] G. Altarelli *et al.*, JHEP **1208**, 021 (2012), arXiv:1205.4670 [hep-ph].
- [51] G. Altarelli, F. Feruglio, and L. Merlo, (2012), arXiv:1205.5133 [hep-ph].
- [52] G. Altarelli and F. Feruglio, Nucl. Phys. B **720**, 64 (2005), arXiv:hep-ph/0504165.
- [53] E. Ma, Phys. Rev. D **73**, 057304 (2006), arXiv:hep-ph/0511133.
- [54] G. Altarelli and F. Feruglio, Nucl. Phys. B **741**, 215 (2006), arXiv:hep-ph/0512103.
- [55] E. Ma, Mod. Phys. Lett. A **22**, 101 (2007), arXiv:hep-ph/0610342.
- [56] F. Bazzocchi, S. Kaneko, and S. Morisi, JHEP **0803**, 063 (2008), arXiv:0707.3032 [hep-ph].
- [57] C. Csaki *et al.*, JHEP **0810**, 055 (2008), arXiv:0806.0356 [hep-ph].
- [58] P. H. Frampton and S. Matsuzaki, (2008), arXiv:0806.4592 [hep-ph].
- [59] G. Altarelli and D. Meloni, J. Phys. G **36**, 085005 (2009), arXiv:0905.0620 [hep-ph].
- [60] F. An *et al.* (Daya Bay), Phys. Rev. Lett. **108**, 171803 (2012), arXiv:1203.1669 [hep-ex].
- [61] J. Ahn *et al.* (RENO), Phys. Rev. Lett. **108**, 191802 (2012), arXiv:1204.0626 [hep-ex].
- [62] G. L. Fogli *et al.*, Phys. Rev. D **86**, 013012 (2012), arXiv:1205.5254 [hep-ph].
- [63] Y. Lin, Nucl. Phys. B **824**, 95 (2010), arXiv:0905.3534 [hep-ph].
- [64] Y. Shimizu, M. Tanimoto, and A. Watanabe, Prog. Theo. Phys. **126**, 81 (2011), arXiv:1105.2929 [hep-ph].
- [65] E. Ma and D. Wegman, Phys. Rev. Lett. **107**, 061803 (2011), arXiv:1106.4269 [hep-ph].
- [66] S. King and C. Luhn, JHEP **1109**, 042 (2011), arXiv:1107.5332 [hep-ph].
- [67] S. Antusch *et al.*, Nucl. Phys. B **856**, 328 (2012), arXiv:1108.4278 [hep-ph].
- [68] M. C. Chen *et al.*, JHEP **1302**, 021 (2013), arXiv:1210.6982 [hep-ph].
- [69] J. Barry and W. Rodejohann, Phys. Rev. D **81**, 119901(E) (2010), arXiv:1003.2385 [hep-ph].
- [70] S. King, JHEP **1101**, 115 (2011), arXiv:1011.6167 [hep-ph].
- [71] S. King and C. Luhn, JHEP **1203**, 036 (2012), arXiv:1112.1959 [hep-ph].
- [72] G. C. Branco *et al.*, Phys. Rev. D **86**, 076008 (2012), arXiv:1203.2646 [hep-ph].

- [73] S. Antusch *et al.*, Nucl. Phys. B **674**, 401 (2003), arXiv:hep-ph/0305273.
- [74] S. Antusch *et al.*, JHEP **0503**, 024 (2005), arXiv:hep-ph/0501272.
- [75] A. Dighe, S. Goswami, and W. Rodejohann, Phys. Rev. D **75**, 073023 (2007), arXiv:hep-ph/0612328.
- [76] A. Dighe, S. Goswami, and P. Roy, Phys. Rev. D **76**, 096005 (2007), arXiv:0704.3735 [hep-ph].
- [77] M. Borah, B. Sharma, and M. K. Das, (2013), arXiv:1304.0164 [hep-ph].
- [78] A. Adulpravitchai, A. Blum, and M. Lindner, JHEP **0909**, 018 (2009), arXiv:0907.2332 [hep-ph].
- [79] J. Berger and Y. Grossman, JHEP **1002**, 071 (2009), arXiv:0910.4392 [hep-ph].
- [80] C. Luhn, JHEP **1103**, 108 (2011), arXiv:1101.2417 [hep-ph].
- [81] P. H. Chankowski and Z. Pluciennik, Phys. Lett. B **316**, 312 (1993), arXiv:hep-ph/9306333.
- [82] K. S. Babu, C. N. Leung, and J. Pantaleone, Phys. Lett. B **319**, 191 (1993), arXiv:hep-ph/9309223.
- [83] P. H. Chankowski, W. Krolikowski, and S. Pokorski, Phys. Lett. B **473**, 109 (2000), arXiv:hep-ph/9910231.
- [84] S. F. King and N. N. Singh, Nucl. Phys. B **591**, 3 (2000), arXiv:hep-ph/0006229.
- [85] S. Antusch *et al.*, Phys. Lett. B **519**, 238 (2001), arXiv:hep-ph/0108005.
- [86] S. Antusch *et al.*, Phys. Lett. B **525**, 130 (2002), arXiv:hep-ph/0110366.
- [87] S. Antusch and M. Ratz, JHEP **0207**, 059 (2002), arXiv:hep-ph/0203027.
- [88] S. Antusch *et al.*, Phys. Lett. B **538**, 87 (2002), arXiv:hep-ph/0203233.