



This is the accepted manuscript made available via CHORUS. The article has been published as:

Third generation in cascade decays

Bhaskar Dutta, Tianjun Li, James A. Maxin, Dimitri V. Nanopoulos, Kuver Sinha, and Joel W. Walker

Phys. Rev. D **91**, 115021 — Published 25 June 2015

DOI: [10.1103/PhysRevD.91.115021](https://doi.org/10.1103/PhysRevD.91.115021)

Third Generation in Cascade Decays

Bhaskar Dutta,¹ Tianjun Li,^{2,3} James A. Maxin,⁴ Dimitri V. Nanopoulos,^{1,5,6} Kuver Sinha,⁷ and Joel W. Walker⁸

¹*George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy,
Texas A&M University, College Station, TX 77843, USA*

²*State Key Laboratory of Theoretical Physics and Kavli Institute for Theoretical Physics China (KITPC),
Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, P. R. China*

³*School of Physical Electronics, University of Electronic Science and Technology of China, Chengdu 610054, P. R. China*

⁴*Department of Physics and Engineering Physics,
The University of Tulsa, Tulsa, OK 74104 USA*

⁵*Astroparticle Physics Group, Houston Advanced Research Center (HARC), Mitchell Campus, Woodlands, TX 77381, USA*

⁶*Academy of Athens, Division of Natural Sciences,
28 Panepistimiou Avenue, Athens 10679, Greece*

⁷*Department of Physics, Syracuse University, Syracuse, NY 13244, USA*

⁸*Department of Physics, Sam Houston State University, Huntsville, TX 77341, USA*

In supersymmetric models with gluinos $m(\tilde{g}) \sim 1000 - 2000$ GeV, new physics searches based on cascade decay products of the gluino are viable at the next run of the LHC. We investigate a scenario where the light stop is lighter than the gluino and both are lighter than all other squarks, and show that its signal can be established using multi b -jet, multi W and/or multi lepton final state topologies. We then utilize both boosted and conventional jet topologies in the final state in conjunction with di-tau production as a probe of the stau-neutralino co-annihilation region responsible for the model's dark matter content. This study is performed in the specific context of one such phenomenologically viable model named No-Scale \mathcal{F} - $SU(5)$.

PACS numbers: 12.60.Jv, 13.85.Hd, 14.80.Ly, 14.60.Fg

I. INTRODUCTION

Weak-scale supersymmetry (SUSY) provides a leading candidate for physics beyond the Standard Model (SM), as it addresses the hierarchy problem, gives gauge coupling unification, and (in R -parity conserving models) provides a robust dark matter (DM) candidate. The search for colored superpartners at the Large Hadron Collider (LHC) has so far yielded null results. The exclusion limits on squark (\tilde{q}) and gluino (\tilde{g}) masses, when they are comparable, are approximately 1.5 TeV at 95% CL with 20 fb^{-1} of integrated luminosity [1–4]. If the squarks are heavy, the gluino mass is constrained to be heavier than approximately 1.2 TeV at 95% CL with 20 fb^{-1} of integrated luminosity [1–5].

There are many interesting scenarios where the squarks are heavy but the gluino mass is about a TeV or so. In this paper we probe one such scenario as a benchmark, namely No-Scale \mathcal{F} - $SU(5)$ [6, 7]. In this model, the Higgs mass receives supplementary contributions from additional TeV scale vector-like fields [8], which may make it easier to obtain a Higgs mass near 126 GeV without tuning in the stop sector. The light stop \tilde{t}_1 is lighter than the gluino in this model, and by an amount that allows for on-shell decays with unity branching ratio for most of the viable parameter space. Similar types of mass spectra may also occur in large volume scenario models [9], and can be arranged in the phenomenological minimal supersymmetric standard model [10]; light stops are also generically of interest with regards to the potential reconciliation of the observed Higgs mass with

modest levels of electroweak fine-tuning [11], specifically in the context of natural SUSY [12–14]. Once gluinos are produced, they may cascade via the stop into the (Wino-like) second lightest neutralino or light chargino, which then decay into the lightest neutralino via stau, e.g. $\tilde{\chi}_2^0 \rightarrow \tilde{\tau}_1 \tau \rightarrow \tau \tau \tilde{\chi}_1^0$. The potentially sizeable mass gap between the first and second neutralino will tend to impart a sizeable kinematic boost to the lead tau production, but the narrow conduit $\Delta M \equiv M_{\tilde{\tau}_1} - M_{\tilde{\chi}_1^0}$ linking the stau and lightest supersymmetric particle (LSP) may render the secondary tau rather soft, and hence more difficult to detect, barring inheritance of a substantive upstream boost. The dark matter content may be obtained thermally by stau and (Bino-like) light neutralino coannihilation in the case of a small $\Delta M \simeq 6$ GeV.

The final state of such a cascade contains b -quarks, τ leptons, missing transverse energy \cancel{E}_T , and also additional light flavor leptons. The scenario can be established by the presence of multiple b -quarks and leptons in the final state. Establishing the coannihilation element would require b 's and τ 's in the final state, where $t\bar{t}$ is the dominant background for such a signal, also containing b 's, τ 's and missing energy. In this paper we first try to establish the scenario from the multi- b , leptons and W 's, so that the existence of a third generation can be surmised. Similar final state event topologies have been studied recently at the LHC [5]. We then try to establish the signal in the coannihilation region by considering two analysis routes, each of which requires at least two τ 's with $P_T > (40, 20)$ GeV in $|\eta| < 2.5$ and large missing energy in the final state, but

with distinct handling of the jets. In the scenario that we will refer to as the “Boosted Event Topology”, two tagging jets $j_{1,2}$ with $P_T > (75, 50)$ GeV in $|\eta| \leq 5.0$ are required, with an opposite hemisphere orientation ($\eta_1 \times \eta_2 < 0$), and an absolute separation in pseudorapidity of $|\eta_2 - \eta_1| > 3.5$. In the scenario that we will refer to as the “Conventional Event Topology”, the two leading jets $j_{1,2}$ in $|\eta| < 2.5$ carry very large transverse momenta $P_T > (400, 200)$ GeV. No vetoes on heavy flavor jets or light lepton flavors are enforced. We will demonstrate the cut flow optimization for all the selections, and calculate the signal to background ratios S/B. A methodology using the boosted event topology has been used in the context of stop searches recently [15] and is found to be very effective for background reduction. A variation of conventional topology has been prescribed in [16] with the requirement that the jets have to be non- b jet to reduce the background. In our case study, however, the signal contains a large population of b -jets arising from decays of the on-shell stop, which makes controlling the $t\bar{t}$ background for establishing the coannihilation region more challenging.

The structure of the paper is as follows. In Section II, the model context is described for this work, followed by event generation and selection in Section III. The primary results and analysis are presented in Sections IV (establishment of the signal), V (di-tau event topologies), and VI (resolving the neutralino-stau system). Conclusions are given in Section VII.

II. NO-SCALE \mathcal{F} - $SU(5)$ MODEL CONTEXT

In order to provide a specific context for long-chain cascades of the described type arising from the gluino decay when the squarks are heavy, this study will reference a class of well-defined models named No-Scale \mathcal{F} - $SU(5)$ (see Ref. [6] and references therein), which combine (i) field content of the Flipped $SU(5)$ grand unified theory (GUT), with (ii) a pair of hypothetical TeV-scale vector-like supermultiplets (“flippons”) of mass M_V derivable within local F-Theory model building, and (iii) the boundary conditions of No-Scale Supergravity (SUGRA).

The signature collider characteristic of this model family is a light stop t_1 , lighter than the gluino \tilde{g} , which is in turn lighter than all other squarks \tilde{q} . Lightness of the gluino is attributable to flippon-induced modifications to evolution of the renormalization group equations (RGEs), specifically a nullification at first-loop of the color-associated beta-function coefficient ($b_3 = 0$); these modifications simultaneously elevate the scale of secondary Flipped $SU(5) \times U(1)_X$ gauge unification into adjacency with M_{Planck} , allowing for phenomenologically consistent implementation of the notoriously delicate No-Scale framework [17, 18]. The gluino to stop plus top decay $\tilde{g} \rightarrow t\bar{t}$ is on-shell, with a 100% branching ratio in the viable parameter space, and is associated with an

extreme prevalence of hadronic jets [7].

The key phenomenological characteristic of this model is proportional rescaling at leading order of the full model spectrum with respect to only a single high scale dimensionful parameter, the unified gaugino mass $M_{1/2}$. The model is only logarithmically sensitive to the specific threshold scale M_V at which the vector-like multiplets actively circulate in loops. Lower order facility is available to exchange small variations of $\tan \beta$, the ratio of up- and down-like Higgs vacuum expectation values, within a narrow range around 20 – 25 for a corresponding fluctuation in the stau-LSP mass gap ΔM between about 5 and 25 GeV. The neutralino LSP dark matter candidate is always dominantly Bino.

This model thereby provides a tightly constrained, yet realistic, testbed for a study of the described type, which may moreover exemplify a formally smooth transition between thermal and non-thermal (or thermal mechanisms with an alternative LSP candidate, namely the gravitino) dark matter scenarios. The \mathcal{F} - $SU(5)$ SUSY spectrum for $M_{1/2} \sim 1$ TeV and $\Delta M \sim 6$ GeV is provided in Table I. This example is in the stau-neutralino coannihilation region, with thermal Bino dark matter providing the observed relic density. The selected mass range is in the vicinity of the exclusion boundary established data from the 7 and 8 TeV Large Hadron Collider (LHC) runs; commencement of collisions near the 14 TeV design energy will actively probe the \mathcal{F} - $SU(5)$ construction at scales above $M_{1/2} = 1$ TeV. The technology to be described is generically applicable for all UV models that feature light gluinos, for example those based alternatively on the $SO(10)$ GUT, and can be employed broadly in the next run of LHC.

TABLE I: Spectrum (in GeV) for $M_{1/2} = 990$ GeV, $M_V = 8044$ GeV, $m_t = 174.4$ GeV, and $\tan \beta = 23.3$. Here, $\Omega_{\text{CDM}} h^2 = 0.1197$, the stau-LSP mass gap is $\Delta M = 6.4$ GeV, and the lightest neutralino is greater than 99% Bino. For other values of $M_{1/2}$, revisions to the complete SUSY spectrum may be very well approximated by a simple proportional rescaling. ΔM may be increased by slightly lowering $\tan \beta$, with minimal additional effect on the spectrum overall.

$\tilde{\chi}_1^0$	213	$\tilde{\chi}_1^\pm$	449	\tilde{e}_R	366	\tilde{t}_1	1104	\tilde{u}_R	1824	m_h	125.1
$\tilde{\chi}_2^0$	449	$\tilde{\chi}_2^\pm$	1463	\tilde{e}_L	989	\tilde{t}_2	1672	\tilde{u}_L	1985	$m_{A,H}$	1590
$\tilde{\chi}_3^0$	1461	$\tilde{\nu}_{e/\mu}$	986	$\tilde{\tau}_1$	220	b_1	1650	\tilde{d}_R	1887	m_{H^\pm}	1592
$\tilde{\chi}_4^0$	1463	$\tilde{\nu}_\tau$	958	$\tilde{\tau}_2$	964	b_2	1789	\tilde{d}_L	1986	\tilde{g}	1328

III. EVENT GENERATION AND SELECTION

Numerical analysis of the parameter interdependencies in our benchmark signal model (No-Scale \mathcal{F} - $SU(5)$) is conducted with **SuSpect 2.34** [19], utilizing a proprietary codebase modification that incorporates the TeV-scale vector-like multiplet RGEs. Signal and standard model (SM) background Monte Carlo event samples, including parton showering and fast detec-

tor simulation, are generated via the standard MADGRAPH5/MADEVENT [20], PYTHIA [21], PGS4 [22] chain. Subclassifications of SUSY two-body production channels, such as $\tilde{g}\tilde{g}$, $\tilde{g}\tilde{q}$, $\tilde{q}\tilde{q}^*$, and etc., are individually simulated for a variety of signal benchmark configurations with 0, 1, or 2 hard jets (generated inclusively) at the matrix element level. There is some approximation here (at a level consistent with other technological and systematic limitations) associated with (for example) the reembedding of certain diagrams for squark production with 0 jets into diagrams for gluino pair production with 2 jets. MADEVENT is configured, in conjunction with PYTHIA, to use MLM matching with kt jets in the “Shower kT” scheme, setting (`ickkw` \Rightarrow 1), (`xqcut` \Rightarrow 200), (`drjj` \Rightarrow 0), (`auto_ptj_mjj` \Rightarrow T), (`QCUT` \Rightarrow 200), and (`SHOWERT` \Rightarrow T). The PGS4 detector simulation employs a standard LHC-appropriate parameter card, with jet clustering performed using the anti-kt algorithm. Given the vital importance of hadronic tau reconstruction to this study, the algorithm native to PGS4 (which is based on the somewhat dated “shrinking cone” treatment employed at CDF) has been entirely replaced by a customized treatment that directly references the explicit decay chain at the PYTHIA level, applying a flat efficiency of 60% and a flat fake rate of 1%. This algorithm is adapted from prior work by Alfredo Gurrola and Kechen Wang, as featured in Ref. [23]. Specifically, the PYTHIA event tree is walked to isolate tau particles, retaining only those which further decay hadronically, and subtracting off the 4-momentum of the invisible tau-neutrino. The resulting tracking information is retained for directionality, while the energy scale is extracted from the hardest calorimeter cluster deposit within 10 degrees (if available), avoiding duplication among tau candidates. The Monte Carlo efficiencies are then applied, with a 40% likelihood for reversion to a (non b-tagged) jet. The algorithms for tagging of heavy flavor jets have also been customized for a higher efficiency. The existing b-tagging in PGS4 is Monte Carlo based, with efficiencies and fake rates parameterized as a high-order polynomial function of the transverse momentum and pseudo-rapidity. We have not deviated from this basic framework, but have instead supplemented the existing (rather low efficiency) polynomial fits with new curves sampled from collaboration reports, e.g. the ATLAS design specification [24] (page 412), with more modern efficiency profiles (see Ref. [7] for additional details), presently adopting a curve with a cumulative efficiency on the order of 75%. Although the PGS4 hard acceptance cutoff in pseudo-rapidity for b-Jets ($|\eta| < 2.0$) is unmodified, the acceptance for generic jet candidates has been extended to $|\eta| < 5.0$. Selection cuts are implemented on the detector-level event simulation within AEACUS 3.6 [25, 26]. Initial event selections for the various targeted final states are made very conservatively, to allow for subsequent optimization.

The strongest signal of new physics in a $m(\tilde{q}) > m(\tilde{g}) > m(\tilde{t}_1)$ type model is expected in association with

extremely long cascade decay chains, featuring a strong four W plus four b heavy flavor jet component [5]. Since the W may decay leptonically (1/3 for three light generations) or hadronically (2/3 for two light generations times 3 colors), the final state will also be profused with leptons and multi-jets. In order to establish the signal, we therefore require at least two b -jets in all cases, while recording the net count of jets, leptons, di-leptons, and missing transverse energy \cancel{E}_T , expecting (i) that events with fewer leptons should have more jets, and (ii) that the dominant $t\bar{t}$ + Jets background may likewise have large jet counts, but should not generally feature very large \cancel{E}_T values. For $t\bar{t}$ + Jets, charge conservation further implies that any dilepton production must be anti-correlated in sign, whereas the independent leptonic decay events are uncorrelated in flavor. The SUSY four W + b signal may readily produce tri-leptons (category III), which are inaccessible, outside of fakes, to $t\bar{t}$ + Jets; this category, which necessarily includes also a like-sign dilepton, should be intrinsically low background. Likewise, the orthogonal categorization of precisely two leptons (category II) with like sign should intrinsically suppress $t\bar{t}$ + Jets, with residual fakes, sign-flips, etc., reduced by a requirement on missing transverse energy \cancel{E}_T . The remaining event subdivisions (category I), i.e. those with 0, 1, or 2 leptons, but no like-sign dilepton, will rely heavily on the missing energy cut for background reduction, but may also feature a much stronger net signal count. Opposite sign di-tau production, which is subsequently studied in detail, is a small subset of this very broad event category (though the final jet requirements will ultimately be quite different). See Ref. [27] for another approach to the category II and III channels at $\sqrt{s} = 8$ TeV.

In Figs. (1,2,3), left-hand panels, we additionally consider the contributions of several other background candidates, including the inclusive production of two vector bosons with 0,1, or 2 Jets, as well as single W - or Z -boson production including 0-4 initial state Jets. It is verified explicitly (with large over-sampling) that no contribution is expected to any of these three signal regions from the vector boson pair plus jets background at any missing energy value above 200 GeV. Likewise, no events are observed to survive cuts in the single vector plus jets backgrounds, although those channels do feature large production cross sections, around two orders larger than the corresponding $t\bar{t}$ + background, and the approximately 3.5 and 10 million events considered for each case remain a substantial under-sampling. In conjunction with the intuition that the targeted final state topologies should prove quite difficult for these alternative backgrounds (especially the single vector scenarios), to replicate, we are persuaded that the $t\bar{t}$ +0-2 Jets channel does indeed constitute a fair proxy for the total background. Some bin smoothing is applied in the referenced figures, and in most subsequent plots.

In the $2j + 2\tau + \cancel{E}_T$ final state, which is vital to probing the question of dark matter coannihilation, we preliminarily select on (i) two or more isolated taus with

$P_T > (40, 20)$ GeV in $|\eta| < 2.5$; (ii) two jets $j_{1,2}$ with $P_T > 20$ GeV in $|\eta| < 2.5$ for the conventional scenario, *OR* two jets $j_{1,2}$ with $P_T > 20$ GeV, $(\eta_1 \times \eta_2 < 0)$, and $|\eta_2 - \eta_1| > 3.5$ in $|\eta| < 5.0$ for the boosted scenario. More specifically, in the latter scenario, the pair satisfying these criteria that has the largest invariant mass is tagged, and additional jets interior to this pair are allowed. The AEACuS instructions (with comments) applicable to these selections (and to the computation and storage of other analysis parameters such as the \cancel{E}_T , jet kinematics, heavy flavor and light lepton counts, di-tau sign orientation, transverse momentum components, and invariant mass) are documented in Card A.

Similarly, in Figs. (8,9), left-hand panels, we additionally consider the inclusive production of two vector bosons with 0,1, or 2 Jets, as well as single W - or Z -boson production including 0-4 initial state Jets. The diboson production mode is quite capable of mimicking the targeted ditau plus jets final state, and a smooth contribution from this background is observed, but it is generically suppressed by at least a magnitude order relative to the $t\bar{t}$ background, and is thus ignorable at leading order. In the single boson production channels, a Z may possibly decay to an opposite sign ditau pair, with tau-neutrino missing transverse momentum (an isolated spike reflecting statistical under-sampling is observed in Fig. (9) at a modest \cancel{E}_T value near 200 GeV), and a jet-to-tau fake may likewise potentially allow contributions from the W . However, these modes are likewise expected to be heavily suppressed at the extreme \cancel{E}_T values to be suggested later in this study. As before, we conclude that the $t\bar{t}+0-2$ Jets channel remains a fair proxy for the total background at this level of resolution, being better topologically fitted to the targeted signal final states, and possessing the added practical benefit of an intermediate cross section, which may be readily oversampled in Monte Carlo. It has additionally been suggested in a study [27] at $\sqrt{s} = 8$ TeV that search channels with leptons (specifically like-sign or trilepton) in conjunction with b-tagged jets may receive a contribution from the very rare standard model process $t\bar{t}V$ that is more important than diboson production. We mention this (along with $t\bar{t}Z$, $t\bar{t}t\bar{t}$, $t\bar{t}H$, and VVV) as an example of various rare backgrounds which are not explored here in detail. In the present case, however, where opposite-sign ditau production is allowed (and in fact targeted), we note that the much more common production of $t\bar{t}$ pairs provides already a suitable topological match, limiting the need for consideration of more exotic modes.

It should be further noted that the so-called “K-factors” associated with $t\bar{t}$ production at next-to-leading order (NLO) may increase event yields by approximately a factor of 2. In earlier studies, e.g. Ref. [28], certain of the present authors have likewise carefully compensated signal yields for models similar to those studied here at NLO using PROSPINO 2.1 [29], finding that a K-factor very near to 2 can likewise be typical, when averaged over production channels (squark pair production

has a smaller K-factor, whereas gluino pair production and gluino-squark processes feature a larger one). These factors typically nearly cancel in a signal-to-background comparison, and their explicit consideration is neglected in the following analysis. However, since they will have some small impact on significances computed with a square-root term, and on absolute event yields, the reader is reminded of their existence.

We close this section with a brief discussion of potential errors associated with the narrow-width approximation, i.e. the assumption that intermediate states are produced nearly on-shell, such that production and decay may be decoupled, and factorized [30]. With respect to the stau-neutralino sector, we have rather direct evidence that the issue is under control, via stability of the production cross sections in each considered channel as the relevant mass gap is widened from 6 to 25 GeV. Another scenario that could be of concern is gluino and heavy squark production, where the system’s mass sum may be several TeV, on par with the likely parton energy threshold of a $\sqrt{s} = 14$ TeV collision. It is our opinion that the standardized Monte Carlo tools are performing acceptably in this regime, but this is nevertheless a potential source of some theoretical uncertainty that the reader may wish to be aware of.

```

1 ***** cut_card.dat v3.6 ***
2 * DiTau, Jets and MET Searches
3 *** Object Reconstruction ***
4 OBJ_JET      = PTM:20, PRM:[0.0,2.5]
5 # Jet candidates with P.T > 20 & |ETA| < 2.5
6 # OBJ_JET      = PTM:20, PRM:[0.0,5.0]
7 # Alternative "VBF" Wide Jets for boosted scenario
8 OBJ_JET_001 = SRC:+000, CUT:[2,UNDEF,-1]
9 # Classifies Leading Jet pair objects
10 # OBJ_JET_001 = SRC:+000, EFF:[-1,3.5,UNDEF,1],
11 # CUT:[2,UNDEF,-1] # Alternative "VBF" Topology
12 # for boosted scenario with Delta ETA > 3.5
13 OBJ_JET_002 = SRC:+001, CUT:[1,UNDEF,-1],
14 OUT:[PTM_001,MAS_001,ETA_001,PHI_001]
15 # Outputs Lead Jet kinematics
16 OBJ_JET_003 = SRC:[+001,-002],
17 OUT:[PTM_002,MAS_002,ETA_002,PHI_002]
18 # Outputs Soft Jet kinematics
19 OBJ_JET_004 = SRC:[+002,+003], HFT:0.5, CUT:0
20 # Counts Heavy Flavor Jets (no cut)
21 OBJ_LEP      = PTM:10, PRM:[0.0,2.5]
22 # Soft Leptons restricted in P.T and ETA
23 OBJ_LEP_001 = SRC:+000, PTM:20, EMT:+3, SDR:[0.3,UNDEF,1],
24 CUT:[2,UNDEF,-1] # Two isolated Taus with P.T > 20
25 OBJ_LEP_002 = SRC:+001, PTM:40, CUT:[1,UNDEF,-1],
26 OUT:PTM_003 # Lead Tau has P.T > 40
27 OBJ_LEP_003 = SRC:[+001,-002], OUT:PTM_004
28 # Object holds the Soft Tau (no cut)
29 OBJ_LEP_004 = SRC:+000, EMT:-3, CUT:0
30 # Counts soft light leptons (no cut)
31 OBJ_DIL_001 = LEP:001, DLS:+1, CUT:0
32 # Counts Like Sign DiTaus (no cut)
33 OBJ_DIL_002 = LEP:001, DLS:-1, CUT:0
34 # Counts Opposite Sign DiTaus (no cut)
35 ***** Event Selection *****
36 EVT_MET      = CUT:0
37 # Outputs Missing Energy (no cut)
38 EVT_OIM_001 = LEP:001, CUT:0
39 # Outputs DiTau Invariant Mass (no cut)
40 *****

```

Card A: AEACuS instruction card for di-Tau, jets and missing transverse energy \cancel{E}_T searches.

IV. ESTABLISHING THE THIRD GENERATION CASCADE SIGNAL

```

1 ***** cut_card.dat v3.6 ***
2 * B-Jets, Jets, Leptons, and MET Searches
3 *** Object Reconstruction ***
4 OBJ_JET      = PTM:20, PRM:[0.0,2.5]
5 # Jet candidates with P_T > 20 & |ETA| < 2.5
6 OBJ_JET_001 = SRC:+000, PTM:40, CUT:8
7 # Force 8 Jets with P_T > 40
8 OBJ_JET_002 = SRC:+001, PTM:80, CUT:4
9 # Force 4 Jets with P_T > 80
10 OBJ_JET_003 = SRC:+002, PTM:200, CUT:2
11 # Force 2 Jets with P_T > 200
12 OBJ_JET_004 = SRC:+003, PTM:400, CUT:1
13 # Force 1 Jet with P_T > 400
14 OBJ_JET_005 = SRC:+002, HFT:0.5, CUT:2
15 # Force 2 b-Jets with P_T > 80
16 OBJ_LEP      = PTM:20, PRM:[0.0,2.5]
17 # Soft Leptons restricted in P_T and ETA
18 OBJ_LEP_001 = SRC:+000, SDR:[0.3,UNDEF,1], CUT:[0,2]
19 # Force 0, 1, or 2 Leptons
20 OBJ_DIL_001 = LEP:001, DLS:+1, CUT:[0,0]
21 # Veto like sign dileptons
22 ***** Event Selection *****
23 EVT_MET      = CUT:700
24 # Cut on Missing Energy below 700 GeV
25 *****

```

Card B: AEACuS instruction card for the Category I event selection with b -jets, jets, leptons, and missing transverse energy \cancel{E}_T . The example may be readily modified for the Category II or III event selections.

The described 2 b -jet signal categories (I,II,III) corresponding to prominent signals associable with models featuring long-chain decay cascades with a light third generation have been established in Monte Carlo simulation. For concreteness, the analysis is performed in the context of the No-Scale \mathcal{F} - $SU(5)$ model described in Section II, although the results may be recast for any construction featuring a similar hierarchy in the mass spectrum. The conclusions are little changed for a b -jet transverse momentum threshold of 40 GeV versus 80 GeV, so the more robust value of 80 is retained. The intuition that something like 8,6,4 hard jets may be respectively expected in the signal for each category is well confirmed, noting that the latter categories exchange jet pair production for lepton production in the W decay. Any jets associated with a squark to gluino transition (typically a 500 GeV to 750 GeV mass gap) are expected to be quite hard. Jets downstream from the stop decay also receive a substantial boost from the mass differential, and all downstream jets may inherit large kinematic boosts, even in decays with less phase space. Requiring $P_T^{1,2} > (400, 200)$ GeV facilitates very robust tagging on the leading jet pair, while dampening background (allowing a lower \cancel{E}_T floor), and retaining excellent signal statistics. Jets 3 and 4 are well resolved at $P_T^{3,4} > 80$ GeV, consistent with the b -jet threshold, whereas any jets required beyond the leading four are better captured with softer threshold around $P_T^{5+} > 40$ GeV. With these cuts in place, the missing transverse energy \cancel{E}_T threshold may be individually optimized for each category, as demonstrated in Figs. (1,2,3).

We will select $\cancel{E}_T > (700, 500, 300)$ GeV, respectively. The AEACuS instructions (with comments) applicable to the Category I final state selections are documented in Card B, and may be readily adapted for application to categories II or III.

The background is found in each case to be extraordinarily well controlled, with excellent signal retention. Categories III (tri-leptons) and II (like-sign di-leptons) appear to be observable up to about $M_{1/2} \sim 1200$ GeV, while the primary category I (all other events) may be probed beyond $M_{1/2} \sim 1400$ GeV, encompassing the majority, if not totality, of the \mathcal{F} - $SU(5)$ model space. The gluino masses in these cases are on the order of 1600 and 1900 GeV, respectively. As demonstrated clearly in Fig. (1), the expected SUSY event yield is a strongly decaying function of $M_{1/2}$, which may be inverted in order to establish the global model mass scale. Since the model is dominantly single parameter, the bulk properties of the spectrum are then fixed, and may be cross-correlated against alternatively designed event selections for consistency, such as the di-tau production channel to be elaborated subsequently.

In Fig. (4), we show that the signal is indeed more rich in heavy flavor jets than the background, even after considering b -fake rates. It should be recalled that these rates also receive a sizeable contribution from the hadronic decays of W to charm/strange. We also show the significance optimization for various $M_{1/2}$ values in ≥ 8 jets + ≥ 4 b -jets + \cancel{E}_T final states. It is clear, in comparison to Fig. (1), that the stronger b -jet requirement allows for a softer \cancel{E}_T cut, although this strategy is not necessarily a more favorable one.

In Fig. (5), we depict the density of hadronic W -bosons reconstructed with an invariant mass inside the 65 to 95 GeV window out of non- b tagged jets, after application of all the Category I cuts. In this limit, the signal is showing more W counts, although a trend toward under representing the expected prevalence of W 's has been observed. In the gluino pair production channel, at $M_{1/2} \sim 1$ TeV, two top/stop pairs are produced, and the stop will produce a top/neutralino with probability $p \sim 0.86$, or a bottom/chargino with probability $(1 - p) \sim 0.14$, such that the expected density of top quarks is $2_t \times (1 - p)^2 + 3_t \times 2p(1 - p) + 4_t \times p^2 \sim 3.7$, each of which should be associated with a final state W -boson. This has been validated in Monte Carlo at the generator level, although is apparently difficult to resolve at the detector level using elementary techniques. Several contributions to the W shortage are trivial, including exclusion of leptonic branching (the neutrino cannot be deconvolved from the large SUSY \cancel{E}_T content), the likelihood of heavy flavor tagging for decays to charm/strange, limits on kinematic acceptance of jets, and smearing of the jet energy resolution. However, difficulty in clearly distinguishing the SUSY hadronic W shape distribution from that of $t\bar{t}$ suggests that a more subtle agent is also at work. Specifically, it would seem to implicate the absorption of distinct partonic chains into combined fat jets, as




FIG. 1: Events with (0,1,2) leptons and no like-sign dilepton (category I) are studied at a luminosity of 30 events per femtobarn. Two heavy-flavor tagged jets with $P_T > 80$ GeV are required, and the leading eight jets (with or without a b-tag) must carry $P_T > (400, 200, 80, 80, 40)$ GeV. Left: Absolute signal and background component event counts are binned as a function of the missing transverse energy \cancel{E}_T . Right: Signal significance relative to the leading $t\bar{t}$ background is evaluated for four signal regions as a function of the \cancel{E}_T cut.




FIG. 2: Events with a like-sign dilepton topology (category II) are studied at a luminosity of 30 events per femtobarn. Two heavy-flavor tagged jets with $P_T > 80$ GeV are required, and the leading four jets (with or without a b-tag) must carry $P_T > (400, 200, 80, 80)$ GeV. Two additional jets (for a total of 6) must carry $P_T > 40$ GeV. Left: Absolute signal and background component event counts are binned as a function of the missing transverse energy \cancel{E}_T . Right: Signal significance relative to the leading $t\bar{t}$ background is evaluated for four signal regions as a function of the \cancel{E}_T cut.

is made more likely in a cascade environment with extremely high jet density, and also when decay products are highly boosted (collimated); it is likely that a jet substructure analysis [31, 32] would improve discrimination, but this will not be considered further here.

V. DI-TAU EVENT RESULTS AND ANALYSIS

Having established a strong third generation signal component in the prior section, we now turn attention to the question of stau/LSP coannihilation, via a focus on di-tau event production. Vector boson fusion (VBF) [23, 33, 34] is a key electroweak process capable of highlighting the light neutralino system at the four-vertex order. The VBF process is inherently suppressed,

but may be emphasized by imposing an event topology that features highly boosted tagging jets with wide separation in pseudorapidity. Also, there are a large variety of strongly-interacting diagrams which may be considered VBF analogs that may likewise give rise to the described event topology, such as those exhibited in Fig. (6). Such diagrams cannot directly invoke the neutralino system, but may still probe the electroweak sector downstream, generating similar di-tau production via long-chain decay cascades, in association with a (potentially substantially) elevated net count of jets. By strength of the interaction and of numbers, such processes may residually swamp their electroweak analogs.

Central considerations of the current section are then the potential role of boosted event topologies in the study of long-chain cascade decays with hadronic di-tau final




FIG. 3: Events with a trilepton topology (category III) are studied at a luminosity of 30 events per femtobarn. Two heavy-flavor tagged jets with $P_T > 80$ GeV are required, and the leading four jets (with or without a b-tag) must carry $P_T > (400, 200, 80, 80)$ GeV. Left: Absolute signal and background component event counts are binned as a function of the missing transverse energy E_T . Right: Signal significance relative to the leading $t\bar{t}$ background is evaluated for four signal regions as a function of the E_T cut.




FIG. 4: Left: Normalized distribution of b -jets retained after soft kinematic grooming in the background, the unified signal, and the isolated gluino pair production channel. The background remains larger than the unified signal by a factor of about 80 at this level, with cross sections of 8.8×10^3 and 1.1×10^2 fb, respectively. Restricting to four or more b -jets produces order parity in the event expectations, reducing the cross sections to 5.5 and 2.0 fb, respectively. The gluino pair production sub-channel yields a somewhat larger density of jets in isolation than the full signal in aggregate, to which it contributes approximately one third of net events at this level. Right: The signal significance is evaluated as a function of the missing transverse energy E_T for the Category I selections, adding a 4 b -jet requirement relative to the selection depicted in Fig. (1).

states, and the comparison of that approach to more conventional requirements on the associated jet content. This treatment should be distinguished from that applied to the much larger superset of “category I” events in Section IV, wherein the primary goal was isolation of the overall strongest signal for all modes of strong production, inclusively. To reiterate, the suggestion is not necessarily that boosted event topologies are a good way to look for taus, and thus equivalently a good way to look for light leptons, but rather, that they might be a reasonable way (albeit one that constitutes a very hard cut) to try and isolate the electroweak sector of the model from the strong sector; if the next-to-lightest superpart-

ner (NLSP) is a stau, then one is led naturally to apply this technique to the study of ditau states.

The signal significance metric $S/\sqrt{1+B}$ has been optimized as a function of the missing energy and transverse momenta of the two relevant jets for each of the two considered event topologies (boosted/conventional), with and without the application of light lepton and heavy flavor jet vetoes. This optimization, which is mildly luminosity and scale dependent, has been performed at an integrated luminosity of 30 events per fb, taking an $\mathcal{F}\text{-}SU(5)$ model benchmark with $M_{1/2} = 1000$ GeV and $\Delta M = 25$ GeV. Including a lepton veto hurts the significance ratio by about 30% for both event topologies.




FIG. 5: Normalized distribution of reconstructed hadronic W 's in the Category I ≥ 8 jets with ≥ 2 b -jets plus large E_T final state for both signal and background. The net signal counts are larger than the background after these hard cuts by a factor of about 65, with cross sections of 2.9 and 4.5×10^{-2} fb, respectively. Restricting to two or more W reconstructions effectively suppresses the background and further reduces the signal cross section by a factor of about 20 to 1.5×10^{-1} fb.

Including a b -jet veto hurts the boosted case by about 10%, but the conventional case by 20 or 30%. This latter difference may be traced in part to the fact that the boosted topology employs jets at large η that cannot be tagged for heavy flavor, such that the veto is less harmful. Both results are consistent with the expectation that long cascades down the squark to gluino to stop to neutralino cascade may fork a large population of both light leptons and b -jets. The possibility of widening the jet pseudorapidity acceptance to $|\eta| < 5.0$ for the conventional event topology has also been examined, and found to be inconsequential with regards both to net signal efficiency and the optimization structure; this observation may be traced to the fact that the two leading jets in P_T are not likely to exist at large pseudorapidity, as these conditions are anti-correlated.

For both scenarios, the most important cut is on missing energy, and values around 700 GeV work very well. In the conventional event topology, the signal significance at $\mathcal{L} = 30/\text{fb}$ is moderately, though not greatly, enhanced by the addition of cuts on the leading jet transverse momenta P_T . This is because the background would already be well controlled by the E_T cut alone and the "+1" regulator term is playing a deciding role in the denominator. Nevertheless signal is not much harmed by pushing the jet momenta substantially (a large E_T will require large jet momenta) and background is thereby suppressed in a much more robust manner. This is supported by the Table II cut flow, presented in terms of the residual fb cross section. Specifically, the addition of a cut $P_T > (400, 200)$ reduces the background by about 150 times, but the signal by only about 2.5 times; this is prior to implementation of any E_T cut whatsoever.




FIG. 6: Feynman diagrams (adapted from MADGRAPH5 [20] output) contributing to strong interaction analogs of the electroweak "Vector Boson Fusion" boosted event topology with di-jet plus $\tilde{g}\tilde{g}$, $\tilde{g}q$, and $q\bar{q}^*$ final states.

TABLE II: Residual effective cross-section (fb) at various cut flow stages for the two described event topologies, as applied to the $M_{1/2} = 1000$ GeV \mathcal{F} - $SU(5)$ benchmark and the $t\bar{t}$ plus 2 Jets inclusive background at the LHC14. The production level cross sections and di-tau event selections are common to both scenarios. Energies and momenta are in GeV.

Selection	$t\bar{t}$ +Jets	Signal
Matched Production	645,000	192
2 τ 's, $P_T > (40, 20)$	2,230	10.1
<i>Conventional Event Topology</i>		
2 Jets, $P_T > (400, 200)$	15	4.2
$E_T > 700$	0.040	1.5
<i>Boosted Event Topology</i>		
2 Jets, $\Delta\eta > 3.5$ & $P_T > (75, 50)$	88.1	0.78
$E_T > 700$	0.005	0.21

For the boosted event topology, increasing P_T of the jets beyond the initial 20 GeV threshold is actually quite detrimental, almost immediately. Note in this case that the E_T contribution may be balanced by more central non-tagged jets with much higher P_T . However, we are




FIG. 7: Normalized signal and background distribution shapes are compared as a function of the missing transverse energy E_T for the boosted and conventional event topologies.




FIG. 8: Di-Tau events are studied at a luminosity of 30 events per femtobarn under application of the conventional event topology. All additional cuts documented in Section I are imposed as a prerequisite. Left: Absolute signal and background component event counts are binned as a function of the missing transverse energy E_T . Right: Signal significance relative to the leading $t\bar{t}$ background is evaluated for four signal regions as a function of the E_T cut.




FIG. 9: Di-Tau events are studied at a luminosity of 30 events per femtobarn under application of the boosted event topology. All additional cuts documented in Section I are imposed as a prerequisite. Left: Absolute signal and background component event counts are binned as a function of the missing transverse energy E_T . Right: Signal significance relative to the leading $t\bar{t}$ background is evaluated for four signal regions as a function of the E_T cut.

concerned that such soft jets are not realistic for tagging in a high pile-up environment, and instead adopt the more conservative bounds $P_T > (75, 50)$, sacrificing approximately two thirds of the signal in the process. The background drops by a similar, if slightly larger, factor close to five. Table II suggests that the hard jet cuts applied in the conventional event topology remove quite a bit less signal and quite a bit more background than the corresponding $\Delta\eta$ requirement imposed in the boosted topology; however, subsequent application of an identical $\cancel{E}_T > 700$ GeV cut on missing transverse energy devastates the boosted topology background component, while removing very little additional signal by comparison.

After all cuts, the signal and background expectations are both larger, by around 7 or 8 times, for the conventional event topology, but the signal to background ratio S/B is essentially identical for the two event topologies. It should be emphasized that both scenarios, and especially the boosted event topology, present such stringent suppression of the $t\bar{t}$ + jets background component, no more than about one event per 25 fb^{-1} of data, that the statistical and systematic reliability of the analysis may become over stretched when extended to the detailed comparison of two extremely small values; nevertheless, the central message appears quite robust: backgrounds are extremely well controlled, and signal has a realistic chance to present itself.

Fig. (7) depicts the unity-normalized $t\bar{t}$ +jets and signal event densities as a function of missing transverse energy \cancel{E}_T for both the boosted and conventional event topologies, with all other cuts applied. The pair of distributions are found to be extremely similar in shape, each exhibiting strong differentiation of signal and background for sufficiently strong \cancel{E}_T cuts. The heavier signal benchmarks extend more prevalently into the extremely large missing energy domain. The boosted event topology does demonstrate a somewhat harder background suppression slope in the distribution tail.

Figs. (8,9), right-hand panels, exhibit the integrated signal significance as a function of the missing transverse energy cut for the boosted and conventional event topologies. The metric $S/\sqrt{1+B}$ is employed, comparing the count of signal events to the Gaussian expectation for fluctuation inherent to the statistical background, with the numeral 1 employed as a low-background regulator. A very mild scale dependence is observed, with lighter models favoring a bit less missing energy; this asymmetry is strongly accentuated when employing the alternative significance metric $S/\sqrt{S+B}$, because lighter models generate a large quantity of signal events, which may exceed or retain parity with the background, such that significance scales with the signal square-root, disfavoring an over-strong cut. Optimization with regard $S/\sqrt{1+B}$ is much less dependent upon the signal event scale, being driven instead primarily by background elimination, which becomes particularly pronounced in the vicinity of $\cancel{E}_T > 500 - 700$ GeV. The apparent tendency of the boosted event topology to favor a lighter \cancel{E}_T cut

is attributable to the previously described reduction in net event counts associated with this scenario, such that the regulator term is dominant at moderate luminosities; nevertheless, the ratio S/B continues to benefit substantially from a harder cut.




FIG. 10: The ratio of boosted to conventional signal to background ratios is displayed as a function of the gaugino scale parameter $M_{1/2}$ for various \cancel{E}_T cut thresholds.

Fig. (10) compares the boosted versus conventional signal to background ratios S/B as a function of the signal event scale for various common \cancel{E}_T cuts. This ratio is observed to be comparable for both event topologies at a missing energy cut around 700 GeV, independent of the model scale.

Fig. (11) depicts the raw signal and $t\bar{t}$ event counts for the boosted and conventional event topologies as a function of the gaugino scale parameter $M_{1/2}$ for various values of the missing energy cut at an integrated luminosity of 300 events per femtobarn. A global suppression factor on the order of 10 is observed for application of the boosted event topology. A dramatic discrimination is observed in the relative impact of background (much stronger) and signal (much milder) suppression with increasing missing energy. For $\cancel{E}_T > 700$ GeV, the \mathcal{F} - $SU(5)$ signal counts exceed the $t\bar{t}$ background counts for event scales across the viable model space, up to $M_{1/2} = 1.5$ TeV and beyond. A minimal signal count of more than 5 events is likewise maintained over this full scale range, even for the boosted topology. For a softer missing energy cut, e.g. $\cancel{E}_T > 300$ GeV, the observed events are expected to become background dominated at a much lighter scale, as low as 1.0 TeV in this case.

VI. THE NEUTRALINO-STAU SYSTEM

The structure of the neutralino-stau mass hierarchy will be strongly imprinted upon the kinematic distribution of invariant masses $M_{\tau\tau}$ for opposite-sign (OS) ditau pairs [16]. The $M_{\tau\tau}$ variable employed here is the invariant mass of the four-vectors associated with the





FIG. 11: Expected $\mathcal{F}\text{-}SU(5)$ signal and $t\bar{t}$ background counts at a luminosity of 300 events per femtobarn are depicted for the boosted and conventional event topologies as a function of the gaugino scale parameter $M_{1/2}$ for various E_T cut thresholds.





FIG. 12: The conventional event topology opposite-sign minus like-sign di-tau invariant mass $M_{\tau\tau}$ distribution shape is depicted for various gaugino mass $M_{1/2}$ scales and mass gaps $\Delta M \equiv M_{\tilde{\tau}_1} - M_{\tilde{\chi}_1^0}$.

hadronic reconstructions described in Section III, taking the hardest pair in the unlikely event of more than two candidates. There is no attempt to further reconstruct leptonic tau decays. After subtracting the density of like-sign (LS) pairs, which serve as a statistical estimator for the density and shape of unassociated opposite sign production, the invariant mass cutoff $M_{\tau\tau}^{\max}$ [16] shown following may become visible, thereby providing a window into the nature of dark matter and its potential compatibility with thermal processes [35, 36] featuring a neutralino LSP that is largely orthogonal to other techniques.

$$M_{\tau\tau}^{\max} = M_{\tilde{\chi}_2^0} \sqrt{1 - \frac{M_{\tilde{\tau}_1}^2}{M_{\tilde{\chi}_2^0}^2}} \sqrt{1 - \frac{M_{\tilde{\chi}_1^0}^2}{M_{\tilde{\tau}_1}^2}} \quad (1)$$

Fig. (12) depicts the opposite-sign (OS) minus like-sign (LS) di-tau invariant mass distribution shape in the conventional event topology after applying a missing energy cut of $E_T > 700$ GeV. In the left-hand pane the universal gaugino mass $M_{1/2}$ is varied as the stau-LSP mass gap

$\Delta M \equiv M_{\tilde{\tau}_1} - M_{\tilde{\chi}_1^0}$ is held fixed at 25 GeV. In the right-hand pane, the mass gap ΔM is fluctuated while holding the gaugino mass $M_{1/2}$ (approximately) constant. The distribution cutoff is visually extended in order to facilitate a numerical comparison with theoretical expectations, as in Table III. Fluctuation of this extension suggests a generic error on the order of 10 GeV in the limit of high statistics. Variation of the bin sizing and smoothing parameterization may induce a more or less comparably sized shift, to be combined in quadrature.

Table III examines the agreement between simulated data of a cascade-rich signal and the theoretical prediction from Eq. (1) for the di-tau invariant mass cutoff $M_{\tau\tau}^{\max}$. For a large mass gap $\Delta M \simeq 25$ GeV, which implies either a non-thermal mechanism for dilution of the Bino relic density or an alternative LSP candidate such as the gravitino, agreement is excellent. Ability to discern a well correlated cutoff persists for the background-polluted sample, although efficacy of the method is somewhat diminished in this case at heavier signal scales. Approaching the neutralino thermal mass gap around $\Delta M = 6$ GeV, where the soft tau element becomes in-




FIG. 13: The distribution of transverse momenta P_T associated with individual hadronic tau elements is depicted in terms of the luminosity-independent shape, and in terms of the absolute event yield at fixed luminosity for various values of the mass gap $\Delta M \equiv M_{\tilde{\tau}_1} - M_{\tilde{\chi}_1^0}$.

TABLE III: Comparison between theoretical prediction and Monte-Carlo observation of the opposite-sign minus like-sign di-tau invariant mass distribution cutoff for \mathcal{F} - $SU(5)$ benchmarks with various gaugino mass $M_{1/2}$ and stau-LSP mass gap ΔM values. The “ $M_{\tau\tau}^{\max}$ ” value is computed by direct application of Eq. (1) to the associated spectrum. The “Visual” entry references the extrapolated linear descents plotted in Fig. (12), where applicable. The “ $2\times\text{Peak}$ ” value is established numerically, averaging over bin counts in the upper third of the corresponding histogram plots (after moderate bin smoothing to wash out noise peaks) and simply doubling the result. The “With $t\bar{t}$ ” column applies this same logic to the \mathcal{F} - $SU(5)$ signal and $t\bar{t}$ +jets SM background combination. A dash is presented when the signal is highly unresolved relative to background. All entries are in units of GeV.

$M_{1/2}$	ΔM	$M_{\tau\tau}^{\max}$	Visual	$2\times\text{Peak}$	With $t\bar{t}$
850	25	153	155	155	154
1000	25	172	175	170	165
1200	25	193	195	196	178
1400	25	213	225	204	-
821	15	123	135	135	139
1000	15	140	150	155	150
1400	16	172	175	183	-
875	6	85	120	110	109
990	6	94	125	125	115
1400	6	111	155	145	-

creasingly difficult to resolve, the cutoff extracted from simulation appears to systematically overestimate the Eq. (1) prediction.

The tabulated elements, as well as Fig. (12), use over-sampled data. In practice, the di-tau invariant mass peak is a difficult measurement requiring high luminosity. For the model samples under consideration, very little is resolvable at a luminosity of 30 fb^{-1} , but benchmarks in the vicinity of $M_{1/2} \sim 1 \text{ TeV}$ present a sufficient event count for peak resolution at 300 fb^{-1} . Reduction of the E_T

threshold substantially improves event yields, but makes discrimination from the background very difficult.

Fig. (13) visually examines the distribution of transverse momenta P_T associated with individual hadronic taus. The left-hand pane plots the luminosity-independent shape of the soft and hard tau elements for $M_{1/2} = 1000 \text{ GeV}$ and a stau-LSP mass gap $\Delta M = 25 \text{ GeV}$. The depicted shape is broadly representative of that observed in other relevant model scenarios. A sharp decline is evident in the transverse momentum population of the softer tau, which is ostensibly associable with the phase space constricted secondary decay $\tilde{\tau}_1 \rightarrow \tilde{\chi}_1^0 + \tau$; this circumstance highlights the criticality of extending detection acceptance for hadronic tau candidates to as low a transverse momentum threshold as is technically possible, and at least to $P_T \sim 20 \text{ GeV}$. The transverse momentum distribution of the lead tau, which is expected to arise from the kinematically broad channel $\tilde{\chi}_2^0 \rightarrow \tilde{\tau}_1 + \tau$, is rather flat; a harder cut here, on the order that currently employed at $P_T > 40$, does not appear detrimental. The right-hand pane depicts the absolute distribution height of the soft tau element at a luminosity of 300 events per femtobarn for various values of ΔM . As expected, this figure suggests that it is substantially more difficult (though not impossible) to effectively probe di-tau production in the case of a narrow stau-LSP mass gap; visibility of the daughter tau is more strongly dependent in this case upon a substantial upstream boosting of the parent stau.

VII. DISCUSSION AND CONCLUSIONS

In this paper we address a scenario where the gluino mass is in the 1 to 2 TeV range, whereas the first two squark masses and sleptons are heavy. Specifically, we have considered the No-Scale \mathcal{F} - $SU(5)$ model, where the presence of additional vector like fields may contribute to the generation of a Higgs mass near 126 GeV. The light

stop \tilde{t}_1 is lighter than the gluino in this model, which allows the gluino to decay on-shell into $\tilde{t}\tilde{t}_1$. The lighter stop may then decay directly into a top plus lightest neutralino, or into a top plus (Wino-like) second lightest neutralino or bottom plus light chargino, which decay in turn to the lightest neutralino via stau. The dark matter content may be obtained thermally by stau and (Bino-like) light neutralino coannihilation in the case of a small $\Delta M \simeq 6$ GeV. The final state of such a cascade contains multiple b -quarks, W 's, τ leptons and missing energy, and additional light flavor leptons, making $t\bar{t}$ the dominant background.

We first established the scenario from the multi- b , leptons and W 's, so that the existence of third generation can be surmised. We showed the signal can be differentiated from the leading background (where we considered $t\bar{t}$ + multijet production with a 1% b -fake rate) by demanding multiple jets, multiple b -quarks, and multiple W 's in the final state, in addition to very large missing transverse energy \cancel{E}_T . We calculated significances and showed that $M_{1/2}$ scales up to 1400 GeV ($M_{\tilde{g}} \sim 1900$ GeV) can be investigated at the 14 TeV LHC with a 30 fb^{-1} luminosity, requiring 8 jets with 2 heavy flavor tags and $\cancel{E}_T > 700$ GeV. In the multi-lepton cases, we showed that a strategy requiring at least 6 jets with a like sign dilepton, or four jets with three leptons, can be utilized in conjunction with missing transverse energy to establish the model. We showed that $M_{1/2}$ up to around 1200 GeV ($M_{\tilde{g}} \sim 1600$ GeV) can be investigated in these channels at the 14 TeV LHC with a 30 fb^{-1} luminosity; sensitivity may be extended to $M_{1/2} \sim 1300$ -1400 GeV ($M_{\tilde{g}} \sim 1750$ -1900 GeV) with a 300 fb^{-1} luminosity.

We investigated the coannihilation region by considering two analysis routes, each of which require at least two τ 's with $P_T > (40, 20)$ GeV in $|\eta| < 2.5$ and large missing energy. In the ‘‘Boosted Event Topology’’, two

tagging jets $j_{1,2}$ with $P_T > (75, 50)$ GeV were required in $|\eta| \leq 5.0$ with an opposite hemisphere orientation ($\eta_1 \times \eta_2 < 0$), and an absolute separation in pseudorapidity of $|\eta_2 - \eta_1| > 3.5$. In the ‘‘Conventional Topology’’, two leading jets $j_{1,2}$ in $|\eta| < 2.5$ carry very large transverse momenta $P_T > (400, 200)$ GeV. To retain the cascade signal, no vetoes on heavy flavor jets or light lepton flavors were enforced. We demonstrated the cut flow optimization that leads to these selections, and calculated the signal to background ratios S/B for both routes. Both event topologies require a large \cancel{E}_T cut around 700 GeV to reduce the background. After all cuts, the signal and background expectations are both larger, by around 7 or 8 times, for the conventional event topology, but the signal to background ratio S/B is essentially identical for the two event topologies. With the optimized cuts, we investigated the small ΔM region and found that gaps of around 6 GeV can be probed for $M_{1/2} \sim 1000$ GeV ($M_{\tilde{g}} \sim 1300$ GeV) at 300 fb^{-1} , although loss of the soft tau somewhat inhibits resolution of the invariant $M_{\tau\tau}^{\text{max}}$ mass in this case; the experimental prospects are improved for larger ΔM GeV (around 15-25 GeV).

VIII. ACKNOWLEDGMENTS

We would like to thank Yu Gao, Teruki Kamon and Nikolay Kolev for helpful discussions. This work is supported in part by DOE Grant No. DE-FG02-13ER42020, NSF Award PHY-1206044, the Natural Science Foundation of China Grant Nos. 10821504, 11075194, 11135003, 11275246, and 11475238, the National Basic Research Program of China (973 Program) under Grant No. 2010CB833000 (TL), NASA Astrophysics Theory Grant NNN12ZDA001N (KS), and the Sam Houston State University Enhancement Research Grant program (JWW).

-
- [1] G. Aad et al. (ATLAS Collaboration), ‘‘Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7 fb^{-1} of $\sqrt{s} = 7$ TeV proton-proton collision data,’’ *Phys.Rev.* **D87**, 012008 (2013), 1208.0949.
 - [2] G. Aad et al. (ATLAS Collaboration), ‘‘Hunt for new phenomena using large jet multiplicities and missing transverse momentum with ATLAS in fb^{-1} of $\sqrt{s} = 7$ TeV proton-proton collisions,’’ (2012), 1206.1760.
 - [3] S. Chatrchyan et al. (CMS Collaboration), ‘‘Search for new physics in the multijet and missing transverse momentum final state in proton-proton collisions at $\sqrt{s} = 7$ TeV,’’ *Phys.Rev.Lett.* **109**, 171803 (2012), 1207.1898.
 - [4] T. A. collaboration, ‘‘Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum and 20.3 fb^{-1} of $\sqrt{s} = 8$ TeV proton-proton collision data,’’ (2013).
 - [5] V. Khachatryan et al. (CMS Collaboration), ‘‘Searches for supersymmetry based on events with b jets and four W bosons in pp collisions at 8 TeV,’’ (2014), 1412.4109.
 - [6] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, ‘‘No-Scale $\mathcal{F}\text{-SU}(5)$ in the Light of LHC, Planck and XENON,’’ *Jour.Phys.* **G40**, 115002 (2013), 1305.1846.
 - [7] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, ‘‘The Ultra-High Jet Multiplicity Signal of Stringy No-Scale $\mathcal{F}\text{-SU}(5)$ at the $\sqrt{s} = 7$ TeV LHC,’’ *Phys.Rev.* **D84**, 076003 (2011), 1103.4160.
 - [8] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, ‘‘A Higgs Mass Shift to 125 GeV and A Multi-Jet Supersymmetry Signal: Miracle of the Flippons at the $\sqrt{s} = 7$ TeV LHC,’’ *Phys.Lett.* **B710**, 207 (2012), 1112.3024.
 - [9] L. Aparicio, M. Cicoli, S. Krippendorff, A. Maharana, F. Muia, et al., ‘‘Sequestered de Sitter String Scenarios: Soft-terms,’’ *JHEP* **1411**, 071 (2014), 1409.1931.
 - [10] M. Cahill-Rowley, J. Hewett, A. Ismail, and T. Rizzo, ‘‘pMSSM Studies at the 7, 8 and 14 TeV LHC,’’ (2013), 1307.8444.
 - [11] H. Baer, V. Barger, P. Huang, A. Mustafayev, and

- X. Tata, “Radiative natural SUSY with a 125 GeV Higgs boson,” *Phys.Rev.Lett.* **109**, 161802 (2012), 1207.3343.
- [12] R. Kitano and Y. Nomura, “Supersymmetry, naturalness, and signatures at the LHC,” *Phys.Rev.* **D73**, 095004 (2006), hep-ph/0602096.
- [13] C. Brust, A. Katz, S. Lawrence, and R. Sundrum, “SUSY, the Third Generation and the LHC,” *JHEP* **1203**, 103 (2012), 1110.6670.
- [14] M. Papucci, J. T. Ruderman, and A. Weiler, “Natural SUSY Endures,” *JHEP* **1209**, 035 (2012), 1110.6926.
- [15] B. Dutta, W. Flanagan, A. Gurrola, W. Johns, T. Kamon, et al., “Probing Compressed Top Squarks at the LHC at 14 TeV,” *Phys.Rev.* **D90**, 095022 (2014), 1312.1348.
- [16] R. L. Arnowitt, B. Dutta, A. Gurrola, T. Kamon, A. Krislock, et al., “Determining the Dark Matter Relic Density in the mSUGRA (X0(1))-tau Co-Annihilation Region at the LHC,” *Phys. Rev. Lett.* **100**, 231802 (2008), 0802.2968.
- [17] J. R. Ellis, D. V. Nanopoulos, and K. A. Olive, “Lower limits on soft supersymmetry breaking scalar masses,” *Phys. Lett.* **B525**, 308 (2002), arXiv:0109288.
- [18] J. Ellis, A. Mustafayev, and K. A. Olive, “Resurrecting No-Scale Supergravity Phenomenology,” *Eur. Phys. J.* **C69**, 219 (2010), 1004.5399.
- [19] A. Djouadi, J.-L. Kneur, and G. Moultaka, “SuSpect: A Fortran code for the supersymmetric and Higgs particle spectrum in the MSSM,” *Comput. Phys. Commun.* **176**, 426 (2007), hep-ph/0211331.
- [20] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer, and T. Stelzer, “MadGraph 5 : Going Beyond,” *JHEP* **1106**, 128 (2011), 1106.0522.
- [21] T. Sjostrand, S. Mrenna, and P. Z. Skands, “PYTHIA 6.4 Physics and Manual,” *JHEP* **05**, 026 (2006), hep-ph/0603175.
- [22] J. Conway et al., “PGS4: Pretty Good (Detector) Simulation,” (2009), URL <http://www.physics.ucdavis.edu/~conway/research/>.
- [23] A. G. Delannoy, B. Dutta, A. Gurrola, W. Johns, T. Kamon, et al., “Probing Supersymmetric Dark Matter and the Electroweak Sector using Vector Boson Fusion Processes: A Snowmass Whitepaper,” (2013), 1308.0355.
- [24] G. Aad et al. (ATLAS), “Expected Performance of the ATLAS Experiment - Detector, Trigger and Physics,” (2009), 0901.0512.
- [25] J. W. Walker, “CutLHCO: A Consumer-Level Tool for Implementing Generic Collider Data Selection Cuts in the Search for New Physics,” (2012), 1207.3383.
- [26] J. W. Walker, “AEACuS 3.4,” (2014), URL <http://www.joelwalker.net/code/aeacus.tar.gz>.
- [27] G. Aad et al. (ATLAS), “Search for supersymmetry at $\sqrt{s}=8$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector,” *JHEP* **1406**, 035 (2014), 1404.2500.
- [28] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, “Correlated Event Excesses in LHC SUSY Searches at 7 & 8 TeV: New Physics or Conspiring Noise?,” *Eur.Phys.J.* **C73**, 2556 (2013), 1302.6579.
- [29] W. Beenakker, R. Hopker, and M. Spira, “PROSPINO: A Program for the production of supersymmetric particles in next-to-leading order QCD,” (1996), hep-ph/9611232.
- [30] D. Berdine, N. Kauer, and D. Rainwater, “Breakdown of the Narrow Width Approximation for New Physics,” *Phys.Rev.Lett.* **99**, 111601 (2007), hep-ph/0703058.
- [31] V. Khachatryan et al. (CMS Collaboration), “Search for massive resonances decaying into pairs of boosted bosons in semi-leptonic final states at $\sqrt{s} = 8$ TeV,” *JHEP* **1408**, 174 (2014), 1405.3447.
- [32] V. Khachatryan et al. (CMS Collaboration), “Search for massive resonances in dijet systems containing jets tagged as W or Z boson decays in pp collisions at $\sqrt{s} = 8$ TeV,” *JHEP* **1408**, 173 (2014), 1405.1994.
- [33] B. Dutta, A. Gurrola, W. Johns, T. Kamon, P. Sheldon, et al., “Vector Boson Fusion Processes as a Probe of Supersymmetric Electroweak Sectors at the LHC,” *Phys.Rev.* **D87**, 035029 (2013), 1210.0964.
- [34] A. G. Delannoy, B. Dutta, A. Gurrola, W. Johns, T. Kamon, et al., “Probing Dark Matter at the LHC using Vector Boson Fusion Processes,” *Phys.Rev.Lett.* **111**, 061801 (2013), 1304.7779.
- [35] R. L. Arnowitt, B. Dutta, T. Kamon, N. Kolev, and D. A. Toback, “Detection of SUSY in the stau-neutralino coannihilation region at the LHC,” *Phys.Lett.* **B639**, 46 (2006), hep-ph/0603128.
- [36] B. Dutta, T. Kamon, A. Krislock, K. Sinha, and K. Wang, “Diagnosis of Supersymmetry Breaking Mediation Schemes by Mass Reconstruction at the LHC,” *Phys.Rev.* **D85**, 115007 (2012), 1112.3966.