



# CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Chiral imprint of a cosmic gauge field on primordial gravitational waves

Jannis Bielefeld and Robert R. Caldwell

Phys. Rev. D **91**, 123501 — Published 1 June 2015

DOI: [10.1103/PhysRevD.91.123501](https://doi.org/10.1103/PhysRevD.91.123501)

# Chiral Imprint of a Cosmic Gauge Field on Primordial Gravitational Waves

Jannis Bielefeld and Robert R. Caldwell

Department of Physics and Astronomy, Dartmouth College,  
6127 Wilder Laboratory, Hanover, New Hampshire 03755 USA

(Dated: March 16, 2015)

A cosmological gauge field with isotropic stress-energy introduces parity violation into the behavior of gravitational waves. We show that a primordial spectrum of inflationary gravitational waves develops a preferred handedness, left- or right-circularly polarized, depending on the abundance and coupling of the gauge field during the radiation era. A modest abundance of the gauge field would induce parity-violating correlations of the cosmic microwave background temperature and polarization patterns that could be detected by current and future experiments.

Since the surprising discovery that parity is violated on the atomic scale by the weak nuclear force [1], searches for broken symmetries have proven to be a remarkably effective technique for uncovering new laws of physics. Here we speculate that parity violation on cosmic scales may be the sign of a dark gauge field. Our theoretical model consists of a classical non-Abelian (Yang-Mills) gauge field which, as we demonstrate, behaves like radiation through cosmic history, but fluctuations of the field couple to gravity in a way that distinguishes between left- and right-handed circularly polarized gravitational waves. We demonstrate the effect of this field on a primordial spectrum of gravitational waves and evaluate its impact on the cosmic microwave background (CMB).

There is an extensive literature on speculative new physics that leads to parity violation in cosmology, e.g. Refs. [2–4], and more specifically in the gravitational sector [5]. But an asymmetry is perfectly compatible with general relativity, without the need to invoke exotic interactions, as Stueckelberg first pointed out [6]. The physics of the weak interaction contains all the necessary elements.

We consider the standard cosmological model with the sole addition of a new gauge field as a toy model, with an action given by

$$S = \int d^4x \sqrt{-g} \left( \frac{1}{2} M_P^2 R + \mathcal{L}_m - \frac{1}{4} F_{I\mu\nu} F^{I\mu\nu} \right) \quad (1)$$

$$F_{\mu\nu}^I \equiv \partial_\mu A_\nu^I - \partial_\nu A_\mu^I - g_{\text{YM}} \epsilon^I_{JK} A_\mu^J A_\nu^K$$

where Greek letters represent space-time indices, lower case Latin letters are spatial indices, and upper case Latin letters  $I \in \{1, 2, 3\}$  are reserved for the SU(2) indices. We assume a *flavor-space locked* configuration for the gauge field, wherein  $A_i^I = \phi(\tau) \delta_i^I$ , so that the directions of the internal group space are aligned with the principle spatial axes of the Robertson-Walker space-time,  $ds^2 = a^2(\tau)(-d\tau^2 + d\vec{x}^2)$  modulo SU(2) transformations. This approach yields an isotropic stress-energy tensor for the gauge field [7], that is invariant under gauge transformations in the invariant SU(2) space. The scalar potential equation of motion is  $\phi'' + 2g_{\text{YM}}^2 \phi^3 = 0$  where the prime indicates derivative with respect to conformal time, with a well-known solution in terms of the




Figure 1. A single right-handed circularly polarized gravitational wave, in this case with wavelength equal to the radius of the surface of last scattering and propagating in the vertical direction, leaves a left-handed spiral pattern of temperature anisotropy and polarization excess on the cosmic microwave background (CMB) sky. In our scenario, the parity reversed wave is damped by the flavor-space locked gauge field, so that the “mirror image” is a right-handed spiral pattern of much weaker temperature and polarization. There is a curl pattern around the zero temperature swath passing diagonally across the equator.

Jacobi elliptic sine-amplitude function [8],  $g_{\text{YM}}\phi(\tau) = c_1 \text{sn}(c_1(\tau - \tau_i) + c_2 | -1)$ . The constants are determined at the initial time  $\tau_i$  as  $c_1^4 = g_{\text{YM}}^2(\phi_i'^2 + g_{\text{YM}}^2 \phi_i^4)$  and  $c_2 = F(\theta | -1)$ , an elliptic integral of the first kind, with  $\text{csc } \theta_i = 1 + \phi_i'^2 / g_{\text{YM}}^2 \phi_i^4$ . The homogeneous, isotropic energy density and pressure of this Yang-Mills fluid are  $\rho_{\text{YM}} = 3p_{\text{YM}} = 3(\phi_i'^2 + g_{\text{YM}}^2 \phi_i^4) / 2a^4 = 3c_1^4 / 2g_{\text{YM}}^2 a^4$ . The gauge field oscillates with period  $\tau = \Gamma(\frac{1}{4})^2 / \sqrt{2\pi} c_1$ , but

its energy density and pressure scale with equation of state  $w = 1/3$ , like radiation. The coupling we consider is exponentially small,  $g_{\text{YM}} \sim \mathcal{O}(H_0/M_P) \sim 10^{-60}$ . An origin for this small coupling as well as the flavor-space locked configuration could be ascribed to an inflationary epoch. For example inflation would dilute the anisotropic field configurations much faster than the isotropic ones. Also given a dilaton-like coupling to the gauge field  $e^\sigma F^2$ , a dilaton can roll high up  $e^\sigma$  without additional cost since  $F^2$  vanishes under equipartition of energy between electric and magnetic modes. This large prefactor can be absorbed into the field and coupling to yield this suppressed constant  $g_{\text{YM}}$ . Inflationary scenarios based on a similar gauge field, but requiring higher order couplings to matter fields, have been studied elsewhere [9, 10].

The parity violation is manifest in the coupling to gravitational waves. When a gravitational wave passes through the gauge field it induces a quadrupolar distortion, squeezing and stretching the stress and energy of the gauge field. However, the gauge field itself possesses a preferred handedness via the right-handed SU(2) structure constants, so the gauge field stress-energy will vibrate in sympathy to a right-handed wave, and with antipathy to a left-handed wave, somewhat like a rattleback top [11]. In detail, we perturb the metric and gauge field  $\delta g_{\mu\nu} = a^2(\tau)h_{\mu\nu}$ ,  $\delta A_{I\mu} = a(\tau)M_P t_{ij}\delta_I^i\delta_\mu^j$  where  $t_{ij}$  and  $h_{ij}$  are transverse, traceless, synchronous tensors, following Refs. [12, 13] that are invariant under generalized covariance and SU(2) gauge transformations. The equations of motion for the Fourier amplitudes of a right circularly polarized gravitational wave traveling in the  $+z$  direction, and the corresponding gauge field fluctuation, are given by

$$\begin{aligned} h_R'' + 2\frac{a'}{a}h_R' + \left[ k^2 + \frac{2}{a^2M_P^2}(g_{\text{YM}}^2\phi^4 - \phi'^2) \right] h_R &= \\ -\frac{2}{aM_P} \left[ (k - g_{\text{YM}}\phi)g_{\text{YM}}\phi^2 t_R + \frac{a'}{a}\phi' t_R + \phi' t_R' \right], \\ t_R'' + 2\frac{a'}{a}t_R' + \left[ k^2 + \frac{a''}{a} - 2kg_{\text{YM}}\phi \right] t_R &= \\ -\frac{2}{aM_P} [(k + g_{\text{YM}}\phi)g_{\text{YM}}\phi^2 h_R - \phi' h_R']. \end{aligned} \quad (2)$$

Upon the exchange  $k \rightarrow -k$ , the equations now describe a left circularly polarized gravitational wave, but with very different consequences for the evolution — circular dichroism — as can be seen by examining the change in the effective mass term  $-2g_{\text{YM}}\phi$  for  $t$  and the coupling between  $h$  and  $t$ . For clarity, we have omitted the anisotropic shear contributed by other species, such as photons and neutrinos, although these effects are included in our cosmic microwave background (CMB) analysis. There is a rich variety of behavior in the evolution of this system, dependent upon the coupling  $g_{\text{YM}}$ , the abundance  $R_{\text{YM}} = \rho_{\text{YM}}/\rho_{\text{rad}}$  during the radiation era, the relative contributions of electric and magnetic field




Figure 2. The gravitational wave energy density spectrum is shown as a function of comoving wavenumber. An ambidextrous, scale free spectrum at an inflationary scale  $H_I = 10^{-5}M_P$  is assumed. Top: The present-day spectrum in the case with  $g_{\text{YM}} = 0$ ,  $R_{\text{YM}} = 0.03$  displays large oscillatory features due to the coupling between the gravitational waves and the gauge field. For comparison, the standard case without the gauge field is shown, as well as the effect of Standard Model particle free-streaming and freeze-outs (taken from Ref. [15]). Bottom: The case  $g_{\text{YM}} = 10^{-60}$ ,  $R_{\text{YM}} = 0.03$  is shown. The long wavelength modes are unaffected given our choice of initial conditions in the background field.

energy,  $\phi'$  and  $g_{\text{YM}}\phi^2$ , and the initial conditions for the perturbations  $h_{R/L}$ ,  $t_{R/L}$ .

The gravitational wave spectral density  $\Omega_{\text{GW}}$  is shown in Fig. 2 for several different cases, where the amplification and periodic modulation due to the gauge field are clearly seen. A WKB solution [14] predicts a peak or dip in the spectral density every five orders of magnitude in  $k$  for  $R_{\text{YM}} = 0.03$  (an energy density comparable to  $\Delta N_\nu \simeq 0.2$ ). In the ambidextrous case  $g_{\text{YM}} = 0$ , shown in the top panel, the negative effective mass in the doubles the gravitational wave amplitude at horizon entry, and the influence of the gauge field on the spectrum is seen to be much larger than the effect of photon and neutrino free-streaming, or brief departures from a pure radiation background when particle species become non-relativistic. In the lower panel, and for the remainder of this article, we investigate a minimal scenario in which the initial field energy of the YM fluid is split equally between the electric and magnetic field,  $\phi' = g_{\text{YM}}\phi^2$ . We further assume equal amplitude scale-free primordial spectra of left- and right-handed gravitational waves. The subsequent background evolution causes  $(g_{\text{YM}}^2\phi^4 - \phi'^2)/a^4$  to become non-zero and grow relative to the energy density  $\rho_{\text{YM}}$  in a time scale given by the period of oscillation of  $\phi$ . If the coupling  $g_{\text{YM}}$  is sufficiently small, as for the case illustrated in Fig. 2, then

this time scale is longer than the age of the universe, and super horizon modes are unaffected by the gauge field. If the coupling is large enough so that the oscillations begin before the present day, then the effective mass will lead to a suppression of order unity, depending on the abundance  $R_{\text{YM}}$ , of all super horizon modes. We note that such suppression, which can also be achieved through a different choice of initial conditions for  $\phi$ ,  $\phi'$  could modify the predictions of the gauge-flation and chromo-natural inflation scenarios [9, 10, 13].

The difference in the evolution for left- and right-circularly polarized waves is primarily due to the effective mass term for the gauge field tensor perturbations,  $-2kg_{\text{YM}}\phi$ , which is tachyonic for right-handed modes. The growth (suppression) of  $t_R$  ( $t_L$ ) is transferred to  $h_R$  ( $h_L$ ) as the mode enters the horizon. Once the relative amplitude is locked in at horizon entry and the fields begin to oscillate rapidly, the slow exchange of amplitude between  $h_{R/L}$  and  $t_{R/L}$  again comes into play. A WKB analysis for sub-horizon modes shows that  $h_{R/L}^2 + t_{R/L}^2 \propto 1/a^2$  and the exchange is oscillatory with similar phase if the background field is not yet oscillatory.

To evaluate the impact of this scenario on the CMB, we implement the scalar and tensor perturbations of the gauge field into CAMB [16]. The gauge field has the biggest impact on tensor correlations. The scalar sector only receives small corrections due to the anisotropic scalar shear of the gauge field. We ignore the vector perturbations which may be shown to decay [14]. Here  $R_{\text{YM}}$  is the ratio between the YM fluid density and the total relativistic energy density. We assume the fraction of critical density in the relativistic fluid is fixed by slightly adjusting the number of neutrino degrees of freedom upon introducing the gauge field. We otherwise assume standard  $\Lambda$ CDM parameters.

The CMB polarization can be decomposed into gradient  $E$ -modes and curl  $B$ -modes. In the tensor sector the gauge field introduces two main effects. First, the left- and right-handed contributions to the  $BB$  spectrum now differ, as shown in Fig. 3. Hence, the temperature and polarization anisotropy due to gravitational waves on roughly degree scales is dominated by a superposition of right-circularly polarized gravitational waves, which imprint left-helical patterns, similar to the display in Fig. 1. Interestingly, the individual contributions deviate strongly from  $\Lambda$ CDM but conspire in a way that puts the combination of both close to the expected standard cosmology result. Second, even though temperature  $T$  and gradient  $E$  polarization are both parity even and curl polarization is parity odd, the parity violation introduced by the YM fluid allows for correlations between  $TB$  and  $EB$  [3]. Detecting these exotic cross correlations is a smoking gun for chiral effects in the universe. Predictions of our model are plotted in Fig. 4.

There are many challenges to detecting the parity-violating cross correlations, not to mention the  $B$ -mode




Figure 3. CMB  $BB$  polarization autocorrelation spectra. The pure left- and right-handed contributions deviate strongly from  $\Lambda$ CDM (dashed) while their sum is closer to it. Solid lines include a gauge field with  $g_{\text{YM}} = 10^{-56}$ ,  $R_{\text{YM}} = 0.1$  and tensor-to-scalar ratio of  $r = 0.1$ . The thin line shows the  $BB$  spectrum for the case  $g_{\text{YM}} = 4 \times 10^{-56}$ .

signal. Galactic foregrounds, magnetic fields, weak lensing, and other systematic effects can all produce a false positive; fortunately there is no fundamental barrier that would prevent a detection that can distinguish a primordial signal. (See Ref. [17] for a recent summary.) But there are other phenomena that could produce a parity-violating signal. First of all, cosmological birefringence (CB) can lead to  $TB$  and  $EB$  power spectra by rotating  $E$  into  $B$  through a novel coupling between electromagnetism and a cosmic pseudoscalar such as quintessence [18]. A second possibility, broadly characterized as chiral gravity, posits a modification of gravity whereby an asymmetry between left- and right-circularly polarized waves is imprinted on the primordial spectrum. The third possibility, as we have shown, is essentially cosmic circular dichroism, whereby the asymmetry develops with time from an initially symmetric primordial spectrum.

Would an actual detection directly point to chiral symmetry breaking on cosmological scales? In Ref. [19] it was shown that the  $TB$  and  $EB$  spectra can be used to distinguish CB effects from chiral physics. As CB rotates the  $E$  into a  $B$  contribution the measured  $B$  spectrum would resemble the  $E$  one which makes this separation into CB and chirality effects feasible. In turn, putting limits on the amplitude of these spectra will put constraints on chiral physics in general and our model in particular.

The deviations seen in the CMB spectra would clearly have an impact on the interpretation of a precision measurement of  $B$  modes [20]. As can be seen from Fig. 3, the gauge field can vary the height of the  $BB$  spectrum at the reionization bump near  $\ell \lesssim 10$  and at the primary acoustic peak near  $\ell \sim 100$  by as much as  $\pm 50\%$ . However, we have the greatest leverage on new physics by focusing on the exotic cross correlations. Hence, we forecast the parameter constraints  $\sigma_{R_{\text{YM}}}$  and  $\sigma_{g_{\text{YM}}}$  using Fisher matrix techniques on  $TB$  and  $EB$ , for which the




Figure 4. CMB temperature and polarization auto- and cross-correlation power spectra. The parity violation allows for  $TB$  and  $EB$  cross correlations. The dashed lines represent the standard  $\Lambda$ CDM cosmology, solid lines include a gauge field with  $g_{\text{YM}} = 10^{-56}$  and  $R_{\text{YM}} = 0.1$  and tensor-to-scalar ratio of  $r = 0.1$ . The  $TB$  and  $EB$  cross correlations only appear when the gauge field is present.

Fisher matrix reads

$$\mathcal{F}_{ij} = \sum_l \sum_{X,Y} \frac{\partial C_l^X}{\partial \theta_i} \frac{\partial C_l^Y}{\partial \theta_j} [\Xi_l^{-1}]_{XY} \quad (3)$$

where  $\vec{\theta} = (R_{\text{YM}}, g_{\text{YM}}) + \vec{\theta}_{\text{cosmo}}$  and  $X, Y = \{TB, EB\}$ . Here, the eight cosmological parameters are  $\vec{\theta}_{\text{cosmo}} = (\omega_b, \omega_c, \omega_\nu, \Omega_K, H_0, w, n_t, r)$ . The Fisher matrix  $\mathcal{F}$  is the inverse of the covariance matrix between  $R_{\text{YM}}$  and  $g_{\text{YM}}$ . The derivatives of the  $C_l^X$  are obtained using CAMB. These derivatives are centered around a fiducial model: we choose  $g_{\text{YM}} = 10^{-56}$ ,  $R_{\text{YM}} = 0.1$  and the standard Planck  $\Lambda$ CDM values for the cosmological parameters [21]. The matrix  $[\Xi_l^{-1}]_{XY}$  is the inverse of the  $TB$ - $EB$  covariance matrix. Details of this calculation can be found in [19]. The parameters used are taken from Ref. [19, 22] and summarized in Tab. I.

The 1D marginalized confidence limits in a scenario in which the Planck satellite measures  $TB$  and  $EB$  correlations are  $\sigma_{g_{\text{YM}}} = 9.5 \times 10^{-57}$  and  $\sigma_{R_{\text{YM}}} = 0.030$ . For the cosmic variance (CV) limited experiment these numbers reduce to  $\sigma_{g_{\text{YM}}} = 3.4 \times 10^{-57}$  and  $\sigma_{R_{\text{YM}}} = 8.1 \times 10^{-3}$ , which would be able to make a detection of  $g_{\text{YM}}$  feasible. The 1- and 2- $\sigma$  contours are plotted in Fig. 5. For

Instrument	$\theta_{\text{FWHM}}$ [arcmin]	NET [ $\mu\text{K}\sqrt{\text{s}}$ ]	$t_{\text{obs}}$ [y]
Planck	7.1	45	2
CV limited	5	0	1.2

Table I. Instrumental parameters for the two experiments considered in this paper. The parameters are the beamwidth  $\theta_{\text{FWHM}}$ , noise-equivalent temperature NET, and observation time  $t_{\text{obs}}$ .




Figure 5. Forecasted 1- and 2- $\sigma$  C.L. contours under the condition that  $TB$  and  $EB$  cross correlations are detected by the respective experiments (indicated by the asterisk \*). The fiducial model is indicated by the black dot which represents  $g_{\text{YM}} = 10^{-56}$  and  $R_{\text{YM}} = 0.1$ .

this fiducial model Planck could make a 2- $\sigma$  detection of  $R_{\text{YM}}$ , but cannot exclude the ambidextrous case since  $g_{\text{YM}} = 0$  lies within its 1- $\sigma$  contour. The future looks brighter for a future satellite mission that gets closer to a CV limited experiment, which could put constraints on the chiral asymmetry; for the fiducial model, the coupling  $g_{\text{YM}}$  could be distinguished from zero at better than the two sigma level. Such a limit could be used to determine whether the gauge field is part of a dark sector that includes dark energy. If dark energy couples to the rolling gauge field, or if the gauge field is dark energy, as in a gauge-flation scenario, then the rate of cosmic acceleration may be linked to the chiral asymmetry.

Concluding, we present a simple model that breaks parity on cosmological scales. We illustrate the impact of this model on the gravitational wave spectrum and the CMB, computing the power spectra along with new  $TB$  and  $EB$  correlations that emerge in parity-breaking models. A detection of one of these correlations could be the sign of a flavor-space locked gauge field.

This work is supported in part by DOE grant DE-SC0010386. We thank Vera Gluscevic and Steve Gubser for useful conversations.

- 
- [1] C. S. Wu, E. Ambler, R. W. Hayward, D. D. Hoppes and R. P. Hudson, Phys. Rev. **105**, 1413 (1957).
- [2] S. M. Carroll, G. B. Field and R. Jackiw, Phys. Rev. D **41**, 1231 (1990).
- [3] A. Lue, L. M. Wang and M. Kamionkowski, Phys. Rev. Lett. **83**, 1506 (1999).
- [4] C. R. Contaldi, J. Magueijo and L. Smolin, Phys. Rev. Lett. **101**, 141101 (2008).
- [5] S. Alexander and N. Yunes, Phys. Rept. **480**, 1 (2009).
- [6] E. C. G. Stueckelberg, Phys. Rev. **106**, 388 (1957).
- [7] C. Armendariz-Picon, JCAP **0407**, 007 (2004).
- [8] P. F. Byrd and M. D. Friedman, "Handbook of elliptic integrals for engineers and scientists," (1971) Springer Verlag, pp. 358.
- [9] A. Maleknejad and M. M. Sheikh-Jabbari, Phys. Lett. B **723**, 224 (2013).
- [10] P. Adshead and M. Wyman, Phys. Rev. Lett. **108**, 261302 (2012).
- [11] [en.wikipedia.org/wiki/Rattleback](https://en.wikipedia.org/wiki/Rattleback)
- [12] E. Dimastrogiovanni and M. Peloso, Phys. Rev. D **87**, no. 10, 103501 (2013).
- [13] R. Namba, E. Dimastrogiovanni and M. Peloso, JCAP **1311**, 045 (2013).
- [14] J. Bielefeld and R. R. Caldwell, *in preparation*.
- [15] Y. Watanabe and E. Komatsu, Phys. Rev. D **73**, 123515 (2006).
- [16] A. Lewis, A. Challinor and A. Lasenby, Astrophys. J. **538**, 473 (2000).
- [17] J. Kaufman, B. Keating and B. Johnson, arXiv:1409.8242 [astro-ph.CO].
- [18] S. M. Carroll, Phys. Rev. Lett. **81**, 3067 (1998).
- [19] V. Gluscevic and M. Kamionkowski, Phys. Rev. D **81**, 123529 (2010).
- [20] P. A. R. Ade *et al.* [BICEP2 Collaboration], Phys. Rev. Lett. **112**, 241101 (2014).
- [21] P. A. R. Ade *et al.* [Planck Collaboration], Astron. Astrophys. **571**, A16 (2014).
- [22] P. A. R. Ade *et al.* [Planck Collaboration], Astron. Astrophys. **571**, A6 (2014) [arXiv:1303.5067 [astro-ph.CO]].