



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Post-merger evolution of a neutron star-black hole binary with neutrino transport

Francois Foucart, Evan O'Connor, Luke Roberts, Matthew D. Duez, Roland Haas, Lawrence E. Kidder, Christian D. Ott, Harald P. Pfeiffer, Mark A. Scheel, and Bela Szilagyi

Phys. Rev. D **91**, 124021 — Published 11 June 2015

DOI: [10.1103/PhysRevD.91.124021](https://doi.org/10.1103/PhysRevD.91.124021)

Post-merger evolution of a neutron star-black hole binary with neutrino transport

Francois Foucart,^{1,2,3} Evan O’Connor,^{4,3} Luke Roberts,^{2,5} Matthew D. Duez,⁶ Roland Haas,^{5,7} Lawrence E. Kidder,⁸ Christian D. Ott,⁵ Harald P. Pfeiffer,^{3,9} Mark A. Scheel,⁵ and Bela Szilagyi⁵

¹*Lawrence Berkeley National Laboratory, 1 Cyclotron Rd, Berkeley, CA 94720, USA*

²*NASA Einstein Fellow*

³*Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, Ontario M5S 3H8, Canada*

⁴*Department of Physics, North Carolina State University, Raleigh, North Carolina 27695, USA; Hubble Fellow*

⁵*TAPIR, Walter Burke Institute for Theoretical Physics, MC 350-17, California Institute of Technology, Pasadena, California 91125, USA*

⁶*Department of Physics & Astronomy, Washington State University, Pullman, Washington 99164, USA*

⁷*Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut, D-14476 Golm, Germany*

⁸*Center for Radiophysics and Space Research, Cornell University, Ithaca, New York, 14853, USA*

⁹*Canadian Institute for Advanced Research, 180 Dundas St. West, Toronto, ON M5G 1Z8, Canada*

We present a first simulation of the post-merger evolution of a black hole-neutron star binary in full general relativity using an energy-integrated general relativistic truncated moment formalism for neutrino transport. We describe our implementation of the moment formalism and important tests of our code, before studying the formation phase of an accretion disk after a black hole-neutron star merger. We use as initial data an existing general relativistic simulation of the merger of a neutron star of mass $1.4M_{\odot}$ with a black hole of mass $7M_{\odot}$ and dimensionless spin $\chi_{\text{BH}} = 0.8$. Comparing with a simpler leakage scheme for the treatment of the neutrinos, we find noticeable differences in the neutron to proton ratio in and around the disk, and in the neutrino luminosity. We find that the electron neutrino luminosity is much lower in the transport simulations, and that both the disk and the disk outflows are less neutron-rich. The spatial distribution of the neutrinos is significantly affected by relativistic effects, due to large velocities and curvature in the regions of strongest emission. Over the short timescale evolved, we do not observe purely neutrino-driven outflows. However, a small amount of material ($3 \times 10^{-4}M_{\odot}$) is ejected in the polar region during the circularization of the disk. Most of that material is ejected early in the formation of the disk, and is fairly neutron rich (electron fraction $Y_e \sim 0.15 - 0.25$). Through r-process nucleosynthesis, that material should produce high-opacity lanthanides in the polar region, and could thus affect the lightcurve of radioactively powered electromagnetic transients. We also show that by the end of the simulation, while the bulk of the disk remains neutron-rich ($Y_e \sim 0.15 - 0.2$ and decreasing), its outer layers have a higher electron fraction: 10% of the remaining mass has $Y_e > 0.3$. As that material would be the first to be unbound by disk outflows on longer timescales, and as composition evolution is slower at later times, the changes in Y_e experienced during the formation phase of the disk could have an impact on nucleosynthesis outputs from neutrino-driven and viscously-driven outflows. Finally, we find that the effective viscosity due to momentum transport by neutrinos is unlikely to have a strong effect on the growth of the magnetorotational instability in the post-merger accretion disk.

PACS numbers: 04.25.dg, 04.40.Dg, 26.30.Hj, 98.70.-f

I. INTRODUCTION

The likely detection of gravitational waves by the advanced LIGO/VIRGO detectors [1, 2] in the coming years will open up an entirely new way to observe the universe, complementing existing electromagnetic and neutrino observations. Mergers of black holes and neutron stars are expected to be among the first and most common sources of gravitational waves to be observed [3]. Beyond the excitement associated with the first gravitational wave detection, binary mergers will provide us with a wealth of information which could constrain the formation and evolution of massive binaries [4, 5], the outcomes of core collapse supernovae, or the properties of the cold, dense neutron-rich matter in the core of neutron stars [6–9].

In the presence of at least one neutron star, the gravitational wave signal may be accompanied by electromagnetic and neutrino emissions. The joint detection of a system by a gravitational wave observatory and an electromagnetic telescope would provide additional information about the location, environment, and parameters of the binary [10]. It could also

help us understand some important astrophysical processes. Neutron star mergers are among the most likely progenitors of short-hard gamma-ray bursts. They may also significantly contribute to the production of heavy elements in the universe, through r-process nucleosynthesis in the neutron-rich material unbound during some mergers [11–15]. This nucleosynthesis could be observed through radioactively powered transients, observable in the optical and/or infrared bands days after the merger (“kilonovae”) [16, 17].

To understand which signals can be emitted by a merger, and how they depend on the initial parameters of the binary, we need numerical simulations in full general relativity: the strong non-linearities in the evolution of the spacetime surrounding the binary make any approximate treatment of gravity unreliable. Most general relativistic simulations focus on a very short period around the time of merger (~ 100 ms for simulations involving neutron stars), when general relativistic effects are important. Over that time period and beyond gravity, the most important physical effects to take into account are the equation of state of neutron star matter, the effects of magnetic fields, and the cooling and composition evolution due to

neutrino-matter interactions. Although numerical simulations of these effects have significantly improved over the last few years, much work is needed to simulate them well-enough to reliably predict the post-merger signals produced by compact binary mergers. The main object of this work is the development and testing of an improved method for the treatment of neutrinos in the SpEC code [18], based on the moment formalism of Thorne [19], and its application to the post-merger evolution of a black hole-neutron star binary.

In terms of the neutron star equations of state, most general relativistic simulations use either a simple Gamma-law equation of state, or parametrized piecewise-polytropic equations of state with a thermal Gamma-law. When modeling the gravitational wave signal up to the point of merger, this should be sufficient: the gravitational wave signal mostly probes a single parameter of the equation of state, its tidal deformability, while the detailed structure of the neutron star appears unimportant [7, 8]. But the equation of state plays a crucial role when simulating the merger and post-merger evolution of the binary [20, 21]. Many systems have been evolved with hot nuclear-theory based equations of state using Newtonian or pseudo-Newtonian potentials (e.g. [22–27]). But only a few general relativistic simulations have used cold [21, 28, 29] or hot [15, 30–33] tabulated, nuclear-theory based equations of state, and of those only the equation of state used in [15] appears consistent with the most recent models for dense neutron-rich matter [34, 35]. In this work, we use a hot, composition dependent, nuclear-theory based equation of state by Lattimer & Swesty [36], which leads to an acceptable mass-radius relationship for neutron stars and allows us to include neutrino-matter interactions, but is not fully consistent with the latest constraints from nuclear experiments.

The simulation of magnetic fields has seen some impressive recent improvements. General relativistic simulations of binary neutron star mergers assuming ideal magnetohydrodynamics have been performed with a number of codes (see e.g. [37] for a recent review of the field), but resolving the growth of magnetic instabilities at an acceptable computational cost remains an open problem [38]. Simulations using force-free or resistive magnetohydrodynamics, needed to properly simulate magnetically dominated regions, have also been performed [39–42]. So far they have, however, mostly studied the pre-merger evolution of the system. Here, we do not take the magnetic fields into account at all, and focus solely on neutrino effects.

The third component needed to simulate compact binaries around the time of merger, neutrinos, is also the most computationally expensive to treat accurately. In theory, for each species of neutrinos, one should evolve the distribution function of neutrinos in both physical and momentum space. The high dimensionality of the problem places it out of reach of current numerical codes and computers. Accordingly, various approximations have been developed - most of them in Newtonian simulations and/or with the aim of studying core collapse supernovae. In general relativistic simulations of binary mergers, a few simulations have used a simple cooling prescription (“leakage”) to model the first-order impact of neutrinos on the cooling of the disk and the composition evolution

of high-density regions [31–33]. The only published result using a more advanced method [15] studied a binary neutron star merger using an energy-integrated (“gray”) version of the moment formalism for radiation transport [19, 43, 44]. The first two moments of the neutrino distribution function (i.e. the energy density and flux of neutrinos) were evolved, and the analytical M1 closure was used to compute the third moment (pressure tensor). In this work, we discuss the implementation of a similar scheme within the SpEC code, and its application to the post-merger evolution of a black hole-neutron star merger. As in [15], we use a gray M1 scheme. We should note that there are significant differences between our implementation of the M1 formalism and the one used in [15], in particular in the treatment of optically thick regions¹, which would make a comparison between the results of the two codes an interesting test of the accuracy of the M1 formalism in the core of the disk or in the presence of a hypermassive neutron star. However, we will not discuss this here, as we are considering a completely different physical setup for the evolution, and the details of the M1 implementation used in [15] have not yet been published.

The focus of this paper is to discuss three different questions. First, we describe the implementation and testing of the M1 formalism in SpEC. An overview of the methods is offered in Sec. II, while the interested reader will find the details of the algorithm and important tests of the code in the Appendices. Second, we simulate the evolution of a black hole-neutron star binary in the critical phase in which the accretion disk is formed, from the moment at which neutrino emission increases due to the heating of the forming disk, to the time at which the disk reaches a quasi-equilibrium state (about 20 ms after merger). We pay particular attention to the temperature and composition evolution, the ejection of matter along the spin axis of the black hole, the geometry of the neutrino radiation, and the potential astrophysical consequences of our results. This is the main focus of Secs. IV–V. And third, we study the impact of using the M1 formalism instead of the simpler leakage scheme, and the impact of the various approximations which have to be made when using a gray scheme. This is the focus of Sec. IV G.

II. MOMENT FORMALISM

We will start here with an overview of the M1 formalism, and of its numerical implementation in SpEC. A more detailed description of many aspects of the algorithm, as well as tests of our implementation, are available in the appendices.

A. Evolution equations

For each neutrino species ν_i , we can describe the neutrinos by their distribution function $f_{(\nu)}(x^\alpha, p^\alpha)$, where $x^\alpha =$

¹ M. Shibata, Y. Sekiguchi, private communication

(t, x^i) gives the time and the position of the neutrinos, and p^α is the 4-momentum of the neutrinos. The distribution function $f_{(\nu)}$ evolves according to the Boltzmann equation

$$p^\alpha \left[\frac{\partial f_{(\nu)}}{\partial x^\alpha} - \Gamma_{\alpha\gamma}^\beta p^\gamma \frac{\partial f_{(\nu)}}{\partial p^\beta} \right] = \left[\frac{df_{(\nu)}}{d\tau} \right]_{\text{coll}}, \quad (1)$$

where the $\Gamma_{\beta\gamma}^\alpha$ are the Christoffel symbols and the right-hand side includes all collisional processes (emissions, absorptions, scatterings). In general, this is a 7-dimensional problem which is extremely expensive to solve numerically. Approximations to the Boltzmann equation have thus been developed for numerical applications. In this work, we consider the moment formalism developed by Thorne [19], in which only the lowest moments of the distribution function in momentum space are evolved. Our code is largely inspired by the implementation of Thorne's formalism into general relativistic hydrodynamics simulations proposed by Shibata *et al.* [43] and Cardall *et al.* [44]. We limit ourselves to the use of this formalism in the ‘‘gray’’ approximation, that is we only consider energy-integrated moments. Although the moment formalism can in theory be used with a discretization in neutrino energies, this makes the simulations significantly more expensive and involves additional technical difficulties in the treatment of the gravitational and velocity redshifts. We will also only consider three independent neutrino species: the electron neutrinos ν_e , the electron antineutrinos $\bar{\nu}_e$, and the heavy-lepton neutrinos ν_x . The latter is the combination of 4 species ($\nu_\mu, \bar{\nu}_\mu, \nu_\tau, \bar{\nu}_\tau$). This merging is justified because the temperatures and neutrino energies reached in our merger calculations are low enough to suppress the formation of the corresponding heavy leptons whose presence would require including the charged current neutrino interactions that differentiate between these individual species.

In the gray approximation, and considering only the first two moments of the distribution function, we evolve for each species projections of the stress-energy tensor of the neutrino radiation $T_{\text{rad}}^{\mu\nu}$. One possible decomposition of $T_{\text{rad}}^{\mu\nu}$ is [43]

$$T_{\text{rad}}^{\mu\nu} = Ju^\mu u^\nu + H^\mu u^\nu + H^\nu u^\mu + S^{\mu\nu}, \quad (2)$$

with $H^\mu u_\mu = S^{\mu\nu} u_\mu = 0$ and u^μ the 4-velocity of the fluid. The energy J , flux H^μ and stress tensor $S^{\mu\nu}$ of the neutrino radiation as observed by an observer comoving with the fluid are related to the neutrino distribution function by

$$J = \int_0^\infty d\nu \nu^3 \int d\Omega f_{(\nu)}(x^\alpha, \nu, \Omega), \quad (3)$$

$$H^\mu = \int_0^\infty d\nu \nu^3 \int d\Omega f_{(\nu)}(x^\alpha, \nu, \Omega) l^\mu, \quad (4)$$

$$S^{\mu\nu} = \int_0^\infty d\nu \nu^3 \int d\Omega f_{(\nu)}(x^\alpha, \nu, \Omega) l^\mu l^\nu, \quad (5)$$

where ν is the neutrino energy in the fluid frame, $\int d\Omega$ denotes integrals over solid angle on a unit sphere in momentum space, and

$$p^\alpha = \nu(u^\alpha + l^\alpha), \quad (6)$$

with $l^\alpha u_\alpha = 0$ and $l^\alpha l_\alpha = 1$. We also consider the decomposition of $T_{\text{rad}}^{\mu\nu}$ in terms of the energy, flux and stress tensor observed by an inertial observer,

$$T_{\text{rad}}^{\mu\nu} = En^\mu n^\nu + F^\mu n^\nu + F^\nu n^\mu + P^{\mu\nu}, \quad (7)$$

with $F^\mu n_\mu = P^{\mu\nu} n_\mu = F^t = P^{t\nu} = 0$, and n^α the unit normal to a $t = \text{constant}$ slice.

We use the 3+1 decomposition of the metric,

$$ds^2 = g_{\alpha\beta} dx^\alpha dx^\beta \quad (8)$$

$$= -\alpha^2 dt^2 + \gamma_{ij}(dx^i + \beta^i)(dx^j + \beta^j), \quad (9)$$

where α is the lapse, β^i the shift, and γ_{ij} the 3-metric on a slice of constant coordinate t . The extension of γ_{ij} to the full 4-dimensional space is the projection operator

$$\gamma_{\alpha\beta} = g_{\alpha\beta} + n_\alpha n_\beta. \quad (10)$$

We similarly define a projection operator onto the reference frame of an observer comoving with the fluid,

$$h_{\alpha\beta} = g_{\alpha\beta} + u_\alpha u_\beta. \quad (11)$$

We can then write equations relating the fluid-frame variables to the inertial frame variables [44]:

$$E = W^2 J + 2W v_\mu H^\mu + v_\mu v_\nu S^{\mu\nu}, \quad (12)$$

$$F_\mu = W^2 v_\mu J + W(g_{\mu\nu} - n_\mu v_\nu) H^\nu + W v_\mu v_\nu H^\nu + (g_{\mu\nu} - n_\mu v_\nu) v_\rho S^{\nu\rho}, \quad (13)$$

$$P_{\mu\nu} = W^2 v_\mu v_\nu J + W(g_{\mu\rho} - n_\mu v_\rho) v_\nu H^\rho + W(g_{\rho\nu} - n_\rho v_\nu) v_\mu H^\rho + (g_{\mu\rho} - n_\mu v_\rho)(g_{\nu\kappa} - n_\nu v_\kappa) S^{\rho\kappa}, \quad (14)$$

using the decomposition of the 4-velocity

$$u^\mu = W(n^\mu + v^\mu), \quad (15)$$

with $v^\mu n_\mu = 0$ and $W = \sqrt{1 + \gamma^{ij} u_i u_j}$.

Evolution equations for $\tilde{E} = \sqrt{\gamma} E$ and $\tilde{F}^i = \sqrt{\gamma} F^i$ can then be written in conservative form:

$$\partial_t \tilde{E} + \partial_j (\alpha \tilde{F}^j - \beta^j \tilde{E}) \quad (16)$$

$$= \alpha (\tilde{P}^{ij} K_{ij} - \tilde{F}^j \partial_j \ln \alpha - \tilde{S}^\alpha n_\alpha),$$

$$\partial_t \tilde{F}_i + \partial_j (\alpha \tilde{P}_i^j - \beta^j \tilde{F}_i) \quad (17)$$

$$= (-\tilde{E} \partial_i \alpha + \tilde{F}_k \partial_i \beta^k + \frac{\alpha}{2} \tilde{P}^{jk} \partial_i \gamma_{jk} + \alpha \tilde{S}^\alpha \gamma_{i\alpha}),$$

where γ is the determinant of γ_{ij} , $\tilde{P}_{ij} = \sqrt{\gamma} P_{ij}$, and \tilde{S}^α includes all collisional source terms.

To close this system of equations, we need two additional ingredients: a prescription for the computation of $P^{ij}(E, F_i)$ (‘‘closure relation’’, which we choose following Minerbo 1978 [45]), and the collisional source terms \tilde{S}^α . In the M1 formalism, the neutrino pressure tensor P^{ij} is recovered as an interpolation between its known limits for an optically thick medium and an optically thin medium with a unique direction of propagation for the neutrinos. We provide details on its computation in Appendix A. For the source

terms \tilde{S}^α , we will consider that the fluid has an energy-integrated emissivity $\bar{\eta}$ due to the charged-current reactions




as well as electron-positron pair annihilation



plasmon decay



and nucleon-nucleon Bremsstrahlung



The inverse reactions are responsible for an energy-averaged absorption opacity $\bar{\kappa}_a$. We also consider an energy-averaged scattering opacity $\bar{\kappa}_s$ due to elastic scattering of neutrinos on nucleons and heavy nuclei. The source terms are then

$$\tilde{S}^\alpha = \sqrt{\gamma} (\bar{\eta} u^\alpha - \bar{\kappa}_a J u^\alpha - (\bar{\kappa}_a + \bar{\kappa}_s) H^\alpha). \quad (23)$$

Details of the choices made for the computation of the gray $\bar{\eta}$, $\bar{\kappa}_a$ and $\bar{\kappa}_s$ are provided in Appendix B. Except for the modifications described in that Appendix, we use the emissivities and opacities proposed by Ruffert et al. [22] for all of the above reactions, except for nucleon-nucleon Bremsstrahlung for which the emissivity is computed following Burrows et al. [46].

B. Numerical scheme

We add the evolution of neutrinos with the moment scheme to the SpEC code [18], which already includes a general relativistic hydrodynamics module [47]. The latest methods used for evolving in SpEC the coupled system formed by Einstein's equation and the general relativistic equations of hydrodynamics are described in [48], Appendix A.

In the M1 formalism, we evolve the variables \tilde{E} and \tilde{F}_i according to Eqs. (16-17), and couple the neutrinos to the fluid evolution. The coupling takes the form

$$\partial_t \tau = \dots + \alpha \tilde{S}^\alpha n_\alpha, \quad (24)$$

$$\partial_t S_i = \dots - \alpha \tilde{S}^\alpha \gamma_{i\alpha}, \quad (25)$$

$$\partial_t (\rho_* Y_e) = \dots - \text{sign}(\nu_i) \alpha \sqrt{\gamma} \frac{\bar{\eta} - \bar{\kappa}_a J}{\langle \epsilon_\nu \rangle}, \quad (26)$$

where ρ_* , $\rho_* Y_e$, τ and S_i are the conservative hydrodynamics variables which are evolved in SpEC,

$$\rho_* = \rho_0 W \sqrt{\gamma}, \quad (27)$$

$$\tau = \rho_* (hW - 1) - \sqrt{\gamma} P, \quad (28)$$

$$S_i = \rho_* h u_i, \quad (29)$$

ρ_0 is the baryon density of the fluid, P its pressure, Y_e its electron fraction, and h its specific enthalpy. $\langle \epsilon_\nu \rangle$ is the weighted average energy of neutrinos, which should be

$$\langle \epsilon_\nu \rangle = \frac{\int_0^\infty [\bar{\eta}(\epsilon_\nu) - \kappa_a(\epsilon_\nu) J(\epsilon_\nu)] d\epsilon_\nu}{\int_0^\infty \frac{\bar{\eta}(\epsilon_\nu) - \kappa_a(\epsilon_\nu) J(\epsilon_\nu)}{\epsilon_\nu} d\epsilon_\nu}, \quad (30)$$

where $\eta(\epsilon_\nu)$ and $\kappa_a(\epsilon_\nu)$ are the emissivity and absorption of neutrinos at energy ϵ_ν . $\text{sign}(\nu_i)$ is 1 for ν_e , -1 for $\bar{\nu}_e$, and 0 for heavy-lepton neutrinos. Lacking the knowledge of the neutrino spectrum, we have to use an approximate form for $\langle \epsilon_\nu \rangle$. If we assume that the neutrinos follow a Fermi-Dirac distribution at the local temperature T of the fluid and with neutrino chemical potential μ_ν , and that the opacity follows $\kappa_a \propto \epsilon_\nu^2$ (as for the dominant charged-current reactions), $\langle \epsilon_\nu \rangle$ is

$$\langle \epsilon_\nu \rangle = \frac{F_5(\eta_\nu)}{F_4(\eta_\nu)} T \quad (31)$$

where

$$F_k(\eta_\nu) = \int_0^\infty \frac{x^k}{1 + \exp(x - \eta_\nu)} dx \quad (32)$$

is the Fermi integral, and $\eta_\nu = \mu_\nu/T$ is the degeneracy parameter. We use this value of $\langle \epsilon_\nu \rangle$ in hot / optically thick regions. Unfortunately, we are not aware of any simple way to make a similar estimate in optically thin regions. In merger simulations, these regions are generally colder than the optically thick regions from which most neutrinos are emitted, and the average energy of neutrinos there is thus higher than what one would expect by assuming equilibrium between the neutrinos and the fluid. We make a first order estimate of the effect of the deviations from the equilibrium spectrum by using

$$\langle \epsilon_\nu \rangle \approx \frac{F_5(\eta_\nu)}{F_4(\eta_\nu)} T \max \left(1, \sqrt{\frac{\langle \epsilon_{\nu,leak}^2 \rangle}{\langle \epsilon_{\nu,fluid}^2 \rangle}} \right). \quad (33)$$

Here, $\langle \epsilon_{\nu,leak}^2 \rangle$ is the global estimate of the average square energy of neutrinos obtained from the simpler leakage scheme [32], and $\langle \epsilon_{\nu,fluid}^2 \rangle$ would be the average square energy of the neutrinos if they obeyed the equilibrium Fermi-Dirac distribution (see Appendix B). In making this approximation, we are helped by the relative homogeneity of the fluid temperature around the neutrinospheres. This implies that there should be only moderate variations of the neutrino spectrum between different points in the optically thin region, making the use of the global average energy $\langle \epsilon_{\nu,leak}^2 \rangle$ better motivated.

For the applications that we are considering here, the back-reaction of the neutrinos onto the fluid evolution is weak (except for transients when we turn on neutrino emission). Accordingly, we separate the hydrodynamics and neutrino evolution. Our evolution scheme thus proceeds as follow:

- Evolve the Einstein equations and the general relativistic hydrodynamics equations, without taking neutrinos into account, over a time step Δt_H chosen as in [48], Appendix A.3.
- Evolve the neutrino radiation, potentially taking multiple time steps, so that the neutrinos are also evolved by Δt_H . The time step used to evolve the neutrinos is chosen as described in Appendix D. Each time step proceeds as follow:

- Reconstruct the fields $E, F_i/E$ at cell faces, using the *minmod* reconstruction method.
- Compute the closure relation at cell faces to get the fluxes $(\alpha \tilde{F}^i - \beta^i \tilde{E})$ and $(\alpha \tilde{P}_j^i - \beta^i \tilde{F}_j)$.
- Use those fluxes to compute the divergence terms in Eqs. (16-17), using the shock-capturing methods and corrections in high optical depth regions described in Appendix C.
- Compute the closure relation at cell centers.
- Compute the gravitational source terms on the right-hand side of Eqs. (16-17) [everything but the terms proportional to \tilde{S}^α] from E and F_i at the beginning of the neutrino step.
- Solve Eqs. (16-17) by treating implicitly the collisional source terms proportional to \tilde{S}^α , following the method described in Appendix D.
- Compute the coupling to the hydrodynamics variables, and update $(\tau, S_i, \rho_* Y_e)$ according to Eqs. (24-26).
- If we have evolved the neutrinos by Δt_H , go back to the GR-Hydro evolution. Otherwise, take the next neutrino time step.

A more complete description of the different steps of this algorithm is provided in Appendices A-D.

III. INITIAL CONDITIONS AND NUMERICAL SETUP

As a first astrophysical application of our code, we consider the disk formation phase of a black hole-neutron star merger from Foucart *et al.* 2014 [32]. This phase is particularly interesting to study with a general relativistic code and neutrino transport because general relativistic effects and the evolution of the metric remain important at this point. Furthermore, due to the high temperatures experienced by the fluid during disk formation, the neutrino luminosity is higher and the composition evolution is faster than at any other time. Finally the fluid is initially very close to the black hole, where relativistic effects cannot be neglected, and it is far from equilibrium. This phase of the evolution would thus not be properly captured by simulations which start from an equilibrium torus configuration, or which model general relativistic effects through the use of pseudo-Newtonian potentials.

In the specific merger that we are considering, the masses of the compact objects before merger are $M_{\text{NS}}^i = 1.4M_\odot$ and $M_{\text{BH}}^i = 7M_\odot$. The initial dimensionless spin of the black hole is $\chi_{\text{BH}}^i = 0.8$, and it is aligned with the orbital angular momentum of the binary. The neutron star is initially non-spinning. This is simulation M14-7-S8 of [32]. We showed in [32] that the disruption of the neutron star results in the ejection of about $0.06M_\odot$ of material, and the formation of an accretion disk of mass $M_{\text{disk}}^f \sim 0.1M_\odot$. The final properties of the black hole are $M_{\text{BH}}^f = 8M_\odot$ and $\chi_{\text{BH}}^f = 0.87$. We use the equation of state of Lattimer & Swesty [36] with nuclear incompressibility parameter $K_0 = 220$ MeV and symmetry energy $S_\nu = 29.3$ MeV, using a table available on

<http://www.stellarcollapse.org> and described in O’Connor & Ott 2010 [49]. For a neutron star of mass $M_{\text{NS}} = 1.4M_\odot$, this results in a neutron star radius $R_{\text{NS}} = 12.7$ km.

The post-merger configuration obtained as a result of the merger is expected to be fairly typical for black hole-neutron star mergers in which the neutron star is disrupted by the black hole. In [32], we studied the range of initial black hole masses $M_{\text{BH}}^i = (7-10)M_\odot$ and neutron stars masses $M_{\text{NS}}^i = (1.2-1.4)M_\odot$ currently deemed most likely from the observation of galactic stellar mass black holes [50, 51] and of neutron stars in compact binary systems [52–54]. For those parameters, a moderate to high initial black hole spin, $\chi_{\text{BH}}^i \geq (0.5-0.9)$, is a requirement for the neutron star to be disrupted by the black hole and thus allow the formation of an accretion disk [55]. But once that condition is satisfied, we showed that the local properties of the disk are largely independent of the binary parameters. Furthermore, 10 ms after merger the mass of the disk is typically $M_{\text{disk}} \sim (0.05-0.15)M_\odot$ and the neutrino luminosities $L_\nu \sim 10^{53}$ erg/s.

In this paper, we want to assess the effects of a better treatment of the neutrinos on the evolution of the disk and of its immediate neighborhood. To do so, we compare simulations using the same leakage scheme as in [31, 32] with simulations using the M1 formalism presented here. The leakage scheme should give us a rough estimate of the cooling of the disk due to the neutrinos and of the evolution of the composition of the high density regions of the disk, but does not include heating or composition evolution due to neutrino absorption in the low-density regions. The M1 formalism should give us a better estimate of both the total luminosity and the effects of neutrino-matter interactions.

We evolve the post-merger accretion disk with the following treatments of the neutrinos:

- A “Leakage” simulation using the same algorithm as in [31, 32].
- A “Leakage” simulation in which the neutrino opacities have been corrected as in our M1 code (to guarantee that the energy density of neutrinos in the optically thick regions is correct, see Appendix B).
- A “M1” simulation using the standard methods described in this paper.
- A “M1” simulation with a simpler treatment of the emissivity in optically thin regions (i.e. without the correction described in Eq. B6), to get a first estimate of the errors due to the use of a gray scheme: we find in our test of the neutrino-fluid interactions in a post-bounce supernova profile (see Appendix E 6) that variations in the results with and without that correction are comparable to the difference between the gray results and the results of an energy-dependent code.
- Three “M1” simulations with a numerical grid covering a larger volume and improved boundary conditions (see below), with and without the correction to the emissivities described in Eq. B6, and with and without the correction to the opacities described in Eq. B8.




FIG. 1. Density and velocity field in the equatorial plane of the black hole at the initial time $t_0 \sim t_{\text{merge}} + 6.1$ ms. For reference, the velocity at the peak of the density distribution is about $0.6c$.

As initial configuration, we consider a snapshot of simulation M14-7-S8 of [32] at $t_0 = t_{\text{merge}} + 6.1$ ms, where t_{merge} is the time at which half of the neutron star material has been accreted by the black hole. This is around the time when an accretion disk is forming and neutrino effects begin to significantly affect its evolution. The initial configuration is shown in Fig. 1. From the disruption of the neutron star to t_0 , the hydrodynamics equations were evolved using two levels of refinement, each with $100^2 \times 50$ grid points and with grid spacings $\Delta x \approx 1.5 \text{ km} \times 2^l$ in the equatorial plane and $\Delta z \approx 0.5 \text{ km} \times 2^l$ in the vertical direction (with $l = 0, 1$, and using the equatorial symmetry to only evolve the $z \geq 0$ region). We note that, due to our choice of map between the grid and the laboratory frame, the resolution in the grid is not uniform. We quote here the resolution close to the black hole, but the grid spacing far from the black hole is actually $\sim 30\%$ coarser. For comparison, the horizon of the black hole has a radius $r_{\text{BH}} \sim 25 \text{ km}$ in the coordinates of our simulation, the peak of the matter density in the accretion disk is initially at $r_{\text{disk}} \sim 50 \text{ km}$, and the scale height of the disk is $H \sim (0.2 - 0.3)r$. The finest grid level thus covers the forming accretion disk. Note that, in SpEC, Einstein’s equations are evolved on a separate pseudospectral grid extending 3700 km away from the center of the black hole. This is why we can use such a small finite difference grid if we only want to study the evolution of the post-merger accretion disk. A smaller grid means that we can maintain a reasonable resolution in the disk at a fairly low computational cost², but also that the unbound material, and some bound material on high-eccentricity orbits, is allowed to leave the grid. By t_0 , we still

² Although such a resolution is only reasonable because we do not include the effects of magnetic fields. It would be much too coarse to capture the effects of MHD instabilities.

have a baryonic mass $M_b = 0.12M_\odot$ on the grid, while we have allowed $0.06M_\odot$ of unbound material and $0.06M_\odot$ of bound material to escape. In [32], we also performed lower resolution simulations following the material in the tidal tail farther out. From these simulations, we can deduce that the material that we allowed to escape would start to fall back onto the disk at $t \sim t_{\text{merge}} + 20$ ms. At later times, material neglected in our simulation may affect the evolution of the disk.

In this work, we thus only evolve the system to $t_f = t_{\text{merge}} + 20$ ms. Another reason to do stop the simulation is that our simulations do not include the effects of magnetic fields, and the only viscosity is due to the finite resolution of our numerical grid. Over longer timescales, turbulent angular momentum transport due to the magnetorotational instability would significantly affect the evolution of the disk (see Sec. IV A). However, we will see that this is long enough for the fluid and the neutrinos to reach a quasi-equilibrium state. This also allows us to study the main differences between simulations using the leakage scheme and simulations using the M1 formalism. Finally, we can get a first estimate of the outflows emitted in the region above the disk and of the effects of interactions between the disk and the tidal tail.

For the evolution between t_0 and t_f , we use an extended finite difference grid. Indeed, although the system still respects the equatorial symmetry after merger, small perturbations which are not equatorially symmetric might grow in the disk due to hydrodynamical instabilities. To make sure that our grid structure does not artificially suppress such perturbations, we remove the assumption of equatorial symmetry. We also want to capture the radial growth of the disk, potential matter outflows, and the evolution of the bound material in the tidal tail remaining on the grid (which is still expanding at $t = t_0$). We thus add a third level of refinement (with the resolution chosen as before, with $l = 2$). Each refinement level now has 100^3 grid points. In three of the M1 simulations, a fourth level is added ($l = 3$), so that we can follow the evolution of low-density material far enough along the direction of the black hole spin axis to extract accurate information about potential disk winds.

We also performed simulations with different numerical resolution, to assess our numerical errors. A detailed discussion of these errors is provided in Appendix F. In summary, we find that the errors due to finite resolution are at most comparable to the errors due to the use of a gray scheme, and much smaller than the errors in the leakage scheme.

As opposed to previous SpEC simulations, we want here to study the behavior of low-density winds. Accordingly, for these last M1 simulations we also made a few modifications to the handling of low-density matter. In general relativistic hydrodynamics simulations, the region surrounding the neutron star or accretion disk is generally filled with lower-density material in order to avoid evolving towards either negative densities or unphysical values of the evolved variables. Additionally, corrections are applied to all regions of baryon density ρ_0 lower than an “atmosphere” threshold ρ_0^{atm} . In our case, those corrections set the temperature to $T = 0.5 \text{ MeV}$ and the spatial components of the 4-velocity to $u_i = 0$, although dif-

ferent prescriptions also work in practice. In previous SpEC simulations with the LS220 equation of state [31, 32], that threshold was set at $\rho_0^{\text{atm}} \sim 10^{6-7} \text{ g/cm}^3$ far away from the black hole. In this work, we are interested in disk outflows with density $\rho_0 \sim 10^{7-8} \text{ g/cm}^3$ close to the outer boundary of our grid. Such outflows cannot be launched with that high an atmosphere threshold. We thus modify our choice of atmosphere threshold to

$$\rho_0^{\text{atm}} = \rho_0^{\text{floor}} + \rho_0^{\text{max}} \left[10^{-5} \left(\frac{2r_{\text{AH}}}{r_{\text{AH}} + r} \right)^2 + 10^{-3} \exp \left(- \left(\frac{r - r_{\text{AH}}}{0.5r_{\text{AH}}} \right)^2 \right) \right], \quad (34)$$

with r_{AH} the grid-coordinate radius of the black hole apparent horizon, which is by construction a sphere in the coordinates of our numerical grid. r is the distance to the center of the black hole in the same coordinates, and $\rho_0^{\text{floor}} \sim 10^5 \text{ g/cm}^3$. With this choice, the atmosphere threshold remains as before close to the black hole, but now drops to $\rho_0^{\text{atm}} \sim \rho_0^{\text{floor}}$ at larger distances, which is sufficient for our current purpose.

Finally, in previous simulations the table containing the equation of state information only covered the range $10^8 \text{ g/cm}^3 < \rho_0 < 10^{16} \text{ g/cm}^3$, and was artificially extended to lower densities using a simple ideal gas equation of state. In this last simulation, we instead use a table going down to 10^5 g/cm^3 . The new table uses 345 logarithmically spaced points to cover that range of densities, instead of 250 for the smaller table. Both tables are also discretized in temperature ($0.01 \text{ MeV} < T < 251 \text{ MeV}$ with 136 logarithmically spaced points) and composition ($0.035 < Y_e < 0.53$ with 50 linearly spaced points). Below 10^8 g/cm^3 , the compositional information of the equation of state is approximated by the nuclear statistical equilibrium of matter at 10^8 g/cm^3 , for the same temperature and electron fraction (see O’Connor & Ott 2010 [49] for details).

IV. DISK EVOLUTION

In the following, we describe the physical results of the simulations. We mainly focus on the M1 simulation using our standard algorithm, including the improved treatment of the low-density regions for the hydrodynamical variables, as it offers the most accurate predictions. The effects of different neutrino treatments are discussed in Sec. IV G.

A. Global properties of the disk

At the beginning of the simulation ($t_0 = t_{\text{merge}} + 6.1 \text{ ms}$), we have about $0.08 M_\odot$ of material in a forming accretion disk extending $\sim 60 \text{ km}$ away from the black hole. As shown in Fig. 1, this material is still far from being circularized or axisymmetric. Another $0.04 M_\odot$ of material is on the grid in an extended tidal tail with a relatively flat density profile. As discussed in the previous section, an additional $0.06 M_\odot$ of

unbound material and $0.06 M_\odot$ of bound material with fallback time longer than $\sim 20 \text{ ms}$ were allowed to escape the grid before t_0 . The initial configuration has a sharp density peak around $r = 45 \text{ km}$. Most of the tail material is cold ($T < 1.5 \text{ MeV}$), while the disk material has a broad temperature distribution, with most of the matter at temperatures $2 \text{ MeV} < T < 12 \text{ MeV}$. The composition of the disk and tail is sharply peaked at $0.05 < Y_e < 0.07$, and the small amount of material at $Y_e > 0.1$ is in the hot regions close to the black hole and rapidly accreted.

Although neutrinos have an important effect on the evolution of the system, to first order, purely hydrodynamical effects dictate the evolution. During the time period considered in this simulation, $5 \text{ ms} - 20 \text{ ms}$ after merger, the circularization of the disk material is the most important effect. At the initial time, most of the disk material is at or close to periastron. Shock heating and the contraction of the disrupted material cause the fluid to heat to its maximum average temperature, $\langle T \rangle = 6.4 \text{ MeV}$ at $t = t_0 + 1.5 \text{ ms}$ ³. Afterwards, the forming disk goes through a damped cycle of expansions and contractions, with a period of about 6 ms . During each expansion period, the temperature of the fluid and the accretion rate decrease. During contractions, shock heating causes the temperature to rise, and the accretion rate increases. Neutrino emissions and absorptions, although not critical to the dynamics, contribute to a smoothing of the temperature distribution and determine the composition of the fluid. Their total luminosity mostly follows the oscillations of the fluid temperature. Energy lost to neutrino emissions and to the accretion of hot material onto the black hole also causes a slower global cooling of the fluid, by about 1 MeV over the 14 ms of evolution.

This is illustrated through snapshots of the simulation taken at $t_0, t_1 = t_0 + 5 \text{ ms}, t_2 = t_0 + 10 \text{ ms}$, and $t_f = t_0 + 14 \text{ ms}$. We plot the fraction of the local mass observed at a given radius in Fig. 2, at a given temperature in Fig. 3, and at a given electron fraction in Fig. 4. All plots are normalized to the total mass on the grid. Due to accretion onto the black hole, the total mass decreases from $0.12 M_\odot$ at t_0 to $0.097 M_\odot$ at t_1 , $0.075 M_\odot$ at t_2 , and $0.069 M_\odot$ at t_f . The first two snapshots t_0 and t_1 , are close to the times of maximum contraction and expansion of the disk, respectively. The third snapshot corresponds to the time at which the temperature becomes largely homogeneous and the electron fraction reaches its quasi-equilibrium value (i.e. nearly equal emission of electron neutrinos and antineutrinos). By the end of the simulation, at t_f , nearly all of the remaining material is in a circularized accretion disk.

Turbulent angular momentum transport due to the magnetorotational instability, which is not modeled here, should eventually take over for purely hydrodynamical effects and cause a viscous spreading of the disk. This should happen over the viscous timescale $t_{\text{visc}} \sim (\alpha \delta^2 \Omega)^{-1}$, where α is the standard viscosity parameter, $\delta = H/r$ and H is the scale height of the disk. For the expected $\alpha \sim 0.01 - 0.1$ and the simulation $\delta \sim 0.2 - 0.3$, we get $t_{\text{visc}} \sim (30 - 700) \text{ ms} >$

³ Here and in the rest of the text, the average fluid properties refer to density-weighted averages




FIG. 2. Radial distribution of the material outside the black hole. We plot the fraction of the total mass within bins of width $\Delta r \approx 3$ km at 4 representative times of the simulation. Most of the matter is initially near the black hole ($r \lesssim 60$ km) and close to periastron. The disk then expands, and later circularizes.

$t_f - t_0 = 14$ ms. We thus expect angular momentum transport due to magnetically-driven turbulence to be fairly unimportant on the timescales considered here. In a disk with $\alpha \sim 0.01 - 0.1$, we can also estimate the viscous accretion rate as

$$\dot{M} \approx 3\alpha\delta^2\Omega_{\text{disk}}M_{\text{disk}}. \quad (35)$$

For $\Omega_{\text{disk}} \approx 1900\text{s}^{-1}$ (at the peak of the density), we get $\dot{M}_{\odot} \approx (0.1 - 3) M_{\odot}/\text{s}$. For comparison, at the end of the simulation, the mass accretion rate has dropped to its lowest value of $\dot{M} \approx 1.5M_{\odot}/\text{s}$. Even at the end of the simulation, the accretion rate due to the circularization of the disk thus remains at least comparable with the expected accretion rate due to the magnetorotational instability. As for the heating of the disk, Newtonian simulations have shown that an isotropic viscosity can heat the disk to maximum temperatures of $T_{\text{max}} \sim 3 \text{ MeV} - 10 \text{ MeV}$ for $\alpha \sim 0.001 - 0.1$ [56]. For the largest viscosities, viscous heating would be relevant to the thermal evolution of the disk towards the end of our simulations. For lower viscosities, it would remain largely irrelevant until the disk cools down, but magnetically-driven turbulence might still hasten the homogenization of the disk temperature. As viscous heating is the main process stopping the cooling of the disk, it is of course always relevant to the long term evolution of the disk.

The observed temperature evolution (Fig. 3) is thus the result of mixing in the fluid, shock heating during the circularization of the disk, and energy transport by neutrinos. Some mixing of fluid elements with different temperatures does occur in purely hydrodynamical simulations, or in simulations using a simple cooling prescription for the neutrinos (leakage runs). But the process is significantly more efficient when using a transport scheme, as now neutrinos can transport energy from hot parts of the disks to cool parts of the disk. Variations in the average temperature of the fluid are due mostly to the oscillations observed as the disk circularizes.




FIG. 3. Fraction of the total mass within temperature bins of width $\Delta T = 0.5$ MeV, plotted at 4 representative times of the simulation. Within 10 ms, the temperature of the disk becomes very homogeneous.

The electron fraction (Fig. 4) reaches its equilibrium composition within about 10 ms. The composition then evolves more slowly as the properties of the disk change, adapting nearly instantaneously to the disk evolution. At the temperatures and electron fractions observed in most of the disk, matter becomes less neutron rich when the disk heats up, and more neutron rich as it cools down. The electron fraction also decreases for denser material. At times beyond 20 ms past merger, the disk is expected to cool and the core of the disk will reneutronize, at least until viscous heating balances neutrino cooling. However, this is not necessarily true for the high- Y_e tail of the distribution which corresponds to cooler, lower density points whose composition has been significantly affected by neutrino irradiation and which evolve fairly slowly at late times. The amount of high- Y_e material can be more easily assessed from Fig. 5, which shows that even after the average Y_e in the disk begins to decrease, the fraction of material with $Y_e \gtrsim 0.3$ actually increases. The production of that high- Y_e material is nearly entirely due to neutrino emissions and absorptions during the simulation: at the beginning of the simulation, 6.1 ms after merger, less than 3% of the material has $Y_e > 0.2$, and even that is mostly due to charged-current reactions before the beginning of the simulation, taken into account with the simpler leakage scheme.

B. Final disk configuration

From the discussion in Sec. IV A, we expect that at t_f the simulated hydrodynamical properties of the accretion disk (density, temperature, composition) are probably a good representation of the state of the system about 20 ms after merger. Magnetically-driven turbulence and magnetically-driven winds could affect the results. But so far simulations have found that for reasonable initial values of the magnetic fields, magnetic outflows only appear more than 100 ms after




FIG. 4. Fraction of the total mass within electron fraction bins of width $\Delta Y_e = 0.01$, plotted at 4 representative times of the simulation. Within 10 ms, the electron fraction of the disk becomes very homogeneous. The tail of higher- Y_e material is due to lower density regions around the disk.




FIG. 5. Fraction of the baryon mass with Y_e below a given value, at 4 representative times of the simulation. We note the slowly growing tail of high- Y_e material even after the average Y_e begins to decrease.

the merger [57], or are not observed at all [58, 59]⁴. As discussed in the previous section, MRI-driven angular momentum transport is also expected to occur on timescales longer than the duration of our simulation. In this section, we thus offer a more detailed description of the disk at the end of the simulation, assuming that it is a fairly good description of a post-merger accretion disk ~ 20 ms after a black hole-neutron star merger.

We first show the density and velocity field in the equatorial plane of the black hole (Fig. 6), and in the vertical plane $y = y_{\text{BH}}$ (Fig. 7). Here, we define the velocity as the “transport velocity” $v_T^i = \frac{\alpha}{W} u^i$, which satisfies $\partial_t \rho_* + \partial_i (\rho_* v_T^i) = 0$. This




FIG. 6. Density and velocity field in the equatorial plane of the black hole at the end of the simulation, 20 ms after merger. For scale, the velocities at the peak of the density distribution are $v \sim 0.5c$.

is a convenient definition of the velocity for visualization purposes, as it directly shows the motion of fluid elements on the grid with all general relativistic coordinate effects taken into account. From Fig. 6 we infer that the disk is well circularized up to $r \sim 100$ km. The velocity in the inner disk is mildly relativistic ($v_T \sim 0.5c$ at $r \sim 60$ km) which, as we shall see, has important effects on the geometry of the neutrino radiation.

Fig. 7 shows some interesting additional features. The core of the disk at the end of our simulation is of moderate geometrical thickness ($H/r \sim 0.2$), and still noticeably asymmetric (note that the density is plotted on a logarithmic scale). There are resolved outflows coming from the contact regions between the disk and the tail (left side of the plot), at densities $\rho_0 \sim 10^8$ g/cm³. At the grid boundary, this material has not reached the escape velocity (it has $u_t > -1$). But most of the outflowing fluid has enough energy to be unbound (i.e. it satisfies the condition $hu_t < -1$, thanks to temperatures $T \sim 2$ MeV). We discuss the disk/tail interactions in more detail in Sec. IV C, and the unbound material in Sec. IV D. For now, we simply note that the outflows, although helped by neutrino heating and radiation pressure, are mostly a consequence of purely hydrodynamical interactions at the disk/tail interface, as can be verified by their existence in the simulations using a leakage scheme. Earlier in the evolution, shocks during the circularization power similar outflows, albeit more axisymmetric and with some material closer to the black hole spin axis.

Similar visualizations for the temperature are shown in Fig. 8 and Fig. 9. Asymmetric features in the temperature distribution remain clearly visible despite the smoothing effect of the neutrino cooling and heating. Nearly all of the material, including the tidal tail, has been heated to $T > 1$ MeV, by a combination of shocks and neutrino absorption. Denser material with $\rho_0 \gtrsim 10^{11}$ g/cm³ is heated to $T \gtrsim 4$ MeV. The relatively small variations of the temperature are one of

⁴ The main difference between the two sets of simulations is the inclusion of an initial dipole magnetic field outside of the neutron star in [57].




FIG. 7. Density (in g/cm^3) and velocity field in the $y = y_{\text{BH}}$ plane at the end of the simulation, 20 ms after merger. For reference, the largest poloidal velocities are about $0.6c$. In the core of the disk, the velocities are mostly directed out of the plane of the visualization.




FIG. 9. Temperature in a plane orthogonal to the equatorial plane of the disk at the end of the simulation, 20 ms after merger. Density contours are shown as thick black lines for $\rho_0 = (10^9, 10^{10}, 10^{11}) \text{g}/\text{cm}^3$.




FIG. 8. Temperature in the equatorial plane of the black hole at the end of the simulation, 20 ms after merger. Density contours are shown as thick black lines for $\rho_0 = (10^9, 10^{10}, 10^{11}) \text{g}/\text{cm}^3$.




FIG. 10. Electron fraction Y_e in the equatorial plane of the black hole at the end of the simulation, 20 ms after merger. Density contours are shown as thick black lines for $\rho_0 = (10^9, 10^{10}, 10^{11}) \text{g}/\text{cm}^3$.

the main reasons a gray scheme for the neutrino radiation is more reliable for merger simulations than in core collapse supernovae, where large temperature gradients exist and small changes in the interactions between the neutrinos and the fluid can significantly modify the dynamics of the system.

Finally, we consider the composition of the fluid, through its electron fraction Y_e , in Fig. 10 and Fig. 11. Being able to reliably predict the electron fraction in low-density regions is one of the main advantage of the use of the moment formalism over the leakage scheme. We see that the core of the disk remains relatively neutron rich ($0.15 \lesssim Y_e \lesssim 0.20$), but material below $\rho_0 \sim 10^{10} \text{g}/\text{cm}^3$ is significantly protonized. About 30% of the total mass is at $Y_e > 0.2$, and about 10% at $Y_e > 0.3$ (see also Fig. 5). More importantly, the less neutron-rich material, which surrounds the neutron-rich core of the disk, is more likely to be unbound by disk winds. Lee *et*

al. [60], Fernandez & Metzger [61, 62] and Just *et al* [63] have shown that 5% – 25% of the matter in accretion disks formed in binary neutron star or black hole-neutron star mergers is eventually unbound, mostly due to viscous heating in the disk. In those 2D simulations, most of the ejecta remains neutron-rich. Its nucleosynthesis output is very close to the output of the more massive dynamical ejecta [61], and the associated radioactively powered electromagnetic transients would also be very similar. However, the initial conditions for these simulations were taken to be $Y_e \sim 0.1$ everywhere, at a time at which the disk is already wider and the composition evolution due to neutrino emission slower than in the simulations presented here. It is possible that a higher initial electron fraction in the outer regions of the disk can have a measurable impact on the properties of kilonovae. We discuss this in more detail in Sec. IV D.




FIG. 11. Electron fraction Y_e in a plane orthogonal to the equatorial plane of the disk at the end of the simulation, 20 ms after merger. Density contours are shown as thick black lines for $\rho_0 = (10^9, 10^{10}, 10^{11}) \text{ g/cm}^3$.

C. Stability and disk/tail interactions

From the beginning of the simulation, strong asymmetries and shocks are present in the disk, and the system is well out of equilibrium. However, as the disk circularizes, it reaches a more stable configuration. By the end of the simulation, although asymmetries remain, fluid elements follow nearly circular trajectories at the expected orbital frequency, up to radii $r \sim 100 \text{ km}$. The core of the disk is convectively stable, and close to equilibrium. Around the disk/tail interface, however, this is not the case. First, the tail material is not circularized, which creates a shear layer at the outer edge of the disk. Additionally, according to the Soldberg-Hoiland criterion [64], the disk is convectively unstable in both the vertical and radial directions. We should note that as this is also the region in which the disk begins to deviate from hydrostatic equilibrium, the Soldberg-Hoiland criterion is not strictly applicable. But the presence of convective regions is clearly observable at all times in the simulation. These instabilities cause the creation of large scale eddies close to the outer edge of the disk, and are strongly correlated with outflows launched above the disk. Given how out of equilibrium the system is, picking a single instability responsible for the creation of the eddies or causally associating these eddies with the outflows is difficult. However, we note that regions in which there no longer are any interactions between the accretion disk and the tidal tail are devoid of both eddies and outflows. We can infer that the ejection of disk material is probably helped by the convection of hotter fluid from the core of the disk to the top of the disk/tail interface. The high neutrino fluxes in that same region probably play a role as well, especially in directing the outflows at late times and setting their composition. Comparing Fig. 7 and Fig. 13, it is clear that the disk outflows are, at the end of the simulation, aligned with the neutrino radiation. This is not the case at earlier times, when unbound material is observed closer to the spin axis of the black hole.

D. Unbound material

The ejection of unbound material by black hole-neutron star and neutron star-neutron star mergers is, from an astrophysical perspective, one of the most important consequences of these mergers. This is because these ejecta are one of the most likely locations for r-process nucleosynthesis to occur [11–13, 65]. Neutron star mergers were long thought to happen too late in the evolution of a galaxy to explain observations of r-process elements [66], but recent studies incorporating updated population synthesis predictions for neutron star binaries are more favorable to the merger process [67–69]. Additionally, the radioactive decay of nuclei in the neutron-rich ejecta during r-process nucleosynthesis powers electromagnetic transients potentially detectable in the optical [70] or, more likely, in the infrared [16, 71, 72]. The luminosity, duration and peak frequency of these electromagnetic signals can significantly vary with the mass, composition, entropy and velocity of the ejecta [17, 71, 73]. In particular, the composition will significantly affect the results of the nucleosynthesis: low Y_e material will produce mostly heavy elements (strong r-process), while high Y_e material will produce more iron-peak elements (weak r-process). The transition occurs at $Y_e \sim 0.25 - 0.3$ for conditions typical of a viscously-driven wind in a post-merger accretion disk [74]. However, nucleosynthesis results also depend on the entropy of the ejected material, and the exact dividing point for the lower entropy ejecta observed in our simulations is not, to our knowledge, known at this point. This difference in the products of r-process nucleosynthesis impacts the lightcurve of radioactively powered electromagnetic transients. Indeed, high-opacity lanthanides are produced in the case of a strong r-process, causing the emission to be fainter, redder and longer lived than in the case of a weak r-process [16, 71, 72].

In black hole-neutron star binaries, matter can be ejected through different processes during and after the merger. First, if the neutron star is disrupted, a mass $M_{\text{ej}} \sim 0.01M_{\odot} - 0.1M_{\odot}$ (depending on the parameters of the binary) is typically unbound during the tidal disruption of the neutron star [32, 48, 75]. The amount of ejected material can even be larger for rapidly spinning, low mass black holes [31, 76]. The tidally ejected material is very neutron rich ($Y_e < 0.1$), cold ($T < 1 \text{ MeV}$), confined close to the equatorial plane, and strongly asymmetric.

After that, material can be ejected during the formation of the accretion disk and at the interface between the disk and the tidal tail. This is the main source of outflows observed in this simulation, and is discussed in more detail below.

Third, disk winds may be triggered by magnetic effects and/or neutrino absorption. Newtonian simulations of binary neutron star mergers using an energy-dependent leakage scheme (with a specifically designed absorption term for neutrinos in optically thin regions) [77] or energy-dependent flux-limited diffusion [78] observed the formation of a neutrino-driven wind within $\sim 100 \text{ ms}$ of the merger, and neutrino driven winds were also observed in two-dimensional simulations of an accretion disk, with initial condition taken from a black hole-neutron star merger [63]. In [77], the observed

outflows were more neutron-rich in the equatorial region, with conditions favorable to a strong r-process, and less neutron-rich in the polar region, with the expectation of a weaker r-process (and bluer kilonovae). In [63], a wide range of electron fraction is observed in all directions, although the polar outflows remain less neutron rich. The disk generated in our black hole-neutron star mergers are likely, over longer timescales than those simulated here, to create winds similar to those observed in [63, 77]. In those studies, about 1% of the disk mass was eventually ejected in the neutrino-driven wind. In black hole-neutron star mergers, this is generally much less material than in the dynamical ejecta: for the configuration considered here, the total mass ejected in the neutrino-driven wind would be $M_{\text{wind}} \sim 10^{-3} M_{\odot}$, while the dynamical ejecta has mass $M_{\text{din}} \sim 0.06 M_{\odot}$. Accordingly, that ejecta is only interesting if its different composition has observable consequences.

Finally, over longer timescales (seconds), and using an idealized initial disk profile, 2D Newtonian simulations have also shown that viscous heating can drive strong outflows in the disk [61, 79]. The total ejected mass is about 5% – 25% of the mass of the disk, with some significant dependence on the spin of the black hole [62]. Starting from idealized initial conditions (equilibrium torus with $Y_e \sim 0.1$), these simulations find that most of the unbound material remains at electron fractions too low to avoid the production of lanthanides – and they thus lead to an electromagnetic signal peaking in the infrared, and to strong r-process nucleosynthesis. For an initial black hole spin similar to our simulation ($\chi_{\text{BH}} = 0.8$), however, 0.1% – 1% of the mass of the disk is ejected in a less neutron-rich outflow, which does not produce any lanthanides and could lead to a ‘blue bump’ in the kilonova lightcurve [62]. More recent results using more accurate neutrino methods also find large ejected masses, of the order of 20% – 25% of the mass of the disk [63]. For the configuration considered here, this would be a mass $M_{\text{vis}} \sim 0.02 M_{\odot}$.

In our simulation, we observe polar outflows during the circularization of the disk, mostly coming from the region in which the accretion disk and the tidal tail interact. The total mass ejected is only $3 \times 10^{-4} M_{\odot} \sim 0.4\% M_{\text{disk}}$, a negligible amount compared to the mass ejected during the disruption of the neutron star ($0.06 M_{\odot}$). But because it is ejected early, and in a direction in which no material has been unbound so far, its effects cannot be neglected. First, material in the polar region can affect the formation and collimation of a jet, if the merger leads to a short gamma-ray burst. Additionally, because this material is ejected before any disk wind, it could obscure the blue component of a kilonova if enough high-opacity lanthanides are formed during r-process nucleosynthesis and the ejected material obscures a significant fraction of the polar regions. The properties of the outflows observed in our simulation vary significantly in time, and are summarized in Table I. The front of the outflow has a relatively low electron fraction ($Y_e \lesssim 0.2$) and specific entropy $s \sim 30 k_B$ per baryon. At later times, the outflows become less neutron rich and colder. By the end of the simulation, we measure $Y_e \gtrsim 0.3$ and $s \sim (10 - 15) k_B$. Overall, about 15% of the ejected material has $Y_e \lesssim 0.2$, and about 15% has

TABLE I. Properties of the post-merger outflows, as measured on the outer boundary of the computational domain. \dot{M} is the mass outflow rate, $\langle Y_e \rangle$ is the average electron fraction, $\langle s \rangle$ is the average specific entropy per baryon, and $\langle \Theta \rangle$ is the average angle between the direction of propagation of the outflowing fluid and the spin axis of the black hole. We list the average properties of the outflow over the last 11ms of evolution (there are not significant outflows at earlier times), as well as average properties of the unbound material crossing the outer boundary of the computational domain at given times. t_{peak} is the time at which the outflow rate is maximal. All values are for our ‘‘standard’’ M1 algorithm. Data required to compute the time-averaged $\langle \Theta \rangle$ was not output by the simulation, so that we only compute $\langle \Theta \rangle$ from snapshots at fixed times.

Time	$\dot{M} (M_{\odot}/s)$	$\langle Y_e \rangle$	$\langle s \rangle$	$\langle \Theta \rangle$
Average	0.032	0.25	$15 k_B$	NA
$t_{\text{peak}} = t_0 + 4 \text{ ms}$	0.131	0.21	$22 k_B$	50°
$t_1 = t_0 + 5 \text{ ms}$	0.087	0.27	$11 k_B$	41°
$t_2 = t_0 + 10 \text{ ms}$	0.005	0.36	$8 k_B$	42°
$t_f = t_0 + 14 \text{ ms}$	0.004	0.42	$11 k_B$	49°

$Y_e \gtrsim 0.3$. Accordingly, we would expect most of the material unbound during disk formation to undergo strong r-process nucleosynthesis, produce lanthanides, and obscure potential optical components of the radioactively powered electromagnetic transient. By the end of the simulation, the mass loss to these outflows has stabilized to about $0.005 M_{\odot}/s$. It is thus conceivable that after the end of our simulation, another $\sim 10^{-4} M_{\odot}$ of less neutron-rich material would be unbound through the same process. The average orientation of the outflows remains around $\langle \Theta \rangle \sim 40^\circ - 50^\circ$, as a large fraction of the material still has a significant velocity orthogonal to the spin axis of the black hole. But while at late times the region along the spin axis of the black hole is largely devoid of matter, at early times some ejecta is observed in that region.

As we only considered a single configuration, and the mass ejected along the polar axis is barely enough to have an effect on the light curve of the radioactively powered transient, it would be dangerous to draw generic conclusions from these results. The properties and total mass of the outflows are indeed likely to depend on the parameters of the binary. But our simulations demonstrate that the effect of polar outflows ejected during the circularization, despite their small mass, should probably be taken into account when modeling those signals. Neutron-rich polar outflows ejected during disk circularization could have an effect on the collimation of relativistic jets, and could obscure the electromagnetic signal from more proton-rich neutrino-driven winds ejected at later times.

E. Neutrino Emission

With our M1 code, we can for the first time examine the spatial distribution of neutrinos in the first 20 ms following a black hole-neutron star merger. To understand the main properties of the neutrino radiation, it is however useful to take a step back and look at a few of the quantities which could already be predicted using our leakage scheme. Indeed, the leakage scheme gives us a good estimate of the optical depth




FIG. 12. Neutrinospheres $\tau = 2/3$ at the end of the simulation, 20 ms after merger. We show the vertical plane $y = y_{\text{BH}}$. The three contours are, from the outside to the inside, the neutrinospheres for ν_e (outer solid line), $\bar{\nu}_e$ (dashed line) and ν_x (inner solid line). The disk is colored according to its baryon density ρ_0 .

in the disk, a useful quantity to understand where the neutrinos effectively decouple from the fluid. For the electron neutrinos, although there are always a few hotter regions in which the optical depth is $\tau_{\nu_e} \sim 10$, the density averaged optical depth is only $\langle \tau_{\nu_e} \rangle \sim 2$. It is even lower for the electron antineutrinos, with $\langle \tau_{\bar{\nu}_e} \rangle \sim 1$, and heavy lepton neutrinos, with $\langle \tau_{\nu_x} \rangle \sim 0.5$. The neutrinospheres, defined as $\tau = 2/3$, are shown for a vertical cut at the end of the simulation in Fig. 12. We see that neutrinos are only trapped in the core of the disk, and mostly free streaming everywhere else.

To illustrate the main properties of the neutrino radiation, we plot in Figs. 13-14 the energy density and “normalized flux” $F_{\text{norm}}^i = \alpha F^i / E - \beta^i$ in vertical and horizontal slices of the disk, for the electron neutrinos. The normalized flux is chosen so that it represents an effective transport velocity for the neutrino energy. In the core of the disk, Fig. 14 shows that the neutrino energy is mostly transported with the fluid, as befits an optically thick region. Outside of the expected neutrinosphere, i.e. for $r \gtrsim 90$ km, the neutrinos transition to free streaming away from the disk. The energy density is maximal close to the inner edge of the disk, in part due to higher temperatures and in part due to gravitational redshifting.

The vertical slice on Fig. 13 shows a few additional features of interest. First, we note the random orientation of the fluxes along the speed axis of the black hole, up to a height $|z| \lesssim 80 M_\odot$. This is a known problematic feature of the M1 approximation, due to the convergence of neutrinos from all around the disk. Beyond this issue, we can also see that the emission at large radii is clearly asymmetric. We find significantly lower energy densities within about 20° of the equatorial plane as well as in the (less reliable) polar regions. This is a purely geometric effect due to the projected shadow of the disk and the relativistic beaming of the neutrinos.

To understand this beaming, we first have to remember that the velocities in the disk are mildly relativistic, with $v \sim 0.5c$. This means that the emission of neutrinos will be focused within a relatively large beam centered on the direction of motion of the fluid, which nearly follows a circular orbit around




FIG. 13. Energy density and normalized flux $(\alpha F^i / E - \beta^i)$ of the electron neutrinos at the end of the simulation, 20 ms after merger. We show the vertical plane $y = y_{\text{BH}}$.




FIG. 14. Energy density and normalized flux $(\alpha F^i / E - \beta^i)$ of the electron neutrinos at the end of the simulation (20 ms after merger), in the equatorial plane of the disk.

the black hole. Only a small fraction of the neutrinos are emitted in the vertical direction. Additionally, most of the neutrinos decouple from the matter around the location of the neutrinosphere, and then free stream away from the disk. Accordingly, if the inner regions of the disk are brighter than its outer regions, as is the case here, the disk casts a shadow along the equatorial plane. This is more clearly illustrated by Fig. 15, which shows the energy density and fluxes for the electron neutrinos on the surface $\tau_{\nu_e} = 0.1$. Most of the energy comes from the inner disk, and neutrinos cannot easily escape along the equatorial plane. The shadow along the equatorial plane observed in Fig. 13 exactly matches the thickness of the disk. Relativistic beaming causes neutrinos to be preferentially emitted nearly tangent to the disk. Hence the neutrino flux is larger just outside of the shadow than at high latitudes. The equatorial shadow is of course not perfect, as some neutrinos are emitted from the outer edge of the neutrinosphere. But this is a relatively small fraction of the total neutrino lu-




FIG. 15. Energy density and normalized fluxes of the electron neutrinos on the $\tau_{\nu_e} = 0.1$ surface.

minosity, as can be seen in Fig. 15.

The radiation fields of the other neutrino species are qualitatively similar, except for the fact that the neutrinosphere is located deeper inside the disk. As shown in Fig. 16, we find that most of the energy emitted in neutrinos goes into electron antineutrinos, at least at early times. The luminosity in ν_e is fairly constant during the evolution, at $L_{\nu_e} \sim 5 - 8 \times 10^{52}$ ergs/s. But the luminosity in $\bar{\nu}_e$ peaks at $L_{\bar{\nu}_e} \sim 5 \times 10^{53}$ ergs/s during the rapid evolution of the electron fraction from its very neutron-rich initial value to the equilibrium value in the disk. Within 5 ms, it decreases to $L_{\bar{\nu}_e} \sim 10^{53}$ ergs/s, and by the end of the simulation it becomes lower than the electron neutrino luminosity, with $L_{\nu_e}^f = 7 \times 10^{52}$ ergs/s and $L_{\bar{\nu}_e}^f = 5 \times 10^{52}$ ergs/s. Finally, there is an equally brief burst of heavy-lepton neutrinos, with $L_{\nu_x} \sim 3 \times 10^{52}$ ergs/s for each species, due to the existence of hot spots early in the disk formation. That luminosity rapidly decreases as the temperature becomes more homogeneous. Within 3 ms, we measure $L_{\nu_x} \sim 5 \times 10^{51}$ ergs/s for each species, and by the end of the simulation, $L_{\nu_x}^f = 10^{51}$ ergs/s per species.

The properties of the neutrino spectrum are not directly measurable within our gray formalism. Nonetheless, it is possible to get reasonable estimates from either the predictions of the leakage scheme, shown on Fig. 17, or the properties of the fluid on the neutrinosphere. The leakage scheme predicts average energies of $\langle \epsilon_{\nu_e} \rangle = 11$ MeV – 13 MeV, $\langle \epsilon_{\bar{\nu}_e} \rangle = 13$ MeV – 15 MeV, and $\langle \epsilon_{\nu_x} \rangle = 14$ MeV – 19 MeV during the last 10 ms of evolution, and higher energies at the beginning of the simulation, when hot spots are present. Neglecting corrections due to the finite chemical potential of neutrinos in the emitting regions and assuming a redshift factor of 2 between the emitting region and the observer, this would correspond to fluid temperatures of roughly 6 MeV, 7 MeV, and 8 MeV. Considering the temperature distribution observed in the disk, the increasing temperature as the neutrinosphere recedes deeper into the disk, and the higher emissivity of high temperature points, this appears fairly reasonable.

F. Neutrino viscosity and the growth of magnetic instabilities

The magnetorotational instability (MRI) is expected to play a crucial role in the growth of magnetic fields in post-merger accretion disk, and may also be important for the generation of jets after a compact binary merger. However, recent work has shown that the transport of momentum by neutrinos can significantly affect the growth timescale and wavelength of the MRI [80, 81]. In the context of protoneutron stars, Guilet *et al.* [81] showed that if the wavelength of the fastest growing mode of the MRI in the absence of neutrinos λ_{MRI} is larger than the neutrino mean free path λ_ν , an effective viscosity from the transport of neutrinos causes the MRI to grow slower than expected for small magnetic fields, but at a fastest-growing wavelength independent of the magnetic field strength (instead of the wavelength decreasing with the magnetic field strength). If instead $\lambda_{\text{MRI}} < \lambda_\nu$, the neutrinos can act as a drag force on the magnetized fluid. For large neutrino energy densities, this slows the growth timescale of the fastest growing mode of the MRI, and slightly increases its wavelength. In accretion disks, and when neutrinos can be modeled through an effective viscosity, a similar increase of the growth timescale of the MRI has been measured [80]. By applying the model of Guilet *et al.* [81] to our accretion disk, we can obtain a first estimate of the expected effect of neutrinos on the growth of the MRI.

First, we need to determine the critical magnetic field B_c at which $\lambda_{\text{MRI}} = \lambda_\nu$, using $\lambda_{\text{MRI}} \sim 2\pi b / \sqrt{b^2 + \rho_0 h}$ and $\lambda_\nu \sim 1 / (\bar{\kappa}_a + \bar{\kappa}_s)$ (where b is the strength of the magnetic field observed by an observer comoving with the fluid). At the highest density points in the disk, we get $B_c \sim 10^{13}$ G, while B_c smoothly increases with decreasing density, to $B_c \sim 10^{15}$ G for $\rho_0 \sim 10^{10}$ g/cm³. These values are larger than the initial magnetic field in most merging neutron stars, but smaller than the expected saturation amplitude of the MRI, thus indicating that both the viscous and neutrino drag regimes might be relevant during the evolution of a post-merger accretion disk.

As far as the neutrino drag is concerned (e.g. for $B < B_c$), its effects on the growth of the MRI should however be negligible. Guilet *et al.* [81] find that the importance of neutrinos on the growth of the MRI in this regime is determined by the value of the dimensionless parameter

$$\frac{\Gamma}{\Omega} = \frac{2(\bar{\kappa}_a + \bar{\kappa}_s)E_\nu}{15\rho_0\Omega}. \quad (36)$$

For $\Gamma/\Omega \gtrsim 1$, neutrinos affect the growth of the MRI. However, in our disk, we find $\Gamma/\Omega \lesssim 0.01$ everywhere. Thus the growth of the MRI should remain unaffected at low magnetic field strengths.

In the viscous regime ($B > B_c$), the importance of neutrino effects is determined by the value of the Elasser number $\epsilon = v_A^2 / (\nu\Omega)$ where v_A is the Alfvén speed and

$$\nu = \frac{2E_\nu}{15\rho_0(\bar{\kappa}_a + \bar{\kappa}_s)} \quad (37)$$

is the effective viscosity due to neutrinos. Viscosity affects the growth of the MRI for $E_\nu \lesssim 1$. For $B \sim B_c$, and the conditions observed in our simulation, we would get $\epsilon \sim 1$ (and the

Elasser number then grows as B^2). Accordingly, within the simple model used here, neutrinos could plausibly affect the growth of the MRI for $B \sim B_c$. But for $\epsilon \sim 1$, these effects are mild, and neutrinos are not expected to have any effect for $B \ll B_c$ or $B \gg B_c$.

G. Impact of the neutrino treatment

Having performed simulations with both a leakage scheme and the M1 formalism, we can now obtain better estimates of the impact that an approximate treatment of the neutrinos has on the evolution of a post-merger accretion disk.

1. Comparison with a leakage scheme

We can first look at differences between the leakage simulations and the M1 simulations. Not surprisingly, the two methods provide reasonable qualitative agreement for the global properties of the high density regions of the disk: the simulations with neutrino leakage capture the formation of the disk, its early protonization and later re-neutronization, and the global temperature evolution due to the initial expansion of the disk, neutrino cooling and shock heating. Lacking neutrino absorption, the leakage simulation however tends to maintain larger temperature differences between neighboring regions of the disk. It underestimates the timescale necessary for the disk to cool down, and the magnitude of its protonization. More quantitatively, the average electron fraction in the leakage simulation is lower by $\Delta Y_e \sim 0.05$, and the average temperature is lower by $\Delta T = (0.5-0.8)$ MeV. Additionally, without neutrino absorption the evolution of the fluid composition in low-density regions is entirely unreliable. The composition of the tidal tail material falling back onto the black hole and of the outflows are thus radically different.

There are also significant differences in the neutrino luminosities, for the electron neutrinos by a factor of two about 10 ms after merger (5 ms after the beginning of the simulation), and by about 30% by the end of the simulation. For heavy-type neutrinos, the difference is typically a factor of 2 – 4, presumably because of the steeper dependence of their emissivities on the temperature of the disk. We show the luminosities for the leakage scheme, the M1 scheme, and the predictions of the leakage scheme for the value of the hydrodynamic variables obtained when evolving the system with the M1 scheme on Fig. 16. The first thing to note is that, since the leakage scheme measures the instantaneous energy loss at each point of the domain while the luminosities in the M1 scheme are measured through the neutrino fluxes at the outer boundary of the computational domain, there is naturally a time shift between the two predictions. Even taking that time shift into account, however, it is quite clear that the electron neutrino luminosity is larger in the leakage scheme, while the heavy-lepton neutrino luminosity is larger in the M1 scheme. The electron antineutrino luminosities cannot be distinguished within the uncertainties due to the time shift in the measurements. Looking at the predictions of the leakage




FIG. 16. Luminosities for the different neutrino species as a function of time. Solid curves show the M1 results. Dashed curves show results using a leakage scheme. Thin dot-dashed curves show the prediction of the leakage scheme, but for the value of the hydrodynamic variables obtained by evolving the disk with the M1 scheme. Heavy-lepton neutrino luminosities ν_x are for all 4 species combined.

scheme within the M1 simulation allows us to differentiate between the error due to the instantaneous estimate of the luminosity, and the error due to the diverging evolution of the hydrodynamics variables. We see that by the end of the simulation, both sources of error are important. The impact of the diverging evolutions of the hydrodynamics variables can also be seen in Fig. 17, which shows the average energy of the neutrinos for the M1 and leakage simulations (computed through the leakage scheme in both cases, as we do not have any information about the neutrino spectrum in the M1 code). We observe differences of $\sim (1-2)$ MeV in the predicted average energy of the neutrinos at late times, for all species. This is comparable to the expected error in the determination of these energies through the leakage scheme.

Overall, these results appear consistent with the expected limitations of a leakage scheme. We also performed leakage simulations with two different methods to compute the opacities, to check whether the differences observed between the leakage and M1 schemes could be due to differences in the estimated neutrino opacities. But the differences between these two simulations were well below the estimated errors in the leakage scheme.

2. Impact of the gray approximation

By looking at M1 simulations using different energy-averaging approximations, we can also attempt to estimate errors related to the use of a gray scheme. We use two different gray approximations which, in our test of neutrino-matter interactions in a post-bounce supernova profile, bracketed the solution obtained with an energy-dependent code (see Appendix E). Clearly, as an error estimate, this is a poor substitute for a comparison with an energy-dependent radiation




FIG. 17. Average neutrino energy for the different neutrino species as a function of time. Solid curves show the results from the M1 simulation. Dashed curves show results using a leakage scheme. Heavy-lepton neutrino energies (ν_x) are for all 4 species combined. Note that for both simulations, the energy is computed using the number and energy source terms from the leakage scheme, as the gray M1 scheme does not contain information about the neutrino spectrum

transport code. Unfortunately, such a simulation remains too costly to perform with the SpEC code at this point. We find that differences in the average properties of the disk, although measurable, are much smaller than the differences between the M1 simulations and the leakage simulations. The average electron fraction varies by $\Delta Y_e < 0.01$, and the average temperature by $\Delta T < 0.1$ MeV. The total neutrino luminosities agree, for all species, within $\sim 20\%$.

We also performed a simulation in which the gray opacities were computed assuming that the neutrinos obey a Fermi-Dirac distribution with temperature T equal to the fluid temperature and equilibrium chemical potential μ_ν , a method which significantly underestimates neutrino opacities: low density regions can have $T \sim 1 - 2$ MeV while neutrinos are mostly emitted in regions with $T \gtrsim 5$ MeV. The resulting differences are significantly smaller than in the test of the neutrino-matter coupling presented at the end of the appendices. In fact, in the core of the disk, the errors are similar to the differences between the two M1 simulations using different prescription for the emissivities. Where the two methods differ, unsurprisingly, is in the composition of the outflows: when assuming an equilibrium distribution at the fluid temperature for the neutrinos, the electron fraction of the outflows is consistently $\Delta Y_e = 0.05$ lower than when using the more realistic neutrino energies computed from the temperature in the emitting regions. This confirms our assumption that the electron fraction in the outflows is significantly affected by neutrino absorption.

3. Limitations of the M1 closure

Finally, when analyzing our results, it is worth noting once more the limitations of the M1 closure. Crossing beams, caustics, and strongly focused beams are known to be problematic when using the M1 closure. Some examples of these issues can be found in the tests presented in Appendix E. Accordingly, the results of our simulations cannot be trusted close to the spin axis of the black hole, where neutrinos from all regions of the disk cross. Because of the relativistic beaming of the neutrinos, the energy density in that region is already low, but unphysical radiation shocks cause an even larger decrease. Unfortunately, the neutrino radiation in the polar region can have important consequences on the post-merger evolution of the system. Indeed, neutrino-antineutrino annihilations into electron-positron pairs deposit energy in this low-density region, and affect the matter density there. The formation of a relativistic jet, desirable if black hole-neutron star mergers are to produce short gamma-ray bursts, is quite sensitive to baryon-loading in the polar regions, but on the other hand could be helped by the energy deposition. The exact impact of the neutrinos in that context remains an open question, which can only be answered with an improved treatment of the neutrinos, and a better understanding of the jet forming mechanism.

V. CONCLUSIONS

We present a new module of the SpEC code [18], allowing us to study the effects of neutrinos within a fully general relativistic - hydrodynamics code. The neutrinos are modeled using the M1 formalism, in which the first two moments of the neutrino distribution function are evolved (energy and fluxes). Although the formalism can in theory be energy-dependent, we limit ourselves to an energy-integrated version of the code, due to the high cost of energy-dependent simulations. We offer here a detailed description of the implementation of the M1 algorithm in SpEC, as well as a series of tests assessing our ability to study the evolution of neutrinos in flat and curved spacetimes, and their interaction with matter.

We also discuss the first simulation with both radiation transport and a general relativistic code of the evolution of an accretion disk produced by a typical black hole-neutron star merger. We use as initial conditions a snapshot of an existing SpEC simulation using a simpler treatment of the neutrinos (leakage scheme) [32], right as the accretion disk begins to form and neutrino effects become important. This provides us with realistic initial conditions for the accretion disk, at least compared to the more commonly used equilibrium tori. As we neglect magnetic fields and all material falling back on the disk on timescales longer than 20 ms, we limit ourselves to a relatively short evolution, up to 20 ms after merger. We find that the evolution of the forming accretion disk is initially dominated by the circularization of the disk material. The disk expands and contracts in a cycle of about 6 ms. These oscillations are the main driver of the evolution of the average disk properties at early times. As the disk circularizes, however,

these oscillations are rapidly damped.

Neutrinos cause a global cooling of the disk, but on a relatively long timescale of order 50 ms. On the timescale of this simulation, neutrinos have however other important effects. First, they drive the composition evolution of the core of the disk. The disk rapidly protonizes, reaching an average electron fraction $Y_e \sim 0.2$ (a larger value than predicted by the leakage scheme [32]). The electron fraction then more slowly decreases as the disk starts to cool down. Second, neutrinos cause changes in the electron fraction of the outer parts of the disk, and of the disk outflows. Although initially neutron rich, by the end of the simulations these regions have $Y_e > 0.3$ ($Y_e \sim 0.4$ in the outflows). These low density regions above the disk would be the first to be unbound by disk winds and viscously-driven outflows [61–63, 79], and their higher initial electron fraction could thus affect the composition of late time outflows, their nucleosynthesis output, and the light curve of associated electromagnetic transients. Third, energy transport by the neutrinos homogenize the temperature distribution in the disk, with cold regions absorbing higher energy neutrinos and hot regions radiating much more strongly. Finally, neutrinos will deposit energy and create electron-positron pairs in the low-density polar regions. This could plausibly affect, positively or negatively, the formation of relativistic jets in that region. Unfortunately, neutrino-antineutrino annihilation in low density regions is not modeled within our formalism. As opposed to what was observed in simulations of protoneutron stars [81], we also estimate that the impact of neutrino transport on the growth of the magnetorotational instability is likely to be small to inexistant in post-merger accretion disks.

We also find that, because the disk is very compact, and forms around a rapidly rotating black hole ($\chi_{\text{BH}} = 0.87$), relativistic effects are significant. Most of the radiation comes from a region with mildly relativistic velocities ($v \sim 0.5c$). Accordingly, the neutrinos are beamed tangentially to the disk, causing strong anisotropies in the neutrino luminosity. Light bending and gravitational redshift also naturally affect the luminosity and neutrino spectrum. The luminosity is low in the equatorial plane, due to the disk shadow, and in the polar regions, due to beaming (although the evolution of polar regions with the M1 closure is unreliable). We note that even though a black hole spin $\chi_{\text{BH}} = 0.87$ might seem large, this should be fairly typical of a black hole-neutron star merger in which an accretion disk forms. Indeed, for the most likely black hole masses, slowly spinning black holes cannot disrupt the neutron star before it plunges into the black hole [55].

A number of simulations using approximate treatments of gravity have also considered the impact of neutrinos on disk evolutions, with methods generally more advanced than contemporary general relativistic simulations (e.g. [25, 56, 63, 77, 82]), and/or capable to evolve the system over longer timescales (e.g. [60–62]). However, direct comparisons with our results are difficult. This is in part because of the importance of relativistic effects for accretion disks around rapidly spinning black holes, and also because simulations which do not include general relativistic effects typically start from either an idealized equilibrium torus, or from the result of a merger in which the neutron star was disrupted by tidal ef-

fects modeled by a pseudo-Newtonian potential, whose realism close to a rapidly spinning black hole is difficult to assess. Our simulation shows that general relativistic effects significantly impact the neutrino radiation and the disk formation, and the forming accretion disk is more compact and hotter than in non-relativistic studies. An important application of our results would in fact be to provide better initial conditions for long-term disk evolutions, or for nucleosynthesis studies requiring a detailed knowledge of the disk structure and composition (e.g. [83]).

Finally, we note that the joint effects of shocks during the disk circularization, instabilities at the disk/tail interface, and neutrino absorption unbinds a small amount of material in the polar regions ($\sim 3 \times 10^{-4} M_{\odot}$). This might seem negligible compared to the material ejected dynamically in the equatorial plane during the disruption of the neutron star ($\sim 0.06 M_{\odot}$). However, this ejecta could be important because it is unbound in a direction in which no material has been ejected so far, and could thus impact the formation of a relativistic jet. Additionally, over longer timescales, we expect neutrino-powered winds to become active and eject material in the polar regions, maybe of the order of 1% of the mass of the disk. Later on, viscously-driven outflows could eject a more significant amount of material (probably 5% – 25% of the disk). But the outflows observed here will be the outermost layer of ejected material in the polar region. Their opacity might affect the properties of electromagnetic transients for observers in directions close to the spin axis of the black hole. From our simulations, it appears that most of the matter in these outflows is too neutron rich and cold to avoid strong r-process nucleosynthesis and the formation of high-opacity lanthanides, and thus that these early outflows could obscure later disk winds. However, we caution that the ejected mass and geometry of the outflows are likely to depend on the parameters of the binary, and thus that it would be dangerous to draw overly generic conclusions from a single initial configuration.

With these results demonstrating the ability of the SpEC code to evolve neutrinos within the moment formalism, we are now in a position to improve on the short simulations presented here. The gray scheme used in this work, which would be inadequate for the study of core collapse supernovae, appears at this point sufficient for the study of neutron star mergers.

ACKNOWLEDGMENTS

The authors wish to thank Brett Deaton, Rodrigo Fernandez, and Dan Kasen for useful discussions over the course of this project, and the members of the SXS collaboration for their suggestions and support. F.F. gratefully acknowledges support from the Vincent and Beatrice Tremaine Postdoctoral Fellowship. Support for this work was provided by NASA through Einstein Postdoctoral Fellowship grants numbered PF4-150122 (F.F.) and PF3-140114 (L.R.) awarded by the Chandra X-ray Center, which is operated by the Smithsonian Astrophysical Observatory for NASA under contract NAS8-03060; and through Hubble Fellowship grant number

51344.001 awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555. The authors at CITA gratefully acknowledge support from the NSERC Canada. M.D.D. acknowledges support through NSF Grant PHY-1402916. L.K. acknowledges support from NSF grants PHY-1306125 and AST-1333129 at Cornell, while the authors at Caltech acknowledge support from NSF Grants PHY-1068881, PHY-1404569, AST-1205732 and AST-1333520, and from NSF CAREER Award PHY-1151197. Authors at both Cornell and Caltech also thank the Sherman Fairchild Foundation for their support.

Computations were performed on the supercomputer Briarée from the Université de Montréal, and Guillimin from McGill University, both managed by Calcul Québec and Compute Canada. The operation of these supercomputers is funded by the Canada Foundation for Innovation (CFI), NanoQuébec, RMGA and the Fonds de recherche du Québec - Nature et Technologie (FRQ-NT). Computations were also performed on the Zwicky cluster at Caltech, supported by the Sherman Fairchild Foundation and by NSF award PHY-0960291. This work also used the Extreme Science and Engineering Discovery Environment (XSEDE) through allocation No. TG-PHY990007N, supported by NSF Grant No. ACI-1053575.

-
- [1] G. M. Harry (for the LIGO Scientific Collaboration), *Class. Quantum Grav.* **27**, 084006 (2010).
- [2] The Virgo Collaboration, “Advanced Virgo Baseline Design,” (2010), [vIR027A09](https://arxiv.org/abs/1007.2655).
- [3] J. Abadie *et al.* (LIGO Scientific Collaboration), *Class. Quant. Grav.* **27**, 173001 (2010), [arXiv:1003.2480 \[gr-qc\]](https://arxiv.org/abs/1003.2480).
- [4] R. O’Shaughnessy, C. Kim, V. Kalogera, and K. Belczynski, *Astrophys. J.* **672**, 479 (2008), [arXiv:astro-ph/0610076](https://arxiv.org/abs/astro-ph/0610076).
- [5] D. Gerosa, M. Kesden, E. Berti, R. O’Shaughnessy, and U. Sperhake, *Phys.Rev.D* **87**, 104028 (2013), [arXiv:1302.4442 \[gr-qc\]](https://arxiv.org/abs/1302.4442).
- [6] J. S. Read, C. Markakis, M. Shibata, K. Uryū, J. D. E. Creighton, and J. L. Friedman, *Phys. Rev. D* **79**, 124033 (2009), [arXiv:0901.3258 \[gr-qc\]](https://arxiv.org/abs/0901.3258).
- [7] J. S. Read, L. Baiotti, J. D. E. Creighton, J. L. Friedman, B. Giacomazzo, K. Kyutoku, C. Markakis, L. Rezzolla, M. Shibata, and K. Taniguchi, *Phys. Rev. D* **88**, 044042 (2013), [arXiv:1306.4065 \[gr-qc\]](https://arxiv.org/abs/1306.4065).
- [8] B. D. Lackey, K. Kyutoku, M. Shibata, P. R. Brady, and J. L. Friedman, *Phys. Rev. D* **89**, 043009 (2014), [arXiv:1303.6298 \[gr-qc\]](https://arxiv.org/abs/1303.6298).
- [9] W. Del Pozzo, T. G. F. Li, M. Agathos, C. Van Den Broeck, and S. Vitale, *Physical Review Letters* **111**, 071101 (2013), [arXiv:1307.8338 \[gr-qc\]](https://arxiv.org/abs/1307.8338).
- [10] B. D. Metzger and E. Berger, *Astrophys. J.* **746**, 48 (2012), [arXiv:1108.6056 \[astro-ph.HE\]](https://arxiv.org/abs/1108.6056).
- [11] L. F. Roberts, D. Kasen, W. H. Lee, and E. Ramirez-Ruiz, *Astrophys. J. Lett.* **736**, L21 (2011), [arXiv:1104.5504 \[astro-ph.HE\]](https://arxiv.org/abs/1104.5504).
- [12] S. Goriely, A. Bauswein, and H.-T. Janka, *Astrophys. J. Lett.* **738**, L32 (2011), [arXiv:1107.0899 \[astro-ph.SR\]](https://arxiv.org/abs/1107.0899).
- [13] O. Korobkin, S. Rosswog, A. Arcones, and C. Winteler, *Monthly Notices of the Royal Astronomical Society* **426**, 1940 (2012), [arXiv:1206.2379 \[astro-ph.SR\]](https://arxiv.org/abs/1206.2379).
- [14] A. Bauswein, S. Goriely, and H.-T. Janka, *ApJ* **773**, 78 (2013), [arXiv:1302.6530 \[astro-ph.SR\]](https://arxiv.org/abs/1302.6530).
- [15] S. Wanajo, Y. Sekiguchi, N. Nishimura, K. Kiuchi, K. Kyutoku, and M. Shibata, *ApJ Lett.* **789**, L39 (2014), [arXiv:1402.7317 \[astro-ph.SR\]](https://arxiv.org/abs/1402.7317).
- [16] M. Tanaka and K. Hotokezaka, *Astrophys.J.* **775**, 113 (2013), [arXiv:1306.3742 \[astro-ph.HE\]](https://arxiv.org/abs/1306.3742).
- [17] J. Barnes and D. Kasen, *ApJ* **775**, 18 (2013), [arXiv:1303.5787 \[astro-ph.HE\]](https://arxiv.org/abs/1303.5787).
- [18] <http://www.black-holes.org/SpEC.html>.
- [19] K. S. Thorne, *MNRAS* **194**, 439 (1981).
- [20] K. Hotokezaka, K. Kyutoku, H. Okawa, M. Shibata, and K. Kiuchi, *Phys. Rev. D* **83**, 124008 (2011), [arXiv:1105.4370 \[astro-ph.HE\]](https://arxiv.org/abs/1105.4370).
- [21] K. Takami, L. Rezzolla, and L. Baiotti, *Phys. Rev. Lett.* **113**, 091104 (2014), [arXiv:1403.5672 \[gr-qc\]](https://arxiv.org/abs/1403.5672).
- [22] M. Ruffert, H.-T. Janka, and G. Schaefler, *Astron. Astrophys.* **311**, 532 (1996), [astro-ph/9509006](https://arxiv.org/abs/astro-ph/9509006).
- [23] S. Rosswog and M. B. Davies, *Mon. Not. Roy. Astr. Soc.* **334**, 481 (2002).
- [24] S. Rosswog, *Astrophys. J.* **634**, 1202 (2005), [astro-ph/0508138](https://arxiv.org/abs/astro-ph/0508138).
- [25] R. Oechslin, H. T. Janka, and A. Marek, *Astron. Astrophys.* **467**, 395 (2007), [arXiv:astro-ph/0611047](https://arxiv.org/abs/astro-ph/0611047).
- [26] A. Bauswein, H.-T. Janka, K. Hebeler, and A. Schwenk, *Phys. Rev. D* **86**, 063001 (2012), [arXiv:1204.1888 \[astro-ph.SR\]](https://arxiv.org/abs/1204.1888).
- [27] A. Bauswein, N. Stergioulas, and H.-T. Janka, *Phys. Rev. D* **90**, 023002 (2014), [arXiv:1403.5301 \[astro-ph.SR\]](https://arxiv.org/abs/1403.5301).
- [28] K. Hotokezaka, K. Kiuchi, K. Kyutoku, H. Okawa, Y.-i. Sekiguchi, M. Shibata, and K. Taniguchi, *Phys. Rev. D* **87**, 024001 (2013), [arXiv:1212.0905 \[astro-ph.HE\]](https://arxiv.org/abs/1212.0905).
- [29] M. Tanaka, K. Hotokezaka, K. Kyutoku, S. Wanajo, K. Kiuchi, Y. Sekiguchi, and M. Shibata, *ApJ* **780**, 31 (2014), [arXiv:1310.2774 \[astro-ph.HE\]](https://arxiv.org/abs/1310.2774).
- [30] M. D. Duez, F. Foucart, L. E. Kidder, C. D. Ott, and S. A. Teukolsky, *Class. Quant. Grav.* **27**, 114106 (2010), [arXiv:0912.3528 \[astro-ph.HE\]](https://arxiv.org/abs/0912.3528).
- [31] M. B. Deaton, M. D. Duez, F. Foucart, E. O’Connor, C. D. Ott, L. E. Kidder, C. D. Muhlberger, M. A. Scheel, and B. Szilagyi, *Astrophys. J.* **776**, 47 (2013), [arXiv:1304.3384 \[astro-ph.HE\]](https://arxiv.org/abs/1304.3384).
- [32] F. Foucart, M. B. Deaton, M. D. Duez, E. O’Connor, C. D. Ott, R. Haas, L. E. Kidder, H. P. Pfeiffer, M. A. Scheel, and B. Szilagyi, *Phys. Rev. D* **90**, 024026 (2014), [arXiv:1405.1121 \[astro-ph.HE\]](https://arxiv.org/abs/1405.1121).
- [33] D. Neilsen, S. L. Liebling, M. Anderson, L. Lehner, E. O’Connor, *et al.*, *Phys. Rev. D* **89**, 104029 (2014), [arXiv:1403.3680 \[gr-qc\]](https://arxiv.org/abs/1403.3680).
- [34] J. M. Lattimer and Y. Lim, *ApJ* **771**, 51 (2013), [arXiv:1203.4286 \[nucl-th\]](https://arxiv.org/abs/1203.4286).
- [35] A. W. Steiner, M. Hempel, and T. Fischer, *Astrophys. J.* **774**, 17 (2013), [arXiv:1207.2184 \[astro-ph.SR\]](https://arxiv.org/abs/1207.2184).
- [36] J. M. Lattimer and F. D. Swesty, *Nucl. Phys.* **A535**, 331 (1991).
- [37] L. Lehner and F. Pretorius, *Ann.Rev.of Astron. & Astroph.* **52**, 661 (2014), [arXiv:1405.4840 \[astro-ph.HE\]](https://arxiv.org/abs/1405.4840).
- [38] K. Kiuchi, K. Kyutoku, Y. Sekiguchi, M. Shibata, and T. Wada, *Phys. Rev. D* **90**, 041502 (2014), [arXiv:1407.2660 \[astro-ph.HE\]](https://arxiv.org/abs/1407.2660).

- [39] K. Dionysopoulou, D. Alic, C. Palenzuela, L. Rezzolla, and B. Giacomazzo, *Phys. Rev. D* **88**, 044020 (2013), arXiv:1208.3487 [gr-qc].
- [40] C. Palenzuela, L. Lehner, M. Ponce, S. L. Liebling, M. Anderson, *et al.*, *Phys. Rev. Lett.* **111**, 061105 (2013), arXiv:1301.7074 [gr-qc].
- [41] M. Ponce, C. Palenzuela, L. Lehner, and S. L. Liebling, *Phys. Rev. D* **90**, 044007 (2014), arXiv:1404.0692 [gr-qc].
- [42] K. Dionysopoulou, D. Alic, and L. Rezzolla, *ArXiv e-prints* (2015), arXiv:1502.02021 [gr-qc].
- [43] M. Shibata, K. Kiuchi, Y. Sekiguchi, and Y. Suwa, *Progress of Theoretical Physics* **125**, 1255 (2011), arXiv:1104.3937 [astro-ph.HE].
- [44] C. Y. Cardall, E. Endeve, and A. Mezzacappa, *Phys. Rev. D* **87**, 103004 (2013), arXiv:1209.2151 [astro-ph.HE].
- [45] G. N. Minerbo, *J.Quant.Sec.Radiat.Transf.* **20**, 541 (1978).
- [46] A. Burrows, S. Reddy, and T. A. Thompson, *Nuc. Phys. A* **777**, 356 (2006).
- [47] M. D. Duez, F. Foucart, L. E. Kidder, H. P. Pfeiffer, M. A. Scheel, and S. A. Teukolsky, *Phys. Rev. D* **78**, 104015 (2008).
- [48] F. Foucart, M. B. Deaton, M. D. Duez, L. E. Kidder, I. Macdonald, C. D. Ott, H. P. Pfeiffer, M. A. Scheel, B. Szilágyi, and S. A. Teukolsky, *Phys. Rev. D* **87**, 084006 (2013), arXiv:1212.4810 [gr-qc].
- [49] E. O'Connor and C. D. Ott, *Class. Quantum Grav.* **27**, 114103 (2010).
- [50] F. Özel, D. Psaltis, R. Narayan, and J. E. McClintock, *Astrophys.J.* **725**, 1918 (2010), arXiv:1006.2834 [astro-ph.GA].
- [51] L. Kreidberg, C. D. Bailyn, W. M. Farr, and V. Kalogera, *Astroph.J.* **757**, 36 (2012), arXiv:1205.1805 [astro-ph.HE].
- [52] I. H. Stairs, *Science* **304**, 547 (2004).
- [53] D. R. Lorimer, *Living Reviews in Relativity* **11**, 8 (2008), arXiv:0811.0762.
- [54] J. M. Lattimer and M. Prakash, arXiv:1012.3208 (2010), arXiv:1012.3208 [astro-ph.SR].
- [55] F. Foucart, *Phys. Rev. D* **86**, 124007 (2012), arXiv:1207.6304 [astro-ph.HE].
- [56] S. Setiawan, M. Ruffert, and H.-T. Janka, *Astron. Astrophys.* **458**, 553 (2006).
- [57] V. Paschalidis, M. Ruiz, and S. L. Shapiro, arXiv:1410.7392 (2014), arXiv:1410.7392 [astro-ph.HE].
- [58] Z. B. Etienne, Y. T. Liu, V. Paschalidis, and S. L. Shapiro, *Phys. Rev. D* **85**, 064029 (2012), arXiv:1112.0568 [astro-ph.HE].
- [59] Z. B. Etienne, V. Paschalidis, and S. L. Shapiro, *Phys. Rev. D* **86**, 084026 (2012), arXiv:1209.1632 [astro-ph.HE].
- [60] W. H. Lee, E. Ramirez-Ruiz, and D. Page, *Astrophys. J.* **632**, 421 (2005), arXiv:astro-ph/0506121.
- [61] R. Fernández and B. D. Metzger, *MNRAS* **435**, 502 (2013), arXiv:1304.6720 [astro-ph.HE].
- [62] R. Fernández, D. Kasen, B. D. Metzger, and E. Quataert, *ArXiv e-prints* (2014), arXiv:1409.4426 [astro-ph.HE].
- [63] O. Just, A. Bauswein, R. Ardevol Pulpillo, S. Goriely, and H.-T. Janka, arXiv:1406.2687 (2014), arXiv:1406.2687 [astro-ph.SR].
- [64] F. H. Seguin, *Astrophys. J.* **197**, 745 (1975).
- [65] J. M. Lattimer and D. N. Schramm, *Astrophys. J.* **210**, 549 (1976).
- [66] D. Argast, M. Samland, F.-K. Thielemann, and Y.-Z. Qian, *A&A* **416**, 997 (2004), astro-ph/0309237.
- [67] F. Matteucci, D. Romano, A. Arcones, O. Korobkin, and S. Rosswog, *MNRAS* **438**, 2177 (2014), arXiv:1311.6980 [astro-ph.GA].
- [68] F. van de Voort, E. Quataert, P. F. Hopkins, D. Kereš, and C.-A. Faucher-Giguère, *MNRAS* **447**, 140 (2015), arXiv:1407.7039.
- [69] S. Shen, R. Cooke, E. Ramirez-Ruiz, P. Madau, L. Mayer, and J. Guedes, *ArXiv e-prints* (2014), arXiv:1407.3796.
- [70] L.-X. Li and B. Paczynski, *Astrophys.J.* **507**, L59 (1998), arXiv:astro-ph/9807272 [astro-ph].
- [71] D. Kasen, N. R. Badnell, and J. Barnes, *ApJ* **774**, 25 (2013), arXiv:1303.5788 [astro-ph.HE].
- [72] D. Grossman, O. Korobkin, S. Rosswog, and T. Piran, .
- [73] R. Surman, G. C. McLaughlin, and N. Sabbatino, *ApJ* **743**, 155 (2011), arXiv:1112.2673 [astro-ph.HE].
- [74] R. Fernández, E. Quataert, J. Schwab, D. Kasen, and S. Rosswog, *ArXiv e-prints* (2014), arXiv:1412.5588 [astro-ph.HE].
- [75] K. Kyutoku, K. Ioka, and M. Shibata, *Phys. Rev. D* **88**, 041503 (2013), arXiv:1305.6309 [astro-ph.HE].
- [76] G. Lovelace, M. D. Duez, F. Foucart, L. E. Kidder, H. P. Pfeiffer, M. A. Scheel, and B. Szilágyi, *Class. Quantum Grav.* **30**, 135004 (2013), arXiv:1302.6297 [gr-qc].
- [77] A. Perego, S. Rosswog, R. M. Cabezón, O. Korobkin, R. Käppeli, A. Arcones, and M. Liebendörfer, *MNRAS* **443**, 3134 (2014), arXiv:1405.6730 [astro-ph.HE].
- [78] L. Dessart, C. D. Ott, A. Burrows, S. Rosswog, and E. Livne, *Astrophys. J.* **690**, 1681 (2009).
- [79] W. H. Lee, E. Ramirez-Ruiz, and Diego-Lopez-Camara, *Astrophys.J.* **699**, L93 (2009), arXiv:0904.3752 [astro-ph.HE].
- [80] Y. Masada and T. Sano, *ApJ* **689**, 1234 (2008), arXiv:0808.2338.
- [81] J. Guilet, E. Mueller, and H.-T. Janka, *ArXiv e-prints* (2014), arXiv:1410.1874 [astro-ph.HE].
- [82] S. Rosswog and M. Liebendörfer, *Mon. Not. Roy. Astron. Soc.* **342**, 673 (2003), arXiv:astro-ph/0302301 [astro-ph].
- [83] R. Surman, O. Caballero, G. McLaughlin, O. Just, and H. T. Janka, *J.Phys.* **G41**, 044006 (2014), arXiv:1312.1199 [astro-ph.SR].
- [84] P. L. Roe, *Annual Review of Fluid Mechanics* **18**, 337 (1986).
- [85] H. Nessyahu and E. Tadmor, *Journal of Computational Physics* **87**, 408 (1990).
- [86] X.-D. Liu, S. Osher, and T. Chan, *Journal of Computational Physics* **115**, 200 (1994).
- [87] G.-S. Jiang and C.-W. Shu, *Journal of Computational Physics* **126**, 202 (1996).
- [88] S. Jin and C. D. Levermore, *Journal of Computational Physics* **126**, 449 (1996).
- [89] E. Audit, P. Charrier, J. Chièze, and B. Dubroca, *ArXiv Astrophysics e-prints* (2002), astro-ph/0206281.
- [90] J. C. McKinney, A. Tchekhovskoy, A. Sadowski, and R. Narayan, *MNRAS* **441**, 3177 (2014), arXiv:1312.6127 [astro-ph.CO].
- [91] J. M. Smit, J. Cernohorsky, and C. P. Dullemond, *A&A* **325**, 203 (1997).
- [92] C. D. Ott, A. Burrows, L. Dessart, and E. Livne, *Astrophys. J.* **685**, 1069 (2008).
- [93] E. Abdikamalov, A. Burrows, C. D. Ott, F. Löffler, E. O'Connor, J. C. Dolence, and E. Schnetter, *Astrophys. J.* **755**, 111 (2012), arXiv:1203.2915 [astro-ph.SR].
- [94] E. O'Connor, *ArXiv e-prints* (2014), arXiv:1411.7058 [astro-ph.HE].

Appendix A: Closure relation

The choice of a closure $P^{ij}(E, F_k)$ is the main approximation used in the moment formalism. In this work, we use the M1 closure, which relies on an interpolation between the expected closure relation in the optically thin and optically thick

regimes. Although we evolve E and F_i , the closure relation is generally easier to express as a function of the fluid frame quantities J and H^μ . Following [43], we define the parameter ζ as

$$\zeta^2 = \frac{H^\alpha H_\alpha}{J^2}, \quad (\text{A1})$$

such that $\zeta \sim 0$ in the optically thick limit and $\zeta \sim 1$ in the optically thin limit. The closure is then

$$P^{ij} = \frac{3\chi(\zeta) - 1}{2} P_{\text{thin}}^{ij} + \frac{3(1 - \chi(\zeta))}{2} P_{\text{thick}}^{ij}. \quad (\text{A2})$$

By default, we use the Minerbo closure [45]

$$\chi(\zeta) = \frac{1}{3} + \zeta^2 \frac{6 - 2\zeta + 6\zeta^2}{15}. \quad (\text{A3})$$

In the optically thin limit, we then use

$$P_{\text{thin}}^{ij} = \frac{F^i F^j}{F^k F_k} E. \quad (\text{A4})$$

Although this expression is exact in the limit $E^2 = F^i F_i$, Shibata *et al.* [43] have shown that it does not respect causality when $E^2 > F^i F_i$. However, proposed alternatives have more serious limitations [43], and P_{thin}^{ij} only matters when $E^2 \sim F^i F_i$.

For the optically thick limit, we choose

$$S_{\text{thick}}^{\mu\nu} = h^{\mu\nu} \frac{J_{\text{thick}}}{3}, \quad (\text{A5})$$

which is equivalent to

$$J_{\text{thick}} = \frac{3}{2W^2 + 1} [(2W^2 - 1)E - 2W^2 F^i v_i], \quad (\text{A6})$$

$$\gamma_\beta^\alpha H_{\text{thick}}^\beta = \frac{F^\alpha}{W} + \frac{W v^\alpha}{2W^2 + 1} [(4W^2 + 1)v_i F^i - 4W^2 E].$$

P_{thick}^{ij} can then be computed from Eq. (14).

In practice, because P^{ij} is a function of ζ , which itself depends on J, H^α , which are themselves functions of P^{ij} , the equations that we just described can only be solved through the use of a root-finding algorithm. We thus define

$$R(\zeta; E, F_i) = \frac{\zeta^2 J^2 - H^\alpha H_\alpha}{E^2}, \quad (\text{A7})$$

and solve for $R(\zeta; E, F_i) = 0$ using a Newton-Raphson algorithm (with ζ initialized at its last computed value at the given point). In Eq. (A7), J and H^μ are computed from E, F_i and P_{ij} , where P_{ij} is now itself a known function of E, F_i and ζ . Because the neutrino moments typically vary slowly, only a few iterations are required to get an accurate estimate of ξ . If cases in which the Newton-Raphson algorithm does not converge in 10 iterations, we fall back onto a simpler bisection method.

Appendix B: Energy integrated source terms

To evolve the energy-integrated moments E and F_i of the neutrino distribution function $f_{(\nu)}$, we need to define the energy-integrated emissivity $\bar{\eta}$ and energy-averaged opacities ($\bar{\kappa}_a, \bar{\kappa}_s$) of the fluid. In theory, these source terms are functions of $f_{(\nu)}$. However, as we only evolve energy-integrated moments, we do not know the neutrino spectrum, and have instead to rely on estimates of $f_{(\nu)}$. There are multiple ways to do so, and we detail the various choices that we have implemented below. We note that a number of those choices rely on information obtained from a simpler leakage scheme for the treatment of neutrino cooling [31] (e.g. average energy of neutrinos, optical depth), so that even when we evolve the neutrino radiation using the moment formalism, we still leave the leakage scheme turned on — but without coupling it to the fluid equations. Although having to mix the leakage scheme and the moment formalism is not ideal, we believe that the improvements observed in the neutrino absorption rates when leveraging information from the leakage scheme about the average energy of neutrinos in optically thin regions (see below) are worth the price of running both schemes together. It may be possible to entirely abandon the leakage scheme if the same information could be reliably extracted by other means. The energy of the neutrinos is largely set by the temperature of the fluid in the region where neutrinos decouple from the matter, i.e. when ζ changes from $\zeta \sim 0$ in the optically thick regions to $\zeta \sim 1$ in the optically thin regions. Doing this reliably, if it is at all possible, would however require some fine-tuning and experimentation.

We can first compute the source terms assuming that the neutrinos obey a Fermi-Dirac distribution with temperature T and chemical potential μ_ν . In that case, we have

$$f_{(\nu)} = \frac{1}{1 + \exp[(\epsilon_\nu - \mu_\nu)/(k_B T)]}, \quad (\text{B1})$$

where ϵ_ν is the neutrino energy, μ_ν the neutrino chemical potential, k_B the Boltzmann constant, and T the temperature of the fluid.

We now have to choose the value of the chemical potentials $\mu_\nu(\rho_0, T, Y_e)$. The value of the chemical potentials for neutrinos in equilibrium with the fluid, μ_ν^{eq} , is taken directly from the equation of state table. We consider two different choices for μ_ν : either we set $\mu_\nu = \mu_\nu^{\text{eq}}$ everywhere, or we make the same choice as in many leakage codes,

$$\mu_\nu = \mu_\nu^{\text{eq}}(1 - e^{-\tau_\nu}), \quad (\text{B2})$$

which is chosen so that $\mu_\nu \rightarrow 0$ in the optically thin region. We will refer to these two choices as ‘equilibrium’ and ‘leakage’ chemical potentials. The choice of μ_ν in the low optical depth regions does not matter much in practice when using our default method for the computation of the source terms (described below), but can have an effect in some of the alternative schemes that we tested.

The energy integrated source terms are then computed as in our leakage code [31] (which is itself based on the GR1D code [49], and previous work by Ruffert *et al.* [22], Rosswog

and Liebendorfer [82] and Burrows et al. [46]): we compute the absorption opacity due to electron and positron capture, the free streaming emission due to e^+e^- pair annihilation, plasmon decay and nucleon-nucleon Bremsstrahlung, and the scattering opacities due to elastic scattering on nucleons and heavy nuclei. We note however that, for the emissivity in optically thick regions, we use the free streaming emissivity, while the leakage scheme replaces the emissivity by the diffusion rate in those regions. Indeed, diffusion through the optically thick regions is handled naturally when using the moment formalism.

In order to guarantee that the neutrinos are in equilibrium with the fluid in the optically thick regions, we then compute the free streaming emission due to charge current reactions and the absorption due to pair processes through an energy-integrated version of Kirchhoff's law. At a given energy ϵ_ν , Kirchhoff's law gives us a relation between the emissivity $\eta(\epsilon_\nu)$, the absorption opacity $\kappa_a(\epsilon_\nu)$, and the equilibrium spectrum of the neutrino radiation $B_\nu(\epsilon_\nu)$ ⁵:

$$\eta(\epsilon_\nu) = \kappa_a(\epsilon_\nu)B_\nu(\epsilon_\nu). \quad (\text{B3})$$

The absorption opacity entering the evolution equations for E and F_i , on the other hand, is the energy averaged

$$\bar{\kappa}_a = \frac{\int_0^\infty \kappa_a(\epsilon_\nu)I(\epsilon_\nu)d\epsilon_\nu}{\int_0^\infty I(\epsilon_\nu)d\epsilon_\nu}, \quad (\text{B4})$$

where $I(\epsilon_\nu)$ is the specific intensity of the neutrino radiation. In our code, when computing $\bar{\kappa}_a$ from $\bar{\eta}$ (for pair processes), or $\bar{\eta}$ from $\bar{\kappa}_a$ (for charged-current reactions), we assume that $I_\nu = B_\nu$. Then, we have

$$\begin{aligned} \bar{\eta} &= \int_0^\infty \eta(\epsilon_\nu)d\epsilon_\nu = \int_0^\infty \kappa_a(\epsilon_\nu)B_\nu(\epsilon_\nu) \\ &\approx \bar{\kappa}_a \int_0^\infty B_\nu(\epsilon_\nu)d\epsilon_\nu. \end{aligned} \quad (\text{B5})$$

As B_ν is a known function of the fluid properties, this equation can easily be enforced. In optically thick regions, this prescription will be accurate, as it maintains the equilibrium between the fluid and the neutrino radiation. In optically thin regions, on the other hand, the neutrino radiation can be far from equilibrium. Computing charged-current interactions from the energy-integrated Kirchhoff's law assuming an equilibrium spectrum can affect the relative emission of electron neutrinos and antineutrinos, and thus the evolution of the electron fraction in low-density regions. We thus consider an alternative to the application of Kirchhoff's law when computing the emissivity of charged-current reactions: we can smoothly interpolate between the value of the emissivity $\bar{\eta}_K$ predicted by the application of Kirchhoff's law and the emissivity $\bar{\eta}_{\text{free}}$ predicted by Ruffert et al. [22] for free emission in an optically thin region, using

$$\bar{\eta} = \bar{\eta}_K f(\tau_\nu) + \bar{\eta}_{\text{free}}[1 - f(\tau_\nu)], \quad (\text{B6})$$

with

$$\begin{aligned} f(\tau_\nu) &= 1 \text{ if } \tau_\nu > 2 \\ &= 0 \text{ if } \tau_\nu < 2/3 \\ &= \frac{\tau_\nu - 2/3}{4/3} \text{ otherwise.} \end{aligned} \quad (\text{B7})$$

Note that even when using this corrected emissivity, the opacity due to charged-current reactions is not modified. Accordingly, this corrected emissivity no longer satisfies Eq. (B5). Although apparently more ad-hoc than the previous method, this prescription gives the best results in our test of neutrino-matter interactions (see Appendix E). This is probably because the energy integrated emissivity $\bar{\eta}_{\text{free}}$ was specifically meant to approximate the emissivity in optically thin regions. We note that the deviations from Kirchhoff's law for the energy integrated $\bar{\kappa}_a$ and $\bar{\eta}_{\text{free}}$ come from the approximations made in computing energy-averaged Pauli blocking factors, as well as from neglecting the electron rest mass and the difference between the neutron and proton masses when integrating over neutrino energies.

Finally, we can go one step further in improving our evolution scheme by noting that for most reactions considered here, including the dominant charged-current reactions, the absorption and scattering cross-sections scale like ϵ_ν^2 . This means that in low-temperature regions, where the fluid is much colder than the neutrinos (which, presumably, are emitted in higher temperature regions), we largely underestimate $\bar{\kappa}_{a,s}$. We can thus apply the correction

$$\bar{\kappa}_{a,s} \rightarrow \bar{\kappa}_{a,s} \left[\max \left\{ 1, \left(\frac{\langle \epsilon_{\nu, \text{leak}}^2 \rangle}{\langle \epsilon_{\nu, \text{fluid}}^2 \rangle} \right) \right\} \right], \quad (\text{B8})$$

where $\langle \epsilon_{\nu, \text{leak}}^2 \rangle$ is the average of ϵ_ν^2 predicted by the leakage scheme (taken over the entire grid) while

$$\epsilon_{\nu, \text{fluid}}^2 = \frac{F_5(\eta_\nu)}{F_3(\eta_\nu)} T^2 \quad (\text{B9})$$

is the average of ϵ_ν^2 for neutrinos in equilibrium with the fluid. The average is weighted by the energy density of neutrinos and not the number density of neutrinos, as we are interested in the energy-averaged $\bar{\kappa}_{a,s}$ given by Eq. (B4). In our tests (see Appendix E), we find that using this corrected value of $\bar{\kappa}_{a,s}$ significantly improves the agreement between the gray scheme used in this work and energy-dependent radiation transport.

We thus leave open three choices when computing the source terms. The chemical potential can be obtained from its equilibrium value or using the 'leakage' prescription. The emissivity of charged-current reactions in the optically thin limit can be obtained from the opacities of the fluid using the energy-integrated version of Kirchhoff's law or from a direct estimate of the emissivity. And finally, the opacities in low temperature regions can be obtained by assuming equilibrium between the neutrinos and the fluid, or by applying a correction for the higher energy of the neutrinos. Our default algorithm is to apply the energy correction to $\bar{\kappa}_{a,s}$, to compute $\bar{\eta}$

⁵ Note that Eq. B3 is always true for charged-current reactions, but only in the equilibrium limit for pair processes.

in optically thin regions from a direct estimate of the emissivity, and to use the equilibrium chemical potential everywhere (although that latest choice has nearly no impact when combined with our prescription for $\bar{\eta}$). We should however note that, even though these choices lead to significant differences in our test based on the evolution of the post-bounce configuration of a core collapse simulation, they are in much better agreement in the case of binary mergers.

Appendix C: Computation of the fluxes at cell faces

When computing the fluxes at cell faces for the evolution of \bar{E} and \bar{F}_i , we first use shock-capturing methods to estimate the value of $(E, F_i/E)$ at cell faces from their value at cell centers. For the simulations presented in this paper, we use the 2nd order *minmod* reconstruction (see e.g. [84, 85]), although higher order reconstruction methods are available in SpEC (the fluid variables at cell faces, for example, are reconstructed using the WENO5 algorithm [86, 87]). These reconstruction methods are used with both left- and right-biased stencils, leaving us with two reconstructed values U_L and U_R for the evolved variable $U = (\bar{E}, \bar{F}_i)$ and \bar{F}_L^i and \bar{F}_R^i for the fluxes $\bar{F}^i = (\alpha\bar{F}^i - \beta^i\bar{E}, \alpha\bar{P}_j^i - \beta^i\bar{F}_j)$. The fluxes at cell interfaces are then approximated by the HLL formula

$$\bar{F} = \frac{c_+\bar{F}_L + c_-\bar{F}_R - c_+c_-(U_R - U_L)}{c_+ + c_-}, \quad (\text{C1})$$

where c_+ and c_- are the absolute values of the largest right- and left-going characteristic speeds of the evolution system (or zero if there is no left/right going characteristic speeds). For the characteristic speeds $\lambda_{1,2}$ at a cell interface along direction d , we use

$$p = \alpha \frac{v^d}{W}, \quad (\text{C2})$$

$$r = \sqrt{\alpha^2 g^{dd}(2W^2 + 1) - 2W^2 p^2}, \quad (\text{C3})$$

$$\lambda_{1,\text{thin}} = -\beta^d - \alpha \frac{|F^d|}{\sqrt{\bar{F}^i \bar{F}_i}}, \quad (\text{C4})$$

$$\lambda_{1,\text{thick}} = \min\left(-\beta^d + \frac{2pW^2 - r}{2W^2 + 1}, -\beta^d + p\right), \quad (\text{C5})$$

$$\lambda_{2,\text{thin}} = -\beta^d + \alpha \frac{|F^d|}{\sqrt{\bar{F}^i \bar{F}_i}}, \quad (\text{C6})$$

$$\lambda_{2,\text{thick}} = \min\left(-\beta^d + \frac{2pW^2 + r}{2W^2 + 1}, -\beta^d + p\right), \quad (\text{C7})$$

$$\lambda_i = \frac{3\chi(\zeta) - 1}{2} \lambda_{i,\text{thin}} + \frac{3(1 - \chi(\zeta))}{2} \lambda_{i,\text{thick}}. \quad (\text{C8})$$

where the characteristic speeds in the optically thick and thin limits are taken from [43] and the interpolation between the two regimes uses the same formula as for the closure.

This prescription works well in optically thin regions. In the optically thick limit, however, these fluxes do not properly reproduce the diffusion rate of the neutrinos through the fluid. We thus apply corrections to the flux \bar{F}_E of \bar{E} [88]

$$\bar{F}_{E,\text{corr}} = a\bar{F}_E + (1 - a)\bar{F}_{E,\text{asym}}, \quad (\text{C9})$$

with

$$a = \tanh \frac{1}{\bar{\kappa} \Delta x^d}, \quad (\text{C10})$$

$$\bar{\kappa}_{i+1/2} = \sqrt{(\bar{\kappa}_\alpha + \bar{\kappa}_s)_i (\bar{\kappa}_\alpha + \bar{\kappa}_s)_{i+1}}, \quad (\text{C11})$$

and where half-integer indices refer to values of the opacities at cell faces while integer indices refer to value of the opacities at cell centers. Here, d is the direction in which we are reconstructing, $\Delta x^d = \sqrt{g_{dd}(\Delta x_{\text{grid}}^d)^2}$ is the proper distance between two grid points along that direction, and Δx_{grid}^d the coordinate grid spacing along that direction. The asymptotic flux in the fluid rest frame, which corresponds to the flux in the diffusion limit, is [19]

$$H_\alpha = -\frac{1}{3\bar{\kappa}} \partial_\alpha J_{\text{thick}}. \quad (\text{C12})$$

Using this and equation A5 in equation 13 gives the asymptotic observer frame flux

$$\begin{aligned} \bar{F}_{E,\text{asym}} = & \sqrt{\gamma} \left(\frac{4}{3} W^2 \alpha v^d J_{\text{thick}} - \beta^d E \right) \\ & - \alpha \frac{W}{3\bar{\kappa}} (\gamma^{di} + v^d v^i) \sqrt{\gamma} \frac{dJ_{\text{thick}}}{dx^i}. \end{aligned} \quad (\text{C13})$$

Numerically, this term can be fairly complex to evaluate. It requires derivatives of the neutrino energy density in the fluid frame along all coordinate directions, estimated at cell faces. In practice, we compute the first and second term of $\bar{F}_{E,\text{asym}}$ separately. For the second term, which models the diffusion of neutrinos, all quantities are estimated by averaging the values at neighboring cell centers, except for $\bar{\kappa}$, which is given by Eq. (C11), and dJ/dx^d , for which we use

$$\left(\frac{dJ}{dx^d} \right)_{i+1/2} = \frac{J_{i+1} - J_i}{\Delta x^d}. \quad (\text{C14})$$

For the first term, which represents the advection of neutrinos with the fluid, we make separate estimates from the left and right states (U_L, U_R). For both states, we also compute the advection speed

$$c_{\text{adv}} = -\beta^d + 4\alpha \frac{W^2}{2W^2 + 1} v^d. \quad (\text{C15})$$

If both speeds are positive, we use the value computed from U_L . If both are negative, we use the value computed from U_R . If their signs disagree, the advection term is set to zero.

A simpler correction is also applied to the flux \bar{F}_F of \bar{F}_i , following [89]

$$\bar{F}_{F,\text{corr}} = A^2 \bar{F}_F + (1 - A^2) \frac{\bar{F}_{F,R} + \bar{F}_{F,L}}{2}, \quad (\text{C16})$$

with

$$A = \frac{1}{\bar{\kappa} \Delta x^d}. \quad (\text{C17})$$

Appendix D: Implicit time stepping

The collisional source terms in equations (16-17) can be very stiff in optically thick regions. If we were to treat those source terms explicitly, we would need to use prohibitively small time steps. Accordingly, we will split the evolution equations into implicit and explicit terms. The variables $(\tilde{E}_{n+1}, \tilde{F}_{i,n+1})$ evolved from $(\tilde{E}_n, \tilde{F}_{i,n})$ by taking a time step dt are given by

$$\begin{aligned} \frac{\tilde{E}_{n+1} - \tilde{E}_n}{dt} + \partial_j(\alpha \tilde{F}_n^j - \beta^j \tilde{E}_n) \\ = \alpha \left(\tilde{P}_n^{ij} K_{ij} - \tilde{F}_n^j \partial_j \ln \alpha - \tilde{S}_{n+1}^\alpha n_\alpha \right), \end{aligned} \quad (D1)$$

$$\begin{aligned} \frac{\tilde{F}_{n+1} - \tilde{F}_n}{dt} + \partial_j(\alpha \tilde{F}_{i,n}^j - \beta^j \tilde{F}_{i,n}) - \alpha \tilde{S}_{n+1}^\alpha \gamma_{i\alpha} \\ = \left(-\tilde{E}_n \partial_i \alpha + \tilde{F}_{k,n} \partial_i \beta^k + \frac{\alpha}{2} \tilde{P}_n^{jk} \partial_i \gamma_{jk} \right). \end{aligned} \quad (D2)$$

Solving this exactly would require the use of a four-dimensional non-linear root-finder algorithm to get $(\tilde{E}_{n+1}, \tilde{F}_{n+1})$. To limit the cost of each time step, we instead use a linear approximation to \tilde{S}_{n+1}^α ,

$$\tilde{S}_{n+1}^\alpha = \sqrt{\gamma \eta} u^\alpha + A^\alpha \tilde{E}_{n+1} + B^{\alpha j} \tilde{F}_{j,n+1}, \quad (D3)$$

where the coefficients A^α and $B^{\alpha j}$ are computed assuming that the closure factor ξ and the angle between the neutrino flux F_i and the fluid velocity v^i are constant. The system then reduces to four linear equations for $(\tilde{E}_{n+1}, \tilde{F}_{i,n+1})$, which can be simply solved by the inversion of a 4×4 matrix at each point and for each neutrino species. To ensure that this procedure is reasonable, however, we need an estimate of the time stepping error – which will help us choose the time step dt . We consider four different types of errors.

- Large fluxes may cause the explicit portion of the time stepping algorithm to be unstable. We thus require that

$$dt < \frac{\alpha_F \tilde{E} + \alpha_{\text{Abs}} \max(\tilde{E})}{|\partial_j(\alpha \tilde{F}^j - \beta^j \tilde{E})|} \quad (D4)$$

at any point where the denominator is positive. For post-merger evolutions, we typically choose $\alpha_F = 0.3$ and $\alpha_{\text{Abs}} = 0.001$. The maximum is taken over the entire computational domain.

- Strong coupling of the neutrino radiation to the fluid can cause large oscillations in the fluid quantities when the radiation transport code is first turned on, causing the evolution to be unstable. These swings, when they occur, are particularly noticeable in the evolution of the electron fraction Y_e . Accordingly, we define

$$\epsilon_Y = \frac{|\Delta Y_e|}{\alpha_{\text{Rel}}} \frac{\rho_0}{\rho_0 + \alpha_{\text{Abs}} \max(\rho_0)}, \quad (D5)$$

where ΔY_e is the change in Y_e over the last neutrino time step due to the coupling between the fluid and the neutrino radiation. For post-merger evolutions, we use $\alpha_{\text{Rel}} = 0.01$.

- The implicit portion of the time step might be inaccurate, for example because of the linearization of \tilde{S}^α . Accordingly, we solve the implicit problem twice: once with a time step dt , and once using two time steps $dt/2$. The explicit terms are kept identical for both time steps, but the linearization of \tilde{S}^α is recomputed at the intermediate point $t + dt/2$. For any variable U , we thus have two estimates U_1 and U_2 for $U(t + dt)$, using respectively one and two intermediate time steps. We can then obtain a second-order accurate estimate $U(t + dt) = 2U_2 - U_1$ and an estimate of the error, $\delta U = U_2 - U_1$. We then define

$$\epsilon_E = \frac{\delta \tilde{E}}{\alpha_{\text{Rel}} \tilde{E} + \alpha_{\text{Abs}} \max(\tilde{E})}, \quad (D6)$$

$$\epsilon_F = \frac{\gamma^{ij} \delta \tilde{F}_i \delta \tilde{F}_j}{\alpha_{\text{Rel}} \tilde{E} + \alpha_{\text{Abs}} \max(\tilde{E})}, \quad (D7)$$

as the errors from the implicit time step.

- Errors in the evolution of \tilde{E} and \tilde{F}_i can cause \tilde{E} to become negative, or the fluxes to violate causality ($\gamma^{ij} \tilde{F}_i \tilde{F}_j > \tilde{E}^2$). To be safe, we additionally define

$$\epsilon'_E = \frac{|\tilde{E}|}{\alpha_{\text{Rel}} \tilde{E} + \alpha_{\text{Abs}} \max(\tilde{E})} \quad (D8)$$

when $\tilde{E} < 0$ and

$$\epsilon'_F = \frac{\gamma^{ij} \tilde{F}_i \tilde{F}_j - \tilde{E}^2}{\alpha_{\text{Rel}} \tilde{E} + \alpha_{\text{Abs}} \max(\tilde{E})} \quad (D9)$$

when $\gamma^{ij} \tilde{F}_i \tilde{F}_j > \tilde{E}^2$. In practice, we find that these errors are generally smaller than ϵ_E, ϵ_F .

After taking a time step, we then define a global error ϵ as the largest error among all points, all neutrino species, and all five errors $\epsilon_F, \epsilon_E, \epsilon_Y, \epsilon'_E, \epsilon'_F$. If the inequality (D4) is not satisfied or if $\epsilon > 10$, we reject the last time step and start again with a new time step given either by (D4) or by

$$dt' = dt \sqrt{\frac{0.9}{\epsilon}}. \quad (D10)$$

Otherwise, we accept the time step and set the next time step for the evolution of neutrinos to be

$$dt' = dt \sqrt{\frac{0.9}{\max(\epsilon, 0.3)}}. \quad (D11)$$

Finally, if dt' is larger than the time step required for the neutrino evolution to catch up with the fluid evolution, we take a time step bringing the two sets of equations to the same time. In practice, for most of the post-merger evolution, the code ends up taking a single neutrino time step for each GR-Hydro time step, as can be expected for a quasi-equilibrium radiation field. The adaptive time stepping algorithm is thus mostly useful to get stable evolutions when the neutrino radiation is




FIG. 18. Energy density for the single beam test, after $\Delta t = 0.4$. The black arrows show the flux F_i .

first turned on, as well as during the plunge of the neutron star into the black hole. Occasionally, taking 2-3 neutrino time steps per hydrodynamical time step can also be necessary when time stepping is limited by the Courant condition, which is more restrictive for neutrinos than for the fluid.

Appendix E: Code Tests

In order to test the ability of our code to perform neutrino transport simulations in compact binary mergers, we perform a series of tests showing that the moment formalism has been properly implemented in SpEC, that it suffers from the same known limitations as in other codes, and finally that the choices made when averaging emissivities and opacities still allow us to reproduce reasonably well the output of an energy-dependent code in a spherically symmetric test evolving a post-bounce configuration taken from a core collapse supernova simulation.

1. Single beam in flat spacetime

The simplest test that we perform is the propagation of a beam of radiation in vacuum and for a flat spacetime. We use a 3-dimensional grid with spacing $\Delta x = 0.01$ and bounds $0 \leq x \leq 1$, $0 \leq y \leq 0.2$ and $0 \leq z \leq 0.2$. The field variables are frozen for $x < 0.1$, with the condition $E = 1$, $F_i = (0.9999999, 0, 0)$ inside of a “beam” confined to $0.05 \leq y \leq 0.15$, $0.05 \leq z \leq 0.15$, and are suppressed by a factor of 10^{-10} in the region outside of the beam.

The results of this test are shown in Fig. 18, at a time $t = t_0 + 0.4$. As expected, the beam propagates in a straight line and at the speed of light. Numerical errors cause a slight widening of the beam, but by only one grid spacing at an energy density around $10^{-3}E_{\text{beam}}$. The beam front is also smoothed upstream of the beam, with an exponential decay of about $10^{-1/2}$ per grid point. This pattern establishes itself quickly as the beam leaves the frozen region, and then propagates at the speed of light with the beam.

We then performed the same test, but for a beam no longer propagating along a coordinate direction. In this case, we freeze the region with $x < 0.03$ or $y > 0.17$, and set $F_i = (0.8, -0.6, 0)$ in the beam region. The results are shown in Fig. 19. The main difference with the aligned beam is that the edges of the beam are not as sharp (while the front of the




FIG. 19. Energy density for the oblique beam test, after $\Delta t = 0.2$. The arrows show the flux F_i .




FIG. 20. Energy density for the shadow test after $\Delta t = 1$. The arrows show the flux F_i .

beam is slightly sharper). The exponential decay of the energy away from the beam is now about the same in all directions.

2. Shadow

We now consider a uniform radiation field with $E = 1$ and $F_i = (0.9999999, 0, 0)$, hitting a sphere of optically thick material. In this test, we set the absorption opacity $\bar{\kappa}_a = 10^6$ within a sphere of radius $r_S = 0.05$ and centered on $\mathbf{c}_S = (0.5, 0.1, 0.1)$. The grid is identical to the one used for the beam tests. The results of the evolution are shown in Fig. 20.

The M1 closure is known to perform well for such a test, in which we expect to see the shadow of the optically thick sphere in the fields for $x > 0.5$ (see e.g. [90]). Indeed, we observe a clear shadow behind the sphere, with the formation of two independent beams each having properties similar to those observed in the previous tests. Our implementation of the M1 formalism thus appears to properly project the shadow of optically thick objects.

3. Single beam in curved spacetime

As our simulations require the evolution of the neutrinos close to a black hole, we want to determine how well the M1 formalism performs in a strongly curved spacetime. To do so, we perform another set of tests on beams propagating in vacuum, but now in a black hole spacetime. These tests are largely inspired from those presented in McKinney




FIG. 21. Energy density in steady-state for a 2D beam in a Kerr-like spacetime, with the beam emitted from $r = 7M - 8M$ and M the mass of the non-spinning central black hole. The solid lines show null geodesics bounding the expected location of the beam. The simulation is performed in full 3D, but the metric and beam are independent of z .

et al. [90] (on their Fig. 13 and Sec. 5.9). The tests in [90] are performed in 2D in spherical polar coordinates, however, while the SpEC code always uses 3D cartesian grids for radiation hydrodynamics. For the sake of comparison, we thus first consider a “Kerr-like” spacetime in which the metric is set by $g_{\mu\nu}(t, x, y, z) = g_{\mu\nu}^{Kerr}(0, x, 0, z)$, with $g_{\mu\nu}^{Kerr}$ the Kerr metric for a non-spinning black hole, in Kerr-Schild coordinates. This metric is unphysical, but allows us to consider effectively 2D beams without having to develop an axisymmetric version of our code. In all tests, the beams are emitted from a region of width $\Delta z = M$ in the $x = 0$ plane, in which $E = 1$ and the flux is chosen so that $F_i F^i = 0.998E^2$ and $\alpha F^i - \beta^i E$ is along the x -axis. We use a grid spacing $\Delta x = 0.2M$ (i.e. the beam is initially 5 grid points wide), where M is the black hole mass.

We first consider a beam emitted from a relatively large distance, $7M < z < 8M$, shown in Fig. 21. The energy density in the beam decreases as the neutrinos get farther from the black hole, in part due to the gravitational redshift (about a 10% effect over the length of the beam on the grid), and in part due to the spreading of the beam caused by the gravitational bending of the radiation. Within the accuracy shown in the previous tests, the beam remains otherwise well contained within the region delimited by the null geodesics defined by $\mathbf{x}(t = 0) = (0, 0, 8M)$, $dr/dt(t = 0) = 0$ and $\mathbf{x}(t = 0) = (0, 0, 7M)$, $dr/dt(t = 0) = 0$ (shown as solid black lines on the figure).

We now move on to a beam initially closer to the black hole, with $3M < z < 4M$. The inner edge of the beam then lies on the unstable circular photon orbit. Both the gravitational redshift effect and the divergence of the null geodesics are much stronger in this case, so that the beam widens and its energy density decreases quickly as the beam orbits the black hole. The results of the simulation are shown in Fig. 22. As before, we can check that the beam remains mostly within the region predicted for free-streaming massless particles.

Finally, we consider the same configuration but in full 3D.




FIG. 22. Same as in Fig. 21, but with the beam emitted from $r = 3M - 4M$. As before, the solid lines trace null-geodesics.




FIG. 23. Same as Fig. 22, but using the full 3-dimensional Kerr-metric and a beam of finite transverse size.

The background metric is now that of a non-spinning black hole in Kerr-Schild coordinates. In 3D, the null geodesics followed by our beam converge towards the equatorial plane, and should all cross on the $z = 0, y = 0$ axis. The numerical results are shown in Fig. 23. As in the 2D test, the widening of the beam is consistent with the predictions obtained by tracing null geodesics. The decrease in energy density due to the widening of the beam in the xy -plane and to the gravitational redshift is however compensated by the collapse of the beam along the z -axis, at least until the beam crosses the x -axis. In the $z < 0$ plane, the beam should rapidly widen again along the y -axis. In practice, however, the beam remains slightly more collimated than expected in that region. Although the qualitative differences are fairly mild in this case, it is a first indication of the problems intrinsic to the use of the M1 for-




FIG. 24. Contours of constant energy density for the crossing beams test. The contours are for 10%, 1% and 0.1% of the original energy density of the beams.

malism, which we will make clearer in the next test.

4. Crossing beams

The problems with converging beams, which we alluded to in the previous section, can be more easily observed if we consider a much simpler setup: two crossing beams in flat space-time. As we do not include any interaction between neutrinos, except indirectly through the effective opacity due to the inverse reactions of thermal processes, the two beams should simply pass through each other. However, this does not occur when using the moment formalism with the M1 closure. We show in Fig. 24 what happens when such a system is evolved: the beams collide and form a wider, single beam along the average direction of propagation of the crossing beams. This is, effectively, due to the difference between the approximate form of the second moment of the distribution function in the analytical M1 closure, and its true second moment.

This inability of the M1 formalism to deal with crossing beams, and the inaccuracies existing in the presence of converging beams, means that the evolution of the moments of the neutrino distribution function in the region along the polar axis of the black hole, as well as close to the inner edge of the disk, is not entirely reliable. The M1 formalism performs very well for diverging free-streaming neutrinos (which is what we see in most of the regions surrounding the disk), and in optically thick regions (as in the core of the disk, see next test). But its performance in regions in which higher moments of the distribution function are required to properly model the neutrino pressure is, by definition, quite poor (see also [90]).

5. Optically thick radiative sphere

The tests presented above are mainly intended to determine how well our code can evolve the neutrino moments for various geometric configurations in the free-streaming limit. In order to evolve neutron stars and the optically thick accretion disks created as a result of neutron star mergers, we also need to determine whether we can properly model optically thick regions and, more importantly maybe, how closely we can re-




FIG. 25. Energy density of neutrinos at $t = 2$ for the emitting sphere tests with $\bar{\kappa} = 5$ and $\bar{\kappa} = 10^{10}$. The numerical results are shown by circles/squares while the solid lines are the analytical predictions.

produce the transition between optically thick and optically thin regions. Indeed, these will determine how well we can predict the neutrino luminosity from neutron stars and accretion disks.

We first consider optically thick radiative spheres, for which we have an analytical solution for the distribution function [91]:

$$f(r, \mu) = b[1 - \exp(-\bar{\kappa}s(r, \mu))] \quad (\text{E1})$$

for a sphere of radius r_s and absorption opacity $\bar{\kappa}$, and for $\mu = \cos(\theta)$ with θ the angle between the neutrino momentum and the radial direction. The function $s(r, \mu)$ is given by

$$\begin{aligned} s(r, \mu) &= r\mu + r_s g(r, \mu) \quad [r < r_s; -1 < \mu < 1], \\ &= 2r_s g(r, \mu) \quad [r \geq r_s; \sqrt{1 - \left(\frac{r_s}{r}\right)^2} < \mu < 1], \\ &= 0 \quad \text{otherwise,} \end{aligned} \quad (\text{E2})$$

with

$$g(r, \mu) = \left(1 - \left(\frac{r}{r_s}\right)^2 (1 - \mu^2)\right). \quad (\text{E3})$$

We can then obtain the exact solution by numerical integration of $f(r, \mu)$ over μ . In this test, we consider two different regimes. First we use a sphere of moderate optical depth ($r_s = 1$, $\bar{\kappa} = 5$), which is fairly typical of the conditions in post-merger accretion disks. Then, we use a sphere of extremely high optical depth ($r_s = 1$, $\bar{\kappa} = 10^{10}$), which provides a sharp neutrinosphere similar to what may be found at the surface of a neutron star. In both cases, we use a 3-dimensional Cartesian grid with grid spacing $\Delta x = 0.04$. The results of the numerical evolutions are compared with the exact solution in Fig. 25. We find good agreement between the two solutions. For the most optically thick case, this is mostly a consequence of the corrections to high optical depth regions described in Appendix C. The configuration with $\bar{\kappa} = 5$, on the other hand, has a very small correction to the fluxes, as $\bar{\kappa}\Delta x = 0.2 < 1$.




FIG. 26. Temperature after 8 ms of evolution for the evolution of a post-bounce configuration, in the region in which the temperature evolution is most affected by the gray approximation.

6. Neutrino emission and coupling with matter in a post-bounce configuration

As a last test of our code, we consider a 1D profile constructed as a spherical average from a 2D, Newtonian, multi-neutrino energy, multi-neutrino angle neutrino transport simulation at 160 ms after core bounce [92]. Since we spherically average the 2D simulation, do not use precisely the same nuclear equation of state and neutrino opacities, and freeze the hydrodynamics, the initial profile is slightly out of equilibrium. Therefore, this provides a good test of the lepton and energy coupling and a venue for comparing with energy-dependent transport. Similar tests were carried on with this profile for testing Monte Carlo transport [93]. We compare the SpEC results with the output of the GR1D code [49]. For this test, GR1D uses an energy-dependent M1 scheme with 12 energy groups, and was itself shown to agree well with the results of full transport codes [94]. During this test, the matter density is fixed and the fluid is assumed to be at rest. But the internal energy and electron fraction of the fluid are coupled to the neutrino evolution. We perform this test with various choices of gray schemes: with the standard SpEC methods described in Appendix B (SpEC-Std), with the average energy of neutrinos always obtained by assuming equilibrium between the neutrinos and the fluid (i.e. ignoring the correction given by Eq. B8, SpEC- T_{fl}), and without the correction B6 to the charged-current emissivities in low-opacity regions (SpEC- η_K). The SpEC simulations, which are performed with the full 3D code assuming octant symmetry, have a fairly coarse resolution of 6 km at the lowest resolution, 3 km at the medium resolution, and 1.5 km at the highest resolution.

We first compare radial profiles of the fluid variables T and Y_e after 8 ms of evolution, shown in Figs. 26-27. The temperature profiles agree well with the GR1D results, as long as we correct the average energy of neutrinos according to




FIG. 27. Electron fraction Y_e after 8 ms of evolution for the evolution of a post-bounce configuration.

Eq. B8. If we do not, the absorption of neutrinos in the low-density regions is widely underestimated – and in particular, the code completely misses the existence of a gain region at $r > 100$ km. With correction B8, the heating in the gain region is reproduced better, but now occurs faster than expected. This is due to an overestimate of the average energy of neutrinos. Whether or not we modify the emissivities according to equation B6 does not appear to affect the temperature evolution.

The evolution of the electron fraction Y_e is more sensitive to the choices made when energy-integrating. With our standard choices, we find results close to GR1D, while the composition is widely inaccurate when neglecting correction B8, and the equilibrium composition can in some regions be wrong by $\Delta Y_e \sim 0.05$ when neglecting correction B6. The grid resolution does not significantly affect these results, except close to $r = 0$.

Finally, we observe over 20 ms the neutrino luminosity extracted on a sphere of radius $r = 250$ km, and compare the average energy-squared $\langle \epsilon_{\text{leak}}^2 \rangle$ predicted by the leakage (and used to compute opacities in colder regions) and the same quantity measured by the energy-dependent M1 code. We find that the luminosities are accurate to $\sim 10\%$ for ν_e and ν_x , and to 20% – 30% for $\bar{\nu}_e$. For comparison, the luminosities predicted by the leakage scheme are off by factors of 2-3 for ν_e and $\bar{\nu}_e$ (but are very good for ν_x), while the errors due to the finite grid resolution are $\sim 5\% - 10\%$. The energies $\sqrt{\langle \epsilon^2 \rangle}$ are typically overestimated by the leakage scheme, by up to 50%. This causes the larger-than-expected heating rate observed in the gain region ($r \gtrsim 90$ km). The reason for these large errors is that the neutrino energies in the hot, highly degenerate matter present at high optical depth are very large and significantly affect the average energy (due to weighting of the electron number emission rate by the square of the neutrino energies). In practice, however, these high energy neutrinos are thermalized as they propagate through the optically thick regions. Significant improvements in $\sqrt{\langle \epsilon^2 \rangle}$ can be obtained

if we define

$$\langle \epsilon^2 \rangle = \frac{\int R_\nu(\mathbf{x}) \langle \epsilon_\nu^2(\mathbf{x}) \rangle \min(1, \exp[\tau_{\text{threshold}} - \tau]) dV}{\int R_\nu(\mathbf{x}) \min(1, \exp[\tau_{\text{threshold}} - \tau]) dV}, \quad (\text{E4})$$

where R_ν is the neutrino number emission predicted by the leakage scheme, and τ is the optical depth. Any value of the threshold $\tau_{\text{threshold}} \sim 1 - 10$ allows us to recover $\sqrt{\langle \epsilon^2 \rangle}$ to within 20% for ν_e and $\bar{\nu}_e$, and within 30% for ν_x . In all cases, the finite resolution error is negligible compared with the error due to the gray approximation. We have checked that these variations are indeed due to the gray approximation, and not to the specific implementation of the M1 scheme in SpEC: an energy-dependent version of the M1 code in SpEC, which can easily be used in this case due to the relatively low computational cost of the test problem and the lack of gravitational redshift or velocity gradients, agrees with the GR1D results.

These results are already much better than if we got the neutrino energies by assuming equilibrium with the fluid: in the gain region, the equilibrium average neutrino energies are $\sqrt{\langle \epsilon^2 \rangle} \sim 5 - 10$ MeV while the actual average neutrino energies are $\sqrt{\langle \epsilon^2 \rangle} \sim 10 - 25$ MeV. And indeed, with this corrected $\langle \epsilon^2 \rangle$, heating in the gain region agrees well with the results of the GR1D simulations. The correction also causes changes to the evolution of Y_e , with results with and without correction B6 to the emissivities now bracketing the GR1D results (with typical errors $\Delta Y_e \sim 0.02$). The simulation without correction B6 still slightly overestimates Y_e around $r \sim 100$ km, while with correction B6 the simulation now slightly underestimates Y_e in that same region.

Such errors would be very significant in the context of core collapse supernova simulations, where properly modeling neutrinos is critical to the explosion mechanism. In compact binary mergers, however, the impact of neutrinos is more modest. The computation of $\langle \epsilon^2 \rangle$ from the leakage scheme is also more accurate in post-merger accretion disks: the temperature does not vary as much within post-merger accretion disks as in the test presented here, the disks are only moderately optically thick ($\tau \leq 10$), and matter within the disks reaches densities of at most $\rho_0 \sim 10^{12}$ g/cm³, at which only electrons are degenerate. Using emissivities with and without correction B6, we can get an estimate of the errors due to the use of a gray scheme. These errors, discussed in the main text of the paper, are much smaller than the errors observed in this test. Using the M1 formalism instead of our leakage scheme then significantly improves our ability to determine the cooling time and composition evolution of the disk, and gives us a reasonable first estimate of the energy deposition in the corona (except along the polar axis of the black hole, where the M1 approximation is unreliable).

Appendix F: Accuracy of the post-merger evolution

Most of the discussion of our simulations of the formation of an accretion disk after a neutron star-black hole merger (Sec. IV) focuses on the qualitative features of the system, and on the differences between various algorithms for the evolu-

tion of neutrinos. In this section, we will discuss the accuracy of these results, and argue that the features of the system discussed in Sec. IV are appropriately resolved by our simulations. We identify four main sources of error. The first is due to the finite resolution of our numerical grid during the simulations. The second is the numerical error in the simulation of the neutron star-black hole merger before we turn on the neutrino transport code, which causes errors in the initial conditions used for our simulations. The third is the approximate treatment of the neutrinos, and in particular the use of a gray scheme and of the M1 closure. And the fourth comes from only turning on the neutrino transport and changing the treatment of low-density material about 6.1 ms after merger. The first two can be estimated through the use of simulations at lower and/or higher resolution. The last two are more difficult to assess, and will only be rigorously measured through the use of more advanced (and more costly) simulations. Until such simulations are available, we have to rely on simpler estimates of these errors.

To determine the importance of the first source of error, we performed simulations of a post-merger accretion disk using 60^3 and 140^3 points for each level of refinement, instead of the 100^3 points used in our "standard" configuration. The higher resolution simulation was only evolved for 1.1ms, to verify that the solution was converging with resolution. Even the simulation using the coarsest grid shows surprisingly good agreement with our standard runs. The average temperature in the disk agrees to $\Delta \langle T \rangle \sim 0.2$ MeV and the electron fraction to $\Delta \langle Y_e \rangle \sim 0.01$. The neutrino luminosities agree to better than 20%, and the average neutrino energies within 0.5 MeV. This is comparable to the differences observed between various choices of gray approximations, and much smaller than the differences between the leakage and M1 simulations. Additionally, the difference between the standard and high resolutions is, for all observed quantities, more than a factor of two smaller than the difference between the low and standard resolutions. Most of those differences arise immediately after the neutrino transport is turned on, while evolution at later times is very similar for all numerical resolutions. Accordingly, we do not expect numerical resolution during the post-merger evolution to be a significant source of error.

The numerical error in the simulations before we turn on neutrino transport can easily be determined from the lower resolution simulations of the same system performed in Foucart *et al.* 2014 [32]. The largest error in these simulations was in the determination of the properties of the dynamical ejecta, which does not concern us here as that material is allowed to escape the numerical grid. At the time at which we begin the simulations with neutrino transport, we otherwise find relative errors of less than 10% in the total mass outside of the black hole, average temperature and average electron fraction in the disk. Considering that the grid spacing used in [32] was about a factor of two coarser than the grid spacing used in the standard simulations of this paper, these are clearly overestimates of the numerical error. The only numerical error which could affect our results is thus the initial temperature of the disk, which generally decreases with resolution. The disk could be, on average, about 0.4 MeV cooler at infinite

resolution. This is only slightly smaller than the difference between leakage and M1 simulations, but does not affect the conclusion of this paper: both the leakage and M1 simulations presented here use the same imperfect initial data. We should also note that even our imperfect initial data remains a much better starting point than the commonly used alternative, an equilibrium torus of constant entropy.

The effect of an approximate treatment of the neutrinos is discussed in Sec. IV G, to which we refer the interested reader. Here we will simply note that our rough estimate of the error due to the use of a gray scheme is comparable to our estimate of the error due to numerical resolution, and much smaller than the differences between simulations using the M1 scheme and the leakage scheme. However, a true measure of the error would require a simulation using an energy-dependent transport scheme without the weaknesses of the M1 closure. Such a simulation is currently out of reach for our code.

Finally, we have to consider the impact of suddenly turning on the neutrino transport code 6.1 ms after merger, and of the use of a higher threshold for the application of atmosphere corrections at earlier times. The only way to rigorously measure this would be to perform a simulation using the M1 code and a lower atmosphere threshold, starting before the disruption of the neutron star. Now that we have a well-tested M1 code and good estimates of the atmosphere thresholds which should be used after merger, we intend to perform such sim-

ulations. However, the disruption of the neutron star and accretion of the neutron star core onto the black hole is the most computationally intensive part of the evolution, while we believe that the simulations performed here capture the most important properties of the merger and post-merger evolution (except for the absence of magnetic fields). Indeed, we begin the simulation around the time at which the neutrino luminosity and the electron fraction begin to significantly increase due to the heating of the disk and the emission of neutrinos. We also find that the disk outflows are not immediately launched when we decrease the atmosphere threshold. At the beginning of the simulation, those outflows are still contained by material from the tidal tail falling onto the forming disk. It is only about 2 ms later that outflows begin to form above the disk. A potentially more important source of error is the transient occurring when the transport code is turned on. To study its effects, we consider two possible initializations of the moments of the neutrino distribution function. First, we initialize them assuming that the neutrinos are in equilibrium with the fluid, in which case the neutrino energy density is overestimated, and the initial transient consists in a transfer of energy from the neutrinos to the fluid and a slight decrease of the average electron fraction. Then, we initialize the neutrino energy density to a negligible value, in which case the transient has the opposite effect. We verify that after about 2 ms the two solutions are in good agreement (at early times, this transient is of course the largest source of error in the simulation).