



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Gauge-invariant implementation of the Abelian-Higgs model on optical lattices

A. Bazavov, Y. Meurice, S.-W. Tsai, J. Unmuth-Yockey, and Jin Zhang

Phys. Rev. D **92**, 076003 — Published 14 October 2015

DOI: [10.1103/PhysRevD.92.076003](https://doi.org/10.1103/PhysRevD.92.076003)

Gauge-invariant implementation of the Abelian Higgs model on optical lattices

A. Bazavov^{1,2}, Y. Meurice¹, S.-W. Tsai², J. Unmuth-Yockey¹, and Jin Zhang²

¹ *Department of Physics and Astronomy, The University of Iowa, Iowa City, Iowa 52242, USA and*

² *Department of Physics and Astronomy, University of California, Riverside, CA 92521, USA*

(Dated: September 14, 2015)

We present a gauge-invariant effective action for the Abelian Higgs model (scalar electrodynamics) with a chemical potential μ on a 1+1 dimensional lattice. This formulation provides an expansion in the hopping parameter κ which we test with Monte Carlo simulations for a broad range of the inverse gauge coupling $\beta_{pl} = 1/g^2$ and small values of the scalar self-coupling λ . In the opposite limit of infinitely large λ , the partition function can be written as a traced product of local tensors which allows us to write exact blocking formulas. Gauss's law is automatically satisfied and the introduction of μ has consequences only if we have an external electric field, $g^2 = 0$ or an explicit gauge symmetry breaking. The time continuum limit of the blocked transfer matrix can be obtained numerically and, for $g^2 = 0$ and a spin-1 truncation, the small volume energy spectrum is identical to the low energy spectrum of a two-species Bose-Hubbard model in the limit of large on-site repulsion. We extend this procedure for finite β_{pl} and derive a spin-1 approximation of the Hamiltonian. It involves new terms corresponding to transitions among the two species in the Bose-Hubbard model. We propose an optical lattice implementations involving a ladder structure.

PACS numbers: 03.67.Ac,05.10.Cc,11.10.Hi,11.15.Ha,37.10.Jk,64.60.De,67.85.-d

I. INTRODUCTION

The lattice formulation of quantum chromodynamics has provided successful treatments of nonperturbative problems involving strongly interacting particles in the context of particle and nuclear physics. However, numerical computations at finite density and real time have remained a major challenge. For this reason, there has been a lot of interest in the possibility of building quantum simulators for lattice gauge theory (LGT) using optical lattices (for recent reviews see Refs. [1, 2]).

When physical degrees of freedom - individual cold atoms or condensates trapped in the optical lattice - play the role of the gauge fields, one needs to make sure that the physical observables are gauge invariant. One way to achieve this goal is to impose Gauss's law at least in some approximation. This question is discussed in the context of Abelian gauge theories in Refs. [3–9]. Another approach that has been advocated [10, 11] is to work directly with a gauge invariant formulation.

A simple example is the correspondence between the Fermi-Hubbard model and a $SU(2)$ gauge theory with one fermion [12]. At strong gauge coupling and small hopping parameter, the quadratic part of the gauge-invariant effective Hamiltonian in one spatial dimension reads

$$H_{eff} \propto \sum_i [M_i M_{i+1} + 2(B_i^\dagger B_{i+1} + B_{i+1}^\dagger B_i)], \quad (1)$$

where M_i and B_i are the gauge invariant operators corresponding to mesons and $SU(2)$ -baryons. A similar effective Hamiltonian can be obtained at second order in degenerate perturbation theory of the Fermi-Hubbard model with strong onsite repulsion.

Effective actions for mesons and baryons can be constructed for $SU(N)$ gauge groups with similar approximations [13]. The computation of the corrections due to

the plaquette interactions is rather involved [14]. Various related techniques have been developed to approach chiral symmetry breaking at strong coupling [15, 16] and can accommodate a chemical potential [17]. Despite this progress, testing the validity of the approximations with numerical methods has remained challenging which suggests to consider simpler examples.

For the Bose-Hubbard model with one species of particle, there is a remarkable level of *quantitative* agreement [18] between state of the art quantum Monte Carlo (MC) calculations and their experimental optical lattice implementations. It would be highly desirable to provide a similar proof of principle for a simple LGT model where calculations at finite density and real time are possible.

It is important to emphasize that to the best of our knowledge there are so far no experimental realizations of any lattice model with dynamical gauge fields. It is thus crucial to start with a simple example as a proof of principle that it is feasible.

In this article, we explain how to establish an approximate *quantitative* correspondence between the well-known Abelian Higgs model (scalar electrodynamics) [19, 20] on a 1+1 dimensional lattice and specific many-body theories that can in principle be realized experimentally on optical lattices. The Abelian Higgs model describes a charged complex scalar field and an Abelian gauge field. Its formulation is reviewed in Sec. II. The model can be studied numerically by various numerical methods in various regions of the parameter space and a first important step is to connect the standard isotropic space-time calculations to the Hamiltonian approach obtained by taking the time continuum limit first and then connecting this Hamiltonian with the Hamiltonian of cold atoms trapped in optical lattices. Our first goal is to achieve some calibration in the gapped, confining phase corresponding to the Mott insulator phase of the cold

atom system. A more ambitious goal is to extend this procedure to the region of parameter space where the gap disappears and a new phase appears.

The exact form of the gauge-invariant effective action for the mesons is derived in Sec. III by integrating over the gauge fields. In this case the constraint enforcing gauge invariance, *i.e.* Gauss's law, is satisfied automatically, while formulations with explicit gauge fields need to impose this non-trivial constraint, which is often done approximately. The absence of fermions permits inexpensive MC simulations at Euclidean time and zero chemical potential. This allows us to test the expansion of the effective action in the hopping parameter for a broad range of the inverse gauge coupling β_{pl} and small values of the scalar self-coupling λ . Good agreement with MC is shown in Sec. IV.

In Sec. V we consider the opposite limit of arbitrarily large λ where the amplitude of the scalar field is frozen to unity. In this limit, all the remaining variables involved in the basic formulation are compact and we can use recently developed Tensor Renormalization Group (TRG) methods [21–24] to write the partition function as a traced product of local tensors with discrete indices. This reformulation allows us to write exact blocking formulas which can be used for numerical purposes. The basic character expansions used in this section are well-known and have been developed in the context of dual formulations [19]. They have also been used for other purposes recently [25, 26].

Gauss's law is automatically implemented and the introduction of a chemical potential has practical consequences only if we have an external electric field, $g^2 = 0$ or an explicit gauge symmetry breaking. In the $g^2 = 0$ case, the practical implementation requires truncations but in contrast to the MC approach there is no sign problem for arbitrary values of μ . We then write the transfer matrix as a product of tensors along a time slice.

The time continuum limit of the blocked transfer matrix can be obtained numerically. In the case $g^2 = 0$, and with a spin-1 truncation (inspired by gauge magnet or gauge link constructions [27, 28]), the small volume energy spectrum is identical to the low energy spectrum of a two-species Bose-Hubbard model in the limit of large onsite repulsion [29, 30]. We calculate numerically the energy spectrum in this limit in Sec. VI. This section reviews the connection between the Abelian Higgs and $O(2)$ models and connects with our previous Hamiltonian construction for the $O(2)$ case, which is an important prerequisite for understanding the material in Sec. VIII.

In Sec. VII, we start with finite values of β_{pl} (non-zero values of g^2) and consider the time continuum limit obtained by extrapolating to infinite β_{pl} while keeping a fixed value for the mass gap. Using the spin-1 approximation we obtain a Hamiltonian which is quite different from the one obtained in Sec. VI where we started with the model at $g^2 = 0$. It involves new terms corresponding to transitions among the two species in the Bose-Hubbard model which we discuss in Sec. VIII. We

then propose an optical lattice implementation involving a ladder structure and discuss further plans to obtain a good correspondence between conventional MC calculations and optical lattice measurements.

II. THE ABELIAN HIGGS MODEL

In this section, we briefly remind the reader of the action for the Abelian Higgs model on a 1+1 space-time lattice of size $N_s \times N_\tau$ and introduce the notations used later. We use x, y etc. for space-time vectors, i, j etc. for the one-dimensional spatial sites and $\hat{\nu} = \hat{s}$ or $\hat{\tau}$ for the unit vectors in space and time, respectively. The gauge fields $U_{x,\hat{\nu}} = \exp(iA_{x,\hat{\nu}})$ are attached to the links. We denote the product of U 's around a plaquette $U_{pl,x}$ where x is the lower left corner of the plaquette in space-time coordinates. We use the notation $\beta_{pl} = 1/g^2$ for the inverse gauge coupling and κ_s (κ_τ) for the hopping coefficient in the space (time) direction. For the potential for the complex scalar field $\phi_x = |\phi_x| \exp(i\theta_x)$, we follow the convention of Ref. [31]. The action reads:

$$S = S_g + S_h + S_\lambda, \quad (2)$$

where the gauge part is

$$S_g = -\beta_{pl} \sum_x \text{Re}[U_{pl,x}], \quad (3)$$

the hopping

$$S_h = -\kappa_\tau \sum_x \left[e^\mu \phi_x^\dagger U_{x,\hat{\tau}} \phi_{x+\hat{\tau}} + e^{-\mu} \phi_{x+\hat{\tau}}^\dagger U_{x,\hat{\tau}}^\dagger \phi_x \right] - \kappa_s \sum_x \left[\phi_x^\dagger U_{x,\hat{s}} \phi_{x+\hat{s}} + \phi_{x+\hat{s}}^\dagger U_{x,\hat{s}}^\dagger \phi_x \right] \quad (4)$$

and the self-interaction

$$S_\lambda = \lambda \sum_x (\phi_x^\dagger \phi_x - 1)^2 + \sum_x \phi_x^\dagger \phi_x. \quad (5)$$

The partition function can then be written as

$$Z = \int D\phi^\dagger D\phi DU e^{-S}. \quad (6)$$

In the hopping part of the action S_h , we can separate the compact and non-compact variables

$$S_h = -2\kappa_\tau |\phi_x| |\phi_{x+\hat{\tau}}| \sum_x \cos(\theta_{x+\hat{\tau}} - \theta_x + A_{x,\hat{\tau}} - i\mu) - 2\kappa_s |\phi_x| |\phi_{x+\hat{s}}| \sum_x \cos(\theta_{x+\hat{s}} - \theta_x + A_{x,\hat{s}}). \quad (7)$$

This equation makes clear that the chemical potential is a constant imaginary gauge field in the time direction and that the Nambu-Goldstone fields θ_x can be eliminated by a gauge transformation

$$A_{x,\hat{\nu}} \rightarrow A_{x,\hat{\nu}} - \theta_{x+\hat{\nu}} + \theta_x, \quad (8)$$

which leaves the plaquette terms unchanged.

III. A GAUGE-INVARIANT EFFECTIVE ACTION

As explained in the introduction, unlike other approaches [2, 4–7, 9, 32–34] we will not try to implement the gauge field on the optical lattice, but rather try to implement a gauge-invariant effective action obtained by integrating over the gauge fields. In this section, we will show that this effective action is a function of the composite, gauge-invariant, meson field which we denote $M_x \equiv \phi_x^\dagger \phi_x$ in order to emphasize the analogy with Eq. (1). In other words,

$$Z = \int D\phi^\dagger D\phi DU e^{-S} = \int DM e^{-S_{eff}(M) - S_\lambda(M)}. \quad (9)$$

For this purpose, we use the Fourier expansion of the Boltzmann weights in terms of the modified Bessel functions I_n , for instance,

$$\begin{aligned} & \exp[2\kappa_\tau |\phi_x| |\phi_{x+\hat{\tau}}| \cos(\theta_{x+\hat{\tau}} - \theta_x + A_{x,\hat{\tau}} - i\mu)] \\ &= \sum_{n=-\infty}^{\infty} I_n(2\kappa_\tau |\phi_x| |\phi_{x+\hat{\tau}}|) \exp[in(\theta_{x+\hat{\tau}} - \theta_x + A_{x,\hat{\tau}} - i\mu)], \end{aligned} \quad (10)$$

and similar expressions for the space hopping and the plaquette interactions. We can then collect all the exponentials involving a given $A_{x,\hat{\nu}}$ and perform the integration over $A_{x,\hat{\nu}}$. This results in Kronecker deltas relating the various Fourier modes. The final result is

$$\begin{aligned} e^{-S_{eff}} = \sum_{\{m_\square\}} & \left[\prod_{\square} I_{m_\square}(\beta_{pl}) \prod_x \left(I_{n_{x,\hat{s}}} (2\kappa_s |\phi_x| |\phi_{x+\hat{s}}|) \right. \right. \\ & \left. \left. \times I_{n_{x,\hat{\tau}}} (2\kappa_\tau |\phi_x| |\phi_{x+\hat{\tau}}|) \exp(\mu n_{x,\hat{\tau}}) \right) \right], \end{aligned} \quad (11)$$

with specific rules to express the link indices $n_{x,\hat{\nu}}$ in terms of the plaquette indices m_\square that we now proceed to explain. Given that for x real, $I_n(x) = I_{-n}(x)$ there are several equivalent ways to label the contributions. We use the convention where time is along the vertical axis. Starting from the lower left corner of the plaquette and moving counterclockwise, the gauge fields in the exponentials come with a plus sign for the first two links and a minus sign for the last two links. For the hopping terms, the gauge fields always come with a positive sign. This results in the rules

$$n_{x,\hat{s}} = m_{below} - m_{above} \quad (12)$$

$$n_{x,\hat{\tau}} = m_{right} - m_{left} \quad (13)$$

where the subscripts such as “below” refers to the plaquette location with respect to the link. This completely fixes the link indices in terms of the plaquette indices and it is easy to check that Eqs. (12-13) guarantee that the link indices automatically satisfy the current conservation imposed by the integration of the θ_x variables. In other words, the m_\square are the dual variables [19].

Models with local $U(1)$ symmetry, in particular, the Abelian Higgs model have been extensively studied in

Ref. [19] in the dual formulation in d dimensions. While the focus of this paper is on 1+1 dimensions, we note that duality transformations, based on (10), allow one to construct a gauge theory of integers in 2+1 dimensions and a Kalb-Ramond type theory [35] in 3+1 dimensions, which should be considered as candidates for quantum simulating higher dimensional models on optical lattices.

Equations (12-13) have simple electromagnetic analogs. First, $n_{x,\hat{\tau}}$ can be interpreted as a charge and m_x as an electric field in the spatial direction. With this Minkowskian interpretation, Eq. (13) enforces Gauss’s law. Second, $n_{x,\hat{\nu}}$ can be interpreted as a two-dimensional current and m_x as a magnetic field normal to the two-dimensional plane. In this Euclidean interpretation, Eqs. (12-13) express the current as the curl of the magnetic field. A right hand rule can be obtained for the following index ordering: time, space, normal direction. A discrete version of Gauss’s theorem guarantees that the sum of the charges on a time slice is the last m on the right minus the first m on the left.

At the lowest order of the strong-coupling expansion we have $\beta_{pl} = 0$ and from $I_n(0) = 0$ for $n \neq 0$, we see that all the indices must be zeros. The effect of the plaquette can be restored perturbatively. This can be organized in an expansion in the hopping parameter. In the isotropic case $\kappa_\tau = \kappa_s = \kappa$ we obtain:

$$\begin{aligned} S_{eff} = \sum_{\langle xy \rangle} & \left(-\kappa^2 M_x M_y + \frac{1}{4} \kappa^4 (M_x M_y)^2 \right) \\ & - 2\kappa^4 \frac{I_1(\beta_{pl})}{I_0(\beta_{pl})} \sum_{\square(xy zw)} M_x M_y M_z M_w + O(\kappa^6) \end{aligned} \quad (14)$$

IV. MONTE CARLO CALCULATIONS

Consider the action (2) for the isotropic case $\kappa_\tau = \kappa_s = \kappa$. We start with the $\beta_{pl} \rightarrow \infty$ limit when all gauge variables are frozen to unity, $U_{x,\hat{\nu}} = 1$. In this limit the expectation value of the hopping term

$$L_\phi = \langle \text{Re} \{ \phi_x^\dagger U_{x,\hat{\nu}} \phi_{x+\hat{\nu}} \} \rangle \quad (15)$$

can be calculated with the hopping parameter expansion, for small κ . It has been derived up to $O(\kappa^5)$ in Ref. [31]. This result can be generalized to $\beta_{pl} < \infty$ by including the appropriate factors of $I_1(\beta_{pl})/I_0(\beta_{pl})$ for the diagrams that involve plaquettes [36].

To check the range of validity of the expansion we perform Monte Carlo simulations at several values of β_{pl} , κ and λ on a 16^2 lattice. To test the $\beta_{pl} \rightarrow \infty$ limit we set $\beta_{pl} = 20$ and for $\lambda = 0.05$ and 0.1 scan the range of $\kappa \in [0.05, 0.30]$. The results for L_ϕ are shown in Fig. 1. The lines represent the expansion at two orders. The expansion starts to break down around $\kappa = 0.15$ at $O(\kappa^3)$ and $\kappa = 0.2$ at $O(\kappa^5)$. At the present, exploratory, stage we use the updating algorithm of Ref. [37] that is applicable at small self-coupling λ .




FIG. 1. L_ϕ at $\beta_{pl} = 20$ for $\lambda = 0.05$ and $\lambda = 0.1$ as function of κ compared with the hopping expansion at $\beta_{pl} = \infty$ at $O(\kappa^3)$ and $O(\kappa^5)$.




FIG. 2. L_ϕ at $\beta_{pl} = 20, 2, 0.2$ and 0.02 for $\lambda = 0.1$ as function of κ compared with the hopping expansion with included dependence on β_{pl} up to $O(\kappa^5)$.

To study the dependence on β_{pl} we focus on the $\kappa \in [0.08, 0.16]$ range at $\lambda = 0.1$ and perform calculations at $\beta_{pl} = 20, 2, 0.2$ and 0.02 . The results for L_ϕ are shown in Fig. 2.

To understand the dependence on β_{pl} better we also calculate L_ϕ for several values of β_{pl} at fixed $\kappa = 0.15$. The results together with the hopping expansion are shown in Fig. 3. There is good agreement, and the dependence on β is weak, compare the scales on Fig. 2 and 3.

At intermediate and large values of the self-coupling λ one cannot consider the quartic term as a perturbation and the updating algorithm of Ref. [37] suffers from low acceptance rate. We are currently implementing the Biased-Metropolis Heatbath algorithm [38] that can provide almost uniform acceptance for arbitrary values of λ . This is achieved by tabulating the proposal probability distribution function. This algorithm has been used for compact variables, such as the ones in $U(1)$ and $SU(2)$




FIG. 3. L_ϕ at fixed $\kappa = 0.15$ as function of β_{pl} for $\lambda = 0.1$ compared with the hopping expansion with included dependence on β_{pl} up to $O(\kappa^5)$.

gauge theory, but has not yet been applied for a non-compact case, such as the scalar field amplitude.

V. THE LARGE λ LIMIT

We now turn to the limit where λ becomes arbitrarily large. In this limit, M_x is frozen to 1, or in other words, the Brout-Englert-Higgs mode becomes infinitely massive. We are then left with compact variables of integration in the original formulation (θ_x and $A_{x,\hat{\nu}}$) and the Fourier expansions described before lead to expressions of the partition function in terms of discrete sums.

As explained in Ref. [22], these sums can be formulated in a compact way using tensorial notations. In order to simplify the equations below we define

$$t_n(z) = \frac{I_n(z)}{I_0(z)}. \quad (16)$$

These normalized Bessel functions have useful properties. For instance $t_n(0) = \delta_{n,0}$. For z non-zero and finite, we have

$$1 = t_0(z) > t_1(z) > t_2(z) > \dots > 0. \quad (17)$$

In addition, for sufficiently large z ,

$$t_n(z) \approx 1 - \frac{n^2}{2z}. \quad (18)$$

Following the general principles of the construction [22], we attach a $B^{(\square)}$ tensor to every plaquette

$$B_{m_1 m_2 m_3 m_4}^{(\square)} = \begin{cases} t_{m_\square}(\beta_{pl}), & \text{if } m_1 = m_2 = m_3 = m_4 = m_\square \\ 0, & \text{otherwise,} \end{cases} \quad (19)$$

a $A^{(s)}$ tensor to the horizontal links

$$A_{m_{above} m_{below}}^{(s)} = t_{|m_{below} - m_{above}|}(2\kappa_s), \quad (20)$$




FIG. 4. The basic B and A tensors (in brown and green, respectively, colors online). The $A^{(s)}$ are associated with the vertical tensors, and the horizontal (spatial) links of the lattice. The $A^{(\tau)}$ are associated with the horizontal tensors, and the vertical (temporal) links of the lattice.

and a $A^{(\tau)}$ tensor to the vertical links

$$A_{m_{left}m_{right}}^{(\tau)} = t_{|m_{left}-m_{right}|}(2\kappa_\tau) e^{(m_{right}-m_{left})\mu}. \quad (21)$$

The partition function can now be written as

$$Z = (I_0(\beta_{pl})I_0(2\kappa_s)I_0(2\kappa_\tau))^V \times \text{Tr} \left[\prod_{s,\tau,\square} A_{m_{above}m_{below}}^{(s)} A_{m_{left}m_{right}}^{(\tau)} B_{m_1m_2m_3m_4}^{(\square)} \right]. \quad (22)$$

The traces are performed by contracting the vertical and horizontal indices as shown in Fig. 4. Note that the tensor $A^{(s)}$ associated with a horizontal space link is represented by a vertical line orthogonal to the link and the tensor $A^{(\tau)}$ associated with a vertical time link is represented by a horizontal line orthogonal to the link.

The traces can also be expressed in terms of a transfer matrix \mathbb{T} which can be constructed in the following way. First, we define a matrix \mathbb{B} as the product of the plaquette tensors on a time slice alternating with the link tensor corresponding to the vertical links in between the plaquettes. There are two natural ways to impose boundary conditions. The first is to connect the last A tensor with the first B tensor (periodic boundary conditions), the second is to impose $m = 0$ for the first and last B tensor (open boundary conditions). However, in both cases, the total charge on the interval has to be zero. A more general option is to allow arbitrary m 's at each end. This would allow us to consider appropriately selected charge sectors. In the following we will focus on the open boundary conditions ($m = 0$ at both ends) and

define

$$\mathbb{B}_{(m_1,m_2,\dots,m_{N_s})(m'_1,m'_2,\dots,m'_{N_s})} = t_{m_1}(2\kappa_\tau)\delta_{m_1,m'_1}t_{m_1}(\beta_{pl}) \times t_{|m_1-m_2|}(2\kappa_\tau)\delta_{m_2,m'_2}t_{m_2}(\beta_{pl})t_{|m_2-m_3|}(2\kappa_\tau) \dots t_{m_{N_s}}(\beta_{pl})t_{m_{N_s}}(2\kappa_\tau). \quad (23)$$

Note that with this choice of open boundary conditions, the chemical potential has completely disappeared. If we had chosen different m 's at the end allowing a total charge Q inside the interval, we would have an additional factor $\exp(\mu Q)$. Another way to get a nontrivial factor involving μ is to introduce an explicit symmetry breaking term of the form, for instance, $b_{\hat{\nu}}(U_{x,\hat{\nu}} + U_{x,\hat{\nu}}^\dagger)$. This brings new independent quantum numbers on the links which introduce local violations of Gauss's law. It may also be argued that if g^2 is exactly zero, there is no electric field and no special reason to impose $m = 0$ at the boundary (see Sec. VI).

We next define a matrix \mathbb{A} as the product

$$\mathbb{A}_{(m_1,m_2,\dots,m_{N_s})(m'_1,m'_2,\dots,m'_{N_s})} = t_{|m_1-m'_1|}(2\kappa_s)t_{|m_2-m'_2|}(2\kappa_s) \dots t_{|m_{N_s}-m'_{N_s}|}(2\kappa_s). \quad (24)$$

With these notations we can construct a symmetric transfer matrix \mathbb{T} . Since \mathbb{B} is diagonal, real and positive, we can define its square root in an obvious way and write the transfer matrix as

$$\mathbb{T} = \sqrt{\mathbb{B}}\mathbb{A}\sqrt{\mathbb{B}}. \quad (25)$$

With this definition, the partition function can be written as

$$Z = (I_0(\beta_{pl})I_0(2\kappa_s)I_0(2\kappa_\tau))^V \text{Tr} [\mathbb{T}^{N_\tau}]. \quad (26)$$

Alternatively, we could diagonalize the symmetric matrix \mathbb{A} and define the (dual) transfer matrix

$$\tilde{\mathbb{T}} = \sqrt{\mathbb{A}}\mathbb{B}\sqrt{\mathbb{A}}. \quad (27)$$

The \mathbb{A} and \mathbb{B} matrices can be constructed by a recursive blocking method similar to those discussed in Ref. [22]. We can construct a new B tensor, called B' , by contracting two B tensors on both sides of a $A^{(\tau)}$ tensor. This process can be iterated and is illustrated in Fig. 5. Using tensorial notation we write

$$B'_{m_3m_6M(m_1,m_2)M'(m'_1,m'_2)} = \sum_{m_4,m_5} B_{m_3m_4m_1m'_1}A_{m_4m_5}^{(\tau)}B_{m_5m_6m_2m'_2}, \quad (28)$$

where the notation $M(m_1, m_2)$ stands for the product state, $M = m_1 \otimes m_2$. One can continue blocking horizontally as above until the desired spatial size is achieved resulting in the matrix \mathbb{B} . By their very nature the B s enforce that this object is diagonal in the collective product-state indices of the upper and lower tensor legs. For practical reasons, truncation methods need to be introduced at the beginning and after each blocking [21, 39].




FIG. 5. Part of the construction of the blocked B' tensor. This shows the contraction of the B and $A^{(\tau)}$ tensors. The dashed lines are the links of the original lattice.




FIG. 6. Graphical representation of the blocking of the A tensors. The vertical tensors are the $A^{(s)}$ and the dashed lines are the links of the original lattice.

Now consider the $A^{(s)}$ matrices which correspond to the horizontal links of the original lattice as in Fig. 6. If one takes their outer product and collects the upper indices into a single product-state index, and does the same for the lower indices, one has another matrix built out of the vertical $A^{(s)}$ tensors. Once again in tensorial notation this can be written as

$$A'_{M(m_1, m_2)M'(m'_1, m'_2)}^{(s)} = A_{m_1 m'_1}^{(s)} A_{m_2 m'_2}^{(s)}. \quad (29)$$

One can continue taking the product of $A^{(s)}$ matrices until the desired spatial size has been reached resulting in the matrix \mathbb{A} .

VI. THE CASE $g^2 = 0$

In this section, we discuss the case where both λ is infinite and g^2 is exactly zero, *i.e.* $\beta_{pl} = \infty$. This limit corresponds to the two-dimensional classical $O(2)$ model, and it has been discussed in the early studies of the phase structure and topological excitations of the Abelian Higgs model, for instance, in Ref. [40, 41].

As explained in Ref. [42–44], the time continuum limit can be achieved by taking the limit $\kappa_\tau \rightarrow \infty$ while keeping the product $\kappa_\tau \kappa_s$ constant. This leads to a Hamiltonian for quantum rotors located at each spatial site and having quantized angular momentum running over positive and negative integers. In order to realize this Hamiltonian on optical lattices, we considered [29] a spin-1 truncation where the angular momentum at each site is restricted to the values 0 and ± 1 . For this model such

truncation is similar in spirit to the dilute-gas approximation [45] and has very small effects on the phase diagram provided that the hopping parameter and the chemical potential are not too large.

Following Refs. [29, 30] with the replacements $\beta_\nu \rightarrow 2\kappa_\nu$, the Hamiltonian for the spin-1 approximation of the $O(2)$ model reads:

$$\hat{H} = \frac{\tilde{U}}{2} \sum_i \left(\hat{L}_{(i)}^z \right)^2 - \tilde{\mu} \sum_i \hat{L}_{(i)}^z - \frac{\tilde{J}}{4} \sum_i \left(\hat{L}_{(i)}^+ \hat{L}_{(i+1)}^- + \hat{L}_{(i)}^- \hat{L}_{(i+1)}^+ \right), \quad (30)$$

with the now dimensionful quantities

$$\tilde{U} \equiv \frac{1}{2\kappa_\tau a}, \quad \tilde{\mu} \equiv \frac{\mu}{a}, \quad \tilde{J} \equiv \frac{2\kappa_s}{a}, \quad (31)$$

and a the time lattice spacing. If we use open boundary conditions, the sums run over all spatial sites in the first two sums and all but the last in the third sum. This approximate Hamiltonian can be matched [29, 30] with the effective Hamiltonian of a Bose-Hubbard model discussed in sec. VIII, that, we expect, can be implemented on optical lattices.

The method discussed in Sec. V allows us to match the spectra at small volume for the classical lattice model and the two-species Bose-Hubbard model, derived to match the Hamiltonian (30) in Ref. [29] with very good accuracy. This section is a warm-up for the more complicated situations discussed in Secs. VII and VIII.

The first step is to understand the matching in the simple situation where \tilde{U} is large compared to $\tilde{\mu}$ and \tilde{J} . In this case, \tilde{U} sets the scale of the mass gap and $\tilde{\mu}$ and \tilde{J} introduce small splittings. We now use energy units where $\tilde{U}=1$. The time continuum limit can be obtained numerically as we keep blocking the transfer matrix in the space direction as discussed in Sec. V. The spectra in blocks of size 2, 4 and 8 in the spin-1 approximation are shown in Fig. 7 for $\tilde{J}=0.1$, $\tilde{\mu} = 0$. A nonzero chemical potential would split the charge conjugated states and make the graph difficult to read. An example will be shown for $L = 2$ in Sec. VIII.

VII. THE TIME CONTINUUM LIMIT AND THE ENERGY SPECTRUM

In this section, we construct the time continuum limit of the transfer matrix \mathbb{T} defined by Eq. (25) starting now with the more general situation where we have β_{pl} finite and increase its value keeping the open boundary conditions ($m = 0$ at both ends). In order to obtain Hamiltonians corresponding to Bose-Hubbard models, a guiding strategy is to assume that the spatial hopping κ_s is proportional to some hopping energy. In the limit $\kappa_s = 0$, the inside matrix \mathbb{A} becomes the identity and the transfer matrix $\mathbb{T} = \sqrt{\mathbb{B}}\mathbb{A}\sqrt{\mathbb{B}}$ becomes \mathbb{B} which is




FIG. 7. $O(2)$ spectra in \tilde{U} units for $L=2, 4$, and 8 , with $\tilde{J}=0.1$, $\tilde{\mu} = 0$. Some higher energy states are not shown on the figure.

diagonal. In this limit, the only way to obtain a time continuum limit, in other words to have \mathbb{T} close to the identity, is to require that *both* κ_τ and β_{pl} become large. At leading order in the inverse of these large parameters, the eigenvalues of \mathbb{T} are

$$\lambda_{(m_1, m_2, \dots, m_{N_s})} = \quad (32)$$

$$1 - \frac{1}{2} \left[\left(\frac{1}{\beta_{pl}} (m_1^2 + m_2^2 + \dots + m_{N_s}^2) + \frac{1}{2\kappa_\tau} (m_1^2 + (m_2 - m_1)^2 + \dots \dots + (m_{N_s} - m_{N_s-1})^2 + m_{N_s}^2) \right) \right]$$

There are two limiting situations: $1 \ll \beta_{pl} \ll \kappa_\tau$ and $1 \ll \kappa_\tau \ll \beta_{pl}$. In the first case, $1/\beta_{pl}$ is the largest coefficient and the low energy states are those with a few m_i nonzero and consequently only a few differences of m 's are nonzero and the second term in Eq. (32) is a perturbation. In the second case, $1/\kappa_\tau$ is the largest coefficient and we could guess that to minimize its contribution, we need to take most of the m 's to be equal and just create a few charges. For a large volume, this can cause the first term of Eq. (32) to dominate, which signals an infrared instability. In electrostatic terms, creating one charge also creates a constant electric field over the entire volume. The linear potential is responsible for the confinement and trying to treat the interaction perturbatively results in infrared problems.

For this reason, in the following, we will only consider the case $1 \ll \beta_{pl} \ll \kappa_\tau$ and set the scale with the (large) gap energy

$$\tilde{U}_P \equiv \frac{1}{a\beta_{pl}}. \quad (33)$$

It is important to distinguish this scale from the \tilde{U} introduced in Sec. VI. \tilde{U}_P is associated with the plaquette

quantum number m (the energy to create a flux tube across one lattice spacing between the opposite charges), while \tilde{U} in Sec. VI is associated with the quantum number n (the energy to create a single charge). In addition, we define the (small) energy scales

$$\tilde{Y} \equiv \frac{1}{2\kappa_\tau a} = \frac{\beta_{pl}}{2\kappa_\tau} \tilde{U}_P, \quad (34)$$

which plays the same role as \tilde{U} in Sec. VI and the space hopping parameter

$$\tilde{X} \equiv \sqrt{2}\beta_{pl}\kappa_s\tilde{U}_P. \quad (35)$$

We are now in position to derive an expression for the Hamiltonian in the spin-1 approximation where the plaquette quantum number m takes values ± 1 and 0 only. The effect of κ_s can be studied by linearization. In the case of two spins, we find that

$$\partial\mathbb{T}/\partial\kappa_s|_{\kappa_s=0} = \sqrt{2}(\bar{L}_{(1)}^x + \bar{L}_{(2)}^x) \quad (36)$$

We use the notation $\bar{L}_{(1)}^x$ to denote the first generator of the spin-1 rotation algebra at the site (1). The notation \bar{L} is used to emphasize that the spin is related to the m quantum numbers attached to the plaquettes in contrast to the spin-1 generators \hat{L} in Sec. VI having a spin related to the charges n attached to the time links. The final form of the Hamiltonian \bar{H} for $1 \ll \beta_{pl} \ll \kappa_\tau$ is

$$\bar{H} = \frac{\tilde{U}_P}{2} \sum_i (\bar{L}_{(i)}^z)^2 + \quad (37)$$

$$\frac{\tilde{Y}}{2} \sum_i' (\bar{L}_{(i)}^z - \bar{L}_{(i+1)}^z)^2 - \tilde{X} \sum_i \bar{L}_{(i)}^x,$$

where \sum_i' is a short notation to include the single terms at the two ends as in Eq. (32), *i.e.* besides $(\bar{L}_{(1)}^z - \bar{L}_{(2)}^z)^2$, $(\bar{L}_{(2)}^z - \bar{L}_{(3)}^z)^2, \dots, (\bar{L}_{(N_s-1)}^z - \bar{L}_{(N_s)}^z)^2$ terms this sum contains $(\bar{L}_{(1)}^z)^2$ and $(\bar{L}_{(N_s)}^z)^2$.

The process outlined in this paper can in principle be generalized for higher spatial dimensions and non-Abelian gauge groups. Tensor formulations and exact blocking procedures have already been worked out in Ref. [22].

VIII. A TWO-SPECIES BOSE-HUBBARD MODEL IMPLEMENTATION

In Refs. [29, 30], we have proposed to match the Hamiltonian of the $O(2)$ model given in Eq. (30) with the two-species Bose-Hubbard Hamiltonian on a linear optical lattice

$$\mathcal{H} = - \sum_{\langle ij \rangle} (t_a a_i^\dagger a_j + t_b b_i^\dagger b_j + h.c.) - \sum_{i,\alpha} (\mu_{a+b} + \Delta_\alpha) n_i^\alpha$$

$$+ \sum_{i,\alpha} \frac{U_\alpha}{2} n_i^\alpha (n_i^\alpha - 1) + W \sum_i n_i^a n_i^b + \sum_{\langle ij \rangle \alpha} V_\alpha n_i^\alpha n_j^\alpha \quad (38)$$

with $\alpha = a, b$ indicating two different species and with $n_i^a = a_i^\dagger a_i$ and $n_i^b = b_i^\dagger b_i$. In this expression, the chemical potential μ_{a+b} is associated with the conservation of $n^a + n^b$ and should not be confused with the chemical potential introduced in the previous section which couples to $n^a - n^b$ and breaks the charge conjugation symmetry. In the limit where $U_a = U_b = U$ and W and $\mu_{a+b} = (3/2)U$ are much larger than any other energy scale, we have the condition $n_i^a + n_i^b = 2$ for the low energy sector. The three states $|2, 0\rangle$, $|1, 1\rangle$ and $|0, 2\rangle$ satisfy this condition and correspond to the three states of the spin-1 projection considered above.

Using degenerate perturbation theory [46, 47], we found [29, 30] that

$$\begin{aligned} \mathcal{H}_{eff} = & \left(\frac{V_a}{2} - \frac{t_a^2}{U} + \frac{V_b}{2} - \frac{t_b^2}{U} \right) \sum_{\langle ij \rangle} L_i^z L_j^z \\ & + \frac{-t_a t_b}{U} \sum_{\langle ij \rangle} (L_i^+ L_j^- + L_i^- L_j^+) + (U - W) \sum_i (L_i^z)^2 \\ & + \left[\left(\frac{pn}{2} V_a + \Delta_a - \frac{p(n+1)t_a^2}{U} \right) \right. \\ & \left. - \left(\frac{pn}{2} V_b + \Delta_b - \frac{p(n+1)t_b^2}{U} \right) \right] \sum_i L_i^z, \quad (39) \end{aligned}$$

where p is the number of neighbors and n is the occupation ($p = 2$, $n = 2$ in the case under consideration). \hat{L} is the angular momentum operator in the representation $n/2$. The effective Hamiltonian (39) without the last term has been studied in [48] and shows a rich phase diagram.

In order to match the Hamiltonian of Eq. (39) with the spin-1 approximation of the $O(2)$ model, Eq. (30), we need to tune the hopping amplitudes as $t_\alpha = \sqrt{V_\alpha U/2}$, obtaining the correspondence $\tilde{J} = 4\sqrt{V_a V_b}$, $\tilde{U} = 2(U - W)$, and $\tilde{\mu} = -(\Delta_a - V_a) + (\Delta_b - V_b)$. Fig. 8 illustrates the matching of the energy spectrum for $\tilde{J}/\tilde{U} = 0.1$ and large U , for small systems (two sites and four sites). In the limit $U \rightarrow \infty$, the matching is excellent. As we lower U , high energy states get a lower energy and can mix with the low energy states. These states may be related to the modes that become infinitely massive when $\lambda \rightarrow \infty$. In 2+1 dimensions, Higgs modes have been identified numerically [49] and experimentally [50] for the one species Hubbard model.

The optical lattice implementation of the two-species Bose-Hubbard Hamiltonian is discussed in Ref. [29] for a ^{87}Rb and ^{41}K Bose-Bose mixture. The on-site interactions (U_α , W) and the hopping amplitudes (t_α) can be tuned to some extent by the parameters of the optical lattice [51, 52]. In addition, an interspecies Feshbach resonance is accessible for this mixture and therefore the interspecies interaction (W) can be further controlled by an external magnetic field. The extended interaction, V_α , is present and small when we consider Wannier gaussian wave functions sitting on nearby lattice sites [53]. Achieving the required extended interaction may be a challeng-




FIG. 8. $O(2)$ with $\tilde{J}/\tilde{U} = 0.1$ and Bose-Hubbard spectra for $L = 2$ with $\tilde{\mu}/\tilde{U} = 0.02$ (top) and $L = 4$ with $\tilde{\mu} = 0$ (bottom).

ing part of the experimental proposal. The strong coupling limit is needed for our perturbation theory treatment to apply, but it is also important to have U significantly larger than the temperature. For the ^{87}Rb - ^{41}K mixture, temperature and recoil energies of the order of 100 nK have been achieved, and values of U that are 10-20 times larger can be reached [52, 54, 55].

For the Hamiltonian \tilde{H} in Eq. (37) corresponding to $1 \ll \beta_{pl} \ll \kappa_\tau$, we need to introduce a new interaction to represent the \tilde{L}^x effects that interchange the $m = 0$ states with the $m = \pm 1$ states. This can be achieved by adding the piece

$$\Delta H = -\frac{t_{ab}}{2} \sum_i (a_i^\dagger b_i + b_i^\dagger a_i). \quad (40)$$

The matching between the two models can be achieved by imposing $t = 0$, $V_a = V_b = -\tilde{Y}/2$ and $t_{ab} = \tilde{X}$. Energy levels obtained numerically with this method have a good matching shown in Fig. 9. Note that if an interspecies nearest neighbor interaction V_{ab} is introduced, it can be incorporated in \mathcal{H}_{eff} by making the substitution $V_a + V_b \rightarrow V_a + V_b - 2V_{ab}$ in the first term of Eq. (39).

This is a very different realization than for the $O(2)$ limit. The presence in the Hamiltonian of the additional term that interchanges the species index, Eq. (40), rules out implementing this Hamiltonian using mixtures of two




FIG. 9. Abelian-Higgs model with $\tilde{X}/\tilde{U}_P = 0.1$, $\tilde{Y}/\tilde{U}_P = 0.1$ and the corresponding Bose-Hubbard spectra for $L = 2$ (top) and $L = 4$ (bottom).

different types of atoms. It could be realized with a single atomic species on a ladder structure with a and b corresponding to the two legs of the ladder. Ladder systems have been realized experimentally by using lattices of double wells [56–60]. The hopping amplitudes can be tuned such that the hopping in the direction along the ladder is negligible, but finite along the rungs, thus exchanging a , b species index at the same rung. An attractive intraspecies interaction ($V_a = V_b = -\tilde{Y}/2$) is also needed, favoring having two atoms in neighboring sides on the same leg of the ladder. This acts as a nearest-neighbor ferromagnetic coupling in the effective one-dimensional spin chain Hamiltonian. For the experimental implementation, an attractive nearest neighbor interaction can be obtained by using cold dipolar atoms or molecules, with dipole moments aligned along the ladder and with inter-rung distance such that the rapidly decaying dipole-dipole interaction between next-nearest-neighbors can be neglected. With this alignment of the moments, the inter-species interaction in the same rung (W) is repulsive. Experiments with ultra cold dipolar quantum gases have been performed with chromium [61], erbium [62, 63], and dysprosium [64–66], which have magnetic moments $6\mu_B$, $7\mu_B$ and $10\mu_B$, respectively, and with polar molecules, such as ^{40}K ^{87}Rb and ^{23}Na ^{40}K [67–

69]. Alternatively, the two boson species in the Bose-Hubbard Hamiltonian could correspond to two hyperfine states of the same atom species, provided these two states support Raman transitions between them so that the species conversion term, Eq. (40), can be created. These transitions have recently been exploited for the creation of synthetic dimensions [70, 71]. Another possible avenue to be explored would be to implement the spin-1 approximation directly on one-dimensional optical lattices by combining Raman lasers and atoms with magnetic sublevels. A possibility to obtain a pseudospin 1 is to use the $F = 1$ hyperfine splitting of ^{87}Rb [71].

IX. CONCLUSIONS

With the remarkable experimental progress in the field of Atomic, Molecular and Optical physics, particularly the loading of trapped ultra-cold atoms onto optical lattices, and the unprecedented levels of control and tunability of their interactions, a wide range of quantum many-body lattice Hamiltonians can be realized and studied.

We have proposed a two-species Bose-Hubbard model that may be used as a quantum simulator for lattice gauge theories. The gauge invariance is built-in and thus does not need to be achieved via fine-tuning, and the correspondence between the proposed Bose-Hubbard Hamiltonian and the Abelian Higgs model can be checked quantitatively.

On the experimental side, various multi-species Bose-Hubbard systems can be created with current cold atom technology. Various possible realizations for the two species that are needed can be explored. On the theory side, the Bose-Hubbard Hamiltonian is amenable to Quantum Monte Carlo calculations and the Abelian Higgs model can be treated with the TRG (and the time-continuum limit obtained) as well as Monte Carlo calculations.

For reasons explained in Ref. [49], the Higgs mode is usually difficult to observe numerically or experimentally. However, for the 2+1 dimensional Bose-Hubbard model, numerical calculations of the scalar susceptibility near the tip of a Mott insulator lobe where the relativistic particle-hole symmetry is restored, show a peak narrowing as one approaches the quantum critical point [49]. Almost simultaneously, an experimental implementation of the model on an optical lattice [50] made possible to observe the corresponding temperature response to a periodic modulation of the lattice depth. However, in 1+1 there are so far no conclusive results. Our setup could contribute to clarify this point in the future.

Another interesting feature of the 1+1 dimensional Abelian Higgs model is a possibility for confinement of fractional charges. The Wilson loop is often used as an observable signaling confinement and it remains to be seen what observables in the optical lattice experiments can be related to it. Following Einhorn and Savit [41], one can introduce a fractional charge by increasing the

basic charge, or, in our case, rescaling the quantum number m . This means that m has to span at least a $[-2, 2]$ range, in other words, at least a spin-2 approximation is required.

For the models we considered here the full spectra can be calculated, at least for small systems. The TRG formulation can very naturally provide time-dependent correlation functions. Therefore, besides the ultimate goal of quantum simulating lattice gauge theories using cold atoms on optical lattices, the correspondence we report here may also provide a way of studying the dynamics of cold atom systems such as response to sudden quenches.

Acknowledgments. We thank N. Gemelke, E. Mueller, Yuzhi Liu, J. Osborn, Li-Ping Yang, and Haiyuan Zou for stimulating discussions and the referee for bringing the point about fractional charges in the Abelian Higgs model to our attention. This article was completed while

Y. M. was attending the workshop INT Program INT-15-1 “Frontiers in Quantum Simulation with Cold Atoms”, he thanks the participants for many interesting discussions. The final version of the manuscript was being written while A. B. and Y. M. attended the workshop “Understanding Strongly Coupled Systems in High Energy and Condensed Matter Physics” at the Aspen Center for Physics (ACP) in May-June 2015, and S.-W. T. acknowledges helpful discussions at ACP during the workshop “Ultra-Cold Quantum Matter with Atoms and Molecules” in July 2015. This research was supported in part by the Department of Energy under Award Numbers DOE grant DE-FG02-05ER41368, DE-SC0010114 and DE-FG02-91ER40664, the NSF under grants DMR-1411345 and PHY11-25915 and by the Army Research Office of the Department of Defense under Award Number W911NF-13-1-0119.

-
- [1] U.-J. Wiese, *Annalen der Physik* **525**, 777 (2013), ISSN 1521-3889.
- [2] E. Zohar, J. I. Cirac, and B. Reznik(2015), arXiv:1503.02312 [quant-ph].
- [3] E. Kapit and E. Mueller, *Phys. Rev. A* **83**, 033625 (Mar 2011).
- [4] E. Zohar and B. Reznik, *Phys.Rev.Lett.* **107**, 275301 (2011), arXiv:1108.1562 [quant-ph].
- [5] E. Zohar, J. I. Cirac, and B. Reznik, *Phys.Rev.Lett.* **109**, 125302 (2012), arXiv:1204.6574 [quant-ph].
- [6] L. Tagliacozzo, A. Celi, A. Zamora, and M. Lewenstein, *Annals Phys.* **330**, 160 (2013), arXiv:1205.0496 [cond-mat.quant-gas].
- [7] K. Kasamatsu, I. Ichinose, and T. Matsui, *Phys.Rev.Lett.* **111**, 115303 (2013), arXiv:1212.4952 [quant-ph].
- [8] D. Banerjee, F.-J. Jiang, P. Widmer, and U.-J. Wiese, *Journal of Statistical Mechanics: Theory and Experiment* **2013**, P12010 (2013).
- [9] Y. Kuno, K. Kasamatsu, Y. Takahashi, I. Ichinose, and T. Matsui(2014), arXiv:1412.7605 [cond-mat.quant-gas].
- [10] Y. Meurice, *PoS LATTICE2011*, 040 (2011), arXiv:1202.1605 [hep-lat].
- [11] Y. Liu, Y. Meurice, and S.-W. Tsai, *PoS LATTICE2012*, 246 (2012), arXiv:1211.4126 [hep-lat].
- [12] E. Dagotto, E. Fradkin, and A. Moreo, *Phys. Rev. B* **38**, 2926 (Aug 1988).
- [13] P. Rossi and U. Wolff, *Nucl.Phys.* **B248**, 105 (1984).
- [14] P. de Forcrand, J. Langelage, O. Philipsen, and W. Unger, *PoS LATTICE2013*, 142 (2014), arXiv:1312.0589 [hep-lat].
- [15] E. Tomboulis, *Phys.Rev.* **D87**, 034513 (2013), arXiv:1211.4842 [hep-lat].
- [16] A. S. Christensen, J. C. Myers, P. D. Pedersen, and J. Rosseel(2015), arXiv:1502.04865 [hep-lat].
- [17] E. Tomboulis, *J.Math.Phys.* **54**, 122301 (2013), arXiv:1304.4678 [hep-lat].
- [18] S. Trotzky, L. Pollet, F. Gerbier, U. Schnorrberger, I. Bloch, N. V. Prokof'Ev, B. Svistunov, and M. Troyer, *Nature Physics* **6**, 998 (Dec. 2010), arXiv:0905.4882 [cond-mat.quant-gas].
- [19] R. Savit, *Rev. Mod. Phys.* **52**, 453 (Apr 1980).
- [20] D. R. T. Jones, J. Kogut, and D. K. Sinclair, *Phys. Rev. D* **19**, 1882 (Mar 1979).
- [21] Z. Y. Xie, J. Chen, M. P. Qin, J. W. Zhu, L. P. Yang, and T. Xiang, *Phys. Rev. B* **86**, 045139 (Jul 2012).
- [22] Y. Liu, Y. Meurice, M. P. Qin, J. Unmuth-Yockey, T. Xiang, Z. Y. Xie, J. F. Yu, and H. Zou, *Phys. Rev. D* **88**, 056005 (Sep 2013).
- [23] A. Denbleyker, Y. Liu, Y. Meurice, M. P. Qin, T. Xiang, Z. Y. Xie, J. F. Yu, and H. Zou, *Phys. Rev. D* **89**, 016008 (Jan 2014).
- [24] J. F. Yu, Z. Y. Xie, Y. Meurice, Y. Liu, A. Denbleyker, H. Zou, M. P. Qin, J. Chen, and T. Xiang, *Phys. Rev. E* **89**, 013308 (Jan 2014).
- [25] B. Bahr, B. Dittrich, F. Hellmann, and W. Kaminski, *Phys. Rev. D* **87**, 044048 (Feb 2013).
- [26] Y. Delgado Mercado, C. Gatteringer, and A. Schmidt, *Phys. Rev. Lett.* **111**, 141601 (Oct 2013).
- [27] P. Orland and D. Rohrlich, *Nucl.Phys.* **B338**, 647 (1990).
- [28] S. Chandrasekharan and U. Wiese, *Nucl.Phys.* **B492**, 455 (1997), arXiv:hep-lat/9609042 [hep-lat].
- [29] H. Zou, Y. Liu, C.-Y. Lai, J. Unmuth-Yockey, L.-P. Yang, A. Bazavov, Z. Y. Xie, T. Xiang, S. Chandrasekharan, S.-W. Tsai, and Y. Meurice, *Phys. Rev. A* **90**, 063603 (Dec 2014).
- [30] A. Bazavov, C.-Y. Lai, S.-W. Tsai, and Y. Meurice(2014), arXiv:1411.4989 [hep-lat].
- [31] J. Heitger, “Numerical simulations of gauge-higgs models on the lattice,” (1997), (Ph. D. thesis).
- [32] D. Banerjee, M. Bögli, M. Dalmonte, E. Rico, P. Stebler, U.-J. Wiese, and P. Zoller, *Phys. Rev. Lett.* **110**, 125303 (Mar 2013).
- [33] E. Zohar, J. I. Cirac, and B. Reznik, *Phys. Rev. Lett.* **110**, 125304 (Mar 2013).
- [34] U. J. Wiese(2013), arXiv:1305.1602 [quant-ph].
- [35] M. Kalb and P. Ramond, *Phys. Rev. D* **9**, 2273 (Apr 1974).
- [36] The details will be presented in a separate publication.
- [37] B. Bunk, *Nucl.Phys.Proc.Suppl.* **42**, 566 (1995).
- [38] A. Bazavov and B. A. Berg, *Phys. Rev. D* **71**, 114506 (2005), arXiv:hep-lat/0503006.

- [39] Y. Meurice, Phys. Rev. B **87**, 064422 (Feb 2013).
- [40] M. B. Einhorn and R. Savit, Phys. Rev. D **17**, 2583 (May 1978).
- [41] M. B. Einhorn and R. Savit, Phys. Rev. D **19**, 1198 (Feb 1979).
- [42] E. Fradkin and L. Susskind, Phys. Rev. D **17**, 2637 (May 1978).
- [43] J. B. Kogut, Rev. Mod. Phys. **51**, 659 (Oct 1979).
- [44] A. M. Polyakov, *Gauge Fields and Strings* (Harwood, Chur, Switzerland, 1987).
- [45] J. M. Kosterlitz and D. J. Thouless, Journal of Physics C: Solid State Physics **6**, 1181 (1973).
- [46] A. B. Kuklov and B. V. Svistunov, Phys. Rev. Lett. **90**, 100401 (Mar. 2003).
- [47] A. Kuklov, N. Prokof'ev, and B. Svistunov, Phys. Rev. Lett. **92**, 050402 (Feb. 2004).
- [48] N. Hasselmann, A. H. Castro Neto, and C. M. Smith, Phys. Rev. B **65**, 220511 (Jun 2002).
- [49] L. Pollet and N. Prokof'ev, Phys. Rev. Lett. **109**, 010401 (Jul 2012).
- [50] M. Endres, T. Fukuhara, D. Pekker, M. Cheneau, P. Schauß, C. Gross, E. Demler, S. Kuhr, and I. Bloch, Nature (London) **487**, 454 (Jul. 2012), arXiv:1204.5183 [cond-mat.quant-gas].
- [51] D. Jaksch, C. Bruder, J. I. Cirac, C. W. Gardiner, and P. Zoller, Phys. Rev. Lett. **81**, 3108 (Oct 1998).
- [52] I. Bloch, J. Dalibard, and W. Zwerger, Rev. Mod. Phys. **80**, 885 (Jul 2008).
- [53] G. Mazzaella, S. M. Giampaolo, and F. Illuminati, Phys. Rev. A **73**, 013625 (Jan. 2006), cond-mat/0509161.
- [54] G. Thalhammer, G. Barontini, L. De Sarlo, J. Catani, F. Minardi, and M. Inguscio, Phys. Rev. Lett. **100**, 210402 (May 2008).
- [55] J. Catani, G. Lamporesi, D. Naik, M. Gring, M. Inguscio, F. Minardi, A. Kantian, and T. Giamarchi, Phys. Rev. A **85**, 023623 (Feb 2012).
- [56] J. Sebby-Strabley, M. Anderlini, P. S. Jessen, and J. V. Porto, Phys. Rev. A **73**, 033605 (Mar. 2006), cond-mat/0602103.
- [57] S. Fölling, S. Trotzky, P. Cheinet, M. Feld, R. Saers, A. Widera, T. Müller, and I. Bloch, Nature (London) **448**, 1029 (Aug. 2007), arXiv:0707.3985 [cond-mat.other].
- [58] Y.-A. Chen, S. D. Huber, S. Trotzky, I. Bloch, and E. Altman, Nature Physics **7**, 61 (Jan. 2011), arXiv:1003.4956 [cond-mat.quant-gas].
- [59] D. Greif, T. Uehlinger, G. Jotzu, L. Tarruell, and T. Esslinger, Science **340**, 1307 (Jun. 2013), arXiv:1212.2634 [cond-mat.quant-gas].
- [60] M. Atala, M. Aidelsburger, M. Lohse, J. T. Barreiro, B. Paredes, and I. Bloch, Nature Physics **10**, 588 (Aug. 2014), arXiv:1402.0819 [cond-mat.quant-gas].
- [61] A. Griesmaier, J. Werner, S. Hensler, J. Stuhler, and T. Pfau, Physical Review Letters **94**, 160401 (Apr. 2005), cond-mat/0503044.
- [62] K. Aikawa, A. Frisch, M. Mark, S. Baier, A. Rietzler, R. Grimm, and F. Ferlaino, ArXiv e-prints(Apr. 2012), arXiv:1204.1725 [cond-mat.quant-gas].
- [63] K. Aikawa, A. Frisch, M. Mark, S. Baier, R. Grimm, and F. Ferlaino, Physical Review Letters **112**, 010404 (Jan. 2014), arXiv:1310.5676 [cond-mat.quant-gas].
- [64] M. Lu, N. Q. Burdick, S. H. Youn, and B. L. Lev, Physical Review Letters **107**, 190401 (Nov. 2011), arXiv:1108.5993 [cond-mat.quant-gas].
- [65] M. Lu, N. Q. Burdick, and B. L. Lev, Physical Review Letters **108**, 215301 (May 2012), arXiv:1202.4444 [cond-mat.quant-gas].
- [66] Y. Tang, N. Q. Burdick, K. Baumann, and B. L. Lev, ArXiv e-prints(Nov. 2014), arXiv:1411.3069 [cond-mat.quant-gas].
- [67] K.-K. Ni, S. Ospelkaus, M. H. G. de Miranda, A. Pe'er, B. Neyenhuis, J. J. Zirbel, S. Kotochigova, P. S. Julienne, D. S. Jin, and J. Ye, Science **322**, 231 (Oct. 2008), arXiv:0808.2963 [quant-ph].
- [68] A. Chotia, B. Neyenhuis, S. A. Moses, B. Yan, J. P. Covey, M. Foss-Feig, A. M. Rey, D. S. Jin, and J. Ye, Physical Review Letters **108**, 080405 (Feb. 2012), arXiv:1110.4420 [physics.atom-ph].
- [69] C.-H. Wu, J. W. Park, P. Ahmadi, S. Will, and M. W. Zwierlein, Physical Review Letters **109**, 085301 (Aug. 2012), arXiv:1206.5023 [cond-mat.quant-gas].
- [70] O. Boada, A. Celi, J. I. Latorre, and M. Lewenstein, Phys. Rev. Lett. **108**, 133001 (Mar 2012).
- [71] A. Celi, P. Massignan, J. Ruseckas, N. Goldman, I. B. Spielman, G. Juzeliūnas, and M. Lewenstein, Phys. Rev. Lett. **112**, 043001 (Jan 2014).