

This is the accepted manuscript made available via CHORUS. The article has been published as:

Structure constant of twist-2 light-ray operators in the Regge limit

Ian Balitsky, Vladimir Kazakov, and Evgeny Sobko

Phys. Rev. D **93**, 061701 — Published 11 March 2016

DOI: [10.1103/PhysRevD.93.061701](https://doi.org/10.1103/PhysRevD.93.061701)

Structure constant of twist-2 light-ray operators in Regge limit

Ian Balitsky^a, Vladimir Kazakov^b, Evgeny Sobko^c

^a *Physics Dept., Old Dominion University, Norfolk VA 23529 & Theory Group, JLAB, 12000 Jefferson Ave, Newport News, VA 23606*

^b *LPT, École Normale Supérieure, 24, rue Lhomond 75005 Paris, France & Université Paris-VI, Place Jussieu, 75005 Paris, France*

^c *DESY Hamburg, Theory Group, Notkestraße 85, 22607 Hamburg, Germany*

We compute the normalized structure constant of three twist-2 operators in $\mathcal{N} = 4$ SYM in the leading BFKL approximation at any N_c . The result is applicable to other gauge theories including QCD.

INTRODUCTION

The problem of high-energy behavior of amplitudes has a long story [1, 2]. One of the most popular approaches is to reduce the gauge theory at high energies to 2+1 effective theory which can be solved exactly or by computer simulations. Unfortunately, despite the multitude of attempts, the Lagrangian for 2+1 QCD at high energies is not written yet. In this context the idea to solve formally the high-energy QCD or $\mathcal{N} = 4$ SYM by calculation of anomalous dimensions and structure constants in the BFKL limit seems to be very promising.

$\mathcal{N} = 4$ SYM is a superconformal theory and its most important physical properties are encoded into the OPE characterized by the spectrum of anomalous dimensions and by the structure constants. While the former is now exactly and efficiently computable at large N_c due to quantum integrability [3], the calculation of the OPE structure constants is these days on a fast track, especially after the ground-breaking all-loop proposal of [4].

In this note we calculate the 3-point correlator of twist-2 operators $\mathcal{O}^j(x) = \text{tr} F_{+i} D_+^{j-2} F_+^i + \text{fermions} + \text{scalars}$ in $\mathcal{N} = 4$ SYM in the BFKL limit [5] when $\omega = j-1 \rightarrow 0$, the 't Hooft coupling $g^2 \equiv \frac{N_c g_{YM}^2}{16\pi^2} \rightarrow 0$ and $\frac{g^2}{\omega}$ fixed, for arbitrary N_c . The symbol '+' in the field-strength tensor F_{+i} means contraction with light-ray vector n_+ and the summation over index 'i' goes over two-dimensional space orthogonal to n_+ and n_- . Since the contribution of *fermions+scalars* is subleading at this limit, including the internal loops, the result is valid for the pure Yang-Mills theory as well. The case of two-point correlator was elaborated in our previous paper [6] where we defined the generalized operators with complex spin as special light-ray operators [7] (regularized as a narrow rectangular Wilson contour called "frame") and calculated their correlator using OPE over Wilson lines [8] with a rapidity cutoff and the BFKL evolution (see Fig. 1). Here we use the same light-ray operators: one along n_+ direction and two along n_- . In this case we should use more general Balitsky-Kovchegov (BK) evolution [9, 10] and the leading BFKL contribution comes from the BK vertex.




FIG. 1. Scheme of computation of 2-point correlator. In the l.h.s., the long sides of regularizing rectangular Wilson frames are stretched along light ray and the short sides in the orthogonal directions. In the r.h.s. we use OPE of frames over color dipoles and compute their correlator, see [6] for details.

LIGHT-RAY OPERATORS AND THEIR RELATION TO LOCAL OPERATORS

The generalization of local operator \mathcal{O}^j for the case of complex spin j was constructed in [6]. It has a form of light-ray operator $\check{\mathcal{S}}^J$ stretched along n_+ direction and realizing the principal series representation of $sl(2|4)$ with conformal spin $J = \frac{1}{2} + i\nu$ which is related to Lorentz spin j as $J = j + 1$. The full regularized operator reads as follows:

$$\check{\mathcal{S}}^{j+1}(x_{1\perp}) = \check{\mathcal{S}}_{gl}^{j+1}(x_{1\perp}) + \frac{i}{2}(j-1)\check{\mathcal{S}}_f^{j+1}(x_{1\perp}) - \frac{1}{2}(j)(j-1)\check{\mathcal{S}}_{sc}^{j+1}(x_{1\perp}), \quad (1)$$

where for example, the regularized gluon operator is:

$$\check{\mathcal{S}}_{gl}^{j+1}(x_{1\perp}) = \lim_{|x_{31\perp}| \rightarrow 0} |x_{13\perp}|^{-\gamma_j} \check{\mathcal{S}}_{gl}^{j+1}(x_{1\perp}, x_{3\perp}),$$

$\check{\mathcal{S}}_{gl}^{j+1}(x_{1\perp}, x_{3\perp}) = \int_{-\infty}^{\infty} \int_{x_{1-}}^{\infty} \frac{dx_{1-} dx_{3-}}{x_{31-}^{j-1}} \text{tr} F_{+i}(x_1) [1, 3]_{\square} F_{+i}(x_3)$ and $x_1 = (x_{1-}, 0, x_{1\perp})$, $x_3 = (x_{3-}, 0, x_{3\perp})$. The anomalous dimension γ_j corresponds to operator $\check{\mathcal{S}}^{j+1}(x_{1\perp})$. Here we introduced the notation $[1, 3]_{\square}$ for rectangular Wilson contour with coordinates x_1, x_3 of two diagonally opposite corners, as in Fig. 1. In the case of even integer Lorentz spin j it can be rewritten as an integral of

local operator $\mathcal{O}^j(x)$ with dimension $\Delta(j)$ along a light ray direction n_+ :

$$\check{\mathcal{S}}^{j+1}(x_\perp)|_{j \in \text{Even}} \sim \int_{-\infty}^{\infty} dx_- \mathcal{O}^j(x) \quad (2)$$

In this case the correlator of two light-ray operators stretched along n_+ and n_- vectors, normalized as $\langle n_+ n_- \rangle = 1$, is just the double integral of two-point correlator of local operators w.r.t. light-ray directions n_\pm :

$$\langle \check{\mathcal{S}}^{j_1+1}(x_\perp) \check{\mathcal{S}}^{j_2+1}(y_\perp) \rangle = \frac{\delta(j_1 - j_2) b_{j_1}}{(|x - y|_\perp^2)^{\Delta(j_1)-1}} \quad (3)$$

In this note, we calculate the correlator of three light-ray operators, restricting ourselves to a particular simple kinematics: one light-ray operator is stretched along n_+ light-ray direction and two other – along n_- . The correlator of 3 light-ray operators can be obtained by integrating the correlator of 3 local operators along these light-rays. The tensor structures of such local correlators are known from general group-theoretical considerations [11], up to a few structure constants depending on the coupling and symmetry charges. The main problem which we are addressing here is the calculation of these non-trivial constants. Remarkably, if the coordinates of all 3 light-rays in the transverse space are restricted to the same line all these structures collapse into a single one [13], with a single overall structure constant which we are going to compute. Note that after a conformal transformation the three points in the transverse space take arbitrary positions.

However, the configuration with two collinear light-ray operators is singular, so we first consider three different light-ray directions n_1, n_2, n_3 and then take the limit $n_2 \rightarrow n_3$. The result of integration along light-rays is quite simple and contains only one unknown overall constant

$$\langle \check{\mathcal{S}}^{j_1+1}(x_\perp) \check{\mathcal{S}}^{j_2+1}(y_\perp) \check{\mathcal{S}}^{j_3+1}(z_\perp) \rangle = C_{\{n_i\}}(\{\Delta_i\}, \{j_i\}) \cdot \frac{\langle n_1 n_2 \rangle^{[j]_{1,2;3}} \langle n_1 n_3 \rangle^{[j]_{1,3;2}} \langle n_2 n_3 \rangle^{[j]_{2,3;1}}}{(|x - y|_\perp^2)^{[\Delta]_{1,2;3}} (|x - z|_\perp^2)^{[\Delta]_{1,3;2}} (|y - z|_\perp^2)^{[\Delta]_{2,3;1}}} \quad (4)$$

where we used a short-hand notation $[a]_{i,j;k} \equiv \frac{1}{2}(a_i + a_j - a_k - 1)$ and $\{a_i\} \equiv \{a_1, a_2, a_3\}$. In what follows, we assume the existence of a good analytic continuation for $C_{\{n_i\}}(\{\Delta(j_i)\}, \{j_i\})$ to non-integer $\{j_i\}$'s. We take the limit $n_1 = n_+, n_2 = n_-, n_3 \rightarrow n_2$ with the normalization $\langle n_+ n_- \rangle = 1$. In BFKL regime $j_i = 1 + \omega_i \rightarrow 1$ we obtain:

$$\begin{aligned} & \langle \check{\mathcal{S}}^{2+\omega_1}(x_\perp) \check{\mathcal{S}}^{2+\omega_2}(y_\perp) \check{\mathcal{S}}^{2+\omega_3}(z_\perp) \rangle = \\ & = \lim_{n_3 \rightarrow n_2 = n_-} \frac{\langle n_2 n_3 \rangle^{\frac{\omega_2 + \omega_3 - \omega_1}{2}}}{\omega_2 + \omega_3 - \omega_1} \times \\ & \times \frac{C_{+--}(\{\Delta_i\}, \{1 + \omega_i\})}{(|x - y|_\perp^2)^{[\Delta]_{1,2;3}} (|x - z|_\perp^2)^{[\Delta]_{1,3;2}} (|y - z|_\perp^2)^{[\Delta]_{2,3;1}}} \quad (5) \end{aligned}$$

where $\Delta_i = \Delta(1 + \omega_i, g^2)$ is given by BFKL spectrum (see below). We explicitly pulled out the denominator $\frac{1}{\omega_2 + \omega_3 - \omega_1}$ because it will emerge in our forthcoming calculation using the BK evolution. We interpret $\lim_{\langle n_2 n_3 \rangle \rightarrow 0} \frac{\langle n_2 n_3 \rangle^{\frac{\omega_2 + \omega_3 - \omega_1}{2}}}{\omega_2 + \omega_3 - \omega_1}$ as a delta function $\delta(\omega_2 + \omega_3 - \omega_1)$ reflecting the boost invariance. In addition, we keep ω_i positive through the paper.

Finally, the structure constant is normalized using the corresponding 2-point correlators:

$$C_{\omega_1, \omega_2, \omega_3} = \frac{C_{+--}(\{\Delta_i\}, \{1 + \omega_i\})}{\sqrt{b_{1+\omega_1} b_{1+\omega_2} b_{1+\omega_3}}} \quad (6)$$

DECOMPOSITION OVER DIPOLES AND BK EVOLUTION

When calculating the two-point correlator [6] we used a point splitting regularization in orthogonal direction, replacing light-rays by infinitely narrow Wilson frames with inserted fields in the corners (see Fig. 1). Now, for the sake of simplicity, we carry out our calculation for pure Wilson frames, related to our operators with zero R -charge in the following way:

$$\begin{aligned} & \partial_{x_{1\perp}} \cdot \partial_{x_{3\perp}} \int \int \frac{dx_{1-} dx_{3-}}{(x_{3-} - x_{1-})^{2+\omega_1}} [x_1, x_3]_\square \rightarrow \\ & \xrightarrow{x_{13\perp} \rightarrow 0, \omega_1 \rightarrow 0} |x_{13\perp}|^{\gamma_{j_1}} c(g_{YM}^2, N_c, \omega_1) \check{\mathcal{S}}^{2+\omega_1}(x_{1\perp}). \quad (7) \end{aligned}$$

The coefficient $c(g_{YM}^2, N_c, \omega_i)$ (denoted below as $c(\omega_i)$) depends on the local regularization procedure and at weak coupling it behaves as $c(\omega_i) \sim \frac{g_{YM}^2}{\omega_i}$, but its explicit form is irrelevant for us because we are going to calculate the normalized structure constant where it cancels. In general, there are a few types of leading twist-2 operators which appear in this decomposition but in the BFKL limit a single one with the smallest anomalous dimension survives. In addition, in the $\omega_i \rightarrow 0$ limit only the term built out of gauge fields alone does contribute [6].

Following the OPE method [8], the pure Wilson frames can be replaced by regularized color dipoles:

$$[x_1, x_3]_\square \rightarrow N(1 - \mathbf{U}^{\sigma_+}(x_{1\perp}, x_{3\perp})) \quad (8)$$

where

$$\mathbf{U}^{\sigma_+}(x_{1\perp}, x_{3\perp}) = 1 - \frac{1}{N} \text{tr}(U_{x_{1\perp}}^{\sigma_+} U_{x_{3\perp}}^{\sigma_+ \dagger}), \quad (9)$$

$$U_{x_\perp}^{\sigma_+} = P \exp[ig_{YM} \int_{-\infty}^{\infty} dx_+ A_-^{\sigma_+}(x)], \quad (10)$$

$$A_\mu^{\sigma_+}(x) = \int d^4 k \theta(\sigma_+ - |k_+|) e^{ikx} A_\mu(k) \quad (11)$$

and σ_+ is a longitudinal cutoff in n_+ direction. Now we

can write:

$$\begin{aligned} \langle S^{2+\omega_1}(x_{1\perp}, x_{3\perp}) S^{2+\omega_2}(y_{1\perp}, y_{3\perp}) S^{2+\omega_3}(z_{1\perp}, z_{3\perp}) \rangle = \\ = -\mathcal{D}_\perp \int_{-\infty}^{\infty} dx_{1-} \int_{x_{1-}}^{\infty} dx_{3-} x_{31-}^{-2-\omega_1} \times \\ \times \int_{-\infty}^{\infty} dy_{1+} \int_{y_{1+}}^{\infty} dy_{3+} y_{31+}^{-2-\omega_2} \int_{-\infty}^{\infty} dz_{1+} \int_{z_{1+}}^{\infty} dz_{3+} z_{31+}^{-2-\omega_3} \times \\ \times \langle \mathbf{U}^{\sigma_{1-}}(x_{1\perp}, x_{3\perp}) \mathbf{V}^{\sigma_{2+}}(y_{1\perp}, y_{3\perp}) \mathbf{W}^{\sigma_{3+}}(z_{1\perp}, z_{3\perp}) \rangle, \end{aligned} \quad (12)$$

$$\text{where } \mathcal{D}_\perp = \frac{N^3(\partial_{x_{1+}} \cdot \partial_{x_{3+}})(\partial_{y_{1+}} \cdot \partial_{y_{3+}})(\partial_{z_{1+}} \cdot \partial_{z_{3+}})}{c(\omega_1)c(\omega_2)c(\omega_3)}.$$

In our kinematics two dipoles \mathbf{V} and \mathbf{W} have zero n_+ projection and in the BFKL approximation they form a "pancake" field configuration in the reference frame related to \mathbf{U} . This means that the rapidity of \mathbf{U} serves as the upper limit for integrations w.r.t. rapidities of \mathbf{V} and \mathbf{W} in our logarithmic approximation. Now we use the BK evolution equation [9, 10] to calculate the quantum average in (12). It gives the evolution of the dipole \mathbf{U}^Y with respect to rapidity $Y = e^\sigma$, namely

$$\sigma \frac{d}{d\sigma} \mathbf{U}^\sigma(z_1, z_2) = \mathcal{K}_{\text{BK}} * \mathbf{U}^\sigma(z_1, z_2), \quad (13)$$

where \mathcal{K}_{BK} is an integral operator having the following form in LO approximation:

$$\begin{aligned} \mathcal{K}_{\text{LO BK}} * \mathbf{U}(z_1, z_2) = \frac{2g^2}{\pi} \int d^2 z_3 \frac{z_{12}^2}{z_{13}^2 z_{23}^2} [\mathbf{U}(z_1, z_3) + \\ + \mathbf{U}(z_3, z_2) - \mathbf{U}(z_1, z_2) - \mathbf{U}(z_1, z_3) \mathbf{U}(z_3, z_2)]. \end{aligned} \quad (14)$$

Evolution of \mathbf{U}^{Y_1} goes from Y_1 to an intermediate Y_0 w.r.t. the linear part of (13), and then the BK vertex acts at Y_0 and generates two dipoles which can be contracted with \mathbf{V}^{Y_2} and \mathbf{W}^{Y_3} . Schematically, it can be written as:

$$\int dY_0 (\mathbf{U}^{Y_1} \rightarrow \mathbf{U}^{Y_0}) \otimes (\text{BK vertex at } Y_0) \otimes \left(\langle \mathbf{U}^{Y_0} \mathbf{V}^{Y_2} \rangle \right) \left(\langle \mathbf{U}^{Y_0} \mathbf{W}^{Y_3} \rangle \right)$$

The linear BFKL evolution of \mathbf{U}^{Y_1} from Y_1 to Y_0 gives:

$$\begin{aligned} \mathbf{U}^{Y_1}(x_1, x_3) = \int d\nu \int d^2 x_0 \frac{\nu_1^2}{\pi^2} E_{\nu_1}(x_{10}, x_{30}) e^{\aleph(\nu_1)Y_{10}} \cdot \\ \cdot \frac{1}{\pi^2} \int \frac{d^2 \gamma d^2 \beta}{|\gamma - \beta|^4} E_{\nu_1}^*(\gamma - x_0, \beta - x_0) \mathbf{U}^{Y_0}(\gamma, \beta), \end{aligned} \quad (15)$$

where we denoted $Y_{ij} \equiv Y_i - Y_j$ and we introduced the function $E_\nu(z_{10}, z_{20}) = \left(\frac{|z_{12}|^2}{|z_{10}|^2 |z_{20}|^2} \right)^{1/2 + i\nu}$ which projects dipoles on the eigenstates of BFKL operator with the eigenvalues $\aleph(\nu) = 4g^2(2\psi(1) - \psi(1/2 + i\nu) - \psi(1/2 - i\nu))$. We take here only the sector $n = 0$, where n is the discrete quantum number of $SL(2, C)$ because it gives the leading contribution.

The non-linear part of BK evolution (13) is described by the following renorm group equation:

$$\begin{aligned} \left. \frac{\partial}{\partial Y} \mathbf{U}^Y(\gamma, \beta) \right|_{Y=Y_0} = \\ = -\frac{2g^2}{\pi} \int d^2 \alpha \frac{|\gamma - \beta|^2}{|\gamma - \alpha|^2 |\beta - \alpha|^2} \mathbf{U}^{Y_0}(\gamma, \alpha) \mathbf{U}^{Y_0}(\alpha, \beta) \end{aligned} \quad (16)$$

Finally, we contract the two emerging dipoles $\mathbf{U}^{Y_0}(\gamma, \alpha)$ and $\mathbf{U}^{Y_0}(\alpha, \beta)$ with $\mathbf{V}^{\sigma_{2+}}(y_{1\perp}, y_{3\perp})$ and $\mathbf{W}^{\sigma_{3+}}(z_{1\perp}, z_{3\perp})$. Thus for the planar contribution we get:

$$\begin{aligned} \langle \mathbf{U}^{Y_1}(x_{1\perp}, x_{3\perp}) \mathbf{V}^{Y_2}(y_{1\perp}, y_{3\perp}) \mathbf{W}^{Y_3}(z_{1\perp}, z_{3\perp}) \rangle_{pl} = \\ = -\frac{2g^2}{\pi} \int dY_0 \int d\nu_1 \int d^2 x_0 \frac{\nu_1^2}{\pi^2} E_{\nu_1}(x_{10}, x_{30}) e^{\aleph(\nu_1)Y_{10}} \times \\ \times \frac{1}{\pi^2} \int \frac{d^2 \alpha d^2 \beta d^2 \gamma}{|\gamma - \beta|^2 |\gamma - \alpha|^2 |\beta - \alpha|^2} E_{\nu_1}^*(\gamma - x_0, \beta - x_0) \cdot \\ \cdot (\langle \mathbf{U}^{Y_0}(\gamma, \alpha) \mathbf{V}^{Y_2}(y_{1\perp}, y_{3\perp}) \rangle \langle \mathbf{U}^{Y_0}(\alpha, \beta) \mathbf{W}^{Y_3}(z_{1\perp}, z_{3\perp}) \rangle + \\ + \langle \mathbf{U}^{Y_0}(\gamma, \alpha) \mathbf{W}^{Y_3}(z_{1\perp}, z_{3\perp}) \rangle \langle \mathbf{U}^{Y_0}(\alpha, \beta) \mathbf{V}^{Y_2}(y_{1\perp}, y_{3\perp}) \rangle) \end{aligned} \quad (17)$$

The last two terms in (17) give the same contribution so it is enough to know the correlators of two dipoles [6]:

$$\begin{aligned} \langle \mathbf{U}^{Y_0}(\gamma, \alpha) \mathbf{V}^{Y_2}(y_{1\perp}, y_{3\perp}) \rangle = \frac{8g^4(1 - N_c^2)}{N_c^4} \int d^2 y_0 \cdot \\ \cdot \int \frac{d\nu_2 \nu_2^2 e^{Y_{02}\aleph(\nu_2)}}{(\frac{1}{4} + \nu_2^2)^2} E_{\nu_2}(\gamma - y_0, \alpha - y_0) E_{\nu_2}^*(y_{10}, y_{30}) \end{aligned} \quad (18)$$

and similarly for $\langle \mathbf{U}^{Y_0}(\alpha, \beta) \mathbf{W}^{Y_3}(z_{1\perp}, z_{3\perp}) \rangle$. It was argued in [6] that we can make the following identification for rapidities in dipole correlators: $Y_{02} = \ln \frac{L_0 y_{31+}}{\Lambda^2}$, $Y_{03} = \ln \frac{L_0 z_{31+}}{\Lambda^2}$, where Λ a cutoff whose precise value is irrelevant in LO. On the other hand, the difference of rapidities of the first dipole and of the BK vertex $Y_{10} = \ln \frac{x_{31-}}{L_0}$ corresponds to BFKL evolution. The integral over $Y_0 = \ln \frac{L_0}{\Lambda}$ goes from Y_1 to $\max(Y_2, Y_3)$. If we plug (17)-(18) into (12) and do the integrals over light ray directions, i.e. over rapidities, we obtain the following planar contribution:

$$\begin{aligned} \langle S^{2+\omega_1}(x_{1\perp}, x_{3\perp}) S^{2+\omega_2}(y_{1\perp}, y_{3\perp}) S^{2+\omega_3}(z_{1\perp}, z_{3\perp}) \rangle_{pl} = \\ = \frac{2^8 g^{10} (N_c^2 - 1)^2}{\pi^3 N_c^8} \delta(\omega_1 - \omega_2 - \omega_3) D_\perp \cdot \\ \cdot \int d\nu_1 \frac{\nu_1^2}{\pi^2} \frac{1}{\omega_2 + \omega_3 - \aleph(\nu_1)} \int \frac{d\nu_2 \nu_2^2}{(\frac{1}{4} + \nu_2^2)^2} \frac{1}{\omega_2 - \aleph(\nu_2)} \cdot \\ \cdot \int \frac{d\nu_3 \nu_3^2}{(\frac{1}{4} + \nu_3^2)^2} \frac{1}{\omega_3 - \aleph(\nu_3)} \int d^2 x_0 d^2 y_0 d^2 z_0 E_{\nu_1}^*(x_{10}, x_{30}) \cdot \\ \cdot E_{\nu_2}^*(y_{10}, y_{30}) E_{\nu_3}^*(z_{10}, z_{30}) \Upsilon_{pl}(\nu_1, \nu_2, \nu_3; x_0, y_0, z_0) \end{aligned} \quad (19)$$

The usual delta-function $\delta(\omega_1 - \omega_2 - \omega_3)$ (see e.g. [14]) is a consequence of boost-invariance as in the formula (5).

Υ_{pl} represents the planar contribution of BK vertex:

$$\begin{aligned} \Upsilon_{pl}(\nu_1, \nu_2, \nu_3; x_0, y_0, z_0) &= \\ &= \int \frac{d^2\alpha d^2\beta d^2\gamma}{|\gamma - \beta|^2 |\gamma - \alpha|^2 |\beta - \alpha|^2} E_{\nu_1}(\beta - x_0, \gamma - x_0) \cdot \\ &\cdot E_{\nu_2}(\alpha - y_0, \gamma - y_0) E_{\nu_3}(\alpha - z_0, \beta - z_0) = \quad (20) \\ &= \frac{\Omega(h_1, h_2, h_3)}{|x_0 - y_0|^{4[h]_{1,2,3}+2} |x_0 - z_0|^{4[h]_{1,3,2}+2} |y_0 - z_0|^{4[h]_{2,3,1}+2}} \end{aligned}$$

where $h_1 = \frac{1}{2} + i\nu_1$, $h_2 = \frac{1}{2} + i\nu_2$, $h_3 = \frac{1}{2} + i\nu_3$ and the function $\Omega(h_1, h_2, h_3)$ was presented in [15].

Remarkably, we can also take into account the non-planar contribution [15, 16], thus providing the finite N_c answer for the BFKL structure constant! It appears as a single extra term Υ_{npl} :

$$\begin{aligned} \Upsilon_{npl}(\nu_1, \nu_2, \nu_3; x_0, y_0, z_0) &= \int \frac{d^2\beta d^2\gamma}{|\gamma - \beta|^4} E_{\nu_1}(\beta - x_0, \\ &\gamma - x_0) E_{\nu_2}(\beta - y_0, \gamma - y_0) E_{\nu_3}(\beta - z_0, \gamma - z_0) = \quad (21) \\ &= \frac{\Lambda(h_1, h_2, h_3)}{|x_0 - y_0|^{4[h]_{1,2,3}+2} |x_0 - z_0|^{4[h]_{1,3,2}+2} |y_0 - z_0|^{4[h]_{2,3,1}+2}} \end{aligned}$$

where $\Lambda(h_1, h_2, h_3)$ was also presented in [15], and the full answer can be obtained from (19) by replacing Υ_{pl} with Υ (see in Fig. 2):

$$\begin{aligned} \Upsilon &= \Upsilon_{pl} - \frac{2\pi}{N_c^2} \Upsilon_{npl} \text{Re}[\psi(1) + \\ &+ \psi(\frac{1}{2} + i\nu_1) - \psi(\frac{1}{2} + i\nu_2) - \psi(\frac{1}{2} + i\nu_3)]. \quad (22) \end{aligned}$$

The integrals over x_0, y_0, z_0 are easily computable, e.g.

$$\begin{aligned} \int d^2x_0 E_{\nu_1}(\beta - x_0, \gamma - x_0) E_{\nu_1}^*(x_{10}, x_{30}) &= \\ &= (\tau^2)^{\frac{1}{2} + i\nu_1} {}_2F_1(\frac{1}{2} + i\nu, \frac{1}{2} + i\nu, 1 + 2i\nu, \tau) \times \\ &\times {}_2F_1(\frac{1}{2} + i\nu, \frac{1}{2} + i\nu, 1 + 2i\nu, \bar{\tau}) \frac{(\frac{1}{4} + \nu^2)^2}{\nu^2} G(\nu) + \\ &+ (\nu \leftrightarrow -\nu), \quad (23) \\ G(\nu) &= \frac{\nu^2}{(\frac{1}{4} + \nu^2)^2} \frac{\pi \Gamma^2(\frac{1}{2} + i\nu) \Gamma(-2i\nu)}{\Gamma^2(\frac{1}{2} - i\nu) \Gamma(1 + 2i\nu)}, \quad (24) \end{aligned}$$

where $\tau = \frac{|x_1 - x_3| |\beta - \gamma|}{|x_1 - \beta| |x_3 - \gamma|}$. In the limit $x_1, x_3 \rightarrow x$ we can replace $\frac{|x_1 - x_3| |\beta - \gamma|}{|x_1 - \beta| |x_3 - \gamma|} \rightarrow \frac{|x_1 - x_3| |\beta - \gamma|}{|x - \beta| |x - \gamma|} \rightarrow 0$. For small τ we close the ν_1 contour in the lower (upper) half-plane for first(second) term, respectively, both of them giving the same contribution. Integrals over α, β, γ in (19) can be reduced to Υ_{pl} represented in [15] in terms of hypergeometric and Meijer G functions, and Υ_{npl} in terms of Γ -functions. Integrals over ν_i can be done by picking up the BFKL poles $\omega_i = \aleph(\nu_i^*)$.

Combining (19), (22) and (23) we come to the final ex-

pression for 3-point correlation function:

$$\begin{aligned} \langle S^{2+\omega_1}(x_{1\perp}, x_{3\perp}) S^{2+\omega_2}(y_{1\perp}, y_{3\perp}) S^{2+\omega_3}(z_{1\perp}, z_{3\perp}) \rangle &= \\ &= -ig^{10} \frac{\delta(\omega_1 - \omega_2 - \omega_3)}{c(\omega_1)c(\omega_2)c(\omega_3)} H. \quad (25) \\ &\cdot \frac{\Psi(\nu_1^*, \nu_2^*, \nu_3^*) |x_{13}|^{\gamma_1} |y_{13}|^{\gamma_2} |z_{13}|^{\gamma_3}}{|x - y|^{2+\gamma_1+\gamma_2-\gamma_3} |x - z|^{2+\gamma_1+\gamma_3-\gamma_2} |y - z|^{2+\gamma_2+\gamma_3-\gamma_1}} \end{aligned}$$

$$\begin{aligned} \text{where } H &= \frac{2^{10}(N_c^2 - 1)^2}{\pi^2 N_c^5} \gamma_1^2 (2 + \gamma_1)^4 (2 + \gamma_2)^2 \times \\ &\times (2 + \gamma_3)^2 \frac{G(\nu_1^*)}{\aleph'(\nu_1^*)} \frac{G(\nu_2^*)}{\aleph'(\nu_2^*)} \frac{G(\nu_3^*)}{\aleph'(\nu_3^*)}, \quad (26) \end{aligned}$$

$\gamma_i = \gamma(1 + \omega_i)$ - anomalous dimension and the coefficient $\Psi(\nu_1^*, \nu_2^*, \nu_3^*)$ can be expressed through the functions $\Omega(h_1, h_2, h_3)$ and $\Lambda(h_1, h_2, h_3)$ defined in (20)-(21) and calculated in [15]:

$$\begin{aligned} \Psi(\nu_1^*, \nu_2^*, \nu_3^*) &= \Omega(h_1^*, h_2^*, h_3^*) - \frac{2\pi}{N_c^2} \Lambda(h_1^*, h_2^*, h_3^*) \cdot \\ &\cdot \text{Re}(\psi(1) - \psi(h_1^*) - \psi(h_2^*) - \psi(h_3^*)), \quad (27) \end{aligned}$$

where $h_i^* = \frac{1}{2} + i\nu_i^* = 1 + \frac{\gamma_i}{2}$.

Our final result for normalized structure constant is:

$$\begin{aligned} C_{\omega_1, \omega_2, \omega_3} &= -i^{1/2} g^4 \frac{2}{\pi^5} \frac{\sqrt{N_c^2 - 1}}{N_c^2} \gamma_1^2 (2 + \gamma_1)^2 \cdot \\ &\cdot \sqrt{\frac{G(\nu_1^*)}{\aleph'(\nu_1^*)} \frac{G(\nu_2^*)}{\aleph'(\nu_2^*)} \frac{G(\nu_3^*)}{\aleph'(\nu_3^*)}} \Psi(\nu_1^*, \nu_2^*, \nu_3^*) \quad (28) \end{aligned}$$

Precising the dependence on parameters $\{\frac{g^2}{\omega_i}\}$, g^2 and N_c we can write: $C_{\omega_1, \omega_2, \omega_3} = g \frac{\sqrt{N_c^2 - 1}}{N_c^2} f(\frac{g^2}{\omega_1}, \frac{g^2}{\omega_2}, \frac{g^2}{\omega_3})$, where f is a function which depends only on the ratios $\{\frac{g^2}{\omega_i}\}$. In the limit $\frac{g^2}{\omega_i} \rightarrow 0$ we get the asymptotics:

$$\begin{aligned} \Omega(h_1^*, h_2^*, h_3^*) &\rightarrow -\frac{16\pi^3}{\gamma_1^2 \gamma_2^2 \gamma_3^2} \cdot [\gamma_1^2 (\gamma_2 + \gamma_3) + \gamma_2^2 (\gamma_1 + \gamma_3) + \\ &+ \gamma_3^2 (\gamma_1 + \gamma_2) + \gamma_1 \gamma_2 \gamma_3] (1 + O(g^2/\omega_i)) \\ \Lambda(h_1^*, h_2^*, h_3^*) &\rightarrow \frac{8\pi^2 (\gamma_1 + \gamma_2 + \gamma_3)}{\gamma_1 \gamma_2 \gamma_3} (1 + O(g^2/\omega_i)) \quad (29) \end{aligned}$$

In this limit $\gamma_i = -\frac{8g^2}{\omega_i} + o(\frac{g^2}{\omega_i})$ and the main contribution to 3-point correlator (28) comes from the planar $\mathcal{O}(g^2)$ term

$$\begin{aligned} C_{\omega_1, \omega_2, \omega_3} &= -ig^2 \frac{\sqrt{N_c^2 - 1}}{\sqrt{2\pi} N_c^2} \frac{1}{\omega_1^{\frac{5}{2}} \omega_2^{\frac{1}{2}} \omega_3^{\frac{1}{2}}} (\omega_1^2 (\omega_2 + \omega_3) + \\ &\omega_2^2 (\omega_1 + \omega_3) + \omega_3^2 (\omega_1 + \omega_2) + \omega_1 \omega_2 \omega_3) (1 + O(g^2)) \quad (30) \end{aligned}$$

whereas the nonplanar one is $\mathcal{O}(g^6)$. It might seem strange that the planar contribution does not start from $\mathcal{O}(g^4)$ terms given by the leading Feynman graphs, e.g.




FIG. 2. The structure of 3-point correlator. Red circles correspond to BFKL propagators (the crossed one has extra multiplier $(\frac{1}{4} + \nu_1^2)$). The blue blob corresponds to the 3-point functions of 2-dimensional BFKL CFT. The triple "Y"-vertices correspond to E -functions. For example vertex with ends labeled as $z_{1\perp}, z_{3\perp}, z_{0\perp}$ corresponds to $E_\nu(z_{10\perp}, z_{30\perp}) = (\frac{|z_{13\perp}|^2}{|z_{10\perp}|^2 |z_{30\perp}|^2})^{1/2 + i\nu_3}$. The $\alpha\beta\gamma$ -triangle in the first, planar, term and $\beta\gamma$ -link in the second, nonplanar, term correspond to triple pomeron vertex.

with 4 gluon vertices. However, in BFKL approximation we should keep $\frac{g^2}{\omega} \gg \omega$ [17]. In addition, when making the point-splitting regularization we have to keep $\frac{g^2}{\omega} |\ln(x_{31\perp}/(x-y))^2| \gg 1$. The limit $|x_{13\perp}|$ has to be taken first, which makes the value $g^2 = 0$ exceptional. This order of limits leads to $\mathcal{O}(g^2)$ behavior of (30).

DISCUSSION

Our result eq.(28), based on BFKL approximation is a rare example of computation of a non-BPS structure constant receiving contributions from all orders in coupling constant, including infinitely many "wrapping" corrections. Moreover, our result is valid at any N_c . Since in the LO BFKL the contributions of all fields but gluons in $\mathcal{N} = 4$ SYM disappear from both the definition of operators and internal loops, the result is applicable to pure YM theory at any N_c , including $N_c = 3$. It would be interesting to apply our structure constants to the OPE at hard scattering in real QCD and to work out the full "dictionary" relating them to the OPE in the 2-dimensional $SL(2, C)$ CFT – the basis of our BFKL computation. It is also not hopeless, though challenging, to compute these structure constants in the NLO approximation in $\mathcal{N} = 4$ SYM. Our present result may serve as an important, all-wrappings test for the future computations of similar quantities in the integrability approaches to planar AdS_5/CFT_4 , such as [4] and the BFKL limit of quantum spectral curve [18].

ACKNOWLEDGMENTS

We thank J. Bartels, S. Caron-Huot, L. Lipatov and V. Schomerus for discussions. Our special thanks to G. Korchemsky who participated in the initial stage of this work. The work of E.S. and V.K. was supported by the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme FP7/2007-2013/ under REA Grant Agreement No 317089 (GATIS). The work of V.K. has received funding from the European Research Council (Programme Ideas ERC-2012-AdG 320769 AdS-CFT-solvable), from the ANR grant StrongInt (BLANC- SIMI- 4-2011) and from the ESF grant HOLOGRAV-09- RNP- 092. The work of I.B. was supported by DOE contract DE-AC05-06OR23177 and by the grant DE-FG02-97ER41028.

-
- [1] B. L. Ioffe, V. S. Fadin and L. N. Lipatov, "Quantum chromodynamics: Perturbative and nonperturbative aspects,"
 - [2] Y. V. Kovchegov and E. Levin, "Quantum chromodynamics at high energy,"
 - [3] N. Beisert, C. Ahn, L. F. Alday, Z. Bajnok, J. M. Drummond, L. Freyhult, N. Gromov and R. A. Janik *et al.*, Lett. Math. Phys. **99** (2012) 3 [arXiv:1012.3982 [hep-th]]; N. Gromov, V. Kazakov, S. Leurent and D. Volin, arXiv:1405.4857 [hep-th].
 - [4] B. Basso, S. Komatsu and P. Vieira, arXiv:1505.06745 [hep-th].
 - [5] V. S. Fadin, E. A. Kuraev and L. N. Lipatov, Phys. Lett. B **60** (1975) 50; I. Balitsky and L. Lipatov, Sov.J.Nucl.Phys. **28** (1978) 822–829.
 - [6] I. Balitsky, V. Kazakov and E. Sobko, arXiv:1310.3752 [hep-th].
 - [7] I. Balitsky and V. M. Braun, Nucl. Phys. B **311** (1989) 541.
 - [8] I. Balitsky, Nucl. Phys. B **463** (1996) 99 [hep-ph/9509348].
 - [9] I. Balitsky, AIP Conf. Proc. **407** (1997) 953 [hep-ph/9706411].
 - [10] Y. V. Kovchegov, Phys. Rev. D **60** (1999) 034008 [hep-ph/9901281]; Y. V. Kovchegov, Phys. Rev. D **61** (2000) 074018 [hep-ph/9905214].
 - [11] M. S. Costa, J. Penedones, D. Poland and S. Rychkov, JHEP **1111** (2011) 071 [arXiv:1107.3554 [hep-th]].
 - [12] V. Kazakov and E. Sobko, JHEP **1306** (2013) 061 [arXiv:1212.6563 [hep-th]].
 - [13] The similar phenomenon was observed in [12]
 - [14] J. Bartels and M. Wusthoff, Z. Phys. C **66** (1995) 157.
 - [15] G. P. Korchemsky, Nucl. Phys. B **550** (1999) 397 [hep-ph/9711277].
 - [16] G. A. Chirilli, L. Szymanowski and S. Wallon, Phys. Rev. D **83** (2011) 014020 doi:10.1103/PhysRevD.83.014020 [arXiv:1010.0285 [hep-ph]].
 - [17] We thank S. Caron-Huot for the discussion on this subject.
 - [18] M. Alfimov, N. Gromov and V. Kazakov, arXiv:1408.2530 [hep-th].