

This is the accepted manuscript made available via CHORUS. The article has been published as:

Measurement of cosmic-ray electrons at TeV energies by VERITAS

A. Archer *et al.* (The VERITAS Collaboration)

Phys. Rev. D **98**, 062004 — Published 20 September 2018

DOI: [10.1103/PhysRevD.98.062004](https://doi.org/10.1103/PhysRevD.98.062004)

Measurement of Cosmic-ray Electrons at TeV Energies by VERITAS

A. Archer,¹ W. Benbow,² R. Bird,³ R. Brose,^{4,5} M. Buchovecky,³ J. H. Buckley,⁶
V. Bugaev,⁶ M. P. Connolly,⁷ W. Cui,^{1,8} M. K. Daniel,² Q. Feng,⁹ J. P. Finley,¹
L. Fortson,¹⁰ A. Furniss,¹¹ G. Gillanders,⁷ M. Hütten,⁵ D. Hanna,⁹ O. Hervet,¹² J. Holder,¹³
G. Hughes,² T. B. Humensky,¹⁴ C. A. Johnson,¹² P. Kaaret,¹⁵ P. Kar,¹⁶ N. Kelley-Hoskins,⁵
M. Kertzman,¹⁷ D. Kieda,¹⁶ M. Krause,⁵ F. Krennrich,¹⁸ S. Kumar,¹³ M. J. Lang,⁷
T. T.Y. Lin,⁹ G. Maier,⁵ S. McArthur,¹ P. Moriarty,⁷ R. Mukherjee,¹⁹ S. O'Brien,²⁰
R. A. Ong,³ A. N. Otte,²¹ A. Petrashyk,¹⁴ M. Pohl,^{4,5} E. Pueschel,⁵ J. Quinn,²⁰ K. Ragan,⁹
P. T. Reynolds,²² G. T. Richards,²¹ E. Roache,² C. Rulten,¹⁰ I. Sadeh,⁵ M. Santander,²³
G. H. Sembroski,¹ D. Staszak,^{24,*} I. Sushch,⁵ S. P. Wakely,²⁴ R. M. Wells,¹⁸
P. Wilcox,¹⁵ A. Wilhelm,^{4,5} D. A. Williams,¹² T. J. Williamson,¹³ and B. Zitzer^{9,†}

(The VERITAS Collaboration)

¹*Department of Physics and Astronomy,
Purdue University, West Lafayette, IN 47907, USA*

²*Fred Lawrence Whipple Observatory,
Harvard-Smithsonian Center for Astrophysics, Amado, AZ 85645, USA*

³*Department of Physics and Astronomy,
University of California, Los Angeles, CA 90095, USA*

⁴*Institute of Physics and Astronomy,
University of Potsdam, 14476 Potsdam-Golm, Germany*

⁵*DESY, Platanenallee 6, 15738 Zeuthen, Germany*

⁶*Department of Physics, Washington University, St. Louis, MO 63130, USA*

⁷*School of Physics, National University of Ireland Galway, University Road, Galway, Ireland*

⁸*Department of Physics and Center for Astrophysics,
Tsinghua University, Beijing 100084, China.*

⁹*Physics Department, McGill University, Montreal, QC H3A 2T8, Canada*

¹⁰*School of Physics and Astronomy,
University of Minnesota, Minneapolis, MN 55455, USA*

¹¹*Department of Physics, California State University - East Bay, Hayward, CA 94542, USA*

¹²*Santa Cruz Institute for Particle Physics and Department of Physics,
University of California, Santa Cruz, CA 95064, USA*

¹³*Department of Physics and Astronomy and the Bartol Research Institute,
University of Delaware, Newark, DE 19716, USA*

¹⁴*Physics Department, Columbia University, New York, NY 10027, USA*

¹⁵*Department of Physics and Astronomy, University of Iowa,
Van Allen Hall, Iowa City, IA 52242, USA*

¹⁶*Department of Physics and Astronomy,
University of Utah, Salt Lake City, UT 84112, USA*

¹⁷*Department of Physics and Astronomy,
DePauw University, Greencastle, IN 46135-0037, USA*

¹⁸*Department of Physics and Astronomy,
Iowa State University, Ames, IA 50011, USA*

¹⁹*Department of Physics and Astronomy,
Barnard College, Columbia University, NY 10027, USA*

²⁰*School of Physics, University College Dublin, Belfield, Dublin 4, Ireland*

²¹*School of Physics and Center for Relativistic Astrophysics,
Georgia Institute of Technology, 837 State Street NW, Atlanta, GA 30332-0430*

²²*Department of Physical Sciences, Cork Institute of Technology, Bishopstown, Cork, Ireland*

²³*Department of Physics and Astronomy,
University of Alabama, Tuscaloosa, AL 35487, USA*

²⁴*Enrico Fermi Institute, University of Chicago, Chicago, IL 60637, USA*

(Dated: August 23, 2018)

Abstract

Cosmic-ray electrons and positrons (CREs) at GeV-TeV energies are a unique probe of our local Galactic neighborhood. CREs lose energy rapidly via synchrotron radiation and inverse-Compton scattering processes while propagating within the Galaxy and these losses limit their propagation distance. For electrons with TeV energies, the limit is on the order of a kiloparsec. Within that distance there are only a few known astrophysical objects capable of accelerating electrons to such high energies. It is also possible that the CREs are the products of the annihilation or decay of heavy dark matter (DM) particles. VERITAS, an array of imaging air Cherenkov telescopes in southern Arizona, USA, is primarily utilized for gamma-ray astronomy, but also simultaneously collects CREs during all observations. We describe our methods of identifying CREs in VERITAS data and present an energy spectrum, extending from 300 GeV to 5 TeV, obtained from approximately 300 hours of observations. A single power-law fit is ruled out in VERITAS data. We find that the spectrum of CREs is consistent with a broken power law, with a break energy at $710 \pm 40_{stat} \pm 140_{syst}$ GeV.

68 INTRODUCTION

69 Despite constituting only a small fraction of the total cosmic-ray flux, cosmic-ray elec-
70 trons and positrons (CREs) provide an important and unique probe of our local Galactic
71 neighborhood. They rapidly lose energy while propagating in the Galaxy via synchrotron
72 radiation and inverse-Compton scattering processes. This limits the propagation distance
73 for TeV electrons to of order ~ 1 kpc [1] [2] [3] and implies that CREs at TeV energies can
74 provide constraints on local cosmic-ray accelerators and diffusion effects.

75 The *Fermi*-LAT [4] collaboration and the AMS-02 collaboration [5], have measured the
76 CRE spectrum up to energies of ~ 1 TeV. More recently, both the DAMPE [6] and CALET [7]
77 collaborations have measured the CRE spectrum to a few TeV with excellent energy reso-
78 lution. At higher energies these instruments run out of statistics due to the combination
79 of the steep CRE spectrum and their relatively small acceptances. Ground-based imaging
80 atmospheric Cherenkov telescopes (IACTs) can extend the CRE spectrum to higher energies
81 due to their large collection areas ($\sim 10^5$ m²). H.E.S.S. [8, 9] and MAGIC [10] have demon-
82 strated this ability and have measured the CRE spectrum up to energies of several TeV.
83 Their results agree with the space-based measurements, within systematic uncertainties, in
84 the energy range where the sensitivities overlap. The combined picture that has emerged is
85 one where the CRE spectrum can be described by a simple power law from ~ 10 GeV up
86 to just below ~ 1 TeV. At higher energy H.E.S.S. sees a spectral steepening¹ while MAGIC
87 data are consistent with a single power law up to ~ 3 TeV, although with larger statistical
88 uncertainties. The DAMPE and CALET data also see a break in the CRE spectrum at ~ 1
89 TeV [7] [6].

90 The inclusive CRE spectrum is understood to have contributions from both electrons
91 and positrons and some instruments are able to separate the two components. The energy
92 dependence of the positron fraction, $e^+/(e^+ + e^-)$, has been measured for energies greater
93 than 10 GeV by the HEAT [11], PAMELA [12], *Fermi*-LAT [13], and AMS-02 [14] collabo-
94 rations. The fraction is found to rise with increasing energy up to ~ 200 GeV, above which it
95 appears to flatten out. Positrons are believed to be produced mainly in interactions between
96 cosmic rays and interstellar gas and this results in a positron fraction that decreases with
97 energy. The unexpected increase could imply the existence of additional local sources such

¹ The H.E.S.S. collaboration has recently reported preliminary results, obtained with higher statistics over
an extended energy range, which support this trend.

as pulsars or supernovae remnants. A more exotic explanation for the excess would be the annihilation or decay of DM particles. More conventional methods such as an improved propagation model [15] or a better accounting of secondary production might explain the results. A full understanding of this situation will require detailed input about both the positron fraction and the CRE spectrum. Given that the CRE spectrum, including its behavior at TeV energies, is such an integral component, it is important for all instruments capable of making such measurements to contribute.

METHODS

VERITAS is an IACT instrument. In contrast to space-borne detectors like *Fermi*-LAT and AMS-02, IACTs do not measure the primary particles (electrons, photons, nuclei) directly. They detect the Cherenkov light generated by charged particles in air showers that result from the impact of the primaries on the upper atmosphere. As such, there is no information on the charge of the primary; electrons and positrons cannot be measured separately. Therefore, CREs in this work refer to the sum of positrons and electrons. Air showers caused by gamma rays and electrons are electromagnetic and are systematically different from the hadronic showers caused by cosmic-ray nuclei. Indeed, these differences are often exploited to separate weak gamma-ray signals from large hadronic backgrounds. However, showers due to electrons are essentially identical to those from gamma rays and there is no practical way to discriminate between these two types of primary particles.

In the following we will use gamma-hadron separation techniques to isolate a gamma-like signal, which will include the electron signal but also a background of gamma rays. To limit the background contribution we will exclude data from regions around known VHE gamma-ray sources. The Galactic plane, a known source of diffuse gamma radiation, is also excluded. There is a limited amount of extra-Galactic diffuse gamma radiation but recent measurements by *Fermi*-LAT [17] up to ~ 800 GeV show that it is of the same order of magnitude as that from discrete sources (mostly blazars) and is described by a power law, with index of about 2.3, and an exponential cut-off at 250 GeV. Based on this *Fermi*-LAT result, we estimate perhaps one out of a thousand CRE-like events is actually a diffuse gamma ray at 250 GeV. This ratio falls off rapidly at higher energies. Contamination from the Fermi bubbles [18] is also possible and we did not explicitly exclude data in those regions.

The spectrum of the Fermi bubbles show an exponential cutoff at ~ 110 GeV, so it is unlikely to affect the results in this work. We will therefore assume in this work that the gamma-ray contamination in our CRE candidate events is negligible.

Separation between hadronic showers and gamma or electron-like showers is not fully efficient. After the application of data selection criteria (cuts) a substantial hadronic background remains, often larger than the signal itself. When measuring gamma-ray sources this background can be determined and subtracted using one of the techniques described in Berge et. al. 2007 [19]. These techniques rely on the presence of a source, known or postulated, in the FOV, surrounded by a source-free region from which to estimate the background. For CREs this is not the case; the intensity is essentially isotropic so there is no way to experimentally estimate the background and one must rely on simulation-based calculations. This introduces additional systematic uncertainties into the CRE estimate.

VERITAS

Data presented here were collected with the VERITAS array, located at the Fred Lawrence Whipple Observatory (FLWO) in southern Arizona, USA ($31^\circ 40'$ N, $110^\circ 57'$ W, 1.3 km a.s.l.). The array comprises four identical telescopes [20], each with a 12-m Davies-Cotton reflector that collects Cherenkov light and directs it onto a camera made up of 499 photomultiplier tubes (PMTs). The telescope FOV has a diameter of 3.5° .

Outputs from the PMTs are continuously digitized by 500 Msps 8-bit flash analog-to-digital converters (FADCs) and written to a buffer. A three-level trigger, requiring a three-fold coincidence of neighboring pixels, each of which passes a discriminator threshold in at least two telescopes, initiates readout of the digitized signals.

VERITAS has been running with four telescopes since 2007. During the summer of 2009 we repositioned one of the telescopes to improve the array sensitivity and in summer 2012 we installed PMTs with higher quantum efficiency to lower the energy threshold [21]. For this study we use data acquired between September 2009 and July 2012, with a narrow zenith angle range. The choice of this period was motivated by the fact that at the time this study started, this was the period of the largest and best-understood data set available. During this period VERITAS had a gamma-ray energy threshold of approximately 100 GeV, an energy resolution of 15-20% between 0.1 and 10 TeV, and an angular resolution (68%

containment) of less than 0.1° at 1 TeV [22]. CRE events have identical resolutions to gamma rays.

Data Selection

Since the CRE flux is isotropic we can use data from all VERITAS observations in our study, subject to the following selection criteria:

- clear weather,
- all telescopes operational,
- zenith angle between 15° and 25° ,
- observation field at least 10° away from the Galactic plane.

These cuts resulted in a data set constituting 296 hours of live time. The range of zenith angles between 15 and 25 degrees is the window containing the most data available. Additionally, the best sensitivity of VERITAS is for zenith angles less than 25 degrees [22]. The restricted zenith angle range and single detector configuration corresponds to where most VERITAS observations were made at the time this study started. Using the restricted range greatly reduced the number of simulations required for this study. The data were processed using standard VERITAS reconstruction software and the resulting events were subjected to further cuts:

- Every telescope should have a good shower image according to the standard VERITAS reconstruction package [29]. This reduces the overall systematic uncertainties in both the energy and position reconstruction of the shower. Above 1 TeV, 95% of simulated CRE events have good shower images in all four telescopes.
- The direction of the reconstructed shower should point to within a degree of the nominal array pointing direction. VERITAS has peak sensitivity using events within one degree of the pointing direction [22].
- The shower axis should intersect the array plane within 200 m of the array center. Roughly 90% of simulated CRE events reconstruct at distances less than this, therefore this cut only removes a small number of potential signal events from the analysis.

- For observation fields that contain known or potential gamma-ray sources, the shower axis should not point within 0.187° of those objects. (The actual cut is on the square of this value, 0.035 deg^2 .) This is wider than the gamma-ray PSF and should remove any significant gamma-ray contamination in the CRE data. The list of known gamma-ray sources was obtained from TeVCat [30]. Potential gamma-ray sources that were the *a priori* targets of VERITAS observations were also excluded.

Some of these cuts are more restrictive than in most VERITAS gamma-ray analyses, resulting in improved data/Monte Carlo agreement with the proton simulations. The cuts were optimized to improve the agreement of diffuse proton simulations with proton data at both the single-telescope image and array-wide levels, and to improve the agreement of gamma-ray simulations with Crab Nebula data. The goal of applying these additional cuts was to reduce the overall systematic uncertainty in the CRE measurement.

Monte Carlo Simulations

As explained previously, extracting a CRE signal requires extensive use of Monte Carlo (MC) simulations. Showers initiated by electrons, protons, and helium nuclei were generated, with arrival directions up to 4° away from the array pointing direction. This range was to cover the VERITAS FOV and it accommodates any edge effects; adding showers from larger angles had no effect. Each particle species was generated with an energy distribution of $dN/dE \sim E^\Gamma$ with $\Gamma = -2$ to increase statistics at higher energies. The proton energy distribution was then reweighed to $\Gamma = -2.7$ to agree with experimental data over the range we are sensitive to [23].

The CORSIKA 6.970 package [24] was used, with the QGSJetII.3 [25] and URQMD 1.3cr [26, 27] event generators, to produce files of ground-level Cherenkov photons. Newer versions of CORSIKA have significantly improved simulations of hadronic interactions at the highest energies, but the energy range here is relatively unaffected. These were processed with the GrISUDet 5.0.0 [28] VERITAS detector modeling program before analysis with the one of the standard VERITAS event-reconstruction packages (EventDisplay [29]). A smaller set of proton-initiated showers was generated using the SIBYLL package [31] and used to test for consistency. Results agreed within systematic uncertainties with those from QGSJetII.3.

Cherenkov light from an electromagnetic (e.g. gamma-initiated) shower produces an approximately elliptical image in the camera plane of an IACT. Images from hadronic (proton-initiated) showers are less regular in shape and can have large fluctuations in their morphology. This forms the basis for all schemes of background rejection in IACT-based gamma-ray astronomy [32]. Traditionally, one characterizes the images using a set of parameters suggested by Hillas [33]. Two of these are the length and width of the ellipse. Images from hadronic showers are typically longer and wider than those from electromagnetic showers.

The standard VERITAS analysis packages make use of image parameters and rely on cuts placed separately on each of them to extract signals; these are called “box cuts”. In this work we make use of boosted decision trees (BDTs) to reject (hadronic) background and retain (electromagnetic) signal. As explained in [34], BDTs are a multivariate analysis technique. They combine several input variables in such a way as to produce a single output variable which describes, in this application, how “electromagnetic-like” a shower is. The BDTs are trained with a combination of data and simulations and provide the ability to utilize non-linear correlations between training variables when classifying data. They have been shown [34] to improve VERITAS sensitivity for a variety of gamma-ray source types compared with the traditional box-cut analyses. Apart from the improved sensitivity offered by BDTs, they are useful because their output is a single number that can be used in likelihood fitting procedures, as shown later in this paper. These methods are essentially the same as those used by the H.E.S.S. and MAGIC collaborations [8–10] for CRE spectra, except for utilizing BDTs instead of Random Forests for hadron discrimination.

The BDTs used in this work were constructed using the Toolkit for Multivariate Data Analysis (TMVA) component [35] of the CERN ROOT package [36]. We used four array-level variables as input to the BDTs:

- mean reduced scaled width (MRSW), [37]
- mean reduced scaled length (MRSL), [37]
- emission height, and
- $\chi^2(E)$, which is defined as the RMS of the energy estimates from the individual telescopes. The mean of the individual telescope energy estimates is accepted as the

reconstructed shower energy.

These variables are described in more detail in [34]. The first two are related to the lengths and widths of the camera images. The third variable is the estimate of the height of maximum development of the shower, made using data from all telescopes. On average, electromagnetic showers have a slightly lower value for this than do hadronic showers. The fourth quantifies how well the different telescopes agree on the energy of the shower. $\chi^2(E)$ tends to be larger for hadronic showers.

The BDTs were trained with a signal data set of 6.4 million simulated diffuse electron events and a background data set consisting of 5.7 million events chosen randomly from the full data set. Both the signal and training samples pass the cuts described in the previous section. Note that the “background” events, chosen in this way, contain an admixture of “signal” events and this compromises the training somewhat. However, since CREs make up a small fraction of the total cosmic-ray flux, this is a higher-order effect. The use of data rather than simulated hadronic showers requires fewer CPU hours and is less dependent on the fidelity of the hadronic simulations.

Independent signal and background data sets were used to check for over-training of the BDTs; no evidence for this was found. The BDTs were also tested by using them for gamma-hadron separation in an analysis of Crab Nebula data. The reconstructed energy spectrum was consistent with previous measurements.

After training, the BDTs were used in classifying events from the data set used for this study. Each shower was assigned a BDT response value, from -1.0 to 1.0 , indicating whether the shower was hadronic (-1.0) or electromagnetic (1.0). Figure 1 shows a plot of BDT responses for the full data set compared with those from a set of simulated proton showers. The agreement is very good except near the limits of the distribution. There are excesses of data over proton simulations near -1.0 and near 1.0 . At -1.0 we expect an excess from helium and higher- Z primaries, particularly since helium makes up $\sim 20\%$ of the overall cosmic-ray flux [39]. We investigated the BDT response for a set of simulated helium events and found it to peak at -1.0 and fall off faster than that for simulated protons, supporting our interpretation of the excess. There is a second excess at the opposite end of the response histogram. This is the CRE signal, which is present in the observational data but not the hadronic background simulations.




FIG. 1. CRE BDT response parameter for the full data set (symbols) and for simulated proton-induced showers (histogram) in the energy range between 630 GeV and 1 TeV. The plot in the inset is the ratio of the two. Hadronic showers are assigned BDT values close to -1.0 while electromagnetic showers are assigned values near 1.0.

In the following, we select events with BDT response values greater than 0.7 and employ a binned extended likelihood fitting method within this region to extract the contributions of electron and proton events to the total. Using the shape of the BDT response distribution from electron and proton simulations, we find the relative contributions that best describe the data. We do not include contributions from helium and higher-Z shower events since they are sufficiently rejected in this region by the BDTs.

RESULTS

We show in Figure 2 the VERITAS CRE energy spectrum between 300 GeV and 5 TeV. These results are also summarized in Table 1. The spectrum steepens at higher energies and is best described by two power laws, with a break between them as one index transitions to

another. The best fit for this break energy is found to be $(710 \pm 40_{stat} \pm 140_{syst})$ GeV, with best-fit spectral indices below (above) this energy of $-3.2 \pm 0.1_{stat}$ ($-4.1 \pm 0.1_{stat}$). The chi-squared per degree-of-freedom (χ^2/dof) of this fit is 9.71/11. Analytically, this fit is described by:

$$F(E) = \begin{cases} k(E/E_b)^{\Gamma_1} & \text{if } E \leq E_b \\ k(E/E_b)^{\Gamma_2} & \text{if } E > E_b \end{cases}, \quad (1)$$

where E_b is the break energy, k is the flux value at the break energy and Γ_1 (Γ_2) is the index below (above) the break energy.

In addition to a broken power law, a single power-law fit was performed, giving a significantly worse fit to the data, with a χ^2/dof of 76.5/13. A power law with an exponential cutoff was also fit to the data, yielding a χ^2/dof of 26.0/12.

The gray band represents the systematic uncertainty, which is dominated by the $\sim 20\%$ uncertainty on the VERITAS absolute energy scale. This translates into a $+64\%/ -33\%$ ($+98\%/ -43\%$) systematic uncertainty for a spectral index of -3.2 (-4.1).

Additional tests on the robustness of the analysis were performed:

- the assumed spectral index for the energies of the simulated protons was varied by $\pm 10\%$ from its nominal value,
- BDT cut values 0.8 and 0.9 were investigated and
- Hadronic-shower rejection was performed using cuts on the MRSW distributions and not the BDT response, effectively testing the entire machinery but not using BDTs.

All changes resulted in final data points consistent with the quoted systematic uncertainties. It should be noted that the proton spectrum is well-measured in this energy range, so 10% is considered very conservative. For the non-BDT analysis a loss of efficiency occurred, resulting in larger statistical uncertainties, and energy bins above 2 TeV could not be recovered.

CONCLUSION

CRE results shown here are consistent with prior ground-based and space-borne measurements at similar energies. This result represents the second ground-based, high-statistics




FIG. 2. Spectrum of CREs between 300 GeV and 5 TeV, as measured by VERITAS along with previously published measurements. Error bars are statistical; systematic uncertainties are indicated by the gray band.

measurement of a break in the CRE spectrum around ~ 1 TeV, seen by H.E.S.S., CALET and DAMPE, but not seen by MAGIC or *Fermi*-LAT. The precise value of this break energy is an important parameter in any successful model of our local CRE environment. Based on the fit of the CRE spectrum in the previous section, two power laws with a break between the indices best describes the data. However, a power law with an exponential cutoff is not completely ruled out. A single power law is ruled out by the VERITAS data.

Several different sources in our local neighborhood have been speculated as the accelerators of electrons at TeV energies, including supernova remnants and pulsars. The decay or annihilation of WIMP DM has also been proposed as the dominant source of CREs and a reason for the positron fraction rising with energy, [38] but it is currently not possible

Energy [TeV]	Energy Bin [TeV]	N_{events}	Electron Fraction	Flux [$\text{cm}^{-2}\text{s}^{-1}\text{TeV}^{-1}$]
0.335	0.316 - 0.355	27529	0.152 ± 0.005	$3.140 \times 10^{-10} \pm 9.790 \times 10^{-12}$
0.376	0.355 - 0.398	21330	0.159 ± 0.006	$2.104 \times 10^{-10} \pm 8.469 \times 10^{-12}$
0.422	0.398 - 0.447	15704	0.187 ± 0.007	$1.491 \times 10^{-10} \pm 5.681 \times 10^{-12}$
0.473	0.447 - 0.501	14787	0.154 ± 0.007	$9.761 \times 10^{-11} \pm 4.728 \times 10^{-12}$
0.531	0.501 - 0.562	11823	0.169 ± 0.007	$7.168 \times 10^{-11} \pm 3.120 \times 10^{-12}$
0.596	0.562 - 0.631	8476	0.175 ± 0.010	$4.480 \times 10^{-11} \pm 2.513 \times 10^{-12}$
0.668	0.631 - 0.708	6480	0.208 ± 0.008	$3.524 \times 10^{-11} \pm 1.372 \times 10^{-12}$
0.749	0.708 - 0.794	5932	0.170 ± 0.009	$2.333 \times 10^{-11} \pm 1.168 \times 10^{-12}$
0.840	0.794 - 0.891	4146	0.164 ± 0.009	$1.376 \times 10^{-11} \pm 7.186 \times 10^{-13}$
0.995	0.891 - 1.122	5547	0.140 ± 0.010	$6.370 \times 10^{-12} \pm 4.700 \times 10^{-13}$
1.253	1.122 - 1.413	2951	0.141 ± 0.014	$2.689 \times 10^{-12} \pm 2.721 \times 10^{-13}$
1.662	1.413 - 1.995	2495	0.107 ± 0.013	$8.620 \times 10^{-13} \pm 1.058 \times 10^{-13}$
2.347	1.995 - 2.818	892	0.111 ± 0.013	$2.233 \times 10^{-13} \pm 3.015 \times 10^{-14}$
3.315	2.818 - 3.981	529	0.055 ± 0.014	$4.645 \times 10^{-14} \pm 1.216 \times 10^{-14}$
4.683	3.981 - 5.623	207	0.167 ± 0.082	$3.621 \times 10^{-14} \pm 1.780 \times 10^{-14}$

TABLE I. CRE Flux with statistical errors. The total number of events in each energy bin and fraction of those events that are CRE-like from the likelihood fit are also given.

to discriminate dark matter models from other sources with the available data [2]. Nearby pulsars with distances less than 1 kpc may also be sources of relativistic electrons and positrons [1][3]. Because of synchrotron and inverse-Compton energy losses, the age of TeV electrons is $\sim 10^5$ years, and decreases with increasing energy. Very few of the known pulsars are capable of accelerating electrons to TeV-scale energies, namely Geminga, Monogem and a handful of others. Breaks in the spectrum at TeV energies are expected as the number of astrophysical sources capable of accelerating CREs to those energies decreases [2]. Refined measurements of CRE spectra from IACTs, including VERITAS and the upcoming CTA observatory [40], should help with understanding the number and distribution of sources capable of accelerating CREs to TeV-scale energies.

ACKNOWLEDGMENTS

This research is supported by grants from the U.S. Department of Energy Office of Science, the U.S. National Science Foundation and the Smithsonian Institution, and by NSERC in Canada. We acknowledge the excellent work of the technical support staff at the Fred Lawrence Whipple Observatory and at the collaborating institutions in the construction and operation of the instrument. Computations were made on the supercomputer Guillimin from McGill University, managed by Calcul Quebec and Compute Canada. The operation of this supercomputer is funded by the Canada Foundation for Innovation (CFI), NanoQuebec, RMGA and the Fonds de recherche du Quebec - Nature et technologies (FRQ-NT).

341 * d.staszak@gmail.com
 342 † bzitzer@physics.mcgill.ca
 343 [1] R. Cowsik et al., *Astrophys. J.*, , 1979
 344 [2] D. Grasso, et al. *Astroparticle Physics*, 32, 140, 2009
 345 [3] M. Pohl, et al. *ApJ*, 507, 327 (1998)
 346 [4] S. Abdollahi et al., *Phys. Rev. D*, 95 (082007), 2017
 347 [5] M. Aguilar et al., *Phys. Rev. Lett.*, 113 (221102), 2014
 348 [6] G. Ambrosi et al., *Nature*, 552, (63), 2017
 349 [7] O. Adriani et al., *Phys. Rev. Lett.*, 119 (181101), 2017
 350 [8] F. Aharonian et al. *Phys. Rev. Lett.*, 101 (261), 2008
 351 [9] F. Aharonian et al. *Astron. Astrophys.*, 508 (561), 2009
 352 [10] D. Borla Tridon et al. Proc. 32nd ICRC, 2011
 353 [11] J. J. Beatty et al., *Phys. Rev Lett.*, 93 (241102), 2004
 354 [12] O. Adriani et al. *Nature* 458 (607), 2009
 355 [13] M. Ackermann et al. *Phys. Rev Lett.*, 108 (011103), 2012
 356 [14] M. Aguilar et al., *Phys. Rev. Lett.*, 113 (121101), 2014
 357 [15] D. Gaggero et al., *Phys. Rev. Lett.*, 101 (021102), 2013
 358 [16] K. Blum et al., *Phys. Rev. Lett.*, 111 (211101), 2013
 359 [17] M. Ackermann et al., *Astrophys. J.*, 799 (86), 2015
 360 [18] M. Ackermann et al. *Astrophys. J.*, 793, (64) 2014
 361 [19] D. Berge, S. Funk, and J. Hinton, *Astron. Astrophys.*, 466 (1219), 2007
 362 [20] J. Holder et al., *Astropart.Phys.* 25 (391),2006
 363 [21] D.B. Kieda et al. Proc. 32nd ICRC, 2011
 364 [22] N. Park et al. Proc. 34th ICRC, 2015
 365 [23] D. Maurin et al. *Astron. Astrophys.*, 569 (A32), 2014
 366 [24] <https://www.ikp.kit.edu/corsika/70.php>
 367 [25] S. Ostapchenko, *Nucl. Phys. B - Proc. Suppl.*, 151 (143), 2005
 368 [26] S. A. Bass et al., *Prog. Part. Nucl. Phys.* 41 (225), 1998
 369 [27] M. Bleicher et al. *J. Phys. G: Nucl. Part. Phys.* 25 (1859), 1999

370 [28] <http://www.physics.utah.edu/gammaray/GrISU/>

371 [29] <https://arxiv.org/abs/1708.04048>

372 [30] <http://tevcat.uchicago.edu/>

373 [31] E.-J. Ahn et al., *Phys. Rev. D*, 80 (094003), 2009

374 [32] D. Fegan, *Journal of Physics G* 23, (9) 1997

375 [33] A. Hillas, Proc. 19th ICRC, 1985

376 [34] M. Krause, E. Pueschel, and G. Maier, *Astropart. Phys*, 89 (1), 2017

377 [35] A. Hoeker et al., PoS ACAT, 040 2007

378 [36] R. Brun and F. Rademakers, *Nucl. Inst. & Meth. in Phys. Res. A*, 389 (81) 1997

379 [37] H. Krawczynski, et al. *Astroparticle Physics*, 25, 380, 2006

380 [38] G. Bertone, et al. *Phys. Reports* , 405, (279) 2005

381 [39] C. Patrignani et al. (Particle Data Group), *Chin. Phys. C*, 40, (100001) 2016

382 [40] <https://www.cta-observatory.org/>