



This is the accepted manuscript made available via CHORUS. The article has been published as:

Phase chimera states on nonlocal hyperrings

Riccardo Muolo, Thierry Njougouo, Lucia Valentina Gambuzza, Timoteo Carletti, and
Mattia Frasca

Phys. Rev. E **109**, L022201 — Published 12 February 2024

DOI: [10.1103/PhysRevE.109.L022201](https://doi.org/10.1103/PhysRevE.109.L022201)

Phase chimera states on non-local hyperrings

Riccardo Muolo,^{1,2,3,*} Thierry Njougouo,^{3,4,5} Lucia Valentina Gambuzza,⁶ Timoteo Carletti,^{2,3} and Mattia Frasca^{6,7}

¹*Department of Systems and Control Engineering, Tokyo Institute of Technology, Tokyo 152-8552, Japan*

²*Department of Mathematics, University of Namur, B5000 Namur, Belgium*

³*naXys, Namur Institute for Complex Systems, University of Namur, B5000 Namur, Belgium*

⁴*Faculty of Computer Science, University of Namur, B5000 Namur, Belgium*

⁵*Department of Electrical and Electronic Engineering, University of Buea, P.O. Box 63, Buea, Cameroon*

⁶*Department of Electrical, Electronics and Computer Science Engineering, University of Catania, 95125 Catania, Italy*

⁷*Istituto di Analisi dei Sistemi ed Informatica “A. Ruberti”, IASI-CNR, 00185 Roma, Italy*

(Dated: January 18, 2024)

Chimera states are dynamical states where regions of synchronous trajectories coexist with incoherent ones. A significant amount of research has been devoted to study chimera states in systems of identical oscillators, non-locally coupled through pairwise interactions. Nevertheless, there is an increasing evidence, also supported by available data, that complex systems are composed by multiple units experiencing many-body interactions, that can be modeled by using higher-order structures beyond the paradigm of classic pairwise networks. In this work we investigate whether phase chimera states appear in this framework, by focusing on a novel topology solely involving many-body, non-local and non-regular interactions, hereby named *non-local d-hyperring*, being $(d + 1)$ the order of the interactions. We present the theory by using the paradigmatic Stuart-Landau oscillators as node dynamics, and show that phase chimera states emerge in a variety of structures and with different coupling functions. For comparison, we show that, when higher-order interactions are “flattened” to pairwise ones, the chimera behavior is weaker and more elusive.

Introduction. Chimera states are an intriguing dynamical phenomenon occurring in systems of coupled oscillators where some units oscillate synchronously, forming coherent domains that exist alongside other domains characterized by incoherent oscillations. Since their first numerical observation by Kaneko [1], the emergence of these patterns has raised the interest of many scholars in nonlinear science. In later studies, these patterns have been identified in experimental setups involving Josephson junctions [2], laser systems [3], mechanical systems [4], electronic circuits [5, 6], nano-electromechanical oscillators [7], neuroscience [8], unihemispheric sleep in birds, mammals, and reptiles [9], as well as various forms of pathological brain states [10]. They have been also found in numerical experiments involving systems of non-locally coupled oscillators, such as Ginzburg-Landau systems, Rössler oscillators, logistic maps [11–15], and identical phase oscillators [16], prior to being named “Chimera states” in [17].

Chimera states have been extensively investigated over the past years, leading to exploration along various intriguing avenues [18, 19]. From a theoretical point of view, there has been considerable effort to characterize the different types of chimera states that may appear: from phase chimeras [16] to amplitude chimeras [20], amplitude mediated chimeras [21], chimera death states [22], traveling chimeras [23], and globally clustered chimeras [24, 25]. Let us observe that the recently defined phase chimera [26] differ from the ones presented in [16], despite bearing the same name. Indeed, by writing the dynamics of the j -th oscillator as $a_j(t)\exp[i(2\pi\Omega_j t + \theta_j)]$, Kuramoto and Battogtokh [16] defined a phase chimera as a state where $a_j(t)$ is constant for all j while the angles $\phi_j = 2\pi\Omega_j t + \theta_j$ can be divided into

two groups, one for which Ω_j is constant and a second one for which Ω_j depends on the node index. Both behaviors can be appreciated after a sufficiently large t . On the other hand, Zajdela and Abrams [26] defined a phase chimera as a state for which both $a_j(t)$ and Ω_j are constant, i.e., do not depend on the node index, and only the phases θ_j exhibit coherent and incoherent behaviors. The latter is the framework we will hereby consider.

Another avenue of research on chimera states has focused on understanding the network topologies and coupling mechanisms that can facilitate their emergence. Although chimera states have been proved to arise in 1D-rings where every node is connected to its two neighbors (a condition named *local coupling* [27, 28]) as well in the “opposite” case of a complete network where each node is connected to all the other ones (*global coupling* [29, 30]), the *non-local coupling* configuration proved to be particularly important for the onset of chimera states. Indeed, a large amount of literature has demonstrated that chimera states are commonly observed within such setting [16, 19]. The non-local coupling corresponds to a 1D-ring where, beside first order neighbors, each node is also connected in a regular way to $2(k - 1) \geq 2$ other ones, i.e., each node is connected to k nearby nodes counted in anti-clockwise manner and k in the other direction. The network is regular by construction, all the nodes having the same degree $2k$, and it has a complete group of automorphisms, i.e., it is invariant for any possible translation. Beside those regular cases, chimera states have been also found on networks with diverse non-regular topologies [31–36]. Nonetheless, the identification of chimera states has been proven to be challenging, given their transient nature [37] and strong dependence on the initial conditions [38]: hence, researchers have also endeavored to propose mechanisms that can lead to the emergence of robust and persistent chimeras [39, 40].

While there have been numerous studies focusing on

* corresponding author: muolo.raa@m.titech.ac.jp

chimera states within network structures, these states have received significantly less attention in systems where the units interact not only in pairs, but also in groups of three or more units. Studying chimera states in the presence of these types of interactions, known as higher-order or many-body [41, 42], is particularly important. In fact, on the one hand, there is an increasing evidence that many systems, e.g., in neuroscience [43–46], naturally exhibit higher-order interactions. On the other hand, several studies on dynamics emerging in hypergraphs and simplicial complexes have already shown a significant impact of higher-order interactions on the collective behavior of the system; the latter include, e.g., studies on synchronization and pattern formation [47–54], random walks [55, 56] and contagion processes [57–61]. Previous works on chimera states in higher-order structures have already lead to several interesting results [7, 62–65]. Chimera states have been found experimentally by considering triadic interactions [7] and, theoretically, while developing a general theory for the study of synchronization patterns in higher-order systems [62, 63]; despite dealing with small hypergraphs and specific settings, these works have pointed out that chimera states may also appear in the presence of higher-order interactions. A more systematic study of the impact of higher-order interactions has been done in [64, 65], where phase oscillators coupled through simplicial complexes are studied; both works show that higher-order interactions, when added to pairwise ones, enhance the likelihood of observing chimera states in the system.

In this work, we further corroborate the claim that chimera states are boosted by the presence of many-body interactions by focusing on a higher-order structure that we call *non-local hyperring* and consists exclusively of higher-order interactions. This is an important difference with respect to the setup studied, for instance, in [65], where simplicial complexes are considered, or in [66], where a generalization of the non-local coupling is extended to higher-order interactions through a continuous limit [67]: in both settings, the effects of pairwise and three-body interactions cannot be fully disentangled. The topology that we consider is non-local and non-regular, i.e., nodes have different hyper degree. We consider a d -uniform hypergraph, where all the hyperedges have the same size, d , and are connected two-by-two, in a periodic structure, by junctions nodes shared by every two consecutive hyperedges. In this way, we have a non-regular extension of the notion of non-local rings to higher-order structures. Since non-local hyperrings only include higher-order interactions, the effect of the latter on the emergence of chimera states can be better identified. Let us observe that, despite the lack of regular topology, the studied systems always admit a global synchronous solution, that thus “competes” with the possible chimera state. For the sake of definiteness, we numerically investigate the behavior of non-local hyperrings when the nodal dynamics is given by the Stuart-Landau oscillator, which has been widely used as a paradigmatic model of oscillatory dynamics to study chimera states [18], given that this system is the normal form of the Hopf-Andronov bifurcation and thus it presents the general features of any limit-cycle oscillator close to such bifurcation [68]. We then com-

pare the results with the ones observed on a pairwise network, obtained by projecting the higher-order structure on a pairwise network: namely, we consider that every node has a non-weighted pairwise connection with all the nodes part of the same hyperedge, i.e., they form a clique. The higher-order setting exhibits a considerable enhancement of chimera patterns, meaning that they are present for a wider range of coupling strengths and they have a longer life span. Finally, we show that not only phase chimera states, but also other interesting dynamical patterns, such as amplitude chimeras [69], can be observed in the novel setting.

Stuart-Landau oscillators on non-local hyperrings. We consider a system made of n interacting Stuart-Landau units, a paradigmatic model of oscillatory dynamics. In the absence of any interaction, each unit j ($j = 1, \dots, n$) of the system is described by the following equations

$$\begin{cases} \dot{x}_j = \lambda x_j - \omega y_j - (x_j^2 + y_j^2) x_j = f(x_j, y_j), \\ \dot{y}_j = \omega x_j + \lambda y_j - (x_j^2 + y_j^2) y_j = g(x_j, y_j), \end{cases} \quad (1)$$

where λ is a bifurcation parameter controlling the onset of a limit cycle of amplitude $\sqrt{\lambda}$, for $\lambda > 0$, and ω is the natural frequency of the oscillator. Let us observe that we assume those parameters to be same for all nodes to ensure the existence of a global synchronous solution.

Here, as in [47, 70], we study nonlinear many-body coupling functions that cannot be decomposed into a combination of weighted pairwise interactions [71]. To model the higher-order interactions we use hyperedges, whose structure can be encoded by using adjacency tensors, that are a generalization of the adjacency matrix for networks [41]. We adopt the convention that a hyperedge involving $(d+1)$ nodes (and, thus, encoding a $(d+1)$ -body interaction) is called a d -hyperedge. Such notation is more common in the literature dealing with simplicial complexes [72], while, in the context of hypergraphs, often a d -hyperedge encodes a d -body interaction [41]. For example, $A^{(3)} = \{A_{i,j,k,l}^{(3)}\}$ is the 3-rd order adjacency tensor, encoding the 4-body interactions, with $A_{i,j,k,l}^{(3)} = 1$ if units i, j, k, l have a group interaction (namely, nodes i, j, k, l are part of the same 3-hyperedge), and 0 otherwise. Using these tensors, the generalized d -degree (or hyperdegree), $k_j^{(d)}$, representing the number of d -hyperedges of which node j is part, can be computed as

$$k_j^{(d)} = \frac{1}{d!} \sum_{j_1, \dots, j_d=1}^N A_{jj_1 \dots j_d}^{(d)}. \quad (2)$$

Finally, we assume that the coupling terms act on both the dynamical variables describing the Stuart-Landau oscillator (1) and involve only the first component of the state vector of the oscillator, as in [6]; let us observe that this working assumption is not restrictive and other couplings can also be considered. Taking into account all these considerations, the model of Stuart-Landau equations coupled via a generic $(d+1)$ -body

interactions reads:

$$\begin{cases} \dot{x}_j = f(x_j, y_j) + \epsilon \sum_{j_1, \dots, j_d=1}^n A_{j, j_1, \dots, j_d}^{(d)} \left(h^{(d)}(x_{j_1}, \dots, x_{j_d}) - h^{(d)}(x_j, \dots, x_j) \right), \\ \dot{y}_j = g(x_j, y_j) + \epsilon \sum_{j_1, \dots, j_d=1}^n A_{j, j_1, \dots, j_d}^{(d)} \left(h^{(d)}(x_{j_1}, \dots, x_{j_d}) - h^{(d)}(x_j, \dots, x_j) \right), \end{cases} \quad (3)$$

where $\epsilon > 0$ is the coupling strength and we assumed that the coupling is diffusive-like [47]. The model encompasses only multi-body interactions (specifically, $(d+1)$ -body interactions encoded in the d -th order adjacency tensor $A^{(d)}$), as we aim to analyze a pure many-body framework.

Let us observe that higher-order interactions are often considered together with pairwise interactions [64, 65]; however, in this setting it is not clear how to distinguish the different types of contributions. For this reason, we decided to consider hypergraphs instead of simplicial complexes, as the latter structures contain interactions of any order, from the highest to the lowest (i.e., order one, which is the pairwise case). Moreover, to the best of our knowledge, structures representing the higher-order counterpart of non-local rings have not yet been considered in previous works on chimera states. Our goal is to fill this gap by considering a ring encompassing m hyperedges of order d . These hyperedges are placed on a circle and labeled in counterclockwise order as $1, 2, \dots, m$. In the simplest case, the m hyperedges share with each other only a single node. Furthermore, each shared node may belong at most to two different hyperedges, meaning that the generic hyperedge j has two shared nodes, one with the hyperedge $(j-1)$ and one with the hyperedge $(j+1)$. The remaining $d-2$ nodes are only part of the hyperedge j . Let us observe that more general structures could be handled as well.




FIG. 1. **Non-local hyperring vs. clique-projected network.** a) A non-local 3-hyperring with $m = 5$ hyperedges and 15 nodes. b) Corresponding clique-projected network obtained by transforming each hyperedge into a clique.

In the following, we will focus on the case of 4-body interactions, but similar results have been obtained for 3-body,

as well as for 5-body and 6-body interactions (see Supplementary Material (SM)). In our model, the coupling functions have been chosen to be cubic and diffusive-like, as previously done in [47], namely

$$h^{(3)}(x_{j_1}, x_{j_2}, x_{j_3}) = x_{j_1} x_{j_2} x_{j_3}. \quad (4)$$

In the SM, we illustrate other choices, i.e., diffusive-like functions that are not cubic. In this specific case, an example of non-local ring with $m = 5$ hyperedges of size 4 is shown in panel a) of Fig. 1. The structure is non-regular, as some nodes are part of two hyperedges, i.e., they have hyperdegree 2 (see Eq. (2)), while others interact only within a single hyperedge, and thus exhibit hyperdegree 1. In general, for a non-local ring with d -body interactions and n nodes, the nodes that are shared by two hyperedges, i.e., nodes $\{1, 1 + (d-1), \dots, n - d - (d-3), n - (d-2)\}$ have generalized d -th degree [73], equal to 2, while all the other nodes have generalized d -th degree equal to 1. In the following, we will be interested in comparing the dynamical behavior of SL oscillators coupled via higher-order structures with the one resulting from a pairwise one. To have a reliable comparison, we decide to work with the network obtained by projecting each hyperedge in a clique, as shown in panel b) of Fig. 1, to obtain the so called clique-projected network (cpn). Namely, each couple of nodes in the hyperedge is connected with an unweighted pairwise link. In this way, we end up with d -cliques connected through junction nodes, that are part of two consecutive cliques, exactly as the hyperedges are disposed in the hyperring. Let us point out that this is not the only possible projection onto a network, as one could also consider projection based on 1-cell complexes, which are topological objects obtained by gluing together cells in different ways. In this sense, they can be viewed as generalizations of simplicial complexes [72] that have recently found applications in several field, from machine learning [74] to synchronization dynamics [75]. In the SM we have analysed a different projection, specifically, mapping any d -hyperedge as a regular polygon having $(d+1)$ vertices. When the topology is pairwise, the coupling functions take only one variable, we thus define $h^{cpn}(x_j) = h^{(3)}(x_{j_1}, x_{j_2}, x_{j_3})|_{x_{j_1}=x_{j_2}=x_{j_3}=x_j}$. In our example of 4-body interactions, the coupling functions (4) can be replaced on the clique-projected networks (cpn) by

$$h^{cpn}(x_\ell) - h^{cpn}(x_j) = x_\ell^3 - x_j^3.$$

To make the difference between the pairwise and higher-order coupling more visible, in the SM, we explicitly write the equations for a node shared between two hyperedges (junction node) and a node belonging to only one hyperedge (non-junction node).

Higher-order vs. pairwise interactions. We now proceed to the analysis of chimera states observed in the non-local hyperring, while also comparing them to the pairwise case. In all the simulations, we have considered hyperrings and networks of $n = 204$ nodes, hence $m = 68$ hyperedges of size 4 (i.e., 3-hyperedges), and we have set initial conditions to $(x_i = +1, y_i = -1)$ for nodes indexes $j \leq n/2$ and $(x_j = -1, y_j = +1)$ for $j > n/2$, as in [18].




FIG. 2. **Non-local hyperring vs. clique-projected network.** Left panels show spatiotemporal patterns a) and time series b) for variables $y_j(t)$ of the SL oscillators coupled with a non-local 3-hyperrings with $n = 204$ nodes and $m = 68$ hyperedges; the emerging dynamical behavior is a phase chimera state with two heads, i.e., there are regions of regular behavior, separated by two regions of decoherence. Right panels c) and d) show the analogous quantities on the clique-projected network. Panels b) and d) show the time series for nodes 50 (blue) and 101 (green). For panels a) and c), on the horizontal axis we set the node index j while in the vertical one, the time. The coupling strength is fixed at $\epsilon = 0.01$ and model parameters are $\lambda = 1$ and $\omega = 1$.




FIG. 3. **Dynamical quantities associated to non-local hyperring and clique-projected network.** Left panels show frequency a), amplitude b) and phase c) of the Stuart-Landau oscillators computed by using the Fast Fourier Transform (FFT) on the complex variables $z_j(t) = a_j(t) \exp[i(2\pi\Omega_j t + \theta_j)]$, as function of the node index, j , at 2500 time units. The left panels refer to quantities whose evolution is determined by a non-local 3-hyperring with $n = 204$ nodes and $m = 68$ hyperedges. The right panels show the analogous quantities for Stuart-Landau oscillators coupled via the clique-projected network. The coupling strength is $\epsilon = 0.01$ and the model parameters are $\lambda = 1$ and $\omega = 1$.

In Fig. 2, we compare the dynamics emerging by using higher-order and pairwise interactions. In panels a) and b), we show the behavior of the system of SL oscillators coupled with a non-local 3-hyperring, while panels c) and d) depict the same system of oscillators, but now coupled via the clique-projected network (pairwise interactions). In panel a), we can appreciate a phase chimera behavior, whereas in panel c) there are no domains with incoherent behaviors. Panels b) and d) show the time series for short and long times for nodes $j = 50$ (blue) and $j = 101$ (green). First of all, we observe that in both cases the orbits have the same amplitude and the same period, hence same frequency. However, in the case of the higher-order coupling, nodes initially synchronized are eventually found in phase opposition (see blue and green curves on panel b)); on the other hand, in the clique-projected network, nodes develop a small phase lag (see blue and green curves on panel d)). We are thus facing phase chimeras [26] in the hyperring, but not on the clique-projected network. To better analyze the obtained pattern, we cast the SL variables into a single complex number, one for each node, $z_j(t) := x_j(t) + iy_j(t)$, and we then rewrite the latter as $z_j(t) = a_j(t)\exp[i(2\pi\Omega_j t + \theta_j)]$, where a_j is the signal amplitude, Ω_j the frequency and θ_j the constant phase.

By using the Fast Fourier Transform and a sliding window setting, we numerically compute these quantities for all nodes after a sufficiently long transient interval. The results reported in Fig. 3 support the claim that we are dealing with phase chimeras according to [26]: indeed, the amplitudes, a_j , and the frequencies, Ω_j , are almost constant, i.e., their values do not depend on the node indexes, while the constant phases, θ_j , are node-dependent. In the case of higher-order coupling (panel c)), the phases can be divided into two regions: a first one, where the phase value does not depend on the node index, and a second one, where the opposite holds true. Moreover, in this second region the phases are scattered in $[-\pi, \pi]$. The clique-projected network has a similar behavior: amplitudes and frequencies have constant values, while the phases exhibit smooth transitions between two “flat” regions, where there is no node dependence of the phase θ_j . The latter difference, which can be quantified through the *normalized total phase variation* (see SM, which includes also Refs. [77–

79]), is indeed the key: while for the system with higher-order coupling we can see a clear chimera behavior, the aforementioned smooth variation cannot be considered a chimera state, as there is no coexistence between coherent and incoherent states. In the SM, we show that such difference between a higher-order topology and a pairwise one persists also when considering different pairwise projections, such as the aforementioned boundary of the cell complex. Although a full correspondence between the two cases cannot be established, as we are dealing with interactions that are intrinsically distinct, the difference is striking and it is consistently observed when the coupling strength is varied. Moreover, it also persists for different hyperrings, such as the 2-, 4- and 5-hyperrings, as shown in the SM numerically and by mean of the normalized total phase variation.

Conclusions. In this work, we have studied the effect of pure many-body interactions on chimera states, showing that the presence of a higher-order topology enhances the emergence of phase chimeras. We have introduced a non-local non-regular (pure) higher-order topology, which we called non-local hyperring, and observed that the comparison between the dynamics on the latter and on the clique-projected network shows a remarkable difference in the behaviors and allows to draw a conclusion that is consistent with the existing literature, namely that higher-order interactions facilitate the emergence of chimera states in systems of coupled oscillators. Moreover, non-local hyperrings allow for the emergence of other phenomena, as we show in the SM, such as a hybrid state of amplitude chimeras and chimera death, and pure chimera death [22]. We believe that the emergence of such type of chimera states, widely studied for pairwise interactions, but until now never observed with higher-order ones, makes our framework promising and will inspire future studies in this direction.

Acknowledgements The authors are grateful to Giovanni Reina, Elio Tuci, Hilda Cerdeira and Hiroya Nakao for useful discussions and feedback. R.M. acknowledges JSPS KAKENHI JP22K11919, JP22H00516, and JST CREST JP-MJCR1913 for financial support. During part of this work, R.M. was supported by a FRIA-FNRS Fellowship, funded by the Walloon Region, Grant FC 33443. T.N. expresses its gratitude to the University of Namur for financial support.

-
- [1] K. Kaneko, *Physica D* **41**, 137 (1990).
 - [2] D. Domínguez and H. Cerdeira, *Phys. Rev. Lett.* **71** (1993).
 - [3] A. M. Hagerstrom, T. E. Murphy, R. Roy, P. Hövel, I. Omelchenko, and E. Schöll, *Nature Physics* **8**, 658 (2012).
 - [4] E. A. Martens, S. Thutupalli, A. Fourriere, and O. Hallatschek, *Proceedings of the National Academy of Sciences* **110**, 10563 (2013).
 - [5] L. Gambuzza, A. Buscarino, S. Chessa, L. Fortuna, R. Meucci, and M. Frasca, *Phys. Rev. E* **9**, 032905 (2014).
 - [6] L. Gambuzza, L. Minati, and M. Frasca, *Chaos, Solitons and Fractals* **138**, 109907 (2020).
 - [7] M. H. Matheny, J. Emenheiser, W. Fon, A. Chapman, A. Salova, M. Rohden, J. Li, M. Hudoba de Badyn, M. Pósfai, L. Duenas-Osorio, *et al.*, *Science* **363**, eaav7932 (2019).
 - [8] T. Chouzeouris, I. Omelchenko, A. Zakharova, J. Hlinka, P. Jiruska, and E. Schöll, *Chaos* **28**, 045112 (2018).
 - [9] N. C. Rattenborg, C. J. Amlaner, and S. L. Lima, *Neuroscience & Biobehavioral Reviews* **24**, 817 (2000).
 - [10] S. Majhi, B. K. Bera, D. Ghosh, and M. Perc, *Physics of life reviews* **28**, 100 (2019).
 - [11] Y. Kuramoto, *Prog. Theor. Phys.* **94** (1995).
 - [12] Y. Kuramoto and H. Nakao, *Phys. Rev. Lett.* **76** (1996).
 - [13] Y. Kuramoto and H. Nakao, *Physica D* **103** (1997).
 - [14] Y. Kuramoto, D. Battogtokh, and H. Nakao, *Phys. Rev. Lett.* **81** (1998).
 - [15] Y. Kuramoto, H. Nakao, and D. Battogtokh, *Physica A* **288** (2000).

- [16] Y. Kuramoto and D. Battogtokh, *Nonlinear Phenom. Complex Syst.* **5** (2002).
- [17] D. M. Abrams and S. H. Strogatz, *Phys. Rev. Lett.* **93**, 174102 (2004).
- [18] A. Zakharova, *Chimera Patterns in Networks. Interplay between Dynamics, Structure, Noise, and Delay* (Springer, 2020).
- [19] O. Omel'chenko, *Nonlinearity* **31**, R121–R164 (2018).
- [20] K. Sathiyadevi, V. Chandrasekar, and D. Senthilkumar, *Phys. Rev. E* **98**, 032301 (2018).
- [21] G. C. Sethia, A. Sen, and G. L. Johnston, *Physical Review E* **88**, 042917 (2013).
- [22] A. Zakharova, M. Kapeller, and E. Schöll, *Phys. Rev. Lett.* **112**, 154101 (2014).
- [23] G. R. Simo, P. Louodop, D. Ghosh, T. Njoungouo, R. Tchitnga, and H. A. Cerdeira, *Phys. Lett. A* **409**, 127519 (2021).
- [24] J. H. Sheeba, V. Chandrasekar, and M. Lakshmanan, *Phys. Rev. E* **81**, 046203 (2010).
- [25] J. H. Sheeba, V. Chandrasekar, and M. Lakshmanan, *Phys. Rev. E* **79**, 055203 (2009).
- [26] E. Zajdela and D. Abrams, arxiv preprint <https://arxiv.org/abs/2308.06190> (2023).
- [27] C. Laing, *Phys. Rev. E* **92**, 050904(R) (2015).
- [28] M. Clerc, S. Coulibaly, M. Ferré, M. García-Ñustes, and R. Rojas, *Phys. Rev. E* **93**, 052204 (2016).
- [29] D. Pazó and E. Montbrió, *Phys. Rev. X* **4**, 011009 (2014).
- [30] A. Yeldesbay, A. Pikovsky, and M. Rosenblum, *Phys. Rev. Lett.* **112**, 144103 (2014).
- [31] I. Omelchenko, A. Provata, J. Hizanidis, E. Schöll, and P. Hövel, *Phys. Rev. E* **91**, 022917 (2015).
- [32] C. Meena, K. Murali, and S. Sinha, *International Journal of Bifurcation and Chaos* **26**, 1630023 (2016).
- [33] B. K. Bera, S. Majhi, D. Ghosh, and M. Perc, *Europhysics Letters* **118**, 10001 (2017).
- [34] R. Muolo, J. O'Brien, T. Carletti, and M. Asllani, *Eur. Phys. J. B* **97**, 6 (2024).
- [35] G. R. Simo, T. Njoungouo, R. Aristides, P. Louodop, R. Tchitnga, and H. A. Cerdeira, *Phys. Rev. E* **103**, 062304 (2021).
- [36] T. Njoungouo, G. R. Simo, P. Louodop, H. Fotsin, and P. K. Talla, *Chaos, Solitons & Fractals* **139**, 110082 (2020).
- [37] M. J. Panaggio and D. M. Abrams, *Nonlinearity* **28**, R67 (2015).
- [38] Z. Faghani, Z. Arab, F. Parastesh, S. Jafari, M. Perc, and M. Slavinec, *Chaos, Solitons and Fractals* **114**, 306 (2018).
- [39] Y. Zhang and A. Motter, *Phys. Rev. Lett.* **126**, 094101 (2021).
- [40] M. Asllani, B. Siebert, A. Arenas, and J. Gleeson, *Chaos* **32**, 013107 (2022).
- [41] F. Battiston, G. Cencetti, I. Iacopini, V. Latora, M. Lucas, A. Patania, J.-G. Young, and G. Petri, *Phys. Rep.* (2020).
- [42] F. Battiston, E. Amico, A. Barrat, G. Bianconi, G. Ferraz de Arruda, B. Franceschiello, I. Iacopini, S. Kéfi, V. Latora, Y. Moreno, M. Murray, T. Peixoto, F. Vaccarino, and G. Petri, *Nat. Phys.* **17**, 1093–1098 (2021).
- [43] G. Petri, P. Expert, F. Turkheimer, R. Carhart-Harris, D. Nutt, P. Hellyer, and F. Vaccarino, *J. Royal Soc. Interface* **11**, 20140873 (2014).
- [44] A. Sizemore, C. Giusti, A. Kahn, J. Vettel, R. Betzel, and D. Bassett, *J. Comp. Neurosci.* **44**, 115 (2018).
- [45] C. Giusti, R. Ghrist, and B. D.S., *J Comput Neurosci* **41**, 1 (2016).
- [46] J. Faskowitz, R. Betzel, and O. Sporns, *Netw. Neurosci.* **6**, 1 (2022).
- [47] L. Gambuzza, F. Di Patti, L. Gallo, S. Lepri, M. Romance, R. Criado, M. Frasca, V. Latora, and S. Boccaletti, *Nat. Comm.* **12**, 1 (2021).
- [48] M. Lucas, G. Cencetti, and F. Battiston, *Phys. Rev. Res.* **2**, 033410 (2020).
- [49] C. Kuehn and C. Bick, *Sci. Adv.* **7**, eabe3824 (2021).
- [50] P. Skardal and A. Arenas, *Phys. Rev. Lett.* **122**, 248301 (2019).
- [51] P. Skardal and A. Arenas, *Comm. Phys.* **3**, 1 (2020).
- [52] I. León and D. Pazó, *Phys. Rev. E* **105**, L042201 (2022).
- [53] T. Carletti, D. Fanelli, and S. Nicoletti, *J. phys. Complex.* **1**, 035006 (2020).
- [54] R. Muolo, L. Gallo, V. Latora, M. Frasca, and T. Carletti, *Chaos Solit. Fractals* **166**, 112912 (2023).
- [55] T. Carletti, F. Battiston, G. Cencetti, and D. Fanelli, *Phys. Rev. E* **101**, 022308 (2020).
- [56] M. Schaub, A. Benson, P. Horn, G. Lippner, and A. Jadbabaie, *SIAM Rev.* **62**, 353 (2020).
- [57] I. Iacopini, G. Petri, A. Barrat, and V. Latora, *Nat. Comm.* **10**, 2485 (2019).
- [58] G. de Arruda, G. Petri, and Y. Moreno, *Phys. Rev. Res.* **2**, 023032 (2020).
- [59] G. St-Onge, H. Sun, A. Allard, L. Hébert-Dufresne, and G. Bianconi, *Phys. Rev. Lett.* **127**, 158301 (2021).
- [60] S. Chowdhary, A. Kumar, G. Cencetti, I. Iacopini, and F. Battiston, *Journal of Physics: Complexity* **2**, 035019 (2021).
- [61] J.-F. de Kemmeter, L. Gallo, F. Boncoraglio, V. Latora, and T. Carletti, arxiv preprint <https://arxiv.org/abs/2305.03879> (2023).
- [62] Y. Zhang, V. Latora, and A. Motter, *Comm. Phys.* **4**, 1 (2021).
- [63] A. Salova and R. D'Souza, arXiv preprint <https://arxiv.org/abs/2107.13712> (2021).
- [64] S. Kundu and D. Ghosh, *Physical Review E* **105**, L042202 (2022).
- [65] X. Li, D. Ghosh, and Y. Lei, *Chaos, Solitons and Fractals* **170**, 113325 (2023).
- [66] C. Bick, T. Böhle, and C. Kuehn, *SIAM J. Appl. Dyn. Syst.* **22** (2023).
- [67] The same setting has been further investigated in [76].
- [68] H. Nakao, *Eur. Phys. J. Spec. Top.* **223**, 2411 (2014).
- [69] A. Zakharova, M. Kapeller, and E. Schöll, in *Journal of Physics: Conference Series*, Vol. 727 (IOP Publishing, 2016) p. 012018.
- [70] L. Neuhausser, A. Mellor, and R. Lambiotte, *Phys. Rev. E* **101**, 032310 (2020).
- [71] Let us observe that the nonlinearity of the coupling functions may not be enough to deal with proper higher-order interactions. In fact, as shown in [54], if the nonlinear functions h are the sum of terms $h(x_\ell) - h(x_j)$, they can still be decomposed onto pairwise ones. In what follows, we chose the coupling functions such that the latter case does not occur.
- [72] G. Bianconi, *Higher-Order Networks: An introduction to simplicial complexes* (Cambridge University Press, 2021).
- [73] Let us point out that labeling of the nodes is arbitrary.
- [74] M. Hajij, K. Istvan, and G. Zamzmi, *NeurIPS* (2020).
- [75] T. Carletti, L. Giambagli, and G. Bianconi, *Phys. Rev. Lett.* **130**, 187401 (2023).
- [76] C. Bick, T. Böhle, and O. Omel'chenko, arxiv preprint <https://arxiv.org/abs/2310.15698> (2023).
- [77] R. Gopal, V. Chandrasekar, A. Venkatesan, and M. Lakshmanan, *Phys. Rev. E* **89**, 052914 (2014).
- [78] S. Osher and L. Rudin, *SIAM J. Numer. Analysis* **27**, 919 (1990).
- [79] T. Rowland, *Total Variation* (MathWorld—A Wolfram Web Resource, <https://mathworld.wolfram.com/TotalVariation.html>, 2023).