

This is the accepted manuscript made available via CHORUS. The article has been published as:

Percolation in interdependent and interconnected networks: Abrupt change from second- to first-order transitions

Yanqing Hu, Baruch Ksherim, Reuven Cohen, and Shlomo Havlin

Phys. Rev. E **84**, 066116 — Published 20 December 2011

DOI: [10.1103/PhysRevE.84.066116](https://doi.org/10.1103/PhysRevE.84.066116)

Percolation in Interdependent and Interconnected Networks: Abrupt Change from Second to First Order Transition

Yanqing Hu^{1,2,3}, Baruch Ksherim², Reuven Cohen^{2,4}, Shlomo Havlin²

^{1.} School of Mathematics Southwest Jiaotong University, Chengdu 610031, China.

^{2.} Department of Physics, Bar-Ilan University, Ramat-Gan 52900, Israel

^{3.} Department of Systems Science, School of Management and Center for Complexity Research, Beijing Normal University, Beijing 100875, China

^{4.} Department of Mathematics, Bar-Ilan University, Ramat-Gan 52900, Israel

Robustness of two coupled networks system has been studied separately only for dependency coupling (S. Buldyrev et. al., Nature, 464, 08932, 2010) and only for connectivity coupling (E. A. Leicht and R. M. D'Souza, arxiv:09070894). Here we study, using a percolation approach, a more realistic coupled networks system where both interdependent and interconnected links exist. We find a rich and unusual phase transition phenomena including *hybrid* transition of mixed first and second order, i.e., discontinuities like in a first order transition of the giant component followed by a continuous decrease to zero like in a second order transition. Moreover, we find unusual discontinuous changes from second order to first order transition as a function of the dependency coupling between the two networks.

PACS numbers: 89.75.Hc, 64.60.ah, 89.75.Fb

Keywords: Cascading failures, Percolation, Coupled networks, Phase transition

I. INTRODUCTION

During the last decade complex networks have been studied intensively, where most of the research was devoted to analyzing the structure and functionality of isolated systems modeled as single non-interacting networks [1–23]. However, most real networks are not isolated, as they either complement other networks (“interconnected networks”), must consume resources supplied by other networks (“interdependent networks”) or both [24–28]. Thus, real networks continuously interact with each other, composing a large complex systems, and with the enhanced development of technology, the coupling between many networks becomes more complex and more significant.

Until now two different types of coupled networks models have been studied. Buldyrev et. al. [29] investigated the robustness of coupled systems with only interdependence links. In these systems, when a node of one network fails, its dependent counterpart node in the other network also fails. They found that this interdependence makes the system significantly more *vulnerable* [29, 30]. At the same time, Leicht and D’Souza [31] studied the case where only connectivity links couple the networks, i.e., “interconnected networks”, and found that the interconnected links make the system significantly more *robust*. However, real coupled networks often contain both types of links, interdependent as well as interconnected links. For example, the airport and the railway networks in Europe are two coupled networks composing a transportation system. In order to arrive to an airport, one usually uses the railway while people arriving to the country by airport usually use the railway. In this system, if the airport is disabled by some strike or accident, the passengers can still use the nearby railway station and travel to their destination or to another airport by train, so the two networks complement each other and are coupled by connectivity links. On the other hand, if the railway network is disabled, the airport traffic is damaged, and if the airport is disabled, the railway traffic is

damaged, so both networks are coupled by dependency links as well. The important characteristics of such systems, is that a failure of nodes in one network carries implications not only for this network, but also on the function of other dependent networks. In this way it is possible to have cascading failures between the coupled networks, that may lead to a catastrophic collapse of the whole system. Nevertheless, small clusters disconnected from the giant component in one network can still function through interconnected links connecting them to the giant component of the other network. Thus, we have two competing effects, the inter-connectivity links *increase* the robustness of the system, while the inter-dependency links *decrease* its robustness. Here we study the competition of the two types of inter-links on the system robustness using a percolation approach, and find unusual types of percolation phase transitions.

II. GENERAL FRAMEWORK

Let us consider a system of two networks, A and B , which are coupled by both dependency and connectivity links (see Fig. 1). The two networks are partially coupled by dependency links, so that a fraction q_A of A -nodes depends on nodes in network B , and a fraction q_B of B -nodes depends on the nodes in network A , with the following two assumptions: a node from one network depends on no more than one node from the other network, and that if node A_i depends on node B_j , then if B_j depends on some A_h , then $h = i$ (Fig. 1). In addition, the connectivity links within each network and between the networks can be described by a set of degree distributions $\{\rho_{k_A, k_{AB}}^A, \rho_{k_B, k_{BA}}^B\}$, where $\rho_{k_A, k_{AB}}^A$ ($\rho_{k_B, k_{BA}}^B$) denotes the probability of an A -node (B -node) to have k_A (k_B) links to other A -nodes (B -node) and k_{AB} (k_{BA}) links towards B -nodes (A -nodes). In this manner we get a two dimensional generating function describing all the connectivity




FIG. 1: (Color online) Two types of inter-links where the dependency links (dashed arrows) are not necessarily bidirectional. The nodes of A and B are randomly connected with *connectivity-links* (full line). The functionality of some of the A nodes (red) open circles *depend* on B-nodes (purple) solid circles and vice versa.

links [31], $\mathcal{G}_0^A(x_A, x_B) = \sum_{k_A, k_{AB}} \rho_{k_A, k_{AB}}^A x_A^{k_A} x_B^{k_{AB}}$, and $\mathcal{G}_0^B(x_A, x_B) = \sum_{k_B, k_{BA}} \rho_{k_B, k_{BA}}^B x_A^{k_{BA}} x_B^{k_B}$.

The cascading process is initiated by randomly removing a fraction $1 - p$ of the A-nodes and all their connectivity links. Because of the interdependence between the networks, the nodes in network B that depend on the removed A-nodes are also removed along with their connectivity links. As nodes and links are removed, each network breaks up into connected components (clusters). We assume that when the network is fragmented, the nodes belonging to the largest component (called the giant component) representing a finite fraction of the network which is still functional, while nodes that are parts of the remaining smaller clusters become dysfunctional, *unless* there exists a path of connectivity-links connecting these small clusters to the largest component of the other network. Since the networks have different topologies, the removal of nodes and related dependency links, is not symmetric in both networks, so that a cascading process occurs, until the system either becomes fragmented or stabilizes with a giant component.

Let $g_A(\varphi, \phi)$ and $g_B(\varphi, \phi)$ be the fraction of A-nodes and B-nodes in the giant components after the percolation process initiated by removing a fraction of $1 - \varphi$ and $1 - \phi$ of networks A and B respectively [11]. The functions $g_A(\varphi, \phi)$ and $g_B(\varphi, \phi)$ depend only on $\mathcal{G}_0^A(x_A, x_B)$ and $\mathcal{G}_0^B(x_A, x_B)$ (for details see Appendix) and the dynamics of the cascading process can be described by the following set of equations,

$$\begin{aligned} \varphi_1 &= p, & \phi_1 &= 1, & P_1^A &= \varphi_1 g_A(\varphi_1, \phi_1), \\ \phi_2 &= 1 - q_B(1 - p g_A(\varphi_1, \phi_1)), & P_2^B &= \phi_2 g_B(\varphi_1, \phi_2), \\ \varphi_2 &= p(1 - q_A(1 - g_B(\varphi_1, \phi_2))), & P_2^A &= \varphi_2 g_A(\varphi_2, \phi_2), \\ \phi_3 &= 1 - q_B(1 - p g_A(\varphi_2, \phi_2)), & P_3^B &= \phi_3 g_B(\varphi_2, \phi_3), \end{aligned} \quad (1)$$

where ϕ_i, φ_i are the remaining fraction of nodes at stage i of the cascade of failures and P_i^A, P_i^B are the corresponding giant components of networks A and B at the cascading stage i ,

respectively. Generally, the n^{th} step is given by the equations,

$$\begin{aligned} \varphi_n &= p(1 - q_A(1 - g_B(\varphi_{n-1}, \phi_n))), \\ \phi_n &= 1 - q_B(1 - p g_A(\varphi_n, \phi_{n-1})), \\ P_n^A &= \varphi_n g_A(\varphi_n, \phi_n), & P_n^B &= \phi_n g_B(\varphi_{n-1}, \phi_n). \end{aligned} \quad (2)$$

By introducing two new notations

$$u_A \equiv g_A(\varphi_\infty, \phi_\infty), \quad u_B \equiv g_B(\varphi_\infty, \phi_\infty), \quad (3)$$

we can write Eqs.(2) at the end of the cascading process, $n \rightarrow \infty$, as

$$\phi_\infty = p(1 - q_A(1 - u_B)), \quad \varphi_\infty = 1 - q_B(1 - p u_A), \quad (4)$$

and the giant components are,

$$\begin{aligned} P_\infty^A &= u_A \phi_\infty = u_A p(1 - q_A(1 - u_B)), \\ P_\infty^B &= u_B \varphi_\infty = u_B(1 - q_B(1 - p u_A)). \end{aligned} \quad (5)$$

III. POISSONIAN DEGREE DISTRIBUTIONS

We consider the case where all degree distributions of the connectivity intra- and inter-links are *Poissonian*, for which the functions u_A and u_B obtain a simple form. Assume \bar{k}_A and \bar{k}_B are the average intra-link degrees in networks A and B, and $\bar{k}_{AB}, \bar{k}_{BA}$ are the average inter-connectivity links degrees between A and B (allowing the case $\bar{k}_{AB} \neq \bar{k}_{BA}$, since the two networks may be of different sizes), we obtain,

$$\begin{aligned} u_A &= 1 - e^{-\bar{k}_A p u_A(1 - q_A(1 - u_B)) - \bar{k}_{AB} u_B(1 - q_B(1 - p u_A))}, \\ u_B &= 1 - e^{-\bar{k}_{BA} p u_A(1 - q_A(1 - u_B)) - \bar{k}_B u_B(1 - q_B(1 - p u_A))}. \end{aligned} \quad (6)$$

Generally, for fixed parameters $\bar{k}_A, \bar{k}_B, \bar{k}_{AB}, \bar{k}_{BA}, q_A, q_B$ and p , it is often impossible to achieve an explicit formula for the giant components P_∞^A and P_∞^B . However, one can still solve Eqs. (6) graphically (numerically) and substitute the numerical solution into Eqs. (5). For simplicity and without loss of generality, we study the case where $\bar{k}_A = \bar{k}_B \equiv \bar{k}$ and $\bar{k}_{AB} = \bar{k}_{BA} \equiv \bar{K}$. Fig. 2a compares the numerical solutions with the simulation results for P_∞^A and P_∞^B as a function of p , showing that the analytical results of Eqs. (5) and (6) are in excellent agreement with the simulations.

A. Partial Dependence

Next we are interested in the properties of the phase transition under random attack, so first we determine the conditions when transition does not occur. This is the case where for a given $q_B < 1$ even if all nodes of network A are removed (*i.e.*, $p = 0$), there still exists a giant component in network B (see circles in Fig. 2a) and no phase transition occurs. For Poissonian degree distributions, it is easy to see, that if after




FIG. 2: (Color online) **a.** Giant components P_A^∞ and P_B^∞ vs. fraction of remaining nodes, p , for $N = 10000$, $\bar{k} = 2$ and $\bar{K} = 1$. Networks A (open symbols) and B (full symbols) are shown for different (q_A, q_B) pairs: (0.8, 0.1) (\circ); (0.1, 0.8) (\diamond); (0.8, 0.8) (\square). The symbols represent simulations and the lines the theory. We see three different types of behaviors: no phase transition (\circ), second order phase transition (\diamond) and first order phase transition (\square). **b.** Phase diagram showing the first order, second order and hybrid phase transition regimes and the boundaries, for $q_B = 1$ and $\bar{k} = 3$. In the second order transition regime, between the two dashed curve (red and blue) there exists a hybrid phase transition regime (details in Fig. 3c). Since the hybrid transition is continuous in the neighborhood of p_c and jump occurs well above p_c , we classify this hybrid phase transition as a second order phase transition.

the removal of all B-nodes that depend on the attacked A-nodes, the new average intra-link degree in network B is less than one, i.e.,

$$\bar{k}_B(1 - q_B) < 1, \quad (7)$$

a phase transition does occur. Therefore, our further analysis is based on condition (7). In addition, from now on, we will set both dependency couplings, q_A and q_B , to be larger than zero.

For a second order phase transition, the giant component decreases continuously to zero at the percolation threshold p_c . Thus, by taking the limit of system (6) at $u_A = u_B = 0$ we obtain the second order threshold,

$$p_c^{II} = \frac{1 - \bar{k}_B(1 - q_B)}{(\bar{k}_A + (\bar{k}_{BA}\bar{k}_{AB} - k_A k_B)(1 - q_B))(1 - q_A)}. \quad (8)$$

In particular for $q_A = 1$ and $0 < q_B < 1$ this threshold becomes

$$p_c^{II} = \frac{1}{\bar{k}_B(1 - q_B)},$$

which together with Eq. (7) implies that $p_c^{II} > 1$, and therefore the phase transition must be of first order at $p_c^I < 1$ (that will be determined later).

When extracting u_B from the first equation of system (6), it can be rewritten as,

$$u_B = -\frac{\log(1 - u_A) + k_A p(1 - q_A)u_A}{k_A p q_A u_A + k_{AB}[1 - q_B(1 - p u_A)]} \equiv H_1(u_A), \quad (9)$$

$$u_B = 1 - e^{-\bar{k}_{BA}u_A p(1 - q_A(1 - u_B)) - \bar{k}_B u_B(1 - q_B(1 - u_A p))} \equiv H_2(u_A).$$

The intersection of the two curves (maximum solution of u_A, u_B) is the solution of the system. When the phase transition is first order and $p = p_c^I$, the curves of Eqs. (9) are tangentially touching at the solution point, where,

$$\left(\frac{dH_1}{du_A} = \frac{dH_2}{du_A}\right)\bigg|_{p=p_c^I}. \quad (10)$$

Obviously, u_A , u_B and p can be treated as variables of Eqs. (9) and (10). Solving these equations, the minimal solution of p and the corresponding maximal u_A and u_B is the solution of the system at criticality.

B. Full Dependence

When networks A and B are fully dependent, i.e., $q_A = q_B = 1$, both networks must be the same size and therefore $\bar{k}_{AB} = \bar{k}_{BA} \equiv \bar{K}$ system (6) yields a simple form

$$u_A = 1 - \exp\{-p u_A u_B (\bar{k}_A + \bar{K})\},$$

$$u_B = 1 - \exp\{-p u_A u_B (\bar{k}_B + \bar{K})\}.$$

The size of the mutual giant component, P_∞ , is thus given by,

$$P_\infty = P_\infty^A = P_\infty^B = p(1 - e^{-P_\infty(\bar{k}_A + \bar{K})})(1 - e^{-P_\infty(\bar{k}_B + \bar{K})}), \quad (11)$$

which is similar to the solution of fully interdependent system [29, 32], where the only difference is that the degrees of networks A and B are now replaced by $\bar{k}_A + \bar{k}_{AB}$ and $\bar{k}_B + \bar{k}_{BA}$, respectively. Thus, interestingly, in a fully interdependent coupled networks system adding connectivity inter-links has the same effect as increasing the intra-degree of the corresponding networks and therefore, in this case, the phase transition must be of first order. From Eqs. (9) and (10), one can get the threshold,

$$p_c^I = \frac{1}{k_A(1 - u_A)[-1 + (1 - u_A)^\alpha - u_A \alpha(1 - u_A)^{\alpha-1}]}, \quad (12)$$

where, $\alpha \equiv (\bar{k}_B + \bar{k}_{BA})/(\bar{k}_A + \bar{k}_{AB})$, and u_A satisfies the equation,

$$u_A = 1 - \exp\left\{\frac{u_A[1 - (1 - u_A)^\alpha]}{(1 - u_A)[-1 + (1 - u_A)^\alpha - u_A \alpha(1 - u_A)^{\alpha-1}]}\right\}. \quad (13)$$

By substituting p_c^{II} from Eq. (8) into Eqs. (9) and (10) and evaluating both u_A and u_B we can derive in the phase diagram, the boundary between the first and second order transitions




FIG. 3: **a.** Size of giant components vs. dependency and connectivity links strength, for $q_B = 1$ and $\bar{k} = 3$. The giant components size at p_c changes from zero to a finite value while changing q_A and \bar{K} . When q_A and \bar{K} are at the boundary of different phase transitions, the jump occurs (see Fig. 2b). **b.** The values of $P_\infty^A(p_c)$ (\circ), $P_\infty^B(p_c)$ (\square) along the boundary for $q_B = 1$ and $\bar{k} = 3$. **c.** An hybrid phase transition, for $q_B = 1$, $q_A = 0.35$, $\bar{k} = 3$ and $\bar{K} = 0.1$. According to Eqs. (5), P_∞^A and P_∞^B have the same properties as u_A and u_B respectively. At $p = p_c^h \approx 0.66$ the values of u_A and u_B jump, and then for lower p values they continuously approach zero. In the inset, simulation and theoretical results are symbols and lines respectively.

(see dashed line in Fig. 2b). An interesting phenomenon, which to the best of our knowledge, has not been observed before, is that when the phase transition changes from first to second order, there are discontinuities (abrupt jumps) of $P_\infty^A(p_c)$, $P_\infty^B(p_c)$ in the phase transition boundary (see Fig. 3a and 3b). The boundary between first and second order phase transition satisfies $p_c^I = p_c^{II}$. Therefore, by replacing p_c^I by p_c^{II} in Eq. (9) and evaluating both u_A and u_B we obtain the boundary, seen in Fig. 3a, between the first and second order transitions. When we reduce the three equations to a single equation, u_A , u_B should always be the maximal non-negative solution in $[0, 1]$. When Eqs. (9) and (10) have more than one solution, we always choose the minimal non-negative value, p_c^{min} , and the corresponding maximal values, u_A^{max} , u_B^{max} as the

physical solution at the threshold. In part of the boundary, $u_A^{max} > 0$ and $u_B^{max} > 0$, and of course $p_c^{min}, u_A = 0, u_B = 0$ are also the solution of the system. This means that there exist two intersections where both satisfy the tangential condition on the boundary (as shown in Fig. 4a). This implies that when the order of the phase transition changes from first to second, $P_\infty^A(p_c)$, $P_\infty^B(p_c)$ are discontinuous (see Fig. 3a and 3b). This phenomenon contrasts most known systems possessing both first and second order transitions. Usually, in physical systems, the first order jump in the order parameter, and other related properties, such as the specific heat, present a continuous change along the transition line when the system changes from first to second order [34].

In addition to the existence of jumps in $P_\infty^A(p_c)$, $P_\infty^B(p_c)$ at the boundary between the first and second order phase transitions, we find also another unusual phenomenon. When one network strongly depends on the other, there exist *hybrid* phase transitions. An hybrid phase transition means that when the attack strength, $1-p$, increases the size of the giant component jumps at p_c^h from a large value to a small value, and then continuously decreases to zero. A similar behavior has been found in bootstrap percolation [33]. Since the second order transition is characterized by a giant component which is continuous in the neighborhood of p_c we regard, the hybrid phase transition regime as a second order phase transition regime (see Fig. 2b). For the hybrid phase transition, there exists a threshold p_c^h at which the jump occurs. For p just below p_c^h , the solution of Eqs. (9) for u_A , u_B , will jump to lower values. After the jump, when p further decreases, u_A and u_B approach to zero continuously which implies that the giant components sizes change to zero continuously (see Fig. 3c).

For the three equations system (Eqs. (9) and (10)), the minimal solution of p^{min} in $[0, 1]$ is p_c (the physical solution). Besides p^{min} , if Eqs. (9) have another solution $p_c^h \in (0, 1)$ and corresponding solution u_A^h , u_B^h , we can find an hybrid phase transition. The set (p^h, u_A^h, u_B^h) means that when p is just below p_c^h , the solutions u_A , u_B of the first two equations of Eqs. (9) will jump to small values. After the jump, when we continue to decrease p towards $p_c = p^{min}$, both u_A and u_B will continuously decrease to zero. For example, for the parameters $q_A = 0.35$, $q_B = 1$, $\bar{k} = 3$ and $\bar{K} = 0.1$, we obtain $p_c \approx 0.556$ and $p_c^h \approx 0.66$. When p is just below 0.66, the giant components drop to smaller positive values like in a first order phase transition. After this discontinuous drop, the giant components' sizes continuously decrease to zero while decreasing p from 0.66 to 0.556, like a second order phase transition (see Fig. 4b).




FIG. 4: (Color online) Tangential conditions. **a.** An abrupt jump on the boundary for $q_A = 0.394, q_B = 0.8$ and $\bar{k} = 3, \bar{K} = 0.2$. Here $p_c^I = p_c^{II} = 0.5464$ which is the threshold of the system. Although both intersections (one of which is at the origin) satisfy the tangential condition, the u_A^{\max}, u_B^{\max} values are the physical solution and the transition is of the first order. **b.** Hybrid transition analysis, for $q_B = 1, q_A = 0.35, \bar{k} = 3$ and $\bar{K} = 0.1$. Here $p_c \approx 0.556$ and $p_c^h \approx 0.66$. The maximal intersection S satisfies the tangential condition. When continuously decreasing p , the solution of the system jumps from the maximal intersection S to the minimal intersection Q and then continuously decreases to zero.

IV. SUMMARY AND CONCLUSION

In summary, we studied the dynamics of the cascading failures process and the steady solutions of failures of the giant components in coupled networks, when both interdependent and interconnected links exist, using a percolation approach. Although our detailed analysis is for ER networks, the theory can be applied to any network systems topology. We find that the existence of inter-connectivity links between interdependent networks, introduces rich and intriguing phenomena through the process of cascading failures. Increasing the strength of interconnecting links can significantly change the transition behavior and often brings up some counterintuitive phenomena, such as changing the transition from second order to first order (as seen in Fig. 2b). We also find an unusual abrupt jump in the boundary between first and second order phase transitions at the criticality. Moreover, when one of the networks strongly depends on the other network, unusual hybrid phase transitions are observed.

We thank Amir Bashan for helpful discussions. This work is partially supported by ONR, DFG, DTRA, EU project Epi-network and the Israel Science Foundation. Y. Hu is supported

by NSFC under Grant No. 60974084, 60534080.

V. APPENDIX: HOW TO GET $g_A(\varphi, \phi)$ AND $g_B(\varphi, \phi)$

We model the percolation process using the branching process approach. Let $\mathcal{G}_0^A(x_A, x_B) = \sum_{k_A, k_{AB}} \rho_{k_A, k_{AB}}^A x_A^{k_A} x_B^{k_{AB}}$, and $\mathcal{G}_0^B(x_A, x_B) = \sum_{k_B, k_{BA}} \rho_{k_B, k_{BA}}^B x_A^{k_{BA}} x_B^{k_B}$, be the degree distributions' generating functions. The probability of following a randomly chosen AB-link connecting an A-node of degree k_A to a B-node with excess k_{AB} degree (i.e., having total A to B degree of $k_{AB}+1$) is proportional to $(k_{AB}+1)\rho_{k_A, k_{AB}}^A$, and the generating function for this distribution is [31],

$$\mathcal{G}_1^{AB}(x_A, x_B) = \sum_{k_A, k_{AB}} \frac{(k_{AB}+1)\rho_{k_A, k_{AB}+1}^A}{\sum_{k'_A, k'_{AB}} k'_{AB} \rho_{k'_A, k'_{AB}}^A} \cdot x_A^{k_A} x_B^{k_{AB}}. \quad (14)$$

Analogously, we construct the other three excess generating functions $\mathcal{G}_1^{AA}(x_A, x_B)$, $\mathcal{G}_1^{BA}(x_A, x_B)$ and $\mathcal{G}_1^{BB}(x_A, x_B)$.

After removing a fraction $1-\varphi$ of nodes in network A, and a fraction $1-\phi$ of nodes in network B, we can set new arguments to the generating functions, so that, x_A and x_B will be replaced by $1-\varphi(1-x_A)$ and $1-\phi(1-x_B)$, respectively [35–37]. Suppose $g_A(\varphi, \phi)$, $g_B(\varphi, \phi)$ are the fractions of A-nodes and B-nodes in the giant components after removal of $1-\varphi$ and $1-\phi$ fractions of networks A and B, respectively. Then we have,

$$g_A(\varphi, \phi) = 1 - \mathcal{G}_0^A(1 - \varphi(1 - f_A), 1 - \phi(1 - f_{BA})), \quad (15)$$

$$g_B(\varphi, \phi) = 1 - \mathcal{G}_0^B(1 - \varphi(1 - f_{AB}), 1 - \phi(1 - f_B)),$$

where,

$$f_A = \mathcal{G}_1^{AA}(1 - \varphi(1 - f_A), 1 - \phi(1 - f_{BA})), \quad (16)$$

$$f_{AB} = \mathcal{G}_1^{AB}(1 - \varphi(1 - f_A), 1 - \phi(1 - f_{BA})),$$

$$f_{BA} = \mathcal{G}_1^{BA}(1 - \varphi(1 - f_{BA}), 1 - \phi(1 - f_B)),$$

$$f_B = \mathcal{G}_1^{BB}(1 - \varphi(1 - f_{BA}), 1 - \phi(1 - f_B)).$$

When all degree distributions of inter and intra networks A and B are Poissonian distributed, all the functions can be more simple. Assume \bar{k}_A and \bar{k}_B are the average intra-links degrees in networks A and B and \bar{k}_{AB} , \bar{k}_{BA} are the average inter-links degrees between A and B (allowing the case $\bar{k}_{AB} \neq \bar{k}_{BA}$, since the network sizes of A and B can be different), then we have $G_0^{AA}(x_A) = e^{\bar{k}_A(x_A-1)}$, $G_0^{AB}(x_B) = e^{\bar{k}_B(x_B-1)}$, $G_0^{BA}(x_A) = e^{\bar{k}_{BA}(x_A-1)}$, $G_0^{BB}(x_B) = e^{\bar{k}_B(x_B-1)}$ and

(17)

$$\begin{aligned} \mathcal{G}_1^{AA}(x_A, x_B) &= \mathcal{G}_1^{AB}(x_A, x_B) = G_0^A(x_A, x_B) = G_0^{AA}(x_A)G_0^{AB}(x_B) \\ \mathcal{G}_1^{BB}(x_A, x_B) &= \mathcal{G}_1^{BA}(x_A, x_B) = G_0^B(x_A, x_B) = G_0^{BA}(x_A)G_0^{BB}(x_B) \end{aligned}$$

Submitting the above equations into systems (15) and (16), we get

$$g_A(\varphi, \phi) = 1 - \exp\{-\bar{k}_A x g_A(\varphi, \phi) - \bar{k}_{AB} y g_B(\varphi, \phi)\}, \quad (18)$$

$$g_B(\varphi, \phi) = 1 - \exp\{-\bar{k}_{BA} x g_A(\varphi, \phi) - \bar{k}_B y g_B(\varphi, \phi)\}.$$

-
- [1] A.-L. Barabási and R. Albert, *Science* **286**, 509 (1999).
 - [2] D. J. Watts and S. H. Strogatz, *Nature (London)* **393**, 440 (1998).
 - [3] R. Albert and A.-L. Barabási, *Rev. Mod. Phys.* **74**, 47 (2002).
 - [4] Y. Hu, Y. Wang, D. Li, S. Havlin and Z. Di, *Phys. Rev. Lett.* **106**, 108701 (2011).
 - [5] C. Song, S. Havlin and H.A. Makse, *Nature (London)* **433**, 392 (2005).
 - [6] R. Pastor-Satorras and A. Vespignani, *Evolution and Structure of the Internet: A Statistical Physics Approach* (Cambridge University Press, 2006).
 - [7] S. N. Dorogovtsev and J. F. F. Mendes, *Evolution of Networks: from Biological nets to the Internet and WWW* (Oxford University Press, New York, 2003).
 - [8] A. Barrat, M. Barthélemy and A. Vespignani, *Dynamical processes on complex networks* (Cambridge University Press, England, 2008).
 - [9] D. S. Callaway, M. E. J. Newman, S. H. Strogatz and D. J. Watts, *Phys. Rev. Lett.* **85**, 5468-5471 (2000).
 - [10] R. Cohen, K. Erez, D. ben-Avraham and S. Havlin, *Phys. Rev. Lett.* **85**, 4626 (2000); **86**, 3682 (2001).
 - [11] M. E. J. Newman, *Networks: An Introduction*. (Oxford University press 2010).
 - [12] S. H. Strogatz, *Nature* **410**, 268 (2001).
 - [13] D. J. Watts, *Proc. Natl. Acad. Sci. USA* **99**, 5766 (2002).
 - [14] M. E. J. Newman, and M. Girvan, *Phys. Rev. E* **69**, 026113 (2004).
 - [15] C. Song, S. Havlin and H. A. Makse, *Nature Physics* **2**, 275 (2006).
 - [16] M. Tumminello, F. Lillo, and R. N. Mantegna, *Europhys. Lett.* **78**, 30006 (2007).
 - [17] G. Caldarelli and A. Vespignani, *Large scale Structure and Dynamics of Complex Webs* (World Scientific, New York, 2007).
 - [18] A. Barrat, M. Barthélemy, and Vespignani, *A. Dynamical Processes on Complex Networks* (Cambridge University Press, Cambridge, 2008).
 - [19] M. Barthélemy, *Spatial networks. Physics Reports* **499**, 1-101 (2011).
 - [20] D. Li, K. Kosmidis, A. Bunde, and S. Havlin, *Nature Physics* **7**, 481-484, (2011).
 - [21] R. Albert, H. Jeong and A. L. Barabási, *Nature* **406**, 378 (2000).
 - [22] R. Milo, S. Shen-Orr, S. Itzkovitz, N. Kashtan, D. Chklovskii, and U. Alon, *Science* **298**, 824- 827 (2002).
 - [23] L. K. Gallos, R. Cohen, P. Argyrakis, A. Bunde and S. Havlin, *Phys. Rev. Lett.* **94**, 188701 (2005).
 - [24] V. Rosato *et al.*, *Int. J. Crit. Infrastruct.* **4**, 63-79, (2008).
 - [25] N. K. Svendsen and S. D. Wolthusen, *Information Security Technical Report*, **21**, page 44-55, (2007).
 - [26] S. M. Rinaldi, J. P. Peerenboom and T. K. Kelly, *IEEE Contr. Syst. Mag.* **21**, 11-25 (2001).
 - [27] J. Peerenboom *et al.*, in *Proc. CRIS/DRM/IIT/NSF Workshop Mitigat. Vulnerab. Crit. Infrastruct. Catastr. Failures* (2001).
 - [28] J. C. Laprie, K. Kanoun and M. Kaaniche, *SAFECOMP-2007*, **4680**, 54-67 (2007).
 - [29] S. V. Buldyrev, R. Parshani, G. Paul, H. E. Stanley and S. Havlin, *Nature*, **464**, 7291 (2010).
 - [30] R. Parshani, S. V. Buldyrev, S. Havlin, *Phys. Rev. Lett.* **105**, 048701 (2010).
 - [31] E. A. Leicht, R. M. D'Souza, *arxiv:0907.0894*.
 - [32] J. Gao, S. V. Buldyrev, S. Havlin and H. Eugene Stanley, *arxiv:1010.5829*.
 - [33] G. J. Baxter, S. N. Dorogovtsev, A. V. Goltsev and J. F. F. Mendes, *Phys. Rev. E* **82**, 011103 (2010).
 - [34] R. K. Pathria, *Statistical Mechanics*. (Elsevier (Singapore) Pte Ltd. 2003).
 - [35] M. E. J. Newman, *Phys. Rev. E* **66**, 016128 (2002).
 - [36] J. Shao, S. V. Buldyrev, R. Cohen, M. Kitsak, S. Havlin and H. E. Stanley, *Europhys. Lett.* **84**, 48004 (2008).
 - [37] J. Shao, S. V. Buldyrev, L. A. Braunstein, S. Havlin and H. E. Stanley, *Phys. Rev. E* **80**, 036105 (2009).