



# CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Generalized thermodynamic relations for a system experiencing heat and mass diffusion in the far-from-equilibrium realm based on steepest entropy ascent

Guanchen Li and Michael R. von Spakovsky

Phys. Rev. E **94**, 032117 — Published 15 September 2016

DOI: [10.1103/PhysRevE.94.032117](https://doi.org/10.1103/PhysRevE.94.032117)

# Generalized thermodynamic relations for a system experiencing heat and mass diffusion in the far-from-equilibrium realm based on steepest entropy ascent

Guanchen Li\* and Michael R. von Spakovsky†

*Center for Energy Systems Research,  
Mechanical Engineering Department  
Virginia Tech, Blacksburg, VA 24061*

(Dated: July 12, 2016)

This paper presents a nonequilibrium thermodynamic model for the relaxation of a local, isolated system in nonequilibrium using the principle of steepest entropy ascent (SEA), which can be expressed as a variational principle in thermodynamic state space. The model is able to arrive at the Onsager relations for such a system. Since no assumption of local equilibrium is made, the conjugate fluxes and forces, which result, are intrinsic to the subspaces of the system's state space and are defined using the concepts of hypoequilibrium state and nonequilibrium intensive properties, which describe the non-mutual equilibrium status between subspaces of the thermodynamic state space. The Onsager relations are shown to be a thermodynamic kinematic feature of the system independent of the specific details of the micro-mechanical dynamics. Two kinds of relaxation processes are studied with different constraints (i.e., conservation laws) corresponding to heat and mass diffusion. Linear behavior in the near-equilibrium region as well as nonlinear behavior in the far-from-equilibrium region are discussed. Thermodynamic relations in the equilibrium and near-equilibrium realm, including the Gibbs relation, the Clausius inequality, and the Onsager relations, are generalized to the far-from-equilibrium realm. The variational principle in the space spanned by the intrinsic conjugate fluxes and forces is expressed via the quadratic dissipation potential. As an application, the model is applied to the heat and mass diffusion of a system represented by a single particle ensemble, which can also be applied to a simple system of many particles. Phenomenological transport coefficients are also derived in the near-equilibrium realm.

## I. INTRODUCTION

The study of nonequilibrium relaxation processes including chemical kinetics, mass diffusion, and heat diffusion is typically accomplished using approaches based on microscopic mechanics or thermodynamics. With the former, system state space is spanned by microstates and the governing equation is based on the dynamics of classical mechanics (e.g., molecular dynamics [1] or kinetic theory [2]), quantum mechanics (e.g., nonequilibrium Green's functions [3] or the quantum Boltzmann equation, i.e., the Uehling-Unlenbeck-Boltzmann equation [4-6]) or a stochastic process (e.g., Monte Carlo simulation of the Ising model [7]). These approaches provide complete information of the microscopic process such as individual particle collisions or quantum state scattering. However, the large amount of very detailed information required inevitably results in large computational burdens, which limit the applicability of these approaches.

Approaches based on thermodynamics, on the other hand, are not similarly burdened and are able to generally capture the features of the relaxation process of interest via, for example, the Onsager relation. Approaches of this type include nonequilibrium thermodynamics [8, 9], linear response functions and the fluctuation-dissipation theorem [10], stochastic thermodynamics

[11], extended irreversible thermodynamics [12], etc. The thermodynamic features captured can be regarded as a coarse graining of the microscopic dynamics or as a pattern in ensemble evolution [13, 14], which computationally is more efficient. However, most of these approaches have limited or no applicability in the far-from-equilibrium realm, since the local or near-equilibrium assumption is needed or analytical solutions are only available at steady state. In addition, their governing equations are phenomenological or stochastic in nature and, thus, do not have a first-principle basis. To address these issues and push the application of thermodynamic principles further into the nonequilibrium realm, it is of great importance to i) find a general and simple description of nonequilibrium state corresponding to a thermodynamic pattern of the microscopic description, ii) fundamentally define the macroscopic properties of any thermodynamic state (i.e., extensive or intensive properties for both equilibrium and nonequilibrium states), and iii) use a thermodynamic governing equation based on first principles.

Steepest-entropy-ascent quantum thermodynamics (SEAQT), which is a first-principle, thermodynamic-ensemble based approach, addresses all of the issues raised above, providing a governing equation able to describe the nonequilibrium process from an entropy generation standpoint. The description of system state is based on the density operator of quantum mechanics or the probability distribution in state space of classical mechanics. The macroscopic properties of entropy, energy, and particle number, which are well defined for any state of any system [15], are

---

\* [guanchen@vt.edu](mailto:guanchen@vt.edu)

† [vonspako@vt.edu](mailto:vonspako@vt.edu)

used to develop the governing equation and describe system state evolution. Recently, this description has been simplified via the concept of hypoequilibrium state [16], which captures the global features of the microscopic description for the relaxation process. In addition, the concept of nonequilibrium intensive properties introduced in [16] based on the concept of hypoequilibrium state enables a complete description of the nonequilibrium evolution of state when combined with the set of nonequilibrium extensive properties. Unlike the intensive property definitions of other nonequilibrium thermodynamic approaches (definitions which require a local-equilibrium, near-equilibrium, or steady state assumption or a phenomenological basis), the definitions in the SEAQT framework are fundamental and available to all nonequilibrium states and are especially suitable for the description of the evolution in state of relaxation processes. Both of these concepts enables the generalization of many equilibrium (or near-equilibrium) thermodynamic relations such as the Gibb's relation, the Clausius inequality, the Onsager relations and the quadratic dissipation potential into the far-from-equilibrium realm.

To describe the SEAQT framework, the paper starts with a derivation in Sec. II.A of the SEAQT relaxation dynamics from the geometry of system state space. Sec II.B distinguishes the kinetic (trajectory) and dynamic (time-dependent) features of the relaxation. Some useful mathematical features of the relaxation process are then presented in Sec. II.C, enabling the definition of the concepts of hypoequilibrium state and nonequilibrium intensive properties. As an alternative to the geometric derivation from state space, the relaxation dynamics of SEAQT can also be derived using a variational principle in system state space as is done in Sec. II.D. Section III then follows with a discussion of mass diffusion in a local, isolated system in nonequilibrium. Subsequently, the transport equations, the Gibbs relation, and the Onsager relations are generalized for the nonequilibrium relaxation process to the far-from-equilibrium realm, and the quadratic dissipation potential is given to complete the link from the variational principle in system state space to that for the conjugate fluxes and/or forces [9]. Section IV then discusses the process of heat diffusion by choosing a set of system constraints different from those used for mass diffusion. In Sec. V, the SEAQT model is applied to the study of the heat and mass diffusion of a simple system consisting of an ideal gas (hydrogen), which can be represented by a single-particle ensemble. Linear behavior in the near-equilibrium realm provides the phenomenological transport equations, and higher-order, nonlinear behavior in the near-equilibrium realm is also studied. Finally, Section VI concludes the paper with some final comments.

## II. THEORY: SEA-QT EQUATION OF MOTION

The basic framework of SEAQT is introduced in this section. This framework describes the relaxation process of a local, isolated system in nonequilibrium based on thermodynamic concepts. To begin with, the equation of motion for such a system is derived in Sec. II.A from the geometric principle of steepest entropy ascent. This is followed in Sec. II.B by a discussion of the kinetics and dynamics of the relaxation process, which enable the study of the thermodynamic trajectory regardless of the details of the microscopic interactions. A description of nonequilibrium state and its evolution for the relaxation process is then given in Sec. II.C using the concepts of hypoequilibrium state and nonequilibrium intensive properties. Finally, in Sec. II.D, a presentation of the variational principle of steepest entropy ascent is used to derive the SEAQT equation of motion for the local system.

### A. SEAQT equation of motion for an isolated system

Based on the discussion by Grmela [13, 14, 17] and Beretta [18, 19], the general form of a nonequilibrium framework is a combination of both irreversible relaxation and reversible symplectic dynamics. If written in the generalized form of the Ginzburg-Landau equation [13, 19], the equation of motion takes the following form:

$$\frac{d}{dt}\gamma(t) = X_{\gamma(t)}^H + Y_{\gamma(t)}^H \quad (1)$$

where  $\gamma(t)$  represents the state evolution trajectory,  $X_{\gamma(t)}^H$  and  $Y_{\gamma(t)}^H$  are functions of the system state  $\gamma(t)$  and represent the reversible symplectic dynamics and irreversible relaxation dynamics, respectively. In the SEAQT framework, system state is represented by the density operator  $\hat{\rho}$ , while  $X_{\gamma(t)}^H$  follows the Schrödinger equation, and  $Y_{\gamma(t)}^H$  is derived from the SEA principle. Thus,

$$\frac{d\hat{\rho}}{dt} = \frac{1}{i\hbar}[\hat{\rho}, \hat{H}] + \frac{1}{\tau(\hat{\rho})}\hat{D}(\hat{\rho}) \quad (2)$$

where  $\tau$  is the relaxation time, which represents the speed of system evolution in Hilbert space and  $\hat{D}$  is the dissipation operator determined via a constrained gradient in Hilbert space. A metric tensor must be specified in the derivation of this dissipation term, since it describes the geometric features of the Hilbert space [18].

In the application of modeling heat and mass diffusion presented here, the system is restricted to the class of dilute-Boltzmann-gas states in which the particles have no quantum correlation between eigenstates and are independently distributed [20]. Such states can be represented by a single-particle density operator

that is diagonal in the basis of the single-particle energy eigenstates. The Hilbert space metric chosen is the Fisher-Rao metric, which is uniform in different dimensions of Hilbert space. Under these conditions, the symplectic Schrödinger term in the equation of motion vanishes. Thus, the focus of this paper is on the irreversible relaxation process only.

A group of energy eigenlevels  $\{\epsilon_i, i = 1, 2, \dots\}$  is determined from the system Hamiltonian. The state of the system can then be represented by a probability distribution among the energy eigenlevels  $\{p_i, i = 1, 2, \dots\}$ , which are the diagonal terms of the density operator. Using the Fisher-Rao metric of the probability space  $\{p_i, i = 1, 2, \dots\}$ , one can define the distance in probability space, which can be used as the state distance. Equivalently, the square root of the probability distribution  $\{x_i, i = 1, 2, \dots\}$  can be used to represent the system state with the result that the Fisher-Rao metric of the probability space becomes the Euclidean metric in the square root of the probability space of  $\{x_i, i = 1, 2, \dots\}$ . The later representation is used in the paper. Both are expressed as follows:

$$\begin{aligned} \text{State: } & \{p_i, i = 1, 2, \dots\}, \\ \text{Distance: } & dl = \frac{1}{2} \sqrt{\sum_i p_i \left(\frac{d \ln p_i}{d\theta}\right)^2} d\theta \end{aligned} \quad (3)$$

$$\begin{aligned} \text{State: } & \{x_i, i = 1, 2, \dots\}, \\ \text{Distance: } & dl = \sqrt{\sum_i (dx_i)^2} \end{aligned} \quad (4)$$

where  $dl$  is the distance between  $p(\theta + d\theta)$  and  $p(\theta)$  or  $x(\theta + d\theta)$  and  $x(\theta)$ , and  $\theta$  is a continuous parameter. An extensive property of the system can then be defined as a function of the state  $\{x_i\}$  such that

$$I = \sum_i x_i^2 \quad (5)$$

$$E = \langle e \rangle = \sum_i \epsilon_i x_i^2 \quad (6)$$

$$S = \langle s \rangle = \sum_i -x_i^2 \ln(x_i^2) \quad (7)$$

where  $\langle \dots \rangle$  indicates the ensemble average. The von Neumann formula for entropy is used, because as shown in [21], it has all the properties required by thermodynamics. The gradient of a given property in state space is then expressed by

$$\mathbf{g}_I = \sum_i \frac{\partial I}{\partial x_i} \hat{e}_i = \sum_i 2x_i \hat{e}_i \quad (8)$$

$$\mathbf{g}_E = \sum_i \frac{\partial E}{\partial x_i} \hat{e}_i = \sum_i 2\epsilon_i x_i \hat{e}_i \quad (9)$$

$$\mathbf{g}_S = \sum_i \frac{\partial S}{\partial x_i} \hat{e}_i = \sum_i [-2x_i - 2x_i \ln(x_i^2)] \hat{e}_i \quad (10)$$

where  $\hat{e}_i$  is the unit vector for each dimension. Furthermore, for an isolated system, the system satisfies

the conservation laws for probability and energy, i.e.,

$$I = \sum_i x_i^2 = 1, \quad E = \sum_i \epsilon_i x_i^2 = \text{constant} \quad (11)$$

The principle of SEA upon which the equation of motion is based is defined as the system state evolving along the direction that at any instant of time has the largest entropy gradient consistent with the conservation constraints. The equation of motion is, thus, given by

$$\frac{d\mathbf{x}}{dt} = \frac{1}{\tau(\mathbf{x})} \mathbf{g}_{S \perp L(\mathbf{g}_I, \mathbf{g}_E)} \quad (12)$$

where  $\tau$ , which is a function of system state, is the relaxation time which describes the speed at which the state evolves in state space in the direction of steepest entropy ascent.  $L(\mathbf{g}_I, \mathbf{g}_E)$  is the manifold spanned by  $\mathbf{g}_I$  and  $\mathbf{g}_E$ , and  $\mathbf{g}_{S \perp L(\mathbf{g}_I, \mathbf{g}_E)}$  is the perpendicular component of the gradient of the entropy to the hyper-surface that conserves the probability and energy. It takes the form of a ratio of Gram determinants expressed as

$$\mathbf{g}_{S \perp L(\mathbf{g}_I, \mathbf{g}_E)} = \frac{\begin{vmatrix} \mathbf{g}_S & \mathbf{g}_I & \mathbf{g}_E \\ (\mathbf{g}_S, \mathbf{g}_I) & (\mathbf{g}_I, \mathbf{g}_I) & (\mathbf{g}_E, \mathbf{g}_I) \\ (\mathbf{g}_S, \mathbf{g}_E) & (\mathbf{g}_I, \mathbf{g}_E) & (\mathbf{g}_E, \mathbf{g}_E) \end{vmatrix}}{\begin{vmatrix} (\mathbf{g}_I, \mathbf{g}_I) & (\mathbf{g}_E, \mathbf{g}_I) \\ (\mathbf{g}_I, \mathbf{g}_E) & (\mathbf{g}_E, \mathbf{g}_E) \end{vmatrix}} \quad (13)$$

where  $(\dots, \dots)$  denotes the scalar product of two vectors in state space. The explicit form of Eq. (13) for  $\{p_j\}$  is, thus, [20]

$$\frac{dp_j}{dt} = \frac{1}{\tau} \frac{\begin{vmatrix} -p_j \ln p_j & p_j & \epsilon_j p_j \\ \langle s \rangle & 1 & \langle e \rangle \\ \langle es \rangle & \langle e \rangle & \langle e^2 \rangle \end{vmatrix}}{\begin{vmatrix} 1 & \langle e \rangle \\ \langle e \rangle & \langle e^2 \rangle \end{vmatrix}} \quad (14)$$

where

$$\langle e^2 \rangle = \sum_i \epsilon_i^2 x_i^2, \quad \langle es \rangle = \sum_i -\epsilon_i x_i^2 \ln(x_i^2) \quad (15)$$

The state representation and the equation of motion can be simplified by combining degenerate energy eigenlevels [16]. The system is defined by a group of different energy eigenlevels  $\{\epsilon_i, i = 1, 2, \dots\}$  and their degeneracy  $\{n_i, i = 1, 2, \dots\}$ . The state of the system is described by a probability distribution among the energy eigenlevels  $\{p_i, i = 1, 2, \dots\}$  or square root of the probability  $\{x_i, i = 1, 2, \dots\}$ . As a result, the equation of motion changes to

$$\frac{dp_j}{dt} = \frac{1}{\tau} \frac{\begin{vmatrix} -p_j \ln \frac{p_j}{n_j} & p_j & \epsilon_j p_j \\ \langle s \rangle & 1 & \langle e \rangle \\ \langle es \rangle & \langle e \rangle & \langle e^2 \rangle \end{vmatrix}}{\begin{vmatrix} 1 & \langle e \rangle \\ \langle e \rangle & \langle e^2 \rangle \end{vmatrix}} \quad (16)$$

where the properties are defined by

$$\begin{aligned} \langle e \rangle &= \sum_i \epsilon_i x_i^2, & \langle s \rangle &= \sum_i -x_i^2 \ln\left(\frac{x_i^2}{n_i}\right) \\ \langle e^2 \rangle &= \sum_i \epsilon_i^2 x_i^2, & \langle es \rangle &= \sum_i -\epsilon_i x_i^2 \ln\left(\frac{x_i^2}{n_i}\right) \end{aligned} \quad (17)$$

### B. Nonequilibrium evolution: Kinetics and Dynamics

In general, the equation of motion for a system with a given group of conservation laws takes the form

$$\frac{dp_j}{dt} = \frac{1}{\tau(\mathbf{p})} D_j(\mathbf{p}) \quad (18)$$

where  $D_j(\mathbf{p})$  is calculated from the conservation laws and the principle of steepest entropy ascent [22]. Specifically, it takes the form of Eq. (14) for an isolate system yielding to mass and energy conservation.

Since for a given initial state of the system, the nonequilibrium thermodynamic path of state evolution is uniquely solved from this equation, Eq. (18), the path can be used to define a new parameter  $\tilde{\tau}$  given by

$$d\tilde{\tau} = \frac{1}{\tau(\mathbf{p}(t))} dt, \quad \text{or} \quad \tilde{\tau} = \int_{\text{path}} \frac{1}{\tau(\mathbf{p}(t'))} dt' = \tilde{\tau}(t) \quad (19)$$

where  $\tilde{\tau}$  is called the dimensionless time. With this time, the independent variable for the equation of motion can be changed so that

$$\frac{dp_j}{d\tilde{\tau}} = D_j(\mathbf{p}) \quad (20)$$

The solution of this equation is written as

$$p_j = p_j(\tilde{\tau}) \quad (21)$$

Independent of how the relaxation time  $\tau$  depends on the real time  $t$  and the state, the equation of motion can always be transformed to Eq. (20) with the parameter change defined by Eq. (19). Furthermore, the evolution of system state follows the same function [Eq. (21)] in  $\tilde{\tau}$ . Physically, this means that the system follows the same trajectory in state space. By using this transformation, the kinetics and dynamics of the system are separated. The former is found via Eqs. (20) and (21) and result in the trajectory in state space based on the parameter  $\tilde{\tau}$  or a constant relaxation time  $\tau$ . The dynamics are found via Eq. (18) and the functional dependence  $\tau = \tau(p)$  [Eq. (19)] and result in the trajectory in state space based on the real time  $t$ .

In the discussion on mass diffusion (Sec. III) and heat diffusion (Sec. IV), it is shown that the kinetics (or its associated trajectory) of the nonequilibrium relaxation results in a generalized Gibbs relation and the Onsager relations in the far-from-equilibrium realm and in linear

phenomenological equations in the near-equilibrium realm, which are independent of the dynamics. The kinetics appears as a system feature or pattern of the thermodynamics in the sense of the GENERIC [13, 14, 17], which is an ensemble or group feature. Information about the mechanical details (e.g., how the particles interact in the system mechanically) can be included in the dynamics (e.g., by how  $\tau$  is chosen) when the state evolution in time  $t$  is studied. The focus of this paper, however, is on the thermodynamic features of the nonequilibrium relaxation so that  $\tau$  is **assumed constant in the applications below**. For a brief discussion on how values for  $\tau$  can be determined at different levels of description, the reader is referred to *Appendix A*. For more discussion on the dynamics of nonequilibrium, especially on how  $\tau$  is chosen using the mechanics and how  $\tau$  is related to the state space geometry, the reader is referred to [18, 23].

### C. Nonequilibrium state and state evolution description: Hypoequilibrium

The solution of the SEA-QT equation of motion exhibits some good properties, which allows for a complete description of nonequilibrium state and the fundamental definition of nonequilibrium temperature. More discussion is presented in reference [16], and an example is provided below. The energy eigenlevels of the system  $\{\epsilon_i, i = 1, 2, 3, \dots\}$  with degeneracy  $\{n_i, i = 1, 2, 3, \dots\}$  can be divided into to  $M$  sets  $\{\epsilon_i^K\}$  (degeneracy  $\{n_i^K\}$ ) with  $i = 1, 2, 3, \dots, K = 1, 2, \dots, M$ , so that the state space of the system (the Hilbert space)  $\mathcal{H}$  can be represented by the sum of  $M$  subspaces  $\mathcal{H}_K$ , with  $K = 1, 2, \dots, M$ , i.e.,

$$\mathcal{H} = \bigoplus_{K=1}^M \mathcal{H}_K \quad (22)$$

To be complete,  $M$  can be infinite. The system state can be represented by the distributions in  $M$  subspace energy eigenlevels  $\{p_i^K, K = 1, \dots, M\}$ . If the probability distribution in one subspace, for example, the  $K$ th subspace yields to the canonical distribution of parameter  $\beta_K$ , the temperature of the  $K$ th subspace is defined to be  $T_K = 1/\beta_K$ , **where the Boltzmann constant is absorbed into temperature unit for simplicity**. Given a way to divide the energy eigenlevels, if the system probability distributions in the  $M$  subspaces are all canonical distribution, the state of the system is called a  $M$ th-order hypoequilibrium state [16], which can be described uniquely by the total probability in each subspace ( $\{p^K = \sum p_i^K, K = 1, \dots, M\}$ ) and the temperature of each of the subspaces ( $\{T_K, K = 1, \dots, M\}$ ). If the initial state of the system is a  $M$ th-order hypoequilibrium state then

$$p_i^K(t=0) = \frac{p^K n_i^K}{Z^K(\beta^K)} e^{-\beta^K \epsilon_i^K}, \quad i = 1, 2, 3, \dots \quad (23)$$



where  $Z^K(\beta^K)$  is the partition function of subspace  $K$  at temperature  $T^K$ . A more general form to represent any nonequilibrium state is given in [22] using the language of quantum mechanics. Li and von Spakovsky [16] have proven that the system retains a  $M$ th-order hypoequilibrium state throughout the nonequilibrium relaxation process if it initially starts out in such a state. The solution to Eq. (18), thus, becomes

$$p_i^K(t) = \frac{p^K(t)}{Z^K(\beta^K(t))} n_i^K e^{-\beta^K(t)\epsilon_i^K}, i = 1, 2, 3, \dots \quad (24)$$

As a result, each subspace has temperature defined fundamentally throughout the entire nonequilibrium relaxation process. This result applies to an isolated system with probability and energy conservation. For a system with a different set of conservation laws, a similar relation exists. However, the general proof is left for a future paper. The proof for a system with heat diffusion only is given in the Appendix B.

#### D. Local variational principle in thermodynamic state space

According to Beretta [20], the equation of motion can be derived from a local variational principle, which can be regarded as the variational form of the steepest entropy ascent principle, i.e.,

$$\begin{aligned} &\text{Maximize: } \dot{S}(\dot{\mathbf{x}}) = (\dot{\mathbf{x}}, \mathbf{g}_S) \text{ subject to} \\ &\dot{E} = (\dot{\mathbf{x}}, \mathbf{g}_E) = 0, \dot{I} = (\dot{\mathbf{x}}, \mathbf{g}_I) = 0, (\dot{\mathbf{x}}, \dot{\mathbf{x}}) = \xi(\mathbf{x}) \\ &\text{with } \delta\dot{\mathbf{x}} \neq 0, \delta\mathbf{x} = 0 \end{aligned} \quad (25)$$

The third constraint on  $\dot{\mathbf{x}}$  indicates that only the direction of  $\dot{\mathbf{x}}$  is of interest. This variational principle is in microscopic state space, which contrasts with the variational principle in the space spanned by conjugate fluxes and forces presented later for the Onsager relations.

### III. THEORY: DIFFUSION IN A NONEQUILIBRIUM SYSTEM

In this section, the theory for mass diffusion in a local, isolated system in nonequilibrium is presented. The Gibbs relation, the entropy generation for a non-quasi-equilibrium process, the Onsager relations and the quadratic dissipation potential are derived based on the concept of hypoequilibrium state and intensive properties. The variational principle using conjugate forces is given at the end of the section.

#### A. Equation of motion for mass diffusion

The mass (or probability) diffusion across energy eigenlevels (or across subspaces) can be studied for an

isolated system in nonequilibrium. Using Eq. (14), one energy eigenlevel in the  $K$ th subspace yields to the following equation of motion:

$$\frac{dp_j^K}{dt} = \frac{1}{\tau} \left( -p_j^K \ln \frac{p_j^K}{n_j^K} - p_j^K \frac{A_2}{A_1} + \epsilon_j p_j^K \frac{A_3}{A_1} \right) \quad (26)$$

where

$$A_1 = \begin{vmatrix} 1 & \langle e \rangle \\ \langle e \rangle & \langle e^2 \rangle \end{vmatrix}, A_2 = \begin{vmatrix} \langle s \rangle & \langle e \rangle \\ \langle es \rangle & \langle e^2 \rangle \end{vmatrix}, A_3 = \begin{vmatrix} \langle s \rangle & 1 \\ \langle es \rangle & \langle e \rangle \end{vmatrix} \quad (27)$$

Summation over all energy eigenlevels in this subspace yields the evolution of the probability in the subspace, represented by  $p^K$ , namely,

$$\frac{dp^K}{dt} = \frac{1}{\tau} \left( -p^K \ln p^K + p^K \langle \tilde{s} \rangle^K - p^K \frac{A_2}{A_1} + p^K \langle \tilde{e} \rangle^K \frac{A_3}{A_1} \right) \quad (28)$$

where  $\langle \cdot \rangle^K$  is the specific property in the  $K$ th subspace. To calculate the specific property of this subspace, the subspace's probability distribution is found from

$$p^K \equiv \sum_j p_j^K, \quad \tilde{p}_j^K \equiv \frac{p_j^K}{p^K} \quad (29)$$

where  $p^K$  is the particle number in the subspace. The specific property is then expressed as

$$\langle \tilde{e} \rangle^K \equiv \sum_j \epsilon_j^K \tilde{p}_j^K \quad (30)$$

$$\langle \tilde{s} \rangle^K \equiv - \sum_j \tilde{p}_j^K \ln \frac{\tilde{p}_j^K}{n_j^K} \quad (31)$$

#### B. Particle number and temperature evolution when the initial state is a hypoequilibrium state

If the system is in a  $M$ th-order hypoequilibrium state, the probability evolution yields Eq. (24). For simplicity, the following definition is made:

$$\alpha^K = \ln Z^K - \ln p^K \quad (32)$$

With this definition, the probability evolution of one energy eigenlevel is given by

$$\begin{aligned} p_i^K(t) &= \frac{p^K(t)}{Z^K(\beta^K(t))} n_i^K e^{-\beta^K(t)\epsilon_i^K} \\ &= n_i^K e^{-\alpha^K(t) - \beta^K(t)\epsilon_i^K} \end{aligned} \quad (33)$$

$\alpha^K$  and  $\beta^K$  are nonequilibrium intensive properties of the  $K$ th subspace, corresponding to the extensive properties  $p^K$  and  $E^K \equiv p^K \langle \tilde{e} \rangle^K$ . Furthermore, by defining

$$\alpha = \frac{A_2}{A_1}, \quad \beta = -\frac{A_3}{A_1} \quad (34)$$

the particle number and energy evolution of the  $K$ th subspace can be acquired from Eq. (26), i.e.,

$$\frac{dp^K}{dt} = \frac{1}{\tau}p^K(\alpha^K - \alpha) + \frac{1}{\tau}p^K\langle\tilde{e}\rangle^K(\beta^K - \beta) \quad (35)$$

$$\frac{d\langle e \rangle^K}{dt} = \frac{1}{\tau}p^K\langle\tilde{e}\rangle^K(\alpha^K - \alpha) + \frac{1}{\tau}p^K\langle\tilde{e}^2\rangle^K(\beta^K - \beta) \quad (36)$$

From Eqs. (24) and (26), the intensive properties  $\alpha^K$  and  $\beta^K$  obey the evolutions (see Appendix C for the derivation)

$$\frac{d\alpha^K}{dt} = -\frac{1}{\tau}(\alpha^K - \alpha) \quad (37)$$

$$\frac{d\beta^K}{dt} = -\frac{1}{\tau}(\beta^K - \beta) \quad (38)$$

The authors prove that  $\alpha$  and  $\beta$  have the physical meaning of intensive properties from measurements of a nonequilibrium state [24]. At stable equilibrium, the intensive properties in any subsystem obey the following relations:

$$\alpha(t = t^{eq}) = \alpha^K(t = t^{eq}) = \alpha^{eq} \quad (39)$$

$$\beta(t = t^{eq}) = \beta^K(t = t^{eq}) = \beta^{eq} \quad (40)$$

### C. Gibbs relation, entropy generation for a non-quasi-equilibrium process, the Onsager relations, and the quadratic dissipation potential in the nonlinear realm

Differential changes of the extensive properties in the  $K$ th subspace are written as

$$dE^K = \sum_i d(\epsilon_j^K p_j^K) \quad (41)$$

$$dS^K = \sum_i \frac{d}{dt}(-p_j^K \ln \frac{p_j^K}{n_j^K}) = \sum_i (-\ln \frac{p_j^K}{n_j^K} - 1) \frac{dp_j^K}{dt} \quad (42)$$

where  $E^K$  and  $S^K \equiv p^K\langle\tilde{s}\rangle^K$  are the energy and entropy in this subspace, respectively.

When a system is in a  $M$ th-order hypoequilibrium state and undergoes a pure relaxation process, a relation for property evolution in one subspace is acquired by using Eq. (33), namely,

$$\frac{dS^K}{dt} = \sum(\epsilon_j^K \beta^K + \alpha^K - 1) \frac{dp_j^K}{dt} = \beta^K \frac{dE^K}{dt} + (\alpha^K - 1) \frac{dp^K}{dt} \quad (43)$$

The proof of this relation for one subspace applies to any differential change (not only the time derivative). Thus, a generalization of the Gibbs relation to the  $K$ th subspace of a system in nonequilibrium is expressed by

$$dS^K = \beta^K dE^K + (\alpha^K - 1) dp^K \quad (44)$$

From the Gibbs relation at stable equilibrium written as

$$dS = \frac{1}{T} dE - \frac{\mu}{T} dN \quad (45)$$

the physical meaning of  $\beta^K$  and  $\alpha^K$  is shown to be

$$\beta^K = \left( \frac{\partial S^K}{\partial E^K} \right)_{p^K} = \frac{1}{T^K} \quad (46)$$

$$\alpha^K - 1 = \left( \frac{\partial S^K}{\partial p^K} \right)_{E^K} = -\frac{\mu^K}{T^K}, \quad \mu^K = \left( \frac{\partial E^K}{\partial p^K} \right)_{S^K} \quad (47)$$

where  $T^K$  is the subspace temperature and  $\mu^K$  is the subspace chemical potential with respect to subspace probability  $p^K$ . The total differential entropy change for the system, which for a pure nonequilibrium relaxation process corresponds to the entropy generation, is

$$\begin{aligned} dS &= \sum_K dS^K = \sum_K \beta^K dE^K + \sum_K (\alpha^K - 1) dp^K \\ &= \sum_K (\beta^K - \beta) dE^K + \sum_K (\alpha^K - \alpha) dp^K \end{aligned} \quad (48)$$

where both energy ( $\sum E^K = 0$ ) and probability ( $\sum p^K = 0$ ) conservations have been applied. **The rate of entropy generation can, thus, be written in terms of the internal fluxes of energy and probability inside the system  $J_E^K = dE^K/dt$  and  $J_p^K = dp^K/dt$  and the conjugate forces  $X_p^K = \beta^K - \beta$  and  $X_E^K = \alpha^K - \alpha$ .** The result is

$$\sigma(\mathbf{J}, \mathbf{X}) = \frac{dS}{dt} = \sum_K X_E^K J_E^K + \sum_K X_p^K J_p^K \quad (49)$$

The Onsager relations are then acquired from Eqs. (35) and (36) in the form of  $\mathbf{J} = \mathbf{\Lambda X}$ , where  $\mathbf{\Lambda}$  is a symmetric and positive definite operator. Thus,

$$J_p^K = \frac{1}{\tau} p^K X_p^K + \frac{1}{\tau} E^K X_E^K \quad (50)$$

$$J_E^K = \frac{1}{\tau} E^K X_p^K + \frac{1}{\tau} \langle e^2 \rangle^K X_E^K \quad (51)$$

The quadratic dissipation potential using force representation [9, 25] is then written as

$$\begin{aligned} \Xi(\mathbf{X}, \mathbf{X}) &= \frac{1}{2} \langle \mathbf{X}, \mathbf{\Lambda X} \rangle = \frac{1}{2\tau} \sum_K [p^K (\alpha^K - \alpha)^2 \\ &\quad + 2E^K (\alpha^K - \alpha)(\beta^K - \beta) + \langle e^2 \rangle^K (\beta^K - \beta)^2] \end{aligned} \quad (52)$$

while the variational principle using force representation is expressed as

$$\delta[\sigma(\mathbf{J}, \mathbf{X}) - \Xi(\mathbf{X}, \mathbf{X})]_{\mathbf{J}} = 0, \quad \mathbf{J} = \text{const}, \quad \delta\mathbf{J} = 0, \quad \delta\mathbf{X} \neq 0 \quad (53)$$

where the  $\sigma(\mathbf{J}, \mathbf{X})$  and  $\Xi(\mathbf{X}, \mathbf{X})$  are given by Eqs. (49) and (52). Furthermore, even though the following constraints apply to the fluxes:

$$\sum_K J_p^K = 0, \quad \sum_K J_E^K = 0 \quad (54)$$

the reciprocity seen in Eqs. (50) and (51) is completely consistent with the Onsager theory since according to Gyarmati [9], *the validity of Onsager's reciprocal*

relations is not influenced by a linear homogeneous dependence valid amongst the fluxes. Therefore, the physical interpretation given here is fully compatible with other investigations [24] and does not require a reformulation in terms of independent fluxes even though this could be done. In addition, it is from the gradient dynamics of the nonequilibrium relaxation process that the entropy generation, the Onsager relations, and the quadratic dissipation potential of a local, isolated system in nonequilibrium have been derived using the geometric principle of SEA as well as the concepts of hypoequilibrium state and nonequilibrium intensive properties. Alternatively, the variational principle of SEA in system state space could be used to arrive at these relations as is done in [22] using the language of quantum mechanics. Of course, these relations also correspond to the variational principle in the space spanned by conjugate forces and fluxes [9].

#### IV. THEORY: HEAT DIFFUSION IN A NONEQUILIBRIUM SYSTEM

A local, isolated system in nonequilibrium with heat diffusion only is considered in this section. This requires a model with a different set of constraints (i.e., the probability redistribution is only allowed in each subspace) than when heat and mass diffusion are both considered. The entropy change of the system and subspaces due to heat diffusion for a non-quasi-equilibrium process is given and the relationship between the SEAQT equation of motion and the phenomenological diffusion equation is presented.

##### A. Equation of motion for heat diffusion

Different from previous forms of the equation of motion, the form for pure heat diffusion yields to a different set of conservation equations. If the system is separated into  $M$  subspaces with energy flow but no probability flow across the subspaces, there are  $M + 1$  conservation laws. System probability conservation is replaced by that for  $M$  individual subspaces. In *Appendix B*, it is proven that the concept of hypoequilibrium state and nonequilibrium temperature are also well defined under these new constraints given by

$$I^K = \sum_i (x_i^K)^2 = p^K, \quad K = 1, 2, \dots, M \quad (55)$$

$$E = \sum_i \epsilon_i x_i^2 = \text{constant} \quad (56)$$

For simplicity, a second-order hypoequilibrium state is studied first. The system is separated into 2 subspaces (subspace  $a$  and subspace  $b$ ) so that the equation of

motion takes the form

$$\frac{dp_j^a}{dt} = \frac{1}{\tau} \begin{vmatrix} p_j^a s_j^a & p_j^a & 0 & \epsilon_j^a p_j^a \\ \langle s \rangle^a & p^a & 0 & \langle e \rangle^a \\ \langle s \rangle^b & 0 & p^b & \langle e \rangle^b \\ \langle es \rangle & \langle e \rangle^a & \langle e \rangle^b & \langle e^2 \rangle \end{vmatrix} \quad (57)$$

where the contribution of each subspace to the total property is defined by

$$\langle s \rangle = \langle s \rangle^a + \langle s \rangle^b, \quad \langle s \rangle^a = \sum p_i^a s_i^a, \quad \langle s \rangle^b = \sum p_i^b s_i^b \quad (58)$$

$$\langle e \rangle = \langle e \rangle^a + \langle e \rangle^b, \quad \langle e \rangle^a = \sum p_i^a \epsilon_i^a, \quad \langle e \rangle^b = \sum p_i^b \epsilon_i^b \quad (59)$$

and where

$$s_j^{a(b)} = -\ln \frac{p_j^{a(b)}}{n_j^{a(b)}} = \tilde{s}_j^{a(b)} - \ln p^{a(b)}, \quad \tilde{s}_j^{a(b)} \equiv -\ln \frac{\tilde{p}_j^{a(b)}}{n_j^{a(b)}} \quad (60)$$

By defining

$$B_1 = \begin{vmatrix} p^a & 0 & \langle e \rangle^a \\ 0 & p^b & \langle e \rangle^b \\ \langle e \rangle^a & \langle e \rangle^b & \langle e^2 \rangle \end{vmatrix}, \quad B_2 = \begin{vmatrix} \langle s \rangle^a & 0 & \langle e \rangle^a \\ \langle s \rangle^b & p^b & \langle e \rangle^b \\ \langle es \rangle & \langle e \rangle^b & \langle e^2 \rangle \end{vmatrix} \\ B_2^b = \begin{vmatrix} \langle s \rangle^b & 0 & \langle e \rangle^b \\ \langle s \rangle^a & p^a & \langle e \rangle^a \\ \langle es \rangle & \langle e \rangle^a & \langle e^2 \rangle \end{vmatrix}, \quad B_3 = \begin{vmatrix} \langle s \rangle^a & p^a & 0 \\ \langle s \rangle^b & 0 & p^b \\ \langle es \rangle & \langle e \rangle^a & \langle e \rangle^b \end{vmatrix} \quad (61)$$

Eq. (57) can be simplified to

$$\frac{dp_j^a}{dt} = \frac{1}{\tau} (p_j^a s_j^a - p_j^a \frac{B_2^a}{B_1} - \epsilon_j^a p_j^a \frac{B_3}{B_1}) \quad (62)$$

Moreover, the equation of motion for the probability distribution in one subspace can also be written in terms of the normalized probability by dividing both sides of Eq. (62) by  $p^a$  so that

$$\frac{d\tilde{p}_j^a}{dt} = \frac{1}{\tau} (\tilde{p}_j^a s_j^a - \tilde{p}_j^a \frac{B_2^a}{B_1} - \epsilon_j^a \tilde{p}_j^a \frac{B_3}{B_1}) \quad (63)$$

Furthermore, if the system is in a second-order hypoequilibrium state initially so that each subspace has a canonical distribution, Eq. (63) can be simplified further to arrive at the form

$$\frac{d\tilde{p}_j^a}{dt} = \frac{1}{\tau} \tilde{p}_j^a [(\tilde{s}_j^a - \langle \tilde{s} \rangle^a) - \beta(\epsilon_j^a - \langle \tilde{e} \rangle^a)] \quad (64)$$

where  $\tilde{s}_j^a$  is defined by Eq. (60) and  $\langle \tilde{s} \rangle^a$  and  $\langle \tilde{e} \rangle^a$  are defined by Eqs. (30) and (31). The parameter  $\beta$  given

$$\beta \equiv \frac{B_3}{B_1} = \frac{p^a \tilde{A}_1^a \beta^a + p^b \tilde{A}_1^b \beta^b}{p^a \tilde{A}_1^a + p^b \tilde{A}_1^b} \quad (65)$$

$$B_1 = p^a p^b (p^a \tilde{A}_1^a + p^b \tilde{A}_1^b) \quad (66)$$

$$B_3 = p^a p^b (p^a \beta^a \tilde{A}_1^a + p^b \beta^b \tilde{A}_1^b) \quad (67)$$



is a weighted average of the inverse temperatures of the subsystems relative to the mole fractions and the energy fluctuation (or nondimensional specific heat at constant volume) of the subspaces written as

$$\tilde{A}_1^{a(b)} = \langle \tilde{e}^2 \rangle^{a(b)} - (\langle \tilde{e} \rangle^{a(b)})^2 = -\frac{\partial \langle \tilde{e} \rangle^{a(b)}}{\partial \beta^{a(b)}} = \frac{C_V^{a(b)}}{(\beta^{a(b)})^2} \quad (68)$$

$$C_V^{a(b)} \equiv \frac{1}{k_b} \frac{\partial \langle \tilde{e} \rangle^{a(b)}}{\partial T^{a(b)}} \quad (69)$$

For the more general case of a  $M$ th-order hypoequilibrium state and the system separated into  $M$  subspaces, equation (63) remains the same but with

$$\beta \equiv \frac{B_3}{B_1} = \frac{\sum_K^M p^K \tilde{A}_1^K \beta^K}{\sum_K^M p^K \tilde{A}_1^K} \quad (70)$$

A given interaction type (e.g., heat diffusion) results in a given relaxation time  $\tau$  (see Sec. II.B), while the ratio  $B_3/B_1$  provides an average temperature based on subspace mole fractions and energy fluctuations. At stable equilibrium,  $\beta = \beta^{eq}$ . Now, if one subspace  $R$  is attached to a reservoir, the evolution of the other subspaces behave according to the equation of motion, Eq. (63), with  $\beta^R$  constant. For example, only part of the energy eigenlevels can absorb energy from the environment. Mathematically, if subspace  $R$  yields to one of two conditions,

$$\forall K \neq R, \quad C_V^R \gg C_V^K, \quad p^R \gg p^K \quad (71)$$

the relation  $\beta = \beta^R$  holds and subspace  $K (\neq R)$  yields to the equation of motion.

$$\frac{d\tilde{p}_j^K}{dt} = \frac{1}{\tau} \tilde{p}_j^K [(-\ln \frac{\tilde{p}_j^K}{n_j^K} - \langle \tilde{s} \rangle^K) - \beta^R (\epsilon_j^K - \langle \tilde{e} \rangle^K)] \quad (72)$$

Note that in this equation, the only parameter related to subspace  $R$  is the reservoir temperature  $\beta^R$ . The energy eigenstructure of subspace  $R$  plays no role!

### B. Property of heat diffusion: non-quasi-equilibrium processes and the second law of thermodynamics

Based on Eq. (62), the total entropy and energy evolution in one subspace can be determined via

$$\frac{dS^K}{dt} = \frac{d\langle s \rangle^K}{dt} = p^K \frac{d\langle \tilde{s} \rangle^K}{dt} = \frac{1}{\tau} p^K (\beta^K - \beta) \beta^K \tilde{A}_1^K \quad (73)$$

$$\frac{dE^K}{dt} = \frac{d\langle e \rangle^K}{dt} = p^K \frac{d\langle \tilde{e} \rangle^K}{dt} = \frac{1}{\tau} p^K (\beta^K - \beta) \tilde{A}_1^K \quad (74)$$

Dividing Eq. (73) by (74) yields

$$\frac{dS^K}{dE^K} = \frac{dS^K}{dt} / \frac{dE^K}{dt} = \beta^K \quad (75)$$

The equation is a generalized form of the differential entropy transfer due to heat diffusion using the nonequilibrium temperature for each subspace, i.e.,

$$dS^K = \beta^K dE^K = \frac{\delta Q^K}{T} \quad (76)$$

Moreover, Eq. (76) can be applied to all kinds of thermodynamic processes and is not limited to quasi-equilibrium processes. This argument comes from the universal definition of nonequilibrium temperature provided in this paper.

## V. MODEL: COMPOSITE SYSTEM IN A NONEQUILIBRIUM STATE

Interacting systems can form a composite nonequilibrium system with the interaction resulting in the nonequilibrium relaxation process for the composite. Using the SEA equation of motion, the state evolution of the composite system can be determined. Proper division of this composite allows the subspaces to be viewed as interacting subsystems within the composite. The behavior of each subsystem, thus, can be studied by an analysis of the state evolution of each subspace. In particular, if two individual subsystems involved in an interaction both have canonical state distributions, and each subsystems energy eigenvectors span one subspace, the composite system is in a second order hypoequilibrium state. On the other hand, if each individual subsystem's state cannot be described by a canonical distribution, a higher-order hypoequilibrium state will be required.

In the following section (Sec. V.A), the SEAQT framework using single particle energy eigenlevels is applied to the study of a simple system. Section V.B then explains the process of subspace division followed in Sec. V.C by a comparison with the phenomenological equations for mass and heat diffusion. In Sec. V.D, the physical details of the system used are described. Finally, in Sec. V.E, the coupling of mass and heat diffusion is modeled and discussed.

### A. Multi-particle classical simple system

Theoretically and in general, the SEAQT equation of motion is applicable to multi-particle systems provided the energy eigenstructure of the system is known [24]. However, for a multi-particle classical simple system, the energy eigenstructure of a single particle and its associated equation of motion can be used to study the system, since all of the particles (or particle groups) have the same energy eigenlevels  $\{\epsilon_i, i = 1, 2, 3, \dots\}$  and degeneracy  $\{n_i, i = 1, 2, 3, \dots\}$ . Thus, the system state can be represented by the particle number (or particle group number) at each energy eigenlevel  $\{m_i, i = 1, 2, 3, \dots\}$ . The mole fraction of particles at the  $i$ th

energy eigenlevel is given by  $y_i = m_i / \sum m_i$ . The extensive property constraints of the system are then

$$M = \sum_i m_i = \text{constant} \quad (77)$$

$$E = \sum_i \epsilon_i m_i = \text{constant} \quad (78)$$

$$S = \sum_i -m_i \ln \frac{y_i}{n_i} \quad (79)$$

Dividing the constraints by the total particle number  $\sum m_i$ , the system state can be represented by the mole fractions  $\{y_i, i = 1, 2, 3, \dots\}$ , which are equivalent to the single-particle probability distribution  $\{p_i, i = 1, 2, 3, \dots\}$ . For an isolated system the constraints become

$$I = \sum_i y_i = \sum_i p_i = \text{constant} \quad (80)$$

$$\langle e \rangle = \frac{E}{M} = \sum_i \epsilon_i y_i = \sum_i \epsilon_i p_i = \text{constant} \quad (81)$$

$$\langle s \rangle = \frac{S}{M} = \sum_i -y_i \ln \frac{y_i}{n_i} = \sum_i -p_i \ln \frac{p_i}{n_i} \quad (82)$$

Here  $\langle \dots \rangle$  represents an average specific property. The constraints and equation of motion reduce to the single-particle case (Eqs. (5) to (7) and (14)) when  $y_i = p_i, i = 1, 2, 3, \dots$

In general, a systems energy eigenstructure and extensive properties are a function of the total particle number of each constituent. This is also true of its specific properties in the presence of a mass interaction or chemical reaction if the system is partitioned and not simple [26], since partitioning influences each partition's energy eigenstructure. Thus, for mass diffusion, the framework outlined here requires an invariant eigenstructure and as a consequence the simple system assumption. With this assumption, particle number no longer influences the specific properties. This same assumption, however, is not required in the case of heat diffusion since the total number of particles for each system partition (i.e., subsystem) does not change.

## B. Interacting systems

It is assumed that a group of observable operators  $\hat{F}$  commuting with the Hamiltonian operator  $\hat{H}$  exists. The degenerate energy eigenlevels of the system can be distinguished by eigenvalues of the observations of  $\hat{F}$ , which have values  $F_1, F_2, \dots, F_M$  so that the system energy eigenlevels can be separated into  $M$  sets  $\{\epsilon_i^K, i = 1, 2, 3, \dots\}$  with degeneracy  $\{n_i^K, i = 1, 2, 3, \dots\}$ , where  $K = 1, \dots, M$ . In each set, every energy eigenlevel represents an eigenstate common to both  $\hat{F}$  and  $\hat{H}$  with the same eigenvalue of  $\hat{F}$ . Eigenstates in each of the sets can be spanned into a subspace of the system state space and can be designated as a subsystem.




FIG. 1. For mass diffusion, mass flow and energy flow are both allowed across the partition. For heat diffusion, only energy flow is allowed.

Practically, by choosing the observable operator, the system can be viewed as a composite system whose subsystems can be properly arranged to study specific phenomena. For the kinetics of a chemical reaction,  $F$  can be chosen to be an observable operator of species, whose eigenvalues are “*Reactant*” and “*Product*”. In heat and mass diffusion, the observable operator  $F$  is chosen to be the relative location to a partition, with eigenvalue of “*left*” (left of the partition) and “*right*” (right of the partition). The partition allows mass and heat diffusion (see Fig. 1). To be more precise, it is assumed that the de Broglie wavelength  $\lambda^d$  is much smaller than the distance  $\delta x$  between the center of the subsystem on the “*left*” and that on the “*right*”. This wavelength represents the classical limit, i.e.,

$$p_v = \sqrt{2mk_bT}, \quad \lambda^d = \frac{h}{p_v} = \frac{h}{\sqrt{2mk_bT}}, \quad \delta x \gg \lambda \quad (83)$$

where  $p_v$  is the expectation value of the particle momentum for a system at temperature  $T$ .

## C. Phenomenological transport equation

With the assumption that the two subspaces of the system are two subsystems at two positions, the phenomenological transport equations of mass and heat diffusion can be derived.

### 1. Mass diffusion

Using Eq. (28), the equations of motion for two subspaces are written as

$$\frac{d \ln p^a}{dt} = -\frac{1}{\tau} \ln p^a + \frac{1}{\tau} (\langle \tilde{s} \rangle^a - \frac{A_2}{A_1} + \langle \tilde{e} \rangle^a \frac{A_3}{A_1}) \quad (84)$$

$$\frac{d \ln p^b}{dt} = -\frac{1}{\tau} \ln p^b + \frac{1}{\tau} (\langle \tilde{s} \rangle^b - \frac{A_2}{A_1} + \langle \tilde{e} \rangle^b \frac{A_3}{A_1}) \quad (85)$$

When the two subspaces of the system have the same eigenstructure and temperature, subtracting Eq. (85)

from Eq. (84) yields

$$\frac{d}{dt}(\ln p^a - \ln p^b) = -\frac{1}{\tau}(\ln p^a - \ln p^b) \quad (86)$$

Substituting the subspace probability (or mole fraction) given by Eq. (87) into Eq. (86) results in

$$p^a = \frac{m^a}{m^a + m^b} \quad p^b = \frac{m^b}{m^a + m^b} \quad (87)$$

$$\frac{d}{dt}(\ln \frac{m^a}{m^b}) = -\frac{1}{\tau}(\ln \frac{m^a}{m^b}) \quad (88)$$

where  $m^{a(b)}$  is the particle number in subspace  $a(b)$  as defined in Sec. V.A. If the global mass distribution is continuous and position A and position B are close enough,

$$m^a = m^b + \Delta m \quad (89)$$

which transforms Eq. (88) into

$$\frac{d}{dt}(\frac{\Delta m}{m^b}) = -\frac{1}{\tau} \frac{\Delta m}{m^b} \quad (90)$$

$$\begin{aligned} J^{a(b)} &\equiv \frac{1}{2A} \frac{d}{dt}(m^a - m^b) = -\frac{m^a - m^b}{2\tau A} = -\frac{\delta x}{2\tau A} \frac{dm}{dx} \\ &= -\frac{(\delta x)^2}{2\tau} \frac{dc}{dx} \end{aligned} \quad (91)$$

where the approximation  $\ln(1+x) \simeq x$  for small  $x$  has been used and higher-order terms dropped to arrive at Eq. (90). In Eq. (91),  $\delta x$  is the distance between two positions,  $A$  is the cross-sectional area of the interacting surface, and  $J^{a(b)}$  is the flux of particle number equal to  $(dm^a/dt)/A$  or  $(-dm^b/dt)/A$  (hence the division by 2 in Eq. (91)). With  $c$  as the concentration, Eq. (91) recovers Fick's law with the diffusion coefficient (diffusivity) given by  $D = \frac{(\delta x)^2}{2\tau}$ . The specific form of  $D$  is directly related to the form of  $\tau$ , which contains the detailed mechanical information (see Sec. V.D for an example). The phenomenological linear equation can be derived without the form of  $\tau$ , which is a pure thermodynamic feature or pattern of the nonequilibrium relaxation process. In addition to results such as these for the near-equilibrium realm, thermodynamic features or patterns in the far-from-equilibrium realm can also be studied using Eqs. (35) and (36) provided the initial state is a hypoequilibrium state. For the case when it is not, Eq. (14) can directly be used as is done, for example, in [16].

## 2. Heat diffusion

For a system in which the only interaction is that of heat diffusion, Eq. (74) captures the energy flow between two subsystems, i.e.,

$$\begin{aligned} J_E^{a(b)} &= -\kappa' \delta x A \frac{dT}{dx} = \frac{1}{\tau} p^a (\beta^a - \beta) \tilde{A}_1^a \\ &= \frac{1}{\tau} \frac{p^a p^b \tilde{A}_1^a \tilde{A}_1^b}{p^a \tilde{A}_1^a + p^b \tilde{A}_1^b} (\beta^a - \beta^b) \end{aligned} \quad (92)$$

where  $J_E^{a(b)}$  is the heat flux or the rate of energy transferred per unit area,  $A$  is the cross-sectional area of the interacting surface,  $T^a$  and  $T^b$  are the temperatures of the two subsystems, and Eq. (65) has been substituted for  $\beta$ . Eq. (92) recovers Fourier's law of heat diffusion (conduction). The thermal conductivity per unit length  $\kappa'$  and the thermal conductivity  $\kappa$  are expressed as

$$\kappa' = \frac{1}{\tau} \frac{p^a p^b \beta^a \beta^b \tilde{A}_1^a \tilde{A}_1^b}{p^a \tilde{A}_1^a + p^b \tilde{A}_1^b} \frac{1}{A}, \quad \kappa = \kappa' \delta x \quad (93)$$

In the near-equilibrium region with the same constituent in both subsystems,

$$p^a = p^b = \frac{1}{2}, \quad C_V^a = C_V^b = C_V \quad (94)$$

and the thermal conductivity per unit length and the thermal conductivity are expressed in terms of the energy fluctuation (or nondimensional specific heat at constant volume) of the subspaces, i.e.,

$$\kappa' = \frac{1}{2\tau} C_V \frac{1}{A}, \quad \kappa = \frac{1}{2\tau} \frac{\delta x}{A} C_V \quad (95)$$

The above formulation is applicable for any kind of interaction resulting in a flow of energy only. Furthermore, if the heat and mass diffusion are affected via the same kind of micro-mechanical interactions such as the collision of particles, it can be assumed that the same  $\tau$  is applicable when the system is in the near-equilibrium region close to the same stable equilibrium point. In this case,  $\kappa = C_V D/V$ , where  $V = A \delta x$  is the volume of the subsystem. This last result is the same as that found from classical transport theory and is a direct consequence of the thermodynamic features of the system minus any direct knowledge of the details of the micro-mechanical interactions taking place.

## D. Mass and heat diffusion of hydrogen

To model the mass and heat diffusion for a specific case, a composite system of hydrogen is set up with two subspaces corresponding to subsystems on two sides of a partition. The energy eigenlevels of the two subsystems together form the energy eigenlevels for the composite system as a whole. Denoting the state space of the subsystem on the "left" by  $\mathcal{H}^a$  and that on the "right" by  $\mathcal{H}^b$ , the composite system state space  $\mathcal{H}$  takes the form

$$\mathcal{H} = \mathcal{H}^a \oplus \mathcal{H}^b \quad (96)$$

The available energy eigenvalues for one subspace ("left" or "right") are constructed from the energy eigenvalues of each degree of freedom for translation and rotation, i.e., from

$$\epsilon^{a(b)} = \epsilon_{t,H_2} + \epsilon_{r,H_2} \quad (97)$$

The translational energy eigenvalue  $\epsilon_t$  uses the form of the infinite potential well, while the rotational energy eigenvalue  $\epsilon_r$  uses the form of the rigid motor. These are expressed as follow:

$$\epsilon_t(n_x, n_y, n_z) = \frac{\hbar^2}{8m} \left( \frac{n_x^2}{L_x^2} + \frac{n_y^2}{L_y^2} + \frac{n_z^2}{L_z^2} \right) \quad (98)$$

$$\epsilon_r(j, m) = \frac{j(j+1)\hbar^2}{2I} \quad (99)$$

where  $n_x$ ,  $n_y$ , and  $n_z$  are the quantum numbers for the translational degrees of freedom,  $j$  and  $m$  are the quantum numbers for the rotational degrees of freedom,  $I$  is the moment of inertia and  $L_x$ ,  $L_y$ , and  $L_z$  are chosen based on the characteristic lengths of the particle container, which are not necessarily the dimensions of each subsystem, since each subsystem can be a local control volume within the particle container. Furthermore, the vibrational energy is not included since as is shown by the authors in [16], at the temperature considered in this study, the vibrational contribution is small. The disassociation energy is also not included via the selection of the proper energy reference. Each combination of quantum numbers and position corresponds to one energy eigenlevel in the subspaces (or subsystems). The composite system energy eigenlevels are formed by all the available energy eigenlevels of the “left” and the “right”.

The evolution in real time can be studied by choosing  $\tau$  based on Eq. (91) or (95) with the help of experimental data or from *ab initio* calculation using quantum mechanics or classical mechanics. Utilizing Eq. (95)

$$\tau = \frac{(\delta x)^2}{2D} \quad (100)$$

where  $\delta x$  is the distance between the centers of the subsystems and  $D$ , which is a function of pressure  $P$  and temperature  $T$ , is the self-diffusion coefficient. For the present study, the dependence of the diffusion coefficient for gases on temperature uses the function provided by the Chapman-Enskog theory. For binary diffusion of  $A$  and  $B$ , the diffusion coefficient is

$$D_{AB} = \frac{0.0018583T^{3/2}\sqrt{1/M_A + 1/M_B}}{P\sigma_{AB}^2\Omega} \quad (101)$$

where  $M_A$  and  $M_B$  are the standard molecular weights,  $P$  [atm] is the pressure,  $T$  [K] the temperature,  $\sigma_{AB}$  [Å] the average collision diameter, and  $\Omega$  is a temperature-dependent collision integral. In the case of self-diffusion for hydrogen, the parameters of  $A$  and  $B$  are both chosen to be that of the hydrogen molecule  $H_2$ , while  $\sigma_{AB}$  and  $\Omega$  are tabulated [27]. For simplicity, we use the self-diffusion coefficient for hydrogen at equilibrium given by  $D = D(P^{eq}, T^{eq})$  to scale our  $\tau$  and provide a time scale for the entropy generation. However, since the focus of our paper is the kinetics (thermodynamic path)

of the diffusion process and not the dynamics, a more detailed discussion of the nonequilibrium effects related to the dynamics of a time-dependent  $\tau$  is left for a future paper [23]. A brief discussion of how values for  $\tau$  can be determined at different levels of description is, nonetheless, given in Appendix A.

A second-order hypoequilibrium state with the subspace division of “left” and “right” is chosen to be the initial condition, which means that the two subspaces (subsystems) are each in a local equilibrium state. The non-quasi-equilibrium process of mass and heat diffusion is studied using Eqs. (26) and (62). For the case when the subsystems are not in states of local equilibrium, the two subspaces can be divided even further. For example, if the “left” subsystem is a  $M$ th-order hypoequilibrium state, this subspace, subspace  $a$ , can be divided into  $M$  subspaces based on the initial condition. However, the evolution of each subspace, regardless of whether or not the subsystem is in a state of local equilibrium, yields to the same form of the equations of motion, Eqs. (26) and (62). For the case considered here, the initial condition is given by Eq. (23), i.e., by

$$p_i^{a(b)}(t=0) = \frac{p^{a(b)}n_i^{a(b)}}{Z^{a(b)}(\beta^{a(b)})} e^{-\beta^{a(b)}\epsilon_i^{a(b)}} \quad (102)$$

The time evolution is acquired by solving Eqs. (26) and (62). For a more general initial condition, such as that for an infinite-order hypoequilibrium state, the equation of motion can be solved using the density of states method developed in [16].

The specific properties of the individual subsystems at a given temperature and volume are expressed as

$$\begin{aligned} Z^{a(b)}(\beta^{a(b)}, V) &= Z^t Z^r = V \left( \frac{m}{2\pi\hbar^2\beta^{a(b)}} \right)^{\frac{3}{2}} \frac{2I}{\beta^{a(b)}\hbar^2} \\ &= C_Z V (\beta^{a(b)})^{-5/2} \end{aligned} \quad (103)$$

$$\langle \tilde{\epsilon} \rangle^{a(b)}(\beta^{a(b)}) = \frac{5}{2} k_b T^{a(b)} = \frac{5}{2\beta^{a(b)}} = \frac{C_V}{\beta^{a(b)}} \quad (104)$$

$$\begin{aligned} \langle \tilde{s} \rangle^{a(b)}(\beta^{a(b)}, V) &= \beta^{a(b)} \langle \tilde{\epsilon} \rangle^{a(b)} + \ln Z^{a(b)} \\ &= -\frac{5}{2} \ln \beta^{a(b)} + \ln V + C_s \\ &= -C_V \ln \beta^{a(b)} + \ln V + C_s \end{aligned} \quad (105)$$

where  $C_Z$  and  $C_s$  are constants determined from Eqs. (103) to (105).

## E. Mass and heat diffusion coupling

For mass diffusion with no temperature difference, as shown in Eq. (38), the temperature difference remains zero, and only the particle number difference changes and Eq. (35) reverts to the linear transport equation in the near-equilibrium realm. The more complex case occurs when mass diffusion takes place in the presence

of a temperature difference in which case coupling effects may be present.

The mass flow between two subsystems is determined by subtracting the probability evolution of one [Eq. (35)] from the other with the result that

$$2J_p^{b \rightarrow a} = \frac{dp^a}{dt} - \frac{dp^b}{dt} = \frac{1}{\tau} p^a (\alpha^a - \alpha) + \frac{1}{\tau} p^a \langle \tilde{\epsilon} \rangle^a (\beta^a - \beta) - \frac{1}{\tau} p^b (\alpha^b - \alpha) - \frac{1}{\tau} p^b \langle \tilde{\epsilon} \rangle^b (\beta^b - \beta) \quad (106)$$

The coupling effects come from the differences in both of the nonequilibrium intensive properties  $\alpha$  and  $\beta$ . To study the effect of temperature on the probability (mass) flow, the two subsystems start from the same initial probability ( $p^a = p^b = p^{eq} = 0.5$ ) but different temperatures. For the case of a perfect gas (i.e., ideal gas with constant specific heats), the final stable equilibrium temperature  $T^{eq}$  is an average of the initial temperatures of the two subsystems, namely,

$$T^{eq} = (T^a + T^b)/2, \quad \xi \equiv \Delta T / T^{eq} \\ T^a = T^{eq} + \Delta T, \quad T^b = T^{eq} - \Delta T, \quad (107)$$

Thus, Eq. (106) can be simplified to

$$J_p^{b \rightarrow a} = \frac{1}{2\tau} p^{eq} [(\langle \tilde{s} \rangle^a - \langle \tilde{s} \rangle^b) - \beta(\langle \tilde{\epsilon} \rangle^a - \langle \tilde{\epsilon} \rangle^b)] \quad (108)$$

where  $\langle \tilde{s} \rangle^{a(b)}$  and  $\langle \tilde{\epsilon} \rangle^{a(b)}$  are the specific entropy and energy of subsystem a(b) defined by Eqs. (30) and (31). Substituting Eqs. (104) and (105) yields

$$J_p^{b \rightarrow a} = \frac{1}{2\tau} p^{eq} [C_V (\ln \frac{1}{\beta^a} - \ln \frac{1}{\beta^b}) - \beta (\frac{C_V}{\beta^a} - \frac{C_V}{\beta^b})] \\ = \frac{C_V p^{eq}}{2\tau} [(\ln T^a - \ln T^b) - \frac{1}{T} (T_a - T_b)] \quad (109)$$

where  $T = 1/(k_b \beta)$  and the relation  $T/T^{eq} = 1 + \xi^2 + O(\xi^4)$  holds when  $p^a = p^b$ . The mass (probability) flux due to a temperature difference is then written as

$$J_p^{b \rightarrow a} = \frac{dp^a}{dt} \frac{C_V p^{eq}}{\tau} \left[ \frac{\Delta T}{T^{eq}} + \frac{\Delta T^3}{(T^{eq})^3} - \frac{\Delta T}{T} \right] \\ = \frac{4}{3} \frac{C_V p^{eq}}{\tau} \left( \frac{\Delta T}{T^{eq}} \right)^3 + O(\lambda^5) \quad (110)$$

and the temperature evolution, Eq. (38), reduces to

$$\frac{dT^a}{dt} = -\frac{1}{\tau} \Delta T \quad (111)$$

In the near-equilibrium realm where only small temperature differences exist ( $\xi \ll 1$ ), higher-order nonlinear temperature difference effects, which influence the probability (mass) flux, are negligible and can, thus, be ignored. For this case, the temperature evolution equation [Eq. (111)] and the probability (mass) evolution equation [Eq. (110)] due to a temperature difference are effectively decoupled. In the far-from-equilibrium realm, however, higher-order temperature




FIG. 2. (color online) Thermodynamic trajectories on a temperature-particle number diagram. The initial probabilities for the two subsystems [black line for subsystem a, green (gray) line for subsystem b] are the same, while the initial temperature of subsystem b is 1500 K, while that for subsystem a is 300 K (solid line), 500 K (dashed line), 800 K (dashed-dotted line) and 1100 K (dotted line), respectively.

difference nonlinearities may be significant in which case coupling effects become important. In Fig. 2, the thermodynamic trajectories of three different cases for which the initial probabilities for the two subsystems are the same are plotted on a temperature-particle number diagram. For each case, the trajectory consists of two lines, one for each subsystem, with each point on each line representing an intermediate state for a given subsystem. The two subsystems start from opposite ends of the two colored lines and evolve towards the common end of the lines, which is the state of stable equilibrium for the system. As can be seen in the figure, when the temperature difference between a and b is small, the maximum of the concentration difference through the evolution approaches zero very quickly. Since the lowest order terms of Eqs. (110) and (111) have different signs, the nonlinear effects of temperature drive the probability (mass) flux towards the higher temperature subsystem. This phenomena can be explained from an entropy generation standpoint. The higher temperature subsystem has a higher specific entropy so that the probability (mass) flux towards it results in entropy generation for the system. On the other hand, the temperature evolution is explained by the fact that the heat diffusion towards the lower temperature subsystem increases the specific entropy in the lower temperature subsystem, which in turn results in entropy generation for the system.

When probability (mass) and temperature differences exist at the same time, it is the combined effect (i.e., the coupling) from the probability and the temperature, which determines the probability flow, since the lower order terms of Eqs. (91) and (110) have opposite signs.






FIG. 3. (color online) Thermodynamic trajectories on a temperature-particle number diagram for the three cases. Case 1:  $p^a = 0.8$ ,  $p^b = 0.2$ ,  $T^a = 1500$  K,  $T^b = 300$  K (dashed-dotted line); Case 2:  $p^a = 0.8$ ,  $p^b = 0.2$ ,  $T^a = 900$  K,  $T^b = 900$  K (solid line); and Case 3:  $p^a = 0.2$ ,  $p^b = 0.8$ ,  $T^a = 1500$  K,  $T^b = 300$  K (dashed line). The black lines are the trajectory of subsystem  $a$ , and the green (gray) lines are the trajectory of subsystem  $b$ .




FIG. 4. (color online) Entropy evolution and entropy generation rate in dimensionless time for the three cases of Fig. 3. Case 1:  $p^a = 0.8$ ,  $p^b = 0.2$ ,  $T^a = 1500$  K,  $T^b = 300$  K (dashed-dotted line); Case 2:  $p^a = 0.8$ ,  $p^b = 0.2$ ,  $T^a = 900$  K,  $T^b = 900$  K (solid line); and Case 3:  $p^a = 0.2$ ,  $p^b = 0.8$ ,  $T^a = 1500$  K,  $T^b = 300$  K (dashed line). The black lines are the entropy evolutions relative to the vertical axis on the left, and the green (gray) lines are the entropy generation rates relative to the vertical axis on the right.

In Fig. 3, the trajectory for Case 1 (initially:  $p^a = 0.8$ ,  $p^b = 0.2$ ,  $T^a = 1500$  K,  $T^b = 300$  K) shows a competition effect between the probability and temperature, while the trajectory of Case 3 (initially:  $p^a = 0.2$ ,  $p^b = 0.8$ ,  $T^a = 1500$  K,  $T^b = 300$  K) shows a cooperation effect. Both of these can be explained via the effects discussed relative

to Fig. 2. In addition, as a validation, Fig. 4 provides the entropy generation rate for the three cases along with the entropy trajectories. All exhibit monotonic increases in the entropy over time. With respect to the entropy generation, Case 1 for which the temperature effect is competitive results in a greater variation in the entropy generation rate than that for the diffusion case without temperature difference (Case 2), while the case for which the temperature effect is cooperative (Case 3) results in a much steeper drop in the entropy generation rate than either of the two other cases.

In order to place the entropy evolution within an actual time scale for the three cases studied, the absolute concentration (or equivalently, the equilibrium pressure) needs to be specified. In particular, assuming an ideal gas and knowing the equilibrium temperature for the three cases (Case 1: 1260 K, Case 2: 900 K, Case 3: 540 K; see Fig. 3), we set the corresponding equilibrium pressures to be 0.014 atm (Case 1), 0.01 atm (Case 2), 0.006 atm (Case 3) in order to arrive at the same particle concentration of  $0.1355$  mole  $m^{-3}$  for all three cases. Using Eq. (100) and (101), the relaxation time for the three cases can be calculated. In Eq. (101), the distance of two subsystems is chosen to be  $10^{-5}$  m, which is on the same order of magnitude as the mean-free-path ( $3.4 \sim 7.8 \times 10^{-5}$  m), so that the system studied exhibits nonequilibrium effects. The mean-free-path is calculated from  $l = \frac{k_B T}{\sqrt{2} \pi d^2 p}$  where  $d$  is the diameter of the gas particles which is set equal to twice the Van der Waals radius of hydrogen ( $2 \times 1.2$  Å). In addition, we verify that the distance of two subsystems is much larger than the de Broglie wave length ( $\sim 10^{-10}$  m) given by Eq. (83).

Using the relaxation times (Case 1:  $4.41 \times 10^{-10}$  s; Case 2  $5.52 \times 10^{-10}$  s; Case 3:  $7.79 \times 10^{-10}$  s) calculated based on the self-diffusion coefficient, Fig. 5 shows the entropy evolution in real time. The three cases have the same concentrations but different equilibrium temperatures so that the scaling of the three curves using the relaxation times is different. However, the curves retain the same features as in Fig. 4. We observe that even though the relaxation time for Case 1 is smaller than that for Case 3 due to its higher equilibrium temperature and that the total entropy generated per particle is smaller as well (i.e.,  $7.76 \times 10^{-23}$   $JK^{-1}particle^{-1}$  for Case 1 as opposed to  $11.86 \times 10^{-23}$   $JK^{-1}particle^{-1}$  for Case 3), Case 1 still takes about the same amount of time as Case 3 to arrive at stable equilibrium. The reason is that the initial entropy generation rates for Case 1 are much smaller than those for Case 3 due to the competition effect of heat and mass diffusion present in the former and the cooperation effect present in the latter.

## VI. CONCLUSION

This paper investigates the relaxation process of local, isolated systems in nonequilibrium using the SEAQT




FIG. 5. (color online) Entropy evolution and entropy generation rate in dimensionless time for the three cases of Fig. 3. Case 1:  $p^a = 0.8$ ,  $p^b = 0.2$ ,  $T^a = 1500$  K,  $T^b = 300$  K (dashed-dotted line); Case 2:  $p^a = 0.8$ ,  $p^b = 0.2$ ,  $T^a = 900$  K,  $T^b = 900$  K (solid line); and Case 3:  $p^a = 0.2$ ,  $p^b = 0.8$ ,  $T^a = 1500$  K,  $T^b = 300$  K (dashed line). The black lines are the entropy evolutions relative to the vertical axis on the left, and the green (gray) lines are the entropy generation rates relative to the vertical axis on the right.

framework. The mass and heat diffusion inside the system, which are the mass and energy redistributions among the system subspaces (or equivalently, among the energy eigenlevels), are described by defining conjugate forces and conjugate fluxes using the concepts of hypoequilibrium state and nonequilibrium intensive properties. These thermodynamic features or patterns of the nonequilibrium relaxation process are used to generalize the Gibbs relation, the Clausius inequality, the Onsager relations, and the quadratic dissipation potential to the far-from-equilibrium realm and for non-quasi-equilibrium processes. In addition, the variational principle in the spaces spanned by conjugate forces and conjugate fluxes is derived from the variational principle in system state space (i.e., from the principle of steepest entropy ascent). As an application, the mass diffusion of a simple system consisting of hydrogen is studied. The study results in decoupled mass and energy transport equations and their associated phenomenological coefficients in the near-equilibrium realm. In far-from-equilibrium realm, the coupling phenomena and the nonlinear effects for mass and energy transport are derived.

From this investigation it is evident that the introduction of the concepts of hypoequilibrium state and nonequilibrium intensive properties into the SEAQT framework provides a novel and fundamental vantage point from which to describe nonequilibrium states and their evolution during a relaxation process. In addition, use of the density of states method developed by the authors in [16] has permitted the wide application of

the SEAQT framework to the study of nonequilibrium systems in which complex, coupled reaction diffusion pathways are modeled and compared with experiment [28, 29]. chemical reaction, mass and heat diffusion. As a complement to the present paper, [24] continues our study of the non-quasi-equilibrium process of two interacting systems and completes our discussion of Onsager type of investigations of the relaxation process with fluxes both inside a nonequilibrium system and those across different systems. All of these studies show SEAQT to be a powerful approach applicable to the study of nonequilibrium phenomena across all temporal and spatial scales.

## ACKNOWLEDGMENT

Funding for this research was provided by the US Office of Naval Research under Award No. N00014-11-1-0266.

## Appendix A: Relaxation time $\tau$

A brief supplementary explanation is given here to describe the range over which the SEA framework can be applied and how the relaxation time  $\tau$  can be chosen. For more detailed discussions on the dynamics of nonequilibrium, especially on how  $\tau$  is chosen using the mechanics and how  $\tau$  is related to the state space geometry, the reader is referred to [18, 23].

Given a well-defined system with a group of constraints (i.e. a ‘level of description’ [13, 30] of system state), Eq. (18) exists to explain the system’s entropy generation, regardless of the spatial and temporal range of system behavior. Beretta has set up several models using the principle of SEA for different spatial and temporal scales by defining the proper state space and constraints [18]. In addition to the ‘level of description’ of system state, the metric and the relaxation time define what are the elementary phenomena [30] involved in the relaxation process, i.e., what the ‘level of phenomena’ is. By elementary phenomena, we mean that the phenomena need not be explained by using a more microscopic process in the model. As an example, when the mass diffusion process in a continuum model is viewed as elementary, it can be described by a diffusion coefficient, even though in a more microscopic framework, such as molecular dynamics or kinetic theory, diffusion can be described or explained by a balance between particle collisions and particle free-flight. Practically, given a model set-up of state space, constraints, and metric, the relaxation time  $\tau$  can be chosen using the experimental data of the given ‘level of phenomena’ (e.g., the diffusion coefficient measurement in Sec. V.D) or calculated from a more microscopic ‘level of phenomena’ (e.g., the Chapman-Enskog theory of Eq. (101)) to recover the dynamics of interest.

At the atomistic level where the ‘level of description’ is at a quantum level, the spin coherence in a study of entropy generation in, for example, a quantum spin system is the ‘level of phenomena’. In this case, the relaxation time can be chosen according to the characteristic time of a quantum-level experiment as is done in [31].

At a mesoscopic level, the ‘level of description’ is the local distribution function; the ‘level of phenomena’ is the local redistribution, which results from the more microscopic collisions or scattering of individual particles; and the relaxation time can be calculated from the collision integral. A comparison of the SEAQT framework with kinetic theory for a local system (i.e., for an infinitesimally small system to which SEAQT is applied) allows the SEA framework to use the relaxation time of kinetic theory. The equation of motion of kinetic theory takes the general form

$$\frac{\partial f}{\partial t} - \{f, H\} = \left( \frac{\partial f}{\partial t} \right)_{col} \quad (\text{A1})$$

where the second term on the left hand side is the Poisson bracket term, which accounts for the reversible part ( $X_\gamma^H$ ) in Eq. (1), while the collision term on the right hand side accounts for the irreversible part ( $Y_\gamma^H$ ) of Eq. (1). In the near-equilibrium region, the BGK approximation with relaxation time  $\tau$  takes the form

$$\frac{\partial f}{\partial t} - \{f, H\} = -\frac{1}{\tau}(f - f^{eq}) \quad (\text{A2})$$

As a comparison, the SEAQT equation of motion can be rewritten in the following form using Hilbert space for state space and the Schrödinger equation for the reversible part:

$$\frac{d\hat{\rho}}{dt} - \frac{1}{i\hbar}[\hat{\rho}, \hat{H}] = \frac{1}{\tau(\hat{\rho})}\hat{D}(\hat{\rho}) \quad (\text{A3})$$

In this equation, the irreversible part on the right hand side is developed thermodynamically and reduces to the right hand side of Eq. (A4) in the near-equilibrium region, i.e.,

$$\frac{d\hat{\rho}}{dt} - \frac{1}{i\hbar}[\hat{\rho}, \hat{H}] = -\frac{1}{\tau(\hat{\rho})}(\hat{\rho} - \hat{\rho}^{eq}) \quad (\text{A4})$$

where  $\hat{\rho}^{eq}$  is diagonal in the energy eigenstate basis and Maxwellian. Thus, in the classical limit, the density operator reduces to the phase space distribution function and the commutator to the Poisson bracket, recovering the BGK equation. Via a comparison of Eq. (A2) and Eq. (A4), the relaxation time in kinetic theory, which is derived *ab initio* from quantum mechanics or classical mechanics (i.e., from the collision integral), is equivalent in the near equilibrium realm to the relaxation time of entropy generation in the SEAQT framework. Thus, practically, when a SEAQT model is set up at a mesoscopic level for a local system, the relaxation time in SEAQT is that of the BGK equation of kinetic

theory. Even in the case of a non-uniform metric, this correspondence exists [23]. As to the ‘level of phenomena’, the choice of  $\tau$  using kinetic theory implies a local relaxation, resulting from more microscopic particle collisions, and is, thus, the elementary phenomenon for this ‘level of description’.

In a macroscopic system at a continuum level where the elementary phenomenon is mass diffusion or heat diffusion (i.e., is the ‘level of phenomena’), SEAQT still sets the ‘level of description’ to be the distribution function among energy eigenlevels. The authors have proven that the SEAQT equation for mass diffusion and heat diffusion can recover the linear transport equation (Sec. V.C), and the simplest way to determine  $\tau$  is to use experimental data of the phenomenological coefficient (Sec. V.D). Moreover, this macroscopic  $\tau$  can also be acquired from the diffusion coefficient calculated from the moment integral of the BGK equation (i.e., the Chapman-Enskog theory) used to determine the mesoscopic  $\tau$  in the previous paragraph and is in this case the result of both local relaxation and convective transport.

## Appendix B: Hypoequilibrium for heat diffusion

In this appendix, it is proven that for a system with heat diffusion only, if the initial state is given by Eq. (23), the system evolution is also given by Eq. (24), and the nonequilibrium temperature is well-defined. The proof follows the same process as in [16] for a system with probability and energy conservations. To show this, Eq. (62) is reformulated such that

$$\frac{d}{dt} \ln \frac{p_j^a}{n_j^a} = \frac{1}{\tau} \left( -\ln \frac{p_j^a}{n_j^a} - \frac{B_2^a}{B_1} + \epsilon_j^a \frac{B_3}{B_1} \right) \quad (\text{B1})$$

where it is noted that  $d(\ln n_j^a)/dt$  is zero and that  $B_1$ ,  $B_2^a$ , and  $B_3$  are the same for all chosen energy eigenlevels  $p_j^a$  from subspace  $a$  and only a function of the entire probability distribution at a given instant of time. Subtracting the equations of motion for the  $i$ th and  $k$ th energy eigenlevels results in

$$\frac{d}{dt} \left( \ln \frac{p_j^a}{n_j^a} - \ln \frac{p_k^a}{n_k^a} \right) = -\frac{1}{\tau} \left( \ln \frac{p_j^a}{n_j^a} - \ln \frac{p_k^a}{n_k^a} \right) + \frac{1}{\tau} \frac{B_3}{B_1} (\epsilon_j^a - \epsilon_k^a) \quad (\text{B2})$$

Defining a new variable

$$W_{jk} = \frac{1}{\epsilon_j^a - \epsilon_k^a} \left( \ln \frac{p_j^a}{n_j^a} - \ln \frac{p_k^a}{n_k^a} \right) \quad (\text{B3})$$

the time evolution of  $W_{jk}$  yields to the ordinary differential equation

$$\frac{dx}{dt} = -\frac{1}{\tau}x + \frac{1}{\tau} \frac{B_3}{B_1} \quad (\text{B4})$$

If  $p_j^a$  and  $p_k^a$  are in the same subsystem for which the initial probability distribution is a canonical one, i.e., if

$$p_j^a(t=0) = \alpha^a n_j^a e^{-\epsilon_j^a \beta^a}, \quad p_k^a(t=0) = \alpha^a n_k^a e^{-\epsilon_k^a \beta^a} \quad (\text{B5})$$

then

$$W_{jk}(t=0) = \frac{1}{\epsilon_j^a - \epsilon_k^a} \left( \ln \frac{p_j^a}{n_j^a} - \ln \frac{p_k^a}{n_k^a} \right) = -\beta^a \quad (\text{B6})$$

For  $\forall p_j^a, p_k^a$  in the same subsystem  $a$ , the time evolution of  $W_{jk}$  yields to the same ordinary differential equation (ODE) with the same initial value, namely,

$$\frac{dx}{dt} = -\frac{1}{\tau}x + \frac{1}{\tau} \frac{B_3}{B_1}, \quad x = W_{jk}(t=0) = -\beta^a \quad (\text{B7})$$

so that the solution of  $W_{jk}$  is the same  $W_{jk}(t) = \beta^a(t)$ . Therefore, the probability distribution in this subsystem maintains the canonical distribution with the parameter  $\beta^a(t)$  given by

$$p_j^a(t) = \alpha^a(t) n_j^a e^{-\epsilon_j^a \beta^a(t)} \quad (\text{B8})$$

In addition, the temperature of the subsystem at time  $t$  is defined by

$$T^a(t) = \frac{1}{k_b \beta^a(t)} \quad (\text{B9})$$

Thus, for a system in a nonequilibrium state, the hypoequilibrium temperature for each subsystem is defined. This temperature can be the same or different from that of any other subsystem. If a system is in a

$M$ th-order hypoequilibrium state, it remains at least of order  $M$  throughout as well as after the evolution, and the probability distribution of each subsystem remains canonical.

### Appendix C: Evolution of intensive properties

In this appendix, the evolutions of intensive properties are given for the system with probability and energy conservations. Eq. (26) is reformulated such that

$$\frac{d}{dt} \ln \frac{p_j^K}{n_j^K} = \frac{1}{\tau} \left( -\ln \frac{p_j^K}{n_j^K} - \frac{A_2}{A_1} + \epsilon_j^a \frac{A_3}{A_1} \right) \quad (\text{C1})$$

Using Eqs. (33) and (34) yields

$$\frac{d}{dt} (-\alpha^K(t) - \beta^K(t) \epsilon_i^K) = \frac{1}{\tau} (\alpha^K(t) + \beta^K(t) \epsilon_i^K - \alpha - \epsilon_j^K \beta) \quad (\text{C2})$$

Subtracting the equations of motion for the  $i$ th and  $j$ th energy eigenlevels results in

$$\begin{aligned} \frac{d}{dt} (-\beta^K(t) \epsilon_i^K + \beta^K(t) \epsilon_j^K) &= \frac{1}{\tau} (\beta^K(t) \epsilon_i^K - \beta^K(t) \epsilon_j^K) \\ &\quad - \frac{1}{\tau} \beta (\epsilon_i^K - \epsilon_j^K) \end{aligned} \quad (\text{C3})$$

If  $i \neq j$ , dividing both sides by  $(\epsilon_i^K - \epsilon_j^K)$  results in the evolution for  $\beta^K$ , namely,

$$\frac{d\beta^K}{dt} = -\frac{1}{\tau} (\beta^K - \beta) \quad (\text{C4})$$

Finally, subtracting Eq. (C4) from (C2) gives the evolution for  $\alpha^K$ , i.e.,

$$\frac{d\alpha^K}{dt} = -\frac{1}{\tau} (\alpha^K - \alpha) \quad (\text{C5})$$

- 
- [1] D. C. Rapaport, *The art of molecular dynamics simulation* (Cambridge university press, 2004).
- [2] S. Chen and G. D. Doolen, *Annual Review of Fluid Mechanics* **30**, 329 (1998).
- [3] P. Vogl and T. Kubis, *Journal of Computational Electronics* **9**, 237 (2010).
- [4] A. Rossani and G. Kaniadakis, *Physica A: Statistical Mechanics and its Applications* **277**, 3492 (2000).
- [5] A. L. Garcia and W. Wagner, *Phys. Rev. E* **68**, 056703 (2003).
- [6] J. M. Torres-Rincon, "Boltzmann-Uehling-Uhlenbeck Equation," in *Hadronic Transport Coefficients from Effective Field Theories* (Springer, 2014) pp. 33–45.
- [7] M. E. J. Newman and G. T. Barkema, *Monte Carlo Methods in Statistical Physics* (Oxford University Press: New York, USA, 1999).
- [8] S. R. de Groot and P. Mazur, *Non-equilibrium thermodynamics* (North-Holland Publication Co., Amsterdam, 1962).
- [9] I. Gyarmati, E. Gyarmati, and W. F. Heinz, *Non-equilibrium thermodynamics* (Springer, 1970).
- [10] R. Kubo, M. Toda, and N. Hashitsume, *Statistical physics II: nonequilibrium statistical mechanics*, Vol. 31 (Springer Science & Business Media, 2012).
- [11] U. Seifert, *Reports on Progress in Physics* **75**, 126001 (2012).
- [12] J. Casas-Vázquez, and G. Lebon, *Extended irreversible thermodynamics* (Springer, 1996).
- [13] M. Grmela and H. C. Öttinger, *Phys. Rev. E* **56**, 6620 (1997).
- [14] H. C. Öttinger and M. Grmela, *Phys. Rev. E* **56**, 6633 (1997).
- [15] E. Zanchini and G. P. Beretta, *Entropy* **16**, 1547 (2014).
- [16] G. Li and M. R. von Spakovsky, *Phys. Rev. E* **93**, 012137 (2016).
- [17] M. Grmela, *Entropy* **16**, 1652 (2014).
- [18] G. P. Beretta, *Phys. Rev. E* **90**, 042113 (2014).

- [19] A. Montefusco, F. Consonni, and G. P. Beretta, *Phys. Rev. E* **91**, 042138 (2015).
- [20] G. P. Beretta, *Phys. Rev. E* **73**, 026113 (2006).
- [21] E. P. Gyftopoulos and E. Çubukçu, *Phys. Rev. E* **55**, 3851 (1997).
- [22] G. P. Beretta, *Reports on Mathematical Physics* **64**, 139 (2009).
- [23] G. Li and M. R. von Spakovsky, “Steepest-entropy-ascent thermodynamics under non-uniform state space geometry,” (unpublished).
- [24] G. Li and M. R. von Spakovsky, “Steepest-entropy-ascent quantum thermodynamic modeling of the far-from-equilibrium interactions between non-equilibrium systems of indistinguishable particle ensembles,” (2016), [arXiv:1601.02703](https://arxiv.org/abs/1601.02703).
- [25] M. Grmela, V. Klika, and M. Pavelka, *Phys. Rev. E* **92**, 032111 (2015).
- [26] E. P. Gyftopoulos and G. P. Beretta, *Thermodynamics: foundations and applications* (Courier Corporation, 2005).
- [27] R. B. Bird, W. E. Stewart, and E. N. Lightfoot, (2002).
- [28] G. Li and M. R. von Spakovsky, in *ASME 2015 International Mechanical Engineering Congress and Exposition* (American Society of Mechanical Engineers, 2015) no. IMECE2015-53726.
- [29] G. Li, M. R. von Spakovsky, F. Shen, and K. Lu, “Multiscale transient and steady state study of the influence of microstructure degradation and chromium oxide poisoning on SOFC cathode performance,” (unpublished).
- [30] G. P. Beretta, J. C. Keck, M. Janbozorgi, and H. Metghalchi, *Entropy* **14**, 92 (2012).
- [31] S. Cano-Andrade, G. P. Beretta, and M. R. von Spakovsky, *Phys. Rev. A* **91**, 013848 (2015).