

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Mixing and the fractal geometry of piecewise isometries

Paul P. Park, Thomas F. Lynn, Paul B. Umbanhowar, Julio M. Ottino, and Richard M. Lueptow

Phys. Rev. E **95**, 042208 — Published 14 April 2017

DOI: [10.1103/PhysRevE.95.042208](https://doi.org/10.1103/PhysRevE.95.042208)

# Mixing and the fractal geometry of piecewise isometries

Paul P. Park and Thomas F. Lynn

*Department of Engineering Sciences and Applied Mathematics,  
Northwestern University, Evanston, Illinois 60208, USA*

Paul B. Umbanhowar

*Department of Mechanical Engineering,  
Northwestern University, Evanston, Illinois 60208, USA*

Julio M. Ottino

*Department of Chemical and Biological Engineering,  
Northwestern University, Evanston, Illinois 60208, USA  
Department of Mechanical Engineering,  
Northwestern University, Evanston, Illinois 60208, USA and  
The Northwestern Institute on Complex Systems (NICO),  
Northwestern University, Evanston, Illinois 60208, USA*

Richard M. Lueptow\*

*Department of Mechanical Engineering,  
Northwestern University, Evanston, Illinois 60208, USA  
Department of Chemical and Biological Engineering,  
Northwestern University, Evanston, Illinois 60208, USA and  
The Northwestern Institute on Complex Systems (NICO),  
Northwestern University, Evanston, Illinois 60208, USA*

(Dated: March 13, 2017)

## Abstract

Mathematical concepts often have applicability in areas that may have surprised their original developers. This is the case with Piecewise Isometries (PWIs) which transform an object by cutting it into pieces that are then rearranged to reconstruct the original object, and which also provide a paradigm to study mixing via cutting and shuffling in physical sciences and engineering. Every PWI is characterized by a geometric structure called the exceptional set,  $E$ , whose complement comprises non-mixing regions in the domain. Varying the parameters that define the PWI changes both the structure of  $E$  as well as the degree of mixing the PWI produces, which begs the question of how to determine which parameters produce the best mixing. Motivated by mixing of yield stress materials, for example granular media, in physical systems, we use numerical simulations of PWIs on a hemispherical shell and examine how the fat fractal properties of  $E$  relate to the degree of mixing for any particular PWI. We present numerical evidence that the fractional coverage of  $E$  negatively correlates with the intensity of segregation, a standard measure for the degree of mixing, which suggests that fundamental properties of  $E$  such as fractional coverage can be used to predict the effectiveness of a particular PWI as a mixing mechanism.

---

\* r-lueptow@northwestern.edu

## I. INTRODUCTION

Mixing by cutting and shuffling, described mathematically as Piecewise Isometries (PWIs), is far less understood than other forms of mixing such as mixing generated by stretching and folding, which is directly applicable to fluids and can be cast in the language of nonlinear dynamics and chaos theory. Nevertheless, mixing with PWIs has potential for application in engineering processes involving granular materials [1–3], yield stress fluids, valved fluid flows [4, 5], or flows with shear-banding materials, as well as for understanding certain notable geophysical phenomena (e.g., imbricate thrust faults [6]). PWIs follow simple rules that generate complex dynamics [7–20]. We focus here on measuring intrinsic features of PWIs and the mixed state of the system in order to establish a relationship between the theoretical properties of PWIs and mixing in practice.

There are various similar definitions of a PWI [22–25] – we define a PWI mapping  $M$  on a domain or space  $S$  as:

**Definition 1.**  $M : S \rightarrow S$  such that the action of  $M$  on each partition element  $P_i$  of  $S$  is a Euclidean isometry (composition of translation, rotation, reflection, *etc.*).  $P_i$ , a closed region (including boundaries), is one of a finite number  $N$  partition elements of  $S$ , i.e.  $\cup_{1 \leq i \leq N} P_i = S$ , and where  $P_i \cap P_j$  is measure-zero for  $i \neq j$ .




FIG. 1. Bottom view of a hemispherical shell (HS) showing (a) the initial location of the pieces to be rearranged,  $P_i$ , and their corresponding boundaries comprising  $\mathcal{D}$  (black and red arcs) and the domain boundary,  $\partial S$  (blue arc) and (b) their subsequent locations after their rearrangement by one iteration of the PWI. Reproduced from Park *et al.* [21] with permission from AIP Publishing.






FIG. 2. PWI operation on the HS for  $(\theta_z, \theta_x) = (45^\circ, 45^\circ)$ . Dashed lines indicate the two rotation axes. (a) Initial condition. (b) Rotation and cut about the  $z$ -axis by  $\theta_z$ . (c) Rotation and cut about the  $x$ -axis by  $\theta_x$ . (d) New state after one iteration. Adapted from Park *et al.* [21] with permission from AIP Publishing.

A lower hemispherical shell (HS) undergoing a PWI transformation is split into four partition elements (bottom view, Figure 1(a)), which are then rearranged into another HS (Figure 1(b)). We let  $\mathcal{D}$  denote the set of all such borders that partition  $S$  (black and red arcs in Figure 1(a)):

**Definition 2.**  $\mathcal{D} = \cup_{i,j} (P_i \cap P_j)$  for  $i \neq j$ .

Note that since each segment of  $\mathcal{D}$  is in the shared domain of two isometries, the map  $M$  is multi-valued on  $\mathcal{D}$ .

In this study we focus on a particular class of PWIs that result from successive rotations about the  $z$  and  $x$ -axes by  $\theta_z$  and  $\theta_x$ , with periodic boundaries at the hemispherical edge (Figure 2). The operation is expressed mathematically by Sturman *et al.* [17], and is applied here for computer simulations of PWIs. This class of PWIs is specified by the ordered pair  $(\theta_z, \theta_x)$ , and approximates the effect of an infinitely thin flowing layer in a spherical granular tumbler [2, 3, 17, 26], which can be reduced to 2D dynamics on the HS. The ordered pair  $(\theta_z, \theta_x)$  will be referred to as a protocol for the remainder of this paper, and it determines the location of  $\mathcal{D}$  (for instance, the set  $\mathcal{D}$  in Figure 1(a) derives from the  $(45^\circ, 45^\circ)$  protocol).

We can observe mixing on the HS when a particular PWI is repeated, with the rate and degree of mixing depending on the protocol. This is demonstrated in Figure 3(b), where the three protocols  $(54^\circ, 54^\circ)$ ,  $(75^\circ, 75^\circ)$  and  $(30^\circ, 15^\circ)$  result in vastly different degrees of mixing after  $m = 150$  iterations for the HS initially colored half red and half light blue in Figure 3(a) (this and all remaining figures with a HS show the bottom view of the HS).




FIG. 3. Bottom view of (a) initial condition of seeded colors on the HS; (b) mixing on the HS for the  $(54^\circ, 54^\circ)$ ,  $(75^\circ, 75^\circ)$  and  $(30^\circ, 15^\circ)$  protocols after  $m = 150$  iterations; and (c)  $E$  for corresponding protocols obtained by tracking the respective  $\mathcal{D}$  for 5000 iterations.

When considering mixing under PWIs, non-mixing regions can be identified by visualizing the exceptional set [24]:

**Definition 3.**  $E = \bigcup_{-\infty < i < \infty} M^i \mathcal{D}$ .

The dynamics on the closure of  $E$  has been shown to be ergodic in other systems [27–30], and we conjecture that this is also the case here, which is why for ease of illustration, we approximate  $E$  with its subset  $E_+$  [21]:

**Definition 4.**  $E_+ = \bigcup_{1 \leq i < \infty} M^i \mathcal{D}$ .

All images representing  $E$  hereafter are approximated with  $E_+$ . We generate  $E_+$  by seeding tracer points on an equivalent substitute of  $\mathcal{D}$  (refer to Park *et al.* [21] for further details) and recording their trajectories for a large number of iterations (Figure 3(c)). The white areas in the HS are non-mixing regions known as cells [31, 32] (we refer to the region as a cell rather than the set of points occupying the region as in Park *et al.* [21]), which form a subset of the complement of the closure of  $E$ . In Figure 3(c) it is evident that the cells

coincide with non-mixing regions in Figure 3(b) for the  $(54^\circ, 54^\circ)$  and  $(75^\circ, 75^\circ)$  protocols. For example, the four large circular cells in Figure 3(c) for the  $(75^\circ, 75^\circ)$  protocol correspond to large light blue and red circular regions in Figure 3(b). For the  $(54^\circ, 54^\circ)$  protocol, there are two adjacent circular regions in Figure 3(b) located near the center and just below that contain both red and light blue colors with a clear divide between the two. This is because the initial seeding of colors overlaps with two of the cells shown in Figure 3(c) and both colors happened to be seeded in each of the cells.

Colors contained inside a cell are said to have a *periodic itinerary* [31, 32] because they travel as a group from one cell to others of the same size before returning to the cell where they originated, but with a rotation. The boundaries between the two colors in the cells for the  $(54^\circ, 54^\circ)$  protocol are slanted rather than in the vertical orientation of the initial condition due to this rotation. The mixing patterns of the  $(54^\circ, 54^\circ)$  and  $(75^\circ, 75^\circ)$  protocols demonstrate that cells are non-mixing because no colors penetrate them and no mixing occurs within them.

While a greater coverage of  $E$  seems to produce qualitatively better mixing in each example in Figure 3, our goal is to show that such a relationship exists more generally based on quantifying the fractional coverage of  $E$  and comparing the coverage to standard measures for the degree of mixing. If greater coverage of  $E$  produces better mixing, this would imply that measures of fundamental properties of  $E$  exist which can be used to judge the general effectiveness of a particular PWI as a mixing mechanism. Quantifying the fractional coverage of  $E$  and the degree of mixing each has its respective complications and factors that need to be considered, which will be mentioned here and addressed more thoroughly in the following sections.

First, we observe that the approximated  $E$  has two distinct and related features: Fine details emerge when  $E$  is viewed at magnified scales (Figure 4), and the apparent coverage depends on the resolution at which  $E$  is measured (described succinctly by Umberger and Farmer as “structure on all scales” and “dependence of the apparent size on the scale of resolution”) [33], all of which are features that characterize fractals. For most of the cases considered here,  $E$  appears to have positive area, which suggests that they are *fat fractals*, as opposed to traditional fractals which are measure-zero, i.e. they have zero area in two dimensions [33].

Note, however, that although  $E$  in Figure 3(c) appears to fill in some regions while




FIG. 4. Magnifying  $E$  for the  $(54^\circ, 54^\circ)$  protocol by  $8\times$  to various scales. Each magnification, from (a) to (b), from (b) to (c), and from (c) to (d) of the red box reveals circular cells at progressively finer scales.

leaving others empty,  $E$  is made up of arcs having length but no thickness (it is a countable union of measure-zero sets). Thus, even regions that are dark blue in Figure 3(c) consist of densely packed curves with zero thickness that are approximated by densely packed points in practice. That  $E$  exhibits fractal characteristics allows us to quantify the area that  $E$  appears to cover using a box counting method [34], but by doing so, we are in effect, “fattening”  $E$  and measuring its closure  $\bar{E}$  [35, 36]. For the remainder of this paper, we use the term “fractional coverage of  $E$ ,” with the understanding that the term refers to the measure of  $\bar{E}$ . Aside from our conjecture that  $E$  is a fat fractal, which gives us a possible method for measuring its coverage, it is significant that the amount of mixing with PWIs could be connected to the fractal characteristics displayed by  $E$  for a particular protocol, given the close relationship between fractals and mixing of fluids [37–46].

Considering a range of protocol values, Figure 5 shows that a variety of intricate patterns

are possible for  $E$ , where  $E$  is approximated at  $m = 5000$  for values of  $\theta_z$  and  $\theta_x$  ranging from  $15^\circ$  to  $90^\circ$  in  $15^\circ$  increments. Only protocols with  $\theta_x \leq \theta_z$  are shown because the global structures of  $E$  for the  $(\theta_1, \theta_2)$  and  $(\theta_2, \theta_1)$  protocols are symmetric (see Appendix A). Further exploration of these patterns for smaller angle increments suggests a rich variety of patterns for  $E$  (Figures 17-19 in Appendix B).

The coverage of  $E$  also varies substantially with the choice of protocol. The coverage ranges from apparently completely filled for the  $(45^\circ, 15^\circ)$  protocol to nearly empty for the  $(90^\circ, 90^\circ)$  protocol. The  $(90^\circ, 90^\circ)$  protocol represents a trivial case where the partition of the HS for the next iteration overlaps perfectly with the reassembled HS after an iteration of the protocol (the four cells have a periodic itinerary of 2 but are period-6 because each cell has three sides that the rotation of the cell aligns to). Visible coverage is affected by the size and number of cells. The observed cells are frequently circular, with the exception of the  $(90^\circ, 90^\circ)$  protocol, whose 3-sided cells appear polygonal, which suggests the possibility of other protocols that share similar features. Circles of different sizes nearly fill the HS




FIG. 5. Visualizing  $E$  with  $m = 5000$  for  $\theta_z$  and  $\theta_x \in [15^\circ, 90^\circ]$  varying in  $15^\circ$  increments (bottom view). Reproduced from Park *et al.* [21] with permission from AIP Publishing.

for the  $(90^\circ, 45^\circ)$  protocol, while circles of similar size nearly fill the HS for the  $(90^\circ, 15^\circ)$  protocol. The  $(90^\circ, 75^\circ)$  protocol has four large circles along with many small circles of similar size that form a band on the HS. Compared to those with  $\theta_z = \theta_x$  or  $\theta_z = 90^\circ$ , the other protocols generally have greater coverage and smaller cells.

It is important to note that while PWIs are “non-mixing” according to mathematical definitions [27–30], it is possible to compare the mixed states of protocols using mechanical definitions. Measuring the degree of mixing is complicated by mechanical factors such as the choice of initial condition of the seeded colors, the number of points used in mixing simulations, and, for most measures of mixing such as the *intensity of segregation* [47] that we used to measure mixedness, the resolution of the grid on which the measure is calculated.

In this paper, we quantify the coverage of  $E$  over the HS using the box counting method [33, 35, 36, 48], and show that for most protocols,  $E$  is well described as a fat fractal. We compare these characteristics to the degree of mixing achieved via numerical simulations, quantified by an established engineering measure known as the intensity of segregation [47], to demonstrate a general relationship between the fat fractal characteristics of  $E$  and mixing with PWIs.

## II. APPROXIMATING $E$

The choice of  $m = 5000$  for approximating  $E$  in Figures 5, and 17-19 is arbitrary, and, consequently, must be addressed to assure that we derive reliable measures of its characteristics. This issue is highlighted in Figure 6 where the “filling” by  $\mathcal{D}$  is shown for various protocols after  $m = 1, 10, 100$ , and  $1000$  iterations. For the  $(15^\circ, 15^\circ)$  protocol, the trajectory of  $\mathcal{D}$  cuts the HS into thin slices initially at  $m = 10$ , but at  $m = 100$  and  $m = 1000$ , the orbit is in close proximity to past trajectories and fills the HS slowly while leaving a band of cells on the diagonal. The  $(54^\circ, 54^\circ)$  protocol features several pentagons initially, which fill in to become large circular cells. At  $m = 100$  and  $m = 1000$ , the  $(54^\circ, 54^\circ)$  protocol fills the HS more evenly, with many small cells in addition to large cells. The trajectory of  $\mathcal{D}$  for the  $(45^\circ, 15^\circ)$  protocol cuts the HS into thin slices like the  $(15^\circ, 15^\circ)$  protocol initially, but fills the HS uniformly without any visible cells at  $m = 1000$  iterations.

The issue of choosing a particular value of  $m$  to approximate  $E$  is highlighted using Figure 6 as an example. For the  $(54^\circ, 54^\circ)$  and  $(45^\circ, 15^\circ)$  protocols, the points that are




FIG. 6. Progression of the approximation of  $E$  as a function of the number of protocol iterations  $m$  (bottom view). Note that  $E$  for the  $(15^\circ, 15^\circ)$  protocol at  $m = 1000$  is not fully developed.

followed after  $m = 1000$  fill the HS more densely with more iterations, but the qualitative structure of  $E$  changes little. In contrast, for the  $(15^\circ, 15^\circ)$  protocol, we conclude that the structure of  $E$  is incomplete after  $m = 1000$  compared to the  $m = 5000$  approximation shown in Figure 5 (bottom left corner). At  $m = 5000$ , the structure appears complete based on how densely the points fill the space.

Two points are evident from Figure 6. First, from a visual perspective, the structure that emerges from the orbit of  $\mathcal{D}$  seems to reach an “asymptotic state” for some sufficiently large value of  $m$ . Second, the value of  $m$  at which the general structure of  $E$  is revealed varies with the protocol. For the  $(45^\circ, 15^\circ)$  protocol,  $m = 1000$  might suffice, whereas  $m \geq 5000$  may be necessary for other protocols such as  $(15^\circ, 15^\circ)$ .

If numerous protocols are to be studied, this leads to the question of how to automate the process of determining  $m$  such that enough information is obtained from the orbit of  $\mathcal{D}$  to generate a reasonable approximation of the coverage of  $E$ . We determine whether the coverage of  $E$  is close to an asymptotic state based on the ability of points to visit new regions of the HS. To achieve this, we use the bottom half of an *isocube* as shown in Figure 7 to discretize the HS in a near-uniform manner (each box in the grid is the same size, and its shape is distorted only slightly when mapped on the HS) [49]. We conclude that an asymptotic state has been reached when the points that are seeded to generate  $E$  fail to



visit a new box in the chosen grid for a specified number of *consecutive* iterations. After each iteration, all boxes containing a point are marked as occupied, and if all seeded points fail to visit a new box for a set number of consecutive iterations, then, for the purpose of this study, we conclude that there is no more information to be gained regarding the coverage of  $E$  at that grid resolution.

Before applying this approach, we briefly describe the grid of the isocube half. Figure 7(a) shows an unfolded isocube half. The blue, orange, purple, and yellow colors have two edges, and when “glued” together with their respective colors, the unfolded isocube half forms the bottom half of a cube. The bold black lines on the outside form the outer edge of the assembled half cube, and the inner bold black lines discretize the domain into 12 quarter faces (we refer to them as quarter faces because the bottom face would be divided into 4 squares). Each quarter face is further discretized with a  $2^n \times 2^n$  grid (fine black lines), such that the total number of boxes on the isocube is  $N = 12 \times 2^n \times 2^n$ . The resolution of the grid will be denoted by the exponent  $n$ . This isocube half can then be mapped to the HS




FIG. 7. (a) An unfolded isocube half whose resolution for a quarter of one face is  $2 \times 2$  (top) and  $2^3 \times 2^3$  (bottom). The total number of boxes is  $N = 12 \times 2^n \times 2^n$ , since there are 12 quarter faces (each quarter face is outlined with bold lines); (b) Isocube half in (a) mapped to the HS as viewed from the bottom.



to form an area-preserving grid as shown in Figure 7(b). All boxes have equal area, and the individual box area goes to 0 in the limit of  $N \rightarrow \infty$ . See Wan *et al.* [49] for details of the isocube transformation.

The grid and its resolution are critical to several aspects of the numerical simulations, such as choosing the number of points to seed for  $\mathcal{D}$ , determining whether enough points are available for obtaining a reliable measure for the coverage of  $E$ , and, in Section IV, seeding points for mixing simulations and measuring the degree of mixing. The role of the grid will be described in detail as we discuss each method.

For determining whether a numerical simulation reaches the asymptotic state for the structure  $E$ , the grid resolution is set to a maximum value of  $n = 10$ . At  $n = 10$ , there are approximately  $1.25 \times 10^7$  boxes that can be visited by points on the HS. We approximate  $\mathcal{D}$  as described in Park *et al.* [21], with two sets of points near the edge of the HS – one set of points is seeded before the  $z$ -axis rotation (blue curve in Figures 1(a) and 2(a)), and the other set of points is seeded before the  $x$ -axis rotation (red curve in Figure 2(c)). The points for each set are seeded in the center of boxes at the very edge of the HS (one point per box), which is approximately 8000 points (there are 8 quarter faces on the edge of the isocube half, and each quarter face has  $2^{10} = 1024$  boxes on one edge). Thus, we follow the orbits of 16000 points across a grid of  $1.25 \times 10^7$  boxes. The structure of  $E$  is considered to be close to an asymptotic state when none of the 16000 points visit a new box for 1000 consecutive iterations.

### III. APPROXIMATING THE FRACTIONAL COVERAGE OF $E$ WITH BOX COUNTING

The approximation of  $E$  is closely tied to measuring its fractional coverage – the number of boxes visited by the end of the simulation divided by the total number of boxes is an approximation of the fractional coverage at a given resolution. Also, if the approximation of  $E$  is judged to have reached an asymptotic state on a high resolution grid ( $n = 10$ ), then there must be enough points on the HS that the coverage of  $E$  can be measured for lower resolutions, i.e.  $n < 10$ , which is additional information that can be used for further analysis. Let  $\Phi_n$  denote the fraction of boxes occupied by  $E$  for a grid discretizing the HS with resolution  $n$ . This is illustrated for the  $(90^\circ, 45^\circ)$  protocol in Figure 8, which shows




FIG. 8. Measuring the fraction of coverage of  $E$  for the  $(90^\circ, 45^\circ)$  protocol at resolutions of  $n \in \{3, 4, 5, 6\}$ . Boxes that contain portions of  $E$  are black (bottom view).

how  $\Phi_n$  changes for different values of  $n$ . A finer grid yields a more “accurate” measure, and this naturally leads to the question of what the appropriate resolution should be to measure  $E$ . If possible, taking  $n \rightarrow \infty$  to determine  $\Phi_\infty$  would be ideal.

Considering the fractional coverage  $\Phi_n$  for six sample protocols and  $2 \leq n \leq 10$ , Figure 9 shows that  $\Phi_n$  generally decays toward an asymptotic value. We fit the points in Figure 9 to a curve to approximate a value for  $\Phi_\infty$ . The choice for the curve is based on the observation that  $E$  typically resembles a fat fractal. Thus, the following equation is used to fit  $\Phi_n$  for each protocol [33, 36]:

$$\Phi_n = \Phi_\infty + Ae^{-Bn}, \quad (1)$$

where  $\lim_{n \rightarrow \infty} \Phi_n = \Phi_\infty$  is the asymptotic fractional coverage.

The *exterior dimension* can be calculated for each  $E$  from the coefficient of  $n$  in Equation 1 [33, 35, 48]:

$$d_{\text{ext}} = 2 - \frac{B}{\ln 2}. \quad (2)$$

We calculate the exterior dimension rather than the traditional Hausdorff dimension, which yields a value of 2 for fat fractals and does not provide information about the multi-scale structure observed for  $E$ .

The built-in trust-region-reflective algorithm [50, 51] in MATLAB is used to fit the data to Equation 1 (starting from  $n = 5$ , because low resolutions give poor approximations to  $\Phi_\infty$ ), which gives the fractional coverage  $\Phi_\infty$  and exterior dimension  $d_{\text{ext}}$  for each case (Table I). The quality of the fit is measured by the  $R^2$  value (goodness-of-fit) which ranges from 0




FIG. 9. Fitting the measured values of  $\Phi_n$  for various protocols. Corresponding  $E$  shown on the right.

to 1, where 1 indicates that there is no difference between the data and fitted values. A constraint that  $\Phi_\infty \geq 0$  and  $B \leq \ln 2$  is also applied to the curve fit based on heuristics (we expect  $\Phi_n \geq 0$  for all  $n$  and  $d_{\text{ext}} \geq 1$  for a fat fractal). Given the nature of the fit and these constraints,  $0 \leq \Phi_\infty \leq 1$  and  $1 \leq d_{\text{ext}} \leq 2$ . As anticipated from visual inspection, the fractional coverage of the exceptional sets on the right of Figure 9 decreases monotonically from close to 1 for the  $(30^\circ, 15^\circ)$  protocol (top) to 0 for the  $(90^\circ, 90^\circ)$  protocol (bottom). In contrast, the connection between the values of  $d_{\text{ext}}$  and  $E$  is not obvious. Note that as expected, for the  $(90^\circ, 90^\circ)$  protocol,  $\Phi_\infty$  is approximately zero and  $d_{\text{ext}} \sim 1$  because  $E$  for this protocol is composed of just two arcs.

For the  $(45^\circ, 15^\circ)$  protocol (not shown in Figure 9)  $E$  appears to cover the entire HS (Figures 5 and 6), and  $\Phi_\infty$  is close to 1, but the  $R^2$  value is relatively low (Table I). This occurs because the  $(45^\circ, 15^\circ)$  protocol generates cells that are nearly all smaller than the box size for a grid of  $n \leq 5$ , which means  $\Phi_n = 1$  for  $n < 5$ . Consequently, a scaling behavior is only observed starting from  $n = 5$  in this case, and higher resolutions ( $n > 10$ ) would be necessary for a better fit to Equation 1. In other cases, the data may not scale as Equation 1

TABLE I. Fitted fat fractal coefficients and goodness-of-fit.

Protocol	$\Phi_\infty$	$d_{\text{ext}}$	$R^2$
(45°, 15°)	0.996	1.674	0.868
(30°, 15°)	0.955	1.201	0.999
(75°, 15°)	0.854	1.223	1.000
(75°, 75°)	0.535	1.563	0.998
(89°, 52°)	0.319	1.077	1.000
(54°, 54°)	0.250	1.660	0.993
(90°, 45°)	0.240	1.322	0.999
(90°, 90°)	0.000	1.014	1.000
(72°, 1°)	0.000	1.921	0.872

immediately, as is illustrated by the dashed lines connecting the data points of  $n < 5$  for (75°, 15°), and (54°, 54°) in Figure 9. The values of  $\Phi_n$  are decreasing for the (75°, 15°) and (54°, 54°) protocols, but an inflection point exists at  $n^* = 3$  and  $n^* = 4$  respectively (uncharacteristic of Equation 1), which can lead to lower  $R^2$  values. In this study,  $\theta_z$  was varied from 1° to 90° in 1° increments, and  $\theta_x$  was varied from 1° to  $\theta_z$  in 1° increments for a given  $\theta_z$  (reasons stated in Section I), for a total of 4095 protocols. Approximately 80% of the protocols had  $R^2 \geq 0.9$ . For ones with  $R^2 < 0.9$ , most, if not all, had  $\Phi_\infty \geq 0.95$ , indicating a very high coverage of the HS by  $E$ .

A key question is the relation between the fractional coverage and fractal dimension. Figure 10 shows the fitted values for  $\Phi_\infty$  and  $d_{\text{ext}}$ . More than 90% of the  $\Phi_\infty$  values are greater than 0.6, and 60% of the  $\Phi_\infty$  values are greater than 0.9. The values of  $d_{\text{ext}}$  are spread relatively uniformly, and there is no obvious relationship between  $\Phi_\infty$  and  $d_{\text{ext}}$ .

#### IV. COMPARING $\Phi_\infty$ TO THE DEGREE OF MIXING

In this section,  $\Phi_\infty$  is compared to the degree of mixing to determine whether the relationship observed based on visual comparison in Section I, i.e. that a greater fractional coverage of  $E$  results in better mixing, is also reflected in standard metrics. To simulate mixing on the HS, the left half of the HS is seeded with points associated with species 1




FIG. 10. No obvious correlation between the asymptotic coverage  $\Phi_\infty$  and exterior dimension  $d_{\text{ext}}$  exists for the 4095 protocols examined.

(red), and the right half with species 2 (light blue), as in Figure 3(a). Points were seeded in the center of each box for a grid resolution of  $n = 8$ , which is approximately  $8 \times 10^5$  points. Let  $\bar{c}$  denote the average concentration of species 1. While concentration is calculated based on the frequency of points divided by the total number, it is meant to represent a fraction of an area of the surface. Here, mixing at each iteration  $m$  is measured on a grid of  $N$  boxes using the intensity of segregation  $\mathcal{I}(m)$  [47]:

$$\mathcal{I}(m) = \frac{1}{\bar{c}(1 - \bar{c})} \left( \frac{1}{N} \sum_{i=1}^N (c_i(m) - \bar{c})^2 \right) \in [0, 1], \quad (3)$$

where  $c_i(m)$  is the local concentration of species 1 in box  $i$  of the grid at iteration  $m$ , and  $\mathcal{I}$  represents the normalized variance in the local concentration of species 1 across all  $N$  boxes. If the domain is uniformly mixed for a given resolution, then  $c_i(m) = \bar{c}$  for all  $i$ , and there is no variance, i.e.  $\mathcal{I} = 0$ . If the domain is completely unmixed (segregated), then  $c_i(m) = 1$  or 0 in each bin, and the variance is at its maximum value of  $\bar{c}(1 - \bar{c})$ , so  $\mathcal{I} = 1$ . Note that for cells with distinct discontinuities, e.g., as shown in Figure 3(b) for the  $(54^\circ, 54^\circ)$  protocol,

if the grid resolution is high enough that the boxes are smaller than the cells, boxes that do not contain the discontinuity will only be occupied by one species. This implies that the local concentration will be 0 or 1, and boxes along the discontinuity will contain a mix of species. We use a grid resolution of  $n = 3$ , which corresponds to  $N = 768$  boxes in the grid. Simulations were run for the same set of 4095 protocols described in Section III.

Note that  $\mathcal{I}$  depends on the resolution of the grid on which it is measured and the total number of points seeded on the HS. For a set number of points, if the grid resolution increases, then the measured segregation will increase as well. If the resolution is such that the number of boxes equals the number of points (i.e. given that each point maintains its distance from neighboring points due to the isometry, only one point should occupy a box), then  $\mathcal{I} = 1$  because the concentration of species 1 in each box will be 0 or 1. On the other hand, if the grid resolution decreases, then the measured segregation will decrease until it reaches 0 when the grid consists of one box. In both extremes,  $\mathcal{I}$  is not a meaningful measure of the mixed state of the system.

A grid resolution of  $n = 3$  and  $8 \times 10^5$  points were chosen in consideration of the issues mentioned above while being mindful of computational limitations. There is a sufficient number of points in each box (approximately 1000 points per box) to assure statistically reasonable results, while there is also a sufficient number of boxes to yield a measurement that reflects the mixed state of the entire HS.

For six sample protocols,  $\mathcal{I}$  decreases over 1000 iterations (Figure 11). After approximately 500 iterations,  $\mathcal{I}$  reaches a “steady” state for the  $(89^\circ, 52^\circ)$ ,  $(54^\circ, 54^\circ)$ ,  $(75^\circ, 75^\circ)$ , and  $(30^\circ, 15^\circ)$  protocols, characterized by an asymptotic trend as  $m$  increases. A steady state also appears likely for the  $(75^\circ, 15^\circ)$  protocol for more than 1000 iterations, but it is unclear if there is a steady state for the  $(72^\circ, 1^\circ)$  protocol. Perhaps what is more interesting is the ordering of  $\mathcal{I}$  for different protocols when compared to the asymptotic fractional coverage  $\Phi_\infty$  for  $E$ . With the exception of the  $(72^\circ, 1^\circ)$  and  $(89^\circ, 52^\circ)$  protocols, the apparent coverage of  $E$  is in ascending order from top to bottom. Because materials in cells do not mix, this suggests that the measure  $\Phi_\infty$  for  $E$  could indicate the ultimate degree of mixing, i.e. a protocol with a higher  $\Phi_\infty$  should lead to better mixing and less segregation than a protocol with a lower  $\Phi_\infty$ .

The  $(72^\circ, 1^\circ)$  protocol is one of possibly several exceptions that behaves unlike the other protocols shown in Figure 11. Its zero  $\Phi_\infty$  value (Table I) suggests that mixing is unlikely




FIG. 11. Intensity of segregation  $\mathcal{I}$  vs. iteration number for six sample protocols, corresponding  $E$  are shown on the right.

for this particular protocol, but its  $\mathcal{I}$ , while oscillating, is steadily decreasing, indicating that mixing is occurring, albeit slowly. It would seem to continue if the simulation were run for longer than 1000 iterations. Thus, the measures  $\Phi_\infty$  and  $\mathcal{I}$  are evidently at odds in this case.

We speculate that this discrepancy comes about because the cells of the  $(72^\circ, 1^\circ)$  protocol are miniscule. The cells appear consistent in size, and each cell is approximately 10 times smaller than a single box of the grid. While it is understood that no mixing occurs within a cell, the cells of the  $(72^\circ, 1^\circ)$  protocol are small enough that they can each be considered an individual particle, in contrast to that for protocols with large cells such as the  $(90^\circ, 90^\circ)$  protocol. The mixing that is measured by  $\mathcal{I}$  results from the rearranging of the miniscule cells of different colors in each box.

The  $(89^\circ, 52^\circ)$  protocol appears to be another exception, compared to the  $(54^\circ, 54^\circ)$  protocol, possibly again relating to cell sizes. While its fractional coverage of  $E$  may be greater ( $\Phi_\infty = 0.319$  compared to 0.250 from Table I), the prominent cells of the  $(89^\circ, 52^\circ)$  protocol are much larger and more numerous than that of the  $(54^\circ, 54^\circ)$  protocol, whose remaining cells other than the several large cells happen to be fairly small. The  $(89^\circ, 52^\circ)$  protocol underperforms compared to what its  $\Phi_\infty$  value would suggest because of its particular

distribution of cell sizes.

While  $\mathcal{I}$  does not reach an asymptotic state by  $m = 1000$  for some protocols, our goal is to determine whether most of the protocols tested indeed reach their respective asymptotes. We take a simple approach to examining this by considering the standard deviation of  $\mathcal{I}$ , denoted  $\sigma(\mathcal{I})$ , over a set span of iterations. For a given protocol, if  $\mathcal{I}$  reaches an asymptotic state, the set of preceding iterations should have similar values of  $\mathcal{I}$ , leading to  $\sigma(\mathcal{I}) \sim 0$ . If




FIG. 12. Distribution of  $\sigma(\mathcal{I})$  for all 4095 protocols over a range of  $s = 50$  iterations at different  $m$ . There is some overlap between distributions for different ranges of iterations – the distribution of later iterations are plotted over earlier ones.



this is the case for most protocols, then the distribution of  $\sigma(\mathcal{I})$  will be narrow and close to 0 for later iterations.

The distributions of  $\sigma(\mathcal{I})$  over a span of 50 iterations ( $s = 50$ ) are shown in Figure 12 for the 4095 protocols at various ranges of iterations. Figure 12(a) shows the distribution of  $\sigma(\mathcal{I})$  on a linear scale. For  $m \in [51, 100]$ , the distribution is wide and centered around 0.045. At higher iterations, the distribution of  $\sigma(\mathcal{I})$  shifts closer to 0, indicating that  $\mathcal{I}$  is essentially constant for most protocols by the end of the simulation consistent with an asymptotic state. Figure 12(b) shows the distribution of  $\sigma(\mathcal{I})$  on a log scale, where most values are in the range of  $10^{-4} \leq \sigma(\mathcal{I}) \leq 10^{-2}$  for  $m \in [951, 1000]$ . There is also a small group of protocols whose values of  $\sigma(\mathcal{I})$  are orders of magnitude smaller than that of the rest of the population. The protocols in this group all have  $\theta_z = 90^\circ$ , i.e. the rotation about the  $z$ -axis only flips the colors from one side to the other, producing no mixing. In theory,  $\mathcal{I} = 1$  for all iterations in this case, which is why  $\sigma(\mathcal{I}) \sim 0$  for all protocols with  $\theta_z = 90^\circ$ . Thus, this group of protocols has values of  $\sigma(\mathcal{I})$  orders of magnitude smaller than that for the rest. This highlights the challenge of accounting for the impact of initial conditions in the study of mixing; however, this topic is beyond the scope of this paper.

Distributions of  $\sigma(\mathcal{I})$  for other spans  $s \in \{5, 10, 25, 100\}$  were also calculated. These distributions also became narrower and tended towards zero for increased numbers of iterations, which reinforces the notion that in general,  $\mathcal{I}$  reaches a steady state by  $m = 1000$  iterations. Distributions of the average change in  $\mathcal{I}$ , denoted  $\overline{\Delta\mathcal{I}}$ , for  $s = 50$  at various ranges of iterations were examined to see if  $\mathcal{I}$  trends in any direction. It was found that  $\overline{\Delta\mathcal{I}}$  has a wide distribution with a negative mean initially, since  $\mathcal{I}$  decreases for most protocols when mixing occurs, but at different rates. At higher iterations, the distribution of  $\overline{\Delta\mathcal{I}}$  is narrower and shifts toward 0, indicating that  $\mathcal{I}$  does not trend in a particular direction after its initial decrease. Again, this indicates that  $\mathcal{I}$  at  $m = 1000$  reaches steady state for most protocols and can be used to compare to  $\Phi_\infty$ , which is an intrinsic value of  $E$ .

Returning now to the relationship between the asymptotic fractional coverage  $\Phi_\infty$  of  $E$  and the traditional measure of mixing, the segregation index  $\mathcal{I}$ , Figure 13 shows a scatter plot of  $\mathcal{I}$  versus  $\Phi_\infty$  at different numbers of iterations. Initially, all protocols start from  $\mathcal{I} = 1$ , but after each iteration,  $\mathcal{I}$  decreases as mixing occurs. The points are broadly scattered for smaller  $m$ , but as  $m$  is increased, they appear to collapse toward a diagonal line. The negative linear correlation between  $\mathcal{I}$  and  $\Phi_\infty$  at  $m = 1000$  confirms the hypothesis




FIG. 13. Comparing  $\mathcal{I}$  to  $\Phi_\infty$  for all 4095 protocols at iterations  $m = 25, 10^2$ , and  $10^3$ . At high  $m$ ,  $\mathcal{I}$  and  $\Phi_\infty$  appear to be negatively, linearly correlated.

that the structure of  $E$ , specifically the asymptotic fractional coverage  $\Phi_\infty$ , can be a useful indicator of the degree of mixing. This is significant given the many factors involved with mixing and measuring the degree of mixing (e.g., species number, concentration, and initial distribution as well as the number of tracer points and the grid geometry and resolution for measuring  $\mathcal{I}$ ). Materials in cells do not mix, and this is reflected in mixing simulations as well. Larger  $\Phi_\infty$  corresponds to a smaller overall size of the non-mixing regions (cells). Thus, a negative correlation between  $\mathcal{I}(m = 1000)$  and  $\Phi_\infty$  results. Therefore, fractional coverage  $\Phi_\infty$  of  $E$  can be a useful indicator for judging a particular protocol's effectiveness at mixing.

There are a few exceptions to the negative linear correlation between  $\Phi_\infty$  and  $\mathcal{I}$ , one of which is the protocols with  $\Phi_\infty \sim 1$  that maintain values of  $\mathcal{I} > 0$  even after 1000 iterations along the right edge of Figure 13. In comparing the points at the three different numbers of iterations as in Figure 13, it is clear that the values of  $\mathcal{I}$  are decreasing, which suggests the rate of mixing may be very slow for these protocols despite the high  $\Phi_\infty$  value. While the fractional coverage of  $E$  predicts the long-term mixing achieved, it cannot predict the rate of mixing. Another exception is when the initial condition and protocol are such that no mixing will occur, e.g. when  $\theta_z = 90^\circ$  and one species initially occupies the left half of the

HS and the other species occupies the right half of the HS as was pointed out earlier. This is the set of points at  $\mathcal{I} \sim 1$  along the top edge in Figure 13 that do not collapse independent of  $m$  or  $\Phi_\infty$ . Finally, there is also a group of protocols near the left edge with  $\Phi_\infty \sim 0$ , but with decreasing values of  $\mathcal{I}$ . These are protocols whose behavior may be similar to the  $(72^\circ, 1^\circ)$  protocol, which has  $\Phi_\infty \sim 0$  and a steadily decreasing  $\mathcal{I}$  for increasing  $m$ . These special cases point to the subtleties of studying mixing with PWIs, but also highlight the fact, that as a general rule,  $\Phi_\infty$  can be used as a first-order type approximation for predicting the degree of mixing for PWIs.

## V. CONCLUSION

When mixing with PWIs, the exceptional set,  $E$ , is an inherent structure on the domain that can be used to predict the overall degree of mixing that will be produced. Specifically, the fractional coverage of  $E$ ,  $\Phi_\infty$ , shows a negative linear correlation with the intensity of segregation  $\mathcal{I}$ , a standard measure for mixedness. Because the complement of  $E$  consists of cells (non-mixing regions), the fractional coverage  $\Phi_\infty$  of the fat fractal geometry of  $E$  represents a fundamental measure that can be used to judge the general mixing effectiveness of a given PWI. The methods employed in this study to obtain this result naturally motivates the question of whether  $\Phi_\infty$  can be predicted purely from protocol values. While the majority of the calculated values of the exterior dimension  $d_{\text{ext}}$  are not equal to 1 or 2, confirming that  $E$  on the HS is usually fractal,  $d_{\text{ext}}$  does not appear to be immediately related to mixing in any practical sense, though this requires further study.

Moving forward, it is important to note that selecting protocols with small non-mixing regions based on  $\Phi_\infty$  is necessary but not sufficient for efficient mixing with PWIs. This is the case because  $E$  is a static feature of PWIs, and only represents the possible degree of mixing as  $m \rightarrow \infty$ . It cannot account for all initial conditions, and it does not yield information about the rate of mixing or the rate at which the structure of the dynamics is achieved [52]. Thus, further work is necessary to determine how initial conditions impact mixing and how characteristics of PWIs are related to the rate of mixing, which may require the development of different methods of analysis.

## ACKNOWLEDGMENTS

We gratefully acknowledge Jack E. Tumblin and Lachlan D. Smith for insightful discussions. This work was supported by National Science Foundation grant CMMI-1435065.

---

- [1] R. Sturman, Adv. Appl. Mech. **45**, 51 (2012).
- [2] G. Juarez, R. M. Lueptow, J. M. Ottino, R. Sturman, and S. Wiggins, Europhys. Lett. **91**, 20003 (2010).
- [3] G. Juarez, I. C. Christov, J. M. Ottino, and R. M. Lueptow, Chem. Eng. Sci. **73**, 195 (2012).
- [4] S. W. Jones and H. Aref, Phys. Fluids **31**, 469 (1988).
- [5] L. D. Smith, M. Rudman, D. R. Lester, and G. Metcalfe, Chaos **26**, 023113 (2016).
- [6] S. E. Boyer and D. Elliott, AAPG Bull. **66**, 1196 (1982).
- [7] S. W. Golomb, SIAM Rev. **3**, 293 (1961).
- [8] M. Keane, Math. Z. **141**, 25 (1975).
- [9] M. Keane, Israel J. Math. **26**, 188 (1977).
- [10] W. A. Veech, J. Anal. Math. **33**, 222 (1978).
- [11] A. Katok, Israel J. Math **35**, 301 (1980).
- [12] H. Masur, Ann. Math. **115**, 169 (1982).
- [13] D. Aldous and P. Diaconis, Am. Math. Mon. **93**, 333 (1986).
- [14] L. N. Trefethen and L. M. Trefethen, Proc. R. Soc. Lond. A **456**, 2561 (2000).
- [15] M. Viana, Rev. Mat. Complut. **19**, 7 (2006).
- [16] A. Avila and G. Forni, Ann. Math. **165**, 637 (2007).
- [17] R. Sturman, S. W. Meier, J. M. Ottino, and S. Wiggins, J. Fluid Mech. **602**, 129 (2008).
- [18] C. F. Novak, J. Mod. Dyn. **3**, 379 (2009).
- [19] H. Hmili, Discret. Contin. Dyn. S. **27**, 1079 (2010).
- [20] M. K. Krotter, I. C. Christov, J. M. Ottino, and R. M. Lueptow, Int. J. Bifurcat. Chaos **22**, 1230041 (2012).
- [21] P. P. Park, P. B. Umbanhowar, J. M. Ottino, and R. M. Lueptow, Chaos **26**, 073115 (2016).
- [22] A. Goetz, in *Trends in Mathematics: Fractals in Graz 2001* (Birkhäuser Verlag, Basel, 2003) p. 135.

- [23] A. Goetz and G. Poggiaspalla, *Nonlinearity* **17**, 1787 (2004).
- [24] X.-C. Fu and J. Duan, *Physica D* **237**, 3369 (2008).
- [25] B. Kahng, *Chaos* **19**, 023115 (2009).
- [26] I. C. Christov, J. M. Ottino, and R. M. Lueptow, *Chaos* **20**, 023102 (2010).
- [27] J. H. Lowenstein and F. Vivaldi, *Nonlinearity* **11**, 1321 (1998).
- [28] J. H. Lowenstein and F. Vivaldi, *Chaos* **10**, 747 (2000).
- [29] B. Kahng, *Ergod. Theor. Dyn. Syst.* **22**, 483 (2002).
- [30] B. Kahng, *Dynam. Syst.* **19**, 245 (2004).
- [31] A. Goetz, *Illinois J. Math.* **44**, 465 (2000).
- [32] A. Goetz, *Nonlinearity* **14**, 205 (2001).
- [33] D. K. Umberger and J. D. Farmer, *Phys. Rev. Lett.* **55**, 661 (1985).
- [34] J. D. Farmer, E. Ott, and J. A. Yorke, *Physica D* **7**, 153 (1983).
- [35] C. Grebogi, S. W. McDonald, E. Ott, and J. A. Yorke, *Phys. Lett. A* **110**, 1 (1985).
- [36] P. Ashwin, *Phys. Lett. A* **232**, 409 (1997).
- [37] K. R. Sreenivasan and C. Meneveau, *J. Fluid Mech.* **173**, 357 (1986).
- [38] J. M. Ottino, C. W. Leong, H. Rising, and P. D. Swanson, *Nature* **333**, 419 (1988).
- [39] E. Ott and T. M. Antonsen Jr., *Phys. Rev. Lett.* **61**, 2839 (1988).
- [40] K. R. Sreenivasan, R. R. Prasad, C. Meneveau, and R. Ramshankar, *Fractals in Geophysics* (Birkhäuser Verlag, Basel, 1989) p. 43.
- [41] J. M. Ottino, F. J. Muzzio, M. Tjahjadi, J. G. Franjione, S. C. Jana, and H. A. Kusch, *Science* **257**, 754 (1992).
- [42] J. Baldyga and J. R. Bourne, *Chem. Eng. Sci.* **50**, 381 (1995).
- [43] P. E. Dimotakis and H. J. Catrakis, *GALCIT Report FM97-1* (1997).
- [44] G. O. Fountain, D. V. Khakhar, and J. M. Ottino, *Science* **281**, 683 (1998).
- [45] D. Perugini and G. Poli, *Earth Planet. Sci. Lett.* **175**, 93 (2000).
- [46] D. Perugini, G. Poli, and R. Mazzuoli, *J. Volcanol. Geoth. Res.* **124**, 255 (2003).
- [47] P. V. Danckwerts, *Appl. Sci. Res.* **3**, 279 (1952).
- [48] P. Ashwin, W. Chambers, and G. Petkov, *Int. J. Bifurcat. Chaos* **7**, 2603 (1997).
- [49] L. Wan, T. Wong, and C. Leung, *IEEE T. Vis. Comput. Gr.* **13**, 720 (2007).
- [50] T. Coleman and Y. Li, *Math. Program.* **67**, 189 (1994).
- [51] T. Coleman and Y. Li, *SIAM J. Optimiz.* **6**, 418 (1996).

[52] S. C. Jana, G. Metcalfe, and J. M. Ottino, J. Fluid Mech. **269**, 199 (1994).

### Appendix A: Reversing Symmetry of $E$ for $(\theta_1, \theta_2)$ and $(\theta_2, \theta_1)$

In Section I we noted that there is a symmetry between the structures of  $E$  for protocols  $(\theta_1, \theta_2)$  and  $(\theta_2, \theta_1)$ . Here we show why this is the case. Consider rotation angles of  $\theta_1 = 90^\circ$ ,  $\theta_2 = 45^\circ$ , and  $\theta_1 = 75^\circ$ ,  $\theta_2 = 15^\circ$  shown in Figure 14. The exceptional set is approximated for protocols  $(\theta_1, \theta_2)$  and  $(\theta_2, \theta_1)$  for both sets of angles in the left and right columns. An image transpose corresponds to reflecting points across the  $z = -x$  diagonal in the coordinate space. By transposing the images of  $E$  for the  $(\theta_1, \theta_2)$  protocol on the left, we visually confirm that the transpose is the same as the images of  $E$  for the  $(\theta_2, \theta_1)$  protocol.

We now investigate whether the progression towards  $E$  is the same for protocols  $(\theta_1, \theta_2)$  and  $(\theta_2, \theta_1)$  once the reflection across  $z = -x$  is employed, as shown in Figure 15 for rotation angles  $\theta_1 = 75^\circ$ ,  $\theta_2 = 45^\circ$ . The progressions of  $(\theta_1, \theta_2)$  and  $(\theta_2, \theta_1)$  protocols are not the same in this case, but what is remarkable is that the global features of  $E$  for both protocols become the same for large  $m$ .

The reason the structures for the  $(75^\circ, 45^\circ)$  and  $(45^\circ, 75^\circ)$  protocols are symmetric for large  $m$  becomes easier to comprehend if the  $(\theta_1, \theta_2)$  protocol is applied backwards in time,




FIG. 14. Comparing  $E$  for  $(\theta_1, \theta_2)$  and  $(\theta_2, \theta_1)$  for two different pairs of  $\theta_1$  and  $\theta_2$  (bottom view). Middle column shows the image transpose of  $E$  for  $(\theta_1, \theta_2)$ , which is a reflection across  $z = -x$  in the coordinate space. The image transpose of  $E$  for  $(\theta_1, \theta_2)$  appears identical to  $E$  for  $(\theta_2, \theta_1)$ .




FIG. 15. Comparing the progression towards  $E$  for  $\theta_1 = 75^\circ$  and  $\theta_2 = 45^\circ$  (bottom view). (Left column) progression for  $(\theta_1, \theta_2)$ , (middle column) image transpose of the left column, and (right column) progression for  $(\theta_2, \theta_1)$ . It is not apparent that the resulting  $E$  from both will be the same.




FIG. 16. Comparing  $\mathcal{D}$  (top) and  $\mathcal{D} \cup M^{-1}(\mathcal{D})$  (bottom) for the  $(75^\circ, 45^\circ)$  protocol (left column), their image transpose (middle column), and the progression towards  $E$  for the  $(45^\circ, 75^\circ)$  protocol at  $m = 1$  and  $2$  (bottom view).

denoted as  $(\theta_1, \theta_2)^{-1}$ , in which case, the rotation by  $\theta_2$  comes before the rotation by  $\theta_1$ , like  $(\theta_2, \theta_1)$  – the only difference being that the rotation by  $\theta_2$  happens on different axes for the two protocols. We compare  $\mathcal{D}$  (top) and  $\mathcal{D} \cup M^{-1}(\mathcal{D})$  (bottom) for the  $(75^\circ, 45^\circ)$  protocol in the left column, its image transpose in the middle column, and the progression for the  $(45^\circ, 75^\circ)$  protocol in the right column in Figure 16 and observe that the two structures are

the same after one is reflected across  $z = -x$ . That the set of inverse iterates of  $\mathcal{D}$  is also dense in  $E$  has been proven for some systems [27–30]. We conjecture that this is also the case here, which is why the structures of  $E$  for  $(\theta_1, \theta_2)$  and  $(\theta_2, \theta_1)$  protocols appear symmetric.

## Appendix B: Taxonomy of $E$

As noted in Section I, different patterns of  $E$  were studied at smaller angular resolutions as shown in Figures 17-19. While the continuous  $(\theta_z, \theta_x)$  parameter space cannot be fully explored, it is useful to understand the sensitivity of  $E$  to the protocol at least in some select areas of the parameter space.

Compared to Figure 5, Figure 17 with rotation angles ranging from  $45^\circ$  to  $60^\circ$  in  $3^\circ$  increments shows less variety in patterns, but patterns along the diagonal remain quite distinct from one another. With rotation angles ranging from  $45^\circ$  to  $50^\circ$  in  $1^\circ$  increments (Figure 18), there appear to be two general features – arrowheads such as those shown for the  $(45^\circ, 45^\circ)$  protocol, and five large circular cells across the diagonal with an additional large cell in the upper left corner of the HS, as shown for the  $(50^\circ, 50^\circ)$  protocol. A transition




FIG. 17.  $E$  for  $\theta_z$  and  $\theta_x \in [45^\circ, 60^\circ]$  varying in  $3^\circ$  increments.






FIG. 18.  $E$  for  $\theta_z$  and  $\theta_x \in [45^\circ, 50^\circ]$  varying in  $1^\circ$  increments.

occurs from the arrowhead feature to the five-cell feature in the parameter space between these two protocols as the cell in the center of the arrowhead pattern becomes larger and dwarfs the cells around it, so that the five arrowheads are replaced by five large circular cells.

For rotation angles ranging from  $45^\circ$  to  $46^\circ$ , the arrowhead feature is the most prominent (Figure 19). Changes in the structure of  $E$  are minimal with  $0.2^\circ$  increments in the  $45^\circ$  to  $46^\circ$  range. For protocols on the diagonal, however, their structures of  $E$  appear to have finer details (smaller cells) than adjacent protocols.




FIG. 19.  $E$  for  $\theta_z$  and  $\theta_x \in [45^\circ, 46^\circ]$  varying in  $0.2^\circ$  increments.