

This is the accepted manuscript made available via CHORUS. The article has been published as:

# Incommensurability and phase transitions in two-dimensional XY models with Dzyaloshinskii-Moriya interactions

Huiping Liu, J. A. Plascak, and D. P. Landau

Phys. Rev. E **97**, 052118 — Published 15 May 2018

DOI: [10.1103/PhysRevE.97.052118](https://doi.org/10.1103/PhysRevE.97.052118)

# Incommensurability and Phase Transitions in Two-dimensional XY models with Dzyaloshinskii-Moriya Interactions

Huiping Liu<sup>1,2</sup>, J. A. Plascak<sup>1,3</sup>, and D. P. Landau<sup>1</sup>

<sup>1</sup>*Center for Simulation Physics, University of Georgia, Athens, GA 30602, USA*

<sup>2</sup>*Department of Physics, Huazhong University of Science and Technology, Wuhan, 430074, P.R.China\* and*

<sup>3</sup>*Departamento de Física, Centro de Ciências Exatas e da Natureza, CCEN, Universidade Federal da Paraíba, Cidade Universitária, 58051-970 João Pessoa, PB - Brazil*

(Dated: April 16, 2018)

The Dzyaloshinskii-Moriya(DM) interaction in magnetic models is the result of a combination of superexchange and spin-orbital coupling, and it can give rise to rich phase transition behavior. In this paper we study ferromagnetic XY-models with DM interaction on two-dimensional  $L \times L$  square lattices using a hybrid Monte Carlo algorithm. In order to match the incommensurability between the resultant spin structure and the lattice due to the DM interaction, a fluctuating boundary condition is adopted. We also define a new kind of order parameter and use finite-size scaling to study the critical properties of this system. We find that a Kosterlitz-Thouless-like phase transition appears in this system and that the phase transition temperature shifts towards higher temperature with increasing DM interaction strength.

PACS numbers: 05.10.Ln; 05.70.Fh; 75.10.Hk

Keywords: 2D XY model, DM Interaction, Phase Transition, Incommensurability.

## I. INTRODUCTION

The XY-model in two dimensions is a prototype model for magnetic spin systems which exhibit continuous symmetry and topological excitations. This model can describe superfluid films [1] and Josephson junction arrays [2] in two dimensions and undergoes a Kosterlitz-Thouless (KT) transition characterized by an exponentially divergent correlation length and in-plane susceptibility [3, 4]. The Kosterlitz-Thouless (KT) transition is driven by the unbinding of pairs of vortices with opposite vorticity at a temperature  $T_{KT}$ .

The Dzyaloshinskii-Moriya (DM) interaction was first proposed by Dzyaloshinskii for explaining weak ferromagnetism in antiferromagnetic compounds [5], and the microscopic basis for this theory was later given by Moriya [6] who extended Anderson's superexchange theory [7] to include the spin-orbit interactions. Arising from the impurities in the system, this interaction is very important in low symmetry crystals while it vanishes in high symmetry crystals. The DM interaction exists in many materials and can lead to some special phenomena. In the superconductor  $\text{LaCu}_2\text{O}_4$ , this interaction induces a slight spin canting out of the  $\text{CuO}_2$  plane [8]. It has been reported that the DM interaction can induce helical spin order in  $\text{Fe}_x\text{Co}_{1-x}\text{Si}$  alloys [9] and the helix period is determined by the ratio of the DM interaction to the spin exchange interaction. In Mn monolayers the adjacent spins are not perfectly antiferromagnetic, but slightly canted, resulting in a spin spiral structure (with chiral order [10]) due to the DM interaction. This non-collinear, or spin spiral, order can be observed by using

spin-polarized scanning tunnelling microscopy and *ab initio* calculations identify that this spin spiral order is stabilized by DM interaction [11]. Using the polarized neutron diffraction method, a non-zero average chirality was obtained in Dy/Y multilayer films [12], which indicates that DM interaction exists in this material. The chirality is due to the lack of the symmetry inversion on the interface. Since the DM interaction plays a key role in these materials, especially these helical magnetic systems, we simulate the 2D XY-model with DM interaction in order to understand the effects of their inclusion. We found that a kind of KT-type phase transition exists in this system when the DM interaction is small [13]. Monte Carlo (MC) simulations [14] provide us with a powerful tool for the study of such complicated systems. While the Metropolis "single spin-flip" algorithm [15] is used widely, there have been many high-resolution MC algorithms developed to improve upon and to speed up the original Metropolis algorithm, e.g. the Swendsen-Wang (SW) algorithm [16].

In this paper we study the two dimensional ferromagnetic XY-model with DM interaction. This model has been recently treated in three dimensions by MC simulations using the standard single spin-flip Metropolis algorithm [17]. It has been shown that the second-order transition continuously transforms to a first-order transition by increasing the value of the DM interaction. This model has also been recently studied in two dimensions through a duality transformation and posterior RG analysis [18]. The system was mapped, in the low temperature limit, into a two-dimensional Coulomb gas model of magnetic vortices with the DM interaction playing the role of an effective electric field. By applying RG analysis, it was claimed that the effective electric field affects the vortex-antivortex pairs and destroys the Kosterlitz-Thouless transition. In this work we treat the same two-

---

\*Electronic address: liuhp@hal.physast.uga.edu

dimensional model by using a hybrid Monte Carlo algorithm which combines the Metropolis algorithm and the Swendsen-Wang algorithm. The uniform magnetization, which is regularly regarded as the order parameter in the 2D XY model [19, 20], disappears in the low temperature region because of the symmetric, undulating arrangement of the spin vectors induced by the DM interaction (see e.g. Fig. 1). In order to describe this system in the low temperature region, we define a new kind of order parameter and use a fluctuating boundary condition (FBC) [21] to match the incommensurability between the spin structure and the lattice size due to the DM interaction. By using finite-size scaling and the Binder (4th-order) cumulant of the order parameter, we can determine the phase transition temperature and, contrary to results by Proskurin et al. [18], we do find a Kosterlitz-Thouless transition as the DM interaction is varied.

This paper is organized as follows. In Sec.II the two-dimensional XY ferromagnetic model with DM interaction is introduced and the hybrid Monte Carlo algorithm is presented. We also introduce the fluctuating boundary condition briefly. In Sec.III, we first illustrate the incommensurability between the spin structure and the lattice due to DM interaction, and use the fluctuating boundary condition to match this incommensurability. Then, a new order parameter is defined and we use scaling of this order parameter with lattice size, and the crossing of Binder cumulants to estimate the transition temperature. Sec.IV concludes the paper with a summary of our results.

## II. MODEL AND METHOD

The XY ferromagnetic model with DM interaction on the two-dimensional  $L \times L$  square lattice can be written as

$$\mathcal{H} = -J \sum_{\langle i,j \rangle} \vec{S}_i \cdot \vec{S}_j - \vec{D} \cdot \sum_{\langle ij \rangle} (\vec{S}_i \times \vec{S}_j) \quad (1a)$$

$$= -J\sqrt{1+d^2} \sum_{\langle i,j \rangle} \cos(\theta_i - \theta_j - \phi), \quad (1b)$$

where  $\vec{S}_i$  is a two-component classical vector of unit length (also known as the planar rotator model), while the DM vector  $\vec{D}$  is taken to be along to the  $z$ -axis. The constants  $J$  and  $D$  are positive and denote the strengths of the nearest-neighbor ferromagnetic coupling and DM interaction, respectively. The angle brackets  $\langle i, j \rangle$  mean that the corresponding sum is over the distinct nearest-neighbor pairs of the lattice.

In going from Eq. (1a) to Eq. (1b) one expands the spin dot product in the first sum, the cross product of the spin variables in the second sum and, after taking the scalar product with the DM interaction, an additional

canonical transformation is performed, namely

$$\begin{aligned} S_i^x &= S_i^{x'} \cos \varphi_i - S_i^{y'} \sin \varphi_i, \\ S_i^y &= S_i^{x'} \sin \varphi_i + S_i^{y'} \cos \varphi_i, \end{aligned} \quad (2)$$

where  $\varphi_i$  is the rotational angle of the primed reference at the  $i$ -th spin around the  $z(=z')$ -axis. The XY format of the Hamiltonian (1b) is eventually recovered by choosing the nearest-neighbor rotational angle difference  $\varphi_i - \varphi_j$  the value  $\varphi_i - \varphi_j = \phi = \arcsin(d/\sqrt{1+d^2})$ , with  $d = D/J$  (a similar approach has been used in reference [22] in treating the quantum version of the system). In this case we have a rescaled exchange interaction  $J\sqrt{1+d^2}$ . Note that here  $\theta_i$  and  $\theta_j$  in (1b) are, respectively, the spin angles of the  $i$ th and  $j$ th sites relative to the original  $x$ -axis direction.

In order to prepare for the simulations, a new transformation can be introduced, namely

$$\theta_i = \theta_i^0 - \frac{\pi}{2}(1 + \sigma_i), \quad (3)$$

where  $\sigma_i$  takes the values  $\pm 1$  and  $\theta_i^0$  denotes a trial angle at the site  $i$ . If  $\sigma_i = -1$ ,  $\theta_i$  just keeps the same value; if  $\sigma_i = 1$ , it means that  $\theta_i$  is rotated by an angle of  $\pi$ . Combining Eqs. (1) and (2) we can obtain an Ising-type Hamiltonian

$$\mathcal{H} = - \sum_{\langle ij \rangle} J_{ij} \sigma_i \sigma_j, \quad (4)$$

where  $J_{ij} = J\sqrt{1+d^2} \cos(\theta_i^0 - \theta_j^0 - \phi)$  is an effective non-uniform Ising coupling between spins. This transformation will be very useful for implementing the Swendsen-Wang algorithm to the model.

We use a hybrid Monte Carlo algorithm which combines the standard Metropolis algorithm and the Swendsen-Wang algorithm to update the spins. At each site a new random orientation for the spin is chosen. The interaction energy between this spin and its nearest neighbors is calculated. If it is lower than the energy of the old state, the new state is accepted; otherwise, it is accepted only with a probability according to the standard Metropolis algorithm. In order to reduce the critical slowing down of simulation the Swendsen-Wang (SW) cluster algorithm is used here. In an initial spin configuration we chose  $\sigma_i = 1$  for all the sites. If  $J_{ij} > 0$ , we put a bond between  $i$ th and  $j$ th sites with a probability  $P(J_{ij}) = 1 - \exp(-2J_{ij}/k_B T)$ , otherwise no bond is set up. After clusters are constructed by putting bonds between spins, every cluster has the same possibility to rotate an angle of  $\pi$  or just to keep the same value. One SW sweep related to Eq.(3) is followed after one Metropolis sweep related to Eq.(1).

In order to match the incommensurability due to the DM interaction, we adopt a fluctuating boundary condition [21]. It is described as following: suppose there is a phase shift  $\Delta$  across the boundary at the same row and column:  $\Delta = \theta_{0,y} - \theta_{L,y}$  and  $\Delta = \theta_{x,0} - \theta_{x,L}$ , for




FIG. 1: (color online) Typical spin configuration at  $T = 0.1$  and  $d = 1$  for  $L=32$  with periodic boundary conditions. The arrows represent the spin orientations.

$0 \leq x \leq L$  and  $0 \leq y \leq L$ . After a compound Monte Carlo sweep to update the spins described above while fixing  $\Delta$ , we use the standard Metropolis to update  $\Delta$  while fixing all the spins.

We perform Monte Carlo simulations on  $L \times L$  lattices, with total sites  $N = L^2$ , to obtain the critical properties of the system. Typically,  $10^4 - 10^5$  hybrid MC steps are discarded for equilibration and  $10^7 - 10^8$  hybrid MC sweeps are then retained for averages. The sampling intervals for measuring system properties varied from 20 to 500 hybrid MC sweeps for different sizes. Multiple independent runs were used to compute statistical errors. Where not shown in the figures, error bars are smaller than the size of the symbols.

### III. RESULTS

As we know, the DM interaction would result in non-zero chirality[12]. According to Eq.(1), there is an angular difference  $\theta_i - \theta_j = \phi$  between the nearest-neighbor spins in the ground state, and  $\phi$  is determined by the ratio of the DM interaction to the spin nearest-neighbor exchange coupling. The nearest-neighbor spins prefer to be parallel to each other in the ground state if there is only exchange coupling, while the DM interaction drives the nearest-neighbor spins to be perpendicular to each other in the ground state. So this angular difference  $\phi$  results in the competition between the DM interaction and the spin exchange coupling. In order to recognize the new order of this system at low temperatures we pick up a random spin configuration when  $d = 1$  at the low

temperature  $T = 0.1$  as Fig. 1 shows (the temperature here is measured in units of  $J/k_B$ , where  $k_B$  is the Boltzmann constant). It is obvious that this angular difference, whose value is close to  $\phi$ , exists between the nearest neighbor spins  $\theta_i$  and  $\theta_j$  both in  $x$ - and in  $y$ -directions. So the nearest neighbor spins in the ground state would not be parallel to each other, as in the ground state of XY model, but rotate through an angle  $\phi$  per site both in  $x$ - and in  $y$ -directions if the spins are on the  $xy$  plane just as Fig. 1 shows. Moreover, the spin arrangement is periodic from the left bottom to the right top and the period is determined by the relative strength of DM interaction  $d$ . This periodic spin arrangement leads to the magnetization being very small at low temperatures, even smaller than at high temperatures.

Moreover, this angular difference  $\phi$  would lead to incommensurability of the system. After  $L$  sites there is a phase shift  $\Delta = L \cdot \phi$  across the boundary in the same row or column. If  $\Delta = 2n\pi$ , where  $n$  is an integer, it means that this phase shift is an integer multiple of  $2\pi$ , the structure is commensurate with the lattice and the periodic boundary condition is reasonable for this system; otherwise the structure is incommensurate with the lattice. In the latter case a periodic boundary condition (PBC) would introduce frustration and increase the energy of the system, or even result in choosing the wrong phase when the system is incommensurate. Thus, we consider a fluctuating boundary condition [21] to match the incommensurability.

How do we confirm the incommensurability of the system? As mentioned above, whether the system is incommensurate or not is dependent upon the lattice size  $L$  and the DM interaction  $d$ . For instance, when  $d = 1$ ,  $\phi = \pi/4$ , and if  $L$  is a positive integral multiple of 8, e.g. 8, 16, 32, etc., the phase shift across the boundary  $\Delta$  is also a integral multiple of  $2\pi$ , so the structure is commensurate with the lattice and the period boundary condition is suitable. But when  $d = 0.5$ ,  $\phi = 0.463648$ , and even if  $L$  is also a positive integer multiple of 8, the system turns out to be incommensurate.

In order to make some sense of the incommensurability of the spin system, we first look at the spin configurations of the incommensurate system. Fig. 2 (top and bottom) shows spin configurations of an incommensurate system for which  $L = 8$ ,  $d = 0.5$  and  $T = 0.1$ , resulting from the use of a fluctuating boundary condition (FBC) and a periodic boundary condition (PBC), respectively.

In fact, in Fig. 2(top) the spin-variation wavelength is larger than the lattice size  $L$  but is allowed by the fluctuating boundary condition, and the system just chooses a phase shift across the boundary according to the balance between the spin structure and the lattice size. When using the periodic boundary condition, we force the spin wavelength to be an integer multiple of the lattice size  $L$ . For example, in Fig. 2(bottom) the wavelength is forced to be equal to  $L$ , and  $\Delta$  must then be equal to zero. So the periodic boundary condition would increase the energy of the incommensurate system. Fig. 3 shows this




FIG. 2: (color online) Typical spin configurations at  $T = 0.1$  and  $d = 0.5$  for  $L=8$  with: (top) fluctuating boundary conditions and (bottom) periodic boundary conditions. The arrows represent the spin orientations.

in a plot of the energies versus temperature for different DM interaction and with different boundary conditions for  $L = 8$ . If  $d = 0.5$ , the system is incommensurate, the energy for PBC is larger than that found using FBC at low temperature. Consequently, FBC is better than PBC when the system is incommensurate. While the system is commensurate, such as  $L = 8$  and  $d = 1$ , PBC and FBC give almost the same answer at low temperature and PBC is suitable in order to reduce the fluctuation and save computing time. Thus, we use the periodic boundary condition if the system is commensurate, and use the fluctuating boundary condition to simulate the incommensurate system.

We thus perform Monte Carlo simulations on  $L \times L$  lattices with  $L$  ranging from  $L = 8$  to  $L = 96$ . Fig. 4 shows the specific heat versus temperature for  $d = 1$  with PBC by using the dissipation fluctuation theorem. The values of specific heat are independent of lattice size and approach  $1/2$  when the temperature  $T$  approaches zero. This is, in fact, expected on general grounds (equipartition theorem), since there is only a single degree of freedom for each spin in our 2D XY model. Moreover, a specific heat peak appears near  $T = 1.5$ , while the corre-




FIG. 3: (color online) Energy  $E/N$ , where  $E = \langle \mathcal{H} \rangle$  and the brackets mean thermal average, versus temperature  $T$  for different DM interactions when  $L = 8$  by using different boundary conditions. The errors are smaller than the symbol sizes. The lines are just guide to the eyes.




FIG. 4: (color online) Specific heat  $C_V/N$  versus temperature  $T$  for different lattice sizes when  $d = 1$  by using periodic boundary conditions. The lines are just guide to the eyes.

sponding transition temperature is near  $T = 1.02$  in the 2D XY model [23]. (It is known that the specific heat peak is above the transition temperature in the 2D-XY model, and we shall see that the same holds true here.) This means that inclusion of the DM interaction drives the transition towards a higher temperature.

The peaks of the specific heat shown in Fig. 4 do not approach a constant value monotonically. After a given value of the lattice size it starts decreasing. In fact, Fig. 5 shows the behavior of the maximum of the specific heat for larger values of the lattice size. The inset in this figure gives a finite-size approach of the form  $C_v(max) = C_v^\infty - A/L$ , where  $C_v^\infty = 1.426(8)$ , comparable to the values obtained for the XY model through a fit  $C_v(max) = C_v^\infty - A/L^\alpha$  with  $C_v^\infty = 1.44$  and  $\alpha = 1.07$  [24]. A similar behavior of the specific heat peak as a function of the lattice size has been recently obtained in the study of

the two-dimensional XY vectorial Blume-Emery-Griffiths model [25].

Since the spins rotate along both the  $x$ - and  $y$ -directions at low temperature if the strength of DM interaction is not very small, the magnetization, which is the regular order parameter in the 2D XY model [19, 20],

$$m = \frac{1}{L^2} \sqrt{\left( \sum_{i=1}^{L^2} \cos[\theta_i - (x_i + y_i)\phi] \right)^2 + \left( \sum_{i=1}^{L^2} \sin[\theta_i - (x_i + y_i)\phi] \right)^2}, \quad (5)$$




FIG. 5: (color online) Peaks of the specific heat as a function of temperature for larger lattice sizes when  $d = 1$  by using periodic boundary conditions. The full line in the inset shows a corresponding fit of the data, as discussed in the text.

where  $(x_i, y_i)$  is the coordinate of the  $i$ th spin and  $x_i, y_i \in [0, L - 1]$ .

According to finite size scaling theory [26, 27] we can analyze the properties of finite systems near the critical temperature of the corresponding infinite system. Using the definition of the order parameter given above, we show plots of the order parameter versus the size  $L$  when  $d = 0.5$  and  $d = 1$  at different temperatures in Fig. 6 and Fig. 7. Both plots show power-law scaling behavior as  $m \propto L^{-x}$  at  $T \leq T_C$  with temperature dependent exponent  $x$ , and there is no power-law behavior when  $T > T_C$ , in agreement with the behavior of an Kosterlitz-Thouless-type phase transition. In addition, the slopes of these straight lines at the “transition temperatures” are  $-0.125$  in both plots. Exactly at the transition temperature  $T = T_C$  the exponent  $x$  is equal to  $\eta/2$  [27], where  $\eta = 1/4$  is the critical exponent. In Fig. 6, when  $T \leq 1.285$ , all the plots are straight lines whose slopes are less than  $-0.125$ , while the plots are not straight lines when  $T > 1.285$ . It is clear that both of the phase transitions are Kosterlitz-Thouless-type and  $x = 0.125$ , and  $T_C = 1.014 \pm 0.001$  for  $d = 0.5$  and  $T_C = 1.284 \pm 0.001$

does not play the role of order parameter in the system. This is because of the periodic, oscillating spin arrangement that results when the DM interaction is included. According to the spin arrangement in the ground state, we define the order parameter as follows




FIG. 6: (color online) Log-log plot of the order parameter  $m$  versus lattice size  $L$  for various temperatures near  $T_C$  when  $d=0.5$  using FBC. The lines are just guide to the eyes. The longer curve corresponds to a straight line with slope  $-0.125$ .

for  $d = 1$ , respectively.

Examining the fourth order cumulant of the order parameter, originally suggested by Binder to analyze Ising model critical properties [28], is an effective way to locate the critical temperature. The Binder cumulant of the order parameter is

$$U_4 = 1 - \langle m^4 \rangle / 3 \langle m^2 \rangle^2, \quad (6)$$

where the angle brackets denote a thermal average. As the system size  $L$  approaches infinity,  $U_4 \rightarrow 0$  for  $T > T_C$  and  $U_4 \rightarrow 2/3$  for  $T < T_C$ . According to the renormalization group theory there is a “fixed point” in the  $U_4$  curves that is independent on the system size  $L$ , and the location of the “fixed point” is the critical point. So, the Binder cumulants are scale independent at the critical point, for large enough lattices, and we can use cumulant crossings for different system sizes to determine




FIG. 7: (color online) Log-log plot of the order parameter  $m$  versus lattice size  $L$  for various temperatures near  $T_C$  when  $d=1$  using PBC. The lines are just guide to the eyes. The longer curve corresponds to a straight line with slope  $-0.125$ .

the phase transition temperature  $T_C$ . Fig. 8 and Fig. 9 show the temperature dependence of Binder cumulants for different lattice sizes for  $d = 1$  and  $d = 0.5$ , respectively. It is obvious that at low temperature the values of  $U_4$  for different sizes of the system rapidly approach the same value. As the temperature increases, however, the curves separate distinctly, especially for larger lattice sizes. Such behavior also coincides with the behavior of a Kosterlitz-Thouless-type phase transition.

One can also note in Figs. 8 and 9 that despite there being a scattering of the cumulant crossings, they are not simply random, with a clear tendency to move to lower temperatures, mainly for smaller systems. This means that the scaling behavior regime for  $U_4$  should be valid for still larger lattices. However, an extrapolation scheme can be used to obtain the critical temperature, in the thermodynamic limit, by resorting to a fit of the data as a function of  $1/L$  for smaller lattices, as was proposed in Binder original paper for the spin-1/2 Ising model [28]. Using this convergence criterium for the temperatures of different lattices, in a similar way as proposed in reference [28], we estimate that  $T_C = 1.013 \pm 0.006$  for  $d = 0.5$  and  $T_C = 1.293 \pm 0.006$  for  $d = 1$ . These values agree with the results obtained by the power-law scaling of the order parameter with lattice size.

However, instead of the fits as a function of  $1/L$ , one can also use the scaling relation for the critical temperature itself (see, for instance, reference [29] for the two-dimensional Ising model). As an example, Fig. 10 shows the crossings with the smaller lattice size  $L = 8$  and the second smallest  $L = 16$ , as a function of the system size, for  $d = 1$  and  $d = 0.5$ . In this case of the Kosterlitz-Thouless transition, the corresponding finite-size-scaling




FIG. 8: (color online) Binder cumulant  $U_4$  of the order parameter versus temperature  $T$  for different lattice sizes when  $d = 0.5$  using FBC. The data symbols, with the corresponding error bars, have been omitted for clarity, except for  $T=1.07$ , in order to give an idea of the error as a function of system size.




FIG. 9: (color online) The Binder cumulant  $U_4$  of the order parameter versus temperature  $T$  for different lattice sizes when  $d=1$  using PBC. The data symbols, with the corresponding error bars, have been omitted for clarity, except for  $T=1.34$ , in order to give an idea of the error as a function of system size.

of the temperature crossings should behave as [30]

$$T_{cross} = T_C + B/(\ln L)^2, \quad (7)$$

where  $T_C$  is the transition temperature in the thermodynamic limit and  $B$  is a non-universal constant. The above relation comes from the fact that the correlation length in the infinite system behaves as  $\xi = \exp(\pi/c(T - T_C)^{1/2})$ , where  $c$  is a non-universal constant, and at  $T = T_{cross}$  one has  $\xi \simeq L$ . Fits of the data in Fig. 10 give, for  $d = 1$ ,  $T_C = 1.297(4)$ , when the smallest lattice size is  $L = 8$ , and  $T_C = 1.294(2)$  when  $L = 16$ . For  $d = 0.5$ , we have  $T_C = 1.022(9)$  with  $L = 8$  and  $T_C = 1.009(9)$  with  $L = 16$ . All the above values are in good agreement






FIG. 10: (color online) Temperature of the crossings of the Binder cumulant of the order parameter as a function of different lattice sizes when  $d = 1$  using PBC (a) and  $d = 0.5$  using FBC (b). The data correspond to considering  $L = 8$  and  $L = 16$  as the smallest lattice. The dashed lines are a fit with Eq. (7). The error bars are smaller than the symbol sizes.

with the previous ones. We can thus overall estimate  $T_C = 1.292(6)$  for  $d = 1$  and  $T_C = 1.013(4)$  for  $d = 0.5$ .

#### IV. SUMMARY

Using Monte Carlo simulations we have studied the thermodynamics and critical properties of a 2D XY model with DM and exchange interactions. Since the DM interaction drives the nearest-neighbor spins to be perpendicular to each other, the spins rotate with respect to each other at low temperature. Hence, the magnetization does not play the role of order parameter in the

2D XY model with DM interaction. The spin rotation in the low temperature regime induces incommensurability between the spin structure and the lattice. Therefore, a fluctuating boundary condition is adopted to match this kind of incommensurability, and a new kind of order parameter is defined to describe the phase transition. We use the power-law behavior of the order parameter with the lattice size and the crossing of the Binder cumulant to estimate the critical temperature. We have determined that there is a Kosterlitz-Thouless-type phase transition in the 2D XY model with DM interaction, and the DM interaction can induce an increase the Kosterlitz-Thouless transition temperature, contrary to what has been recently obtained by mapping the model to a two-dimensional Coulomb gas and by applying RG analysis [18]. We have not considered here the corresponding order parameter susceptibility because the size dependence of the susceptibility should really be the same as the size dependence of the order parameter, since they are computed from the same quantity.

#### Acknowledgments

The authors would like to thank Jian-Sheng Wang, Lin Yi, and Stefan Schnabel for stimulating discussions. This research was supported by a Fellowship from China Scholarship Council, by the Brazilian agencies CAPES and CNPq, and by NSF grant #DMR – 0810223. This study was also supported in part by resources and technical expertise from the Georgia Advanced Computing Resource Center (GACRC), a partnership between the University of Georgia's Office of the Vice President for Research and Office of the Vice President for Information Technology. We would like to address invaluable help from Shan-Ho Tsai and Zhuofei Hou regarding the use of the GACRC computing facilities at UGA.

- 
- [1] T. Ohta and D. Jasnow, Phys. Rev. B **20**, 139 (1979).
  - [2] D. J. Resnick, J. C. Garland, J. T. Boyd, S. Shoemaker, R. S. Newrock, Phys. Rev. Lett. **47**, 1542 (1981).
  - [3] J. M. Kosterlitz and D. J. Thouless, J. Phys. C: Solid State Phys. **6**, 1183 (1973).
  - [4] J. M. Kosterlitz, J. Phys. C: Solid State Phys. **7**, 1046 (1974).
  - [5] I. Dzyaloshinskii, J. Phys. Chem. Solids. **4**, 241 (1958).
  - [6] T. Moriya, Phys. Rev. Lett. **4**, 228 (1960); T. Moriya, Phys. Rev. **120**, 91 (1960).
  - [7] P. W. Anderson, Phys. Rev. **115**, 2 (1959).
  - [8] D. Coffey, K. S. Bedell and S. A. Trugman, Phys. Rev. B **42**, 6509 (1990).
  - [9] M. Uchida, Y. Onose, Y. Matsui, *et al*, Science **311**, 359 (2006).
  - [10] M. Bode, M. Heide, K. V. Bergmann, *et al* Nature **447**, 190 (2007).
  - [11] P. Ferriani, K. V. Bergmann, E. Y. Vedmedenko, *et al*, Phys. Rev. Lett. **101**, 027201 (2008).
  - [12] S. V. Grigoriev, Yu. O. Chetverikov, D. Lott, A. Schreyer, Phys. Rev. Lett. **100**, 197203 (2008).
  - [13] Y. Sun, L. Yi, X. Zhao, *et al*, Solid State Commun. **144**, 61 (2007).
  - [14] D. P. Landau and K. Binder, *A Guide to Monte Carlo Simulation in Statistical Physics* 3rd Edition, (Cambridge University Press, Cambridge, 2009).
  - [15] N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, *et al.*, Journal of Chemical Physics, **21**, 1087 (1953).
  - [16] R. H. Swendsen and J. S. Wang, Phys. Rev. Lett. **58**, 86 (1987).
  - [17] A. M. Belemuk and S. M. Stishov, Phys. Rev. B **95**, 224433 (2017).
  - [18] I. Proskurin, A. S. Ovchinnikov, and J.-i. Kishine, J. Phys.: Conf. Ser. **903** 012062 (2017).
  - [19] M. S. S. Challa and D. P. Landau, Phys. Rev. B **33**, 437 (1986).
  - [20] E. Rastelli, S. Regina, and A. Tassi, Phys. Rev. B **69**, 174407 (2004); Phys. Rev. B **70**, 174447 (2004).
  - [21] P. Olsson, Phys. Rev. Lett. **73**, 3339 (1994).
  - [22] F. C. Alcaraz and W. F. Wreszinski, J. Stat. Phys. **58** 45



- (1990).
- [23] J. Tobochnik and G. V. Chester, Phys. Rev. B **20**, 3761 (1979).
  - [24] Johannes E. Van Himbergen and Sudip Chakravarty, Phys. Rev. B **23** 359 (1981).
  - [25] J. B. Santos-Filho and J. A. Plascak, Phys. Rev. E **96** 032141 (2017);
  - [26] M. E. Fisher, in *Proceeding of the International Summer School “Enrico Fermi” 1970, Course 51*, Varenna, Italy, edited by M. S. Green (Academic, New York, 1971).
  - [27] M. S. S. Challa and D. P. Landau, Phys. Rev. B **33**, 437 (1986).
  - [28] K. Binder, Z. Phys. B: Condens. Matter **43**, 119 (1981).
  - [29] J. A. Plascak, Alan M. Ferrenberg, and D. P. Landau Phys. Rev. E **65** 066702 (2002).
  - [30] D. P. Landau and K. Binder, *A guide to Monte Carlo simulations in Statistical Physics*, Cambridge University Press, 4th Edition, Cambridge (2014).