



This is the accepted manuscript made available via CHORUS. The article has been published as:

# Electrohydrodynamic generation of atmospheric turbulence

Yuan Yao and Jesse Capecelatro

Phys. Rev. Fluids **4**, 123701 — Published 18 December 2019

DOI: [10.1103/PhysRevFluids.4.123701](https://doi.org/10.1103/PhysRevFluids.4.123701)

# Electrohydrodynamic generation of atmospheric turbulence

Yuan Yao\* and Jesse Capecelatro

*Department of Mechanical Engineering,  
University of Michigan, Ann Arbor, MI 48105*

(Dated: October 14, 2019)

## Abstract

Ionization produced by cosmic rays and atmospheric radioactivity creates charged short life-time aerosol droplets in the upper troposphere. Inhomogeneities in the spatial distribution of aerosol droplets lead to time varying electric fields and space charge, which can often be amplified by more than three orders of magnitude in extreme conditions, such as thunderstorms. The nonlinear coupling between ionized air, charged aerosol droplets, and the background electric field can result in electrohydrodynamic (EHD) body forces that augment atmospheric turbulence. In this paper, a theoretical and numerical study on the electrohydrodynamic generation of atmospheric turbulence under fair weather and thunderstorm conditions is presented. Linear stability shows that coupling between ionized air and a background electric field acts to increase turbulent kinetic energy (TKE) in the upper troposphere, albeit over long time durations. Direct simulations of charged droplets in homogeneous shear flow demonstrate a nonlinear feedback mechanism capable of accelerating the growth rate. Streamwise velocity gradients induce fluctuations in droplet concentration and electric potential, resulting in a body force that generates vertical velocity fluctuations. Pressure strain then transfers this energy to turbulent fluctuations in the streamwise direction and the process repeats. This feedback mechanism was found to augment TKE at late stages of the shear layer growth.

---

\* yyaoaa@umich.edu

## I. INTRODUCTION

Ionization produced by cosmic rays and atmospheric radioactivity creates charged short life-time aerosols that have significant spatial and temporal variations in the troposphere [1–4]. In extreme conditions, such as thunderstorms, this variability has been observed to amplify the local electric field strength in the atmosphere by more than three orders of magnitude compared to typical fair weather conditions [5, 6]. In the presence of a background electric field, gradients in space charge generate an electrohydrodynamic (EHD) body force if the gas is ionized. The EHD force that arises due to coupling between an electric field, ionized air, and charged aerosols has been shown experimentally to produce instabilities that trigger the onset of turbulent motion in quiescent aerosol clouds [7]. While EHD instabilities are known to induce turbulence in ionized fluids, such as electrolytic cells [8], liquid films [9], and electrostatic precipitators [10], and similar instabilities due to magnetohydrodynamics exist in plasmas [11–13], its role in augmenting atmospheric turbulence is less established.

Recent studies have shown that detailed interactions between turbulence and aerosol droplets can give rise to preferential concentration of charged particles capable of self-inducing atmospheric electricity [14, 15], which might amplify the aforementioned effects. Depending on the sign and magnitude of the charge each particle carries, these self-induced electrical fields can either mitigate or enhance the level of clustering [15–17]. Preferential concentration describes the accumulation of inertial (heavy) particles or droplets away from highly vortical regions of the turbulent flow [18–22]. When the time scale of the particles is on the same order as the turbulent eddies, particles are directed by coherent vortical structures to create non-homogeneities in concentration and the onset of clusters. In the presence of gravity, preferential concentration by turbulence has been observed to cause particles to accumulate near the downward moving side of vortices where the fluid velocity aligns with the direction of gravity [23–25]. The gravitational settling of aerosol particles can be enhanced by this preferential sweeping mechanism by as much as 50% [24]. The collision rate between monodisperse particles is also considerably lower when gravity is present [26].

Wind shear (mean velocity gradient) ranging from  $\mathcal{O}(10^{-3})/\text{s}$  to  $\mathcal{O}(1)/\text{s}$  is common in the atmosphere [27–29]. When particles or droplets are placed within a mean velocity gradient, the anisotropy of the velocity fluctuations induces a preferential orientation of particle clusters and modifies their turbulent transport characteristics [30, 31]. Recently,

a new preferential concentration-driven instability has been identified for inertial particles suspended in sheared turbulence [32], showing that two-way coupling between the phases in the presence of mean shear and gravity can lead to the onset of turbulence.

The motivating hypothesis of the present study is that charged aerosols within the atmosphere couple nonlinearly with the background electric field and ionized air, resulting in the production of TKE. We refer to this as a three-way coupling process, whereby velocity fluctuations induce non-homogeneities in droplet concentration, resulting in gradients in electric potential that increase the electric field strength and in turn amplifies the gas-phase velocity fluctuations. We seek to identify under what conditions EHD-induced turbulence is relevant. To this end, a linear stability analysis is performed for sheared flow in the presence of a background electric field to understand the stability criteria, relevant non-dimensional numbers, and time scales associated with EHD-induced turbulence. Then, direct numerical simulations are carried out with relevant atmospheric parameters at 6 km altitude to understand the effect of added nonlinearity by aerosol dynamics on turbulence generation. The TKE budget and growth rate are then reported to assess the significance of EHD for a range of particle loadings.

## II. LINEAR STABILITY OF THE UPPER TROPOSPHERE

### A. Governing equations

In this work, the fluid dynamics in the troposphere is modeled according to the incompressible Navier–Stokes equations, given by

$$\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} = -\frac{1}{\rho} \nabla p + \nu \nabla^2 \mathbf{u} + \frac{1}{\rho} \mathbf{F}_e - \Gamma z \frac{\partial \mathbf{u}}{\partial x} - \Gamma w \mathbf{e}_x, \quad (1)$$

where  $\rho$  is the gas-phase density,  $\mathbf{u} = [u \ v \ w]^\top$  is the velocity,  $p$  is pressure,  $\nu$  is the kinematic viscosity, and  $\mathbf{F}_e$  is an electrohydrodynamic (EHD) body force. The last two terms on the right-hand side of Eq. (1) are introduced to account for mean velocity gradients due to wind shear, where  $\Gamma$  is the shear rate,  $z$  is the vertical position and  $\mathbf{e}_x$  is the unit vector in the streamwise direction.

Within the upper troposphere, the electric permittivity is close to the vacuum permittivity ( $\epsilon \approx \epsilon_0 = 8.854 \times 10^{-12}$  F/m) and can be assumed constant. As a consequence, the EHD

body force can be expressed as  $\mathbf{F}_e = q_f \mathbf{E}$ , where the electric field,  $\mathbf{E} = -\nabla\phi$ , is determined by solving a Poisson equation for the electric potential  $\phi$ , given by

$$\nabla^2\phi = -(q_f + q_p)/\epsilon, \quad (2)$$

which accounts for both the effect of space charge density in the fluid,  $q_f$ , and charge density of aerosol droplets,  $q_p$ . In the remainder of this study, the terms aerosols, droplets, and particles will be used interchangeably. Within the first 20 km of the troposphere, there exists a vertical gradient in air conductivity that gives rise to finite space charge density [33, 34], even under fair weather conditions (see Fig. 1). We model the transport of space charge as

$$\frac{\partial q_f}{\partial t} + \nabla \cdot (\mu_I q_f \mathbf{E} - D_I \nabla q_f + q_f \mathbf{u} + q_f \Gamma z \mathbf{e}_x) = 0, \quad (3)$$

where  $\mu_I \approx 10^{-4} \text{ m}^2 \text{ V}^{-1} \text{ s}^{-1}$  is the ion mobility at 6 km altitude based on balloon-borne measurements [35] and  $D_I = \mu_I k_B T / e$  is the molecular diffusion coefficient based on the Einstein relation [36]. Here,  $k_B$  is the Boltzmann constant,  $T$  is temperature, and  $e = 1.6 \times 10^{-19} \text{ C}$  is the elementary charge. Within the first few kilometers of the atmosphere  $D_I \approx 2 \times 10^{-6} \text{ m}^2 \text{ s}^{-1}$ .




FIG. 1: Fair weather atmospheric conditions [37] used to define the background state in the stability analysis.

## B. Linear stability analysis

The flow quantities  $q_f$ ,  $\phi$ ,  $p$ ,  $\mathbf{u}$  and  $\mathbf{E}$  can be decomposed into a base state and a fluctuating component as  $A = \bar{A} + A'$ , where

$$A' = A_0(z) \exp(i\kappa_x x + i\kappa_y y) \exp(\omega\tau), \quad (4)$$

with  $A_0$  the fluctuation amplitude,  $\kappa_x$  and  $\kappa_y$  the horizontal wave numbers,  $\omega$  the growth rate, and  $\tau = (\mu_I \bar{\phi} t)/L^2$  is the time scale normalized over a vertical span  $L$ . Linearizing (1)–(3) and enforcing  $\nabla \cdot \mathbf{u} = 0$ , the following set of ODEs of total order 7 can be obtained to find solutions for the non-dimensional fluctuation amplitudes of vertical velocity,  $f(z) = w_0(z)L/(\mu_I \bar{\phi}(z))$ , and potential,  $g(z) = \phi_0(z)/\bar{\phi}(z)$ , according to

$$\begin{aligned} \left( \frac{\omega}{\text{Pr}} - (D^2 - \alpha^2) \right) (D^2 - \alpha^2) f &= -R \frac{d\bar{q}_f}{dz} \frac{L^3}{\epsilon \bar{\phi}} \alpha^2 g \\ &- R \frac{\bar{E}L}{\bar{\phi}} (D^2 - \alpha^2) \alpha^2 g - S \hat{z} \beta (D^2 - \alpha^2) f - 2S \beta D f, \end{aligned} \quad (5)$$

$$\begin{aligned} -\frac{\epsilon \bar{E}}{L^2} (D^2 - \alpha^2) Dg + \frac{d\bar{q}_f}{dz} Dg + \frac{\epsilon \bar{\phi}}{L^3} \omega (D^2 - \alpha^2) g \\ + \frac{2\bar{q}_f}{L} (D^2 - \alpha^2) g = \frac{d\bar{q}_f}{dz} f. \end{aligned} \quad (6)$$

Here,  $\hat{z} = z/L$ ,  $\beta = L d/dx$  and  $D = L d/dz$  are the normalized derivative operators and  $\alpha = L \sqrt{\kappa_x^2 + \kappa_y^2}$  is the normalized wavenumber. This analysis yields the following non-dimensional numbers

$$R = \frac{\epsilon \bar{\phi}}{\mu \mu_I}, \quad \text{Pr} = \frac{\mu}{\rho \bar{\phi} \mu_I}, \quad S = \frac{\rho \Gamma L^2}{\mu} \quad (7)$$

that dictate stability and growth rate. Details on the derivation of the linearized equations and associated boundary conditions employed in the analysis are given in Appendix B, and a similar formulation can be found in Schneider and Watson [38] for a charged dielectric liquid film.  $R$  is sometimes referred to as the electric Rayleigh number [10] and represents the competition between turbulence production due to charge density fluctuations and dissipation due to viscosity. When  $R$  is greater than a critical value  $R_c$ , velocity fluctuations will increase with the Coulomb force, and below  $R_c$  the EHD-induced turbulence is damped by viscosity. Pr is the electric Prandtl number, a measure of electric time scale to the hydrodynamic time scale [38].  $S$  is the shear Reynolds number that characterizes the importance of mean shear inertial forces to viscous forces over the vertical span  $L$ . Note that since  $\bar{\phi}$  is also a function of  $L$ , all three non-dimensional numbers are functions of  $L$ .

### C. Effects of vertical span and shear rate

The ODEs reported in Sec. II B are solved numerically using a 4-th order Runge–Kutta scheme for a vertical thickness of the atmosphere  $z_0 \leq z \leq z_0 + L$  with a base altitude  $z_0 = 6$  km. We first consider base state values corresponding to fair weather conditions within the troposphere [37] (see Fig. 1). Figure 2 shows the minimum values of  $R_c$  such that the flow remains stable over all wave numbers,  $\alpha$ , as a function of vertical span  $L$ . It can immediately be seen that the flow becomes more unstable as the vertical span increases. It is interesting to note that  $R \approx 100$  in the troposphere, which corresponds to a vertical span of  $L \approx 1$  km, consistent with typical cloud thicknesses observed at those elevations [39].




FIG. 2: Critical value of the electric Rayleigh number as a function of vertical span  $L$  for different shear Reynolds numbers  $S$ . The system is unconditionally stable when  $L \leq 0.707$  km (denoted by the vertical dashed line).

Wind shear is also known to play an important role in atmospheric dynamics, and is responsible for Kelvin-Helmholtz instabilities that influence the turbulence energy budget [40]. Figure 2 shows that the system is unconditionally stable for  $L \leq 0.707$  km regardless of wind shear, but for  $L > 0.707$  km,  $R_c$  decreases with increasing shear rate, demonstrating shear acts to destabilize the flow, as expected. Even modest values of wind shear, i.e.  $\Gamma > 1 \times 10^{-3} \text{ s}^{-1}$ , will lead to values of  $S$  significantly larger than those considered in Fig. 2, affirming that atmospheric electricity (even in fair weather conditions) is a destabilizing

mechanism in the troposphere. We note that the stability analysis neglects gravity, and thus these findings isolate the role of electric body forces on atmospheric stability.

Of key interest in the present study is the rate at which small disturbances in the velocity field grow due to EHD. The growth rate,  $\omega$ , is obtained using prescribed values of  $\alpha$ ,  $R$  and  $Pr$ . Using typical fair weather conditions ( $Pr \approx 2.5 \times 10^{-6}$ ), Fig. 3 shows that the maximum growth of vertical velocity fluctuations,  $w'$ , by EHD is slow, taking approximately  $10^9$  s to amplify the fluctuations by two orders of magnitude. Using values relevant to thunderstorm conditions ( $\bar{E} = 3 \times 10^5$  V/m and  $\bar{\phi} = 2 \times 10^8$  V), similar growth is achieved in approximately  $10^3$  s. However, despite this enhanced growth rate, the time scales are still relatively slow, even as the shear rate varies by several orders of magnitude. It is important to note that the present linear stability analysis does not take into account detailed aerosol dynamics that are known to be significant in turbulent flows. Numerical simulations that explicitly take these dynamics into account will be presented in the following section.




FIG. 3: Normalized growth of vertical velocity fluctuations under varying shear rates predicted by linear stability analysis under fair weather (solid fill) and thunderstorm (hatch fill) conditions.



### III. NUMERICAL SIMULATIONS

#### A. Simulation configuration

Direct numerical simulations are performed to study the role of nonlinearities associated with aerosol dynamics not accounted for in the linear stability analysis. We consider charged particles suspended in homogeneous shear flow in the presence of a background electric field (see Fig. 4). The flow field and space charge are initialized using simple turning wave vectors referred to as Kelvin modes. These wave vectors are initially oriented as upstream waves that rotate as the flow is being sheared and transform into downstream waves. Kelvin modes are known to first experience a transient algebraic growth (a bypass transition) to turbulence and decay after they pass the turning point to downstream waves [41, 42]. Nonlinear effects may couple with the initial transient growth leading to a ‘bootstrapping’ mechanism [43]. If the electrostatic time scale is significantly smaller than the fluid time scale, heterogeneity in charge distribution may give rise to local EHD body forces during the transient period and prevent or postpone the fluid TKE from decaying.




FIG. 4: Simulation configuration for  $\Phi_v = 1 \times 10^{-4}$ . Gray scale shows fluid velocity magnitude, ranging from 0 (black) to its maximum value (white). Particles are colored by charge: positive (red) and negative (blue).

We consider a triply-periodic domain of length 0.256 m in each direction, discretized using  $256^3$  grid points. The maximum amount of charge each particle can carry is dictated by the Rayleigh limit,  $Q_R = \pi \sqrt{8\epsilon\gamma d_p^3}$ , where  $\gamma = 0.0738$  N/m is the surface tension coefficient

for water droplets and  $d_p$  is the particle diameter. The particle charge is taken to be  $Q_p = \pm 0.1Q_R$  and the average particle volume fraction varies between  $0 \leq \Phi_v \leq 2 \times 10^{-3}$ , corresponding to typical aerosol properties observed in thunderstorm conditions [37, 44]. The root-mean-square EHD fluctuations,  $(q_f E)_{\text{rms}} \equiv \sqrt{\langle (q_f E - \bar{q}_f \bar{E})^2 \rangle}$ , is fixed as the particle loading varies, where angled brackets denote a volume average. Given that  $\bar{E}$  is constant and  $\bar{q}_f = 0$ ,  $(q_f E)_{\text{rms}} = \langle q_f^2 E'^2 \rangle$ . Since the self-induced electric field  $E'$  is proportional to the local particle-phase charge density  $q_p$ ,  $(q_f E)_{\text{rms}} \sim \langle q_f^2 q_p^2 \rangle$  is held constant for each simulation. This is done to avoid a trivial solution (e.g., increasing the effect of EHD with increasing volume fraction) and isolate the effect of particle dynamics on TKE production. Therefore, as  $\Phi_v$  increases,  $q_f$  decreases accordingly (see Table I).

## B. Numerics

The fluid-phase equations (1)–(3) are solved on a staggered grid with second-order spatial accuracy for both the convective and viscous terms, and the second-order accurate semi-implicit Crank–Nicolson scheme is used for time advancement [45]. Periodic shear boundary conditions are enforced in the  $z$ -direction using a discretely conservative algorithm recently proposed by Kasbaoui *et al.* [42]. Individual aerosol particles are tracked in a Lagrangian manner [46] governed by Newton’s second law of motion

$$\frac{d\mathbf{x}_p^{(i)}}{dt} = \mathbf{v}_p^{(i)} \quad \text{and} \quad m_p \frac{d\mathbf{v}_p^{(i)}}{dt} = \mathbf{F}_{\text{drag}}^{(i)} + \mathbf{F}_c^{(i)}, \quad (8)$$

where  $\mathbf{x}_p^{(i)}$  and  $\mathbf{v}_p^{(i)}$  are the instantaneous position and velocity of particle ‘ $i$ ’, respectively, and  $m_p$  is its mass. The drag force is modeled according to the standard drag correlation of Schiller and Naumann [47] as

$$\mathbf{F}_{\text{drag}}^{(i)} = \frac{m_p}{\tau_p} (\mathbf{u}[\mathbf{x}_p^{(i)}] - \mathbf{v}_p^{(i)}) (1 + 0.15 \text{Re}_p^{0.687}), \quad (9)$$

where  $\mathbf{u}[\mathbf{x}_p^{(i)}]$  is the fluid velocity interpolated to the center position of the  $i$ -th particle,  $\text{Re}_p = |\mathbf{u}[\mathbf{x}_p^{(i)}] - \mathbf{v}_p^{(i)}| d_p / \nu$  is the particle Reynolds number, and  $\tau_p = \rho_p d_p^2 / (18\mu\nu)$  is the particle response time. The electrostatic force is given by

$$\mathbf{F}_c^{(i)} = Q_p^{(i)} \mathbf{E}[\mathbf{x}_p^{(i)}], \quad (10)$$

where  $Q_p^{(i)}$  is the charge of the  $i$ -th particle and  $\mathbf{E}[\mathbf{x}_p^{(i)}]$  is the electric field at the particle position. The particle concentrations are low enough such that inter-particle collisions and

direct two-way coupling due to interphase momentum exchange can be neglected. Instead, aerosols affect the gas phase indirectly via their charge distribution.

The aerosol charge density,  $q_p$ , is obtained by projecting the particle charge to the Eulerian grid according to

$$q_p(\mathbf{x}, t) = \sum_{i=1}^{N_p} Q_p^{(i)} \mathcal{W}(\mathbf{x} - \mathbf{x}_p^{(i)}(t)), \quad (11)$$

where  $\mathcal{W}$  is a second-order accurate polynomial weighting function and  $N_p$  is the number of particles. A second-order Runge–Kutta scheme is used for updating each particle’s position and velocity. Further details on the numerical implementation can be found in Yao and Capecelatro [15].

Because the simulation domain is periodic,  $q_f$  and  $\mathbf{E}$  must be decomposed into mean and fluctuating components and only  $q'_f$  and  $\mathbf{E}'$  are solved for. Equation (2) can be rewritten as  $\nabla^2 \phi' = -(q'_f + q_p)/\epsilon$  and the electric field fluctuation can be solved via  $\mathbf{E}' = -\nabla \phi'$ . We note that particles are assigned equal and opposite charge such that  $\langle q_p \rangle = 0$  and the Poisson equation is well posed. The total electric field is then reconstructed as  $\mathbf{E} = \mathbf{E}' + \overline{\mathbf{E}}$ , and similarly  $q_f = q'_f + \overline{q}_f$ , with  $\overline{\mathbf{E}} = [0 \ 0 \ \overline{E}]^\top$ . In each simulation,  $\overline{q}_f = 0$  and the amplitude of  $q'_f$  is prescribed as  $q_{f0}$ . Parameters are chosen to correspond to thunderstorm conditions at an altitude of 6 km as summarized in Table. I.

$\Phi_v$	Mean volume fraction	0	$1 \times 10^{-4}$	$4 \times 10^{-4}$	$1 \times 10^{-3}$	$2 \times 10^{-3}$
$N_p$	Number of particles	0	$3.2 \times 10^6$	$1.3 \times 10^7$	$3.2 \times 10^7$	$6.4 \times 10^7$
$q_{f0}$ [C/m <sup>3</sup> ]	Space charge fluctuation	$2 \times 10^{-8}$	$1 \times 10^{-8}$	$5 \times 10^{-9}$	$3.2 \times 10^{-9}$	$2.2 \times 10^{-9}$
$d_p$	Particle diameter	100 $\mu\text{m}$				
$Q_p$	Particle charge	$\pm 7.27 \times 10^{-13}$ C				
$\rho_p/\rho_f$	Density ratio	1000				
$\overline{E}$	Background electric field	$3.0 \times 10^5$ V/m				

TABLE I: Parameters used in the homogeneous shear simulations.  $q_{f0}$  is varied with  $\Phi_v$  such that  $(q_f E)_{\text{rms}} \sim \sqrt{\langle q_p^2 q_f^2 \rangle}$  is held constant. All simulations are performed with conditions of  $d_p/\Delta x = 0.1$  and  $\Gamma = 1.5$ .

### C. Evolution and growth of TKE

The aerosol droplets are initially randomly distributed throughout the domain. As shown in Fig. 5, horizontal wind shear combined with the background electric field leads to the onset of particle clustering in high strain regions, which eventually generates fluctuations in fluid velocity via coupling with the charge density. Particle clustering arises from a similar mechanism identified by Kasbaoui *et al.* [32], except here the body force is due to the electric field instead of gravity, and the momentum coupling exists through the charge density instead of the drag force. In the absence of charged droplets, the initial Kelvin modes are rotated and decay without introducing any fluctuations. Gas-phase velocity fluctuations become more intense with increasing droplet loading. To identify the mechanisms responsible for this, we evaluate the evolution of Reynolds stresses, governed by

$$\frac{d\langle u'_i u'_j \rangle}{dt} = \mathcal{P}_{ij} + \mathcal{R}_{ij} - \varepsilon_{ij} + \mathcal{E}_{ij}, \quad (12)$$

where  $\mathcal{P}_{ij}$ ,  $\mathcal{R}_{ij}$  and  $\varepsilon_{ij}$  denote mean-gradient production, pressure strain, and viscous dissipation, respectively [48] given by

$$\mathcal{P}_{ij} \equiv -\langle u'_i u'_k \rangle \frac{\partial \langle u_j \rangle}{\partial x_k} - \langle u'_j u'_k \rangle \frac{\partial \langle u_i \rangle}{\partial x_k}, \quad (13)$$

$$\mathcal{R}_{ij} \equiv \left\langle \frac{p'}{\rho} \left( \frac{\partial u'_i}{\partial x_j} + \frac{\partial u'_j}{\partial x_i} \right) \right\rangle, \quad (14)$$

$$\varepsilon_{ij} \equiv 2\nu \left\langle \frac{\partial u'_i}{\partial x_k} \frac{\partial u'_j}{\partial x_k} \right\rangle, \quad (15)$$

and

$$\rho \mathcal{E}_{ij} = \langle E_j \rangle \langle q'_f u'_i \rangle + \langle E_i \rangle \langle q'_f u'_j \rangle + \langle q'_f E'_j u'_i \rangle + \langle q'_f E'_i u'_j \rangle \quad (16)$$

are contributions from EHD. In the present configuration,  $\langle E_1 \rangle = \langle E_2 \rangle = 0$  and  $\langle E_3 \rangle = \overline{E}$ . Also note the last two terms on the right-hand side of Eq. (16) containing triply correlations were found to be negligible compare to the first two terms. Taking one-half the trace of (12) yields a transport equation for the TKE, defined as  $\langle u'_i u'_i \rangle / 2$  (repeated indices imply summation). The contributions from EHD,  $\mathcal{E}_{ii} / 2$ , contain two terms.  $\mathcal{E}_P = \overline{E} \langle q'_f w' \rangle / \rho$  represents production due to the background electric field and correlation between charge density and vertical velocity fluctuations. The second contribution,  $\mathcal{E}_T = \langle q'_f E'_i u'_i \rangle / \rho$  is found to be negligible in the present study.




FIG. 5: Temporal evolution of the gas and particle phases with volume fraction increasing from top to bottom and time increasing from left to right. Color scheme same as Fig. 4.

The temporal evolution of TKE is shown in Fig. 6. All cases exhibit similar algebraic growth at early times. An analytic solution for this early time behavior is provided in Appendix A, which is shown to be independent of the aerosol distribution. It can immediately be seen that increasing particle loading enhances TKE post the initial algebraic growth ( $t \gtrsim 1$  s). By transferring the charge density from the carrier phase to aerosol particles, the TKE is amplified by as much as a factor of 6 for the highest loading under consideration. It should be noted that the size of the simulation domain is constrained by the high computational cost in tracking  $\mathcal{O}(10^8)$  individual particles. Based on the linear stability analysis performed in Sec. II C, larger domain sizes should result in greater enhancement of TKE. Moreover, a

scaling analysis by Di Renzo and Urzay [14] demonstrated that while clustering occurs at the microscale (at the same scale as the smallest turbulent eddies), the self-induced electric field can span much larger scales. Thus, the effects reported here are anticipated to be greater under more relevant length scales.




FIG. 6: Normalized TKE evolution for particle loading  $\Phi_v = 0$  (—),  $1 \times 10^{-4}$  ( $\cdots$ ),  $4 \times 10^{-4}$  ( $-\cdot-$ ),  $1 \times 10^{-3}$  (—) and  $2 \times 10^{-3}$  (- -).

#### D. Reynolds stress budget

After the initial algebraic growth is complete, turbulence modulation by charged aerosols is apparent. The temporal evolution of Reynolds stresses are shown in Fig. 7. When aerosol particles are absent ( $\Phi_v = 0$ ), the vertical component of Reynolds stress,  $\langle w'w' \rangle$ , initially grows due to production by EHD ( $\mathcal{E}_{33} = 2\overline{E}\langle q'_f w' \rangle / \rho$ ). This term then contributes to mean-gradient production ( $\mathcal{P}_{13} = -\langle w'w' \rangle \Gamma$ ) that increases  $\langle u'w' \rangle$ . Next, the energy is transferred to the streamwise component  $\langle u'u' \rangle$ . Because there is no mechanism to sustain TKE when charged aerosol droplets are absent, all three components decay to zero at late times due to viscous dissipation, as shown in Fig. 6. When charged droplets are introduced, the Reynolds stresses follow a similar trend at early times and deviate at later times. Only the vertical component  $\langle w'w' \rangle$  approaches zero, while  $\langle u'w' \rangle$  levels off to a negative value, and the streamwise component,  $\langle u'u' \rangle$ , is seen to increase. This effect is enhanced with increasing

particle volume fraction as shown in Figs. 7(b)–7(d).




FIG. 7: Evolution of the Reynolds stresses normalized by  $w_0^2$  for (a)  $\Phi_v = 0$ , (b)  $1 \times 10^{-4}$ , (c)  $4 \times 10^{-4}$  and (d)  $1 \times 10^{-3}$ .

The Reynolds stress budgets are reported for  $\Phi_v = 1 \times 10^{-4}$  in Fig. 8 to further understand how energy is transferred between the separate components. We note that the trends reported here are consistent at other values of  $\Phi_v$ . The initial sinusoidal fluctuation in space charge and vertical velocity together with the background electric field generate a non-zero EHD source  $\mathcal{E}_{33} = 2\overline{E}\langle q'_f w' \rangle / \rho$ . As a result, the vertical component of Reynolds stress,  $\langle w'w' \rangle$ , grows due to this production terms (see Fig. 8(a)). A significant portion of this energy is also transferred to the corresponding pressure strain,  $\mathcal{R}_{33} = \langle 2p'(\partial w' / \partial z) / \rho \rangle$ , and consequently  $\langle w'w' \rangle$  decays to zero at late times.

The EHD production term,  $\mathcal{E}_{33}$ , then decays as a result of diffusion of space charge, at which time pressure strain,  $\mathcal{R}_{13} = \langle p'(\partial u' / \partial z + \partial w' / \partial x) / \rho \rangle$ , acts to redistribute the




FIG. 8: Budget of (a)  $\langle w'w' \rangle$ , (b)  $\langle u'w' \rangle$  and (c)  $\langle u'u' \rangle$  normalized by  $\Gamma w_0^2$  for  $\Phi_v = 1 \times 10^{-4}$ . (d)  $\langle u'u' \rangle$  normalized by  $\Gamma w_0^2$  in the absence of particles.

energy to  $\langle u'w' \rangle$ . Fig. 8(b) reveals that while the streamwise velocity fluctuations increase, a new EHD source term,  $\mathcal{E}_{13} = \overline{E} \langle q_f' u' \rangle / \rho$ , emerges, further augmenting  $\langle u'w' \rangle$ . At late time, pressure strain continues to transfer energy from  $\langle u'w' \rangle$ , similar to  $\langle w'w' \rangle$ , and the corresponding mean-gradient production,  $\mathcal{P}_{13} = -\langle w'w' \rangle \Gamma$ , is negative and acts as a sink of  $\langle u'w' \rangle$ . As a result,  $\langle u'w' \rangle$  eventually decays and approaches a negative value when production and dissipation balance.

Fig. 8(c) shows the budget for the streamwise component of the Reynolds stress,  $\langle u'u' \rangle$ . As energy is transferred to the streamwise components,  $\langle u'u' \rangle$  is first augmented by the corresponding pressure strain,  $\mathcal{R}_{11} = \langle 2p'(\partial u' / \partial x) / \rho \rangle$ , during the early stage, then mean-gradient production,  $\mathcal{P}_{11} = -2\langle u'w' \rangle \Gamma$ , acts as a source when  $\langle u'w' \rangle$  reaches a negative equilibrium that later sustains  $\langle u'u' \rangle$ . Note that unlike the other two components of Reynolds



stress, the streamwise component always has non-zero production and therefore continues to increase even at late time. Comparing Figs. 8(c) and 8(d), it can be seen that this non-zero equilibrium state of production ( $\mathcal{P}_{11}$ ) only occurs when charged aerosols are present.

A schematic summarizing the energy transfer is illustrated in Fig. 9. As depicted by the solid lines, a background electric field combined with fluctuations in velocity and space charge yields production of TKE that is eventually dissipated into heat. In the absence of charged droplets, no other mechanism exists to further produce TKE. However, when aerosol droplets are present, turbulence can be sustained through two additional processes: (i) turbulence preferentially concentrates charged particles, resulting in fluctuations in electric potential that self-induces an electric field, enhancing  $\mathcal{E}_{33}$  and TKE; and (ii) the space charge fluctuation induced by the electric field contributes to another EHD source term,  $\mathcal{E}_{13}$ , which enhances  $\langle u'w' \rangle$ , contributing to mean-gradient production ( $\mathcal{P}_{11}$ ) that further augments TKE. This three-way coupling gives rise to a feedback loop that enhances TKE beyond the initial algebraic growth.

#### IV. CONCLUSION

We presented a theoretical and numerical study on the electrohydrodynamic generation of atmospheric turbulence under fair weather and thunderstorm conditions. The focus of this study was on the three-way coupling between turbulence, charged aerosol droplets, and a background electric field. A canonical flow configuration was presented to simulate the upper troposphere, in which a random distribution of charged particles was placed in homogeneous shear flow with a uniform electric field.

Linear stability analysis was performed using base state parameters relevant to fair weather conditions in the upper troposphere. Atmospheric stability was characterized by the electric Rayleigh number, which was found to be a function of vertical span and shear rate. The stability analysis revealed that the atmospheric fluid dynamics are most unstable over a vertical span consistent with typical cloud thicknesses observed at these elevations. The growth rate in velocity fluctuations was estimated using both fair weather and thunderstorm parameters, which was found to be slow even under thunderstorm conditions.

Direct numerical simulations revealed a nonlinear feedback mechanism that acts to accelerate turbulence production under thunderstorm conditions. It was found that horizontal




FIG. 9: Energy transfer due to three-way coupling between turbulence, charged particles, and a background electric field. The dashed lines represent energy transfer when aerosol particles are present. Fluctuations in vertical velocity and space charge give rise to an EHD source term,  $\mathcal{E}_{33}$ , that produces TKE. In the absence of charged aerosol particles, TKE is eventually dissipated to heat. When particles are present, preferential concentration by the turbulence generates inhomogeneities in particle distribution and consequently gradients in electric potential. This further contributes to the EHD source term  $\mathcal{E}_{33}$  in addition to  $\mathcal{E}_{13}$  that generates TKE via mean gradient production.

wind shear combined with EHD body forces led to the onset of particle clustering that generated fluctuations in charge density. These fluctuations acted to augment turbulent kinetic energy (TKE) after the initial algebraic growth. TKE was initially produced in the vertical direction and was later transferred to the streamwise (shear) components. The streamwise velocity gradients acted to induce fluctuations in aerosol concentration and consequently generated gradients in electric potential, resulting in an increase in the EHD body force and the process repeated. TKE was enhanced by up to a factor of six as the particle loading increased.

The present numerical and theoretical analysis demonstrated that charged aerosols in the atmosphere can couple nonlinearly with the background electric field to enhance the

onset of TKE. It is important to note that other effects may be at play as well. Particles are capable of transferring charge among each other or acquire charge through triboelectric effects [49–52]. In addition, the droplet size may vary due to evaporation and condensation. Additional instabilities may also arise when gravity is present, which may add to the TKE augmentation studied here. In the present study, we isolate the effect of charged aerosols on turbulence generation, and therefore these effects were not included. However, the Eulerian–Lagrangian simulation framework can be extended to account for these kinetic processes in TKE augmentation in future work.

## V. ACKNOWLEDGEMENT

The authors wish to thank Dr. Housseem Kasbaoui for valuable discussions and providing the homogeneous shear code. The computing resources and assistance provided by the staff of Advanced Research Computing at The University of Michigan, Ann Arbor is greatly appreciated. This work was supported in part by the Office of Naval Research (ONR).

### Appendix A: Analytical solution of the initial algebraic growth

In homogeneous shear flows, the initial algebraic growth is known to sustain large amplification of perturbation energy that may introduce nonlinear effects and lead to onset of turbulence [53–57]. Because conventional linear stability analysis fails to accurately predict the transient growth, other techniques, such as non-normal modal analysis [43, 58, 59], or the WentzelKramersBrillouin (WKB) method [60, 61], have been used in the past. In the present analysis, we consider the method of characteristics. By neglecting the particle phase, Eqs. (1) and (3) are rewritten along characteristics in Fourier space, together with the wavevector characteristic equations, resulting in the following three coupled ODEs

$$\begin{aligned}\frac{d\boldsymbol{\kappa}}{dt} &= -\nabla \mathbf{u}_b^T \cdot \boldsymbol{\kappa} \\ \frac{d\hat{q}_f}{dt} &= -i\mu_I \bar{E} \kappa_3 \hat{q}_f - D_I \kappa^2 \hat{q}_f \\ \frac{d\hat{w}}{dt} &= \frac{2\Gamma \kappa_1 \kappa_3}{\kappa^2} \hat{w} - \nu \kappa^2 \hat{w} + \left(1 - \frac{\kappa_3^2}{\kappa^2}\right) \frac{\hat{q}_f \bar{E}}{\rho},\end{aligned}\tag{A1}$$

where the hat denotes the Fourier transform of a quantity,  $\boldsymbol{\kappa} = [\kappa_1 \ \kappa_2 \ \kappa_3]^T$  is the wavevector, and  $\nabla \mathbf{u}_b = \Gamma \mathbf{e}_1 \mathbf{e}_3^T$  is the base state velocity gradient tensor. The vertical velocity can be

determined by assuming  $w'(t, \mathbf{x}) = \hat{w}(t) \exp(i\boldsymbol{\kappa}(t) \cdot \mathbf{x})$ . The remaining velocity components can be obtained by enforcing continuity. As shown in Fig. 10, the second-order algebraic growth closely matches the numerical simulations from Sec. III, confirming that charged aerosols do not affect the initial algebraic growth rate.




FIG. 10: Normalized TKE evolution for particle loading  $\Phi_v = 0$  (—),  $1 \times 10^{-4}$  (⋯),  $4 \times 10^{-4}$  (— · —),  $1 \times 10^{-3}$  (—), and  $2 \times 10^{-3}$  (— · —). Theoretical solution obtained from Eq. (A1) (○) are shown for comparison. The vertical dashed line indicates the turning point from upstream and downstream waves.

The second-order algebraic initial growth rate can be approximated analytically from Eq. (A1). Solving the wavevector characteristic equation yields  $\kappa_1 = \kappa_0 = 2\pi/L$  and  $\kappa_3 = (1 - \Gamma t) \kappa_0$ . With this, the magnitude of  $\hat{q}_f$  can easily be obtained by solving the first-order linear ODE in Eq. (A1)

$$\hat{q}_f(t) = \hat{q}_{f0} \exp\left(-\frac{1}{3} D_I \kappa_0^2 t (\Gamma^2 t^2 - 3\Gamma t + 6)\right). \quad (\text{A2})$$

Solving  $\hat{q}_f$ , the third ODE for velocity results in a first-order linear equation and can be approximated via a Taylor series expansion according to

$$\begin{aligned} \frac{d\hat{w}}{dt} &= \frac{1}{1 + (1 - \Gamma t)^2} \frac{\hat{q}_{f0} \bar{E}}{\rho} \exp\left(-\frac{1}{3} D_I \kappa_0^2 t (\Gamma^2 t^2 - 3\Gamma t + 6)\right) \\ &\approx \frac{1}{1 + (1 - \Gamma t)^2} \frac{\hat{q}_{f0} \bar{E}}{\rho} (1 - 2D_I \kappa_0^2 t + D_I \kappa_0^2 (\Gamma + 2D_I \kappa_0^2) t^2). \end{aligned} \quad (\text{A3})$$

The first two terms on the right-hand-side can be neglected since they are orders of magnitude smaller than the last term. Integrating Eq. (A3) yields

$$\begin{aligned}\hat{w}(t) &= \hat{w}_0 - \frac{\hat{q}_{f0}\overline{E}}{\rho\Gamma^2}(\Gamma - 2D_I\kappa_0^2)\left(\tan^{-1}(1 - \Gamma t) - \frac{\pi}{4}\right) \\ &\approx \hat{w}_0 + \frac{\hat{q}_{f0}\overline{E}}{2\rho\Gamma}(\Gamma - 2D_I\kappa_0^2)t.\end{aligned}\tag{A4}$$

By applying continuity, the streamwise velocity can be expressed as

$$\hat{u}(t) = -\frac{\hat{w}(t)\kappa_3}{\kappa_1} = -\hat{w}_0(1 - \Gamma t) - \frac{\hat{q}_{f0}\overline{E}}{2\rho\Gamma}(\Gamma - 2D_I\kappa_0^2)(1 - \Gamma t)t.\tag{A5}$$

With this, the TKE can be approximated by

$$\text{TKE}(t) = \frac{1}{2}(\hat{w}^2 + \hat{u}^2) \approx \text{TKE}_0 + \frac{\hat{q}_{f0}^2\overline{E}^2}{2\rho^2\Gamma^2}(\Gamma - 2D_I\kappa_0^2)^2 t^2 = \mathcal{O}(t^2),\tag{A6}$$

which implies when  $t$  is small, the initial growth of TKE is second-order. This growth rate is also consistent with the well-known quadratic-scaling law for the transient energy growth in uniform shear flow [57, 62].

## Appendix B: Linearization and non-dimensionalization of the governing equations

We start with the governing equations (1) and (3) and neglect the convective term in the momentum equation and diffusion term in the charge transport equation due to their relatively small magnitudes

$$\left(\rho\frac{\partial}{\partial t} - \mu\nabla^2\right)\mathbf{u} = q_f\mathbf{E} - \nabla p - \rho\Gamma z\frac{\partial\mathbf{u}}{\partial x} - \rho\Gamma w\mathbf{e}_x,\tag{B1}$$

$$\frac{\partial q_f}{\partial t} + \nabla \cdot (\mu_I q_f \mathbf{E} + q_f \mathbf{u} + q_f \Gamma z \mathbf{e}_x) = 0.\tag{B2}$$

The flow quantities  $q_f$ ,  $\phi$ ,  $p$ ,  $\mathbf{u}$  and  $\mathbf{E}$  are decomposed into a base state and a fluctuating component as  $A = \overline{A} + A'$ , where

$$A' = A_0(z) \exp(i\kappa_x x + i\kappa_y y) \exp(\omega\tau),\tag{B3}$$

with  $A_0$  the fluctuation amplitude,  $\kappa_x$  and  $\kappa_y$  the horizontal wave numbers,  $\omega$  the growth rate, and  $\tau = (\mu_I \overline{\phi} t)/L^2$  is the time scale normalized over a vertical span  $L$ . Consider (B1) first at a base case where  $\overline{\mathbf{u}} = [\overline{u} \ \overline{v} \ \overline{w}]^T = 0$ , then (B1) reduces to

$$\overline{q}_f \overline{\mathbf{E}} = \nabla \overline{p},\tag{B4}$$

by subtracting (B1) by (B4), we have

$$\left(\rho \frac{\partial}{\partial t} - \mu \nabla^2\right) \mathbf{u} = (q_f \mathbf{E})' - \nabla p' - \rho \Gamma z \frac{\partial \mathbf{u}}{\partial x} - \rho \Gamma w \mathbf{e}_x, \quad (\text{B5})$$

for incompressible flow, we can take divergence of (B5) and apply  $\nabla \cdot \mathbf{u} = 0$  to arrive

$$\nabla^2 p' = \nabla \cdot (q_f \mathbf{E})'. \quad (\text{B6})$$

Now consider the vertical component of (B5)

$$\left(\rho \frac{\partial}{\partial t} - \mu \nabla^2\right) w' = (q_f E_3)' - \frac{\partial}{\partial z} p' - \rho \Gamma z \frac{\partial w'}{\partial x}, \quad (\text{B7})$$

we can take the Laplacian of (B7) and incorporate (B6)

$$\left(\rho \frac{\partial}{\partial t} - \mu \nabla^2\right) \nabla^2 w' = \nabla^2 (q_f E_3)' - \frac{\partial}{\partial z} (\nabla \cdot (q_f \mathbf{E})') - \rho \Gamma \nabla^2 \left(z \frac{\partial w'}{\partial x}\right), \quad (\text{B8})$$

where

$$\begin{aligned} \frac{\partial}{\partial z} (\nabla \cdot (q_f \mathbf{E})') &= \frac{\partial}{\partial z} (\nabla \cdot (q_f' \bar{\mathbf{E}}) + \nabla \cdot (\bar{q}_f \mathbf{E}')) \\ &= \bar{E} \frac{\partial^2 q_f'}{\partial z^2} + \frac{d^2 \bar{q}_f}{dz^2} E_3' + \frac{d \bar{q}_f}{dz} \frac{\partial E_3'}{\partial z} + 2 \frac{q_f'}{\epsilon} \frac{d \bar{q}_f}{dz} + 3 \frac{\bar{q}_f}{\epsilon} \frac{\partial q_f'}{\partial z}, \end{aligned} \quad (\text{B9})$$

$$\begin{aligned} \nabla^2 (q_f E_3)' &= \nabla^2 (q_f' \bar{E}) + \nabla^2 (\bar{q}_f E_3') \\ &= \bar{E} \nabla^2 q_f' + \frac{q_f'}{\epsilon} \frac{d \bar{q}_f}{dz} + 2 \frac{d q_f'}{dz} \frac{d \bar{E}}{dz} + \bar{q}_f \nabla^2 E_3' + E_3' \frac{d^2 \bar{q}_f}{dz^2} + 2 \frac{d \bar{q}_f}{dz} \frac{d E_3'}{dz}, \end{aligned} \quad (\text{B10})$$

and

$$\rho \Gamma \nabla^2 \left(z \frac{\partial w'}{\partial x}\right) = \rho \Gamma z \frac{\partial}{\partial x} (\nabla^2 w') + 2 \rho \Gamma \frac{\partial}{\partial x} \left(\frac{\partial w'}{\partial z}\right). \quad (\text{B11})$$

Combining (B8)-(B11) gives

$$\begin{aligned} \left(\rho \frac{\partial}{\partial t} - \mu \nabla^2\right) \nabla^2 w' &= \frac{\partial^2 q_f'}{\partial z^2} \bar{E} - \frac{q_f'}{\epsilon} \frac{d \bar{q}_f}{dz} + \frac{\bar{q}_f}{\epsilon} \frac{\partial q_f'}{\partial z} + \bar{q}_f \nabla^2 E_3' + \frac{d \bar{q}_f}{dz} \frac{\partial E_3'}{\partial z} + \bar{E} \nabla^2 q_f' \\ &\quad - \rho \Gamma z \frac{\partial}{\partial x} (\nabla^2 w') - 2 \rho \Gamma \frac{\partial}{\partial x} \left(\frac{\partial w'}{\partial z}\right). \end{aligned} \quad (\text{B12})$$

Using  $q_f' = -\epsilon \nabla^2 \phi'$  and  $E_3' = -d\phi'/dz$ , (B12) can be rewritten in terms of potential

$$\left(\rho \frac{\partial}{\partial t} - \mu \nabla^2\right) \nabla^2 w' = \epsilon \bar{E} \nabla^2 \nabla_1^2 \phi' + \frac{d \bar{q}_f}{dz} \nabla_1^2 \phi' - \rho \Gamma z \frac{\partial}{\partial x} (\nabla^2 w') - 2 \rho \Gamma \frac{\partial}{\partial x} \left(\frac{\partial w'}{\partial z}\right), \quad (\text{B13})$$

where  $\nabla_1^2 = \partial^2/\partial x^2 + \partial^2/\partial y^2$ . Similarly (B2) can be linearized the same way in the vertical direction to obtain

$$-\epsilon \bar{E} \frac{\partial}{\partial z} \nabla^2 \phi' + \frac{d \bar{q}_f}{dz} \frac{\partial \phi'}{\partial z} + \frac{\epsilon}{\mu_I} \frac{\partial}{\partial t} \nabla^2 \phi' + 2 \bar{q}_f \nabla^2 \phi' = \frac{1}{\mu_I} \frac{d \bar{q}_f}{dz} w'. \quad (\text{B14})$$

Recall that  $\phi' = \phi_0(z) \exp(i\kappa_x x + i\kappa_y y) \exp(\omega\tau)$  and  $w' = w_0(z) \exp(i\kappa_x x + i\kappa_y y) \exp(\omega\tau)$ , so  $\nabla^2 \phi' = (-\kappa_x^2 + \kappa_y^2 + (d/dz)^2)\phi'$  and  $\nabla^2 w' = (-\kappa_x^2 + \kappa_y^2 + (d/dz)^2)w'$ , therefore the equations can be non-dimensionalized with  $\nabla^2 = (D^2 - \alpha^2)/L^2$ ,  $\nabla_1^2 = -\alpha^2/L^2$  where  $\alpha = L\sqrt{\kappa_x^2 + \kappa_y^2}$  and  $D = Ld/dz$ ,  $\partial/\partial t = \partial/\partial\tau (\mu_I \bar{\phi}/L^2) = \omega (\mu_I \bar{\phi}/L^2)$ ,  $\partial/\partial x = \beta/L$ ,  $z = \hat{z}L$ ,  $f(z) = w_0(z)L/(\mu_I \bar{\phi}(z))$ ,  $g(z) = \phi_0(z)/\bar{\phi}(z)$ . All the exponential terms eventually cancel out, and we arrive at

$$\begin{aligned} \left(\frac{\omega}{\text{Pr}} - (D^2 - \alpha^2)\right)(D^2 - \alpha^2)f &= -R \frac{d\bar{q}_f}{dz} \frac{L^3}{\epsilon \bar{\phi}} \alpha^2 g \\ -R \frac{\bar{E}L}{\bar{\phi}} (D^2 - \alpha^2) \alpha^2 g &- S \hat{z} \beta (D^2 - \alpha^2) f - 2S \beta D f, \end{aligned} \quad (\text{B15})$$

$$\begin{aligned} -\frac{\epsilon \bar{E}}{L^2} (D^2 - \alpha^2) D g &+ \frac{d\bar{q}_f}{dz} D g + \frac{\epsilon \bar{\phi}}{L^3} \omega (D^2 - \alpha^2) g \\ &+ \frac{2\bar{q}_f}{L} (D^2 - \alpha^2) g = \frac{d\bar{q}_f}{dz} f, \end{aligned} \quad (\text{B16})$$

where

$$R = \frac{\epsilon \bar{\phi}}{\mu \mu_I}, \quad \text{Pr} = \frac{\mu}{\rho \bar{\phi} \mu_I}, \quad S = \frac{\rho \Gamma L^2}{\mu}. \quad (\text{B17})$$

Equations (B15) and (B16) are linear equations of order four and three, respectively. Therefore, four boundary conditions for the initial velocity fluctuation,  $w_0(z)$ , and three boundary conditions for the initial potential fluctuation,  $\phi_0(z)$ , are required to obtain a unique solution. For the velocity fluctuation, we impose  $w_0(z_0) = \partial w_0(z_0)/\partial z = 0$  and  $w_0(z_0 + L) = \partial w_0(z_0 + L)/\partial z = 0$  based on no-slip boundary conditions and incompressibility of the fluid. For the potential fluctuation, it is assumed that the initial electric field fluctuations at the upper and lower boundaries are zero such that it is not violating Gauss's law, i.e.  $\partial \phi_0(z_0)/\partial z = \partial \phi_0(z_0 + L)/\partial z = 0$ . The last boundary condition is determined by recognizing that the electrical conductivity decreases significantly with altitude in the lower atmosphere [37], and therefore we assume the lower boundary has much larger electrical conductivity than the upper boundary such that relatively the initial potential fluctuation at the lower boundary is negligible, i.e.  $\phi_0(z_0) = 0$ .

The boundary conditions are then transformed into the final dimensionless form:  $f(z_0) = f'(z_0) = g(z_0) = g'(z_0) = 0$  for the lower boundary and  $f(z_0 + L) = f'(z_0 + L) = g'(z_0 + L) = 0$  for the upper boundary. To ensure numerical stability, we follow Appendix A in Schneider and Watson [38] to solve the ODEs numerically with these boundary conditions using a shooting method. We also observed that changing boundary conditions for the potential

(e.g., employing a Neumann condition for the electric field  $g''(z_0) = g''(z_0 + L) = 0$ ) has little effect on the critical Rayleigh number.

- 
- [1] B. A. Tinsley, R. P. Rohrbaugh, M. Hei, and K. V. Beard, “Effects of image charges on the scavenging of aerosol particles by cloud droplets and on droplet charging and possible ice nucleation processes,” *Journal of the Atmospheric Sciences* **57**, 2118–2134 (2000).
  - [2] F. Yu and R. P. Turco, “From molecular clusters to nanoparticles: Role of ambient ionization in tropospheric aerosol formation,” *Journal of Geophysical Research: Atmospheres* **106**, 4797–4814 (2001).
  - [3] S. N. Tripathi and R. G. Harrison, “Enhancement of contact nucleation by scavenging of charged aerosol particles,” *Atmospheric Research* **62**, 57–70 (2002).
  - [4] B. A. Tinsley, “Scavenging of condensation nuclei in clouds: dependence of sign of electroscavenging effect on droplet and CCN sizes,” in *Proceedings, International Conference on Clouds and Precipitation* (2004) p. 248.
  - [5] M. J. Rycroft, S. Israelsson, and C. Price, “The global atmospheric electric circuit, solar activity and climate change,” *Journal of Atmospheric and Solar-Terrestrial Physics* **62**, 1563–1576 (2000).
  - [6] D. Singh, V. Gopalakrishnan, R. P. Singh, A. K. Kamra, S. Singh, V. Pant, R. Singh, and A. K. Singh, “The atmospheric global electric circuit: an overview,” *Atmospheric Research* **84**, 91–110 (2007).
  - [7] W. A. Hoppel and S. G. Gathman, “Charge transport through an aerosol cloud,” *Journal of Applied Physics* **41**, 1971–1977 (1970).
  - [8] R. Bruinsma and S. Alexander, “Theory of electrohydrodynamic instabilities in electrolytic cells,” *The Journal of Chemical Physics* **92**, 3074–3085 (1990).
  - [9] G. Tomar, V. Shankar, A. Sharma, and G. Biswas, “Electrohydrodynamic instability of a confined viscoelastic liquid film,” *Journal of Non-newtonian Fluid Mechanics* **143**, 120–130 (2007).
  - [10] P. Atten, F. M. J. McCluskey, and A. C. Lahjomri, “The electrohydrodynamic origin of turbulence in electrostatic precipitators,” *IEEE Transactions on Industry Applications* , 705–711 (1987).



- [11] E. M. Lifschitz and L. P. Pitajewski, “Physical kinetics,” in *Textbook of Theoretical Physics*. 10 (1983).
- [12] A. V. Timofeev and B. N. Shvilkin, “Drift-dissipative instability of an inhomogeneous plasma in a magnetic field,” *Soviet Physics Uspekhi* **19**, 149 (1976).
- [13] A. A. Galeev, S. S. Moiseev, and R. Z. Sagdeev, “The theory of the stability of non-uniform plasma and anomalous diffusion,” *Journal of Nuclear Energy. Part C, Plasma Physics, Accelerators, Thermonuclear Research* **6**, 645 (1964).
- [14] M. Di Renzo and J. Urzay, “Aerodynamic generation of electric fields in turbulence laden with charged inertial particles,” *Nature Communications* **9** (2018).
- [15] Y. Yao and J. Capecelatro, “Competition between drag and Coulomb interactions in turbulent particle-laden flows using a coupled-fluid–Ewald-summation based approach,” *Physical Review Fluids* **3**, 034301 (2018).
- [16] A. U. Karnik and J. S. Shrimpton, “Mitigation of preferential concentration of small inertial particles in stationary isotropic turbulence using electrical and gravitational body forces,” *Physics of Fluids* **24**, 073301 (2012).
- [17] J. Lu, H. Nordsiek, E. W. Saw, and R. A. Shaw, “Clustering of charged inertial particles in turbulence,” *Physical Review Letters* **104**, 184505 (2010).
- [18] J. K. Eaton and J. R. Fessler, “Preferential concentration of particles by turbulence,” *International Journal of Multiphase Flow* **20**, 169–209 (1994).
- [19] S. Elghobashi and G. C. Truesdell, “Direct simulation of particle dispersion in a decaying isotropic turbulence,” *Journal of Fluid Mechanics* **242**, 655–700 (1992).
- [20] P. J. Ireland, A. D. Bragg, and L. R. Collins, “The effect of Reynolds number on inertial particle dynamics in isotropic turbulence. Part 1. Simulations without gravitational effects,” *Journal of Fluid Mechanics* **796**, 617–658 (2016).
- [21] J. P. L. C. Salazar, J. De Jong, L. Cao, S. H. Woodward, H. Meng, and L. R. Collins, “Experimental and numerical investigation of inertial particle clustering in isotropic turbulence,” *Journal of Fluid Mechanics* **600**, 245–256 (2008).
- [22] J. Chun, D. L. Koch, S. L. Rani, A. Ahluwalia, and L. R. Collins, “Clustering of aerosol particles in isotropic turbulence,” *Journal of Fluid Mechanics* **536**, 219–251 (2005).
- [23] M. R. Maxey, “The gravitational settling of aerosol particles in homogeneous turbulence and random flow fields,” *Journal of Fluid Mechanics* **174**, 441–465 (1987).

- [24] L. P. Wang and M. R. Maxey, “Settling velocity and concentration distribution of heavy particles in homogeneous isotropic turbulence,” *Journal of Fluid Mechanics* **256**, 27–68 (1993).
- [25] A. Aliseda, A. Cartellier, F. Hainaux, and J. C. Lasheras, “Effect of preferential concentration on the settling velocity of heavy particles in homogeneous isotropic turbulence,” *Journal of Fluid Mechanics* **468**, 77–105 (2002).
- [26] P. J. Ireland, A. D. Bragg, and L. R. Collins, “The effect of Reynolds number on inertial particle dynamics in isotropic turbulence. Part 2. Simulations with gravitational effects,” *Journal of Fluid Mechanics* **796**, 659–711 (2016).
- [27] K. A. Browning, T. W. Harrold, and J. R. Starr, “Richardson number limited shear zones in the free atmosphere,” *Quarterly Journal of the Royal Meteorological Society* **96**, 40–49 (1970).
- [28] A. A. Townsend, “Excitation of internal waves in a stably-stratified atmosphere with considerable wind-shear,” *Journal of Fluid Mechanics* **32**, 145–171 (1968).
- [29] K. H. Lloyd, C. H. Low, and R. A. Vincent, “Turbulence, billows and gravity waves in a high shear region of the upper atmosphere,” *Planetary and Space Science* **21**, 653–661 (1973).
- [30] P. Gualtieri, F. Picano, and C. M. Casciola, “Anisotropic clustering of inertial particles in homogeneous shear flow,” *Journal of Fluid Mechanics* **629**, 25–39 (2009).
- [31] C. Nicolai, B. Jacob, and R. Piva, “On the spatial distribution of small heavy particles in homogeneous shear turbulence,” *Physics of Fluids* **25**, 083301 (2013).
- [32] M. H. Kasbaoui, D. L. Koch, G. Subramanian, and O. Desjardins, “Preferential concentration driven instability of sheared gas–solid suspensions,” *Journal of Fluid Mechanics* **770**, 85–123 (2015).
- [33] R. G. Harrison and M. H. P. Ambaum, “Enhancement of cloud formation by droplet charging,” *Proceedings of the Royal Society of London A* **464**, 2561–2573 (2008).
- [34] L. Zhou and B. A. Tinsley, “Production of space charge at the boundaries of layer clouds,” *Journal of Geophysical Research: Atmospheres* **112** (2007).
- [35] J. M. Rosen, D. J. Hofmann, and W. Gringel, “Measurements of ion mobility to 30 km,” *Journal of Geophysical Research: Atmospheres* **90**, 5876–5884 (1985).
- [36] E. W. McDaniel and E. A. Mason, *The mobility and diffusion of ions in gases* (Wiley, 1973).
- [37] H. Volland, *Atmospheric electrodynamics*, Vol. 11 (Springer Science & Business Media, 2013).
- [38] J. M. Schneider and P. K. Watson, “Electrohydrodynamic stability of space-charge-limited currents in dielectric liquids. I. Theoretical study,” *Physics of Fluids* **13**, 1948–1954 (1970).

- [39] J. Wang, W. B. Rossow, and Y. Zhang, “Cloud vertical structure and its variations from a 20-yr global rawinsonde dataset,” *Journal of Climate* **13**, 3041–3056 (2000).
- [40] J. C. Wyngaard and O. R. Coté, “The budgets of turbulent kinetic energy and temperature variance in the atmospheric surface layer,” *Journal of the Atmospheric Sciences* **28**, 190–201 (1971).
- [41] L. Kelvin, “Stability of fluid motion: rectilinear motion of viscous fluid between two parallel plates,” *Philosophical Magazine* **24**, 188–196 (1887).
- [42] M. H. Kasbaoui, R. G. Patel, D. L. Koch, and O. Desjardins, “An algorithm for solving the Navier–Stokes equations with shear-periodic boundary conditions and its application to homogeneously sheared turbulence,” *Journal of Fluid Mechanics* **833**, 687–716 (2017).
- [43] L. N. Trefethen, A. E. Trefethen, S. C. Reddy, and T. A. Driscoll, “Hydrodynamic stability without eigenvalues,” *Science* **261**, 578–584 (1993).
- [44] E. R. Mansell and C. L. Ziegler, “Aerosol effects on simulated storm electrification and precipitation in a two-moment bulk microphysics model,” *Journal of the Atmospheric Sciences* **70**, 2032–2050 (2013).
- [45] O. Desjardins, G. Blanquart, G. Balarac, and H. Pitsch, “High order conservative finite difference scheme for variable density low Mach number turbulent flows,” *Journal of Computational Physics* **227**, 7125–7159 (2008).
- [46] J. Capecelatro and O. Desjardins, “An Euler–Lagrange strategy for simulating particle-laden flows,” *Journal of Computational Physics* **238**, 1–31 (2013).
- [47] L. Schiller and A. Naumann, “A drag coefficient correlation,” *Zeit. Ver. Deutsch. Ing.* **77**, 318–320 (1935).
- [48] S. B. Pope, *Turbulent flows* (Cambridge University Press, 2000).
- [49] J. Lowell and A. C. Rose-Innes, “Contact electrification,” *Advances in Physics* **29**, 947–1023 (1980).
- [50] S. Matsusaka, H. Maruyama, T. Matsuyama, and M. Ghadiri, “Triboelectric charging of powders: A review,” *Chemical Engineering Science* **65**, 5781–5807 (2010).
- [51] H. Grosshans and M. V. Papalexandris, “Direct numerical simulation of triboelectric charging in particle-laden turbulent channel flows,” *Journal of Fluid Mechanics* **818**, 465–491 (2017).
- [52] X. Jin and J. S. Marshall, “The role of fluid turbulence on contact electrification of suspended particles,” *Journal of Electrostatics* **87**, 217–227 (2017).

- [53] T. Gebhardt and S. Grossmann, “Chaos transition despite linear stability,” *Physical Review E* **50**, 3705 (1994).
- [54] A. Schmiegell and B. Eckhardt, “Fractal stability border in plane Couette flow,” *Physical Review Letters* **79**, 5250 (1997).
- [55] B. Gayen and M. Alam, “Algebraic and exponential instabilities in a sheared micropolar granular fluid,” *Journal of Fluid Mechanics* **567**, 195–233 (2006).
- [56] P. Sagaut and C. Cambon, *Homogeneous turbulence dynamics*, Vol. 10 (Springer, 2008).
- [57] M. Malik, J. Dey, and M. Alam, “Linear stability, transient energy growth, and the role of viscosity stratification in compressible plane Couette flow,” *Physical Review E* **77**, 036322 (2008).
- [58] K. M. Butler and B. F. Farrell, “Three-dimensional optimal perturbations in viscous shear flow,” *Physics of Fluids* **4**, 1637–1650 (1992).
- [59] P. J. Schmid, “Nonmodal stability theory,” *Annual Review of Fluid Mechanics* **39**, 129–162 (2007).
- [60] E. J. Hinch, *Perturbation methods* (Cambridge University Press, 1991).
- [61] C. M. Bender and S. A. Orszag, *Advanced mathematical methods for scientists and engineers I: Asymptotic methods and perturbation theory* (Springer Science, 2013).
- [62] M. Malik, M. Alam, and J. Dey, “Nonmodal energy growth and optimal perturbations in compressible plane Couette flow,” *Physics of Fluids* **18**, 034103 (2006).