

CHCRUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Phases of active matter composed of multicellular magnetotactic bacteria near a hard surface

Alexander Petroff, Alejandra Rosselli-Calderon, Ben Roque, and Pradeep Kumar Phys. Rev. Fluids **7**, 053102 — Published 27 May 2022 DOI: 10.1103/PhysRevFluids.7.053102

Phases of Active Matter Composed of Multicellular Magnetotactic Bacteria Near a Hard Surface

Alexander Petroff¹, Alejandra Rosselli-Calderon¹, Ben Roque¹, and Pradeep Kumar²

¹Department of Physics, Clark University, Worcester, MA 01610, USA

²Department of Physics, University of Arkansas, Fayetteville, AR 72701, USA

(Dated: May 10, 2022)

We investigate the motion of *Multicellular Magnetotactic Bacteria* (MMB) near a hard surface. Directing MMB towards a flat surface causes them the accumulate about the wall as an active gas. MMB are exponentially distributed about the surface with a penetration depth determined by the typical distance an MMB swims before it aligns with the ambient field. Increasing the magnetic field past a critical value, at which the penetration depth is of order the size of an MMB, causes the active gas to condense into a two-dimensional active fluid. Measurements of MMB motion in this phase reveal that contact with the surface enhances rotational diffusion by a factor of eight and that velocity fluctuations are Laplace distributed. The fluid undergoes a percolation transition at a critical magnetic field. Supercritical active fluids exhibit dynamic voids, which limit the relaxation of density fluctuations. We describe the possible ecological significance of these results.

I. INTRODUCTION

Understanding the strategies by which microbes navigate complex environments gives insight into their ecology and evolution and provides inspiration for engineers [1–4]. A variety of microbes living in sediment have independently evolved [5] the ability to navigate the labyrinthine pore spaces in which they live using the geomagnetic field [6]. Because, at high latitudes, Earth's magnetic field has a substantial radial component, swimming along magnetic field lines allows cells to position themselves in the continuously changing vertical chemical gradients typical of natural sediment [5]. However, if cells exclusively swim along field lines they are quickly trapped when they inevitably collide with an obstacle (e.g., a grain of sand). The strategies that allow magnetotactic cells to balance motion along field lines with obstacle avoidance remain poorly understood. To elucidate the physical basis of these strategies, we investigate the motion of one type of magnetotactic bacteria near a hard surface [7].

We focus on a group of δ -proteobacteria commonly called *Multicellular Magnetotactic Bacteria* (MMB), which have been assigned the provisional name *Candidatus* Magnetoglobus multicellularis [8]. These cells are the only known obligately multicellular bacteria [9]. These cells exclusively live in spherical colonies (see Fig. 1) composed of a monolayer of ~ 50 bacteria [10, 11]. 95% of colonies have diameters between 3.8 μ m and 7.3 μ m with and mean diameter of 5.3 μ m. Fewer that 1% of colonies, which are close of dividing (Fig. 1(e)), can reach diameters of 10.8 μ m. Cells in an MMB colony cooperate to push the colony along magnetic field lines with a typical speed of $U = 75 \,\mu$ m/s. Each cell in an MMB colony precipitates magnetic crystals, either magnetite [12] or greigite [13], which cause the colony to align its average magnetic moment with the ambient magnetic field. The outer surface of each cell is covered with about 30 flagella [14, 15]. The constituent cells exert an average force that is parallel to within a few degrees of the net magnetic moment [16]. The colonies swim through the fluid without tumbling, though a few colonies rarely ($\approx 1/60 \,\mathrm{s}^{-1}$) reverse their motion to briefly swim against the imposed magnetic field [17].

Two torques guide an MMB colony through the confined geometry of the pore space. The magnetic torque on a colony causes it to align with the ambient magnetic field *B* at a rate $\gamma = mB/8\pi\mu a^3$, where *m* is the magnetic moment, μ is the viscosity of water, and *a* is the colony radius. MMB, moving with speed *U*, swim a typical distance $\ell = 8\pi\mu a^3 U/mB$ as they align with the magnetic field. When a colony approaches to within a distance $\sim a$ of a surface, hydrodynamic interactions with the surface can turn the colony [18–20].

We tune the ratio of the length scales ℓ/a over which these two processes act by varying B. At low magnetic fields, for which $\ell/a \sim 30$, colonies quickly escape into the bulk fluid, where the magnetic field turn them back to towards the surface. This process of collision and alignment causes colonies to accumulate near a surface like an active gas. Increasing the magnetic field such that $\ell/a \sim 1$ causes this active gas to condense on the surface as an active two-dimensional fluid. Examining the dynamics of this active fluid reveals that contact with a surface increases the rotational diffusion coefficient by a factor of eight.

II. RESULTS

A. Enrichment of MMB

MMB have not been grown in pure culture. We enrich them from sediment taken from a shallow pool in a Massachusetts salt marsh (41°34′34.2″ N, 70°38′21.4″ W) between late Spring and early Fall and maintained in the lab. We lightly stir the sediment and place a neodymium magnet near the sediment surface [14, 21]. After twenty minutes, we collect 1 ml of water near the magnet surface. This material is placed in a vial, mixed lightly, and placed next to a magnet for an additional twenty minutes. By collecting the 0.1 ml of water near the magnet surface, we find thousands of MMB colonies, which represent the overwhelming majority of cells.

B. MMB Motion in the Bulk Fluid

We begin by characterizing the motion of free-swimming MMB in an applied magnetic field [17]. We inoculate MMB colonies into a rectangular microfluidic chamber ($1.5 \text{ cm} \times 1 \text{ cm} \times 0.015 \text{ cm}$), composed primarily of Polydimethylsiloxane (PDMS) and sealed on one side with a glass slide. We focus a microscope (Olympus X71, $20 \times$ objective) on a plane halfway between the top and bottom of the chamber. The entire experiment is performed inside a 3-axis Helmholtz coil (Walker Scientific, 76 cm coils), which creates a magnetic field **B** as large as $600 \,\mu\text{T}$ with $1 \,\mu\text{T}$ precision; the magnetic field varies over the field of view by less than 0.03%.

We initially apply a constant magnetic field of $256 \,\mu\text{T}$ and measure the angle $\theta(t)$ between the velocity U of each colony and B at time t. MMB swim with average speed $U = 75 \,\mu\text{m/s}$, although individual colonies show substantial variability in the range of $45 \,\mu\text{m/s}$ to $134 \,\mu\text{m/s}$ (95% confidence interval). Although their average direction of motion is parallel to the applied field, rotational diffusion causes their direction to slightly fluctuate with $\langle \theta^2 \rangle = 0.17 \,\text{rad}^2$. By switching the magnetic coils through



Figure 1. Multicellular Magnetotactic Bacteria live in spherical colonies composed of about 50 bacteria that grow and move like a single organism [22]. (a)–(f) As the cells in a colony divide, the colony grows, elongates, and divides symmetrically. The scale bar is 4 μ m. Panels (a)–(f) are modified slightly from Ref. [11] for legibility. Panel (g) shows the angle θ between the colony velocity U and the ambient magnetic field B averaged over several hundred colonies as they turn to align with the field. The average alignment rate γ is fit for each magnetic field. The solid black line is the solution of Eq. (1).

which current flows, we rotate the field by 90° and measure $\theta(t)$ as MMB align with the new field [see Fig. 1(g), supplemental video SV1, and supplemental Fig. S1 [23]].

To measure the magnetic moment m and rotational diffusion coefficient D_r of MMB colonies, we compare these trajectories to the two-dimensional motion of a colony that swims at constant speed parallel to its magnetic moment, which passively rotates to align with the ambient field [24, 25]. Combining this deterministic motion with rotational Brownian motion, gives the angular velocity

$$\frac{\mathrm{d}\theta}{\mathrm{d}t} = -\gamma\sin(\theta) + \sqrt{2D_r}\eta(t),\tag{1}$$

where η is white noise with $\langle \eta(t) \rangle = 0$ and $\langle \eta(t)\eta(t') \rangle = \delta(t-t')$. The colony swims with velocity

$$\mathbf{U} = U\left(\cos(\theta)\hat{x} + \sin(\theta)\hat{y}\right),\tag{2}$$

where \hat{x} and \hat{y} are unit vectors parallel and perpendicular to the applied field, respectively. As shown in Fig. 1(g), the measured trajectories of swimming MMB colonies are in good agreement with this model. At 256 μ T, we find $\gamma = 1.24 \pm 0.02 \text{ s}^{-1}$. The rotational diffusion coefficient $D_r = \langle \theta^2 \rangle \gamma = 0.212 \pm 0.003 \text{ rad}^2/\text{s}$ is measured from the trajectories before the field was rotated. Its value is roughly similar that of the eukaryotic alga *Chlamydomonas reinhardtii* (0.4 rad²/\text{s}) [18]. By fitting the trajectories of 128 individual MMB colonies, we find the natural variability of magnetic moments in the sample to range [26] from $3.9 \times 10^{-15} \text{ Am}^2$ to $10.9 \times 10^{-15} \text{ Am}^2$ (95% confidence interval).

C. An Active Gas of MMB Near a Surface

Next, we investigate how the motion of MMB changes as they are scattered from a surface. We apply a field to direct MMB to collide normal to the chamber wall. We begin by considering a null model of colony-surface interference. If the surface principally acts to cancel the normal component of the colony velocity, the rate k_e at which MMB escape is the rate at which rotational diffusion turns the colony away from the surface. Given a correlation time γ^{-1} of fluctuations, $k_e \approx \gamma \operatorname{erf_c}\left(\sqrt{\pi^2 \gamma/8D_r}\right)$. At a magnetic field of 0.29 mT, the expected rate of escape $k_e = 7 \times 10^{-5} \,\mathrm{s^{-1}}$. We therefore expect that, once a colony collides with a surface, it will remain trapped for several hours.

In contrast to this naïve estimate, when a colony collides with the chamber wall, it is only in contact for a fraction of a second before it escapes back into the bulk fluid [Fig. 2(a) and supplemental video SV2 [23]], where it realigns with the ambient field. The measured concentration n of colonies, normalized by the concentration at the surface n_0 , at a distance X from the wall [Fig. 2(b)] is consistent with an exponential decay. As we increase the magnitude of the magnetic field, the exponential profiles become more sharp as colonies are confined more closely to the wall. The penetration depth λ of colonies into the bulk fluid is proportional to the turning length ℓ ; we find $\lambda/\ell = 1.5 \pm 0.04$. We conclude that colonies scatter after a collision with a surface into the bulk fluid, where they realign with the ambient field and collide with the surface again.



Figure 2. MMB scatter from a surface. (a) A snapshot of an experiment (B = 0.29 mT) is shown. Red circles show positions of colonies near the chamber wall (black line). The blue curve shows the trajectory of a colony that collided with the wall. (b) Colonies are distributed exponentially about the wall. The distance X from the wall is normalized by ℓ . The black line shows the best fit exponential.

D. Formation of a Two-Dimensional Active Fluid

These results imply that at a critical magnetic field $B = 8\pi\mu a^2/m \sim 4$ mT, the turning radius ℓ shrinks to the size of a single MMB colony. At such a high magnetic field, we expect the active gas of free-swimming MMB colonies to condense on the surface as a two-dimensional phase of active matter.

We modify our experiments to visualize this state of active matter [see Fig. 3(a)]. To generate the relatively high magnetic fields required, we wrap a $40 \times$ microscope objective with 1000 turns of copper wire. We generate fields between 0.9 mT and 4.0 mT measured at the focal distance of the objective. We inoculate a chamber composed of a glass slide and coverslip separated by a 250 μ m plastic spacer (Gene Frame, ThermoFisher Scientific, USA). We focus the microscope to just below the height of the coverslip.

As we increase the magnetic field, we observe the emergence of a two-dimensional fluid phase [25, 27]. At high magnetic field (B > 2.0 mT), MMB colonies display roughly hexagonal order [28–31] [see Figs. 3(b) and 6 supplementary video[23] SV3] with many vacancies. We measure the concentration of colonies and extract their positions, velocities, and trajectories over the course of one minute. The concentration of colonies on the surface increases linearly with *B* to an area fraction of 0.61 at B = 3.2 mT, where it saturates. Removing the magnetic field causes colonies to rapidly escape to the bulk fluid, implying that the magnetic field produced by the MMB is insufficient to trap colonies [25].

E. Dynamics of Individual MMB in an Active Fluid

We begin our analysis by studying the motion of isolated colonies as they move over the two-dimensional glass surface. A colony may move laterally over the surface either because it exerts a force or it is advected by the flows generated by a nearby colony [29]. To disentangle these two components of motion we examine the trajectories of isolated colonies shortly before they collide. We find [Fig. 4(a)] that the the root-mean squared distance $\langle r^2 \rangle$ initially grows quadratically with time, consistent with ballistic motion, up to a time τ_B , after which $\langle r^2 \rangle$ grows linearly in time. The timescale for ballistic motion τ_B increases in proportion to γ^{-1} , indicating that colonies tumble for as long as their magnetic moment is misaligned from the ambient field. We conclude that the colonies move over the surface like non-interacting hard Brownian disks and that the influence of advection by neighboring colonies is imperceptible.

Fitting the effective two-dimensional diffusion coefficient D_{eff} to these trajectories [32], we find reasonably large values of D_{eff} [Fig. 4(b)], which decrease with the applied magnetic field. At the lowest magnetic field, $D_{\text{eff}} = 104 \pm 3 \,\mu\text{m}^2/\text{s}$.

Next, we use these measurements to infer the rotational diffusion coefficient of colonies near the surface. The in-plane diffusion coefficient $D_{\text{eff}} = \langle v_t^2 \rangle / 2\gamma$ [33], where v_t is the speed tangent to the chamber wall. From Eq. (2), colonies move laterally with velocity $v_t = U \sin(\theta)$. At high magnetic field, $\langle v_t^2 \rangle \approx U^2 \langle \theta^2 \rangle = U^2 D_r / \gamma$. Note that increasing the magnetic field decreases both the time scale for ballistic motion and the typical speed (see supplemental [23] Fig. S3). Combining these effects, $D_{\text{eff}} = U^2 D_r / 2\gamma^2$. Fitting the measured diffusion coefficients at various magnetic fields to this model [Fig. 4(b)], we find $D_r = 1.6 \text{ rad}^2/\text{s}$. We conclude that the rotational diffusion coefficient of colonies near a surface is eight times greater than that of freely-swimming colonies ($D_r = 0.212 \text{ rad}^2/\text{s}$).

To understand how the increase in rotational diffusion is reflected in the motion of colonies, we measure the distribution of instantaneous velocities v_t for all the colonies in the field of view, including those in contact with others. At the lowest magnetic



Figure 3. MMB are directed to collide normal to a coverslip by an applied magnetic field. (a) A schematic of the experiment is shown. The microscope is focused one colony radius below the top surface. (b) At the highest magnetic field examined (4 mT), colonies localize on the glass surface, where they diffuse laterally and collide with one another. The contact network is shown to highlight the local hexagonal order of the active fluid and the many vacancies, dislocations, and voids. The scale bar is $10 \,\mu$ m.

field examined, v_t is Laplace distributed [Fig. 5], in stark contrast to the Gaussian fluctuations predicted by Eq. (1) and observed in the motion of free-swimming colonies (see supplemental [23] Fig. S2). As the magnetic field increases, the distribution becomes Gaussian for small v_t , but the exponential tails remain.

This anomalous behavior can be understood as a consequence of the multicellular nature of the colonies. Equations (1) and (2) implicitly assume that the coordinated action of the cells in a colony can be approximated as a single point force. In reality, each cell exerts a force on the colony. Consider the effect of a single cell on the motion of the colony. By Eq. (1), an individual cell produces velocity fluctuations that are normally distributed $\mathcal{N}(v_t, w)$ with zero mean and variance w. As the component of the force exerted by different cells tangent to the wall varies with the position of the cell in the colony, the motion of the colony is a weighted sum over the forces exerted by each cell. Approximating the sum over the several tens of cells in a colony as an integral and assuming an exponential distribution $p(w) = e^{-w/2b^2}/2b^2$ of weights [34], the weighted sum

$$P(v_t) = \int_0^\infty \mathcal{N}(v_t, w) p(w) \mathrm{d}w = \frac{1}{2b} e^{-|v_t|/b}$$
(3)

is Laplace distributed.

While this analysis is promising, a model is needed to explain why the weighting function p(w) would be exponentially distributed. We speculate that collisions between the hundreds of flagella covering the colony and the surface could be dynamically important in reorienting the colony. The length of the flagella and the diameter of the colony may set the scale of the exponential decay.

F. Dynamics of the Active Fluid

We now provide a brief over view of the large-scale dynamics of the active fluid, which will be the focus of a future paper. As shown in Fig. 6(e), active fluids undergo a percolation transition [35]. This transition arises as the the concentration of MMB trapped on the surface increases with the applied magnetic field. From each frame of the recorded videos, we extract the center of each colony and reconstruct the contact network. To account for the polydisperse size distribution of colonies, we infer contacts between colonies that are separated by less than twice the average colony diameter. We measure the distribution in the number N of colonies in connected clusters. The variance in cluster size—averaged over the one minute experiment—increases sharply and exhibits a maximum at the critical magnetic field $B_c \approx 2.0 \text{ mT}$ [Fig. 6(e)]. At this magnetic field a spanning cluster emerges. The weight P_s of the spanning cluster increases with magnetic field. The packing fraction 0.46 of colonies at the critical magnetic field is substantially less than the value predicted [36] for hard disks (0.673), suggesting slight adhesion [37] or Motility-Induced phase separation [38] cause colonies to cluster.

We measure the pair-correlation function and structure factor of the active fluids at each magnetic field [39, 40]. The paircorrelation function g(r) is consistent with that of a fluid; as the magnetic field increases g(r) develops secondary and tertiary peaks [see Fig. 6(f)], which suggests that the active fluid becomes more structured with the increase in magnetic field. Moreover, we find that the structure factor $S(q) = \langle \rho(q)\rho(-q) \rangle / N$ —where $\rho(q)$ is the Fourier transform of the density and N is the number of colonies—is sharply peaked at low wave number q [see Fig. 6(g)], indicating that the fluid is compressible.

The most striking feature of the active fluids at the highest magnetic field is the persistence of large voids [Fig 6(d) and



Figure 4. At high magnetic field, MMB diffuse laterally over the chamber wall. (a) The measured distance **r** between isolated colonies at a time -t before they collide is consistent with the motion of hard Brownian disks; we fit the 2D diffusion coefficient D_{eff} at each magnetic field. The solid line is quadratic for $-t < \tau_{\text{B}}$ and then increases linearly with slope $16D_{\text{eff}}$ [32]. (b) D_{eff} decreases with increasing magnetic field. The red line shows the fit to $D_{\text{eff}} = \alpha B^{-2}$.



Figure 5. The probability density functions of tangential velocities of colonies are shown for a range of magnetic fields. The number of instantaneous measurements from which the distributions is calculated range from 6.5×10^5 (B = 0.9 mT) to 1.6×10^6 (B = 4.0 mT). Normal and Laplace distributions are shown as guides to the eye.

supplemental [23] video SV3]. Voids form continuously as colonies escape from the glass surface. These voids diffuse through the active fluid, coarsen as they merge, and diminish as escaped colonies collide with the coverslip. The average area of voids evolve as a random walk. The instantaneous changes in void areas are—like the velocity of individual colonies—Laplace distributed (see Fig. 7). A time series of the average void area shows a 1/f spectrum.

To characterize the effect of voids on the dynamics of the active fluid at the highest magnetic field, we measure the intermediate scattering function [39]

$$F(q,t) = \frac{\langle \rho(q,t)\rho(-q,0)\rangle}{\langle \rho(q,0)^2 \rangle}.$$
(4)

Fig. 6(h) shows F(q, t) for three values of q at the highest magnetic field. On scales similar to the size of a colony $(2\pi/q = 1.5 \text{ colony diameters})$, density fluctuations decay diffusively. The relaxation of density fluctuations on the scale of $2\pi/q = 23 \text{ colony diameters}$ reaches a plateau over the time scale of our measurements and the system is presumably arrested either by a jamming transition (area fraction $\phi = 0.6$) or possibly very slow glassy relaxation. At intermediate length scales $(2\pi/q = 11 \text{ colony diameters})$ similar to the size of voids, the density fluctuations appear to relax logarithmically. A similar logarithmic decay is observed for Brownian particles diffusing in an evolving pore space [41]. This similarity suggests that relaxation is limited by the time it takes large voids to form, through which colonies can diffuse. In future work, we plan to extend the time scale of the experiment to see if the slow relaxation is truly logarithmic or if it is an intermediate state in a two-step decay.



Figure 6. Colonies self organize into active fluids. Increasing the magnetic field increases the concentration of colonies but decreases their effective temperature. Representative snapshots show the geometry of the active fluids at magnetic fields (a) 0.9 mT, (b) 2.0 mT, (c) 2.8 mT, and (d) 4.0 mT. Blue curves and red curves show the boundaries of finite clusters and spanning clusters, respectively. Scale bars are $10 \mu \text{m}$. (e) The fluids undergo a percolation transition at a critical magnetic field $B_c \approx 2 \text{ mT}$. Near B_c , the spanning cluster rapidly appears and disappears as colonies reorganize. The distribution of cluster sizes N and weight of the spanning cluster are averaged in time. The (f) pair correlation function g(r) and (g) structure factor S(q) are consistent with a compressible fluid. (h) The intermediate scattering function is shown for B = 4.0 mT. All distances r and wave numbers q are measured relative to the average colony diameter.

III. DISCUSSION AND CONCLUSION

We have experimentally investigated the motion of MMB near a surface. At relatively low magnetic fields, MMB accumulate within a typical distance ℓ from a surface like an active gas. At a critical magnetic field, at which the typical distance ℓ a colony swims into the bulk fluid decreases to the diameter of a colony, the active gas condenses into an active two-dimensional fluid. Tracking individual colonies in the active fluid reveals that the rotational diffusion coefficient of the colonies increases by a factor of eight when they are in contact with a surface and velocity fluctuations are Laplace distributed rather than Normally distributed. As the applied magnetic field increases, the concentration of colonies in the active fluid increases, which leads to a percolation transition. Density fluctuations on the scale of the void space of the fluid appear to relax logarithmically in time.

We turn now to the possible ecological and evolutionary significance of these results. In the natural environment, where the ambient geomagnetic field is $B_{\text{geo}} = 51.9 \,\mu\text{T}$, the turning radius of the colony $\ell \approx 300 \,\mu\text{m} \gg a$. Consequently, colonies accumulate about obstructions as an active gas. This random motion could facilitate navigation through the labyrinthine pore space. MMB use the the geomagnetic field to navigate the vertical chemical gradients in the top millimeters sediment [5]. To successfully navigate through the pose space, MMB must balance motion along field lines with motion around obstacles. We define the scattering number $\text{Sc} = \gamma/k_{\text{sc}}$ as the ratio of the rate γ at which MMB align with the magnetic field to the rate k_{sc} at which colonies orient their motion in random directions. If $\text{Sc} \gg 1$, colonies move deterministically along field lines. Such colonies are unable to navigate the complex pore space because they become trapped as soon as they collide with a surface. If $\text{Sc} \ll 1$, colonies move diffusively through the sediment and cannot align with the ambient field. Thus, natural selection may tune the phenotype of colonies such that $\text{Sc} \sim 1$. Having found that MMB scatter upon collision with a surface, we estimate the scattering rate from the time U/ℓ_p require for an MMB to swim one pore diameter $\ell_p \approx 0.1$ mm. Indeed, in the local geomagnetic field B_{geo} , $\text{Sc} = mB_{\text{geo}}U/8\pi\mu a^3 \ell_p \approx 1/3$. This scaling can be rewritten to highlight the reflection of environment on the phenotype as

$$\frac{mU}{a^3} \sim \frac{8\pi\mu\ell_p}{B_{\rm geo}},\tag{5}$$

where the left hand side describes the phenotype of the colony, upon which natural selection can act, and the right hand side is determined by the environment.



Figure 7. The two-dimensional active fluid formed by colonies at the highest magnetic field examined (4 mT) shows a prevalence of large voids. The typical size of a void $\approx 200 \mu m^2$ is much larger than the cross-sectional area of a colony $40 \mu m^2$. Voids form as colonies escape the surface. The voids diffuse through the fluid and merge (see SV3.avi [23]). To characterize these dynamics we measure the average area $\langle A_{\text{void}} \rangle$ of voids that are entirely surrounded by colonies. (a) Like the velocity fluctuations of individual colonies, instantaneous fluctuations of void areas are Laplace distributed. The average change is zero, consistent with the persistence of voids. (b) The power spectrum of the time series shown in the first panel shows a 1/f dependence (red curve).

IV. ACKNOWLEDGMENTS

We would like to thank A. Libchaber, C. Modes, A. Kudrolli, H. Gould, and O. Devauchelle for their comments. We are grateful to J. Norton, M. Hanout, and C. Mitchell for their assistance. R. Hatzenpichler provided invaluable suggestions for the enrichment of MMB. This work was supported by the American Chemical Society (ACS PRF 60426-DNI2) and partially by the National Science Foundation (NSF PHY-2042150). Bruce Weiller provided additional funding for A.R.C.

V. APPENDIX

Tracking the motion of swimming colonies allows us to measure their magnetic moment and rotational diffusion coefficient and characterize the dynamics of active fluids. We track colonies in two experiments. In the first experiment, colonies swim in a microfluidic chamber. The applied magnetic field is orthogonal to the optical path used to visualize the colonies. In the second experiment, colonies are directed towards a coverslip. The optical path is parallel to the magnetic field. These differences require different approaches to identify the positions of colonies in each frame.

Colonies swimming in the microfluidic device appear as dark circles on a bright background (see videos [23] SV1.avi and SV2.avi). The large contrast difference between free-swimming colonies and the background makes tracking the colonies relatively simple. We first average all the frames from the video to find the background light intensity. Subtracting the background intensity from each frame highlights the positions of the colonies. We then apply an intensity threshold and extract the centroid of each dark disk. The trajectories of the centroids are extracted using the Matlab function "assignDetectionsToTracks."

In the second experiment, colonies move laterally over the chamber surface. In these experiments, the scattering of light through the individual cells that compose the colonies give each colony a textured appearance when it is in focus (see video [23] SV3.avi). Colonies swimming below the focal plane appear as bright disks with dark halos. The textured appearance of colonies and their close proximity to each other makes the extraction of their positions challenging. We identify the position of colonies in three steps. We first average all of the frames in the video to find the background illumination, which we subtract from each frame. Next, to highlight the texture of each colony, we measure the local standard deviation of the image; at each point in the frame we measure the standard deviation of pixel intensities in a neighborhood of one half of a colony radius. Finally, we identify circular components in the image of the local standard deviation by using the Matlab function "imfindcircles," which computes the circular Hough transform. We adjust the parameters in this procedure to eliminate false positive. Comparing the positions of colonies identified by this method to the original images, we find a rate of about 15% of false negatives and very few false positives. We use these instantaneous measurement of positions to calculate the contact network, pair correlation function, scattering function, and intermediate scattering function (see main text figure 6). We extract the trajectories of colonies from their positions using the Matlab function "assignDetectionsToTracks." Doing so allows us to calculate their instantaneous velocities and measure the distance between isolated colonies immediately before a collision (see main text figures 4 and 5).

To test the sensitivity of these results to the considerable number of false negatives, we developed a complementary method to identify connected clusters of colonies. Again, we begin by subtracting the background image from each frame to remove inhomogeneities in illumination. Next, we apply an intensity threshold to identify regions of the image that are occupied by colonies. While the colonies are textured, the average intensity is darker than the background. The resulting binary image includes all of the colonies in the image as well as the dark halos of colonies swimming below the focal plane. Next, we find the

connected components formed by colonies in contact with each other. To do so, we first dilate the binary image using a circular neighborhood of three pixels (about one tenth the colony radius). After dilating the image three times, we then erode the binary image three times using the same circular filter. This process of dilation and erosion fills in small holes and connects colonies that are close together. The perimeters of these clusters are shown in Fig. 6 and supplemental video [23] SV3. Computing the weight of the spanning cluster and the variance of cluster areas shows the same percolation transition at a critical magnetic field 2 mT as found from the reconstructed contact network.

- [1] Tom Fenchel, "Microbial behavior in a heterogeneous world," Science 296, 1068–1071 (2002).
- [2] HH Wensink, V Kantsler, RE Goldstein, and J Dunkel, "Controlling active self-assembly through broken particle-shape symmetry," Physical Review E 89, 010302 (2014).
- [3] Hugo Wioland, Francis G Woodhouse, Jörn Dunkel, and Raymond E Goldstein, "Ferromagnetic and antiferromagnetic order in bacterial vortex lattices," Nature physics 12, 341–345 (2016).
- [4] Jan Cammann, Fabian Jan Schwarzendahl, Tanya Ostapenko, Danylo Lavrentovich, Oliver Bäumchen, and Marco G Mazza, "Emergent probability fluxes in confined microbial navigation," Proceedings of the National Academy of Sciences 118 (2021).
- [5] Richard P Blakemore, "Magnetotactic bacteria," Annual Reviews in Microbiology 36, 217–238 (1982).
- [6] Nicolas Waisbord, Amin Dehkharghani, and Jeffrey S Guasto, "Fluidic bacterial diodes rectify magnetotactic cell motility in porous environments," Nature communications 12, 1–9 (2021).
- [7] Carolina N Keim, Daniel Mendes da Silva, Roger Duarte de Melo, Daniel Acosta-Avalos, Marcos Farina, and Henrique Lins de Barros, "Swimming behavior of the multicellular magnetotactic prokaryote *Candidatus* Magnetoglobus multicellularis near solid boundaries and natural magnetic grains," Antonie van Leeuwenhoek **114**, 1899–1913 (2021).
- [8] Fernanda Abreu, Juliana Lopes Martins, Thaís Souza Silveira, Carolina Neumann Keim, Henrique GP Lins de Barros, Frederico J Gueiros Filho, and Ulysses Lins, "*Candidatus* Magnetoglobus multicellularis, a multicellular, magnetotactic prokaryote from a hypersaline environment," International Journal of Systematic and Evolutionary Microbiology 57, 1318–1322 (2007).
- [9] Nicholas A Lyons and Roberto Kolter, "On the evolution of bacterial multicellularity," Current opinion in microbiology 24, 21–28 (2015).
- [10] Christopher T Lefèvre and Dennis A Bazylinski, "Ecology, diversity, and evolution of magnetotactic bacteria," Microbiol. Mol. Biol. Rev. 77, 497–526 (2013).
- [11] Carolina N Keim, Juliana L Martins, Fernanda Abreu, Alexandre Soares Rosado, Henrique Lins de Barros, Radovan Borojevic, Ulysses Lins, and Marcos Farina, "Multicellular life cycle of magnetotactic prokaryotes," FEMS Microbiology Letters 240, 203–208 (2004).
- [12] Ulysses Lins, Carolina N Keim, Flávia F Evans, Marcos Farina, and Peter R Buseck, "Magnetite (Fe₃O₄) and greigite (Fe₃S₄) crystals in multicellular magnetotactic prokaryotes," Geomicrobiology Journal **24**, 43–50 (2007).
- [13] Fernanda Abreu, Karen T Silva, Marcos Farina, Carolina N Keim, and Ulysses Lins, "Greigite magnetosome membrane ultrastructure in Candidatus Magnetoglobus multicellularis," Int Microbiol 11, 75–80 (2008).
- [14] Frank G Rodgers, Richard P Blakemore, Nancy A Blakemore, Richard B Frankel, Dennis A Bazylinski, Denise Maratea, and Christine Rodgers, "Intercellular structure in a many-celled magnetotactic prokaryote," Archives of Microbiology 154, 18–22 (1990).
- [15] Karen Tavares Silva, Fernanda Abreu, Fernando P Almeida, Carolina Neumann Keim, Marcos Farina, and Ulysses Lins, "Flagellar apparatus of south-seeking many-celled magnetotactic prokaryotes," Microscopy research and technique 70, 10–17 (2007).
- [16] Fernando P Almeida, Nathan B Viana, Ulysses Lins, Marcos Farina, and Carolina N Keim, "Swimming behaviour of the multicellular magnetotactic prokaryote candidatus magnetoglobus multicellularis under applied magnetic fields and ultraviolet light," Antonie Van Leeuwenhoek 103, 845–857 (2013).
- [17] Michael Greenberg, Karl Canter, Inga Mahler, and Adam Tornheim, "Observation of magnetoreceptive behavior in a multicellular magnetotactic prokaryote in higher than geomagnetic fields," Biophysical journal 88, 1496–1499 (2005).
- [18] Knut Drescher, Jörn Dunkel, Luis H Cisneros, Sujoy Ganguly, and Raymond E Goldstein, "Fluid dynamics and noise in bacterial cell-cell and cell-surface scattering," Proceedings of the National Academy of Sciences 108, 10940–10945 (2011).
- [19] Eric Lauga and Thomas R Powers, "The hydrodynamics of swimming microorganisms," Reports on Progress in Physics 72, 096601 (2009).
- [20] Enkeleida Lushi, Vasily Kantsler, and Raymond E Goldstein, "Scattering of biflagellate microswimmers from surfaces," Physical Review E 96, 023102 (2017).
- [21] Orr H Shapiro, Roland Hatzenpichler, Daniel H Buckley, Stephen H Zinder, and Victoria J Orphan, "Multicellular photo-magnetotactic bacteria," Environmental microbiology reports **3**, 233–238 (2011).
- [22] Carolina N Keim, Marcos Farina, and Ulysses Lins, "Magnetoglobus, magnetic aggregates in anaerobic environments," Microbe-American Society for Microbiology **2**, 437 (2007).
- [23] See Supplemental Material at [URL will be inserted by publisher] for three videos and three additional figures. Video SV1: MMB colonies turn in a rotated magnetic field. Video SV2: MMB colonies collide with and escape from a hard surface at three magnetic fields. Video SV3: Colonies self organize into active fluids. Figure S1: Trajectories of colonies as they align with an imposed magnetic field. Figure S2: The orientations of a free swimming colony about an imposed magnetic field are normally distributed. Figure S3: Instantaneous velocity fluctuations of colonies in an active decrease inversely with the magnitude of the applied magnetic field.
- [24] A Kalmijn, "Biophysics of geomagnetic field detection," IEEE Transactions on Magnetics 17, 1113–1124 (1981).
- [25] CJ Pierce, H Wijesinghe, E Mumper, BH Lower, SK Lower, and R Sooryakumar, "Hydrodynamic interactions, hidden order, and emergent collective behavior in an active bacterial suspension," Physical review letters **121**, 188001 (2018).
- [26] Marcelo Perantoni, Darci MS Esquivel, Eliane Wajnberg, Daniel Acosta-Avalos, Geraldo Cernicchiaro, and Henrique Lins de Barros,

"Magnetic properties of the microorganism Candidatus Magnetoglobus multicellularis," Naturwissenschaften 96, 685–690 (2009).

- [27] Xiao-Lun Wu and Albert Libchaber, "Particle diffusion in a quasi-two-dimensional bacterial bath," Physical review letters 84, 3017 (2000).
- [28] Ingmar H Riedel, Karsten Kruse, and Jonathon Howard, "A self-organized vortex array of hydrodynamically entrained sperm cells," Science 309, 300–303 (2005).
- [29] Alexander P Petroff, Xiao-Lun Wu, and Albert Libchaber, "Fast-moving bacteria self-organize into active two-dimensional crystals of rotating cells," Physical review letters 114, 158102 (2015).
- [30] Jeremie Palacci, Stefano Sacanna, Asher Preska Steinberg, David J Pine, and Paul M Chaikin, "Living crystals of light-activated colloidal surfers," Science 339, 936–940 (2013).
- [31] Tzer Han Tan, Alexander Mietke, Hugh Higinbotham, Junang Li, Yuchao Chen, Peter J Foster, Shreyas Gokhale, Jörn Dunkel, and Nikta Fakhri, "Development drives dynamics of living chiral crystals," arXiv preprint arXiv:2105.07507 (2021).
- [32] AP Petroff and A Libchaber, "Nucleation of rotating crystals by thiovulum majus bacteria," New Journal of Physics 20, 015007 (2018).
- [33] Howard C Berg, Random walks in biology (Princeton University Press, 2018).
- [34] Bo Wang, James Kuo, Sung Chul Bae, and Steve Granick, "When brownian diffusion is not gaussian," Nature materials **11**, 481–485 (2012).
- [35] Dietrich Stauffer and Ammon Aharony, Introduction to percolation theory (Taylor & Francis, 2018).
- [36] Stephan Mertens and Cristopher Moore, "Continuum percolation thresholds in two dimensions," Physical Review E 86, 061109 (2012).
- [37] Mark A Miller and Daan Frenkel, "Competition of percolation and phase separation in a fluid of adhesive hard spheres," Physical review letters **90**, 135702 (2003).
- [38] Michael E Cates and Julien Tailleur, "Motility-induced phase separation," Annu. Rev. Condens. Matter Phys. 6, 219–244 (2015).
- [39] Jean-Pierre Hansen and Ian R McDonald, Theory of simple liquids (Elsevier, 1990).
- [40] N de Macedo Biniossek, Hartmut Löwen, Th Voigtmann, and Frank Smallenburg, "Static structure of active brownian hard disks," Journal of Physics: Condensed Matter 30, 074001 (2018).
- [41] Tatjana Sentjabrskaja, Emanuela Zaccarelli, Cristiano De Michele, Francesco Sciortino, Piero Tartaglia, Thomas Voigtmann, Stefan U Egelhaaf, and Marco Laurati, "Anomalous dynamics of intruders in a crowded environment of mobile obstacles," Nature communications 7, 1–8 (2016).