

This is the accepted manuscript made available via CHORUS. The article has been published as:

Search for High Mass Resonances Decaying to Muon Pairs in $\sqrt{s}=1.96$ TeV $pp[\text{over } \gamma]$ Collisions

T. Aaltonen *et al.* (CDF Collaboration)

Phys. Rev. Lett. **106**, 121801 — Published 21 March 2011

DOI: [10.1103/PhysRevLett.106.121801](https://doi.org/10.1103/PhysRevLett.106.121801)

Search for High Mass Resonances Decaying to Muon Pairs in $\sqrt{s} = 1.96$ TeV $p\bar{p}$ Collisions

T. Aaltonen,²² B. Álvarez González^v,¹⁰ S. Amerio,⁴³ D. Amidei,³³ A. Anastassov,³⁸ A. Annovi,¹⁸ J. Antos,¹³ G. Apollinari,¹⁶ J.A. Appel,¹⁶ A. Apresyan,⁴⁸ T. Arisawa,⁵⁸ A. Artikov,¹⁴ J. Asaadi,⁵³ W. Ashmanskas,¹⁶ B. Auerbach,⁶¹ A. Aurisano,⁵³ F. Azfar,⁴² W. Badgett,¹⁶ A. Barbaro-Galtieri,²⁷ V.E. Barnes,⁴⁸ B.A. Barnett,²⁴ P. Barria^{cc},⁴⁶ P. Bartos,¹³ M. Bauce^{aa},⁴³ G. Bauer,³¹ F. Bedeschi,⁴⁶ D. Beecher,²⁹ S. Behari,²⁴ G. Bellettini^{bb},⁴⁶ J. Bellinger,⁶⁰ D. Benjamin,¹⁵ A. Beretvas,¹⁶ A. Bhatti,⁵⁰ M. Binkley*,¹⁶ D. Bisello^{aa},⁴³ I. Bizjak^{gg},²⁹ K.R. Bland,⁵ B. Blumenfeld,²⁴ A. Bocci,¹⁵ A. Bodek,⁴⁹ D. Bortoletto,⁴⁸ J. Boudreau,⁴⁷ A. Boveia,¹² B. Brau^a,¹⁶ L. Brigliadori^z,⁶ A. Brisuda,¹³ C. Bromberg,³⁴ E. Brucken,²² M. Bucciantonio^{bb},⁴⁶ J. Budagov,¹⁴ H.S. Budd,⁴⁹ S. Budd,²³ K. Burkett,¹⁶ G. Busetto^{aa},⁴³ P. Bussey,²⁰ A. Buzatu,³² C. Calancha,³⁰ S. Camarda,⁴ M. Campanelli,³⁴ M. Campbell,³³ F. Canelli¹²,¹⁶ A. Canepa,⁴⁵ B. Carls,²³ D. Carlsmith,⁶⁰ R. Carosi,⁴⁶ S. Carrillo^k,¹⁷ S. Carron,¹⁶ B. Casal,¹⁰ M. Casarsa,¹⁶ A. Castro^z,⁶ P. Catastini,¹⁶ D. Cauz,⁵⁴ V. Cavalieri^{cc},⁴⁶ M. Cavalli-Sforza,⁴ A. Cerri^f,²⁷ L. Cerrito^q,²⁹ Y.C. Chen,¹ M. Chertok,⁷ G. Chiarelli,⁴⁶ G. Chlachidze,¹⁶ F. Chlebana,¹⁶ K. Cho,²⁶ D. Chokheli,¹⁴ J.P. Chou,²¹ W.H. Chung,⁶⁰ Y.S. Chung,⁴⁹ C.I. Ciobanu,⁴⁴ M.A. Ciocci^{cc},⁴⁶ A. Clark,¹⁹ G. Compostella^{aa},⁴³ M.E. Convery,¹⁶ J. Conway,⁷ M. Corbo,⁴⁴ M. Cordelli,¹⁸ C.A. Cox,⁷ D.J. Cox,⁷ K. Cranmer,³⁷ F. Crescioli^{bb},⁴⁶ C. Cuena Almenar,⁶¹ J. Cuevas^v,¹⁰ R. Culbertson,¹⁶ D. Dagenhart,¹⁶ N. d'Ascenzo^t,⁴⁴ M. Datta,¹⁶ P. de Barbaro,⁴⁹ S. De Cecco,⁵¹ G. De Lorenzo,⁴ M. Dell'Orso^{bb},⁴⁶ C. Deluca,⁴ L. Demortier,⁵⁰ J. Deng^c,¹⁵ M. Deninno,⁶ F. Devoto,²² M. d'Errico^{aa},⁴³ A. Di Canto^{bb},⁴⁶ B. Di Ruzza,⁴⁶ J.R. Dittmann,⁵ M. D'Onofrio,²⁸ S. Donati^{bb},⁴⁶ P. Dong,¹⁶ M. Dorigo,⁵⁴ T. Dorigo,⁴³ K. Ebina,⁵⁸ A. Elagin,⁵³ A. Eppig,³³ R. Erbacher,⁷ D. Errede,²³ S. Errede,²³ N. Ershaidat^y,⁴⁴ R. Eusebi,⁵³ H.C. Fang,²⁷ S. Farrington,⁴² M. Feindt,²⁵ J.P. Fernandez,³⁰ C. Ferrazza^{dd},⁴⁶ R. Field,¹⁷ G. Flanagan^r,⁴⁸ R. Forrest,⁷ M.J. Frank,⁵ M. Franklin,²¹ J.C. Freeman,¹⁶ Y. Funakoshi,⁵⁸ I. Furic,¹⁷ M. Gallinaro,⁵⁰ J. Galyardt,¹¹ J.E. Garcia,¹⁹ A.F. Garfinkel,⁴⁸ P. Garosi^{cc},⁴⁶ H. Gerberich,²³ E. Gerchtein,¹⁶ S. Giagu^{ee},⁵¹ V. Giakoumopoulou,³ P. Giannetti,⁴⁶ K. Gibson,⁴⁷ C.M. Ginsburg,¹⁶ N. Giokaris,³ P. Giromini,¹⁸ M. Giunta,⁴⁶ G. Giurgiu,²⁴ V. Glagolev,¹⁴ D. Glenzinski,¹⁶ M. Gold,³⁶ D. Goldin,⁵³ N. Goldschmidt,¹⁷ A. Golossanov,¹⁶ G. Gomez,¹⁰ G. Gomez-Ceballos,³¹ M. Goncharov,³¹ O. González,³⁰ I. Gorelov,³⁶ A.T. Goshaw,¹⁵ K. Goulianatos,⁵⁰ A. Gresele,⁴³ S. Grinstein,⁴ C. Grossi-Pilcher,¹² R.C. Group,⁵⁷ J. Guimaraes da Costa,²¹ Z. Gunay-Unalan,³⁴ C. Haber,²⁷ S.R. Hahn,¹⁶ E. Halkiadakis,⁵² A. Hamaguchi,⁴¹ J.Y. Han,⁴⁹ F. Happacher,¹⁸ K. Hara,⁵⁵ D. Hare,⁵² M. Hare,⁵⁶ R.F. Harr,⁵⁹ K. Hatakeyama,⁵ C. Hays,⁴² M. Heck,²⁵ J. Heinrich,⁴⁵ M. Herndon,⁶⁰ S. Hewamanage,⁵ D. Hidas,⁵² A. Hocker,¹⁶ W. Hopkins^g,¹⁶ D. Horn,²⁵ S. Hou,¹ R.E. Hughes,³⁹ M. Hurwitz,¹² U. Husemann,⁶¹ N. Hussain,³² M. Hussein,³⁴ J. Huston,³⁴ G. Introzzi,⁴⁶ M. Iori^{ee},⁵¹ A. Ivanov^o,⁷ E. James,¹⁶ D. Jang,¹¹ B. Jayatilaka,¹⁵ E.J. Jeon,²⁶ M.K. Jha,⁶ S. Jindariani,¹⁶ W. Johnson,⁷ M. Jones,⁴⁸ K.K. Joo,²⁶ S.Y. Jun,¹¹ T.R. Junk,¹⁶ T. Kamon,⁵³ P.E. Karchin,⁵⁹ Y. Katoⁿ,⁴¹ W. Ketchum,¹² J. Keung,⁴⁵ V. Khotilovich,⁵³ B. Kilminster,¹⁶ D.H. Kim,²⁶ H.S. Kim,²⁶ H.W. Kim,²⁶ J.E. Kim,²⁶ M.J. Kim,¹⁸ S.B. Kim,²⁶ S.H. Kim,⁵⁵ Y.K. Kim,¹² N. Kimura,⁵⁸ M. Kirby,¹⁶ S. Klimenko,¹⁷ K. Kondo,⁵⁸ D.J. Kong,²⁶ J. Konigsberg,¹⁷ A.V. Kotwal,¹⁵ M. Kreps,²⁵ J. Kroll,⁴⁵ D. Krop,¹² N. Krumbach^l,⁵ M. Kruse,¹⁵ V. Krutelyov^d,⁵³ T. Kuhr,²⁵ M. Kurata,⁵⁵ S. Kwang,¹² A.T. Laasanen,⁴⁸ S. Lami,⁴⁶ S. Lammel,¹⁶ M. Lancaster,²⁹ R.L. Lander,⁷ K. Lannon^u,³⁹ A. Lath,⁵² G. Latino^{cc},⁴⁶ I. Lazzizzera,⁴³ T. LeCompte,² E. Lee,⁵³ H.S. Lee,¹² J.S. Lee,²⁶ S.W. Lee^w,⁵³ S. Leo^{bb},⁴⁶ S. Leone,⁴⁶ J.D. Lewis,¹⁶ C.-J. Lin,²⁷ J. Linacre,⁴² M. Lindgren,¹⁶ E. Lipeles,⁴⁵ A. Lister,¹⁹ D.O. Litvintsev,¹⁶ C. Liu,⁴⁷ Q. Liu,⁴⁸ T. Liu,¹⁶ S. Lockwitz,⁶¹ N.S. Lockyer,⁴⁵ A. Loginov,⁶¹ D. Lucchesi^{aa},⁴³ J. Lueck,²⁵ P. Lujan,²⁷ P. Lukens,¹⁶ G. Lungu,⁵⁰ J. Lys,²⁷ R. Lysak,¹³ R. Madrak,¹⁶ K. Maeshima,¹⁶ K. Makhoul,³¹ P. Maksimovic,²⁴ S. Malik,⁵⁰ G. Manca^b,²⁸ A. Manousakis-Katsikakis,³ F. Margaroli,⁴⁸ C. Marino,²⁵ M. Martínez,⁴ R. Martínez-Ballarín,³⁰ P. Mastrandrea,⁵¹ M. Mathis,²⁴ M.E. Mattson,⁵⁹ P. Mazzanti,⁶ K.S. McFarland,⁴⁹ P. McIntyre,⁵³ R. McNultyⁱ,²⁸ A. Mehta,²⁸ P. Mehtala,²² A. Menzione,⁴⁶ C. Mesropian,⁵⁰ T. Miao,¹⁶ D. Mietlicki,³³ A. Mitra,¹ H. Miyake,⁵⁵ S. Moed,²¹ N. Moggi,⁶ M.N. Mondragon^k,¹⁶ C.S. Moon,²⁶ R. Moore,¹⁶ M.J. Morello,¹⁶ J. Morlock,²⁵ P. Movilla Fernandez,¹⁶ A. Mukherjee,¹⁶ Th. Muller,²⁵ P. Murat,¹⁶ M. Mussini^z,⁶ J. Nachtman^m,¹⁶ Y. Nagai,⁵⁵ J. Naganoma,⁵⁸ I. Nakano,⁴⁰ A. Napier,⁵⁶ J. Nett,⁵³ C. Neu,⁵⁷ M.S. Neubauer,²³ J. Nielsen^e,²⁷ L. Nodulman,² O. Norniella,²³ E. Nurse,²⁹ L. Oakes,⁴² S.H. Oh,¹⁵ Y.D. Oh,²⁶ I. Oksuzian,⁵⁷ T. Okusawa,⁴¹ R. Orava,²² L. Ortolan,⁴ S. Pagan Griso^{aa},⁴³ C. Pagliarone,⁵⁴ E. Palencia^f,¹⁰ V. Papadimitriou,¹⁶ A.A. Paramonov,² J. Patrick,¹⁶ G. Pauletta^{ff},⁵⁴ M. Paulini,¹¹ C. Paus,³¹ D.E. Pellett,⁷ A. Penzo,⁵⁴ T.J. Phillips,¹⁵ G. Piacentino,⁴⁶ E. Pianori,⁴⁵ J. Pilot,³⁹ K. Pitts,²³ C. Plager,⁹ L. Pondrom,⁶⁰ K. Potamianos,⁴⁸ O. Poukhov*,¹⁴

F. Prokoshin^{x, 14} A. Pronko,¹⁶ F. Ptohos^{h, 18} E. Pueschel,¹¹ G. Punzi^{bb, 46} J. Pursley,⁶⁰ E. Quinlan,⁸ A. Rahaman,⁴⁷ V. Ramakrishnan,⁶⁰ N. Ranjan,⁴⁸ I. Redondo,³⁰ P. Renton,⁴² M. Rescigno,⁵¹ F. Rimondi^{z, 6} L. Ristori^{45, 16} A. Robson,²⁰ T. Rodrigo,¹⁰ T. Rodriguez,⁴⁵ E. Rogers,²³ S. Rolli,⁵⁶ R. Roser,¹⁶ M. Rossi,⁵⁴ F. Rubbo,¹⁶ F. Ruffini^{cc, 46} A. Ruiz,¹⁰ J. Russ,¹¹ V. Rusu,¹⁶ A. Safonov,⁵³ W.K. Sakamoto,⁴⁹ Y. Sakurai,⁵⁸ L. Santiff,⁵⁴ L. Sartori,⁴⁶ K. Sato,⁵⁵ V. Saveliev^{t, 44} A. Savoy-Navarro,⁴⁴ P. Schlabach,¹⁶ A. Schmidt,²⁵ E.E. Schmidt,¹⁶ M.P. Schmidt*,⁶¹ M. Schmitt,³⁸ T. Schwarz,⁷ L. Scodellaro,¹⁰ A. Scribano^{cc, 46} F. Scuri,⁴⁶ A. Sedov,⁴⁸ S. Seidel,³⁶ Y. Seiya,⁴¹ A. Semenov,¹⁴ F. Sforza^{bb, 46} A. Sfyrla,²³ S.Z. Shalhout,⁷ T. Shears,²⁸ P.F. Shepard,⁴⁷ M. Shimojima^{s, 55} S. Shiraishi,¹² M. Shochet,¹² I. Shreyber,³⁵ A. Simonenko,¹⁴ P. Sinervo,³² A. Sissakian*,¹⁴ K. Sliwa,⁵⁶ J.R. Smith,⁷ F.D. Snider,¹⁶ A. Soha,¹⁶ S. Somalwar,⁵² V. Sorin,⁴ P. Squillaciotti,¹⁶ M. Stancari,¹⁶ M. Stanitzki,⁶¹ R. St. Denis,²⁰ B. Stelzer,³² O. Stelzer-Chilton,³² D. Stentz,³⁸ J. Strologas,³⁶ G.L. Strycker,³³ Y. Sudo,⁵⁵ A. Sukhanov,¹⁷ I. Suslov,¹⁴ K. Takemasa,⁵⁵ Y. Takeuchi,⁵⁵ J. Tang,¹² M. Tecchio,³³ P.K. Teng,¹ J. Thom^{g, 16} J. Thome,¹¹ G.A. Thompson,²³ E. Thomson,⁴⁵ P. Ttito-Guzmán,³⁰ S. Tkaczyk,¹⁶ D. Toback,⁵³ S. Tokar,¹³ K. Tollefson,³⁴ T. Tomura,⁵⁵ D. Tonelli,¹⁶ S. Torre,¹⁸ D. Torretta,¹⁶ P. Totaro^{ff, 54} M. Trovato^{dd, 46} Y. Tu,⁴⁵ F. Ukegawa,⁵⁵ S. Uozumi,²⁶ A. Varganov,³³ F. Vázquez^{k, 17} G. Velev,¹⁶ C. Vellidis,³ M. Vidal,³⁰ I. Vila,¹⁰ R. Vilar,¹⁰ J. Vizán,¹⁰ M. Vogel,³⁶ G. Volpi^{bb, 46} P. Wagner,⁴⁵ R.L. Wagner,¹⁶ T. Wakisaka,⁴¹ R. Wallny,⁹ S.M. Wang,¹ A. Warburton,³² D. Waters,²⁹ M. Weinberger,⁵³ W.C. Wester III,¹⁶ B. Whitehouse,⁵⁶ D. Whiteson,⁸ A.B. Wicklund,² E. Wicklund,¹⁶ S. Wilbur,¹² F. Wick,²⁵ H.H. Williams,⁴⁵ J.S. Wilson,³⁹ P. Wilson,¹⁶ B.L. Winer,³⁹ P. Wittich^{g, 16} S. Wolbers,¹⁶ H. Wolfe,³⁹ T. Wright,³³ X. Wu,¹⁹ Z. Wu,⁵ K. Yamamoto,⁴¹ J. Yamaoka,¹⁵ T. Yang,¹⁶ U.K. Yang^{p, 12} Y.C. Yang,²⁶ W.-M. Yao,²⁷ G.P. Yeh,¹⁶ K. Yi^{m, 16} J. Yoh,¹⁶ K. Yorita,⁵⁸ T. Yoshida^{j, 41} G.B. Yu,¹⁵ I. Yu,²⁶ S.S. Yu,¹⁶ J.C. Yun,¹⁶ A. Zanetti,⁵⁴ Y. Zeng,¹⁵ and S. Zucchelli^{z, 6}

(CDF Collaboration[†])

¹Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China

²Argonne National Laboratory, Argonne, Illinois 60439, USA

³University of Athens, 157 71 Athens, Greece

⁴Institut de Fisica d'Altes Energies, ICREA, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain

⁵Baylor University, Waco, Texas 76798, USA

⁶Istituto Nazionale di Fisica Nucleare Bologna, ^zUniversity of Bologna, I-40127 Bologna, Italy

⁷University of California, Davis, Davis, California 95616, USA

⁸University of California, Irvine, Irvine, California 92697, USA

⁹University of California, Los Angeles, Los Angeles, California 90024, USA

¹⁰Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain

¹¹Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

¹²Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA

¹³Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia

¹⁴Joint Institute for Nuclear Research, RU-141980 Dubna, Russia

¹⁵Duke University, Durham, North Carolina 27708, USA

¹⁶Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

¹⁷University of Florida, Gainesville, Florida 32611, USA

¹⁸Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy

¹⁹University of Geneva, CH-1211 Geneva 4, Switzerland

²⁰Glasgow University, Glasgow G12 8QQ, United Kingdom

²¹Harvard University, Cambridge, Massachusetts 02138, USA

²²Division of High Energy Physics, Department of Physics,

University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland

²³University of Illinois, Urbana, Illinois 61801, USA

²⁴The Johns Hopkins University, Baltimore, Maryland 21218, USA

²⁵Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany

²⁶Center for High Energy Physics: Kyungpook National University,

Daegu 702-701, Korea; Seoul National University, Seoul 151-742,

Korea; Sungkyunkwan University, Suwon 440-746,

Korea; Korea Institute of Science and Technology Information,

Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757,

Korea; Chonbuk National University, Jeonju 561-756, Korea

²⁷Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

²⁸University of Liverpool, Liverpool L69 7ZE, United Kingdom

²⁹University College London, London WC1E 6BT, United Kingdom

³⁰Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain

³¹Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

- ³²Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto, Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
- ³³University of Michigan, Ann Arbor, Michigan 48109, USA
- ³⁴Michigan State University, East Lansing, Michigan 48824, USA
- ³⁵Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- ³⁶University of New Mexico, Albuquerque, New Mexico 87131, USA
- ³⁷New York University, New York, New York 10003, USA
- ³⁸Northwestern University, Evanston, Illinois 60208, USA
- ³⁹The Ohio State University, Columbus, Ohio 43210, USA
- ⁴⁰Okayama University, Okayama 700-8530, Japan
- ⁴¹Osaka City University, Osaka 588, Japan
- ⁴²University of Oxford, Oxford OX1 3RH, United Kingdom
- ⁴³Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, ^{aa}University of Padova, I-35131 Padova, Italy
- ⁴⁴LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
- ⁴⁵University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁴⁶Istituto Nazionale di Fisica Nucleare Pisa, ^{bb}University of Pisa, ^{cc}University of Siena and ^{dd}Scuola Normale Superiore, I-56127 Pisa, Italy
- ⁴⁷University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
- ⁴⁸Purdue University, West Lafayette, Indiana 47907, USA
- ⁴⁹University of Rochester, Rochester, New York 14627, USA
- ⁵⁰The Rockefeller University, New York, New York 10065, USA
- ⁵¹Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, ^{ee}Sapienza Università di Roma, I-00185 Roma, Italy
- ⁵²Rutgers University, Piscataway, New Jersey 08855, USA
- ⁵³Texas A&M University, College Station, Texas 77843, USA
- ⁵⁴Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, ^{ff}University of Trieste/Udine, I-33100 Udine, Italy
- ⁵⁵University of Tsukuba, Tsukuba, Ibaraki 305, Japan
- ⁵⁶Tufts University, Medford, Massachusetts 02155, USA
- ⁵⁷University of Virginia, Charlottesville, VA 22906, USA
- ⁵⁸Waseda University, Tokyo 169, Japan
- ⁵⁹Wayne State University, Detroit, Michigan 48201, USA
- ⁶⁰University of Wisconsin, Madison, Wisconsin 53706, USA
- ⁶¹Yale University, New Haven, Connecticut 06520, USA

We present a search for a new narrow, spin-1, high mass resonance decaying to $\mu^+\mu^- + X$, using a matrix element based likelihood and a simultaneous measurement of the resonance mass and production rate. In data with 4.6 fb^{-1} of integrated luminosity collected by the CDF detector in $p\bar{p}$ collisions at $\sqrt{s} = 1960 \text{ GeV}$, the most likely signal cross section is consistent with zero at 16% confidence level. We therefore do not observe evidence for a high mass resonance, and place limits on models predicting spin-1 resonances, including $M > 1071 \text{ GeV}/c^2$ at 95% confidence level for a Z' boson with the same couplings to fermions as the Z boson.

PACS numbers: 12.60.-i, 13.85.Rm, 14.80.-j

*Deceased

[†]With visitors from ^aUniversity of Massachusetts Amherst, Amherst, Massachusetts 01003, ^bIstituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, ^cUniversity of California Irvine, Irvine, CA 92697, ^dUniversity of California Santa Barbara, Santa Barbara, CA 93106 ^eUniversity of California Santa Cruz, Santa Cruz, CA 95064, ^fCERN, CH-1211 Geneva, Switzerland, ^gCornell University, Ithaca, NY 14853, ^hUniversity of Cyprus, Nicosia CY-1678, Cyprus, ⁱUniversity College Dublin, Dublin 4, Ireland, ^jUniversity of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017, ^kUniversidad Iberoamericana, Mexico D.F., Mexico, ^lIowa State University, Ames, IA 50011, ^mUniversity of Iowa, Iowa City, IA 52242, ⁿKinki University, Higashi-Osaka City, Japan 577-8502, ^oKansas State University,

We report a search for a narrow spin-1 resonance (Z') with decays to muon pairs and a mass between 130 GeV/c^2 and $\approx 1 \text{ TeV}/c^2$. Such a resonance is pre-

Manhattan, KS 66506, ^pUniversity of Manchester, Manchester M13 9PL, England, ^qQueen Mary, University of London, London, E1 4NS, England, ^rMuons, Inc., Batavia, IL 60510, ^sNagasaki Institute of Applied Science, Nagasaki, Japan, ^tNational Research Nuclear University, Moscow, Russia, ^uUniversity of Notre Dame, Notre Dame, IN 46556, ^vUniversidad de Oviedo, E-33007 Oviedo, Spain, ^wTexas Tech University, Lubbock, TX 79609, ^xUniversidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile, ^yYarmouk University, Irbid 211-63, Jordan, ^{gg}On leave from J. Stefan Institute, Ljubljana, Slovenia,

dicted generically in models with additional gauge groups [1], a feature of many extensions to the standard model of particle physics (SM). Typical examples are Little Higgs models [2] and the next-to-minimal supersymmetric model [3]; models can also be framed generically to cover a large set of potential signals [4].

Current 95% confidence level (CL) lower limits on the mass of a Z' boson with the same couplings to fermions as the Z boson (Z'_{SM}) are 1030 GeV/c^2 from a search of CDF dimuon data with 2.3 fb^{-1} of integrated luminosity [5], and 1023 GeV/c^2 [6] and 963 GeV/c^2 [7] from respective searches in dielectron data from D0 in 5.4 fb^{-1} and from CDF in 2.5 fb^{-1} , respectively.

This Letter reports a new search of the CDF dimuon data, superceding Ref. [5], with several significant enhancements: twice the integrated luminosity, a matrix element based likelihood providing an approximately 20% relative increase in cross section sensitivity at large Z' mass, and a new statistical approach designed to maximize simultaneously both discovery potential and mass exclusion limits in searches for new physics. This new approach directly fits the data to a single Z' boson hypothesis in the plane of boson mass and signal cross section, and can be applied to any search for a hypothetical new particle of unknown mass.

We use a data sample corresponding to an integrated luminosity of 4.6 fb^{-1} , collected with the CDF II detector [8], a general purpose detector designed to study $p\bar{p}$ collisions at the Fermilab Tevatron. The tracking system consists of a cylindrical open-cell drift chamber and silicon microstrip detectors in a 1.4 T magnetic field parallel to the beam axis. The silicon detectors provide tracking information for pseudorapidity $|\eta| < 2$ [9] and are used to reconstruct collision and decay points. The drift chamber surrounds the silicon detectors and gives full coverage in the central pseudorapidity region ($|\eta| < 1$). For the muon kinematics relevant to this search, the tracker provides an invariant mass resolution of $\delta M/M^2 \approx 17\%/\text{TeV}/c^2$. Electromagnetic and hadronic calorimeters surrounding the tracking system measure energy from particle showers and minimum ionizing particles such as muons. Drift chambers and scintillators located outside the calorimeters detect muons in the central pseudorapidity region $|\eta| < 1$.

Events used for this search are selected online with a trigger requiring a muon candidate with $p_T > 18 \text{ GeV}/c$. The event selection is unchanged from the previous search [5], requiring at least two oppositely charged muons with $p_T > 30 \text{ GeV}/c$ and no identified cosmic rays [10]. Each muon is required to have calorimeter deposits consistent with a minimum ionizing particle and have a track fully within the fiducial volume of the drift chamber. At least one muon must have an associated muon-chamber track.

The dominant and irreducible background is standard model Z/γ^* production with subsequent decay to muons.

We model this background using events generated at leading order by PYTHIA [11] with the CTEQ6L [12] parton distribution functions (PDF). Events are reweighted by the ratio of next-to-next-to-leading order to leading-order cross sections as a function of dimuon mass [4]. We use the standard CDF simulation [13] to model the detector response to the particles in the event. The overall normalization of this background is derived from data near the Z boson pole mass, $70 < M_{\mu\mu} < 110 \text{ GeV}/c^2$.

The remaining background contributions are small relative to standard model sources of dimuons. Decays of WW (1%) and $t\bar{t}$ (1%) to dimuons are described using PYTHIA and the CDF detector simulation, normalized to NLO cross sections [14]. The background from objects misidentified as muons (0.5%) is estimated using the distribution of calorimeter energy in the vicinity of each candidate muon in the sample [15]. Cosmic ray backgrounds ($< 0.01\%$) are modeled using identified cosmic ray events, weighted by the probability to survive the cosmic ray removal algorithm [10].

The observed spectrum of dimuon invariant mass ($M_{\mu\mu}$) is in good agreement with the total SM plus cosmic-ray prediction, as shown in Fig. 1. We expect 1851 ± 90 events with $M_{\mu\mu} > 130 \text{ GeV}/c^2$, and observe 1813 events.

We model the production and decay of a spin-1 resonance Z'_{SM} using MADEVENT [16], assuming fermionic couplings equal to those of the Z boson. Initial-state QCD radiation, hadronization, and showering are modelled with PYTHIA. The acceptance for Z' boson events varies with invariant mass, increasing from 20% at $200 \text{ GeV}/c^2$ to nearly 40% at $1 \text{ TeV}/c^2$; for pole masses above $M_{Z'} \approx 1 \text{ TeV}/c^2$, the resonance peak is suppressed due to lack of initial-state partons with sufficient momentum, leading to a drop in mean invariant mass and acceptance.

The previous CDF search [5] used a binned likelihood fit with bins uniform in the inverse of the reconstructed dimuon mass to extract the best-fit signal cross section at a range of masses. That approach weighted events in the same reconstructed inverse mass bin equally, independent of expected track resolutions. We improve on this method by using an unbinned likelihood that includes a theoretical model of the full kinematics of the event and the event-by-event knowledge of the muon p_T resolution. Well-measured events have narrower likelihoods, reflecting the higher quality of their information and making a stronger impact on the fitted result. The likelihood is then used both to set limits on the Z' cross section as a function of mass and to construct a 2D interval that gives a well-defined discovery condition. The likelihood is given by

$$L(M_{Z'}, s_{Z'}) = \frac{(N_{bg+Z'})^N e^{-(N_{bg+Z'})}}{N!} \prod_i^N L(x_i | M_{Z'}, s_{Z'}),$$

where $M_{Z'}$ is the Z' pole mass, N is the number of




FIG. 1: Top: The distributions of $M_{\mu\mu}$ for data with 4.6 fb^{-1} of integrated luminosity (triangles) and expected SM backgrounds (histograms) with two example Z'_{SM} signals. Bottom: The relative difference between observed and expected data, as a function of dimuon mass. Error bars show statistical uncertainty.

events in the data, $N_{Z'}(N_{bg+Z'})$ is the number of events expected due to Z' signal (and background), x_i represents the observed kinematics of the i -th event, and $s_{Z'} = N_{Z'}/(N_{bg+Z'})$. The dependence of the per-event likelihood is given by

$$L(x_i|M_{Z'}, s_{Z'}) = s_{Z'} L_{Z'}(x_i|M_{Z'}) + (1 - s_{Z'}) L_{Z/\gamma^*}(x_i),$$

and the likelihood $L_{Z'}$ is calculated by integrating the matrix element for Z' production convolved with PDFs and the detector resolution functions:

$$L_{Z'}(x_i = p_1, p_2, \sigma_{p_{T1}}, \sigma_{p_{T2}}, N_{\text{jets}}|M_{Z'}) =$$

$$\int d\Phi(q_1, q_2) |\mathcal{M}_{Z'}(q_1, q_2, M_{Z'})|^2 f_{PDF}^p f_{PDF}^{\bar{p}}$$

$$\times T(p_1, q_1, \sigma_{p_{T1}}) T(p_2, q_2, \sigma_{p_{T2}}) \times P_{PT}(q_1 + q_2, N_{\text{jets}}),$$

where $p_{1,2}$ represent the four-vectors of the two measured muons, $q_{1,2}$ represent the unknown four-vectors of the two true muons, Φ represents phase space for the true muons, \mathcal{M} is the matrix element, f_{PDF} is the parton distribution function [12]. $T(p, q, \sigma_{p_T})$ is the transfer function that parametrizes the detector resolution as a function of the measured uncertainty σ_{p_T} , extracted from simulated data; the function uses two Gaussian functions in $1/p_T^\mu$, to describe the bulk and tail regions. P_{PT} is the probability density function for p_T of the $\mu\mu$ system, parameterized in the number of jets (N_{jets}) with $E_T^{\text{jet}} > 15 \text{ GeV}$ and $|\eta^{\text{jet}}| < 2.5$, extracted from simulated samples with initial- and final-state radiation; the function uses two exponential functions in $p_T^{\mu\mu}$ to describe the bulk and tail regions. $L_{Z'}$ is evaluated numerically using importance-sampling techniques. The distributions of σ_{p_T} for the Z' signal and dominant Z/γ^* background are the same in the phase space region near the hypothesized Z' mass, thus they do not affect the likelihood ratio ordering. An analogous expression is used for L_{Z/γ^*} , which describes the likelihood for the dominant Z/γ^* background. In simulated experiments, maximization of $L(x_i|M_{Z'}, s_{Z'})$ faithfully recovers the correct values of $(M_{Z'}, s_{Z'})$.

We analyze the resulting likelihood in two ways. First, we aim to discover the regions in $(M_{Z'}, s_{Z'})$ that are consistent with the data and inconsistent with the SM, making no assumptions about the relationship between $M_{Z'}$ and $s_{Z'}$. We refer to this as the *2D interval* analysis. Second, we wish to set limits on the Z' mass in specific models. In that case, we perform a *raster scan*, in which we choose a set of values of $M_{Z'}$, and at each point derive limits on $s_{Z'}$. Together with a prediction for $s_{Z'}(M_{Z'})$ in a specific theory, we can use the raster scan to place lower limits on $M_{Z'}$.

The 2D interval is constructed via the unified ordering scheme [17] in two dimensions, resonance mass and cross section (see Fig. 2). At each test point in the $(M_{Z'}, s_{Z'})$ space, we calculate the ratio of the likelihood at the test point to the likelihood at the best-fit point where the likelihood is maximized. To determine which test points are consistent with the data at a given confidence level, we perform pseudo-experiments drawn from fully-simulated samples over a grid of mass and Z' signal cross-section; these pseudo-experiments allow us to determine the fraction of pseudo-experiments which have a larger likelihood ratio. Test points which lie between the grid of simulated samples are interpolated between neighboring grid points. The pseudo-experiments include all backgrounds and interference effects between the Z and Z' , as well as variation of the nuisance parameters from systematic uncertainties described below. This approach is well designed for discovery, as it tests the background hypothesis exactly once. The significance of an observation corresponds to the first confidence-level contour that includes a signal rate of zero. It also provides a summary

of Z' models consistent with the data without relying on specific model details. In the 2D interval analysis, the best-fit signal cross section of $\sigma = 26 \text{ fb}$ occurs at a resonance mass of $M = 199 \text{ GeV}/c^2$ but is consistent with the standard model at 16% confidence level; in 84% of simulated experiments with no Z' events, we observe a more significant excess.




FIG. 2: Observed 95% CL limits in 1D raster scan (top) and 2D intervals (bottom) for data with 4.6 fb^{-1} of integrated luminosity. The solid circle indicates the best-fit value (\hat{M}, \hat{s}); in regions below each line, 25, 50, 68, 95, and 99% of pseudo-experiments drawn from the point (M, s) yield a likelihood ratio $L(M, s)/L(\hat{M}, \hat{s})$ greater or equal to the observed ratio in data. A 16% region (not shown) is the smallest region that still includes the standard model.

The raster scan is the traditional approach used in many analyses, including the previous Z' search [5]. It provides mass limits on theories that enforce a relationship between the signal fraction and the resonance mass, and is appropriate if outside information indicates a particular mass is interesting. In the presence of a significant excess above the background-only hypothesis, one must account for the number of possible Z' masses, each of which tests the background-only hypothesis: the *look-elsewhere effect*. The 2D analysis needs no such correction. While the resulting discovery significance of the




FIG. 3: Observed 95% CL limits for data with 4.6 fb^{-1} of integrated luminosity expressed as limits on the up and down type charges c_u and c_d [4]. The solid and dotted lines show possible models in $U(1)_{B-XL}$ and $U(1)_{10+x5}$ groups, respectively. The dashed lines show the range for models in the $U(1)_{q+xu}$ group.

corrected raster scan will be correct, one will still be left with an interval in the signal fraction at every mass. In contrast, in the presence of a signal, the 2D analysis will provide a range of masses that is consistent with the signal.

TABLE I: Mass limits on specific spin-1 Z' models [4] in data with 4.6 fb^{-1} of integrated luminosity at 95% confidence level.

Model	Z'_l	Z'_{sec}	Z'_N	Z'_{ψ}	Z'_{χ}	Z'_{η}	Z'_{SM}
Mass Limit (GeV/ c^2)	817	858	900	917	930	938	1071

Dominant systematic uncertainties [5] include uncertainties on the PDFs and the dependence of the next-to-leading order cross section on the dimuon invariant mass. These weaken the final limits by 5-10% depending on mass. Additional uncertainties are the level of initial state radiation and muon acceptance at large transverse momentum.

The raster scan in mass allows us to set strong limits on specific models of Z' production; see Fig. 2 and Table I. The production cross section times branching fraction to the dimuon final state is determined by the couplings of the fermions to the Z' . Figure 3 shows how mass limits depend on the charges of the up- and down-type fermions to the $U(1)$ group associated with the Z' . Table I shows the limits for the specific models described in Ref [4].

In conclusion, we have applied the matrix-element-based likelihood technique to a search for new spin-1 resonances decaying to muon pairs, set the strong limits on the resonance cross section and mass, and introduced a statistical analysis approach that is useful for this analysis as well as for potential LHC discoveries.

We acknowledge useful conversations with Tim Tait. We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucléaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

Phys. **58**, 247 (2007).

- [3] M. Cvetic and P. Langacker, Phys. Rev. D **54**, 3570 (1996); M. Cvetic *et al.*, Phys. Rev. D **56**, 2861 (1997); E. Keith and E. Ma, Phys. Rev. D **56**, 7155 (1997); P. Langacker and J. Wang, Phys. Rev. D **58**, 115010 (1998).
- [4] M. Carena, A. Daleo, B. A. Dobrescu, and T. M. P. Tait, Phys. Rev. D **70**, 093009 (2004).
- [5] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **102**, 091805 (2009).
- [6] V. Abazov *et al.* (D0 Collaboration), arXiv:1008.2023v1 (2010).
- [7] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **102**, 031801 (2009).
- [8] D. E. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005); A. Abulencia *et al.* (CDF Collaboration), J. Phys. G **34**, 2457 (2007); T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **77**, 112001 (2008).
- [9] CDF uses a cylindrical coordinate system with the z axis along the proton beam axis. Pseudorapidity is $\eta \equiv -\ln(\tan(\theta/2))$, where θ is the polar angle relative to the proton beam direction, and ϕ is the azimuthal angle while $p_T = |p| \sin \theta$, $E_T = E \sin \theta$.
- [10] A. V. Kotwal, H. K. Gerberich, and C. Hays, Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 110 (2003).
- [11] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135** (2001) 238; T. Sjöstrand, Comput. Phys. Commun. **82**, 74 (1994).
- [12] J. Pumplin *et al.* (CTEQ Collaboration), J. High. Energy Phys. 07 (2002) 012.
- [13] T. Affolder *et al.* (CDF Collaboration), Nucl. Instrum. Methods **A447**, 1 (2000).
- [14] J. M. Campbell and R. K. Ellis, Phys. Rev. D **60**, 113006 (1999); R. Bonciani, S. Catani, M. L. Mangano, and P. Nason, Nucl. Phys. B **529**, 424 (1998); M. Cacciari *et al.*, J. High Energy Phys. 04 (2004) 068.
- [15] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **93**, 142001 (2004).
- [16] J. Alwall *et al.* J. High Energy Phys 07 (2007) 028
- [17] G. J. Feldman and R. D. Cousins, Phys. Rev. D **57**, 3873 (1998).

-
- [1] P. Langacker, Rev. Mod. Phys. **81**, 1199 (2009), and references therein.
 - [2] N. Arkani-Hamed, A. G. Cohen, and H. Georgi, Phys. Lett. B **513**, 232 (2001); T. Han, H. E. Logan, B. McElrath and L.-T. Wang, Phys. Rev. D **67**, 095004 (2003); T. Han, H. E. Logan, and L.-T. Wang, J. High Energy Phys. **01**, 099 (2006); M. Perelstein, Prog. Part. Nucl.