



# CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Suppression of Back-to-Back Hadron Pairs at Forward Rapidity in d+Au Collisions at $\sqrt{s_{NN}}=200$ GeV

A. Adare *et al.* (PHENIX Collaboration)

Phys. Rev. Lett. **107**, 172301 — Published 18 October 2011

DOI: [10.1103/PhysRevLett.107.172301](https://doi.org/10.1103/PhysRevLett.107.172301)

**Suppression of back-to-back hadron pairs at forward rapidity in  $d+Au$  collisions  
at  $\sqrt{s_{NN}} = 200$  GeV**

A. Adare,<sup>11</sup> S. Afanasiev,<sup>26</sup> C. Aidala,<sup>39</sup> N.N. Ajitanand,<sup>55</sup> Y. Akiba,<sup>50,51</sup> H. Al-Bataineh,<sup>45</sup> J. Alexander,<sup>55</sup>  
A. Angerami,<sup>12</sup> K. Aoki,<sup>32,50</sup> N. Apadula,<sup>56</sup> Y. Aramaki,<sup>10</sup> E.T. Atomssa,<sup>33</sup> R. Averbeck,<sup>56</sup> T.C. Awes,<sup>46</sup>  
B. Azmoun,<sup>5</sup> V. Babintsev,<sup>21</sup> M. Bai,<sup>4</sup> G. Baksay,<sup>17</sup> L. Baksay,<sup>17</sup> K.N. Barish,<sup>6</sup> B. Bassalleck,<sup>44</sup> A.T. Basye,<sup>1</sup>  
S. Bathe,<sup>6,51</sup> V. Baublis,<sup>49</sup> C. Baumann,<sup>40</sup> A. Bazilevsky,<sup>5</sup> S. Belikov,<sup>5,\*</sup> R. Belmont,<sup>60</sup> R. Bennett,<sup>56</sup>  
A. Berdnikov,<sup>53</sup> Y. Berdnikov,<sup>53</sup> J.H. Bhom,<sup>63</sup> D.S. Blau,<sup>31</sup> J.S. Bok,<sup>63</sup> K. Boyle,<sup>56</sup> M.L. Brooks,<sup>35</sup> H. Buesching,<sup>5</sup>  
V. Bumazhnov,<sup>21</sup> G. Bunce,<sup>5,51</sup> S. Butsyk,<sup>35</sup> S. Campbell,<sup>56</sup> A. Caringi,<sup>41</sup> C.-H. Chen,<sup>56</sup> C.Y. Chi,<sup>12</sup> M. Chiu,<sup>5</sup>  
I.J. Choi,<sup>63</sup> J.B. Choi,<sup>8</sup> R.K. Choudhury,<sup>3</sup> P. Christiansen,<sup>37</sup> T. Chujo,<sup>59</sup> P. Chung,<sup>55</sup> O. Chvala,<sup>6</sup>  
V. Cianciolo,<sup>46</sup> Z. Citron,<sup>56</sup> B.A. Cole,<sup>12</sup> Z. Conesa del Valle,<sup>33</sup> M. Connors,<sup>56</sup> M. Csanád,<sup>15</sup> T. Csörgő,<sup>29</sup>  
T. Dahms,<sup>56</sup> S. Dairaku,<sup>32,50</sup> I. Danchev,<sup>60</sup> K. Das,<sup>18</sup> A. Datta,<sup>39</sup> G. David,<sup>5</sup> M.K. Dayananda,<sup>19</sup> A. Denisov,<sup>21</sup>  
A. Deshpande,<sup>51,56</sup> E.J. Desmond,<sup>5</sup> K.V. Dharmawardane,<sup>45</sup> O. Dietzsch,<sup>54</sup> A. Dion,<sup>25</sup> M. Donadelli,<sup>54</sup>  
O. Drapier,<sup>33</sup> A. Drees,<sup>56</sup> K.A. Drees,<sup>4</sup> J.M. Durham,<sup>56</sup> A. Durum,<sup>21</sup> D. Dutta,<sup>3</sup> L. D’Orazio,<sup>38</sup> S. Edwards,<sup>18</sup>  
Y.V. Efremenko,<sup>46</sup> F. Ellinghaus,<sup>11</sup> T. Engelmore,<sup>12</sup> A. Enokizono,<sup>46</sup> H. En’yo,<sup>50,51</sup> S. Esumi,<sup>59</sup> B. Fadem,<sup>41</sup>  
D.E. Fields,<sup>44</sup> M. Finger,<sup>7</sup> M. Finger, Jr.,<sup>7</sup> F. Fleuret,<sup>33</sup> S.L. Fokin,<sup>31</sup> Z. Fraenkel,<sup>62,\*</sup> J.E. Frantz,<sup>56</sup> A. Franz,<sup>5</sup>  
A.D. Frawley,<sup>18</sup> K. Fujiwara,<sup>50</sup> Y. Fukao,<sup>50</sup> T. Fusayasu,<sup>43</sup> I. Garishvili,<sup>57</sup> A. Glenn,<sup>34</sup> H. Gong,<sup>56</sup> M. Gonin,<sup>33</sup>  
Y. Goto,<sup>50,51</sup> R. Granier de Cassagnac,<sup>33</sup> N. Grau,<sup>12</sup> S.V. Greene,<sup>60</sup> G. Grim,<sup>35</sup> M. Grosse Perdekamp,<sup>22</sup>  
T. Gunji,<sup>10</sup> H.-Å. Gustafsson,<sup>37,\*</sup> J.S. Haggerty,<sup>5</sup> K.I. Hahn,<sup>16</sup> H. Hamagaki,<sup>10</sup> J. Hamblen,<sup>57</sup> R. Han,<sup>48</sup> J. Hanks,<sup>12</sup>  
E. Haslum,<sup>37</sup> R. Hayano,<sup>10</sup> X. He,<sup>19</sup> M. Heffner,<sup>34</sup> T.K. Hemmick,<sup>56</sup> T. Hester,<sup>6</sup> J.C. Hill,<sup>25</sup> M. Hohlmann,<sup>17</sup>  
W. Holzmann,<sup>12</sup> K. Homma,<sup>20</sup> B. Hong,<sup>30</sup> T. Horaguchi,<sup>20</sup> D. Hornback,<sup>57</sup> S. Huang,<sup>60</sup> T. Ichihara,<sup>50,51</sup>  
R. Ichimiya,<sup>50</sup> Y. Ikeda,<sup>59</sup> K. Imai,<sup>32,50</sup> M. Inaba,<sup>59</sup> D. Isenhower,<sup>1</sup> M. Ishihara,<sup>50</sup> M. Issah,<sup>60</sup> A. Isupov,<sup>26</sup>  
D. Ivanishev,<sup>49</sup> Y. Iwanaga,<sup>20</sup> B.V. Jacak,<sup>56,†</sup> J. Jia,<sup>5,55</sup> X. Jiang,<sup>35</sup> J. Jin,<sup>12</sup> B.M. Johnson,<sup>5</sup> T. Jones,<sup>1</sup> K.S. Joo,<sup>42</sup>  
D. Jouan,<sup>47</sup> D.S. Jumper,<sup>1</sup> F. Kajihara,<sup>10</sup> J. Kamin,<sup>56</sup> J.H. Kang,<sup>63</sup> J. Kapustinsky,<sup>35</sup> K. Karatsu,<sup>32</sup> M. Kasai,<sup>52,50</sup>  
D. Kawall,<sup>39,51</sup> M. Kawashima,<sup>52,50</sup> A.V. Kazantsev,<sup>31</sup> T. Kempel,<sup>25</sup> A. Khanzadeev,<sup>49</sup> K.M. Kijima,<sup>20</sup> J. Kikuchi,<sup>61</sup>  
A. Kim,<sup>16</sup> B.I. Kim,<sup>30</sup> D.J. Kim,<sup>27</sup> E.J. Kim,<sup>8</sup> Y.-J. Kim,<sup>22</sup> E. Kinney,<sup>11</sup> Á. Kiss,<sup>15</sup> E. Kistenev,<sup>5</sup> L. Kochenda,<sup>49</sup>  
B. Komkov,<sup>49</sup> M. Konno,<sup>59</sup> J. Koster,<sup>22</sup> A. Král,<sup>13</sup> A. Kravitz,<sup>12</sup> G.J. Kunde,<sup>35</sup> K. Kurita,<sup>52,50</sup> M. Kurosawa,<sup>50</sup>  
Y. Kwon,<sup>63</sup> G.S. Kyle,<sup>45</sup> R. Lacey,<sup>55</sup> Y.S. Lai,<sup>12</sup> J.G. Lajoie,<sup>25</sup> A. Lebedev,<sup>25</sup> D.M. Lee,<sup>35</sup> J. Lee,<sup>16</sup> K.B. Lee,<sup>30</sup>  
K.S. Lee,<sup>30</sup> M.J. Leitch,<sup>35</sup> M.A.L. Leite,<sup>54</sup> X. Li,<sup>9</sup> P. Lichtenwalner,<sup>41</sup> P. Liebing,<sup>51</sup> L.A. Linden Levy,<sup>11</sup> T. Liška,<sup>13</sup>  
A. Litvinenko,<sup>26</sup> H. Liu,<sup>35</sup> M.X. Liu,<sup>35</sup> B. Love,<sup>60</sup> D. Lynch,<sup>5</sup> C.F. Maguire,<sup>60</sup> Y.I. Makdisi,<sup>4</sup> A. Malakhov,<sup>26</sup>  
M.D. Malik,<sup>44</sup> V.I. Manko,<sup>31</sup> E. Mannel,<sup>12</sup> Y. Mao,<sup>48,50</sup> H. Masui,<sup>59</sup> F. Matathias,<sup>12</sup> M. McCumber,<sup>56</sup>  
P.L. McGaughey,<sup>35</sup> N. Means,<sup>56</sup> B. Meredith,<sup>22</sup> Y. Miake,<sup>59</sup> T. Mibe,<sup>28</sup> A.C. Mignerey,<sup>38</sup> K. Miki,<sup>59</sup> A. Milov,<sup>5</sup>  
J.T. Mitchell,<sup>5</sup> A.K. Mohanty,<sup>3</sup> H.J. Moon,<sup>42</sup> Y. Morino,<sup>10</sup> A. Morreale,<sup>6</sup> D.P. Morrison,<sup>5</sup> T.V. Moukhanova,<sup>31</sup>  
T. Murakami,<sup>32</sup> J. Murata,<sup>52,50</sup> S. Nagamiya,<sup>28</sup> J.L. Nagle,<sup>11</sup> M. Naglis,<sup>62</sup> M.I. Nagy,<sup>29</sup> I. Nakagawa,<sup>50,51</sup>  
Y. Nakamiya,<sup>20</sup> K.R. Nakamura,<sup>32</sup> T. Nakamura,<sup>50</sup> K. Nakano,<sup>50</sup> S. Nam,<sup>16</sup> J. Newby,<sup>34</sup> M. Nguyen,<sup>56</sup>  
M. Nihashi,<sup>20</sup> R. Nouicer,<sup>5</sup> A.S. Nyanin,<sup>31</sup> C. Oakley,<sup>19</sup> E. O’Brien,<sup>5</sup> S.X. Oda,<sup>10</sup> C.A. Ogilvie,<sup>25</sup> M. Oka,<sup>59</sup>  
K. Okada,<sup>51</sup> Y. Onuki,<sup>50</sup> A. Oskarsson,<sup>37</sup> M. Ouchida,<sup>20</sup> K. Ozawa,<sup>10</sup> R. Pak,<sup>5</sup> V. Pantuev,<sup>23,56</sup> V. Papavassiliou,<sup>45</sup>  
I.H. Park,<sup>16</sup> S.K. Park,<sup>30</sup> W.J. Park,<sup>30</sup> S.F. Pate,<sup>45</sup> H. Pei,<sup>25</sup> J.-C. Peng,<sup>22</sup> H. Pereira,<sup>14</sup> V. Peresedov,<sup>26</sup>  
D.Yu. Peressounko,<sup>31</sup> R. Petti,<sup>56</sup> C. Pinkenburg,<sup>5</sup> R.P. Pisani,<sup>5</sup> M. Proissl,<sup>56</sup> M.L. Purschke,<sup>5</sup> H. Qu,<sup>19</sup> J. Rak,<sup>27</sup>  
I. Ravinovich,<sup>62</sup> K.F. Read,<sup>46,57</sup> K. Reygers,<sup>40</sup> V. Riabov,<sup>49</sup> Y. Riabov,<sup>49</sup> E. Richardson,<sup>38</sup> D. Roach,<sup>60</sup> G. Roche,<sup>36</sup>  
S.D. Rolnick,<sup>6</sup> M. Rosati,<sup>25</sup> C.A. Rosen,<sup>11</sup> S.S.E. Rosendahl,<sup>37</sup> P. Rukoyatkin,<sup>26</sup> P. Ružička,<sup>24</sup> B. Sahlmueller,<sup>40</sup>  
N. Saito,<sup>28</sup> T. Sakaguchi,<sup>5</sup> K. Sakashita,<sup>50,58</sup> V. Samsonov,<sup>49</sup> S. Sano,<sup>10,61</sup> T. Sato,<sup>59</sup> S. Sawada,<sup>28</sup> K. Sedgwick,<sup>6</sup>  
J. Seele,<sup>11</sup> R. Seidl,<sup>22,51</sup> R. Seto,<sup>6</sup> D. Sharma,<sup>62</sup> I. Shein,<sup>21</sup> T.-A. Shibata,<sup>50,58</sup> K. Shigaki,<sup>20</sup> M. Shimomura,<sup>59</sup>  
K. Shoji,<sup>32,50</sup> P. Shukla,<sup>3</sup> A. Sickles,<sup>5</sup> C.L. Silva,<sup>25</sup> D. Silvermyr,<sup>46</sup> C. Silvestre,<sup>14</sup> K.S. Sim,<sup>30</sup> B.K. Singh,<sup>2</sup>  
C.P. Singh,<sup>2</sup> V. Singh,<sup>2</sup> M. Slunečka,<sup>7</sup> R.A. Soltz,<sup>34</sup> W.E. Sondheim,<sup>35</sup> S.P. Sorensen,<sup>57</sup> I.V. Sourikova,<sup>5</sup>  
P.W. Stankus,<sup>46</sup> E. Stenlund,<sup>37</sup> S.P. Stoll,<sup>5</sup> T. Sugitate,<sup>20</sup> A. Sukhanov,<sup>5</sup> J. Sziklai,<sup>29</sup> E.M. Takagui,<sup>54</sup>  
A. Taketani,<sup>50,51</sup> R. Tanabe,<sup>59</sup> Y. Tanaka,<sup>43</sup> S. Taneja,<sup>56</sup> K. Tanida,<sup>32,50,51</sup> M.J. Tannenbaum,<sup>5</sup> S. Tarafdar,<sup>2</sup>  
A. Taranenko,<sup>55</sup> H. Themann,<sup>56</sup> D. Thomas,<sup>1</sup> T.L. Thomas,<sup>44</sup> M. Togawa,<sup>51</sup> A. Toia,<sup>56</sup> L. Tomášek,<sup>24</sup> H. Torii,<sup>20</sup>  
R.S. Towell,<sup>1</sup> I. Tserruya,<sup>62</sup> Y. Tsuchimoto,<sup>20</sup> C. Vale,<sup>5</sup> H. Valle,<sup>60</sup> H.W. van Hecke,<sup>35</sup> E. Vazquez-Zambrano,<sup>12</sup>  
A. Veicht,<sup>22</sup> J. Velkovska,<sup>60</sup> R. Vértesi,<sup>29</sup> M. Virius,<sup>13</sup> V. Vrba,<sup>24</sup> E. Vznuzdaev,<sup>49</sup> X.R. Wang,<sup>45</sup> D. Watanabe,<sup>20</sup>  
K. Watanabe,<sup>59</sup> Y. Watanabe,<sup>50,51</sup> F. Wei,<sup>25</sup> R. Wei,<sup>55</sup> J. Wessels,<sup>40</sup> S.N. White,<sup>5</sup> D. Winter,<sup>12</sup> C.L. Woody,<sup>5</sup>

R.M. Wright,<sup>1</sup> M. Wysocki,<sup>11</sup> Y.L. Yamaguchi,<sup>10</sup> K. Yamaura,<sup>20</sup> R. Yang,<sup>22</sup> A. Yanovich,<sup>21</sup> J. Ying,<sup>19</sup>  
S. Yokkaichi,<sup>50,51</sup> Z. You,<sup>48</sup> G.R. Young,<sup>46</sup> I. Younus,<sup>44</sup> I.E. Yushmanov,<sup>31</sup> W.A. Zajc,<sup>12</sup> S. Zhou,<sup>9</sup> and L. Zolin<sup>26</sup>

(PHENIX Collaboration)

- <sup>1</sup>Abilene Christian University, Abilene, Texas 79699, USA  
<sup>2</sup>Department of Physics, Banaras Hindu University, Varanasi 221005, India  
<sup>3</sup>Bhabha Atomic Research Centre, Bombay 400 085, India  
<sup>4</sup>Collider-Accelerator Department, Brookhaven National Laboratory, Upton, New York 11973-5000, USA  
<sup>5</sup>Physics Department, Brookhaven National Laboratory, Upton, New York 11973-5000, USA  
<sup>6</sup>University of California - Riverside, Riverside, California 92521, USA  
<sup>7</sup>Charles University, Ovocný trh 5, Praha 1, 116 36, Prague, Czech Republic  
<sup>8</sup>Chonbuk National University, Jeonju, 561-756, Korea  
<sup>9</sup>China Institute of Atomic Energy (CIAE), Beijing, People's Republic of China  
<sup>10</sup>Center for Nuclear Study, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan  
<sup>11</sup>University of Colorado, Boulder, Colorado 80309, USA  
<sup>12</sup>Columbia University, New York, New York 10027 and Nevis Laboratories, Irvington, New York 10533, USA  
<sup>13</sup>Czech Technical University, Zikova 4, 166 36 Prague 6, Czech Republic  
<sup>14</sup>Dapnia, CEA Saclay, F-91191, Gif-sur-Yvette, France  
<sup>15</sup>ELTE, Eötvös Loránd University, H - 1117 Budapest, Pázmány P. s. 1/A, Hungary  
<sup>16</sup>Ewha Womans University, Seoul 120-750, Korea  
<sup>17</sup>Florida Institute of Technology, Melbourne, Florida 32901, USA  
<sup>18</sup>Florida State University, Tallahassee, Florida 32306, USA  
<sup>19</sup>Georgia State University, Atlanta, Georgia 30303, USA  
<sup>20</sup>Hiroshima University, Kagamiyama, Higashi-Hiroshima 739-8526, Japan  
<sup>21</sup>IHEP Protvino, State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, 142281, Russia  
<sup>22</sup>University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA  
<sup>23</sup>Institute for Nuclear Research of the Russian Academy of Sciences, prospekt 60-letiya Oktyabrya 7a, Moscow 117312, Russia  
<sup>24</sup>Institute of Physics, Academy of Sciences of the Czech Republic, Na Slovance 2, 182 21 Prague 8, Czech Republic  
<sup>25</sup>Iowa State University, Ames, Iowa 50011, USA  
<sup>26</sup>Joint Institute for Nuclear Research, 141980 Dubna, Moscow Region, Russia  
<sup>27</sup>Helsinki Institute of Physics and University of Jyväskylä, P.O.Box 35, FI-40014 Jyväskylä, Finland  
<sup>28</sup>KEK, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan  
<sup>29</sup>KFKI Research Institute for Particle and Nuclear Physics of the Hungarian Academy of Sciences (MTA KFKI RMKI), H-1525 Budapest 114, POBox 49, Budapest, Hungary  
<sup>30</sup>Korea University, Seoul, 136-701, Korea  
<sup>31</sup>Russian Research Center "Kurchatov Institute", Moscow, 123098 Russia  
<sup>32</sup>Kyoto University, Kyoto 606-8502, Japan  
<sup>33</sup>Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS-IN2P3, Route de Saclay, F-91128, Palaiseau, France  
<sup>34</sup>Lawrence Livermore National Laboratory, Livermore, California 94550, USA  
<sup>35</sup>Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA  
<sup>36</sup>LPC, Université Blaise Pascal, CNRS-IN2P3, Clermont-Fd, 63177 Aubiere Cedex, France  
<sup>37</sup>Department of Physics, Lund University, Box 118, SE-221 00 Lund, Sweden  
<sup>38</sup>University of Maryland, College Park, Maryland 20742, USA  
<sup>39</sup>Department of Physics, University of Massachusetts, Amherst, Massachusetts 01003-9337, USA  
<sup>40</sup>Institut für Kernphysik, University of Muenster, D-48149 Muenster, Germany  
<sup>41</sup>Muhlenberg College, Allentown, Pennsylvania 18104-5586, USA  
<sup>42</sup>Myongji University, Yongin, Kyonggido 449-728, Korea  
<sup>43</sup>Nagasaki Institute of Applied Science, Nagasaki-shi, Nagasaki 851-0193, Japan  
<sup>44</sup>University of New Mexico, Albuquerque, New Mexico 87131, USA  
<sup>45</sup>New Mexico State University, Las Cruces, New Mexico 88003, USA  
<sup>46</sup>Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA  
<sup>47</sup>IPN-Orsay, Université Paris Sud, CNRS-IN2P3, BP1, F-91406, Orsay, France  
<sup>48</sup>Peking University, Beijing, People's Republic of China  
<sup>49</sup>PNPI, Petersburg Nuclear Physics Institute, Gatchina, Leningrad region, 188300, Russia  
<sup>50</sup>RIKEN Nishina Center for Accelerator-Based Science, Wako, Saitama 351-0198, Japan  
<sup>51</sup>RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, New York 11973-5000, USA  
<sup>52</sup>Physics Department, Rikkyo University, 3-34-1 Nishi-Ikebukuro, Toshima, Tokyo 171-8501, Japan  
<sup>53</sup>Saint Petersburg State Polytechnic University, St. Petersburg, 195251 Russia  
<sup>54</sup>Universidade de São Paulo, Instituto de Física, Caixa Postal 66318, São Paulo CEP05315-970, Brazil  
<sup>55</sup>Chemistry Department, Stony Brook University, SUNY, Stony Brook, New York 11794-3400, USA  
<sup>56</sup>Department of Physics and Astronomy, Stony Brook University, SUNY, Stony Brook, New York 11794-3400, USA  
<sup>57</sup>University of Tennessee, Knoxville, Tennessee 37996, USA  
<sup>58</sup>Department of Physics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan  
<sup>59</sup>Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan

<sup>60</sup>*Vanderbilt University, Nashville, Tennessee 37235, USA*  
<sup>61</sup>*Waseda University, Advanced Research Institute for Science and Engineering, 17 Kikui-cho, Shinjuku-ku, Tokyo 162-0044, Japan*  
<sup>62</sup>*Weizmann Institute, Rehovot 76100, Israel*  
<sup>63</sup>*Yonsei University, IPAP, Seoul 120-749, Korea*

Back-to-back hadron pair yields in  $d+Au$  and  $p+p$  collisions at  $\sqrt{s_{NN}} = 200$  GeV were measured with the PHENIX detector at the Relativistic Heavy Ion Collider. Rapidity separated hadron pairs were detected with the trigger hadron at pseudorapidity  $|\eta| < 0.35$  and the associated hadron at forward rapidity (deuteron direction,  $3.0 < \eta < 3.8$ ). Pairs were also detected with both hadrons measured at forward rapidity; in this case the yield of back-to-back hadron pairs in  $d+Au$  collisions with small impact parameters is observed to be suppressed by a factor of 10 relative to  $p+p$  collisions. The kinematics of these pairs is expected to probe partons in the Au nucleus with low fraction  $x$  of the nucleon momenta, where the gluon densities rise sharply. The observed suppression as a function of nuclear thickness,  $p_T$ , and  $\eta$  points to cold nuclear matter effects arising at high parton densities in the nucleus probed by the deuteron.

PACS numbers: 25.75.Dw

Nuclear effects on quark and gluon densities of nucleons bound in nuclei can be studied in collisions of deuteron and gold nuclei at the Relativistic Heavy Ion Collider. At  $\sqrt{s_{NN}} = 200$  GeV the hadron yields in the forward rapidity (deuteron going) direction were observed to be suppressed for  $d+Au$  collisions relative to  $p+p$  collisions [1–3]. However, the mechanism for the suppression was not firmly established and may indicate novel QCD effects in nuclei. Competing theoretical approaches include initial state energy loss [4, 5], parton recombination [6], shadowing effects [7, 8] and gluon saturation [9].

Back-to-back dijet yields were proposed as an additional observable to distinguish better between competing mechanisms. The Color Glass Condensate (CGC) framework [10] predicts that quarks and gluons scattering at forward angles (large rapidity) will interact coherently off gluons at low  $x$  in the gold nucleus. As a result, the rate of observed recoiling jets is expected to be suppressed in  $d+Au$  collisions compared to  $p+p$ , and angular broadening of the back-to-back correlation of jets is predicted [11, 12]. Dihadron correlation measurements were used [13, 14] successfully in  $d+Au$  collisions to select dijet production based on the back-to-back peak at  $\Delta\phi=\pi$  between trigger hadrons and their associated hadrons. Dihadron correlation measurements with varying kinematic constraints (transverse momentum  $p_T$  and rapidity) probe different  $x$  ranges in the nucleus. In particular, measurements at forward rapidity are thought to probe small  $x$  values in the Au nucleus.

In this Letter we report results on the suppression in  $d+Au$  relative to  $p+p$  collisions of inclusive  $\pi^0$ 's and back-to-back cluster- $\pi^0$  pairs in the forward rapidity region, and for back-to-back  $\pi^0$ - $\pi^0$  or hadron- $\pi^0$  pairs separated in rapidity. The data were obtained from  $p+p$  and  $d+Au$  runs in 2008 with the PHENIX detector and include a new electromagnetic calorimeter, the Muon Piston Calorimeter (MPC) with an acceptance of  $3.0 < \eta < 3.8$  in pseudorapidity and  $0 < \phi < 2\pi$ . The clusters are reconstructed from the energy deposit of photons in individual MPC towers. The MPC comprises 220  $PbWO_4$  towers of  $20.2X_0$  depth, with lateral dimensions of  $2.2 \times 2.2$  cm<sup>2</sup>, and is located 220 cm along the beam axis from the nominal interaction point.

The  $d+Au$  sample is separated into four centrality classes, 0–20% (most central), 20–40%, 40–60%, and 60–88% (most peripheral), based on charge deposited in the backward (gold direction) beam-beam counter (BBC,  $3.0 < -\eta < 3.9$ ). We determine the average number of binary collisions  $\langle N_{coll} \rangle$  from a Glauber model [3] and a simulation of the BBC;  $\langle N_{coll} \rangle$  values are  $15.1 \pm 1.0$ ,  $10.2 \pm 0.70$ ,  $6.6 \pm 0.44$ , and  $3.2 \pm 0.19$ , respectively.

The charged hadron ( $h^\pm$ ) and  $\pi^0$  analysis in the midrapidity region  $|\eta| < 0.35$  is identical to that for previous measurements by PHENIX [15, 16]. For the analysis in the forward rapidity region, a fiducial cut is applied ensuring that clusters are fully reconstructed within the MPC acceptance. Photon candidates are identified in the MPC by comparing cluster candidates to the expected shower profile for photons. The shower profile for photons is determined from the PHENIX GEANT3 [17] based detector simulation, PISA, which was tuned to reproduce MPC test beam data. For the pair analyses reported here, the associated particles in the MPC are  $\pi^0$ 's, which are identified by reconstructing the mass of their decay photon pairs. The  $\pi^0$  yield is obtained after subtraction of the combinatoric background of uncorrelated photon pairs. The shape of the background was determined from  $p+p$  PYTHIA 6.4 [18] and  $d+Au$  HIJING [19] events that are subsequently processed through PISA. The  $p_T$ -dependent systematic uncertainty on the associated  $\pi^0$  yield extraction procedure is estimated to be 2–5% for  $p+p$  and  $d+Au$ .

The closeness of the MPC to the collision vertex and the high energy of particles emitted in the forward direction make it difficult to reconstruct photon pairs from  $\pi^0$  decays at high  $p_T$ . For example at  $p_T = 1$  GeV/ $c$ , approximately 30% of the photon cluster pairs are merged and cannot be reconstructed separately in the MPC. To extend the  $p_T$




FIG. 1: (color online) Pedestal-subtracted  $\pi^0$ - $\pi^0$  per-trigger correlation functions for, as indicated,  $p+p$ ,  $d+Au$  peripheral (60–88% centrality) and  $d+Au$  central (0–20% centrality) collisions at  $\sqrt{s_{NN}} = 200$  GeV; the associated  $\pi^0$ 's of  $p_T = 0.5$ – $0.75$  GeV/ $c$  are measured at forward rapidity ( $3.0 < \eta < 3.8$ ) and the triggered  $\pi^0$ 's are measured at midrapidity ( $|\eta| < 0.35$ ) for the indicated  $p_T$  ranges. The subtracted pedestal values,  $b_0$ , are also indicated.




FIG. 2: (color online) Pedestal-subtracted cluster- $\pi^0$  per-trigger correlation functions measured at forward rapidity ( $3.0 < \eta < 3.8$ ) for, as indicated,  $p+p$ ,  $d+Au$  peripheral (60–88% centrality) and  $d+Au$  central (0–20% centrality) collisions at  $\sqrt{s_{NN}} = 200$  GeV; the correlation functions are for associated  $\pi^0$ 's of  $p_T = 0.5$ – $0.75$  GeV/ $c$  and trigger clusters over the indicated  $p_T$  ranges. Systematic uncertainties of up to 30% on the near side ( $|\Delta\phi| < 0.5$ ) are not shown. The subtracted pedestal values,  $b_0$ , are also indicated.

range and the pair yield, single electromagnetic clusters are used as trigger particles to construct cluster- $\pi^0$  dihadron pairs in the MPC. These trigger clusters are treated assuming that they are all  $\pi^0$ 's. However, PYTHIA studies indicate that  $\gtrsim 80\%$  of these trigger clusters are from  $\pi^0$ 's with the rest being dominantly single photons from asymmetric decays of  $\eta$  mesons or direct photons; thus, according to these studies a relatively small contamination remains. The cluster energy was corrected to the true  $\pi^0$  energy to account for the merging effects of the two photons from  $\pi^0$  decay. These corrections were determined by embedding Monte Carlo generated  $\pi^0$ 's into real data, as well as from PYTHIA tuned to match the data.

Figure 1 shows the azimuthal angle correlations between midrapidity and forward-rapidity  $\pi^0$  pairs, per  $\pi^0$  trigger detected at midrapidity, in  $p+p$ , peripheral  $d+Au$ , and central  $d+Au$  collisions for varying trigger  $\pi^0$   $p_T$ . Figure 2 shows the same correlations for trigger clusters where the cluster- $\pi^0$  pairs are both detected at forward rapidity. The constant pedestal,  $b_0$ , was subtracted from the correlation function. The correlations were corrected for the forward  $\pi^0$  detection efficiency and for the combinatoric background beneath the  $\pi^0$  peaks in the photon-pair invariant mass spectra. This background is determined by measurement of the azimuthal correlations for photon-pair mass selections adjacent to the  $\pi^0$  mass window and from studies of simulated jet events from PYTHIA events processed through PISA.




FIG. 3: (color online) Relative yield  $J_{dA}$  versus  $\langle N_{\text{coll}} \rangle$  for forward rapidity ( $3.0 < \eta < 3.8$ )  $\pi^0$ 's paired with (left) midrapidity ( $|\eta| < 0.35$ ) hadrons and  $\pi^0$ 's and (right) forward rapidity ( $3.0 < \eta < 3.8$ ) cluster- $\pi^0$  pairs for the indicated combinations of  $p_T$  ranges. Also plotted as inverted solid triangles are the values of the forward  $\pi^0$   $R_{dA}$ . Around each data point the vertical bars indicate statistical uncertainties and the open boxes indicate point-to-point systematic uncertainties. The gray bar at the left in each panel represents a global systematic scale uncertainty of 9.7%. Additional centrality dependent systematic uncertainties of 7.5%, 5.1%, 4.1%, and 4.8% for the peripheral to central bins, respectively, are not shown. The  $\langle N_{\text{coll}} \rangle$  values within a centrality selection are offset from their actual values for visual clarity (see text for actual  $\langle N_{\text{coll}} \rangle$  values).

For the mid-forward correlations (Fig. 1), due to the large pseudorapidity gap of  $\Delta\eta \sim 3.3$  between the hadrons, only an away-side peak ( $\Delta\phi = \pi$ ) is seen. For the forward-forward correlations a near-side peak ( $\Delta\phi = 0$ ) is also present (see Fig. 2). The yields and widths of the correlated pairs are extracted by fits to an away-side Gaussian signal shape plus a constant background ( $b_0$ ). The fit to the forward-forward correlations has an additional Gaussian signal for the near-side peak. The pedestal is determined from a fit in the mid-forward correlations and is consistent with the pedestal level found based on the assumption that the signal yield is 0 at the minimum of the correlation function - Zero Yield At Minimum (ZYAM) [20]. In the forward-forward correlations the ZYAM pedestal is used in the yield extraction. Additional systematic uncertainties of up to 30% (not shown in Fig. 2) are ascribed to the near-side peak due to corrections for resonance decays that contaminate the jet signal, and due to the acceptance loss around the trigger particle of  $\Delta\phi \times \Delta\eta \approx 0.5 \times 0.5$  rad, resulting from the minimum separation cut of one tower between cluster peaks in the MPC. The acceptance loss gives rise to the decrease observed for the near side peak.

Figures 1 and 2 show that the away-side peak for  $d+Au$  central collisions is suppressed compared to  $p+p$  collisions and peripheral  $d+Au$  collisions. This effect is large for the mid-forward correlations (Fig. 1) and becomes even larger when both particles are required to be in the forward rapidity region (Fig. 2).

For the mid-forward correlations, within their large uncertainties the Gaussian widths of the away-side correlation peak remain the same between  $p+p$  and central  $d+Au$  and the broadening predicted in the CGC framework in reference [11] is not observed. For example, in  $d+Au$  central collisions,  $\sigma = 0.93 \pm 0.09^{\text{stat}} \pm 0.139^{\text{syst}}$  for  $p_T^{\text{fwd}} = 1.25$  GeV/c and trigger particle momentum  $2.5 < p_T^t < 3.0$  GeV/c, while  $\sigma = 0.97 \pm 0.07^{\text{stat}} \pm 0.08^{\text{syst}}$  for  $p+p$  collisions. For the forward-forward correlations, the measurement does not discern whether there is appreciable broadening between  $d+Au$  and  $p+p$  collisions, as the ZYAM pedestal determination can bias the widths to smaller values.

The observed suppression is quantified by studying the relative yield,  $J_{dA}$  [21], of correlated back-to-back hadron pairs in  $d+Au$  collisions compared to  $p+p$  collisions scaled with  $\langle N_{\text{coll}} \rangle$ ,

$$J_{dA} = I_{dA} \times R_{dA}^t = \frac{1}{\langle N_{\text{coll}} \rangle} \frac{\sigma_{dA}^{\text{pair}} / \sigma_{dA}}{\sigma_{pp}^{\text{pair}} / \sigma_{pp}}, \quad (1)$$

where  $R_{dA}^t = (1/\langle N_{\text{coll}} \rangle) \cdot (\sigma_{dA}^t / \sigma_{dA}) / (\sigma_{pp}^t / \sigma_{pp})$  is the usual nuclear modification factor for trigger particles  $t$ , and  $\sigma$ ,  $\sigma^t$ , and  $\sigma^{\text{pair}}$  are the cross sections (or normalized yields) for the full event selection, trigger particle event selection,

and dihadron pair event selection.  $I_{dA}$  is the ratio of conditional hadron yields,  $CY$ , for  $d+Au$  and  $p+p$  collisions:

$$CY = \frac{\int d(\Delta\phi)[dN/d(\Delta\phi) - b_0]}{N^t \times \epsilon^a \times \Delta\eta^a \times \Delta p_T^a}, \quad (2)$$

with the acceptance corrected dihadron correlation function  $dN/d(\Delta\phi)$ , the number of trigger particles  $N^t$ , the detection efficiency for the associated particle  $\epsilon^a$  and the level of the uncorrelated pedestal in the correlation functions  $b_0$ . The integral is taken over the Gaussian fit of the away side peak. The  $J_{dA}$  uncertainties include a systematic uncertainty from the ZYAM pedestal subtraction. In determining this uncertainty it was assumed that changes between  $d+Au$  and  $p+p$  in the Gaussian away-side width remain below a factor two. This upper limit is based on the small observed changes in width in the mid-forward correlations and the correlations studied previously with the PHENIX muon spectrometers [14]. The  $J_{dA}$  is calculated from the measured  $I_{dA}$  and  $R_{dA}^t$  for the forward-rapidity trigger correlations with the new  $\pi^0$   $R_{dAu} = R_{dAu}^t$  determined in the MPC. For the midrapidity trigger correlations, published values for  $R_{dA}$  from the 2003 RHIC run [15, 16] were used.

Figure 3 presents  $J_{dA}$  versus  $\langle N_{coll} \rangle$  for forward-rapidity  $\pi^0$ 's paired with midrapidity hadrons and  $\pi^0$ 's, and for  $\pi^0$ 's and clusters paired at forward rapidity. The  $J_{dA}$  decreases with increasing number of binary collisions,  $\langle N_{coll} \rangle$ , or equivalently with increasing nuclear thickness. The suppression also increases with decreasing particle  $p_T$  and is significantly larger for forward-forward hadron pairs than for mid-forward pairs. The observed suppression of  $J_{dA}$  versus nuclear thickness,  $p_T$  and  $\eta$  points to large cold nuclear matter effects arising at high parton densities in the nucleus probed by the deuteron, consistent with predictions from CGC [12]. This trend is seen more clearly in Fig. 4 where  $J_{dA}$  is plotted versus  $x_{Au}^{frag} = (\langle p_{T1} \rangle e^{-\langle \eta_1 \rangle} + \langle p_{T2} \rangle e^{-\langle \eta_2 \rangle}) / \sqrt{s_{NN}}$  for all pair selections in  $\eta$  and  $p_T$ . In the case of  $2 \rightarrow 2$  parton scattering, where two final state hadrons carry the full parton energy,  $z=1$ , the variable  $x_{Au}^{frag}$  would be equal to  $\langle x_{Au} \rangle$ , which is the average momentum fraction of the struck parton in the Au nucleus.




FIG. 4: (color online)  $J_{dA}$  versus  $x_{Au}^{frag}$  for peripheral (60–88%) and central (0–20%)  $d+Au$  collisions at  $\sqrt{s_{NN}} = 200$  GeV. The statistical error bars and systematic uncertainty boxes are the same as in Fig. 3. Above  $x_{Au}^{frag} > 10^{-3}$ , some data points were offset from their true  $x_{Au}^{frag}$  to avoid overlap. The leftmost point in each group of three is at the correct  $x_{Au}^{frag}$ .

Because the fragmentation hadrons on average carry a momentum fraction  $\langle z \rangle < 1$ ,  $x_{Au}^{frag}$  will be smaller than  $\langle x_{Au} \rangle$ . Based on previous studies by PHENIX at midrapidity, the mean fragmentation  $\langle z \rangle$  is expected to be between 0.5–0.75 [22]. In general the theoretical extraction of  $x_{Au}$  from the measured  $p_T$  and  $\eta$  will differ from the leading order QCD picture of  $2 \rightarrow 2$  processes used above. Also, at modest  $p_T$ 's the interpretation of the measured correlation functions as high energy  $2 \rightarrow 2$  parton scattering accessing low  $x$  may be limited by contributions from processes with small momentum transfer  $Q^2$ . Future theoretical analysis will be necessary to evaluate these and other contributions

from different nuclear effects [4–10] on the observed large suppression in  $J_{dA}$ . These analyses could additionally be complicated by the presence of hadron pairs originating from multiparton interactions (MPI) [23] that might not probe gluon structure at low  $x_{Au}$ .

In summary, measurements of the inclusive  $\pi^0$  yield at forward rapidity, of the back-to-back correlated yield of cluster- $\pi^0$  pairs in the forward rapidity region, and of the correlated yield of forward rapidity  $\pi^0$ 's with midrapidity  $\pi^0$ 's or hadrons in  $p+p$  and  $d+Au$  collisions at  $\sqrt{s_{NN}} = 200$  GeV were presented. The correlated yields of back-to-back pairs were analyzed for various kinematic selections in  $p_T$  and rapidity. The forward-central pair measurements show no increase in the azimuthal angular correlation width within experimental uncertainties. The correlated yield of back-to-back pairs in  $d+Au$  collisions is observed to be substantially suppressed relative to  $p+p$  collisions with a suppression that is observed to increase with decreasing impact parameter selection and for pairs probing more forward rapidities.

We thank the staff of the Collider-Accelerator and Physics Departments at Brookhaven National Laboratory and the staff of the other PHENIX participating institutions for their vital contributions. We acknowledge support from the Office of Nuclear Physics in the Office of Science of the Department of Energy, the National Science Foundation, Abilene Christian University Research Council, Research Foundation of SUNY, and Dean of the College of Arts and Sciences, Vanderbilt University (U.S.A), Ministry of Education, Culture, Sports, Science, and Technology and the Japan Society for the Promotion of Science (Japan), Conselho Nacional de Desenvolvimento Científico e Tecnológico and Fundação de Amparo à Pesquisa do Estado de São Paulo (Brazil), Natural Science Foundation of China (P. R. China), Ministry of Education, Youth and Sports (Czech Republic), Centre National de la Recherche Scientifique, Commissariat à l'Énergie Atomique, and Institut National de Physique Nucléaire et de Physique des Particules (France), Ministry of Industry, Science and Technologies, Bundesministerium für Bildung und Forschung, Deutscher Akademischer Austausch Dienst, and Alexander von Humboldt Stiftung (Germany), Hungarian National Science Fund, OTKA (Hungary), Department of Atomic Energy and Department of Science and Technology (India), Israel Science Foundation (Israel), National Research Foundation and WCU program of the Ministry Education Science and Technology (Korea), Ministry of Education and Science, Russian Academy of Sciences, Federal Agency of Atomic Energy (Russia), VR and the Wallenberg Foundation (Sweden), the U.S. Civilian Research and Development Foundation for the Independent States of the Former Soviet Union, the US-Hungarian Fulbright Foundation for Educational Exchange, and the US-Israel Binational Science Foundation.

---

\* Deceased

† PHENIX Spokesperson: [jacak@skipper.physics.sunysb.edu](mailto:jacak@skipper.physics.sunysb.edu)

- [1] I. Arsene et al. (BRAHMS Collaboration), *Phys. Rev. Lett.* **93**, 242303 (2004).
- [2] J. Adams et al. (STAR Collaboration), *Phys. Rev. Lett.* **97**, 152302 (2006).
- [3] S. S. Adler et al. (PHENIX Collaboration), *Phys. Rev. Lett.* **94**, 082302 (2005).
- [4] I. Vitev, *Phys. Rev. C* **75**, 064906 (2007).
- [5] L. Frankfurt and M. Strikman, *Phys. Lett.* **B645**, 412 (2007).
- [6] R. C. Hwa, C. B. Yang, and R. J. Fries, *Phys. Rev. C* **71**, 024902 (2005).
- [7] V. Guzey, M. Strikman, and W. Vogelsang, *Phys. Lett.* **B603**, 173 (2004).
- [8] J. Qiu and I. Vitev, *Phys. Lett.* **B632**, 507 (2006).
- [9] L. Gribov, E. Levin, and M. Ryskin, *Phys.Rept.* **100**, 1 (1983).
- [10] L. McLerran and R. Venugopalan, *Phys. Rev. D* **49**, 3352 (1994).
- [11] D. Kharzeev, E. Levin, and L. McLerran, *Nucl. Phys.* **A748**, 627 (2005).
- [12] C. Marquet, *Nucl. Phys.* **A796**, 41 (2007).
- [13] S. S. Adler et al. (PHENIX Collaboration), *Phys. Rev. C* **73**, 054903 (2006).
- [14] S. S. Adler et al. (PHENIX Collaboration), *Phys. Rev. Lett.* **96**, 222301 (2006).
- [15] S. S. Adler et al. (PHENIX Collaboration), *Phys. Rev. C* **77**, 014905 (2008).
- [16] S. S. Adler et al. (PHENIX Collaboration), *Phys. Rev. Lett.* **98**, 172302 (2007).
- [17] R. Brun et al. (1987), CERN-DD-EE-84-1.
- [18] T. Sjostrand, S. Mrenna, and P. Z. Skands, *JHEP* **0605**, 026 (2006).
- [19] X.-N. Wang and M. Gyulassy, *Phys. Rev. D* **44**, 3501 (1991).
- [20] N. N. Ajitanand et al., *Phys. Rev. C* **72**, 011902 (2005).
- [21] A. Adare et al. (PHENIX Collaboration), *Phys. Rev. C* **78**, 014901 (2008).
- [22] S. S. Adler et al. (PHENIX Collaboration), *Phys. Rev. D* **74**, 072002 (2006).
- [23] M. Strikman and W. Vogelsang, *Phys. Rev. D* **83**, 034029 (2011).