

This is the accepted manuscript made available via CHORUS. The article has been published as:

Measurements of the Angular Distributions in the Decays $B \rightarrow K^{(*)} \mu^{+} \mu^{-}$ at CDF

T. Aaltonen *et al.* (CDF Collaboration)

Phys. Rev. Lett. **108**, 081807 — Published 24 February 2012

DOI: [10.1103/PhysRevLett.108.081807](https://doi.org/10.1103/PhysRevLett.108.081807)

Measurements of the Angular Distributions in the Decays $B \rightarrow K^{(*)}\mu^+\mu^-$ at CDF

T. Aaltonen,²² B. Álvarez González^{v,10} S. Amerio,⁴² D. Amidei,³³ A. Anastassov,³⁷ A. Annovi,¹⁸ J. Antos,¹³ G. Apollinari,¹⁶ J.A. Appel,¹⁶ A. Apresyan,⁴⁷ T. Arisawa,⁵⁶ A. Artikov,¹⁴ J. Asaadi,⁵² W. Ashmanskas,¹⁶ B. Auerbach,⁵⁹ A. Aurisano,⁵² F. Azfar,⁴¹ W. Badgett,¹⁶ A. Barbaro-Galtieri,²⁷ V.E. Barnes,⁴⁷ B.A. Barnett,²⁴ P. Barria^{ee,45} P. Bartos,¹³ M. Bauce^{cc,42} G. Bauer,³¹ F. Bedeschi,⁴⁵ D. Beecher,²⁹ S. Behari,²⁴ G. Bellettini^{dd,45} J. Bellinger,⁵⁸ D. Benjamin,¹⁵ A. Beretvas,¹⁶ A. Bhatti,⁴⁹ M. Binkley^{*,16} D. Bisello^{cc,42} I. Bizjak^{ii,29} K.R. Bland,⁵ C. Blocker,⁷ B. Blumenfeld,²⁴ A. Bocci,¹⁵ A. Bodek,⁴⁸ D. Bortoletto,⁴⁷ J. Boudreau,⁴⁶ A. Boveia,¹² B. Brau^{a,16} L. Brigliadori^{bb,6} A. Brisuda,¹³ C. Bromberg,³⁴ E. Brucken,²² M. Bucciantonio^{dd,45} J. Budagov,¹⁴ H.S. Budd,⁴⁸ S. Budd,²³ K. Burkett,¹⁶ G. Busetto^{cc,42} P. Bussey,²⁰ A. Buzatu,³² S. Cabrera^{x,15} C. Calancha,³⁰ S. Camarda,⁴ M. Campanelli,³⁴ M. Campbell,³³ F. Canelli^{12,16} A. Canepa,⁴⁴ B. Carls,²³ D. Carlsmith,⁵⁸ R. Carosi,⁴⁵ S. Carrillo^{k,17} S. Carron,¹⁶ B. Casal,¹⁰ M. Casarsa,¹⁶ A. Castro^{bb,6} P. Catastini,¹⁶ D. Cauz,⁵³ V. Cavaliere^{ee,45} M. Cavalli-Sforza,⁴ A. Cerri^{f,27} L. Cerrito^{q,29} Y.C. Chen,¹ M. Chertok,⁸ G. Chiarelli,⁴⁵ G. Chlachidze,¹⁶ F. Chlebana,¹⁶ K. Cho,²⁶ D. Chokheli,¹⁴ J.P. Chou,²¹ W.H. Chung,⁵⁸ Y.S. Chung,⁴⁸ C.I. Ciobanu,⁴³ M.A. Ciocci^{ee,45} A. Clark,¹⁹ D. Clark,⁷ G. Compostella^{cc,42} M.E. Convery,¹⁶ J. Conway,⁸ M. Corbo,⁴³ M. Cordelli,¹⁸ C.A. Cox,⁸ D.J. Cox,⁸ F. Crescioli^{dd,45} C. Cuenca Almenar,⁵⁹ J. Cuevas^{v,10} R. Culbertson,¹⁶ D. Dagenhart,¹⁶ N. d'Ascenzo^{t,43} M. Datta,¹⁶ P. de Barbaro,⁴⁸ S. De Cecco,⁵⁰ G. De Lorenzo,⁴ M. Dell'Orso^{dd,45} C. Deluca,⁴ L. Demortier,⁴⁹ J. Deng^{c,15} M. Deninno,⁶ F. Devoto,²² M. d'Errico^{cc,42} A. Di Canto^{dd,45} B. Di Ruzza,⁴⁵ J.R. Dittmann,⁵ M. D'Onofrio,²⁸ S. Donati^{dd,45} P. Dong,¹⁶ T. Dorigo,⁴² K. Ebina,⁵⁶ A. Elagin,⁵² A. Eppig,³³ R. Erbacher,⁸ D. Errede,²³ S. Errede,²³ N. Ershaidat^{aa,43} R. Eusebi,⁵² H.C. Fang,²⁷ S. Farrington,⁴¹ M. Feindt,²⁵ J.P. Fernandez,³⁰ C. Ferrazza^{ff,45} R. Field,¹⁷ G. Flanagan^{r,47} R. Forrest,⁸ M.J. Frank,⁵ M. Franklin,²¹ J.C. Freeman,¹⁶ I. Furic,¹⁷ M. Gallinaro,⁴⁹ J. Galyardt,¹¹ J.E. Garcia,¹⁹ A.F. Garfinkel,⁴⁷ P. Garosi^{ee,45} H. Gerberich,²³ E. Gerchtein,¹⁶ S. Giagu^{gg,50} V. Giakoumopoulou,³ P. Giannetti,⁴⁵ K. Gibson,⁴⁶ C.M. Ginsburg,¹⁶ N. Giokaris,³ P. Giromini,¹⁸ M. Giunta,⁴⁵ G. Giurgiu,²⁴ V. Glagolev,¹⁴ D. Glenzinski,¹⁶ M. Gold,³⁶ D. Goldin,⁵² N. Goldschmidt,¹⁷ A. Golossanov,¹⁶ G. Gomez,¹⁰ G. Gomez-Ceballos,³¹ M. Goncharov,³¹ O. González,³⁰ I. Gorelov,³⁶ A.T. Goshaw,¹⁵ K. Goulianos,⁴⁹ A. Gresele,⁴² S. Grinstein,⁴ C. Grosso-Pilcher,¹² R.C. Group,¹⁶ J. Guimaraes da Costa,²¹ Z. Gunay-Unalan,³⁴ C. Haber,²⁷ S.R. Hahn,¹⁶ E. Halkiadakis,⁵¹ A. Hamaguchi,⁴⁰ J.Y. Han,⁴⁸ F. Happacher,¹⁸ K. Hara,⁵⁴ D. Hare,⁵¹ M. Hare,⁵⁵ R.F. Harr,⁵⁷ K. Hatakeyama,⁵ C. Hays,⁴¹ M. Heck,²⁵ J. Heinrich,⁴⁴ M. Herndon,⁵⁸ S. Hewamanage,⁵ D. Hidas,⁵¹ A. Hocker,¹⁶ W. Hopkins^{g,16} D. Horn,²⁵ S. Hou,¹ R.E. Hughes,³⁸ M. Hurwitz,¹² U. Husemann,⁵⁹ N. Hussain,³² M. Hussein,³⁴ J. Huston,³⁴ G. Introzzi,⁴⁵ M. Iori^{gg,50} A. Ivanov^{o,8} E. James,¹⁶ D. Jang,¹¹ B. Jayatilaka,¹⁵ E.J. Jeon,²⁶ M.K. Jha,⁶ S. Jindariani,¹⁶ W. Johnson,⁸ M. Jones,⁴⁷ K.K. Joo,²⁶ S.Y. Jun,¹¹ T.R. Junk,¹⁶ T. Kamon,⁵² P.E. Karchin,⁵⁷ Y. Kato^{n,40} W. Ketchum,¹² J. Keung,⁴⁴ V. Khotilovich,⁵² B. Kilminster,¹⁶ D.H. Kim,²⁶ H.S. Kim,²⁶ H.W. Kim,²⁶ J.E. Kim,²⁶ M.J. Kim,¹⁸ S.B. Kim,²⁶ S.H. Kim,⁵⁴ Y.K. Kim,¹² N. Kimura,⁵⁶ S. Klimenko,¹⁷ K. Kondo,⁵⁶ D.J. Kong,²⁶ J. Konigsberg,¹⁷ A. Korytov,¹⁷ A.V. Kotwal,¹⁵ M. Kreps,²⁵ J. Kroll,⁴⁴ D. Krop,¹² N. Krumnack^{l,5} M. Kruse,¹⁵ V. Krutiyov^{d,52} T. Kuhr,²⁵ M. Kurata,⁵⁴ S. Kwang,¹² A.T. Laasanen,⁴⁷ S. Lami,⁴⁵ S. Lammel,¹⁶ M. Lancaster,²⁹ R.L. Lander,⁸ K. Lannon^{u,38} A. Lath,⁵¹ G. Latino^{ee,45} I. Lazzizzera,⁴² T. LeCompte,² E. Lee,⁵² H.S. Lee,¹² J.S. Lee,²⁶ S.W. Lee^{w,52} S. Leo^{dd,45} S. Leone,⁴⁵ J.D. Lewis,¹⁶ C.-J. Lin,²⁷ J. Linacre,⁴¹ M. Lindgren,¹⁶ E. Lipeles,⁴⁴ A. Lister,¹⁹ D.O. Litvintsev,¹⁶ C. Liu,⁴⁶ Q. Liu,⁴⁷ T. Liu,¹⁶ S. Lockwitz,⁵⁹ N.S. Lockyer,⁴⁴ A. Loginov,⁵⁹ D. Lucchesi^{cc,42} J. Lueck,²⁵ P. Lujan,²⁷ P. Lukens,¹⁶ G. Lungu,⁴⁹ J. Lys,²⁷ R. Lysak,¹³ R. Madrak,¹⁶ K. Maeshima,¹⁶ K. Makhoul,³¹ P. Maksimovic,²⁴ S. Malik,⁴⁹ G. Manca^{b,28} A. Manousakis-Katsikakis,³ F. Margaroli,⁴⁷ C. Marino,²⁵ M. Martínez,⁴ R. Martínez-Ballarín,³⁰ P. Mastrandrea,⁵⁰ M. Mathis,²⁴ M.E. Mattson,⁵⁷ P. Mazzanti,⁶ K.S. McFarland,⁴⁸ P. McIntyre,⁵² R. McNulty^{i,28} A. Mehta,²⁸ P. Mehtala,²² A. Menzione,⁴⁵ C. Mesropian,⁴⁹ T. Miao,¹⁶ D. Mietlicki,³³ A. Mitra,¹ H. Miyake,⁵⁴ S. Moed,²¹ N. Moggi,⁶ M.N. Mondragon^{k,16} C.S. Moon,²⁶ R. Moore,¹⁶ M.J. Morello,¹⁶ J. Morlock,²⁵ P. Movilla Fernandez,¹⁶ A. Mukherjee,¹⁶ Th. Muller,²⁵ P. Murat,¹⁶ M. Mussini^{bb,6} J. Nachtman^{m,16} Y. Nagai,⁵⁴ J. Naganoma,⁵⁶ I. Nakano,³⁹ A. Napier,⁵⁵ J. Nett,⁵⁸ C. Neu^{z,44} M.S. Neubauer,²³ J. Nielsen^{e,27} L. Nodulman,² O. Norniella,²³ E. Nurse,²⁹ L. Oakes,⁴¹ S.H. Oh,¹⁵ Y.D. Oh,²⁶ I. Oksuzian,¹⁷ T. Okusawa,⁴⁰ R. Orava,²² L. Ortolan,⁴ S. Pagan Griso^{cc,42} C. Pagliarone,⁵³ E. Palencia^{f,10} V. Papadimitriou,¹⁶ A.A. Paramonov,² J. Patrick,¹⁶ G. Pauletta^{hh,53} M. Paulini,¹¹ C. Paus,³¹ D.E. Pellett,⁸ A. Penzo,⁵³ T.J. Phillips,¹⁵ G. Piacentino,⁴⁵ E. Pianori,⁴⁴ J. Pilot,³⁸ K. Pitts,²³ C. Plager,⁹ L. Pondrom,⁵⁸ K. Potamianos,⁴⁷ O. Poukhov^{*,14} F. Prokoshin^{y,14} A. Pronko,¹⁶ F. Ptohos^{h,18} E. Pueschel,¹¹ G. Punzi^{dd,45} J. Pursley,⁵⁸ A. Rahaman,⁴⁶ V. Ramakrishnan,⁵⁸ N. Ranjan,⁴⁷ I. Redondo,³⁰ P. Renton,⁴¹ M. Rescigno,⁵⁰

F. Rimondi^{bb},⁶ L. Ristori⁴⁵,¹⁶ A. Robson,²⁰ T. Rodrigo,¹⁰ T. Rodriguez,⁴⁴ E. Rogers,²³ S. Rolli,⁵⁵ R. Roser,¹⁶ M. Rossi,⁵³ F. Ruffini^{ee},⁴⁵ A. Ruiz,¹⁰ J. Russ,¹¹ V. Rusu,¹⁶ A. Safonov,⁵² W.K. Sakumoto,⁴⁸ L. Santi^{hh},⁵³ L. Sartori,⁴⁵ K. Sato,⁵⁴ V. Saveliev^t,⁴³ A. Savoy-Navarro,⁴³ P. Schlabach,¹⁶ A. Schmidt,²⁵ E.E. Schmidt,¹⁶ M.P. Schmidt*,⁵⁹ M. Schmitt,³⁷ T. Schwarz,⁸ L. Scodellaro,¹⁰ A. Scribano^{ee},⁴⁵ F. Scuri,⁴⁵ A. Sedov,⁴⁷ S. Seidel,³⁶ Y. Seiya,⁴⁰ A. Semenov,¹⁴ F. Sforza^{dd},⁴⁵ A. Sfyrila,²³ S.Z. Shalhout,⁸ T. Shears,²⁸ P.F. Shepard,⁴⁶ M. Shimojima^s,⁵⁴ S. Shiraishi,¹² M. Shochet,¹² I. Shreyber,³⁵ A. Simonenko,¹⁴ P. Sinervo,³² A. Sissakian*,¹⁴ K. Sliwa,⁵⁵ J.R. Smith,⁸ F.D. Snider,¹⁶ A. Soha,¹⁶ S. Somalwar,⁵¹ V. Sorin,⁴ P. Squillacioti,¹⁶ M. Stanitzki,⁵⁹ R. St. Denis,²⁰ B. Stelzer,³² O. Stelzer-Chilton,³² D. Stentz,³⁷ J. Strologas,³⁶ G.L. Strycker,³³ Y. Sudo,⁵⁴ A. Sukhanov,¹⁷ I. Suslov,¹⁴ K. Takemasa,⁵⁴ Y. Takeuchi,⁵⁴ J. Tang,¹² M. Tecchio,³³ P.K. Teng,¹ J. Thom^g,¹⁶ J. Thome,¹¹ G.A. Thompson,²³ E. Thomson,⁴⁴ P. Ttito-Guzmán,³⁰ S. Tkaczyk,¹⁶ D. Toback,⁵² S. Tokar,¹³ K. Tollefson,³⁴ T. Tomura,⁵⁴ D. Tonelli,¹⁶ S. Torre,¹⁸ D. Torretta,¹⁶ P. Totaro^{hh},⁵³ M. Trovato^{ff},⁴⁵ Y. Tu,⁴⁴ N. Turini^{ee},⁴⁵ F. Ukegawa,⁵⁴ S. Uozumi,²⁶ A. Varganov,³³ E. Vataga^{ff},⁴⁵ F. Vázquez^k,¹⁷ G. Velez,¹⁶ C. Vellidis,³ M. Vidal,³⁰ I. Vila,¹⁰ R. Vilar,¹⁰ M. Vogel,³⁶ G. Volpi^{dd},⁴⁵ P. Wagner,⁴⁴ R.L. Wagner,¹⁶ T. Wakisaka,⁴⁰ R. Wallny,⁹ S.M. Wang,¹ A. Warburton,³² D. Waters,²⁹ M. Weinberger,⁵² H. Wenzel,¹⁶ W.C. Wester III,¹⁶ B. Whitehouse,⁵⁵ D. Whiteson^c,⁴⁴ A.B. Wicklund,² E. Wicklund,¹⁶ S. Wilbur,¹² F. Wick,²⁵ H.H. Williams,⁴⁴ J.S. Wilson,³⁸ P. Wilson,¹⁶ B.L. Winer,³⁸ P. Wittich^g,¹⁶ S. Wolbers,¹⁶ H. Wolfe,³⁸ T. Wright,³³ X. Wu,¹⁹ Z. Wu,⁵ K. Yamamoto,⁴⁰ J. Yamaoka,¹⁵ T. Yang,¹⁶ U.K. Yang^p,¹² Y.C. Yang,²⁶ W.-M. Yao,²⁷ G.P. Yeh,¹⁶ K. Yi^m,¹⁶ J. Yoh,¹⁶ K. Yorita,⁵⁶ T. Yoshida^j,⁴⁰ G.B. Yu,¹⁵ I. Yu,²⁶ S.S. Yu,¹⁶ J.C. Yun,¹⁶ A. Zanetti,⁵³ Y. Zeng,¹⁵ and S. Zucchelli^{bb6}

(CDF Collaboration[†])

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

²*Argonne National Laboratory, Argonne, Illinois 60439, USA*

³*University of Athens, 157 71 Athens, Greece*

⁴*Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

⁵*Baylor University, Waco, Texas 76798, USA*

⁶*Istituto Nazionale di Fisica Nucleare Bologna, ^{bb}University of Bologna, I-40127 Bologna, Italy*

⁷*Brandeis University, Waltham, Massachusetts 02254, USA*

⁸*University of California, Davis, Davis, California 95616, USA*

⁹*University of California, Los Angeles, Los Angeles, California 90024, USA*

¹⁰*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

¹¹*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

¹²*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*

¹³*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

¹⁴*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

¹⁵*Duke University, Durham, North Carolina 27708, USA*

¹⁶*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*

¹⁷*University of Florida, Gainesville, Florida 32611, USA*

¹⁸*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

¹⁹*University of Geneva, CH-1211 Geneva 4, Switzerland*

²⁰*Glasgow University, Glasgow G12 8QQ, United Kingdom*

²¹*Harvard University, Cambridge, Massachusetts 02138, USA*

²²*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*

²³*University of Illinois, Urbana, Illinois 61801, USA*

²⁴*The Johns Hopkins University, Baltimore, Maryland 21218, USA*

²⁵*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*

²⁶*Center for High Energy Physics: Kyungpook National University,*

Daegu 702-701, Korea; Seoul National University, Seoul 151-742,

Korea; Sungkyunkwan University, Suwon 440-746,

Korea; Korea Institute of Science and Technology Information,

Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757,

Korea; Chonbuk National University, Jeonju 561-756, Korea

²⁷*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

²⁸*University of Liverpool, Liverpool L69 7ZE, United Kingdom*

²⁹*University College London, London WC1E 6BT, United Kingdom*

³⁰*Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain*

³¹*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

³²*Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia,*

- Canada V5A 1S6; University of Toronto, Toronto, Ontario,
Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
- ³³University of Michigan, Ann Arbor, Michigan 48109, USA
- ³⁴Michigan State University, East Lansing, Michigan 48824, USA
- ³⁵Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- ³⁶University of New Mexico, Albuquerque, New Mexico 87131, USA
- ³⁷Northwestern University, Evanston, Illinois 60208, USA
- ³⁸The Ohio State University, Columbus, Ohio 43210, USA
- ³⁹Okayama University, Okayama 700-8530, Japan
- ⁴⁰Osaka City University, Osaka 588, Japan
- ⁴¹University of Oxford, Oxford OX1 3RH, United Kingdom
- ⁴²Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, ^{cc}University of Padova, I-35131 Padova, Italy
- ⁴³LPNHE, Universite Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
- ⁴⁴University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁴⁵Istituto Nazionale di Fisica Nucleare Pisa, ^{dd}University of Pisa,
- ^{ee}University of Siena and ^{ff}Scuola Normale Superiore, I-56127 Pisa, Italy
- ⁴⁶University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
- ⁴⁷Purdue University, West Lafayette, Indiana 47907, USA
- ⁴⁸University of Rochester, Rochester, New York 14627, USA
- ⁴⁹The Rockefeller University, New York, New York 10065, USA
- ⁵⁰Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1,
- ⁹⁹Sapienza Università di Roma, I-00185 Roma, Italy
- ⁵¹Rutgers University, Piscataway, New Jersey 08855, USA
- ⁵²Texas A&M University, College Station, Texas 77843, USA
- ⁵³Istituto Nazionale di Fisica Nucleare Trieste/Udine,
I-34100 Trieste, ^{hh}University of Trieste/Udine, I-33100 Udine, Italy
- ⁵⁴University of Tsukuba, Tsukuba, Ibaraki 305, Japan
- ⁵⁵Tufts University, Medford, Massachusetts 02155, USA
- ⁵⁶Waseda University, Tokyo 169, Japan
- ⁵⁷Wayne State University, Detroit, Michigan 48201, USA
- ⁵⁸University of Wisconsin, Madison, Wisconsin 53706, USA
- ⁵⁹Yale University, New Haven, Connecticut 06520, USA

We report an indirect search for non-standard model physics using the flavor-changing neutral current decays $B \rightarrow K^{(*)}\mu^+\mu^-$. We reconstruct the decays and measure their angular distributions, as a function of $q^2 = M_{\mu\mu}^2$, where $M_{\mu\mu}$ is the dimuon mass, in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using a data sample corresponding to an integrated luminosity of 6.8 fb^{-1} . The transverse polarization asymmetry $A_T^{(2)}$ and the time-reversal-odd charge-and-parity asymmetry A_{im} are measured for the first time, together with the K^* longitudinal polarization fraction F_L and the muon forward-backward asymmetry A_{FB} for the decays $B^0 \rightarrow K^{*0}\mu^+\mu^-$ and $B^+ \rightarrow K^{*+}\mu^+\mu^-$. The $B \rightarrow K^{*}\mu^+\mu^-$ forward-backward asymmetry in the most sensitive kinematic regime, $1 \leq q^2 < 6 \text{ GeV}^2/c^2$, is measured to be $A_{FB} = 0.29_{-0.23}^{+0.20}(\text{stat}) \pm 0.07(\text{syst})$, the most precise result to date. No deviations from the standard model predictions are observed.

PACS numbers: 13.25 Hw, 13.20 He

The decays $B \rightarrow K^{(*)}\mu^+\mu^-$ [1], which proceed via the flavor-changing neutral current process $b \rightarrow s\mu^+\mu^-$, are among the most promising probes of the standard

*Deceased

[†]With visitors from ^aUniversity of Massachusetts Amherst, Amherst, Massachusetts 01003, ^bIstituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, ^cUniversity of California Irvine, Irvine, CA 92697, ^dUniversity of California Santa Barbara, Santa Barbara, CA 93106 ^eUniversity of California Santa Cruz, Santa Cruz, CA 95064, ^fCERN, CH-1211 Geneva, Switzerland, ^gCornell University, Ithaca, NY 14853, ^hUniversity of Cyprus, Nicosia CY-1678, Cyprus, ⁱUniversity College Dublin, Dublin 4, Ireland, ^jUniversity of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017, ^kUniversidad Iberoamericana, Mexico D.F., Mexico, ^lIowa State University, Ames, IA 50011, ^mUniversity of Iowa, Iowa City, IA 52242, ⁿKinki University, Higashi-Osaka City, Japan 577-8502, ^oKansas State University, Manhattan, KS 66506, ^pUniversity of Manchester, Manchester M13

9PL, England, ^qQueen Mary, University of London, London, E1 4NS, England, ^rMuons, Inc., Batavia, IL 60510, ^sNagasaki Institute of Applied Science, Nagasaki, Japan, ^tNational Research Nuclear University, Moscow, Russia, ^uUniversity of Notre Dame, Notre Dame, IN 46556, ^vUniversidad de Oviedo, E-33007 Oviedo, Spain, ^wTexas Tech University, Lubbock, TX 79609, ^xIFIC(CSIC-Universitat de Valencia), 56071 Valencia, Spain, ^yUniversidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile, ^zUniversity of Virginia, Charlottesville, VA 22906, ^{aa}Yarmouk University, Irbid 211-63, Jordan, ⁱⁱOn leave from J. Stefan Institute, Ljubljana, Slovenia,

model (SM) and its extensions [2]. These decays are suppressed in the SM since they occur through higher order amplitudes involving quantum loops. Such amplitudes can receive competing contributions from unknown massive virtual particles or new couplings, which may significantly alter the decay kinematics with respect to the SM predictions. Multi-body final states further enrich the phenomenology of these decays, offering sensitivity to a broad class of SM extensions through measurement of angular distributions as a function of $q^2 \equiv M_{\mu\mu}^2 c^2$, where $M_{\mu\mu}$ is the dimuon invariant mass.

The full differential decay distribution for decays $B \rightarrow K^*(892)\mu^+\mu^- \rightarrow K\pi\mu^+\mu^-$ is described in terms of four kinematic variables: the angle θ_μ between the μ^+ (μ^-) direction and the direction opposite to the B (\bar{B}) meson in the dimuon rest frame, the angle θ_K between the kaon direction and the direction opposite to the B meson in the K^* rest frame, the angle ϕ between the two planes formed by the dimuon and the K - π systems, and q^2 . The decay distribution is a function of eight independent angular coefficients, each functions of q^2 , that are physics observables to be determined experimentally. A simultaneous determination of all the coefficients requires signal event samples of much larger size than currently available. Following [3–6], we project the decay distribution into three simpler relations each involving just one of the angles:

$$\begin{aligned} \frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta_K} &= \frac{3}{2} F_L \cos^2\theta_K + \frac{3}{4} (1 - F_L)(1 - \cos^2\theta_K), \\ \frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta_\mu} &= \frac{3}{4} F_L (1 - \cos^2\theta_\mu) + \frac{3}{8} (1 - F_L)(1 + \cos^2\theta_\mu) \\ &\quad + A_{FB} \cos\theta_\mu, \\ \frac{1}{\Gamma} \frac{d\Gamma}{d\phi} &= \frac{1}{2\pi} \left[1 + \frac{1}{2} (1 - F_L) A_T^{(2)} \cos 2\phi + A_{im} \sin 2\phi \right], \end{aligned} \quad (1)$$

where Γ is the partial decay width. These one-dimensional relations are functions of a subset of four angular observables, each of which depends on q^2 : A_{FB} , the muon forward-backward asymmetry; F_L , the K^* longitudinal polarization fraction; $A_T^{(2)}$, the transverse polarization asymmetry [3, 5]; and A_{im} , the time-reversal-odd charge-and-parity asymmetry (T -odd CP asymmetry) [4, 6].

BABAR [7], Belle [8], and CDF [9] have reported measurements of A_{FB} and F_L in the $B \rightarrow K^*\ell^+\ell^-$ decay modes. All experiments find A_{FB} to be larger than the SM expectation, but so far none has had sufficient sensitivity to be conclusive. However, Belle reports that the cumulative difference of their measurement from the SM prediction corresponds to 2.7σ [8]. With the current sample, CDF has sensitivity comparable to Belle's. A similar discrepancy in CDF data would provide a strong indication of physics beyond the standard model (BSM).

In this Letter, we report measurements of F_L , A_{FB} , and for the first time use the angle ϕ to access $A_T^{(2)}$ and

A_{im} in the decay $B \rightarrow K^*\mu^+\mu^-$. We use a sample of $p\bar{p}$ collisions at a center-of-mass energy of $\sqrt{s} = 1.96$ TeV, corresponding to 6.8 fb^{-1} integrated luminosity collected with the CDF II detector. The measurement updates and supersedes an earlier analysis [9]. We include the $B^+ \rightarrow K^{*+}\mu^+\mu^-$ decay, reconstructed for the first time in hadronic collisions, and improve the $B^0 \rightarrow K^{*0}\mu^+\mu^-$ signal selection by 9% with almost the same background rejection, along with the added data. The resulting 82% increase in signal statistics reduces the uncertainties on A_{FB} and F_L by 50% and enables the first measurement of $A_T^{(2)}$ and A_{im} . We also update the measurement of A_{FB} in $B^+ \rightarrow K^+\mu^+\mu^-$ decays. Measurements of branching ratios are reported in another Letter [10].

The differential decay distribution of $B \rightarrow K^*\mu^+\mu^-$ is calculated in an operator product expansion [11]. In the SM, only the Wilson coefficients $C_{7,9,10}^{\text{eff}}$ are relevant. Each of the angular observables, which include different combinations of Wilson coefficients, has unique sensitivity to specific features of BSM models. The measurement of F_L could constrain BSM models in which couplings to the K^* helicity states are different from the SM ones [3]. The observables A_{FB} , $A_T^{(2)}$, and A_{im} are especially sensitive since the hadronic uncertainties cancel in the asymmetry. The asymmetry of the muon direction between the forward ($\cos\theta_\mu > 0$) and backward ($\cos\theta_\mu < 0$) directions in the dimuon rest frame, A_{FB} , is expected to be small at low q^2 and large and positive at high q^2 in the SM. The BSM contributions can change the magnitude and the sign of A_{FB} . The observables $A_T^{(2)}$ and A_{im} have recently been proposed as powerful probes of SM extensions that involve weak interactions of particles with positive chirality (right-handed currents).

In the SM $A_T^{(2)}$ is accurately predicted to be approximately zero at low q^2 and negative at high q^2 [12]. However a broad class of BSM models, which involve the CP -conserving right-handed currents, expect $A_T^{(2)}$ reaching values up to ± 1 [3, 5, 13]. The SM predicts A_{im} to be close to zero for all accessible values of q^2 . Deviations could be observed in the case of CP -violating right-handed currents [6].

The $B^+ \rightarrow K^+\mu^+\mu^-$ angular distribution is simpler than the $B \rightarrow K^*\mu^+\mu^-$ distribution. Although A_{FB} is the only observable accessible in this decay, it provides complementary sensitivity to BSM models with respect to $B \rightarrow K^*\mu^+\mu^-$ measurements. The asymmetry expected in the SM is quite small over the entire range of q^2 [14], but could be enhanced if scalar- or tensor-type BSM contributions are present [13].

The reconstruction of the $B \rightarrow K^{(*)}\mu^+\mu^-$ decays starts with a dimuon sample selected by the online trigger system [15] of the CDF II detector [16]. The trigger uses information from muon detectors and the central drift tracking chamber [17]. The chamber provides 96 samplings of charged-particle's trajectories (tracks)

between radii of 40 and 137 cm, allowing an accurate determination of particles' momenta. The CMU and CMX muon drift chambers [18] cover the pseudorapidity regions $|\eta| < 0.6$ and $0.6 < |\eta| < 1.0$, respectively [19]. The CMP muon system is located radially behind the CMU and an additional steel absorber, and covers $|\eta| < 0.6$. The dimuon trigger requires a pair of oppositely charged particles with momenta transverse to the beamline $p_T > 1.5 \text{ GeV}/c$, that are also identified in the CMU or CMX chambers. At least one of the muon tracks in the pair is required to be a CMU muon. The trigger also requires either of the following criteria; the dimuon pair satisfies $L_{xy} > 100 \mu\text{m}$, where the transverse decay length L_{xy} is the flight distance between the dimuon vertex and the event primary vertex projected onto the dimuon momentum vector; or one of the muon candidates has $p_T > 3.0 \text{ GeV}/c$ and is identified by both the CMU and CMP chambers. The other detector subsystems relevant for this analysis are discussed in Ref. [20].

Each offline track is required to satisfy the standard quality requirements to ensure well measured momenta and decay vertices [9]. The decay length and mass of each dimuon pair are calculated after a vertex fit. Dimuons are required to have q^2 values outside the ranges of $8.68 < q^2 < 10.09 \text{ GeV}^2/c^2$ and $12.86 < q^2 < 14.18 \text{ GeV}^2/c^2$ [9], to reject J/ψ and $\psi(2S)$ decays, typically reconstructed with $14 \text{ MeV}/c^2$ mass resolution. The dimuon pair is then combined with tracks forming a $K^{*0} \rightarrow K^+\pi^-$ candidate to form a $B^0 \rightarrow K^{*0}\mu^+\mu^-$ candidate, or a $K^{*+} \rightarrow K_S^0(\rightarrow \pi^+\pi^-)\pi^+$ candidate to form a $B^+ \rightarrow K^{*+}\mu^+\mu^-$ candidate. Loose particle identification requirements are applied to the K^+ candidate [10]. The K_S^0 , K^{*0} , and K^{*+} candidates are required to have masses consistent with the known values [21], and to have $p_T > 1 \text{ GeV}/c$. The K_S^0 is also required to decay in a vertex displaced from the dimuon vertex. In the K^{*0} reconstruction, we choose the $K^+\pi^-$ assignment that yields the mass closer to the known K^{*0} mass. This is correct 92% of the times, as shown by simulation. The reconstructed B candidates are required to have $p_T > 4 \text{ GeV}/c$. To further optimize the selection, a neural network (NN) classifier [22] is trained for each channel using simulated signal and background sampled from B mass sidebands ($0.1\text{--}0.36 \text{ GeV}/c^2$ higher than the known B mass [21]) in data. The optimized NN threshold is determined to minimize the statistical uncertainty of the angular observables, using 15-18 discriminating observables including p_T of B meson and daughter particles, K^* and K_S^0 masses, vertex fit parameters, and muon identification quality [9]. No angular bias due to the choice of the NN threshold is found in simulation or in $B \rightarrow J/\psi(\rightarrow \mu^+\mu^-)K^{(*)}$ control samples.

The signal yield is obtained from an unbinned maximum likelihood fit to the mass distribution with a probability density function (PDF) consisting of Gaussian




FIG. 1: Invariant mass of $B^0 \rightarrow K^{*0}\mu^+\mu^-$ (left) and $B^+ \rightarrow K^{*+}\mu^+\mu^-$ (right), with fit results overlaid.

distributions determined from simulation for the signal and a linear background. The $5.18\text{--}5.70 \text{ GeV}/c^2$ fit range avoids the $5.0\text{--}5.18 \text{ GeV}/c^2$ region, dominated by background from multi-body B decays. The signal region is defined within $\pm 40 \text{ MeV}/c^2$ from the known B mass [21]. In this range, the contributions from similar decays such as $B^0 \rightarrow \rho^0\mu^+\mu^-$, and $B^+ \rightarrow \pi^+\mu^+\mu^-$ are negligible due to mis-reconstruction and rates smaller than our signal rates. The 1% $B_s^0 \rightarrow \phi\mu^+\mu^-$ contamination to the $B^0 \rightarrow K^{*0}\mu^+\mu^-$ signal is included in the systematic uncertainties.

We find 234 ± 19 $B^+ \rightarrow K^+\mu^+\mu^-$, 164 ± 15 $B^0 \rightarrow K^{*0}\mu^+\mu^-$, and 20 ± 6 $B^+ \rightarrow K^{*+}\mu^+\mu^-$ events (Fig. 1). The uncertainties are statistical only. We divide the events into six bins in q^2 . Two semi-inclusive bins are included with ranges covering theoretically well-controlled regions. We obtain the signal yields (Tabs. I, II, and III) in the individual q^2 ranges by fitting the mass as described above.

To extract the quantities F_L , A_{FB} , $A_T^{(2)}$, and A_{im} , we perform likelihood fits to the distributions of $\cos\theta_K$, $\cos\theta_\mu$, and ϕ for candidates in each q^2 range and with B mass in the signal region. The signal fractions are fixed to the results of the mass fits. Signal PDFs for angular distributions are formed from Eq. (1), including detector acceptance and the K - π interchange, estimated using simulation. The incorrect K - π assignment in the $K^{*0} \rightarrow K^+\pi^-$ decay distorts the signal mass distribution and swaps the sign of $\cos\theta_\mu$. This is modeled with an additional signal-like term in the likelihood. The contribution from decays with non-resonant K - π pairs is expected to be small [3] and neglected in the fit. The angular PDF for the background are modeled using the same B mass sidebands used for NN training.

The values of F_L in individual q^2 ranges are extracted from fits to the $\cos\theta_K$ distributions and then used as fixed inputs in the fits of A_{FB} , $A_T^{(2)}$, and A_{im} . The asymmetry A_{FB} ($A_T^{(2)}$ and A_{im}) is obtained from fits to the $\cos\theta_\mu$ (ϕ) distributions.




FIG. 2: Measurements of forward-backward asymmetry A_{FB} in the decay (a) $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ and (b) $B^+ \rightarrow K^{*+} \mu^+ \mu^-$ as a function of dimuon mass squared q^2 . Points are the fit results from data. The solid curves are the SM expectation [23]. The dotted curve is the $C_7 = -C_7^{\text{SM}}$ expectation suggested by some BSM models. Hatched regions are excluded resonant (charmonium) decay regions.

The $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ results are listed in Table I, and the forward-backward asymmetry A_{FB} is illustrated in Fig. 2(a) as a function of q^2 . To increase sensitivity, we also fit the combined $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ and $B^+ \rightarrow K^{*+} \mu^+ \mu^-$ modes (Table II and Fig. 3), assuming they have the same decay dynamics. The determinations of A_{FB} and F_L are among the most precise from a single experiment. However, the shape of the A_{FB} distribution as a function of q^2 , which carries the majority of the discriminating information [12], does not deviate from the SM expectation and cannot yet provide conclusive discrimination between the SM and the inverted C_7 scenario which is suggested by the Belle results [8]. In the most sensitive kinematic regime, $1 \leq q^2 < 6 \text{ GeV}^2/c^2$, we find $A_{FB} = 0.29^{+0.20}_{-0.23}(\text{stat}) \pm 0.07(\text{syst})$, consistent with the SM prediction of $A_{FB}^{\text{SM}} = 0.022 \pm 0.028$ [5], and the Belle result of $A_{FB}^{\text{Belle}} = 0.26^{+0.27}_{-0.30}(\text{stat}) \pm 0.07(\text{syst})$ [8]. The polarization in the same q^2 range, $F_L = 0.69^{+0.19}_{-0.21}(\text{stat}) \pm 0.08(\text{syst})$, is also consistent with the corresponding SM prediction, $F_L^{\text{SM}} = 0.73^{+0.02}_{-0.03}$ [5], and the Belle result of $F_L^{\text{Belle}} = 0.67 \pm 0.23(\text{stat}) \pm 0.05(\text{syst})$ [8].

Tables I and II and Fig. 3(c, d) show the results of the first measurement of $A_T^{(2)}$ and A_{im} . These results explore new regions of BSM parameter space providing, in combination with other observables, initial discriminating information between different classes of BSM models.

In the $B^+ \rightarrow K^{*+} \mu^+ \mu^-$ fit for A_{FB} we assume no scalar term [14] and set $F_L = 1$ in Eq. 1. The results are the most precise from a single experiment and are consistent with the SM predictions (Fig. 2(b) and Tab. III).

The sources of systematic uncertainty include the estimation of detector acceptance, signal fraction estimation and shape modeling of events in the signal window, feed-down background from other B decays, trigger efficiency and bias modeling, incorrect K - π assignment in the $K^{*0} \rightarrow K^+ \pi^-$ decay, and fitting bias. The largest




FIG. 3: Measurements of (a) longitudinal K^* polarization fraction F_L , (b) forward-backward asymmetry A_{FB} , (c) transverse polarization asymmetry $A_T^{(2)}$, and (d) T -odd CP asymmetry A_{im} in the combined decay mode $B \rightarrow K^* \mu^+ \mu^-$.

contributions to A_{FB} , F_L , and A_{im} are from uncertainties on the signal fraction in the signal window. For the A_{FB} , $A_T^{(2)}$, and A_{im} measurements, we also consider the additional uncertainty from the statistical uncertainty on F_L , which gives the largest contribution to the $A_T^{(2)}$ uncertainty.

In summary, we have reconstructed the decays $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ and $B^+ \rightarrow K^{*+} \mu^+ \mu^-$, and report the measurements of the three one-dimensional projections of their full angular distributions. The decay $B^+ \rightarrow K^{*+} \mu^+ \mu^-$ is reconstructed for the first time in hadron collisions. We have measured the muon forward-backward asymmetry A_{FB} , the K^* longitudinal polarization fraction F_L , the transverse polarization asymmetry $A_T^{(2)}$, and the T -odd CP asymmetry A_{im} as a function of q^2 . Measurements of $A_T^{(2)}$ and A_{im} are reported for the first time. We also have measured A_{FB} using the decay $B^+ \rightarrow K^{*+} \mu^+ \mu^-$. All results are among the most precise from a single experiment to date. We do not observe a discrepancy from the SM as reported by Belle, although our results are consistent with all recent measurements from B -factory experiments [7, 8].

We thank C. Bobeth, J. Matias, and W. Altmannshofer for valuable suggestions. Our special gratitude goes to D. van Dyk, who provided the tool to obtain theoretical predictions. We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported

TABLE I: Summary of $B^0 \rightarrow K^{*0}\mu^+\mu^-$ fit results. The first (second) uncertainty is statistical (systematic).

q^2 (GeV ² /c ²)	$N(K^{*0}\mu^+\mu^-)$	F_L	A_{FB}	$A_T^{(2)}$	A_{im}
[0.00, 2.00)	30.7 ± 4.7	$0.31^{+0.17}_{-0.16} \pm 0.02$	$-0.37^{+0.27}_{-0.32} \pm 0.11$	$-0.8 \pm 0.7 \pm 0.3$	$0.37^{+0.31}_{-0.33} \pm 0.08$
[2.00, 4.30)	14.2 ± 4.2	$0.35^{+0.26}_{-0.24} \pm 0.03$	$0.30^{+0.32}_{-0.36} \pm 0.17$	$1.4^{+2.0}_{-1.1} \pm 1.2$	$-0.80^{+0.48}_{-0.29} \pm 0.13$
[4.30, 8.68)	31.3 ± 7.4	$0.60^{+0.17}_{-0.18} \pm 0.05$	$-0.08^{+0.22}_{-0.21} \pm 0.03$	$1.8^{+1.6}_{-1.7} \pm 1.5$	$0.03^{+0.34}_{-0.34} \pm 0.06$
[10.09, 12.86)	38.4 ± 7.6	$0.40^{+0.16}_{-0.16} \pm 0.02$	$0.42^{+0.17}_{-0.21} \pm 0.10$	$-1.0^{+0.9}_{-0.8} \pm 0.5$	$0.47^{+0.26}_{-0.28} \pm 0.09$
[14.18, 16.00)	31.6 ± 4.7	$0.32^{+0.14}_{-0.14} \pm 0.03$	$0.40^{+0.18}_{-0.21} \pm 0.07$	$0.4 \pm 0.8 \pm 0.2$	$0.15^{+0.25}_{-0.26} \pm 0.01$
[16.00, 19.30)	20.7 ± 4.8	$0.16^{+0.22}_{-0.18} \pm 0.06$	$0.66^{+0.18}_{-0.26} \pm 0.19$	$-0.9 \pm 0.8 \pm 0.4$	$-0.30^{+0.36}_{-0.35} \pm 0.14$
[0.00, 4.30)	44.2 ± 6.5	$0.33^{+0.14}_{-0.14} \pm 0.02$	$-0.08^{+0.21}_{-0.20} \pm 0.05$	$-0.2 \pm 0.6 \pm 0.1$	$-0.02^{+0.28}_{-0.28} \pm 0.01$
[1.00, 6.00)	23.8 ± 6.5	$0.60^{+0.21}_{-0.23} \pm 0.09$	$0.36^{+0.46}_{-0.28} \pm 0.11$	$1.6^{+1.8}_{-1.9} \pm 2.2$	$-0.02^{+0.40}_{-0.40} \pm 0.03$

TABLE II: Summary of combined $B \rightarrow K^*\mu^+\mu^-$ fit results. $N(K^{*0}\mu^+\mu^-)$ is taken from Table I.

q^2 (GeV ² /c ²)	$N(K^{*+}\mu^+\mu^-)$	F_L	A_{FB}	$A_T^{(2)}$	A_{im}
[0.00, 2.00)	2.5 ± 1.6	$0.30^{+0.16}_{-0.16} \pm 0.02$	$-0.35^{+0.26}_{-0.23} \pm 0.10$	$-1.0^{+0.7}_{-0.6} \pm 0.4$	$0.21^{+0.30}_{-0.31} \pm 0.10$
[2.00, 4.30)	1.3 ± 1.8	$0.37^{+0.24}_{-0.25} \pm 0.10$	$0.29^{+0.32}_{-0.35} \pm 0.15$	$1.3^{+2.4}_{-1.2} \pm 0.9$	$-0.72^{+0.46}_{-0.36} \pm 0.21$
[4.30, 8.68)	3.9 ± 3.5	$0.68^{+0.15}_{-0.17} \pm 0.09$	$0.01^{+0.20}_{-0.20} \pm 0.09$	$3.4^{+1.9}_{-2.1} \pm 3.6$	$0.11^{+0.31}_{-0.32} \pm 0.09$
[10.09, 12.86)	6.0 ± 2.8	$0.47^{+0.14}_{-0.14} \pm 0.03$	$0.38^{+0.16}_{-0.19} \pm 0.09$	$-1.8^{+0.9}_{-0.8} \pm 0.8$	$0.32^{+0.25}_{-0.26} \pm 0.06$
[14.18, 16.00)	1.6 ± 1.8	$0.29^{+0.14}_{-0.13} \pm 0.05$	$0.44^{+0.18}_{-0.21} \pm 0.10$	$0.2 \pm 0.8 \pm 0.2$	$0.19^{+0.24}_{-0.26} \pm 0.04$
[16.00, 19.30)	4.1 ± 2.3	$0.20^{+0.19}_{-0.17} \pm 0.05$	$0.65^{+0.17}_{-0.18} \pm 0.16$	$-0.7 \pm 0.8 \pm 0.3$	$-0.20^{+0.33}_{-0.33} \pm 0.09$
[0.00, 4.30)	3.8 ± 2.4	$0.33^{+0.14}_{-0.13} \pm 0.03$	$-0.08^{+0.21}_{-0.20} \pm 0.05$	$-0.3 \pm 0.6 \pm 0.1$	$-0.10^{+0.27}_{-0.26} \pm 0.10$
[1.00, 6.00)	5.0 ± 3.0	$0.69^{+0.19}_{-0.21} \pm 0.08$	$0.29^{+0.20}_{-0.23} \pm 0.07$	$1.7 \pm 2.2 \pm 2.5$	$0.09^{+0.34}_{-0.35} \pm 0.18$

TABLE III: Summary of $B^+ \rightarrow K^+\mu^+\mu^-$ fit results.

q^2 (GeV ² /c ²)	$N(K^+\mu^+\mu^-)$	A_{FB}
[0.00, 2.00)	18.6 ± 5.6	$0.13^{+0.42}_{-0.43} \pm 0.07$
[2.00, 4.30)	40.3 ± 6.7	$0.32^{+0.15}_{-0.16} \pm 0.05$
[4.30, 8.68)	68.5 ± 10.5	$0.01^{+0.13}_{-0.10} \pm 0.01$
[10.09, 12.86)	43.5 ± 7.1	$-0.03^{+0.11}_{-0.10} \pm 0.04$
[14.18, 16.00)	35.9 ± 5.7	$-0.05^{+0.09}_{-0.11} \pm 0.03$
[16.00, 23.00)	28.9 ± 6.3	$0.09^{+0.17}_{-0.13} \pm 0.03$
[0.00, 4.30)	57.8 ± 8.8	$0.31^{+0.16}_{-0.16} \pm 0.04$
[1.00, 6.00)	74.5 ± 9.6	$0.13^{+0.09}_{-0.09} \pm 0.02$

by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slo-

vak R&D Agency; and the Academy of Finland.

- [1] $K^{(*)}$ denotes K^\pm , $K^*(892)^0$, or $K^*(892)^\pm$. Charge-conjugate modes are implied throughout this Letter.
- [2] T. Hurth and M. Nakao, Ann. Rev. Nucl. Part. Sci. **60**, 645 (2010).
- [3] F. Kruger and J. Matias, Phys. Rev. D **71**, 094009 (2005).
- [4] U. Egede *et al.*, J. High Energy Phys. 11 (2008) 032.
- [5] S. Descotes-Genon *et al.*, J. High Energy Phys. 06 (2011) 099.
- [6] C. Bobeth, G. Hiller, and G. Piranishvili, J. High Energy Phys. 07 (2008) 106; W. Altmannshofer *et al.*, J. High Energy Phys. 01 (2009) 019.
- [7] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **79**, 031102 (2009).
- [8] J. T. Wei *et al.* (Belle Collaboration), Phys. Rev. Lett. **103**, 171801 (2009).
- [9] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **106**, 161801 (2011).
- [10] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **107**, 201802 (2011).
- [11] K. G. Wilson, Phys. Rev. **179**, 1499 (1969).
- [12] C. Bobeth, G. Hiller, and D. van Dyk, J. High Energy Phys. 07 (2010) 098.
- [13] A. K. Alok *et al.*, arXiv:1008.2367 (2010).
- [14] C. Bobeth, G. Hiller, and G. Piranishvili, J. High Energy Phys. 12 (2007) 040.
- [15] E. J. Thomson *et al.*, IEEE Trans. Nucl. Sci. **49**, 1063 (2002).

- [16] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
- [17] A. A. Affolder *et al.* (CDF Collaboration), Nucl. Instrum. Methods A **526**, 249 (2004).
- [18] G. Ascoli *et al.*, Nucl. Instrum. Methods A **268**, 33 (1988); T. Dorigo (CDF Collaboration), Nucl. Instrum. Methods A **461**, 560 (2001).
- [19] We use a cylindrical coordinate system in which θ is the polar angle with respect to the proton beamline and pseudorapidity $\eta \equiv -\ln(\tan \theta/2)$.
- [20] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **102**, 242002 (2009), and references therein.
- [21] K. Nakamura *et al.* (Particle Data Group), J. Phys. G **37**, 075021 (2010).
- [22] M. Feindt, arXiv:physics/0402093 (2004).
- [23] We draw theory curves using the EOS code by D. van Dyk *et al.* (<http://project.het.physik.tu-dortmund.de/eos/>). However, for illustration purposes, we extend these curves into kinematical regions ($q^2 < 1$, $7 < q^2 < 8.68$, $10.09 < q^2 < 12.86$) where their reliability is known to be approximate.