



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Spin and Majorana Polarization in Topological Superconducting Wires

Doru Sticlet, Cristina Bena, and Pascal Simon

Phys. Rev. Lett. **108**, 096802 — Published 1 March 2012

DOI: [10.1103/PhysRevLett.108.096802](https://doi.org/10.1103/PhysRevLett.108.096802)

Spin and Majorana polarization in topological superconducting wires

Doru Sticlet,¹ Cristina Bena,^{1,2} and Pascal Simon¹

¹*Laboratoire de Physique des Solides, CNRS UMR-8502,*

Univ. Paris Sud, 91405 Orsay Cedex, France

²*Institute de Physique Théorique, CEA/Saclay,*

Orme des Merisiers, 91190 Gif-sur-Yvette Cedex, France

Abstract

We study a one-dimensional wire with strong Rashba and Dresselhaus spin-orbit coupling (SOC), which supports Majorana fermions when subject to a Zeeman magnetic field and in proximity of a superconductor. Using both analytical and numerical techniques we calculate the electronic spin texture of the Majorana end states. We find that the spin polarization of these states depends on the relative magnitude of the Rashba and Dresselhaus SOC components. Moreover, we define and calculate a local “Majorana polarization” and “Majorana density” and argue that they can be used as order parameters to characterize the topological transition between the trivial system and the system exhibiting Majorana bound modes. We find that the local “Majorana polarization” is correlated to the transverse spin polarization, and we propose to test the presence of Majorana fermions in a 1D system by a spin-polarized density of states measurement.

PACS numbers: 73.20.-r, 74.78.Fk, 73.63.Nm

Keywords:

Introduction—Majorana fermions have been attracting a lot of interest recently in the light of the discovery of new materials with topological-insulator properties [1]. These atypical fermionic particles have been predicted long time ago by E. Majorana as real solutions of the Dirac equation [2]. Many condensed-matter systems such as Pfaffian states in fractional quantum Hall (FQH) systems [3], chiral p -wave superconductors [4] (like strontium ruthenate[5]), nodal superconductors under certain conditions [6], ultracold fermionic atoms with laser-field-generated spin-orbit interactions [7], the surface of 3D strong topological insulators [8], as well as semiconductor/superconductor heterostructures [9] have been proposed as platforms supporting Majorana fermions. Among the possible heterostructures, one-dimensional (1D) systems with a strong spin-orbit coupling such as InAs and InSb wires [10], subject to Zeeman magnetic field and in proximity of a superconductor (SC), can exhibit Majorana fermions at their extremities [11, 12]. Various proposals have been made to detect the Majorana states including interferometry [13], noise [14] and spectroscopy measurements [15], etc. However, while a direct confirmation of the existence of the Majorana states would constitute an important step for fields such as quantum computation [16], they have not been detected experimentally so far.

Majorana modes for two dimensional spin-triplet topological superconductors have been shown to exhibit an Ising-like spin density that may allow their detection via coupling to a magnetic impurity[17]. Along similar lines, we propose a method to detect the Majorana states in 1D topological semiconducting wire spectroscopically, using spin-polarized scanning tunneling microscopy (STM). We generalize the model in Refs. 11, 12 to include both Rashba and Dresselhaus spin-orbit interactions, and we show that the resulting Majorana bound states exhibit a characteristic spin texture. In particular we find that the component of the spin polarization in the transverse spin-plane (orthogonal to the direction of the magnetic field) is non-zero solely in the topological phase, and its orientation is determined by the relative weight of the SOC components. Moreover, we introduce a “Majorana pseudospin” local order parameter and define two new local quantities denoted Majorana polarization and Majorana density which quantify locally the Majorana character of a state. We show that the transverse spin polarization is related to the Majorana polarization and we propose that its measurement via spin-polarized STM will allow one to directly visualize the Majorana fermionic states, and thus to test the topological character of a one-dimensional system.

Model— We consider a semiconducting wire oriented along the x -direction, and in prox-

imity to a s -wave superconductor. Due to bulk inversion asymmetry, semiconducting wires can exhibit along with the Rashba SO interaction analyzed in Refs. 12, 18, a Dresselhaus SO interaction of the same order of magnitude[19] (~ 0.1 eV Å). The Bogoliubov-de Gennes (BdG) Hamiltonian for the infinite wire with both types of SOC can be written as

$$H = \int \Psi^\dagger \mathcal{H} \Psi dx, \quad \Psi^\dagger = (\psi_\uparrow^\dagger, \psi_\downarrow^\dagger, \psi_\downarrow, -\psi_\uparrow),$$

$$\mathcal{H} = \left(\frac{p^2}{2m} - \mu + \alpha p \sigma_y + \beta p \sigma_x \right) \tau_z + V_z \sigma_z - \Delta \tau_x. \quad (1)$$

σ 's and τ 's are the usual Pauli matrices acting respectively in the spin and particle-hole spaces. The chemical potential is denoted by μ , V_z is the Zeeman field, Δ is the induced superconducting pairing and α (β) characterize the strengths of the Rashba (Dresselhaus) SOC components. The presence of the Dresselhaus term only trivially modifies the spectrum for the translationally invariant system[12]

$$E^2 = \xi^2 + (\alpha^2 + \beta^2)p^2 + V_z^2 + \Delta^2$$

$$\pm 2(\xi^2(\alpha^2 + \beta^2)p^2 + \xi^2 V_z^2 + \Delta^2 V_z^2)^{1/2}, \quad (2)$$

with $\xi = p^2/2m - \mu$. A careful analysis of this model shows that the conditions for the existence of the topological phase supporting Majorana fermions are unaffected by the Dresselhaus SO interactions, $V_z^2 > \Delta^2 + \mu^2$. It is interesting to note however that Majorana bound states can exist even in the absence of the Rashba term, when Dresselhaus SO interactions are present. Most importantly, the spin texture of the Majorana states is influenced by the presence of the Dresselhaus term. To support this claim we present an analytical study of the wavefunctions corresponding to the Majorana bound states, and we complement it by a numerical study of the corresponding lattice model.

Analytical solution—It has been shown that Majorana bound states can arise at the interface between trivial and topological regions of a one dimensional wire, for example by considering a position dependent chemical potential [12, 20]. Thus, by choosing a chemical potential μ_1 for $x \in [0, L]$, such that $\mu_1^2 < V_z^2 - \Delta^2$, and a $\mu_0^2 > V_z^2 - \Delta^2$ outside this interval, one obtains a finite-size topological region inside a topologically trivial phase. The chemical potentials are chosen such that the $p = 0$ gap, $\Delta - \sqrt{V_z^2 - \mu^2}$, is much smaller than the superconducting gap Δ , which allows one to obtain analytical solutions to the problem by linearizing the Hamiltonian in p . We assume that $L \gg 1$, such that the problem can be

solved independently at the two ends. Thus, ignoring the finite-size effects, the condition to have zero-energy solutions bound at the two interfaces yields the allowed values for the momenta, $\sqrt{\alpha^2 + \beta^2}k_j^\pm = \Delta \pm \sqrt{V_z^2 - \mu_j^2}$, $j \in \{0, 1\}$. Consequently, the Majorana solution at the left boundary $\psi(x \sim 0)$ can be written as

$$\begin{cases} \kappa \mathbf{u}_1(\mu_1) e^{k_1^- x}, & x > 0, \\ \frac{\kappa}{2} \left[\left(1 + \frac{\tan \phi_1}{\tan \phi_0}\right) \mathbf{u}_1(\mu_0) e^{k_0^- x} \right. \\ \left. + \left(1 - \frac{\tan \phi_1}{\tan \phi_0}\right) \mathbf{u}_2(\mu_0) e^{k_0^+ x} \right], & x < 0. \end{cases} \quad (3)$$

Similarly, the Majorana solution at the right boundary $\psi(x \sim L)$ is

$$\begin{cases} \kappa \mathbf{u}_3(\mu_1) e^{-k_1^-(x-L)}, & x < L \\ \frac{\kappa}{2} \left[\left(1 + \frac{\tan \phi_1}{\tan \phi_0}\right) \mathbf{u}_3(\mu_0) e^{-k_0^-(x-L)} \right. \\ \left. + \left(1 - \frac{\tan \phi_1}{\tan \phi_0}\right) \mathbf{u}_4(\mu_0) e^{-k_0^+(x-L)} \right], & x > L \end{cases} \quad (4)$$

with $e^{i\phi_j} = 1/\sqrt{2}(\sqrt{1 + \mu_j/V_z} + i\sqrt{1 - \mu_j/V_z})$, and the Majorana eigenvectors are given by

$$\begin{aligned} \mathbf{u}_1(\mu_j)^T &= (\cos \phi_j e^{i\vartheta}, -\sin \phi_j, \sin \phi_j e^{i\vartheta}, \cos \phi_j), \\ \mathbf{u}_2(\mu_j)^T &= (\cos \phi_j e^{i\vartheta}, \sin \phi_j, -\sin \phi_j e^{i\vartheta}, \cos \phi_j), \\ \mathbf{u}_3(\mu_j)^T &= -(\cos \phi_j e^{i\vartheta}, \sin \phi_j, \sin \phi_j e^{i\vartheta}, -\cos \phi_j), \\ \mathbf{u}_4(\mu_j)^T &= (-\cos \phi_j e^{i\vartheta}, \sin \phi_j, \sin \phi_j e^{i\vartheta}, \cos \phi_j). \end{aligned} \quad (5)$$

Here $e^{i\vartheta} = (\alpha + i\beta)/\sqrt{\alpha^2 + \beta^2}$ allows one to define a two-dimensional spin-orbit vector in the transverse plane, $\mathbf{e}_\vartheta = (\cos \vartheta, \sin \vartheta)$. Note that the obtained wavefunctions are indeed Majorana fermions respecting the reality condition through the phase choice $(\vartheta + \pi)/2$ for the complex coefficient κ . The magnitude of κ is determined from the normalization conditions of the wave functions and is of the order of $(\sqrt{V_z^2 - \mu_1^2} - \Delta/\sqrt{\alpha^2 + \beta^2})^{1/2}$.

The electronic local spin polarization $\mathbf{s}(x)$ of a given four-component (two spin \times electron/hole) Nambu state $|\psi\rangle$ can be calculated by evaluating the expectation values $s_i(x) = \langle \psi | \sigma_i \frac{\tau_0 + \tau_z}{2} | \psi \rangle$ (the $\tau_0 + \tau_z$ insures that we only take into account the electronic, and not the hole degrees of freedom). This prescription yields for the Majorana fermionic states at $x = 0$

and $x = L$:

$$\begin{aligned} \mathbf{s}(0) &= \frac{|\kappa|^2}{2}(-\sin(2\phi_1)\cos\vartheta, \sin(2\phi_1)\sin\vartheta, \cos(2\phi_1)) \\ \mathbf{s}(L) &= \frac{|\kappa|^2}{2}(\sin(2\phi_1)\cos\vartheta, -\sin(2\phi_1)\sin\vartheta, \cos(2\phi_1)). \end{aligned} \quad (6)$$

The above results show that the spin z -components are equal at the ends of the wire, while the transverse spin polarization is equal in magnitude and opposite. Its direction is fixed solely by the relative weight of the Rashba and Dresselhaus SOC components, $s_y/s_x = -\beta/\alpha$. It would be interesting to see whether this results also holds in presence of interactions [18, 21].

Definitions—A general wave function written in Nambu basis described in Eq. (1) as $(u_\uparrow, u_\downarrow, v_\downarrow, v_\uparrow)$ can be recast in a Majorana basis $(\gamma_{1\uparrow}, \gamma_{2\uparrow}, \gamma_{1\downarrow}, \gamma_{2\downarrow})$, where

$$(\gamma_{1\delta}, \gamma_{2\delta}) = \frac{1}{\sqrt{2}}(\psi_\delta^\dagger e^{-i\varphi/2} + \psi_\delta e^{i\varphi/2}, i(\psi_\delta^\dagger e^{-i\varphi/2} - \psi_\delta e^{i\varphi/2})), \quad (7)$$

and φ is a phase characterizing the particular choice of Majorana basis. We can focus simply on the spin up part of the wave function and define the Majorana polarization P_M^\uparrow as the difference between the probability to have a $\gamma_{1\uparrow}$ Majorana and the probability to have a $\gamma_{2\uparrow}$ Majorana, $P_M^\uparrow = P_{\gamma_{1\uparrow}} - P_{\gamma_{2\uparrow}}$. Note that $P_{\gamma_{1/2}}$ but not P_M , have been also introduced in Ref. 18. For the generalized (φ not fixed) Majorana basis, it reads $P_M^\uparrow = -2\text{Re}[u_\uparrow v_\uparrow^* e^{-i\varphi}]$. The phase φ defines a ‘‘Majorana polarization axis’’ such that P_M^\uparrow can be interpreted as a vector (we can denote this vector as ‘‘Majorana pseudospin’’). As such it is decomposed on the axis defined by φ into $P_{M_x}^\uparrow$ ($\varphi = 0$) = $-2\text{Re}[u_\uparrow v_\uparrow^*]$ and $P_{M_y}^\uparrow$ ($\varphi = \pi/2$) = $-2\text{Im}[u_\uparrow v_\uparrow^*]$.

The same procedure can be repeated for the spin down component. This leads to the definition of the full Majorana polarization, $P_{M_i} = P_{M_i}^\uparrow + P_{M_i}^\downarrow$ with $i \in \{x, y\}$. We should note that the absolute value of the polarization vector $\mathcal{P}_M = \sqrt{P_{M_x}^2 + P_{M_y}^2}$, which we denote Majorana density, is not dependent on the choice of (x, y) axes in the Majorana space. For further reference the *Majorana polarization* and *density* are

$$\begin{aligned} P_{M_x} &= 2\text{Re}[u_\downarrow v_\downarrow^* - u_\uparrow v_\uparrow^*], & P_{M_y} &= 2\text{Im}[u_\downarrow v_\downarrow^* - u_\uparrow v_\uparrow^*], \\ P_M &= 2|u_\downarrow v_\downarrow^* - u_\uparrow v_\uparrow^*|. \end{aligned} \quad (8)$$

To emphasize the utility of such definitions the following properties should be noted. For a wave function containing only electronic (or hole) degrees of freedom, the Majorana

polarization is always zero. It equally vanishes for a conventional s -wave superconductor where the wave function has the symmetry $u_{(\downarrow,\uparrow)} = v_{(\uparrow,\downarrow)}^*$. A non-zero value for the Majorana density is a necessary, but not sufficient condition to have Majorana fermions, and identifies the low energy regime in which the model yields an effective p -wave type superconductivity. At zero energy this quantity is maximal and predicts Majorana bound states at the edges. Inspecting the Majorana polarization of these states allows one to unambiguously identify that the two Majorana fermions are of different type. The most important property is that such definitions allow one to explore locally (on-site in the discretized system) the structure of the wave function, and its Majorana character.

For the above analytical solutions, the Majorana polarization vectors $\mathbf{P}_M = (P_{M_x}, P_{M_y})$ yield

$$\mathbf{P}_M(0) = -\mathbf{P}_M(L) = -|\kappa|^2(\cos \vartheta, \sin \vartheta \cos(2\phi_1)) \quad (9)$$

The Majorana polarization vectors are opposite for the two Majorana wave functions at the two ends. Besides, one can identify a relation between the spin polarization vector and the Majorana polarization. When μ , Δ , V_z are fixed, P_{M_x} is proportional to s_x , while P_{M_y} is proportional to s_y . Thus, when only Rashba/Dresselhaus SOC is present, the total transverse spin polarization is proportional to the Majorana polarization, with a proportionality constant which depends on the chemical potential and the applied Zeeman field. When both components of the SOC are present, the Majorana polarization and the transverse spin polarization vectors are no longer collinear but the correlations between these two quantities are still qualitatively preserved.

Numerical model—For the numerical study, we consider a tight-binding formulation of BdG Hamiltonian in Eq. (1)

$$H = \sum_j \Psi_j^\dagger [(\mu - t)\tau_z + V_z\sigma_z - \Delta\tau_x] \Psi_j - \frac{1}{2} \left[\Psi_j^\dagger (t + i\alpha\sigma_y + i\beta\sigma_x)\tau_z \Psi_{j+1} + \text{h.c.} \right] \quad (10)$$

with the Nambu basis $\Psi_j^\dagger = (c_{j\uparrow}^\dagger, c_{j\downarrow}^\dagger, c_{j\downarrow}, -c_{j\uparrow})$. The sum is performed over $N = 100$ sites in the system. We work in units of $t = 1$ and we consider the lattice constant $l = 1$ and $\hbar = 1$. Numerical simulations are done for typical values of the parameters $V_z = 0.4$, $\Delta = 0.3$, $\mu = 0$.

By exact diagonalization we have access to the local density of states, and the local spin-polarized density of states along the x , y , and z directions. For example the local (site n)

electronic i -spin polarization density is given by

$$\rho_{\sigma_i,n} = \sum_{j=1}^{4N} \delta(\omega - E_j) \langle \Psi_n^{(j)} | \sigma_i \frac{\tau_0 + \tau_z}{2} | \Psi_n^{(j)} \rangle \quad (11)$$

where E_j is the j^{th} eigenvalue of H and $|\Psi_n^{(j)}\rangle$ is the site n component of the j^{th} eigenvector, $|\Psi_n^{(j)}\rangle = (u_{n\uparrow}^{(j)}, u_{n\downarrow}^{(j)}, v_{n\downarrow}^{(j)}, v_{n\uparrow}^{(j)})$. We can thus use the definitions in Eq. (8) to calculate the local Majorana polarization and density. For example, the local (n site) Majorana x -polarization can be written as

$$P_{M_x,n} = \sum_{j=1}^{4N} \delta(\omega - E_j) 2\text{Re}[u_{n\downarrow}^{(j)} v_{n\downarrow}^{*(j)} - u_{n\uparrow}^{(j)} v_{n\uparrow}^{*(j)}]. \quad (12)$$

Numerically, we implement the δ functions as very thin Gaussians of width $\approx 0.0001\hbar v_F/l$.

Numerical results—As expected, the exact diagonalization of H recovers the Majorana bound states at the ends of the wire. In Fig. (1) we present the x and z components of the spin polarization, as well as the Majorana polarization when the only considered SOC is Rashba. The symmetrical situation, with only the Dresselhaus component of the SOC present, is analyzed in the Supplementary Material (SM). We can see that the zero-energy Majorana wavefunctions are extended over a small number of edge sites, while exhibiting strongly damped spatial oscillations. The transverse electronic spin polarization is opposite at the two ends of the wire; when only the Rashba term is present only the x -component of the transverse spin is non-zero. The z -spin polarization is the same at the two ends of the wire. While Eq. (6) predicts that the z -polarization vanishes for $\mu_1 = 0$, this is not the case in the numerical calculation. This can be understood by the presence of an effective μ due to the neglected kinetic term, which to leading order, contributes to $\langle p^2 \rangle \approx O((\Delta - V_z)^2)$, which creates a *negative* effective potential as in the numerical results. Moreover, this effective chemical potential is responsible for the spatial (quickly damped) oscillations of the spin polarization observed numerically. Although these oscillations are not captured by the continuum limit calculations, one can however check that the ratio $s_{y,i}/s_{x,i}$ depends only on the spin-orbit couplings in agreement with Eq. (6).

The numerical results for the Majorana polarization presented in Fig. (1) also follow closely Eq. (9). Thus, the values of the Majorana polarization are always opposite at the two ends of the wire, and only P_{M_x} is non-zero when only the Rashba component is present. Also, as expected, P_{M_x} is proportional in this case to the x -spin polarization. Moreover, while we do not present this here in detail, when both the Rashba and Dresselhaus SOC

components are present, the peculiar dependence of the Majorana polarization in Eq. (9) on the $\cos(2\phi_1)$ term indicates that the Majorana polarization vector should exhibit a spatial precession in Majorana space; we have verified numerically that this is indeed the case.

We examine if Majorana polarization is a good order parameter to characterize the topological transition. This is done by varying one of the parameters (Δ, V_z, μ) to drive the system in a trivial phase. In Fig. (2), we vary V_z , and indeed we see that the system becomes trivially gapped (no Majorana bound states) for $V_z \leq \Delta$. The inset describes the dependence of the half-wire integral of the Majorana polarization for one of the lowest-energy states as a function of V_z (an integral of 0.5 is equivalent to a “full” Majorana state). The Majorana polarization decreases smoothly to zero below the critical value of V_z . The same phenomenon can be observed in the second panel, where we plot the spatial distribution of the Majorana polarization as a function of V_z . We have noted that the transition becomes sharper when increasing the size of the system. The same qualitative features are obtained when Δ and μ are varied across the topological transition (the dependence on μ is presented in SM). This shows that the Majorana polarization (and density) is a good local order parameter for the topological transition at $V_z^2 = \Delta^2 + \mu^2$. This suggests that the Majorana polarization can be used to investigate disordered wires[22], and indeed, as shown in SM, in the presence of disorder it exhibits interesting features such as a weak polarization of the low energy bulk states[23]. Moreover, the spin and Majorana polarization exhibit similar spatial structure, even in the presence of disorder.

Conclusion—To summarize, we have found that the Majorana end states are oppositely spin-polarized in the transverse spin-plane, and the direction of polarization depends on the relative weight of the Rashba and Dresselhaus SOC in the wire. Moreover, we have proposed a new wave-function-based measure of the Majorana character of a system, which we denote Majorana polarization. We have seen that this quantity is related to the electronic spin polarization and we have proposed to test the Majorana character of a 1D system using spin-polarized STM measurements. While the density of states measurements can only give information about the existence of a localized state at a given energy, without telling anything about its Majorana character, such a spin-polarized measurement can make the difference between a Majorana excitation or a non-topological localized state.

Acknowledgements

We would like to thank J. Alicea, S. Chakravarty, M. Franz, L. Glazman, A. Kitaev, L.-K.

Lim, F. Piéchon, S. Raghu, G. Refael for interesting discussions. C.B. has been funded by the ERC Starting Independent Researcher Grant NANO-GRAPHENE 256965, in part by the National Science Foundation under Grant No. 1066293, and acknowledges the hospitality of the Aspen Center for Physics where part of this work has been done.

- [1] J. E. Moore, *Nature* **464**, 194 (2010); M. Z. Hasan and C. L. Kane, *Rev. Mod. Phys.* **82**, 3045 (2010); X.-L. Qi and S.-C. Zhang, *Rev. Mod. Phys.* **83**, 1057-1110 (2011).
- [2] E. Majorana, *Il Nuovo Cimento* (1924-1942) **14**, 171 (1937).
- [3] G. Moore and N. Read, *Nucl. Phys. B* **360**, 362 (1991); C. Nayak, and F. Wilczek, *Nucl. Phys. B* **479**, 529 (1996).
- [4] D. A. Ivanov, *Phys. Rev. Lett.* **86**, 268 (2001).
- [5] A. P. Mackenzie and Y. Maeno, *Rev. Mod. Phys.* **75**, 657 (2003).
- [6] M. Sato and S. Fujimoto, *Phys. Rev. Lett.* **105**, 217001 (2010).
- [7] M. Sato, Y. Takahashi, and S. Fujimoto, *Phys. Rev. Lett.* **103**, 020401 (2009); L. Jiang *et al.*, *Phys. Rev. Lett.* **106**, 220402 (2011).
- [8] L. Fu and C. L. Kane, *Phys. Rev. Lett.* **100**, 096407 (2008); L. Fu and C. L. Kane, *Phys. Rev. B* **79**, 161408 (2009).
- [9] J. D. Sau *et al.*, *Phys. Rev. Lett.* **104**, 040502 (2010); J. D. Sau, S. Tewari, R. M. Lutchyn, T. D. Stanescu, and S. Das Sarma, *Phys. Rev. B* **82**, 214509 (2010); J. Alicea, *Phys. Rev. B* **81**, 125318 (2010).
- [10] S. Nadj-Perge *et al.*, *Nature* **468**, 1084 (2010); H. A. Nilsson *et al.*, *Nano Letters* **9**, 3151 (2009); V. Aleshkin *et al.*, *Semiconductors* **42**, **828** (2008), ISSN 1063-7826.
- [11] R. M. Lutchyn, J. D. Sau, and S. Das Sarma, *Phys. Rev. Lett.* **105**, 077001 (2010).
- [12] Y. Oreg, G. Refael, and F. von Oppen, *Phys. Rev. Lett.* **105**, 177002 (2010);
- [13] P. Bonderson, A. Kitaev, and K. Shtengel, *Phys. Rev. Lett.* **96**, 016803 (2006); A. Stern and B. I. Halperin, *Phys. Rev. Lett.* **96**, 016802 (2006); A. R. Akhmerov, J. Nilsson, and C. W. J. Beenakker, *Phys. Rev. Lett.* **102**, 216404 (2009); J. D. Sau, S. Tewari, and S. Das Sarma, *Phys. Rev. B* **84**, 085109 (2011).
- [14] K. T. Law, P. A. Lee, and T. K. Ng, *Phys. Rev. Lett.* **103**, 237001 (2009); S. B. Chung, X.-L. Qi, J. Maciejko, and S.-C. Zhang, *Phys. Rev. B* **83**, 100512 (2011).

- [15] C. J. Bolech and E. Demler, Phys. Rev. Lett. **98**, 237002 (2007), Y. Tanaka, T. Yokoyama, and N. Nagaosa, Phys. Rev. Lett. **103**, 107002 (2009); J. Linder *et al.*, Phys. Rev. Lett. **104**, 067001 (2010).
- [16] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Rev. Mod. Phys. **80**, 1083 (2008).
- [17] R. Shindou, A. Furusaki, and N. Nagaosa, Phys. Rev. B **82**, 180505(R) (2010); R. Zitko and P. Simon, Phys. Rev. B **84**, 195310 (2011)
- [18] E. M. Stoudenmire, J. Alicea, O. A. Starykh, and M. P.A. Fisher, Phys. Rev. B **84**, 014503 (2011).
- [19] J. Schliemann, J. C. Egues, and D. Loss, Phys. Rev. Lett. **90**, 146801 (2003).
- [20] L. Jiang *et al.*, arXiv:1107.4102.
- [21] S. Gangadharaiah, B. Braunecker, P. Simon, and D. Loss, Phys. Rev. Lett. **107**, 036801 (2011).
- [22] see e.g. A. C. Potter and P. A. Lee, Phys. Rev. Lett. **105**, 227003 (2010); R. M. Lutchyn, T. D. Stanescu, and S. Das Sarma, Phys. Rev. Lett. **106**, 127001 (2011); A. R. Akhmerov *et al.*, Phys. Rev. Lett. **106**, 057001 (2011); P. W. Brouwer, M. Duckheim, A. Romito, and F. von Oppen, Phys. Rev. B **84**, 144526 (2011).
- [23] D. Sticlet, C. Bena, and P. Simon, in preparation.

Figures

FIG. 1: The spin polarization along the z and x directions, and the Majorana polarization P_{M_x} , as a function of energy and position, $\Delta = 0.3$, $V_z = 0.4$, $\alpha = 0.2$, $\beta = 0$ and $\mu = 0$.

FIG. 2: First panel: lowest-energy eigenvalues and the half-wire integral of the Majorana polarization (inset) as a function of V_z . Second panel: Majorana polarization of the lowest-energy state as a function of position and V_z . Parameters: $\Delta = 0.3$, $\mu = 0$, $\beta = 0$, and $\alpha = 0.2$




Figure 1a LJ13510 15Dec2011




Figure 1b LJ13510 15Dec2011




Figure 1c LJ13510 15Dec2011




Figure 2a

LJ13510 15Dec2011

