

This is the accepted manuscript made available via CHORUS. The article has been published as:

Approaching Universality in Weakly Bound Three-Body Systems

Vladimir Roudnev and Michael Cavagnero

Phys. Rev. Lett. **108**, 110402 — Published 12 March 2012

DOI: [10.1103/PhysRevLett.108.110402](https://doi.org/10.1103/PhysRevLett.108.110402)

Approaching universality in weakly-bound three-body systems

Vladimir Roudnev and Michael Cavagnero

Department of Physics and Astronomy, University of Kentucky, Lexington, KY 40506-0055

Atom-dimer scattering below the three-body break-up threshold is studied for a system of three identical bosons. The atom-dimer scattering length and the energy of the most weakly-bound three-body state are shown to be strongly correlated. An appropriate rescaling of the observables reveals the subtlety of the correlation, and serves to identify universal trends in the unitary limit of divergent two-body scattering length. The correlation provides a new quantitative measure of the degree of universality in three-body systems with short-ranged interactions, as well as a consistency check of effective field theories and other theoretical models. PACS numbers: 03.65.Nk, 03.65.Ge, 34.50.Cx, 36.40.-c

Among the most striking results of recent experimentation with supercooled atomic gases is the demonstration that trap loss rates are extraordinarily sensitive to few-body interactions within a trapped many-body system. The experiments of many groups [1] show signatures of few-body correlations within a trapped ensemble of Bose alkali vapors at nano-Kelvin temperatures. This discovery has stimulated a large number of theoretical and experimental investigations, particularly in the unitarity limit of a divergent two-body scattering length, where the system has no natural length scale beyond that of the trapping potential.

It is widely anticipated that in the vicinity of unitarity, supercooled Bose gases display universal collective properties. However, criteria for the onset of universality and measures of the degree of universality have not yet emerged. The primary purpose of this Letter is to provide such a measure through a thorough investigation of atom-dimer scattering for a wide variety of two-body short-ranged potentials and the zero-range interaction model [2].

Weakly-bound few-body systems with relatively large two-body scattering lengths have long been studied in nuclear physics. A salient example is a result published by Phillips in 1968 [3] which puzzled nuclear theorists for more than a decade. Phillips compared results of calculations of the neutron-deuteron scattering length and the energy of the triton bound state made with different two-body potentials. He found that, unlike two-body scattering in which the energy of the last bound state scales as the inverse squared scattering length ($E \sim -1/a^2$), the bound state energy of the triton (n-n-p) appears approximately proportional to the scattering length of the neutron-deuteron collision. The data illustrating this correlation is shown in Figure 1.

Many subsequent works [4] confirmed that a strong linear correlation between these two observables exists. Two decades later a simple and elegant explanation of the Phillips line was given by Efimov and Tkachenko [5], who simply noted that the apparent linearity derives from the fact that the two-body potentials used sample only a small portion of the space of scattering parameters; that is, they yield similar values for the neutron-deuteron




FIG. 1. The Phillips line, showing the unexpected linear correlation between the triton bound state energy and the neutron-deuteron scattering length (data from ref. [5]).

scattering lengths.

While the origin of Phillips' observation is now well understood, modern experiments with ultracold gases are able to sample a much wider range of scattering parameters by magnetic field tuning through a Fano-Feshbach resonance. This suggests that the correlation between two- and three-body parameters can be investigated in far greater detail than in earlier nuclear physics studies. Theoretical and experimental works related to universality in ultracold Bose gases have been concentrated on three-body recombination in the close vicinity of the three-body threshold. In this work we discuss the properties of the three-body system at the two-body threshold and below.

In this work we introduce a new parametrization of the relationship between the three-body (atom-dimer) scattering length and the energy of the last three-body bound state for the specific case of three identical Bosons. This relationship is referred to below as the modified Phillips line. In contrast to the original Phillips line, this relationship is found to be linear over a large range of interaction parameters and more directly reflects the well known threshold law for single-channel scattering, namely $E \approx \frac{\hbar^2}{ma^2}$. It also provides a simple test for universality of three-body systems.

As pointed out by Efimov and Tkachenko [5], for a




FIG. 2. a) Phillips line for three ^4He atoms: the He_3 binding energy as a function of the $He - He_2$ scattering length based on four commonly used two-body potentials: TTY [6], LM2M2 [7], HFD-B [8] and HFDBFCI1 [9]. Note the approximately linear relationship (dashed line). b) Modified Phillips line for three ^4He atoms for the same set of potentials. Note an improved linearity.

weakly-bound three-body state the approximate correlation $E_2 - E_3 \approx \hbar^2/(2m_{12}a_3^2)$ should hold, where E_2 and E_3 are the energies of the dimer and trimer bound states (relative to the three-body break-up threshold), m_{12} is the reduced mass of the particle-dimer system and a_3 is the particle-dimer scattering length. A more transparent representation of the strength of the correlation is obtained by rewriting this equation in terms of the scaled dimensionless variables

$$\begin{aligned} \alpha &\equiv a_3 \sqrt{-2m_{12}E_2}/\hbar \\ \omega &\equiv 1/\sqrt{E_3/E_2 - 1} \end{aligned} \quad (1)$$

The variable α can be thought of as a dimensionless scattering length, and ω characterizes the three-body binding energy. If the three-body state nearest to the threshold is deeply bound, ω is small; large values of ω indicate the existence of a weakly-bound three-body state.

Using a recently developed three-body code, we have strenuously tested the well studied case of three bosonic ^4He atoms [12], and so we will use the helium trimer as a first illustration of our improved parameterization. The traditional Phillips line for helium trimer states calculated with four commonly used two-body potentials is plotted in Figure 2a. Figure 2b shows the modified Phillips line obtained using the suggested scaling with the dimer binding energy. Note that the scaling results in an improved fit.

Equation $E_2 - E_3 \approx \hbar^2/(2m_{12}a_3^2)$ derives from the fact that atom-dimer scattering is dominated by the pole of the t -matrix corresponding to the near-threshold state of the trimer. Generally speaking, we should expect the linear relation of the rescaled parameters α and ω to hold provided that no other poles of the t -matrix are relevant




FIG. 3. (Color online) A modified Phillips line plotted with rescaled parameters for a wide variety of two-body parameters and several alternative two-body potentials. Red crosses mark the results for realistic He-He potentials (see Fig. 2)

in the energy range of interest. We have tested this assertion for systems with three indistinguishable bosons by studying the correlation for a variety of two-body potentials with widely adjusted two-body scattering lengths and binding energies.

Figure 3 illustrates the correlation for six different families of two-body potentials. The potentials include the TTY potential [6] with an artificial coupling constant, the family of Bargmann potentials [10] with fixed small effective range and a scattering length varying from 2 to 250 atomic units, the Bargmann potential with an effective range simulating the He-He interaction, the family of Bargmann potentials with varying asymptotic normalizing constant, and the MTV potential [11] ("symmetric model" for the triton) with a varying coupling constant. The near-linearity over a wide range of ω is apparent in the figure, though deviations can be seen, especially for small values of the 3-body scattering length, magnified in the inset of Figure 3, where the complexity of the correlation is revealed. An interesting aspect of the plot, which we had not anticipated, is the emergence of a universal behavior as the 3-body scattering length approaches zero.

To better understand the complex correlation revealed in Figure 3, we show in Figure 4 the same correlation plot using a single two-body potential (the Bargmann potential with the effective range $r_0 = 1$ a.u.), but with large variations of the two-body scattering length. We have also shown the three-body data generated by solving the regularized Skornyakov/Ter-Martirosyan equations (STM) [2, 13] (which are equivalent to Faddeev equations for a zero-range interaction model).

We can identify two distinctive regimes: a universal regime and a strong coupling regime. In the strong coupling regime (characterized by relatively small two-body scattering length) the shape of the modified Phillips line depends on the potential (inset Fig. 3) and, typically, forms an elongated half-loop above the universal curve. First, as the two-body scattering length increases, the

points on the Phillips line move right along the correlation plot reaching a local maximum, and then turn left as they approach the universal regime.

In the universal regime, as the two-body scattering length increases, the points along the Phillips line move left along the universal curve. The shape of the universal correlation plot stems from the interplay of poles of the three-body t -matrix corresponding to the formation of near-threshold bound and virtual states.

We can distinguish three characteristic parts of the universal curve. The linear part at the right end corresponds to large positive particle-dimer scattering length and a very shallow three-body bound state. The diverging part at the left end of the correlation plot corresponds to large negative particle-dimer scattering length and indicates the formation of a near-threshold virtual state. With increasing two-body scattering length this virtual state turns into a bound state, and the corresponding point on the modified Phillips line jumps to the far end of the positive linear part of the universal curve. There is also a transitional nonlinear regime, when the scaled particle-dimer scattering length α is small, and both poles contribute to the shape of the universal curve.

In order to understand the shape of the universal curve better, we use the Faddeev formalism which treats the interactions within two-body subsystems explicitly [14]. Let us start from a simple analysis of the spectrum of the Faddeev operator for identical bosons

$$\hat{K}(E) = \hat{G}_2(E)V(\hat{P}^+ + \hat{P}^-),$$

where $\hat{G}_2(E) = (\hat{H}_0 + V - E)^{-1}$ is the Green's function for the three-body channel Hamiltonian, \hat{H}_0 is the Hamiltonian for three free particles, V is the two-body pairwise potential and \hat{P}^\pm are Jacobi coordinate transformation operators. The equation for the component of the scattering wave function ¹ then reads

$$[1 + \hat{K}(E + i0)]\Phi = -K(E + i0)\chi_0,$$

where χ_0 stands for the atom-dimer plane wave and Φ behaves asymptotically as a three-dimensional spherical wave [15]. The component of the bound state wave function satisfies the homogeneous equation

$$[1 + \hat{K}(E)]\Phi = 0.$$

The Faddeev operator $\hat{K}(E)$ for short-range potentials has a discrete spectrum with eigenvalues $\lambda_n(E)$. For $E \leq E_2$ the eigenvalues $\lambda_n(E)$ are real.

At the two-body threshold $E = E_2$ the Faddeev operator can be approximated by a sum of projectors on




FIG. 4. The modified Phillips line for the Bargmann potential and the zero-range model. For the Bargmann potential the effective range is held fixed at 1 a.u., while the scattering length varies from 2 to 460 a.u. . The values of the two-body scattering length a_2 are shown along the curve.

the states corresponding to eigenvalues λ^+ and λ^- closest to -1 . The particle-dimer scattering length can then be expressed as

$$a_3 = \frac{a^+}{1 + \lambda^+(E_2)} + \frac{a^-}{1 + \lambda^-(E_2)}, \quad (2)$$

where a^+ and a^- are some real positive coefficients corresponding to the residues of the scattering length at the poles $\lambda^\pm(E) = -1$. These residues depend on the energy parameter very smoothly and can be approximated by constants a^+ and a^- . There are two possible situations: 1) one of the eigenvalues $-\lambda^+(E_2)$ or $\lambda^-(E_2)$ is very close to the critical value $\lambda = -1$ and the corresponding term gives the major contribution to the particle-dimer scattering length; 2) both of the terms contribute substantially. The first case is responsible for the “linear” and “singular” parts of the $\alpha(\omega)$ universal correlation plot. The second case corresponds to the intermediate regime of small scaled particle-dimer scattering length α .

This interpretation is illustrated in Fig. 5, where we show the particle-dimer scattering length (top) and the three-body operator spectrum (bottom) as a function of the two-body scattering length. The poles in the three-body scattering length correspond to the eigenvalue of the three-body operator crossing the critical value $\lambda(E_2) = -1$, where a new three-body bound state is formed.

A close fit to the universal part of the modified Phillips line is reproduced by the simple empirical formula

$$\alpha = \frac{\alpha_1}{\frac{1}{\omega} - \frac{1}{\omega_0}} + \omega + \alpha_0 \quad (3)$$

with $\alpha_1 = 5.5$, $\omega_0 = 0.419$ and $\alpha_0 = 4$. The first term here is responsible for the description of the large negative scattering length regime, the other two terms fit the large positive scattering length regime. Equation 3 can be used in practical calculations to estimate the trimer binding energy from the scattering length. A simpler

¹ The three-body wave function can be recovered from the component as $\Psi = (1 + P^+ + P^-)(\Phi + \chi_0)$.




FIG. 5. Atom-dimer scattering length and the spectrum of the kernel of the Faddeev (a) or STM (b) equation at the two-body threshold as a function of the two-body scattering length. a) Bargmann potential with $r_0 = 1$ a.u., b) zero-range interaction model.

empirical fit $\alpha = \omega + \frac{3}{2}(1 - 1/\omega^2)$ is suitable for positive three-body scattering length.

Careful analysis of the modified Phillips line reveals that in the universal, potential-independent regime the scattering length for the atom-dimer collision approaches zero when $E_3/E_2 \approx 2.54$. The other special point on the plot corresponds to divergent negative scattering length, which corresponds to the universal ratio $E_3/E_2 \approx 6.7$.

The result demonstrates a novel form of universality in weakly bound systems. Simple in physical nature, the modified Phillips line provides an important test for numerical and theoretical analysis of 3-body systems. It affords the opportunity to check bound-state and scattering results for consistency and classifies 3-body systems according to distinct dynamical regimes. It also provides an opportunity to check estimates of 3-body bound states and atom-dimer scattering lengths for internal consistency; such as those obtained with regularized zero-range potential models [16]. Correlation plots similar to the modified Phillips line for bosons can be constructed for three-body systems with nontrivial spin-isospin structure and nonidentical particles. Here we shall only mention that the data shown in Figure 1 is consistent with the universal part of the modified Phillips line constructed on the base of the zero-range interaction model [13].

The calculations presented here have been performed using an original code for solving Faddeev equations [17, 18]. The numerically effective computational kernel quickly solves the system of Faddeev equations for bound or scattering states (from a few seconds to a few minutes, depending on the desired numerical accuracy and physical parameters of the system). The code is available from the authors by request and will be available online in the near future.

ACKNOWLEDGEMENTS

Authors thank Dr. Kolganova for stimulating discussions and independent testing of our three-body code. This work is supported by NSF grant PHY-0903956.

- [1] T. Kraemer *et al.*, *Nature* **440**, 315 (2006); M. Zaccanti *et al.*, *Nature Phys.* **5**, 586 (2009); S.E. Pollack, D. Dries, and R.G. Hulet, *Science* **326**, 1683 (2009).
- [2] G.V. Skorniyakov and K.A. Ter-Martirosyan, *Sov. Phys. JETP* **4**, 648 (1957).
- [3] A.C. Phillips *Nucl. Phys A* **107**, 209 (1968)
- [4] N. J. McGurk and H. Fiedeldey, *Zeitschrift fr Physik A*, **274**, 365-372 (1975); I. R. Afnan and J. M. Read, *Phys. Rev. C* **12**, 293305 (1975); E. Huttel, W. Arnold, H. Baumgart, H. Berg and G. Clausnitzer, *Nuclear Physics A*, **406**, 443 (1983); S. Sofianos, H. Fiedeldey and N. J. McGurk *Physics Letters B*, **68**, 117 (1977); J. L. Friar, B. F. Gibson, G. L. Payne and C. R. Chen, *Phys. Rev. C* **30**, 1121 (1984)
- [5] Vitaly Efimov and E.G. Tkachenko, *Phys. Lett. B*, **157**, 108 (1985); Vitaly Efimov and E.G. Tkachenko, *Few-Body Systems*, **4**, 71 (1988)
- [6] K.T.Tang, J.P.Toennies and C.L.Yiu, *Phys. Rev. Lett.* **74**, 1546 (1995).
- [7] Ronald A. Aziz and Martin J. Slaman, *J. Chem Phys.* **94**, 8047 (1991).
- [8] R. A. Aziz, F. R. W. McCourt and C. C. K. Wong, *Mol. Phys.* **61**, 1487 (1987).
- [9] A. R. Janzen and R. A. Aziz, *J. Chem. Phys.* **107**, 914 (1997)
- [10] R. Newton, *Phys. Rev.*, **105**, 763 (1957)
- [11] R. A. Malfliet and J. A. Tjon, *Nucl. Phys.* **127A**, 161 (1969).
- [12] Th. Cornelius and W. Glöckle, *J. Chem. Phys.* **85**, 3906 (1986); E. A. Kolganova, A. K. Motovilov, and W. Sandhas, *Physics of Particles and Nuclei*, **40** 206 (2009); V. Roudnev and S. Yakovlev, *Chem. Phys. Lett.* **328**, 97106 (2000); V. A. Roudnev, S. L. Yakovlev, and S. A. Sofianos, *Few-Body Systems* **37**, 179196 (2005); B. D. Esry, C. D. Lin, and C. H. Greene, *Phys. Rev. A* **54**, 394 (1996); E. Nielsen, D. V. Fedorov, and A. S. Jensen, *J. Phys. B* **31**, 4085 (1998); D. Blume and C. H. Greene, *J. Chem. Phys.* **113**, 2145 (2000); Viatcheslav Kokouline and Franoise Masnou-Sewuws, *Phys. Rev. A* **73**, 012702 (2006); Moses Salci, Evgeny Yarevsky, Sergey B. Levin, Nils Elander, *Int. J. Quant. Chem.*, **107**, 464 (2007).
- [13] V.F. Kharchenko, *Sov. J. Nucl. Phys.* **16** 173 (1973).
- [14] Erich W. Schmidt and Horst Ziegelmann, *The Quantum Mechanical Three-Body Problem*, Pergamon Press, 1974.
- [15] L. D. Faddeev, S. P. Merkuriev, *Quantum scattering theory for several particle systems*, Dordrecht: Kluwer Academic Publishers, (1993).
- [16] Eric Braaten, H.-W. Hammer, *Phys. Rep.* **428** 259 (2006).
- [17] V. Roudnev and Michael Cavagnero, *Comp. Phys. Comm.* **182**, 2099 (2011).
- [18] V. Roudnev and M. Cavagnero, *J. Phys. B* **45** 025101 (2012).