

This is the accepted manuscript made available via CHORUS. The article has been published as:

# Ionization-Grating-Induced Nonlinear Phase Accumulation in Spectrally Resolved Transient Birefringence Measurements at 400 nm

J. H. Odhner, D. A. Romanov, E. T. McCole, J. K. Wahlstrand, H. M. Milchberg, and R. J.  
Levis

Phys. Rev. Lett. **109**, 065003 — Published 7 August 2012

DOI: [10.1103/PhysRevLett.109.065003](https://doi.org/10.1103/PhysRevLett.109.065003)

# Ionization-Grating-Induced Nonlinear Phase Accumulation in Spectrally-Resolved Transient Birefringence Measurements at 400 nm

J. H. Odnher,<sup>1,2</sup> D. A. Romanov,<sup>2,3</sup> E. T. McCole,<sup>1,2</sup> J. K. Wahlstrand,<sup>4</sup> H. M. Milchberg,<sup>4</sup> and R. J. Levis<sup>1,2,\*</sup>

<sup>1</sup>*Department of Chemistry, Temple University Philadelphia, PA 19122*

<sup>2</sup>*Center for Advanced Photonics Research, Temple University Philadelphia, PA 19122*

<sup>3</sup>*Department of Physics, Temple University Philadelphia, PA 19122*

<sup>4</sup>*Institute for Research in Electronics and Applied Physics,  
University of Maryland, College Park, MD 20742*

We report experimental confirmation of the ionization-grating induced transient birefringence predicted by Wahlstrand and Milchberg [1] and discuss its impact on the higher-order Kerr effect (HOKE) interpretation by Loriot, *et al.* of pump-probe transient birefringence measurements made at 800 nm [2]. Measurement of the transient birefringence in air at 400 nm shows a negative contribution to the index of refraction at zero-delay for frequencies within the pump bandwidth, the degenerate case, and no negative contribution for frequencies exceeding the pump bandwidth, the non-degenerate case. Our findings suggest that a re-evaluation of the HOKE hypothesis of Loriot, *et al.* is necessary.

In a recent report, results from a pump-probe transient birefringence measurement were interpreted as evidence for the significance of higher-order terms in the expansion of the nonlinear index of refraction,  $n_{NL}$ , in argon, oxygen, and nitrogen [2, 3]. The so-called higher-order Kerr effect (HOKE) was purported to be responsible for the saturation and subsequent sign inversion of the pump-probe birefringence at zero-delay, measured in a heterodyned polarization gate geometry in [2]. Subsequently, the hypothesis of a negative HOKE contribution to the refractive index at high intensities was used to re-interpret the underlying physics of dynamic spatial replenishment of ultrashort laser pulses (filamentation) [4] as being dominated by saturation of the Kerr response over a wide range of input parameters [2, 3, 5]. The proposed HOKE implies that plasma-induced defocusing does not act as the primary mechanism responsible for counterbalancing Kerr-induced self-focusing in a gas phase filament for short pulses in the near-IR.

Wahlstrand, *et al.* [1] proposed an alternative interpretation of the transient birefringence measurements of Loriot, *et al.* [2], wherein the nonlinear processes in an ionization-induced grating impose a phase shift on the probe pulse through two-beam coupling, leading to an effective birefringence in the medium. A spatially stationary modulation (grating) of the ionization front in the medium originating from optical interference between degenerate pump and probe beams results in negative nonlinear phase accumulation in the probe polarization component parallel to the pump laser, with no phase accumulation in the perpendicular polarization component. Thus, for a probe pulse that is linearly polarized at an angle of 45 degrees with respect to the pump polarization (as in the Loriot measurement), a net nonlinear polarization rotation occurs while the pump and probe beams are temporally overlapped in the medium. The intensity-dependence of the ionization-induced birefringence is predicted to scale as  $I_e^m$  in the multi-photon regime, where  $I_e$

is the pump intensity and  $m$  is the number of photons required to ionize the medium at the laser wavelength (with degenerate pump and probe pulses). The theory predicts that non-degenerate pump and probe beams do not result in an ionization-induced birefringence due to the lack of a stationary ionization grating. In a first approximation, this occurs when the phase velocity of the interference pattern,  $|\delta|/|\mathbf{k}_e - \mathbf{k}_p|$ , traverses the characteristic grating distance  $d = 2\pi/|\mathbf{k}_e - \mathbf{k}_p|$  on the time scale of the laser pulse,  $\tau$ . This suggests that when  $\pi/\delta > \tau$ , nonlinear phase accumulation can occur in the parallel probe polarization, leading to birefringence. In these expressions  $\delta$  is the difference in frequency between the pump ( $e$ ) and probe ( $p$ ) beams and  $\mathbf{k}$  is the wavevector of the respective pulses. For degenerate beams the grating is stationary and net nonlinear phase accumulation can occur; for non-degenerate beams the shifting grating will prevent net nonlinear phase accumulation through destructive interference. If the birefringence observed by Loriot, *et al.* were due to the optical Kerr effect based on neutral-atomic or molecular properties, the polarization rotation should be observed regardless of the frequency of the probe beam with respect to a particular pump frequency [3, 6], excepting the existence of significant dispersion in the higher-order nonlinear refractive index, which is not expected (cf. [7]).

Two previous investigations revealed no evidence for HOKE-induced saturation of the nonlinear refractive index. One was a two-color, 800 nm/400 nm pump-probe cross defocusing measurement [8]. The other used (non-degenerate) single-shot supercontinuum spectral interferometry [9] to measure the nonlinearity directly. However, no direct comparison of two-color and degenerate transient birefringence experiments has been performed using the transient birefringence apparatus of Loriot, *et al.* Furthermore, no experiments have investigated transient birefringence with small detuning between pump and probe pulses, where any possibility of

detrimental dispersion is effectively eliminated. Here, we present spectrally-resolved transient birefringence measurements in air at 400 nm for the degenerate and non-degenerate pump-probe cases using such an apparatus. We demonstrate that no sign inversion is observed where pump and probe are non-degenerate and that the intensity dependence of the observed birefringence in the degenerate case is consistent with the ionization-induced grating model.

The transient birefringence of air was measured using a pump-probe scheme similar to the one described in [2]. An 800 nm, 50 fs laser pulse was split into pump and probe pulses. The probe beam was attenuated with a neutral density filter and a half-wave plate/Glan-laser polarizer pair. A second half-wave plate was used to control the polarization of the probe after the polarizer and a 10- $\mu\text{m}$  Type-II BBO was used to double the probe pulse. The probe beam was then reflected off of multiple dielectric mirrors coated for 400 nm to remove the 800 nm fundamental. The second wave plate and the BBO were rotated to minimize the transmission through a second Glan-laser polarizer (acting as an analyzer) placed after the interaction region and before the spectrometer (USB4000, Ocean Optics) used for signal detection. The pump intensity is independently controlled using a half-wave plate and polarizer. The pump beam was doubled using a 1 mm Type-I BBO and reflected off of multiple 400 nm dielectric high reflectors to remove the fundamental. A second half-wave plate before the BBO was used in combination with the BBO to set the pump polarization at an angle of  $45^\circ$  with respect to the probe. The two beams were focused with an  $f=20\text{-cm}$  lens into ambient air, where they crossed at a small ( $\sim 3^\circ$ ) angle. After the interaction region the probe beam was collimated and sent through a quarter-wave plate before the analyzer and spectrometer. The quarter-wave plate was used to generate a local oscillator for heterodyned measurements by allowing the transmission of a small portion of the probe beam through the analyzer. By rotating the quarter-wave plate  $\pm 1^\circ$  with respect to the incident probe beam polarization axis and measuring the birefringence with positive and negative local oscillator phase contributions the sign of the birefringence is obtained when the two measurements are subtracted. The heterodyne signal can be written as [2]:  $S_{\text{het}}(t) \propto I_p(t) \otimes (\Delta n(t) + \mathcal{P})^2 - I_p(t) \otimes (\Delta n(t) - \mathcal{P})^2$ , where  $I_p(t)$  is the probe intensity profile,  $\Delta n(t) = n_{\parallel}(t) - n_{\perp}(t)$  is the difference in the refractive index along the pump polarization axis and the refractive index along the axis perpendicular to the pump polarization,  $\mathcal{P}$  is the static birefringence induced by the quarter-wave plate, and  $\otimes$  is the convolution operator. The amplitude of the heterodyne birefringence signal is thus proportional to  $n_{\parallel}(t) - n_{\perp}(t)$ . An electronic stage in the probe path was used to control the delay between the pump and probe pulses and the quarter-revolutions of nitrogen and oxygen were used to cal-




FIG. 1. (a) The heterodyned, time- and frequency-resolved birefringence of air measured at  $I_{\text{pump}} = 54 \text{ TWcm}^{-2}$ . Inset: measured pump (dashed) and probe (solid) spectra. (b) The heterodyned, time- and frequency-resolved birefringence of air calculated using Eq. (2) (see text for details).

ibrate the position of zero-delay. The pump intensity was calculated from the pump beam parameters ( $\tau_{\text{pump}} \approx 75 \text{ fs}$ ,  $w_0 = 17 \mu\text{m}$ ). The heterodyned birefringence signal is spectrally-resolved to provide a measurement of the phase accumulation as a function of probe frequency.

Doubling the pump and probe beams with BBO crystals of different thicknesses results in a difference in spectral bandwidth between the two pulses due to the more stringent phase-matching conditions in the thicker crystal. The full-width at half maximum (FWHM) bandwidth of the frequency-doubled pump is 5.2 nm, while the doubled probe FWHM is 9.5 nm. This difference in spectral bandwidth between the pump and the probe pulses allows us to investigate the effects of degeneracy and non-degeneracy on the time-resolved birefringence signal. The broader probe bandwidth provides degenerate birefringence signal where it spectrally overlaps the pump and non-degenerate birefringence signal where there is no spectral overlap with the pump. Figure 1(a) shows the spectrally-resolved transient birefringence of air measured at a pump peak intensity of  $54 \text{ TWcm}^{-2}$ . The intensity experienced by the probe will actually be smaller due to spatial averaging resulting from the non-collinear beam geometry. The effective intensity is expected to be on the order of that reported in [3] ( $\sim I_{\text{pump}}/1.7$ ), corresponding to an intensity of  $32 \text{ TWcm}^{-2}$  for the data shown in Figure 1(a).

The data presented in Figure 1(a) shows that the birefringence undergoes a sign inversion at pump-probe degeneracy similar to that observed in the measurements presented in [2], while the birefringence in regions where there is no spectral overlap between the pump and the probe pulses (i.e. under non-degenerate conditions) exhibits no sign inversion. The sign of the non-degenerate birefringence signal is positive on both the high and low energy sides of the spectrum with respect to the nominal pump central wavelength ( $\sim 400.7 \text{ nm}$ ), suggesting

that pump photons scattered in the probe direction from the induced grating and vice versa (two-beam coupling) play a negligible role in generating the non-degenerate birefringence signal in the present experiment. The contribution of two-beam coupling to the transient birefringence has been discussed previously [3] and will not be revisited here. The positive sign of the birefringence for near-degenerate pump and probe wavelengths is inconsistent with a HOKE interpretation of the data. It is unlikely that dispersion in the nonlinear refractive index would be so large as to prevent a sign inversion at  $(\omega - \omega_0)/\omega_0 \sim \pm 0.7\%$  change in the photon energy both above and below degeneracy.

To gain insight into the effect of frequency detuning between the pump and probe pulses on the transient birefringence, we revisit the theory developed in [1]. The total electron density generated by the pump and probe fields,  $N_e(\mathbf{r}, t)$  can be expressed in terms of a general ionization rate model,  $W(E_{\text{tot}}(t))$ , that depends only on the total laser field amplitude:

$$\frac{N_e(\mathbf{r}, t)}{N_0} = \int_{-\infty}^t dt' W(E_e(t')) \quad (1)$$

$$+ \frac{1}{2} \left( e^{i\mathbf{q}\mathbf{r}} \int_{-\infty}^t dt' E_p(t') W'(E_e(t')) e^{-i\delta t'} + c.c. \right),$$

where  $E_e$  and  $E_p$  are the envelope functions of the excitation or pump laser field and the probe laser field, respectively;  $\mathbf{q} = \mathbf{k}_e - \mathbf{k}_p$  is the grating wavevector;  $W'(E_e)$  is the derivative of  $W(E_e)$  with respect to  $E_e$ ; and the carrier frequencies are retained in the expressions for the pump and probe electric fields, leading to the term  $\exp(-i\delta t)$  in the grating contribution. The first integral in Eq. (1) corresponds to the smoothly varying electron density generated by the pump temporal envelope and the second integral in Eq. (1), corresponding to the contribution leading to ionization grating-induced nonlinear phase accumulation from the pump and probe together, integrates to zero unless  $\delta$  is smaller than the inverse lifetime of the ionization-induced grating formed by  $E_p$  and  $W'(E_e)$ . That is, if there is no spectral overlap between the pump and probe pulses, then no stationary grating is formed through ionization and there is no selective nonlinear phase accumulation in the component of the probe pulse parallel to the pump polarization. If the ionization model is assumed to be multi-photon, i.e.  $W(E) = \sigma_m |E|^{2m}$ , then according to Eq.(1) the amplitude of the ionization-induced change in the refractive index should scale as  $I_e^{m-1}$ , while the nonlinear source term for the transient birefringence should scale as  $I_e^m$ . For ionization of oxygen at 400 nm, a multi-photon ionization model adequately describes ionization in the pump intensity range described here, leading to the prediction of a fourth-order scaling of the negative birefringence signal with pump intensity.

To further investigate the dependence of the negative




FIG. 2. The intensity dependence of the integrated transient birefringence signal at zero-delay (squares), during the permanent alignment (circles), and during the oxygen quarter-revival (triangles). The lines represent fits to the data (see text for details).

birefringence on pump laser intensity, the transient birefringence was measured as a function of laser energy and integrated over the spectrum to yield an equivalent signal to that measured in [2]. Figure 2 displays the temporally-integrated birefringence amplitude over the region of sign inversion at zero-delay (squares), in the region of field-free permanent alignment after zero-delay (circles), and in the vicinity of the oxygen quarter revival (triangles) as a function of pump intensity, respectively. The representative error bar in Figure 2 reflects uncertainty in the intensity calculated from measured beam parameters (est. error of 10% pulse duration and  $\pm 0.5 \mu\text{J}$ ) and applies to all data points. The field-free alignment and revival data are fit with exponentials of the form:  $A \times x^B$ , and the zero-delay signal is fit by:  $A \times x^B + C \times x$ . We observe linear ( $B = 1.06$ ) and quadratic ( $B = 2.02$ ) behavior of the integrated oxygen revival and permanent alignment data, respectively, confirming that saturation of the alignment does not occur [10] and, more importantly, indicating that a decrease in the pump intensity due to a shift in the focal position (resulting from Kerr self-focusing and/or plasma defocusing) does not significantly impact the current experiment even though measurements were performed at atmospheric pressure. The integrated zero-delay data, fit with linear and fourth-order contributions, is in agreement with the predictions of the ionization-induced birefringence model of [1].

In the experimental geometry of the present investigation, the detected signal is given by  $I_{\text{het}}(\omega) \propto 2\text{Re}[E_{LO}(\omega) \times E_s(\omega)]$  (by analogy with the time-domain expression for the total heterodyne birefringence signal in [11]), where  $E_{LO}(\omega)$  and  $E_s(\omega)$  are the Fourier components of the local oscillator and signal fields, respectively. For carrier-frequency-degenerate pump and probe pulses crossing at a small angle in the medium, the detected

heterodyne birefringence signal can be approximated as:

$$I_s(\omega) \propto \text{Re} \left[ e^{i\omega t_d} E_p(\omega) i z \int_{-\infty}^{\infty} dt e^{i\omega t} \left( -\frac{k_p n_2}{2n_0} E_p^*(t - t_d) \left( \frac{2}{3} I_e(t) + \frac{3}{2} \int_{-\infty}^t dt' R(t - t') I_e(t') \right) \right. \right. \\ \left. \left. + 4\beta e^{iz\beta f(t)} E_e^*(t) \int_{-\infty}^t dt' E_p^*(t' - t_d) E_e(t') I_e(t')^{m-1} \right) \right], \quad (2)$$

where the first term in the Fourier transform describes the contribution from the electronic and rotational Kerr response of the medium, the second term describes the contribution from the ionization-induced grating, and  $z$  is the interaction length. Here,

$$\beta = \frac{2\pi k_p}{n_0^2} \frac{N_0 \sigma_m}{N_c}, \quad f(t) = \int_{-\infty}^t d\tau I_e(\tau)^m,$$

and  $N_c$  represents the critical plasma density at the carrier frequency. Equation (2) retains only the leading terms in the parameter  $z\beta f(\infty)$ ; inclusion of additional terms leads to small corrections in (2). The rotational contribution has been approximated with the response function  $R(t - t_d)$  [12] rather than with a full calculation as in [2], and multi-photon ionization has been assumed. The spectrally-resolved transient birefringence calculated using Equation (2) with Gaussian input pulses ( $\tau_{pu} = 75$  fs,  $\tau_{pr} = 37$  fs) and  $I_e = 32$  TWcm $^{-2}$  is shown in Figure 1(b). Because of the uncertainty in the value of  $n_2$  at 400 nm (taken to be  $n_2 = 5.36 \times 10^{-19}$  Wcm $^{-2}$  [13]) and in the ionization rate, the Raman/Kerr contribution has been scaled to match the experimental proportions relative to the ionization-induced grating contribution. The simulation captures the essential features of the experimental data that distinguish it from a purely Kerr-driven response, namely the positive sidebands on the leading edge of the pump pulse. Spectral- and temporal- integration of the pure ionization-induced birefringence as a function of intensity yielded a fourth-order power dependence, and in combination with a linear contribution (accounting for Kerr and Raman effects) agrees with the data shown in Figure 2.

We next consider the possibility of HOKE term contributions to the measured signal and also the ramifications of the ionization-induced birefringence on the HOKE theory. In the UV (i.e. at  $\leq 400$  nm), plasma generation is expected to dominate over the HOKE indices [3, 6], which would lead to suppression of the sign inversion of the birefringence were it due solely to the HOKE terms. Ettoumi, *et al.* calculate the Kerr-canceling intensity in air at 400 nm to be  $\sim 32$  TWcm $^{-2}$  based on a generalization of the Miller formulæ [14], but our data shows inversion well below this intensity. For comparison, inversion of the integrated temporal profile in our data, occurs at 32.1  $\mu$ J input energy, which corresponds to  $I_e/1.7 = 28$

TWcm $^{-2}$ . We note that subsequent work has cast doubt on the validity of the generalized Miller formulæ [15], suggesting that the disagreement between the calculation in [6] and our data can only be qualitatively discussed. Regardless, the measurements presented here show comparable inversion of the transient birefringence as seen in [2] at 800 nm, suggesting that the sign inversion of the transient birefringence is unlikely to be a distinct indication of HOKE-dominated propagation.

To conclude, we have presented experimental results consistent with the existence of an ionization-grating-induced contribution to the transient birefringence of air measured in a pump-probe configuration similar to [2]. We measured the dependence of the sign inversion of the birefringence on pump intensity at 400 nm and showed that it scales as  $I_e^m$ , where  $m$  is the number of photons required to ionize in the multi-photon limit. We also report that the ionization-induced transient birefringence signal occurs only at degeneracy and that non-degeneracy between the pump and probe pulses results in no sign inversion of the birefringence. This is in accord with the prediction of Wahlstrand, *et al.* The difference in the sign of the birefringence between degenerate and near-degenerate pump and probe wavelengths cannot be explained by the HOKE model. Our findings cast doubt on the interpretation of the transient birefringence measured by Loriot, *et al.* as originating purely from higher-order terms in the Kerr nonlinearity and show that the ionization-induced grating contribution to the induced birefringence must be accounted for in order to correctly interpret the results reported in [2].

RJL acknowledges the Office of Naval Research Grant No. N00014-10-0293, the Air Force Office of Scientific Research Grant No. FA9550-10-1-0561, and the National Science Foundation Grant No. CHE0957694 for support. HMM acknowledges the National Science Foundation, the U.S. Department of Energy, the Office of Naval Research, and the Lockheed Martin Corporation for support.

---

\* rjlevis@temple.edu

[1] J. K. Wahlstrand and H. M. Milchberg, Optics Letters, **36**, 3822 (2011).

- [2] V. Lorient, E. Hertz, O. Faucher, and B. Lavorel, *Optics Express*, **17**, 13429 (2009).
- [3] V. Lorient, P. Béjot, W. Ettoumi, Y. Petit, J. Kasparian, S. Henin, E. Hertz, B. Lavorel, O. Faucher, and J.-P. Wolf, *Laser Physics*, **21**, 1319 (2011).
- [4] M. Mlejnek, E. M. Wright, and J. V. Moloney, *Optics Letters*, **23**, 382 (1998).
- [5] P. Béjot, J. Kasparian, S. Henin, V. Lorient, T. Vieillard, E. Hertz, O. Faucher, B. Lavorel, and J.-P. Wolf, *Physical Review Letters*, **104**, 103903 (2010).
- [6] W. Ettoumi, P. Béjot, Y. Petit, V. Lorient, E. Hertz, O. Faucher, B. Lavorel, J. Kasparian, and J.-P. Wolf, *Physical Review A*, **82**, 033826 (2010).
- [7] D. P. Shelton and J. E. Rice, *Chemical Review*, **94**, 3 (1994).
- [8] Y. Feng, H. Pan, J. Liu, C. Chen, J. Wu, and H. Zeng, *Optics Express*, **19**, 2852 (2011).
- [9] J. K. Wahlstrand, Y.-H. Cheng, Y.-H. Chen, and H. M. Milchberg, *Physical Review Letters*, **107**, 103901 (2011).
- [10] V. Renard, M. Renard, A. Rouzée, S. Guérin, H. R. Jauslin, B. Lavorel, and O. Faucher, *Physical Review A*, **70**, 033420 (2004).
- [11] B. Lavorel, H. Tran, E. Hertz, O. Faucher, P. Joubert, M. Motzkus, T. Buckup, T. Lang, H. Skenderovi, G. Knopp, P. Beaud, and H. M. Frey, *Comptes Rendus Physique*, **5**, 215 (2004).
- [12] E. T. J. Nibbering, G. Grillon, M. A. Franco, B. S. Prade, and A. Mysyrowicz, *Journal of the Optical Society of America B*, **14**, 650 (1997).
- [13] Y. E. Geints and A. A. Zemlyanov, *Applied Physics B*, **101**, 735 (2010).
- [14] W. Ettoumi, Y. Petit, J. Kasparian, and J.-P. Wolf, *Optics Express*, **18**, 6613 (2010).
- [15] G. Stegeman, M. G. Kuzyk, D. G. Papazoglou, and S. Tzortzakis, *Optics Express*, **19**, 22486 (2011).