

This is the accepted manuscript made available via CHORUS. The article has been published as:

Emergent Critical Phase and Ricci Flow in a 2D Frustrated Heisenberg Model

Peter P. Orth, Premala Chandra, Piers Coleman, and Jörg Schmalian

Phys. Rev. Lett. **109**, 237205 — Published 4 December 2012

DOI: [10.1103/PhysRevLett.109.237205](https://doi.org/10.1103/PhysRevLett.109.237205)

Emergent Critical Phase and Ricci Flow in a 2D Frustrated Heisenberg Model

Peter P. Orth,¹ Premala Chandra,² Piers Coleman,^{2,3} and Jörg Schmalian^{1,4}

¹*Institute for Theory of Condensed Matter, Karlsruhe Institute of Technology (KIT), 76131 Karlsruhe, Germany*

²*Center for Materials Theory, Rutgers University, Piscataway, New Jersey 08854, USA*

³*Hubbard Theory Consortium and Department of Physics,*

Royal Holloway, University of London, Egham, Surrey TW20 0EX, UK

⁴*DFG Center for Functional Nanostructures, Karlsruhe Institute of Technology, 76128 Karlsruhe, Germany*

(Dated: September 14, 2012)

We introduce a two-dimensional frustrated Heisenberg antiferromagnet on interpenetrating honeycomb and triangular lattices. Classically the two sublattices decouple, and “order from disorder” drives them into a coplanar state. Applying Friedan’s geometric approach to nonlinear sigma models, we obtain the scaling of the spin-stiffnesses governed by the Ricci flow of a 4D metric tensor. At low temperatures, the relative phase between the spins on the two sublattices is described by a six-state clock model with an emergent critical phase.

PACS numbers: 75.10.-b, 75.10.Jm

A remarkable discovery of recent years is that frustrated two dimensional Heisenberg models can evade the Hohenberg-Mermin-Wagner theorem [1, 2] via the development of long-range discrete order driven by short-range thermal spin fluctuations: such discrete long-range order develops despite the persistence of a finite spin correlation length, leading to a finite temperature Ising (\mathbb{Z}_2) or \mathbb{Z}_3 Potts phase transition [3–9]. This phenomenon is well-established in the J_1 - J_2 Heisenberg model on the square lattice and has recently been realized in iron-based superconductors [10]. An interesting question motivated by this discovery is whether it can be generalized to higher \mathbb{Z}_p ($p \geq 5$) order. If one can show, in addition, that these emergent discrete degrees of freedom are described by a p -state clock model [11, 12], the unique situation arises that a Heisenberg spin system exhibits two Berezinskii-Kosterlitz-Thouless (BKT) transitions which bracket a critical phase. In a system of discrete Ising spins, such a scenario was reported to occur on the triangular lattice [13, 14].

In this Letter, we introduce such a Heisenberg model defined on interpenetrating honeycomb and triangular lattices (Fig. 1(a)) with nearest-neighbor antiferromagnetic coupling. This model may be realized with cold spinful atoms in optical lattices, where it arises naturally in the limit of large on-site interactions [15–18]. Another promising experimental route is to employ STM techniques for nano-fabrication and spin-resolved read-out of stacked triangular and honeycomb monolayers of magnetic atoms like Cr or Co [19–22]. For classical spins the two sublattices are decoupled giving rise to an $\text{SO}(3) \times \text{O}(3)/\text{O}(2)$ order parameter. “Order from disorder” [23, 24] drives the two sublattices into a coplanar spin configuration [25] with an $\text{SO}(3) \times \text{U}(1)$ order parameter and a six-fold in-plane potential. In the coplanar state we explicitly show that the $\text{U}(1)$ degrees of freedom decouple to form an emergent \mathbb{Z}_6 clock model with an intermediate power-law phase. This non-trivial decoupling

of the $\text{U}(1)$ phase is essential for the critical phase to occur.

A novel aspect of our work is that we apply Friedan’s coordinate-independent approach to nonlinear sigma models [26] to the scaling of the spin-stiffness. In this approach the configurations of the 2D spin system correspond to a worldsheet of a string evolving in four-dimensions, where the metric is determined by the components of the antiferromagnetic stiffness and its renormalization corresponds to a Ricci flow of the metric tensor. We also note the decoupling of our $\text{U}(1)$ phase can be viewed as a toy model for the compactification of a four-dimensional string-theory.

Specifically we study the antiferromagnetic Heisenberg model on a decorated 2D triangular lattice (cf. Fig. 1(a));




FIG. 1. (color online) (a) Heisenberg model on the “windmill” lattice. (b) Definition of angles α and β describing the relative orientation of magnetic order on triangular and honeycomb lattice. (c-d) Angle dependent free energy correction δF from thermal and quantum spin fluctuations for parameters $J_{hh} = J_{tt} = 1$, $J_{th} = 0.4$, $T = 1$. Panel (d) is for fixed $\beta = \pi/2$.

the associated Bravais lattice has three basis sites per unit cell at positions $\mathbf{b}_t = a_0(0, 2/\sqrt{3})$, $\mathbf{b}_A = (0, 0)$ and $\mathbf{b}_B = a_0(0, 1/\sqrt{3})$ where indices A, B label the two honeycomb sites. We set the lattice constant $a_0 = 1$. The Hamiltonian is $H = H_{tt} + H_{AB} + H_{tA} + H_{tB}$ with

$$H_{ab} = J_{ab} \sum_{j=1}^{N_L} \sum_{\{\delta_{ab}\}} \mathbf{S}_a(r_j) \cdot \mathbf{S}_b(r_j + \delta_{ab}), \quad (1)$$

where $\mathbf{S}_a(r_j)$ denote spin operators at Bravais lattice site j and basis site $a \in \{t, A, B\}$. The vectors $\{\delta_{ab}\}$ point between nearest-neighbors of sublattices a, b . We assume in the following that the spin exchange couplings within the same sublattice are larger than the inter-sublattice coupling $J_{th} < J_{tt}, J_{hh}$, where $J_{th} \equiv J_{tA} = J_{tB}$ and $J_{hh} \equiv J_{AB}$. For decoupled lattices $J_{th} = 0$, the classical ground state on the bipartite honeycomb lattice is the usual Néel state, while spins on the triangular lattice arrange in a 120° configuration [27]. Although the exchange fields between the two sublattices exactly cancel for this configuration even for $J_{th} > 0$, quantum and thermal fluctuations depend on the relative orientation of the magnetization on the two sublattices. The uniaxial magnetic order on the honeycomb lattice is described by a normal vector $\mathbf{n}(x)$, which points along the magnetization on sublattice A . The biaxial order on the triangular lattice is characterized by a triad of orthonormal vectors $\{\mathbf{t}_j(x)\}$ with $j = 1, 2, 3$. Equivalently, it may be expressed by an orthogonal matrix $t = (\mathbf{t}_1, \mathbf{t}_2, \mathbf{t}_3) \in SO(3)$. We take the vectors $\mathbf{t}_{1,2}$ to span the plane of the magnetization on the triangular lattice. The relative order between the two sublattices is thus determined by two angles α and β , that are defined in Fig. 1(b).

Symmetry considerations dictate the form of the long-wavelength action which takes the form of a nonlinear sigma model (NLSM)

$$S = \int d^2x \left(\frac{K}{2} (\partial_\mu \mathbf{n})^2 + \sum_{j=1}^3 \frac{K_j}{2} (\partial_\mu \mathbf{t}_j)^2 \right) + S_c. \quad (2)$$

The action contains the usual gradient terms of the $O(3)/O(2)$ and the $SO(3)$ NLSM for the order parameter on the honeycomb and triangular lattice. The bare spin stiffnesses K, K_j can be derived in a $1/S$ -expansion and read $K = 2J_{hh}S^2/T$, $K_1 = K_2 = \sqrt{3}J_{tt}S^2/4T$ and $K_3 = 0$ [28–30]. In addition, the action in Eq. (2) contains two potential terms, generated by short-wavelength spin fluctuations (“order from disorder”) [23, 24]

$$S_c = \frac{1}{2} \int d^2x (\gamma \cos^2 \beta + \lambda \sin^6 \beta \sin^2(3\alpha)). \quad (3)$$

A positive $\gamma > 0$ favors coplanarity, whereas $\gamma < 0$ induces \mathbf{n} to be perpendicular to the plane of the triangular magnetization. The six-fold anisotropy term λ is relevant only for $\gamma > 0$.

Heuristically, we expect $\gamma > 0$, favoring coplanarity: spins on the honeycomb lattice can minimize their energy by aligning themselves perpendicular to the fluctuation Weiss field from the triangular lattice [24]. To confirm this reasoning, we have performed a Holstein-Primakov spin wave analysis of Eq. (1). Our results for the fluctuation correction to the free energy for arbitrary angles α and β between the two sublattices are given in Fig. 1(c-d) and show that $\gamma > 0$. For small J_{th} we find $\gamma = (J_{th}/\bar{J})^2 A_\gamma (J_{tt}/J_{hh}, \bar{J}/T)$ is the dominant term in the potential, while $\lambda = (J_{th}/\bar{J})^6 A_\lambda (J_{tt}/J_{hh}, \bar{J}/T)$, where $\bar{J} = \sqrt{J_{tt}J_{hh}}$ and the $A_{\gamma,\lambda}$ are functions that depend weakly on J_{tt}/J_{hh} .

As temperature is reduced, the two sublattices enter a *coplanar regime*. The temperature scale for this crossover is easily determined from standard scaling arguments and yields the coplanar crossover temperature

$$T_{cp} \simeq \frac{J_{hh}S^2}{1 + \ln(1/\gamma)/4\pi} \quad (4)$$

in case where $J_{hh} < J_{tt}$. In the opposite regime $J_{tt} < J_{hh}$ we obtain an implicit expression for T_{cp} that also approaches zero only logarithmically for $\gamma \rightarrow 0$. The crossover temperature in Eq. (4) follows from the known flow equation $\frac{d}{dl}K = -1/2\pi$ for the stiffness K with running cutoff $\Lambda(l) = a_0^{-1}e^{-l}$ and the flow of the coplanar potential amplitude $\gamma(l) = \gamma \exp(2l)$ that is determined by its engineering dimension. While the spin stiffnesses are reduced at longer length-scales, the potential term grows, and scaling stops when $\gamma(l_\gamma) = 1$, which defines a length-scale $a_\gamma = a_0 e^{l_\gamma} \simeq a_0 (J_{hh}/J_{th})^2$. The coplanar crossover takes place when this length-scale is comparable to the shorter of the magnetic correlation lengths on the two sublattices. From the known flow equation of the $O(3)/O(2)$ and the $SO(3)$ NLSM further follows that the stiffnesses of the triangular lattice approach an isotropic fixed point [31]. The six-fold symmetric potential $\propto \lambda$ flows to larger values, yet due to $\lambda \ll \gamma$ holds that $\lambda(l_\gamma) \simeq \mathcal{O}(J_{th}^4) \ll 1$.

Once the two sublattices are coplanar, their dynamics are intimately connected. To describe this regime we impose a hard-core constraint: $\mathbf{n} \perp \mathbf{t}_3$, i.e., $\beta = \pi/2$. It is now convenient to introduce a second triad $\mathbf{h}_{1,2,3}$, defining an $SO(3)$ matrix $h = (\mathbf{h}_1, \mathbf{h}_2, \mathbf{h}_3)$ that describes the magnetic order on the honeycomb lattice with $\mathbf{h}_1 = \mathbf{n}$. The coplanar constraint is expressed as $t = hU$ where $U = \exp(i\alpha\tau_3)$ determines the relative in-plane orientation of the two sublattices, defined by the angle α . We describe h in terms of three Euler angles, $h = e^{-i\phi\tau_3} e^{-i\theta\tau_1} e^{-i\psi\tau_3}$. Here, the τ_a satisfy the $SU(2)$ algebra $[\tau_a, \tau_b] = i\epsilon_{abc}\tau_c$ and take the adjoint form $(\tau_a)_{bc} = i\epsilon_{bac}$. The coplanar system is thus determined by an $SO(3) \times U(1)$ order parameter, defined by three Euler angles and a single relative phase α .

To analyze this coupled problem we write the action

in the form $S = S_X + S_c$, where

$$S_X = \frac{1}{2} \int d^2x g_{ij} [X(x)] \partial_\mu X^i(x) \partial_\mu X^j(x) \quad (5)$$

with coordinates $X = (\phi, \theta, \psi, \alpha)$ and stiffness tensor

$$g = \begin{pmatrix} g^{SO(3)} & \mathcal{K}^T \\ \mathcal{K} & I_\alpha \end{pmatrix}, \quad (6)$$

where

$$g_{ij}^{SO(3)} = \begin{pmatrix} (I_1 s_\psi^2 + I_2 c_\psi^2) s_\theta^2 + I_3 c_\theta^2 & (I_1 - I_2) c_\psi s_\theta s_\psi & I_3 c_\theta \\ (I_1 - I_2) c_\psi s_\theta s_\psi & I_1 c_\psi^2 + I_2 s_\psi^2 & 0 \\ I_3 c_\theta & 0 & I_3 \end{pmatrix},$$

with $s_{X^j} = \sin X^j$ and $c_{X^j} = \cos X^j$. In our system we find $I_1 = K_1 + K_3$, $I_2 = K_1 + K_3 + K$, $I_3 = 2K_1 + K$, which are set by the stiffnesses of the two sublattices at $l = l_\gamma$. The $U(1)$ degree of freedom α has an initial stiffness $I_\alpha = 2K_1(l_\gamma)$ and is coupled to the non-Abelian $SO(3)$ sector by the term $\mathcal{K} = \frac{\kappa}{2}(c_\theta, 0, 1)$ in the four-dimensional metric, where $\kappa = 4K_1(l_\gamma)$. The six-fold potential $S_c(\beta = \frac{\pi}{2}) = \frac{1}{2} \lambda \int d^2x \sin^2(3\alpha)$ is a small but relevant perturbation to S_X . At length-scales where λ is small, the anisotropy S_c and the gradient term S_X (5) is the action of a classical string in a four dimensional space with coordinates $X(x)$ at the two dimensional worldsheet point x , with metric tensor $g_{ij}[X]$. Under coordinate transformations $X_i \rightarrow X'_i$, S_X in Eq. (5) is invariant, with transformed metric $g'_{lm} = g_{ij} \frac{\partial X^i}{\partial X'^l} \frac{\partial X^j}{\partial X'^m}$. Like Einstein's theory of gravity, this covariance tells us that the long-wavelength action S_X is co-ordinate independent and only depends on the geometric aspects of the mapping $X(x)$ of the worldsheet to the compact four-dimensional space of the coordinate X . The renormalization group flow of the metric tensor must also be covariant under co-ordinate transformations, and following the geometric formulation of the NLSM by Friedan [26], to two loop order takes the form

$$\frac{dg_{ij}}{dl} = \frac{1}{2\pi} R_{ij} - \frac{1}{8\pi^2} R_i{}^{klm} R_{jklm}, \quad (7)$$

where R^{iklm} is the Riemann curvature tensor and $R_{ij} = R^k{}_{ikj}$ is the Ricci tensor [32]. This expression defines a generalized Ricci flow [33]. The Riemann tensor is determined by the Christoffel symbols $\Gamma_{jk}^i = \frac{1}{2} g^{il} (g_{jl,k} + g_{kl,j} - g_{jk,l})$ as $R^k{}_{lij} = \Gamma_{lj,i}^k - \Gamma_{li,j}^k + \Gamma_{ni}^k \Gamma_{lj}^n - \Gamma_{nj}^k \Gamma_{li}^n$. The flow equations of our five coupling constants I_j , I_α and κ follow from Eq. (7).

A key insight into the low energy phase diagram is obtained by noting the coupling term \mathcal{K} can be eliminated via a coordinate transformation $\psi \rightarrow \psi' = \psi + r\alpha$ with $r = \kappa/2I_3$. This yields a metric g in Eq. (6) with $\mathcal{K} = 0$, $I_\alpha \rightarrow I'_\alpha = I_\alpha - \kappa^2/4I_3$ yet with $g^{SO(3)}$ that depends on the $U(1)$ phase α via the above shift of the Euler angle ψ . This gauge transformation to the appropriate center of

mass coordinates allows for clear criteria when the $U(1)$ sector of the theory decouples from the $SO(3)$ sector: if either $|I_1 - I_2| \ll \sqrt{I_1 I_2}$ or $r \ll 1$ it follows that $g^{SO(3)}$ becomes independent of α and the $U(1)$ phase decouples from the dynamics of the non-collinear magnetic degrees of freedom. The first criterion follows from the fact that $g^{SO(3)}$ is independent of ψ if $I_1 = I_2$, while the second criterion implies that the shift in ψ is negligible. From Eq. (7) follows after a lengthy but straightforward calculation that $I_{1,2,3}$ flow to an isotropic fixed point, while the dimensionless variable r follows the flow equation (for simplicity we only list the one loop result, the two loop correction does not change our conclusions):

$$\frac{dr}{dl} = -r \frac{(I_1 - I_2)^2}{4\pi I_1 I_2 I_3}. \quad (8)$$

Thus, if the initial anisotropy $|I_1 - I_2| = K$ is weak, which happens for $J_{hh} \ll J_{tt}$, the coupling r does not change much. The $SO(3)$ sector, however, quickly becomes isotropic in the 1-2-plane leading to a decoupling of the $U(1)$ phase. On the other hand, in the limit of strong anisotropy for $J_{hh} \gg J_{tt}$, where $|I_1 - I_2|$ is not small, we find that r vanishes rapidly. In both cases follows that the phase angle α emerges as an independent degree of freedom. The β -function for the reduced phase stiffness $I'_\alpha = I_\alpha - \kappa^2/4I_3$ follows from Eq. (7) as

$$\frac{dI'_\alpha}{dl} = \beta_\alpha = \frac{(I_1 - I_2)^2 r^2}{4\pi I_1 I_2}, \quad (9)$$

and does, as expected, approach zero once either of the two decoupling conditions are fulfilled. Thus, perturbatively no renormalization of the stiffness I'_α takes place. An interesting aspect of the decoupling follows from the Ricci scalar $R = g^{ij} R_{ji}$:

$$R = R^{SO(3)} - \frac{1}{2\pi I'_\alpha} \beta_\alpha \quad (10)$$

where $R^{SO(3)} = \sum_{j=1}^3 \left(I_j^{-1} - \frac{1}{2I_1 I_2 I_3} I_j^2 \right)$ is the Ricci scalar of the $SO(3)$ sector. Once the decoupling takes place, $\beta_\alpha \rightarrow 0$ and the $U(1)$ sector becomes flat. On the other hand $R \rightarrow R^{SO(3)}$ grows under renormalization since the stiffnesses I_j decrease. Thus, we arrive at a flat one dimensional sector weakly coupled to an three-dimensional manifold with large curvature. This "curling-up" and asymptotic decoupling of a subspace may serve as a toy model for compactification.

Since in both limits $J_{hh} \ll J_{tt}$ and $J_{hh} \gg J_{tt}$ follows that the decoupling emerges rapidly, we find that λ , whose flow is governed by $\frac{d}{dl} \lambda = (2 - 9/\pi I'_\alpha) \lambda$, is still small. The resulting low-energy theory corresponds to $S = S_{SO(3)} + S_{\mathbb{Z}_6}$ with

$$S_{\mathbb{Z}_6} = \frac{1}{2} \int d^2x [(I'_\alpha (\partial_\mu \alpha)^2 + \lambda \cos(6\alpha))]. \quad (11)$$




FIG. 2. (a) Schematic phase diagram. (b) Coplanar RG flow of the variables I'_α (blue), $\bar{I} = (I_1 I_2 I_3)^{1/3}$ (green dashed), $(I_2 - I_1)/\bar{I}$ (red), and r (pink dotted). Curves are normalized to initial values at l_γ . Upper panel is for $J_{tt} \gg J_{hh}$ with $J_{hh} = 1$, $J_{tt} = 5$, $J_{th} = 0.4$, $T = 0.6$, and initial values $\bar{I} = 5.3$, $(I_2 - I_1)/\bar{I} = 0.27$, $r = 0.82$, $I'_\alpha = 1.2$. Decoupling is due to $(I_1 - I_2)/\bar{I} \rightarrow 0$. Lower panel is for $J_{hh} \gg J_{tt}$ with $J_{hh} = 5$, $J_{tt} = 1$, $J_{th} = 0.4$, $T = 0.5$, and initial values $\bar{I} = 4.5$, $(I_2 - I_1)/\bar{I} = 2.1$, $r = 0.11$, $I'_\alpha = 1.1$. Decoupling is due to $r \rightarrow 0$.

This is the well-known six state clock model that exhibits two consecutive BKT transitions [11]: one at $T_{\text{BKT}}^>$ that separates a high temperature disordered phase from a low temperature critical phase, where correlations $\langle \exp[i(\alpha(x) - \alpha(x'))] \rangle$ decay with a power-law in $|x - x'|$; a second at $T_{\text{BKT}}^<$ where the \mathbb{Z}_6 symmetry is spontaneously broken, leading to true long-range order with $\alpha = n\pi/3$ ($n \in \{1, \dots, 6\}$). It is crucial that the decoupling of the $U(1)$ phase occurs first, otherwise the $SO(3)$ sector would screen the long-range interactions between topological defects – vortices at $T_{\text{BKT}}^>$ or domain walls at $T_{\text{BKT}}^<$ that are responsible for the BKT transitions and the intermediate critical phase.

Following the RG program of the BKT problem for Eq. (11) we need to take into account that the size of the vortex is now given by the coplanar lengthscale $a_\gamma \gg a_0$ [34, 35]. We determine the vortex unbinding transition temperature $T_{\text{BKT}}^>$ implicitly via

$$I'_\alpha(T_{\text{BKT}}^>)^{-1} = \frac{\pi}{2 + 4\pi y(T_{\text{BKT}}^>)} \quad (12)$$

with fugacity $y = e^{-S_c} a_\gamma^2 / a_0^2$ and core action $S_c \simeq \pi \{1 + \min(K, K_t)\}$. From Eq. (12) we predict that $T_{\text{BKT}}^> \lesssim T_{\text{cp}}$, *i.e.*, the BKT transition is only numerically smaller than the coplanar crossover temperature. The system enters the critical phase soon after it becomes coplanar. Similarly, it follows from Ref. [11] that $T_{\text{BKT}}^<$ and $T_{\text{BKT}}^>$ are of the same order of magnitude.

In summary we have presented a 2D Heisenberg model on a decorated triangular lattice where short wavelength thermal fluctuations select long-range \mathbb{Z}_6 order. This is preceded in temperature by an emergent critical phase that is framed by two BKT transitions. We have written the action of this model as a classical 4D string theory where the spin stiffness is determined by the metric tensor of the manifold; the scaling equations are then extracted as components of the resulting Ricci flow. We note that the decoupling of the $U(1)$ degree of freedom corresponds to a dimensional reduction of the analogous string theory and thus to a toy model of compactification. Finally we note that the emergence of massless modes in collective mode massive theories could have interesting implications for two-dimensional field theories.

We acknowledge useful discussions with S. T. Carr, R. Fernandes, E. J. König, D. Nelson, V. Oganesyan, P. Ostrovsky, N. Perkins, J. Reuther, S. Sondhi, and O. Sushkov. PPO is supported by a Young Investigator Group of the KIT. This work was supported by DOE grant DE-FG02-99ER45790 (P. Coleman) and SEPNET (PC, PC and JS). PC, PC and JS acknowledge the hospitality of Royal Holloway, University of London where this work was begun.

Note added.— After obtaining these results we learned of two recent studies: one on a Kitaev-Heisenberg model, where an emergent \mathbb{Z}_6 -symmetry results from a conceptually different mechanism [36], a second on itinerant systems where an emergent \mathbb{Z}_4 Potts model appears [37].

-
- [1] P. C. Hohenberg, Phys. Rev., **158**, 383 (1967).
 - [2] N. D. Mermin and H. Wagner, Phys. Rev. Lett., **17**, 1307 (1966).
 - [3] P. Chandra, P. Coleman, and A. I. Larkin, Phys. Rev. Lett., **64**, 88 (1990).
 - [4] L. Capriotti, A. Fubini, T. Roscilde, and V. Tognetti, Phys. Rev. Lett., **92**, 157202 (2004).
 - [5] C. Weber, L. Capriotti, G. Misguich, F. Becca, M. Elhadj, and F. Mila, Phys. Rev. Lett., **91**, 177202 (2003).
 - [6] A. Mulder, R. Ganesh, L. Capriotti, and A. Paramekanti, Phys. Rev. B, **81**, 214419 (2010).
 - [7] C. L. Henley, Annu. Rev. Cond. Mat. Phys., **1**, 179 (2010).
 - [8] M. E. Zhitomirsky, Phys. Rev. B, **78**, 094423 (2008).
 - [9] G.-W. Chern and R. Moessner, arXiv:1207.4752v1 [cond-mat.str-el] (2012).
 - [10] R. M. Fernandes, L. H. VanBebber, S. Bhattacharya, P. Chandra, V. Keppens, D. Mandrus, M. A. McGuire, B. C. Sales, A. S. Sefat, and J. Schmalian, Phys. Rev. Lett., **105**, 157003 (2010).
 - [11] J. V. José, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson, Phys. Rev. B, **16**, 1217 (1977).
 - [12] G. Ortiz, E. Cobanera, and Z. Nussinov, Nucl. Phys. B, **854**, 780 (2012), ISSN 0550-3213.
 - [13] D. Blankschtein, M. Ma, A. N. Berker, G. S. Grest, and C. M. Soukoulis, Phys. Rev. B, **29**, 5250 (1984).

- [14] S. V. Isakov and R. Moessner, Phys. Rev. B, **68**, 104409 (2003).
- [15] L.-M. Duan, E. Demler, and M. D. Lukin, Phys. Rev. Lett., **91**, 090402 (2003).
- [16] M. Lubasch, V. Murg, U. Schneider, J. I. Cirac, and M.-C. Bañuls, Phys. Rev. Lett., **107**, 165301 (2011).
- [17] S. Trotzky, P. Cheinet, S. Fölling, M. Feld, U. Schnorrberger, A. M. Rey, A. Polkovnikov, E. A. Demler, M. D. Lukin, and I. Bloch, Science, **319**, 295 (2008).
- [18] Y.-A. Chen, S. Nascimbène, M. Aidelsburger, M. Atala, S. Trotzky, and I. Bloch, Phys. Rev. Lett., **107**, 210405 (2011).
- [19] H. C. Manoharan, C. P. Lutz, and D. M. Eigler, Nature (London), **403**, 512 (2000).
- [20] K. K. Gomes, W. Mar, K. W., F. Guinea, and H. C. Manoharan, Nature (London), **483**, 306 (2012).
- [21] C. L. Gao, W. Wulfskel, and J. Kirschner, Phys. Rev. Lett., **101**, 267205 (2008).
- [22] M. Waśniowska, S. Schröder, P. Ferriani, and S. Heinze, Phys. Rev. B, **82**, 012402 (2010).
- [23] J. Villain, J. Phys (Paris), **38** (1977).
- [24] C. L. Henley, Phys. Rev. Lett., **62**, 2056 (1989).
- [25] J. T. Chalker, P. C. W. Holdsworth, and E. F. Shender, Phys. Rev. Lett., **68**, 855 (1992).
- [26] D. Friedan, Phys. Rev. Lett., **45**, 1057 (1980).
- [27] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, U.K., 1999).
- [28] F. D. M. Haldane, Phys. Rev. Lett., **50**, 1153 (1983).
- [29] T. Dombre and N. Read, Phys. Rev. B, **39**, 6797 (1989).
- [30] S. Chakravarty, B. I. Halperin, and D. R. Nelson, Phys. Rev. B, **39**, 2344 (1989).
- [31] P. Azaria, B. Delamotte, and T. Jolicoeur, Phys. Rev. Lett., **64**, 3175 (1990).
- [32] We found the one-loop part also using Polyakov scaling.
- [33] R. S. Hamilton, J. Differential Geom., **17**, 255 (1982).
- [34] P. M. Chaikin and T. C. Lubensky, *Principles of condensed matter physics* (Cambridge University Press, Cambridge, U.K., 1995).
- [35] J. M. Fellows, S. T. Carr, C. A. Hooley, and J. Schmalian, [arXiv:1205.1333v1](https://arxiv.org/abs/1205.1333v1) (2012).
- [36] C. Price and N. B. Perkins, arXiv:1205:3967 (2012).
- [37] G.-W. Chern, R. M. Fernandes, R. Nandkishore, and A. V. Chubukov, arXiv:1203.5776 (2012).